Research Article

Role of collagen-like protein BclA3 in the assembly of the exosporium layer of *Clostridioides*

difficile spores.

Marjorie Pizarro-Guajardo [2,3], César Ortega-Lizárraga[1], Francisca Cid-Rojas[1,2,3], Ana Inostroza[1]

and Daniel Paredes-Sabja [1,2,3*].

[1]Microbiota-Host Interactions and Clostridia Research Group, Facultad de Ciencias de la Vida,

Universidad Andrés Bello, Santiago, Chile.

[2]Millenium Nucleus in the Biology of the Intestinal Microbiota, Santiago, Chile.

[3] Department of Biology, Texas A&M University, TX, USA.

Running Title: *Clostridium difficile* BclA3 characterization.

Key Words: *Clostridium difficile*, Spore, BclA3.

[*]Corresponding author:

Dr. Daniel Paredes-Sabja, Department of Biology, Texas A&M University, College Station, TX,

77843, USA. E-mail: dparedes-sabja@bio.tamu.edu

Abstract

Newly formed spores are essential for persistence of *C. difficile* in the host, transmission to a new susceptible host (Deakin *et al.*, 2012b) and recurrence of CDI. BclA3 and BclA2 Spore surface proteins are expressed during sporulation under the control of mother-cell specific sigma factors of the RNA polymerase, SigE and SigK. Deletion of *bclA3* leads to spores with an electron-dense exosporium layer that lacks bump-like structures in the electron-dense layer and hair-like projections, both structures typically found in the wild type spore. Therefore, in this work, we have addressed the role of the exosporium collagen-like BclA3 glycoprotein in the assembly of the exosporium layer. Immunogold labelling of BclA2$_{CTD}$ and BclA3$_{CTD}$ indicates that both proteins are located in the hairs, with BclA2 located outermost of BclA3. Absence of BclA3 leads to spores with no hair-like projections, and absence of bumps in thick exosporium spores, a phenotype also expressed in by the deletion of the collagen-like region of BclA3. Overall, these results provide insights into the role of BclA3 in the assembly of the exosporium layer of *C. difficile* spores.

Introduction

Clostridioides difficile is the leading cause of nosocomial antibiotic-associated diarrhea (Martin *et al.*, 2016, Rupnik *et al.*, 2009). Newly formed spores are essential for persistence of *C. difficile* in the host, transmission to a new susceptible host (Deakin *et al.*, 2012b) and recurrence of CDI, which is a major problem with an incidence of 25 to 65% of patients, associated to increase severity of symptoms and time in hospital settings (Lessa *et al.*, 2015). The surface of *C. difficile* spores play an important role in host-spore interaction (Mora-Uribe *et al.*, 2016, Calderón-

Romero *et al.*, 2018, Paredes-Sabja & Sarker, 2012). In the spore of epidemic *C. difficile* strains,

such as RT27 and RT78, the most external layer, the exosporium, is surrounded by an

ultrastructure defined as hair-like projection, which appears as a layer of ~80 nm of disheveled

hairs, differently to the hair-like nap in *B. anthracis* spore, where all hairs are extended to the

exterior (Thompson *et al.*, 2011). However, hair-like extensions are absent in 630 strain spores

(Paredes-Sabja *et al.*, 2014). The role of this structure in pathogenesis is unknown.

Spore surface proteins are expressed during sporulation under the control of mother-cell

specific sigma factors of the RNA polymerase, SigE and SigK (Fimlaid *et al.*, 2013). By

sonication of *C. difficile* spores, trypsin digestion of extracted proteins and MS/MS, spore coat

and exosporium proteins have been identified (Díaz-González *et al.*, 2015). Some of the surface

spore proteins are collagen-like proteins, characterized by a large inner domain with several

GXY repeated pattern, flanked by globular NTD and CTD (Díaz-González *et al.*, 2015, Pizarro-

Guajardo *et al.*, 2014). Collagen-like proteins are present in several spore structures, such as

appendages in *Clostridium teaniosporium* (Steichen *et al.*, 2003) and external nap in *Bacillus*

anthracis (Steichen *et al.*, 2003), where collagen-like proteins tend to form trimers by the

globular carboxi-terminal domain (CTD) that allow the formation of triple helical collagen-like

structures (Steichen *et al.*, 2003, Yu *et al.*, 2014).

In *C. diffi*cile genome, three homologues of the *bclA* loci are present in most *C. difficile*

strains (Pizarro-Guajardo *et al.*, 2014). However, most epidemically-relevant strains belonging to

ribotype 027 have only full-length *bclA2* and *bclA3* open reading frames, while a nonsense

mutation allows the expression of a pseudogenized *bclA1* (Pizarro-Guajardo *et al.*, 2014,

Phetcharaburanin *et al.*, 2014, Hosseini, 2018). A previous study demonstrated that insertional

inactivation of *bclA1* and *bclA2* genes lead to spores with a defectively assembled electron-dense

exosporium layer in strain 630, but normal morphology for the mutant in *bclA3*

(Phetcharaburanin *et al.*, 2014). For epidemic strain R20291, our results indicate that a deletion

of *bclA3* leads to spores with an electron-dense exosporium layer that lacks bump-like structures

in the electron-dense layer and hair-like projections, both structures typically found in the wild

type spore (Castro-Córdova *et al.*, 2020). The gene *bclA3* is the second element in a bicistronic

operon, where the first cistron, *sgtA*, has been characterized as a glycosyltransferase enzyme that

is required for the glycosylation of collagen-like fragments of BclA3 (Aubry *et al.*, 2020, Strong

et al., 2014).

Therefore, in this work, we have addressed the role of the exosporium collagen-like

BclA3 glycoprotein in the assembly of the exosporium layer. Immunogold labelling of BclA2$_{CTD}$

and BclA3$_{CTD}$ indicates that both proteins are located in the hairs, with BclA2 located outermost

of BclA3. Absence of BclA3 leads to spores with no hair-like projections, and absence of bumps

in thick exosporium spores, a phenotype also expressed in by the deletion of the collagen-like

region of BclA3. Overall, these results provide insights into the role of BclA3 in the assembly of

the exosporium layer of *C. difficile* spores.

Materials and Methods

Bacterial strains and growth conditions. *C. difficile* strains (see Supplementary Table 1) were

routinely grown at 37 °C under anaerobic conditions in a Bactron III-2 anaerobic chamber

(Shellab, USA) in BHIS medium: 3.7% weight vol−1 brain heart infusion broth (BD, USA)

supplemented with 0.5% weight vol−1 yeast extract (BD, USA) and 0.1% weight vol−1 L-

cysteine (Merck, USA) or on BHIS agar plates. Escherichia coli strains were routinely grown

aerobically at 37 °C under aerobic conditions with shaking at $1 \times g$ in Luria-Bertani medium

(BD, USA), supplemented with 25 μg mL−1 chloramphenicol (Merck, USA), where appropriate.

C. difficile **mutant construction of mutant strain in the CLR domain in** *bclA3* **gene**

(***bclA3*$_{\Delta CLR}$**)**.** In order to construct a R20291 *C. difficile* strains which lacks the CLR domain in

bclA3 gene (*bclA3*$_{\Delta CLR}$), *C. difficile* genome FN545816 from the EMBL/GenBank databases was

employed for primer design. In-frame deletions were made by allelic exchange using *pyrE* alleles

(Ng *et al.*, 2013). To remove the CLR segment from *bclA3* locus, left homology arm was

constructed by amplification of 1676 bp fragment including 1529 nt upstream *bclA3* start codon,

first 138 nt which codify 46 aa corresponding to amino-terminal domain (NTD), plus 9 nt

corresponding to the first GXX of collagen-like region, flanked by 5' SbfI and 3' BamHI

restriction sites, which was cloned in those restriction sites into plasmid pMTL-YN4. For right

homology arm, a 1042 nt fragment, starting at 1516 nt downstream start codon, was amplified

with 5' BamHI and 3' AscI restriction sites, and cloned in those restriction sites into pMTL-YN4

containing left homology arm. To verify the correct construction of the plasmids, all constructs

were Sanger sequenced.

The plasmid obtained were transformed into *E. coli* CA434 and mated with *C. difficile*

R20291 *ΔpyrE* (Ng *et al.*, 2013). *C. difficile* transconjugants were selected by sub-culturing on

BHIS agar containing 15 µg/mL thiamphenicol (Sigma-Aldrich USA), 8 µg/mL cefoxitin

(Sigma-Aldrich USA) and 250 µg/mL cefoxitin, and re-streaked five times. In order to select for

plasmid excision, the single-crossover mutants identified were streaked onto CDMM (Karasawa

et al., 1995) with 1.5% agar supplemented with 2 mg/mL 5-Fluoroorotic acid (US Biological,

USA) and 5 µg/mL uracil (Sigma-Aldrich USA). Confirmation of plasmid excision was made by

negative selection in BHIS-15% thiamphenicol plates. The isolated FOA-resistant and

thiamphenicol sensitive colonies were screened using the primer pair P664 (FP-bclA3-detect) /

P665 (RP-bclA3-detect) for the *bclA3* mutant. The mutant was whole genome sequenced to

confirm the genetic background and that no additional SNPs were introduced during the genetic

manipulation. For correction of the *pyrE* mutation, transconjugants with pMTL-YN2C were

streaked onto minimal media without uracil or FOA supplementation, and developed colonies

were analysed further. To verify correct insertion, genomic DNA was extracted and sequenced.

Spore purification of *C. difficile* R20291. *C. difficile* strains used in this study are listed in

Table 1. To produce pure spores, *C. difficile* was routinely grown under anaerobic conditions on

BHIS broth (3.7% Brain Heart Infusion supplemented with 0.5% yeast extract, 1% cysteine). 16

h cultures were diluted to 1:500 dilution and 100 ul were plated onto 70:30 (63 g Bacto peptone,

3.5 g protease peptone, 11.1 g BHI medium, 1.5 g yeast extract, 1.06 g Tris, 0.7 g NH_4SO_4, 15 g

agar per litter) or TY (3% Trypticase Soy-0.5% yeast extract) agar plate and incubated at 37 °C

for 7 days. After incubation, plates were scraped up with 1 ml of ice-cold sterile water. Spores

were gently washed five times (resuspension in ice-cold sterile water and centrifugation at

16,000 × g for 5 min). Next, spores were gently loaded onto a 1.5 ml tube containing 300 μl of

45% Nycodenz solution and centrifuged for 40 min at 16,000 × g. After centrifugation, the

supernatant was removed, and the spore pellet was washed five times (Calderón-Romero *et al.*,

2018). The spores were counted in a Neubauer chamber and concentration adjusted at 5×10^9

spores/ml to store at -80°C until use.

Transmission electronic microscopy (TEM). Spores $(2x10^8)$ were fixed with 3%

glutaraldehyde on 0.1 M cacodylate buffer (pH 7.2), incubated overnight at 4°C, and stained for

30 min with 1% tannic acid. Samples were further embedded in a Spurr resin (Paredes-Sabja *et*

al., 2012). Thin sections of 90 nm were obtained with a microtome, placed on glow discharge

carbon-coated grids and double lead stained with 2% uranyl acetate and lead citrate. Spores were

analyzed with a Philips Tecnai 12 Bio Twin microscope at Unidad de Microscopía Avanzada at

the Pontificia Universidad Católica de Chile. ImageJ software was employed to calculate

thickness of spore layers.

Immunogold. To locate BclA2 and BclA3 in the spore, 4×10^7 R20291 spores were incubated in

PBS–0.2% BSA containing 1:200 mouse serum from immunized animal with BclA2$_{CTD}$ and

BclA3CTD, respectively, for 1 hr at 37 ℃. The excess of antibody was eliminated by three

cycles of centrifugation at 13,000 × g for 5 min and resuspension in PBS–0.1% BSA. Spore

suspensions were then incubated for 1 h with 1:20 donkey anti-mouse IgG antibody coupled to

12-nm gold particles (Sigma G7777) in PBS–1% BSA for 1 hr at RT, and were washed by

centrifugation at 13,000 × g for 5 min. Subsequently, samples were fixed and processed, as

described. ImageJ software was employed to calculate distance from spore surface to gold label.

Results

Even length of the filaments in thick and thin exosporium spores. Hair-like projections are filament ultrastructures located at the surface of most hypervirulent *C. difficile* spores (Pizarro-Guajardo *et al.*, 2016a, Pizarro-Guajardo *et al.*, 2016b, Paredes-Sabja *et al.*, 2014, Escobar-Cortes *et al.*, 2013). As previously described, *C. difficile* sporulation process generates two distinctive exosporium morphotypes, thick and thin exosporium (Pizarro-Guajardo *et al.*, 2016a, Pizarro-Guajardo *et al.*, 2016b, Paredes-Sabja *et al.*, 2014), which differs in the electron-dense layer; while a ~20 nm electron-dense layer surrounds the coat spore in thin exosporium spore, the electron-dense layer is made of bumps that can reach ~100 nm . Despite this difference, both morphotypes share the hair-like projection ultrastructure (Pizarro-Guajardo *et al.*, 2016b, Pizarro-Guajardo *et al.*, 2016a). It remains unclear if the site of anchorage of these filaments is the coat with the electron-dense bumps being formed surrounding these filaments, or otherwise, the filaments are located on the surface of the bumps, in the thick exosporium spore.

In order to identify the assembly of hair-projections in the spore surface, we revisited micrograph of R20291 spores and measure the length of these filaments in both type of spores, and for thick exosporium spores we compared the filament length in the highest and lowest point of the bump. The filament length is similar in thick and thin exosporium spore, of 96 nm in thin exosporium spores. For thick exosporium spores, the filament is 91nm in the highest point of the bump and 89 nm in the lowest point, with no significant difference (Figure 1A). In the other hand, the electron-dense layer in thick exosporium spores varies from 70 nm to 27 nm between the highest and lowest point of the bump (Figure 1B). Interestingly, the lowest point of the bumps has an electron-dense thickness similar to what is found in thin exosporium spore, of 20 nm. These results are visually observable in an electron-micrograph. Al these findings suggest

that hairs are assembled over the electron-dense layer of the exosporium and over the conformed

bumps in the case of thick exosporium spores.

Collagen-like proteins BclA2 and BclA3 are part of the hair-like projections of *C. difficile*

spores. R20291 strain encodes two collagen-like proteins, BclA2 and BclA3 (Pizarro-Guajardo

et al., 2014). These proteins have a central collagen-like region, a similar size CTD, and a

different size NTD (i.e. BclA2 has a shorter NTD compared to BclA3, 5 versus 46 amino acid-

length) (Figure 2a). We have previously shown that in strain 630, which has a functional BclA1

protein, the NTD seems to be oriented in an inward orientation, with the CTD oriented outwards

(Pizarro-Guajardo *et al.*, 2014). In addition, transmission electron micrographs of *bclA3* mutant

spores demonstrates that BclA3 is required for the formation of the hair-like projections (Castro-

Córdova *et al.*, 2021); however, it is unclear whether these hair-like projections are formed by

BclA2 and/or BclA3.

To solve this question, we employed antibodies against $BclA2_{CTD}$ and $BclA3_{CTD}$ to tag

these proteins in *C. difficile* spore in an immunogold assay. Immunoelectron micrographs

demonstrate that both antibodies labelled the tip of the hair-like projections in both exosporium

morphotypes (thick and thin exosporium) (Figure 2B and Figure 2C), indicating that BclA2 and

BclA3 are part of the hair-like projections, and that both CTD are oriented outwards. Next, we

quantified the distance of the labelled CTD of BclA2 and BclA3 to the electron dense surface of

spores, which indicates that BclA2 could have a location 25 nm outer that BclA3 (Figure 2D).

This outer position of $BclA2_{CTD}$ respect to $BclA3_{CTD}$ is independent of the exosporium type.

Thick and thin-exosporium layer *C. difficile* spores lack the hair-like projections in the

absence of BclA3. Our previous work demonstrated that inactivation of BclA3 lead to the

absence of hair-like projections. While in the wild type strain, hair-like projections and

exosporium electron-dense bumps were present in thick and thin exosporium spores; as

previously described (Castro-Córdova *et al.*, 2020), Δ*bclA3* spores exhibited no hair-like

projections in thick and thin exosporium spores (Figure 3A). Notably, we identified in thick

exosporium spores a smooth electron-dense layer without bumps (Figure 3A). Upon

complementation of the Δ*bclA3* mutant strain with *bclA3* into the *pyrE* loci, we observed that

spores of the Δ*bclA3/bclA3+* strain restored the hair-like projections and electron dense bump

formation (Figure 3A, B), demonstrating that BclA3 is required for the formation of the hair-like

projections and electron-dense bumps of the exosporium layer. Upon analysis of the spore-coat

thickness, we observed that BclA3 had no impact in the thickness of the spore-coat (Figure 3A

and 3B). Collectively, these results demonstrate that the role of BclA3 in external layers

morphology is restricted to the morphology of the exosporium, due to the absence of alterations

in spore coat layer. Specifically, BclA3 plays a key morphological role in the formation of the

electron-dense bumps and hair-like projections.

Lack of BclA3 leads to enrichment of *C. difficile* spores with appendages. Another feature of

C. difficile spores that has been recently described is the presence of appendages in the spore

poles (Pizarro-Guajardo *et al.*, 2020, Antunes *et al.*, 2018). Appendages are defined as a

prolongation of the exosporium in one polar region of the spore (Pizarro-Guajardo *et al.*, 2020,

Antunes *et al.*, 2018), which is dependent on CdeM and is related to faster germination (Antunes

et al., 2018). In order to evaluate impact of BclA3 in the appendage frequency, we used phase

contrast microscopy to characterize populations of wild type and Δ*bclA3* mutant spores. Phase

contrast microscopy shows polar appendage in two different morphologies: appendage type 1, a

polar prolongation, and appendage type 2, diffuse and round prolongation (Figure 4A). In the

wild type strain, most of the spores (84%) had no appendage, while the rest of the spores were

distributed between App-1 (8.9%) and App-2 (6.4%). In contrast, for the *ΔbclA3* mutant, the

spores lacking appendage decreased to 59%, while spores with App-1 and App-2 increased to

18% and 23%, respectively (Figure 4B). Notably, restoration of *bclA3* restores the spore

proportions of appendages to similarly levels as wild type strain (Figure 4B), indicating that

presence of BclA3 in the exosporium spore favors the spore type without appendage.

The collagen-like region of BclA3 is required for the formation of the hair-like projections

and electron-dense bumps of the exosporium layer. Work in *Bacillus anthracis* has

demonstrated that the length of the hairs of the exosporium layers depended on the collagen-like

region (CLR) of the BclA glycoprotein (Sylvestre *et al.*, 2003). Since our results demonstrate

that the hair-like projections are formed by BclA2 and BclA3, but *bclA3* is essential for the

formation of the hair-like projections, we asked if the CLR of *C. difficile* BclA3 is required for

the formation of these projections. To assess this, we constructed a *C. difficile* strain with a

deletion in the CLR region of the *blcA3* gene (*bclA3*$_{\Delta CLR141-1509}$) (Figure 5A). Transmission

electron micrographs demonstrate that while long hair-like projections are exhibited in wild type

spores, mutant strain lacking BclA3 CLR domain did not exhibit hair-like projections or

electron-dense bumps in the exosporium layer; *bclA3*$_{\Delta CLR141-1509}$ spores resembled *ΔbclA3* spore

ultrastructural phenotype (Figure 5B). These results demonstrate that the CLR of BclA3 is

essential for the proper formation of the hair-like projection structures and bump formation in the

exosporium of *C. difficile* spores.

Discussion.

During CDI, *C. difficile* spore formation is essential in the recurrence of the disease (Deakin *et al.*, 2012a), and the underlying mechanisms that correlate *C. difficile* spore persistence and recurrence of the disease stared to be unveiled. In this sense, the outermost layer of the spore, the exosporium, is critical to understand CDI pathogenesis and developing of novel therapies. Recently, a novel mechanism for interaction between the spore and the intestinal mucosa has been described, in which BclA3 is essential for the formation of the hair-like projections, but we know little of the role in exosporium assembly and spore properties (Castro-Córdova *et al.*, 2020). Therefore, BclA3 has major contribution in *C. difficile* persistence and initiation of recurrence, that suggest that blocking its interactions with the host environment could improve the clinical outcome of recurrence. In this work, we further dissected the morphogenic role of collagen-like proteins BclA3 in the exosporium of *C. difficile* spore. Previous work on the role of BclA3 in C. difficile spore assembly made on the strain 630 cannot be complementary with these results, due to the lack of hairs in 630 strain.

The first major finding of this work is the labeling of $BclA2_{CTD}$ and $BclA3_{CTD}$ at the tip of the hair-like extension, that indicates that hair-like extensions are made by both, BclA2 and BclA3. A mutant $\Delta bclA3$ strain spore has no capability to form hair-like extensions (Castro-Córdova *et al.*, 2020), indicating that the presence of only BclA2 is not sufficient to build the hair-like projection structure. However, it is described that BclA2 can form collagen-like structures susceptible to digestion by collagenase (Pizarro-Guajardo *et al.*, 2014). As only BclA3 is essential for hair-like projection structure, and $BclA2_{CTD}$ was tagged in the structure, it suggests that BclA2 is capable to adhere to hair-like structures once they are formed by BclA3.

Whether BclA2 is forming collagen-like structure with BclA3 after BclA3 spore assembly

remains unknown. The location of $BclA2_{CTD}$ more external than $BclA3_{CTD}$ suggest that BclA3 is

necessary to assemble of BclA2 in the collagen structure, which is supported by the lack of hair

when BclA2 is transcribed but not *bclA3* in the mutant Δ*bclA3*. Detection of $BclA2_{CTD}$ in the

spore Δ*bclA3* suggest that BclA2 can assemble in the spore independently of BclA3, but in this

assemble is not capable of structure hairs.

Bump formation in exosporium seems to be a consequence of hair structure, since all

mutations affecting BclA3 *(Δbclא3*, lack of collagen-like repeat domain) impair the hair

formation, but also prevent the bumps structure to form in the exosporium. It is interesting to

observe that mutant strain lacking the collagen rich region of BclA3, the CLR, lacks the hair-like

extension but the exosporium shows a reduced bump exosporium, which suggest that a slightly

development of exosporium bump, but not all, is developed in the presence of $BclA3_{NTD\text{-}CTD}$

only. It is described that only NTD of BclA3 and BclA2 are sufficient to anchor the protein to

the spore (Pizarro-Guajardo *et al.*, 2014), which suggest that CTD domain of both proteins is

exposed to the surface, as confirmed by immunogold experiments in this work.

5. Acknowledgements

This work was supported by FONDECYT Regular 1191601, FONDECYT Regular 1151025,

and Millennium Science Initiative Program–NCN17_093, and Fondo de Fomento al Desarrollo

Científico y Tecnológico (FONDEF) ID18|10230 to M.P-G and D.P-S. Support was also

provided by start-up funds from Texas A&M University to D.P-S.

References

Antunes, W., Pereira, F.C., Feliciano, C., Saujet, L., Vultos, T.d., Couture-Tosi, E., Pechine, S., Bruxelle, J.-F., Janoir, C., Melo, L., Brito, P., Martin-Verstraete, I., Serrano, M., Dupuy, B., and Henriques, A.O. (2018) Structure and assembly of a *Clostridioides difficile* spore polar appendage. *bioRxiv*: 468637.

Aubry, A., Zou, W., Vinogradov, E., Williams, D., Chen, W., Harris, G., Zhou, H., Schur, M.J., Gilbert, M., Douce, G.R., and Logan, S.M. (2020) *In vitro* Production and Immunogenicity of a *Clostridium difficile* Spore-Specific BclA3 Glycopeptide Conjugate Vaccine. *Vaccines (Basel)* **8**.

Calderón-Romero, P., Castro-Córdova, P., Reyes-Ramírez, R., Milano-Céspedes, M., Guerrero-Araya, E., Pizarro-Guajardo, M., Olguín-Araneda, V., Gil, F., and Paredes-Sabja, D. (2018) *Clostridium difficile* exosporium cysteine-rich proteins are essential for the morphogenesis of the exosporium layer, spore resistance, and affect *C. difficile* pathogenesis. *PLOS Pathogens* **14**: e1007199.

Castro-Córdova, P., Mora-Uribe, P., Reyes-Ramírez, R., Cofré-Araneda, G., Orozco-Aguilar, J., Brito-Silva, C., Mendoza-León, M.J., Kuehne, S.A., Minton, N.P., Pizarro-Guajardo, M., and Paredes-Sabja, D. (2020) *Clostridioides difficile* spore-entry into intestinal epithelial cells contributes to recurrence of the disease. *Nature Communications*: Accepted.

Castro-Córdova, P., Mora-Uribe, P., Reyes-Ramírez, R., Cofré-Araneda, G., Orozco-Aguilar, J., Brito-Silva, C., Mendoza-León, M.J., Kuehne, S.A., Minton, N.P., Pizarro-Guajardo, M., and Paredes-Sabja, D. (2021) Entry of spores into intestinal epithelial cells contributes to recurrence of *Clostridioides difficile* infection. *Nat Commun* **12**: 1140.

Deakin, L.J., Clare, S., Fagan, R.P., Dawson, L.F., Pickard, D.J., West, M.R., Wren, B.W., Fairweather, N.F., Dougan, G., and Lawley, T.D. (2012a) The *Clostridium difficile spo0A* gene is a persistence and transmission factor. *Infect Immun* **80**: 2704-2711.

Deakin, L.J., Clare, S., Fagan, R.P., Dawson, L.F., Pickard, D.J., West, M.R., Wren, B.W., Fairweather, N.F., Dougan, G., and Lawley, T.D. (2012b) The *Clostridium difficile spo0A* Gene Is a persistence and transmission factor. *Infection and Immunity* **80**: 2704-2711.

Díaz-González, F., Milano, M., Olguin-Araneda, V., Pizarro-Cerda, J., Castro-Córdova, P., Tzeng, S.-C., Maier, C.S., Sarker, M.R., and Paredes-Sabja, D. (2015) Protein composition of the outermost exosporium-like layer of *Clostridium difficile* 630 spores. *Journal of Proteomics* **123**: 1-13.

Escobar-Cortes, K., Barra-Carrasco, J., and Paredes-Sabja, D. (2013) Proteases and sonication specifically remove the exosporium layer of spores of *Clostridium difficile* strain 630. *J Microbiol Methods* **93**: 25-31.

Fimlaid, K.A., Bond, J.P., Schutz, K.C., Putnam, E.E., Leung, J.M., Lawley, T.D., and Shen, A. (2013) Global analysis of the sporulation pathway of *Clostridium difficile*. *PLoS Genet* **9**: e1003660.

Hosseini, S., (2018) Biological Containment of Recombinant Spores and Dissemination of Pathogenic Spores. In: School of Biological Sciences. United Kingdom: Royal Holloway, University of London., pp. 216.

Karasawa, T., Ikoma, S., Yamakawa, K., and Nakamura, S. (1995) A defined growth medium for Clostridium difficile. *Microbiology* **141 (Pt 2)**: 371-375.

Lessa, F.C., Mu, Y., Bamberg, W.M., Beldavs, Z.G., Dumyati, G.K., Dunn, J.R., Farley, M.M.,
Holzbauer, S.M., Meek, J.I., Phipps, E.C., Wilson, L.E., Winston, L.G., Cohen, J.A.,
Limbago, B.M., Fridkin, S.K., Gerding, D.N., and McDonald, L.C. (2015) Burden of
Clostridium difficile infection in the United States. *N Engl J Med* **372**: 825-834.
Martin, J.S., Monaghan, T.M., and Wilcox, M.H. (2016) *Clostridium difficile* infection:
epidemiology, diagnosis and understanding transmission. *Nat Rev Gastroenterol Hepatol*
13: 206-216.
Mora-Uribe, P., Miranda-Cardenas, C., Castro-Cordova, P., Gil, F., Calderon, I., Fuentes, J.A.,
Rodas, P.I., Banawas, S., Sarker, M.R., and Paredes-Sabja, D. (2016) Characterization of
the adherence of *Clostridium difficile* spores: the integrity of the outermost layer affects
adherence properties of spores of the epidemic strain R20291 to components of the
intestinal mucosa. *Front Cell Infect Microbiol* **6**: 99.
Ng, Y.K., Ehsaan, M., Philip, S., Collery, M.M., Janoir, C., Collignon, A., Cartman, S.T., and
Minton, N.P. (2013) Expanding the repertoire of gene tools for precise manipulation of
the *Clostridium difficile* genome: allelic exchange using *pyrE* alleles. *PLoS One* **8**:
e56051.
Paredes-Sabja, D., Cofre-Araneda, G., Brito-Silva, C., Pizarro-Guajardo, M., and Sarker, M.R.
(2012) *Clostridium difficile* spore-macrophage interactions: spore survival. *PLoS One* **7**:
e43635.
Paredes-Sabja, D., and Sarker, M.R. (2012) Adherence of *Clostridium difficile* spores to Caco-2
cells in culture. *Journal of Medical Microbiology* **61**: 1208-1218.
Paredes-Sabja, D., Shen, A., and Sorg, J.A. (2014) *Clostridium difficile* spore biology:
sporulation, germination, and spore structural proteins. *Trends Microbiol* **22**: 406-416.
Phetcharaburanin, J., Hong, H.A., Colenutt, C., Bianconi, I., Sempere, L., Permpoonpattana, P.,
Smith, K., Dembek, M., Tan, S., Brisson, M.C., Brisson, A.R., Fairweather, N.F., and
Cutting, S.M. (2014) The spore-associated protein BclA1 affects the susceptibility of
animals to colonization and infection by *Clostridium difficile*. *Mol Microbiol* **92**: 1025-
1038.
Pizarro-Guajardo, M., Calderon-Romero, P., Castro-Cordova, P., Mora-Uribe, P., and Paredes-
Sabja, D. (2016a) Ultrastructural Variability of the Exosporium Layer of *Clostridium*
difficile Spores. *Appl Environ Microbiol* **82**: 2202-2209.
Pizarro-Guajardo, M., Calderon-Romero, P., and Paredes-Sabja, D. (2016b) Ultrastructure
Variability of the Exosporium Layer of *Clostridium difficile* Spores from Sporulating
Cultures and Biofilms. *Appl Environ Microbiol* **82**: 5892-5898.
Pizarro-Guajardo, M., Calderón, P., Romero-Rodriguez, A., and Paredes-Sabja, D. (2020)
Characterization of exosporium layer variability of *Clostridioides difficile* spores in the
epidemically relevant strain R20291. *Front Microbiol* **11**: 1345.
Pizarro-Guajardo, M., Olguin-Araneda, V., Barra-Carrasco, J., Brito-Silva, C., Sarker, M.R., and
Paredes-Sabja, D. (2014) Characterization of the collagen-like exosporium protein,
BclA1, of *Clostridium difficile* spores. *Anaerobe* **25**: 18-30.
Rupnik, M., Wilcox, M.H., and Gerding, D.N. (2009) *Clostridium difficile* infection: new
developments in epidemiology and pathogenesis. *Nat Rev Microbiol* **7**: 526-536.
Steichen, C., Chen, P., Kearney, J.F., and Turnbough, C.L. (2003) Identification of the
immunodominant protein and other proteins of the *Bacillus anthracis* exosporium. *J*
Bacteriol **185**: 1903-1910.

Strong, P.C., Fulton, K.M., Aubry, A., Foote, S., Twine, S.M., and Logan, S.M. (2014)
Identification and characterization of glycoproteins on the spore surface of *Clostridium*
difficile. *J Bacteriol* **196**: 2627-2637.
Sylvestre, P., Couture-Tosi, E., and Mock, M. (2003) Polymorphism in the collagen-like region
of the Bacillus anthracis BclA protein leads to variation in exosporium filament length. *J*
Bacteriol **185**: 1555-1563.
Thompson, B.M., Binkley, J.M., and Stewart, G.C. (2011) Current physical and SDS extraction
methods do not efficiently remove exosporium proteins from Bacillus anthracis spores. *J*
Microbiol Methods **85**: 143-148.
Yu, Z., An, B., Ramshaw, J.A., and Brodsky, B. (2014) Bacterial collagen-like proteins that form
triple-helical structures. *J Struct Biol* **186**: 451-461.

Figure legends

Figura 1. Even length of hair-like projection layer in thick and thin exosporium spores. A)

Hair-like projection length and B) electron-dense layer thickness in thick (blue) and thin (black)

exosporium spores. T-test. Each dot represent one spore, as the average of 4 measures on each

spore. Thick-high: filament length from the highest point of the bump until the end of the

filament. Thick-Low: filament length from the lowest point of the bump until the end of the

filament. C) Schematic representation of ultrastructure in the spore external layer for thick and

thin Exosporium spores. TEM micrograph of the structures. Bar scale: 200 nm. a, hair-like

projection; b, electron-dense layer, c, coat.

Figure 2. Immunogold labeling of hair-like extensions trough BclA2 and BclA3 targeting in

R20291 spores. A) Schematic representation of domains in BclA2 and BclA3. Red: NTD, green:

CTD, yellow: collagen like repeats domain. Numbers indicate the aminoacidic position where

domains end. B) Targeting of hair-like extensions with anti-BclA2 antibody and C) anti-BclA3

antibody, in wild type R20291. Right panel is a zoom of left panel. Bar scale: 200 nm. Black

squared dot shows the zoomed area. D) Distance from the surface of the spore to the gold label.

Every dot represent one spore, as the average distance of 1 to 9 gold particles to the electron-

dense surface. Statistical test: Unpaired t-test.

Figure 3. BclA3 have a role in hair-like ultrastructure in R20291 spores. A) Purified spores

from R20291, *ΔbclA3* and complemented. *C. difficile* strains were fixed and analyzed by

transmission electron microscopy. Spores from wild type, *ΔbclA3* and *ΔbclA3/bclA3+* were

classified as thick or thin exosporium. Scale bar: 200nm. B) Thickness of each layer (hair,

exosporium and coat) was measured at 6 different points (n = 7; 6 measures per spore). 2-way

ANOVA, Turkey's multiple comparison test.

Figure 4. Appendage occurrence in R20291 and bclA3 mutant spores. A) Example of phase-

contrast microscopy spores showing no-appendage (No-app), appendage type 1 (App-1) and

appendage type 2 (App-2). Scale bar: 1 μm. B) Percentage of appendage type. N = 727 wild

type, 985 *bclA3-*, 832 *bclA3-/bclA3+*. 2-way ANOVA, Multiple comparison test.

Figure 5. Lack of collagen-like domain in bclA3 gene impairs the hair formation and

impacts exosporium morphology. A) Schematic representation of reverse R20291 DNA strand

in wild type strain and mutant in CLR portion of *bclA3* gene (*bclA3* $\Delta CLR_{142-1509}$) in reverse

strand. B) TEM imagen of thick and thin exosporium spores of *bclA3* $\Delta CLR_{142-1509}$, showing

hair-exosporium lack and lack of bumps. Scale bar: 200 nm.

Tables

Table S1. Primers used in this work to create SNAP fusions.

Table S2. Plasmids and strains used in this work

Figure 1

Figure 2

A

B Anti-BclA2$_{CTD}$ **C** Anti-BclA3$_{CTD}$ **D**

Figure 3

A

Thick Exosporium   Thin Exosporium

Wild type

Δ*bclA3*

Δ*bclA3/bclA3*+

B

Thick exosporium spore

Thin exosporium spore

Length (nm)

Hair   Exosporium   Coat

Wild type
Δ*bclA3*
Δ*bclA3/bclA3*+

Figure 4

A

B

Figure 5

A

Wild type

bclA3 ΔCLR$_{141-1509}$

0.5 kb

B

Thick Exosporium   Thin Exosporium

Wild type

bclA3 ΔCLR$_{141-1509}$