1

The green tea catechin EGCG provides proof-of-concept for a pan-coronavirus entry inhibitor

Emmanuelle V. LeBlanc[1] and Che C. Colpitts[1,#]

[1]Department of Biomedical and Molecular Sciences, Queen's University, Kingston, ON K7L 3N6, Canada

[#]Correspondence: che.colpitts@queensu.ca

Abstract

The COVID-19 pandemic caused by the severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) has emphasized the serious threat to human health posed by emerging coronaviruses. Effective antiviral countermeasures are urgently needed as vaccine development against an emerging coronavirus takes time, even in the best-case scenario. The green tea catechin, epigallocatechin gallate (EGCG), has broad spectrum antiviral activity. We demonstrate here that EGCG prevents murine and human coronavirus infection and blocks the entry of lentiviral particles pseudotyped with spike proteins from bat or highly pathogenic coronaviruses. We show that EGCG treatment reduces coronavirus attachment to target cell surfaces. Our results demonstrate the potential for the development of pan-coronavirus attachment inhibitors.

Key words

SARS-CoV-2, coronavirus, entry inhibitor, natural product

Introduction

The novel severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) has infected hundreds of millions of people and has resulted in millions of deaths around the world. The COVID-19 pandemic, caused by SARS-CoV-2, underscores the severe health threat posed by emerging coronaviruses (CoVs), and makes evident the paucity of antivirals with extended spectrums of activity. SARS-CoV-2 is the third highly pathogenic CoV to emerge in the 21st century. The abundance and diversity of SARS-related CoVs in bats points to the likelihood of future spillover into human populations [1]. In addition, seasonally circulating common cold CoVs continually burden human health [2], and cause severe disease in elderly populations [3,4], without effective preventative or therapeutic strategies. In order to develop inhibitors that are effective against such a broad range of endemic, pandemic, and emerging CoVs, a conserved step of viral infection needs to be targeted.

Most human viruses, including CoVs, initiate primary attachment to cell surfaces through low-affinity but high-avidity interactions with glycans. Heparan sulfate proteoglycans are necessary attachment receptors for the human alphacoronavirus NL63 (HCoV-NL63) [5], as well as many other human viruses, such as herpes simplex virus (HSV) [6,7], hepatitis C virus (HCV) [8,9] and human immunodeficiency virus (HIV) [10]. Recently, heparan sulfate has also been identified as a necessary co-factor for SARS-CoV-2 infection [11,12]. Alternatively, other viruses, including influenza A virus (IAV) and betacoronaviruses HCoV-OC43 and HCoV-HKU1 require sialic acid-containing glycans for attachment and entry [13–15]. Regardless of the glycan subtype, glycan binding is critical to the primary attachment of many different viruses, facilitating their specific high-affinity interaction with less abundant protein receptors, such as ACE2 for HCoV-

NL63, SARS-CoV-1 and SARS-CoV-2 [16,17]. For some viruses, like IAV, glycans serve as the

only known entry receptor.

Blocking viral attachment to cellular glycans is a demonstrated approach to inhibit diverse

viral infections. Epigallocatechin gallate (EGCG) is the most bioactive polyphenolic compound

from green tea with antiviral activity against diverse DNA and RNA viruses, including HSV-1,

HCV, and IAV [18]. The broad antiviral spectrum of EGCG has been attributed to its ability to

inhibit the primary attachment of unrelated viruses by competing with both heparan sulfate and

sialic acid for virion binding [19]. However, whether EGCG inhibits coronavirus infection by

similar mechanisms remains unknown.

Here, we show that EGCG inhibits infectivity of murine, bat, and human CoVs. Cell

surface binding of multiple human CoVs is blocked by EGCG, suggesting that this natural product

inhibits a conserved step in CoV attachment, such as initial binding to glycans. These findings

demonstrate that blocking primary attachment is a potential antiviral strategy to prevent infection

by diverse CoVs.

Material and Methods

Compounds

Epigallocatechin-3-gallate (EGCG) was purchased from Thermo Fisher Scientific (CAS 989-51-

5). Heparin was obtained from Carbosynth Ltd (YH09354).

Plasmids

Lentiviral pseudoparticles were produced using plasmids encoding HIV-1 gag/pol (BEI Resources

NR-52517), tat (BEI Resources NR-52518), rev (BEI Resources NR-52519), a luciferase-

encoding lentiviral genome (BEI Resources NR-52516) and spike plasmids. The SARS-CoV-2

spike expression plasmid was obtained from Dr. Raffaele De Francesco (Addgene plasmid #

155297). The MERS-CoV spike expression plasmid was kindly provided by Dr. Stefan Pöhlmann

(Göttingen, Germany). The SARS-CoV-1 (Tor2 strain, GenBank accession no. **NC_004718.3**) and

WIV1-CoV spike (GenBank accession no. **KC881007.1**) expression plasmids were synthesized

by Genscript and codon-optimized for human expression. The lentiviral vector expressing human

ACE2 was obtained from Dr. Sonja Best (Addgene plasmid # 154981).

Cells and viruses

HEK293T/17 (ATCC ACS-4500), L929 (ATCC CCL-1), Huh7 (JCRB0403), HCT-8 (ATCC

CCL-244), 17Cl-1 (BEI Resources NR-53719) and Vero E6 (ATCC CRL-1586) cells were

cultured in Dulbecco's minimal essential medium (DMEM) with 10% FBS, 50 U/mL penicillin,

and 50 μg/mL streptomycin at 37°C in 5% CO_2. A549 (BEI Resources NR-52268) cells were

cultured in Hams F-12K (Kaighn's) medium with 10% FBS, 50 U/mL penicillin, and 50 μg/mL

streptomycin, and 10 μg/mL blasticidin (for A549-ACE2 cells) at 37°C in 5% CO_2. Calu-3 (ATCC

HTB-55) cells were cultured in minimal essential medium (MEM) with 10% FBS, 1 mM sodium

pyruvate, 1X non-essential amino acid solution, 50 U/mL penicillin, and 50 μg/mL streptomycin

at 37°C in 5% CO_2.

Human coronavirus 229E (HCoV-229E) and HCoV-OC43 were obtained from BEI

Resources (NR-52726 and NR-52725). For HCoV-229E, Huh7 cells were infected with a

multiplicity of infection (MOI) of 0.01 plaque-forming units (pfu)/cell and incubated at 33°C in

5% CO_2 until full cytopathic effects (CPE) were observed (~5 days post-infection). The

supernatants were collected, and cellular debris was pelleted by centrifugation at 1,000 × g for 10

min. The resulting supernatant was then aliquoted, and the viral stocks were stored at −80°C. For

HCoV-OC43, HCT-8 cells were infected (MOI of 0.01) and incubated at 33°C in 5% CO_2 until

full CPE were observed (~5 days post-infection). Similarly, supernatants were collected, cellular

debris was pelleted by centrifugation at 1,000 × g for 10 min and the resulting supernatant was

then aliquoted and stored at −80°C.

Green fluorescent protein-expressing murine hepatitis virus (MHV A59-GFP) was kindly

provided by Dr. Volker Thiel (Institute of Virology and Immunology, Bern, Switzerland) [20].

VSV-SARS-CoV-2 S was a gift from Dr. Sean Whelan (Washington University School of

Medicine in St. Louis, USA) [21]. MHV A59-GFP and VSV-SARS-CoV-2 were propagated as

described [20,21].

Infectivity assays

Huh7 cells ($1.6 × 10^6$ cells/well in 6-well plates) were infected with approximately 100 PFU of

HCoV-229E pre-exposed for 10 min at 37°C to EGCG or dimethyl sulfoxide (DMSO) vehicle in

DMEM. Inocula were removed 2 h later, and monolayers were overlaid with 1.2%

carboxymethylcellulose in DMEM containing 2% FBS. Infected cells were incubated at 33°C in

5% CO_2 until 4 days post-infection, when they were fixed with 10% formalin and stained with 1%

(wt/vol) crystal violet–10% (vol/vol) ethanol in H_2O for counting plaques.

Huh7 and A549 cells ($4 × 10^5$ cells/well in 24-well plates) were infected with

approximately 200 focus-forming units (FFU) of HCoV-OC43, pre-exposed to EGCG or DMSO

vehicle for 10 min at 37°C in DMEM. Inocula were removed 2 h later, and monolayers were

overlaid with 1.2% microcrystalline colloidal cellulose (Sigma-Aldrich 435244) in DMEM

containing 2% FBS. Infected cells were incubated at 33°C in 5% CO_2 for 3 days, then fixed and

processed for immunofluorescence to detect OC43 nucleoprotein. Cells were incubated with

primary mouse IgG anti-coronavirus group antibody MAB9013 (Millipore Sigma; diluted 1:500)

for 1 h at room temperature, followed by addition of secondary Alexa Fluor 488 anti-mouse IgG

Fab 2 antibody (Cell Signaling Technology 4408S) for 1 h at room temperature. Foci were

visualized and counted under a fluorescence microscope (Nikon Eclipse Ts2).

MHV A59-GFP virions pre-exposed to EGCG or the DMSO vehicle control for 10 min at

37°C in DMEM were used to infect L929 cell monolayers. Inocula were removed 1 h later, and

monolayers were overlaid with 1.2% carboxymethylcellulose in DMEM containing 2% FBS.

Infected cells were incubated at 33°C in 5% CO_2 for 24 h, then fixed prior to counting of GFP-

positive cells under the fluorescence microscope.

VSV-SARS-CoV-2 S virions pre-exposed to EGCG or the DMSO vehicle control for 10

min at 37°C in DMEM were used to infect Huh7, A549-ACE2 or Calu-3 cell monolayers in

duplicate wells of a 96-well plate. Infected cells were incubated at 33°C in 5% CO_2 for 24 h, then

fixed with 10% formalin. GFP signals were captured by Nikon Eclipse Ts2, processed using the

Python Imaging Library to find the mean grey value, and plotted as percentage of inhibition.

In all experiments, half maximal inhibitory concentrations (IC_{50}) were calculated by

nonlinear regression analysis using GraphPad Prism (version 9.0; GraphPad Software, Inc.).

Pseudoparticle entry assays

Lentiviral pseudoparticles were generated in HEK293T/17 cells by co-transfection using

Lipofectamine 2000 (Invitrogen 11668-019) [22]. The CoV spike expression plasmids were co-

transfected with a luciferase-encoding lentiviral genome, and plasmids encoding HIV-1 gag/pol,

tat, and rev [22]. Cell culture supernatants containing coronavirus spike pseudotyped lentiviral

particles were collected at 48 and 72 hours post-transfection, pooled and filtered through a 0.45

μm filter, and stored at -80°C.

Pseudoparticles were incubated for 10 min at 37°C with serially diluted EGCG and used

to infect Huh7 or A549-ACE2 cells in triplicate in 96-well plates. Infectivity was determined after

3 days by luminescence following addition of BrightGlo reagent (Promega PR-E2620). IC_{50} were

calculated by nonlinear regression analysis using GraphPad Prism (version 9.0; GraphPad

Software, Inc.).

Cell toxicity assay

Cells (50,000 per well in 96-well plate) were treated with EGCG or DMSO in DMEM and

incubated at 37°C in 5% CO_2 for 2h. To mimic the conditions used in the infectivity and

pseudoparticle assays, media was removed, cells were washed with PBS, and supplemented with

normal culture media of DMEM with 10% FBS, 50 U/mL penicillin, and 50 μg/mL streptomycin.

After 72 h, cell viability was assessed by alamarBlue cell viability reagent (Invitrogen DAL1025)

on a fluorescence plate reader (SpectraMax ID3 Multimode Plate reader).

Time-of-addition experiments

Near-confluent Huh7 and A549 cell monolayers in 12 or 24-well plates were pre-treated with

EGCG or DMSO vehicle for 1 h at 37°C in DMEM. Cells were washed with DMEM and infected

with 50-200 PFU of HCoV-229E or HCoV-OC43 for 2 h at 33°C. Alternatively, Huh7 and A549

cells were first infected with HCoV-229E or HCoV-OC43 for 2 h at 33°C. After removing the

inocula, the infected cells were overlaid with 1.2% microcrystalline colloidal cellulose in DMEM-

2% FBS containing EGCG or DMSO. Infectivity was assessed by plaquing efficiency or

immunostaining.

Binding assays

OC43 (4 x 10^4 PFU; MOI 0.05) or VSV-SARS-CoV-2 (2 x 10^5 PFU; MOI 0.5, chosen to match

the conditions of the infectivity assay) were exposed for 10 min at 37°C to EGCG or DMSO in

DMEM and adsorbed onto pre-chilled Huh7 and A549 cells for 1 h at 4°C. After three washes

with cold phosphate-buffered saline (PBS), cells were lysed and RNA was isolated using the

Monarch Total RNA Miniprep Kit (NEB T2010S), according to the manufacturer's instructions.

The RNA was transcribed to cDNA using the High Capacity cDNA Reverse Transcription Kit

(Applied Biosystems LS4368814). Quantitative real-time PCR was performed on the QuantStudio

3 (Applied Biosystems) instrument, using PowerTrack SYBR Green Master Mix (Applied

Biosystems LSA46110), with primers to detect actin (F: 5'-CTGGGAGTGGGTGGAGGC-3', R:

5'-TCAACTGGTCTCAAGTCAGTG-3')   and   OC43   N   gene   (F: 5'-

CCCAAGCAAACTGCTACCTCTCAG-3', R: 5'-GTAGACTCCGTCAATATCGGTGCC-3')

or   VSV   N   gene   (F: 5'-GATAGTACCGGAGGATTGACGACTA-3',   R: 5'-

TCAAACCATCCGAGCCATTC-3'). Following normalization to actin, the percentage of binding

was expressed relative to binding of virions treated with the DMSO control.

Results

EGCG inhibits murine and human coronavirus infection.

We first tested the effect of EGCG on the infectivity of human common cold coronaviruses HCoV-

229E (an alphacoronavirus) and HCoV-OC43 (a betacoronavirus). HCoV-229E and HCoV-OC43

virions were treated with DMSO or varying concentrations of EGCG for 10 min at 37°C and used

to inoculate infection-susceptible human hepatoma Huh7 or human lung epithelial A549 cell

monolayers. After 2 h of infection, inocula were removed and cells were overlaid with plaquing

media. Infectivity was assessed by plaquing efficiency on Huh7 for HCoV-229E or by

immunofluorescence focus-forming assay on Huh7 and A549 cells for HCoV-OC43. EGCG

inhibited infectivity of HCoV-229E (IC_{50} = 0.77 μM) and HCoV-OC43 (IC_{50} = 0.49-0.56 μM) at

sub- to low micromolar concentrations (**Figure 1A**), similar to other viruses highly susceptible to

inhibition by EGCG, such as HSV-1 [19]. We next tested the inhibitory activity of EGCG against

a murine CoV, murine hepatitis virus (MHV-A59), on murine fibroblast L929 cell monolayers.

We observed that EGCG inhibited MHV-A59 infection with similar potency as for the human

common cold CoVs (IC_{50} = 0.71 μM) (**Figure 1B**) and inhibited the formation of syncytia (**Figure**

1C). Consistent with previous reports, we found EGCG to have minimal effect on cell viability at

the low micromolar range in which antiviral activity is exerted (**Figure 1D**).

Entry of highly pathogenic coronaviruses is blocked by EGCG.

We produced luciferase-encoding lentiviral particles pseudotyped with SARS-CoV-1, SARS-

CoV-2 and MERS-CoV spike (S) proteins to investigate the effect of EGCG on entry of highly

pathogenic coronaviruses. We also generated lentiviral particles pseudotyped with the spike

protein of WIV1-CoV, a bat coronavirus that binds to human ACE2 [23]. Huh7 cells were

inoculated with CoV S pseudoparticles pre-exposed to EGCG or DMSO for 10 min at 37°C.

Inocula were removed after 2 h and replaced with fresh DMEM. After 72 h, cells were lysed and

luminescence was measured. EGCG inhibited SARS-CoV-1, SARS-CoV-2, MERS-CoV and

WIV1-CoV pseudoparticle entry with IC_{50} of 2.5, 48.6, 13.7 and 3.8 μM, respectively (**Figure**

2A). To evaluate the antiviral effect in a more relevant cell type, we generated A549 cells stably

expressing human ACE2, the receptor for SARS-CoV-1 and SARS-CoV-2, and isolated a clone

with high levels of ACE2 expression and enhanced susceptibility to SARS-CoV-2 S

pseudoparticle infection (**Supplemental Figure 1**). We found that EGCG also inhibited entry of

SARS-CoV-1, SARS-CoV-2, and WIV1-CoV pseudoparticles into A549-ACE2 B9 cells, with

IC_{50} of 15.9, 24.0 and 24.5 µM, respectively (**Figure 2A**).

Figure 1. EGCG pre-treatment inhibits diverse CoV infections. Pre-treatment of authentic
HCoV-229E and HCoV-OC43 **(A)** or murine coronavirus MHV-A59 **(B)** with EGCG for 10 min
at 37°C prevented infection of Huh7, A549 or L292 cells, respectively. Mean values with standard
deviation of three independent experiments are plotted. **(C)** A reduction in syncytia (black arrows)
was observed for EGCG-treated MHV-A59. Scale bar: 200 µm. **(D)** Cell exposure to EGCG during
the 2 h infection had minimal effect on cell viability. Mean values with standard deviation of two
independent experiments with duplicates are plotted.

We next assessed the effect of EGCG on replication-competent vesicular stomatitis virus

(VSV) virions expressing GFP and SARS-CoV-2 S instead of the VSV glycoprotein (VSV-SARS-

CoV-2) [21]. Huh7, A549-ACE2, and lung epithelial Calu-3 cells were infected with EGCG-

treated VSV-SARS-CoV-2 virions (MOI of 5). Infected cells were imaged by fluorescence

microscopy after 24 h. Similar to the pseudoparticle assays, EGCG inhibited VSV-SARS-CoV-2

S infection with IC_{50} ranging from 11.45 to 15.40 μM, depending on the cell type (**Figure 2B**),

with minimal cytotoxicity up to 100 μM (**Supplemental Figure 2**). We next evaluated whether

the effect of EGCG was dependent on MOI. Using the same assay, but with MOI of 0.5, we found

that EGCG inhibited VSV-SARS-CoV-2 infection with IC_{50} ranging from 0.14 to 1.13 μM. These

findings demonstrate that the inhibitory activity of EGCG depends on the MOI (**Figure 2B**),

consistent with the virions as the proposed target of EGCG [19]. This also provides a potential

explanation for the difference in IC_{50} for authentic viruses investigated at low MOI (<0.001)

(**Figure 1A-B**) compared to lentiviral pseudoparticles (**Figure 2A**), for which p24 ELISA data for

lentiviral particles pseudotyped with SARS-CoV-2 spike in our laboratory provide an estimate of

approximately 1.7×10^6 transducing units per mL, resulting in an apparent MOI of 3.4.

EGCG acts on virions to inhibit coronavirus binding to cell surfaces.

To confirm that EGCG acts on CoV virions, and not on cells, we performed time-of-addition

experiments. Cells were pre-treated with EGCG or DMSO 1 h at 37°C, at which point media was

removed and cells were infected with HCoV-229E or HCoV-OC43. Infectivity was determined by

plaque assay or immunostaining. Alternatively, cells were infected, and EGCG- or DMSO-

containing medium was added 2 h post-infection. In either condition, no inhibition of infectivity

was observed until the highest concentration of 100 μM EGCG (**Figure 3A**). To test if EGCG

inhibits CoV attachment, we performed binding assays in which cells were incubated with pre-

treated virions at 4°C for 1 h to allow for attachment, but not fusion or entry. After cell washing,

bound virions were quantified by RT-qPCR (**Supplemental Figure 3**). Pre-chilled Huh7 or Calu-

3 cells were inoculated with HCoV-OC43 or VSV-SARS-CoV-2 treated with DMSO, 10 μM

EGCG, or 20 μg/mL heparin, as a control attachment inhibitor [11]. As expected, heparin partially

blocked VSV-SARS-CoV-2 binding. We also observed that heparin reduced HCoV-OC43

binding, which unlike other coronavirus does not have a known protein receptor but depends on

sialic acid, and, as recently reported, glycosaminoglycans for entry [24]. EGCG similarly reduced

VSV-SARS-CoV-2 binding to Huh7 and Calu-3 cells (**Figure 3B-C**).

Figure 2. EGCG inhibits entry of highly pathogenic CoVs. (A) EGCG treatment reduced entry
of lentiviral particles pseudotyped with S from SARS-CoV-1, MERS-CoV, SARS-CoV-2, or
WIV1-CoV. Mean values with standard error of the mean of three independent experiments each
done in triplicates are plotted. **(B)** EGCG displayed MOI-dependent inhibition of VSV-SARS-
CoV-2 in Huh7, A549-ACE2 B9 and Calu-3. Mean values with standard deviation of three
independent experiments each in duplicates are plotted. Representative images are shown. Scale
bar: 200 μm.

Figure 3. EGCG inhibits CoV attachment to cells. (A) Cell pre-treatment or post-229E and OC43 infection treatment does not restrict infection to similar extents as virion pre-treatment. Mean values with standard deviation of three independent experiments are plotted. (B) Virions were pre-treated with EGCG or heparin for 10 min at 37°C then cooled on ice and added to pre-chilled Huh7 (B) or Calu-3 cells (C) for 1 hour on ice. Attached virus was quantified by RT-qPCR after washing cells with PBS three times. Mean values with standard deviation from qPCR triplicates of two independent experiments are plotted.

Discussion

Green tea catechins, especially EGCG, exert antiviral effects against diverse viruses. We show that

EGCG broadly inhibits CoV infections, and that its activity is due, at least in part, to blocking CoV

attachment to target cells. Infection by authentic murine and human CoVs, as well as entry of

lentivirus particles pseudotyped with spike proteins of highly pathogenic or potentially emerging

CoVs [23], was inhibited by EGCG treatment. Interestingly, EGCG was recently shown to

significantly inhibit authentic SARS-CoV-2 infection, and time-of-addition experiments suggested

that EGCG at least partially blocks SARS-CoV-2 entry [25], consistent with our findings. In this

study, we expand the scope of EGCG activity to other CoVs, including murine and bat CoVs, and demonstrate the physiological relevance of the inhibitory activity of EGCG on CoV attachment and infection of lung epithelial cells.

A variety of mechanisms for the antiviral activity of EGCG have previously been described. EGCG inhibits the primary attachment of diverse viruses including VSV, IAV, HCV, HSV-1 and vaccinia virus [19]. EGCG is proposed to act directly on HCV virions to block attachment in a genotype independent manner, with no alteration in HCV (co)-receptor expression nor any effect on viral replication, assembly, or release [26,27]. As for HBV, EGCG may inhibit entry [28], as well as replication, with proposed roles in impairing HBV replicative intermediates during DNA synthesis [29], diminishing the transcriptional activation of the HBV core promoter [30], and enhancing lysosomal acidification to adversely affect HBV replication [31]. Alternatively, EGCG is suggested to prevent the binding of HIV glycoprotein 120 to its receptor molecule CD4 on cells [32], and to act as an HIV reverse transcriptase inhibitor [33]. Finally, recent studies have identified *in vitro* inhibitory effects of EGCG against the SARS-CoV-2 3C-like protease [34–37]. However, our time-of-addition experiments with HCoV-229E and HCoV-OC43, as well as those of others with SARS-CoV-2 [25], strongly suggest that EGCG predominantly inhibits authentic CoV infection by blocking entry, although other activities might be exerted at higher concentrations. Furthermore, we show that EGCG reduces cell surface binding of HCoV-OC43 and VSV-SARS-CoV-2, in a similar manner to heparin, which has been demonstrated to competitively block virion attachment [11].

While COVID-19 vaccine development and deployment are ongoing, extensive efforts are underway to identify antibodies, peptides, and small molecules to prevent SARS-CoV-2 entry [38]. To protect against emerging CoVs, the identification of broad-spectrum entry inhibitors is a

priority. Recently, lactoferrins have been described to disrupt CoV primary attachment, mediated

by heparan sulfate interactions, conferring antiviral activity against multiple CoVs *in vitro*,

demonstrating the potential for a pan-coronavirus inhibitor [39]. Small molecules continue to be

preferred as drugs in general due to superior pharmacokinetic properties and simpler synthesis

[40]. As with other green tea catechins, EGCG does not accumulate at high levels, is unstable

under physiological conditions, and is rapidly metabolized [41–43]. However, the observation that

a small molecule exerts pan-coronavirus activity provides the basis for the development of entry

inhibitors that could provide an effective antiviral strategy to protect against future emerging CoV

infections.

Funding information

This work was supported by a Natural Sciences and Engineering Research Council of Canada

Discovery Grant, a Banting Research Foundation Discovery Award, as well as by the Canadian

Foundation for Innovation John R. Evans Leaders Fund, and Queen's University. E.V.L. is

supported by a Vanier Canada Graduate Scholarship.

Acknowledgements

The following reagents were obtained through BEI Resources (NIAID, NIH): *Homo sapiens* A549

Lung Carcinoma Cells (NR-52268); Murine 17CL1 Cell Line (NR-53719); SARS-Related

Coronavirus 2, Wuhan-Hu-1Spike-Pseudotyped Lentiviral Kit (NR-52948); Human Coronavirus

OC43 (NR-52725) and 229E (NR-52726). We thank Gerard O'Leary for Python processing scripts,

Dr. Selena Sagan (McGill University) for providing coronavirus infection protocols, and Dr.

Chantelle Capicciotti (Queen's University) for helpful discussions.

References

1. Li, W.; Shi, Z.; Yu, M.; Ren, W.; Smith, C.; Epstein, J.H.; Wang, H.; Crameri, G.; Hu, Z.;

Zhang, H.; et al. Bats Are Natural Reservoirs of SARS-Like Coronaviruses. *Science.*

2005, *310*, 676–679, doi:10.1126/science.1118391.

2. Walsh, E.E.; Shin, J.H.; Falsey, A.R. Clinical impact of human coronaviruses 229E and

OC43 infection in diverse adult populations. *J. Infect. Dis.* **2013**, *208*, 1634–1642,

doi:10.1093/infdis/jit393.

3. Patrick, D.M.; Petric, M.; Skowronski, D.M.; Guasparini, R.; Booth, T.F.; Krajden, M.;

McGeer, P.; Bastien, N.; Gustafson, L.; Dubord, J.; et al. An Outbreak of Human

Coronavirus OC43 Infection and Serological Cross-reactivity with SARS Coronavirus.

Can. J. Infect. Dis. Med. Microbiol. **2006**, *17*, 330–336, doi:10.1155/2006/152612.

4. Hand, J.; Rose, E.B.; Salinas, A.; Lu, X.; Sakthivel, S.K.; Schneider, E.; Watson, J.T.

Severe respiratory illness outbreak associated with human coronavirus NL63 in a long-

term care facility. *Emerg. Infect. Dis.* **2018**, *24*, 1964–1966, doi: 10.3201/eid2410.180862.

5. Milewska, A.; Zarebski, M.; Nowak, P.; Stozek, K.; Potempa, J.; Pyrc, K. Human

coronavirus NL63 utilizes heparan sulfate proteoglycans for attachment to target cells. *J.*

Virol. **2014**, *88*, 13221–13230, doi:10.1128/JVI.02078-14.

6. WuDunn, D.; Spear, P.G. Initial interaction of herpes simplex virus with cells is binding

to heparan sulfate. *J. Virol.* **1989**, *63*, 52–58, doi:10.1128/JVI.63.1.52-58.1989.

7. Lycke, E.; Johansson, M.; Svennerholm, B.; Lindahl, U. Binding of herpes simplex virus

to cellular heparan sulphate, an initial step in the adsorption process. *J. Gen. Virol.* **1991**,

72, 1131–1137, doi:10.1099/0022-1317-72-5-1131.

8. Germi, R.; Crance, J.-M.; Garin, D.; Guimet, J.; Lortat-Jacob, H.; Ruigrok, R.W.H.;

Zarski, J.-P.; Drouet, E. Cellular glycosaminoglycans and low density lipoprotein receptor

are involved in hepatitis C virus adsorption. *J. Med. Virol.* **2002**, *68*, 206–215,

doi:10.1002/jmv.10196.

9.   Morikawa, K.; Zhao, Z.; Date, T.; Miyamoto, M.; Murayama, A.; Akazawa, D.; Tanabe,

J.; Sone, S.; Wakita, T. The roles of CD81 and glycosaminoglycans in the adsorption and

uptake of infectious HCV particles. *J. Med. Virol.* **2007**, *79*, 714–723,

doi:10.1002/jmv.20842.

10.   Patel, M.; Yanagishita, M.; Roderiquez, G.; Bou-Habib, D.C.; Oravecz, T.; Hascall, V.C.;

Norcorss, M.A. Cell-Surface Heparan Sulfate Proteoglycan Mediates HIV-1 Infection of

T-Cell Lines. *AIDS Res. Hum. Retroviruses* **1993**, *9*, 167–174,

doi:10.1089/aid.1993.9.167.

11.   Clausen, T.M.; Sandoval, D.R.; Spliid, C.B.; Pihl, J.; Perrett, H.R.; Painter, C.D.;

Narayanan, A.; Majowicz, S.A.; Kwong, E.M.; McVicar, R.N.; et al. SARS-CoV-2

Infection Depends on Cellular Heparan Sulfate and ACE2. *Cell* **2020**, *183*, 1043-

1057.e15, doi:10.1016/j.cell.2020.09.033.

12.   Zhang, Q.; Chen, C.Z.; Swaroop, M.; Xu, M.; Wang, L.; Lee, J.; Wang, A.Q.; Pradhan,

M.; Hagen, N.; Chen, L.; et al. Heparan sulfate assists SARS-CoV-2 in cell entry and can

be targeted by approved drugs in vitro. *Cell Discov.* **2020**, *6*, 80, doi:10.1038/s41421-020-

00222-5.

13.   Krempl, C.; Schultze, B.; Herrler, G. Analysis of cellular receptors for human coronavirus

OC43. *Adv Exp Med Bio.* **1995**, 380, 371–4, doi:10.1007/978-1-4615-1899-0_60.

14.   Huang, X.; Dong, W.; Milewska, A.; Golda, A.; Qi, Y.; Zhu, Q.K.; Marasco, W.A.; Baric,

R.S.; Sims, A.C.; Pyrc, K.; et al. Human Coronavirus HKU1 Spike Protein Uses O -

Acetylated Sialic Acid as an Attachment Receptor Determinant and Employs

Hemagglutinin-Esterase Protein as a Receptor-Destroying Enzyme. *J. Virol.* **2015**, *89*,

7202–13, doi:10.1128/jvi.00854-15.

15.   Weis, W.; Brown, J.H.; Cusack, S.; Paulson, J.C.; Skehel, J.J.; Wiley, D.C. Structure of

the influenza virus haemagglutinin complexed with its receptor, sialic acid. *Nature* **1988**,

333, 426–431, doi:10.1038/333426a0.

16.   Hofmann, H.; Pyrc, K.; van der Hoek, L.; Geier, M.; Berkhout, B.; Pöhlmann, S. Human

coronavirus NL63 employs the severe acute respiratory syndrome coronavirus receptor for

cellular entry. *Proc. Natl. Acad. Sci.* **2005**, *102*, 7988–7993,

doi:10.1073/pnas.0409465102.

17.   Hoffmann, M.; Kleine-Weber, H.; Schroeder, S.; Krüger, N.; Herrler, T.; Erichsen, S.;

Schiergens, T.S.; Herrler, G.; Wu, N.-H.; Nitsche, A.; et al. SARS-CoV-2 Cell Entry

Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease

Inhibitor. *Cell* **2020**, *181*, 271-280.e8, doi:10.1016/j.cell.2020.02.052.

18.   Xu, J.; Xu, Z.; Zheng, W. A Review of the Antiviral Role of Green Tea Catechins.

Molecules **2017**, *22*, 1337, doi:10.3390/molecules22081337.

19.   Colpitts, C.C.; Schang, L.M. A Small Molecule Inhibits Virion Attachment to Heparan

Sulfate- or Sialic Acid-Containing Glycans. *J. Virol.* **2014**, *88*, 7806–7817,

doi:10.1128/JVI.00896-14.

20.   Coley, S.E.; Lavi, E.; Sawicki, S.G.; Fu, L.; Schelle, B.; Karl, N.; Siddell, S.G.; Thiel, V.

Recombinant mouse hepatitis virus strain A59 from cloned, full-length cDNA replicates to

high titers in vitro and is fully pathogenic in vivo. *J. Virol.* **2005**, *79*, 3097–3106,

doi:10.1128/JVI.79.5.3097-3106.2005.

21. Case, J.B.; Rothlauf, P.W.; Chen, R.E.; Liu, Z.; Zhao, H.; Kim, A.S.; Bloyet, L.-M.; Zeng,

Q.; Tahan, S.; Droit, L.; et al. Neutralizing Antibody and Soluble ACE2 Inhibition of a

Replication-Competent VSV-SARS-CoV-2 and a Clinical Isolate of SARS-CoV-2. *Cell*

Host Microbe **2020**, *28*, 475-485.e5, doi:10.1016/j.chom.2020.06.021.

22. Crawford, K.H.D.; Eguia, R.; Dingens, A.S.; Loes, A.N.; Malone, K.D.; Wolf, C.R.; Chu,

H.Y.; Tortorici, M.A.; Veesler, D.; Murphy, M.; et al. Protocol and Reagents for

Pseudotyping Lentiviral Particles with SARS-CoV-2 Spike Protein for Neutralization

Assays. *Viruses.* **2020**, *12*, 513, doi:10.1007/978-1-4615-1899-0_60.

23. Ge, X.-Y.; Li, J.-L.; Yang, X.-L.; Chmura, A.A.; Zhu, G.; Epstein, J.H.; Mazet, J.K.; Hu,

B.; Zhang, W.; Peng, C.; et al. Isolation and characterization of a bat SARS-like

coronavirus that uses the ACE2 receptor. *Nature* **2013**, *503*, 535–538,

doi:10.1038/nature12711.

24. Wang, R.; Simoneau, C.R.; Kulsuptrakul, J.; Bouhaddou, M.; Travisano, K.A.; Hayashi,

J.M.; Carlson-Stevermer, J.; Zengel, J.R.; Richards, C.M.; Fozouni, P.; et al. Genetic

Screens Identify Host Factors for SARS-CoV-2 and Common Cold Coronaviruses. *Cell*

2021, *184*, 106-119.e14, doi:10.1016/j.cell.2020.12.004.

25. Henss, L.; Auste, A.; Schürmann, C.; Schmidt, C.; von Rhein, C.; Mühlebach, M.D.;

Schnierle, B.S. The green tea catechin epigallocatechin gallate inhibits SARS-CoV-2

infection. *J. Gen. Virol.* **2021**, *102*, doi:10.1099/jgv.0.001574.

26. Calland, N.; Albecka, A.; Belouzard, S.; Wychowski, C.; Duverlie, G.; Descamps, V.;

Hober, D.; Dubuisson, J.; Rouillé, Y.; Séron, K. (−)-Epigallocatechin-3-gallate is a new

inhibitor of hepatitis C virus entry. *Hepatology* **2012**, *55*, 720–729,

doi:10.1002/hep.24803.

27. Ciesek, S.; von Hahn, T.; Colpitts, C.C.; Schang, L.M.; Friesland, M.; Steinmann, J.;

Manns, M.P.; Ott, M.; Wedemeyer, H.; Meuleman, P.; et al. The green tea polyphenol,

epigallocatechin-3-gallate, inhibits hepatitis C virus entry. *Hepatology* **2011**, *54*, 1947–

1955, doi:10.1002/hep.24610.

28. Huang, H.-C.; Tao, M.-H.; Hung, T.-M.; Chen, J.-C.; Lin, Z.-J.; Huang, C. (−)-

Epigallocatechin-3-gallate inhibits entry of hepatitis B virus into hepatocytes. *Antiviral*

Res. **2014**, *111*, 100–111, doi:10.1016/j.antiviral.2014.09.009.

29. He, W.; Li, L.X.; Liao, Q.J.; Liu, C.L.; Chen, X.L. Epigallocatechin gallate inhibits HBV

DNA synthesis in a viral replication - inducible cell line. *World J. Gastroenterol.* **2011**,

17, 1507–1514, doi:10.3748/wjg.v17.i11.1507.

30. Xu, J.; Gu, W.; Li, C.; Li, X.; Xing, G.; Li, Y.; Song, Y.; Zheng, W. Epigallocatechin

gallate inhibits hepatitis B virus via farnesoid X receptor alpha. *J. Nat. Med.* **2016**, *70*,

584–591, doi:10.1007/s11418-016-0980-6.

31. Zhong, L.; Hu, J.; Shu, W.; Gao, B.; Xiong, S. Epigallocatechin-3-gallate opposes HBV-

induced incomplete autophagy by enhancing lysosomal acidification, which is unfavorable

for HBV replication. *Cell Death Dis.* **2015**, *6*, e1770–e1770, doi:10.1038/cddis.2015.136.

32. Williamson, M.P.; McCormick, T.G.; Nance, C.L.; Shearer, W.T. Epigallocatechin

gallate, the main polyphenol in green tea, binds to the T-cell receptor, CD4: Potential for

HIV-1 therapy. *J. Allergy Clin. Immunol.* **2006**, *118*, 1369–1374,

doi:10.1016/j.jaci.2006.08.016.

33. Li, S.; Hattori, T.; Kodama, E.N. Epigallocatechin gallate inhibits the HIV reverse

transcription step. *Antivir. Chem. Chemother.* **2011**, *21*, 239–243, doi:10.3851/IMP1774.

34. Jang, M.; Park, R.; Park, Y.-I.; Cha, Y.-E.; Yamamoto, A.; Lee, J.I.; Park, J. EGCG, a

444 green tea polyphenol, inhibits human coronavirus replication in vitro. *Biochem. Biophys.*

445 *Res. Commun.* **2021**, *547*, 23–28, doi:10.1016/j.bbrc.2021.02.016.

446 35. Jang, M.; Park, Y.-I.; Cha, Y.-E.; Park, R.; Namkoong, S.; Lee, J.I.; Park, J. Tea

447 Polyphenols EGCG and Theaflavin Inhibit the Activity of SARS-CoV-2 3CL-Protease *In*

448 *Vitro*. *Evidence-Based Complement. Altern. Med.* **2020**, *2020*, 5630838,

449 doi:10.1155/2020/5630838.

450 36. Chiou, W.-C.; Chen, J.-C.; Chen, Y.-T.; Yang, J.-M.; Hwang, L.-H.; Lyu, Y.-S.; Yang,

451 H.-Y.; Huang, C. The inhibitory effects of PGG and EGCG against the SARS-CoV-2 3C-

452 like protease. *Biochem. Biophys. Res. Commun.* **2021**, *S0006-291X*, 32299–3,

453 doi:10.1016/j.bbrc.2020.12.106.

454 37. Du, A.; Zheng, R.; Disoma, C.; Li, S.; Chen, Z.; Li, S.; Liu, P.; Zhou, Y.; Shen, Y.; Liu,

455 S.; et al. Epigallocatechin-3-gallate, an active ingredient of Traditional Chinese

456 Medicines, inhibits the 3CLpro activity of SARS-CoV-2. *Int. J. Biol. Macromol.* **2021**,

457 *176*, 1–12, doi:10.1016/j.ijbiomac.2021.02.012.

458 38. Xiu, S.; Dick, A.; Ju, H.; Mirzaie, S.; Abdi, F.; Cocklin, S.; Zhan, P.; Liu, X. Inhibitors of

459 SARS-CoV-2 Entry: Current and Future Opportunities. *J. Med. Chem.* **2020**, *63*, 12256–

460 12274, doi:10.1021/acs.jmedchem.0c00502.

461 39. Hu, Y.; Meng, X.; Zhang, F.; Xiang, Y.; Wang, J. The in vitro antiviral activity of

462 lactoferrin against common human coronaviruses and SARS-CoV-2 is mediated by

463 targeting the heparan sulfate co-receptor. *Emerg. Microbes Infect.* **2021**, *10*, 317–330,

464 doi:10.1080/22221751.2021.1888660.

465 40. Ngo, H.X.; Garneau-Tsodikova, S. What are the drugs of the future? *Medchemcomm*

466 **2018**, *9*, 757–758, doi:10.1039/c8md90019a.

467   41.   Lambert, J.D.; Lee, M.-J.; Lu, H.; Meng, X.; Hong, J.J.J.; Seril, D.N.; Sturgill, M.G.;

468   Yang, C.S. Epigallocatechin-3-Gallate Is Absorbed but Extensively Glucuronidated

469   Following Oral Administration to Mice. *J. Nutr.* **2003**, *133*, 4172–4177,

470   doi:10.1093/jn/133.12.4172.

471   42.   Chow, H.-H.S.; Cai, Y.; Hakim, I.A.; Crowell, J.A.; Shahi, F.; Brooks, C.A.; Dorr, R.T.;

472   Hara, Y.; Alberts, D.S. Pharmacokinetics and Safety of Green Tea Polyphenols after

473   Multiple-Dose Administration of Epigallocatechin Gallate and Polyphenon E in Healthy

474   Individuals. *Clin. Cancer Res.* **2003**, *9*, 3312–3319.

475   43.   Smith, T.J. Green tea polyphenols in drug discovery: a success or failure? *Expert Opin.*

476   *Drug Discov.* **2011**, *6*, 589–595, doi:10.1517/17460441.2011.570750.