**A conserved *trans* regulatory loop involving an odorant binding protein controls male mating behavior in flies**

Pablo J Delclos[†,1], Kiran Adhikari[†,1], Oluwatomi Hassan[1], Alexander A Oderhowho[1], Vyshnika Sriskantharajah[1,2], Tammie Trinh[1,3], Richard P Meisel[1]

[†]Co-first author/equal contribution

Affiliation: [1]Department of Biology & Biochemistry, University of Houston, Houston, TX, USA

[2]Current affiliation: University of Texas Health Science Center School of Biomedical Informatics, Houston, TX, USA

[3]Current affiliation: University of Texas School of Dentistry, Houston, TX, USA

## ABSTRACT

A major goal in evolutionary biology is to understand how natural variation is maintained in sexually selected and sexually dimorphic traits. Hypotheses to explain genetic variation in sexually selected traits include context-dependent fitness effects, epistatic interactions, and pleiotropic constraints. The house fly, *Musca domestica*, is a promising system to investigate how these factors affect polymorphism in sexually selected traits. Two common Y chromosomes ($Y^M$ and $III^M$) segregate as stable polymorphisms in natural house fly populations, appear to be locally adapted to different thermal habitats, and differentially affect male mating success. Here, we perform a meta-analysis of RNA-seq data which identifies genes encoding odorant binding proteins (in the *Obp56h* family) as differentially expressed between the heads of males carrying $Y^M$ and $III^M$. Differential expression of *Obp56h* has been associated with variation in male mating behavior in *Drosophila melanogaster*. We find differences in male mating behavior between house flies carrying the Y chromosomes that are consistent with the relationship between male mating behavior and expression of *Obp56h* in *D. melanogaster*. We also find that male mating behaviors in house fly are affected by temperature, and the same temperature differentials further affect the expression of *Obp56h* genes. However, we show that temperature-dependent effects cannot explain the maintenance of genetic variation for male mating behavior in house fly. Using a network analysis and allele-specific expression measurements, we find evidence that the house fly $III^M$ chromosome is a *trans* regulator of *Obp56h* gene expression. Moreover, we find that *Obp56h* disproportionately affects the expression of genes on the *D. melanogaster* chromosome that is homologous to the house fly $III^M$ chromosome. This provides evidence for a conserved *trans* regulatory loop involving *Obp56h* expression that affects male mating behavior in flies. The complex regulatory architecture controlling *Obp56h* expression suggests that variation in male mating behavior could be maintained by epistasis or pleiotropic constraints.

## INTRODUCTION

Sexual selection occurs when there is heritable variation in reproductive success that arises from competition for access to mates (Jones 2016). Sexual selection can shape phenotypic variation within and among species (Seehausen and van Alphen 1999; Greene et al. 2000; Brooks and Endler 2001; van Doorn and Weissing 2002; Rueffler et al. 2006). If a trait is under strong sexual selection, one may expect the fixation of alleles that increase reproductive success. In contrast to this expectation, substantial genetic variation can exist in strongly sexually selected traits (e.g., the lek paradox), and explaining this variation has been a long-standing goal in evolutionary biology (Kirkpatrick and Ryan 1991; Tomkins et al. 2004). Addressing this goal includes determining the genetic basis of sexually selected traits, as well as the environmental drivers that affect variation in these traits (Jia et al. 2000; Ingleby et al. 2010).

Three potential, and non-exclusive, causes of variation in sexually selected traits are epistasis, pleiotropic constraints, and context-dependent fitness effects. Epistatic and pleiotropic constraints are signatures of a complex gene regulatory architecture underlying a trait, and both are predicted to be important genetic mechanisms in maintaining genetic variance, particularly for fitness-related traits (Fenster et al. 1997; Merilä and Sheldon 1999; Arnqvist et al. 2014). For example, although Y chromosome genotype is a major determinant of male fitness in *Drosophila melanogaster*, those fitness effects depend on the genetic background (Chippindale and Rice 2001). Epistatic interactions between Y-linked and X-linked or autosomal alleles can therefore reduce heritable variation for male fitness. Pleiotropy creates genetic covariation amongst unrelated traits, which can similarly reduce the response to selection and thereby allow genetic variation in sexually selected traits to be maintained (Fitzpatrick 2004; Chenoweth and McGuigan 2010). For instance, *Drosophila* cuticular hydrocarbons (CHCs) are lipid compounds used for both chemical communication and resistance to various environmental stressors, including desiccation (Chung and Carroll 2015). Disrupted expression of individual genes responsible for CHC production in *D. melanogaster* can result in significant alterations to both mating behaviors and ecologically relevant CHCs (Marcillac et al. 2005; Shorter et al. 2016). Lastly, context-dependence, in the form of genotype-by-environment interactions, may also maintain variation when environments fluctuate substantially (Kokko and Heubel 2008). However, the combined effects of ecological and genetic variation on sexually selected and sexually dimorphic traits is especially unresolved, with conflicting evidence for the relative importance of sex-by-genotype-by-environment interactions (Delph et al. 2011; Connallon 2015; Allen et al. 2017; Lasne et al. 2018; Connallon et al. 2019; Ruzicka et al. 2020).

The house fly, *Musca domestica*, is a promising system to investigate the factors that maintain genetic variation for sexually selected traits. House fly has a polygenic sex determination system, in which multiple male and female determining loci segregate as polymorphisms in natural

populations (Hamm et al. 2015). This polymorphism has remained stable in natural house fly
populations since at least the mid-20th century (Kozielska et al. 2008; Meisel et al. 2016). Two
common proto-Y chromosomes ($III^M$ and $Y^M$) are distributed along latitudinal clines across
multiple continents, with $III^M$ most common in the south and $Y^M$ most common in the north
(Franco et al. 1982; Tomita and Wada 1989; Hamm et al. 2005). This clinal distribution suggests
that geographically heterogeneous selection pressures are responsible for maintaining polygenic
sex determination (Levene 1953; Hedrick et al. 1976). Consistent with this prediction,
seasonality in temperature is the best predictor of the frequencies of the proto-Y chromosomes
across populations (Feldmeyer et al. 2008). The $Y^M$ and $III^M$ chromosomes also affect male
mating success, with $III^M$ males exhibiting an advantage over $Y^M$ males (Hamm et al. 2009).
Together, the geographic distribution of the proto-Y chromosomes and their effects on male
mating behavior raise the possibility that polymorphic house fly proto-Y chromosomes affect
sexually selected traits in a context-dependent (i.e., environment-specific) manner.

Here, we utilized the house fly system to test the potential roles of context-dependence and
genetic architecture (epistasis and pleiotropy) in the maintenance of genetic variation of a
sexually selected trait. To those ends, we tested for genotype-by-temperature interactions
affecting differences in male mating behavior between $Y^M$ and $III^M$ males. We also used
RNA-seq data to investigate the genetic basis of differences in mating performance between $Y^M$
and $III^M$ males (Meisel et al. 2015; Son et al. 2019; Adhikari et al. 2021). Our results implicate
genes encoding odorant binding proteins (Obps) in an evolutionarily conserved *trans* regulatory
loop involving a chromosome that independently became sex-linked in both house fly and
*Drosophila*.

**METHODS**

***RNA-seq differential gene expression analysis***

We analyzed published RNA-seq data from *M. domestica* male heads (NCBI Gene Expression
Omnibus accessions GSE67065, GSE126685, and GSE126689, shown in Table S1). The
RNA-seq data include nine $Y^M$ and fifteen $III^M$ samples (Meisel et al. 2015; Son et al. 2019). We
aligned RNA-Seq reads to house fly transcripts from genome assembly v2.0.2 and annotation
release 102 (Scott et al. 2014) using kallisto in single-end read mode (Bray et al. 2016). All
RNA-seq reads were single-end, and we set the average fragment length to 300 bp and standard
deviation to 30 bp for all samples.

We tested for differentially expressed (DE) genes between males with a $Y^M$ chromosome and
males with a $III^M$ chromosome using a combination of DESeq2 (Love et al. 2014), sva (Leek et
al. 2012), and limma (Ritchie et al. 2015). Read counts per gene that passed an initial threshold
filter (0.5 counts per million in at least 4 samples) were normalized by variance stabilizing
transformation in DESeq2. To remove batch effects across data sets, we used the sva package to

identify and estimate surrogate variables that adjust for latent sources of variation (e.g., batch
effects). To identify genes DE between $Y^M$ and $III^M$ males, we used the lmFit() function in limma
to fit a linear model comprised of male type ($Y^M$ vs $III^M$) and our surrogate variables as fixed
effects, and read counts as the response variable. We then computed contrasts between male
types and calculated test statistics using the eBayes() function. Genes below a false discovery
rate (FDR) adjusted $p$ value of 0.05 were categorized as DE (Benjamini and Hochberg 1995).

### Weighted gene co-expression network analysis

We used weighted gene co-expression network analysis (WGCNA) to identify modules of house
fly genes whose expression correlates with male type ($Y^M$ or $III^M$) on normalized read count data
that were adjusted for batch effects in sva (Langfelder and Horvath 2008). For all pairs of genes
with variable expression across samples, we calculated Pearson's correlation coefficient across
all samples. We created an unsigned correlation matrix and adjusted the soft-threshold value ($\beta$)
to which among-gene covariances are exponentially raised to ensure a scale-free topology (this
resulted in $\beta = 7$), thereby creating a weighted network of gene expression. An unsigned matrix
allows us to identify connected genes whose expression is either positively or negatively
correlated. Within this topological overlapping network (Li and Horvath 2007), genes were
hierarchically clustered, and modules were identified based on the degree of similarity among
genes. We used a merging threshold of 0.2, with a minimum module size of 30 and a mean
connectivity threshold of greater than or equal to 0.7. We used the default parameters of
WGCNA for the rest of the analyses. We then correlated module eigengene values for a given
module across samples via Pearson's correlation and identified modules differentially regulated
between male types at FDR-adjusted $p < 0.05$.

To visualize WGCNA genetic covariance results among modules significantly associated with
male type, we exported final co-expression networks to Cytoscape (Shannon et al. 2003). We
attached information on $\log_2$ fold-change in expression between $III^M$ and $Y^M$ males, as well as
chromosomal location, to the network as metadata so this information could be visualized. To
identify genes that may have more central functions within and across our significant modules,
we ranked genes in descending order based on intramodular connectivity (kWithin), calculated in
WGCNA. Hub genes identified by intramodular connectivity are generally functionally
important genes within a module (Langfelder et al. 2013).

We further analyzed among-gene connections involving a family of odorant-binding protein
genes (*Obp56h*). Specifically, to identify genes that may regulate or be regulated by genes within
the family, we calculated a "connection score" $C_i$ for every gene $i$ as follows:

$$C_i = \sum_{j=1}^{n} a_{ij}|F_j| \, ,$$

where $a_{ij}$ represents the adjacency (the Pearson correlation coefficient raised to the soft-threshold power $\beta$) between gene $i$ and *Obp56h* gene $j$, and $F_j$ represents the $\log_2$ fold-change in expression between $III^M$ and $Y^M$ males for *Obp56h* gene $j$. This weighted product ensured that connections with *Obp56h* genes that are more differentially expressed between male types were prioritized in calculating a gene's connection score. Genes were then ranked by $C_i$ to identify candidate genes that may be strongly tied to *Obp56h* gene expression. Genes with the 100 highest *Obp56h* connection scores were classified as "central genes". We tested for chromosomal enrichment among these central genes using Fisher's exact tests (comparing the number of central and non-central genes on a focal chromosome with the number of central and non-central genes on all other chromosomes) to determine whether the expression of *Obp56h* genes (which are all located on the *M. domestica* chromosome V) might be involved in *trans* regulation with genes located on the $III^M$ proto-Y chromosome.

### *Gene ontology enrichment analysis*

To identify gene ontology (GO) classes and molecular pathways that are enriched among DE genes, across gene modules identified in WGCNA, or among central genes co-expressed with *Obp56h* genes, we used the BiNGO plug-in within Cytoscape (Maere et al. 2005). We identified *D. melanogaster* orthologs for each house fly gene within a given gene list via NCBI blastx best hits (with default parameters) and used the *D. melanogaster* gene name as input. We identified GO terms that are significantly enriched in BiNGO for biological processes, cellular components, and molecular function.

### *Allele-specific expression analysis*

We tested for differential expression of house fly chromosome III genes between the allele on the $III^M$ chromosome and the allele on the standard third chromosome in $III^M$ males. To do so, we followed methods as in previous studies (Meisel et al. 2017; Son and Meisel 2021), which use the Genome Analysis Toolkit (GATK) best practices workflow for single nucleotide polymorphism (SNP) calling to identify sequence variants in our RNA-Seq data (McKenna et al. 2010). We focused our analysis on libraries that were sequenced from head tissue of male house flies that comprise a CS genetic background (Meisel et al. 2015; Son et al. 2019; Adhikari et al. 2021). We used STAR (Dobin et al. 2013) to align reads from a total of 30 head libraries (15 $III^M$ and 15 $Y^M$ libraries) to the house fly reference genome (Musca_domestica-2.0.2). We then followed the same methods and applied the same parameters as we have done previously to identify SNPs and genotype individual strains (Meisel et al. 2017; Son and Meisel 2021). We

performed separate joint genotyping for each house fly strain within a given experiment (a total
of 4 III$^M$ and 4 Y$^M$ strain-by-experimental batch combinations).

We use the following approach to differentiate between III$^M$ and standard chromosome III alleles.
We first identified SNPs in the exonic regions of the top "hub" genes within a WGCNA module
that mapped to house fly chromosome III. We selected SNPs in those genes that are
heterozygous in III$^M$ males and homozygous in Y$^M$ males. We used the genotype of these SNPs
in Y$^M$ males (which possess two standard third chromosome alleles) to determine the standard
chromosome III allele. The allele not found in Y$^M$ genotypes was assigned to the III$^M$
chromosome. We also identified positions where III$^M$ males appear monoallelic for an allele not
found in Y$^M$ males. These positions that exhibit a complete bias for a III$^M$ allele are suggestive of
monoallelic expression of the III$^M$ allele (i.e., no expression from the III allele).

We tested for differences in expression of III$^M$ and standard chromosome III alleles by following
best practices for comparing allele-specific expression  (Castel et al. 2015). First, for each
strain-by-experimental batch combination, we calculated the normalized read depth at each
variable site as the number of mapped reads at that site divided by the total number of mapped
reads throughout the genome. At each variable site, we used Wilcoxon rank sum tests to make
three different pairwise comparisons per site. First, we compared normalized read depths
between III$^M$ and III alleles in III$^M$ males (III$^M$-III). Second, we compared the read depths of the
III$^M$ allele in III$^M$ males with the normalized read depth of both III alleles in Y$^M$ males (III$^M$-Y$^M$).
Third, we compared the read depths of the III allele in III$^M$ males with the normalized read depth
of both III alleles in Y$^M$ males (III-Y$^M$). We set a threshold of significance at $p < 0.05$ for all
comparisons.

**Drosophila melanogaster *RNA-seq data analysis***

We analyzed RNA-seq results reported in a previous study (Shorter et al. 2016) to determine how
knockdown of *Obp56h* affects gene expression in *D. melanogaster*. Shorter et al. (2016)
identified DE genes between *Obp56h* knockdown and control samples. This analysis was done
separately in males and females, and in separate tissue samples within a given sex (head or the
remaining body). We conducted GO enrichment analysis, as described above, on the list of DE
genes in *D. melanogaster* male head tissue upon *Obp56h* knockdown.

We tested if an excess of  DE genes (between *Obp56h* knockdown and controls) are found on the
*D. melanogaster* X chromosome, which is homologous to house fly chromosome III (Foster et
al. 1981; Weller and Foster 1993). This chromosome is known as Muller element A across flies
(Meisel and Scott 2018; Schaeffer 2018). *Obp56h* is located on *D. melanogaster* chromosome
2R (Muller element C), which is homologous to house fly chromosome V. We used Fisher's
exact tests (comparing the number of X and non-X chromosome genes that are DE in a given

tissue within a given sex with the number of X and non-X chromosome genes that are not DE) to
determine whether *Obp56h* knockdown in *D. melanogaster* results in the disproportionate
differential expression of X chromosome genes in male heads, male bodies, female heads, or
female bodies.

We also tested if the same genes are DE between III$^M$ vs Y$^M$ house flies and *Obp56h* knockdown
vs control *D. melanogaster*. Using NCBI blastx best hits, we identified 20 *M. domestica*
transcripts that are orthologous to *D. melanogaster* genes that are DE upon knockdown of
*Obp56h* (11 matches to upregulated *D. melanogaster* genes, and 9 matches to downregulated *D.*
*melanogaster* genes). We compared the mean log$_2$ fold-changes between Y$^M$ and III$^M$ house fly
males for those 20 genes to 10,000 random subsets of log$_2$ fold-change values taken from our
data (10,000 subsamples without replacement of 11 genes to test for an excess of positive log$_2$
fold-change values, and 10,000 subsamples of 9 genes to test for an excess of negative log$_2$
fold-change values; see Additional Files for R script). We assessed significance by calculating
the proportion of replicated subsamples that generated a mean log$_2$ fold-change value more
extreme than our observed mean.

### *Competitive mating assays*

We performed competitive mating experiments in which two different house fly males were
combined with a single female, and we recorded the "winning" male (i.e., the one who
successfully mated with the female), similar to what was done previously (Hamm et al. 2009). In
these experiments, we used the same two house fly strains as in Hamm et al. (2009): a III$^M$ strain
called CS and a Y$^M$ strain called IsoCS. These two strains have a common genetic background
(CS), and only differ in which proto-Y chromosome they carry. Both strains are represented in
the RNA-seq data we analyzed (Meisel et al. 2015; Son et al. 2019), and IsoCS was also
included in a previous RNA-seq study comparing the effects of proto-Y chromosome and
temperature on gene expression (Adhikari et al. 2021). Our experiment differed from previous
work because we reared larvae from each strain at either 18°C and 29°C, whereas Hamm et al.
(2009) worked with flies raised at 28°C. We used the same larval wheat bran diet as done
previously, and we fed adults an *ad libitum* supply of water and an *ad libitum* 1:1 mixture of
dry-milk:sugar. This is also the same diet and rearing protocol used for the flies in the RNA-seq
datasets that we analyzed (Meisel et al. 2015; Son et al. 2019). Male flies were isolated from
females within ~1 hour of eclosion, and each sex was kept separately to ensure that flies had not
mated prior to the experiment.

We carried out two distinct competitive mating experiments: 1) inter-strain competition between
males with different genotypes (i.e., Y$^M$ vs III$^M$) that were reared at the same temperature (363
successful mating trials out of 490 total attempts across 27 experimental batches); and 2)
intra-strain mating between males with the same genotype that were reared at different

temperatures (104 successful mating trials out of 129 total attempts across 7 batches). When we competed flies with different genotypes raised at the same temperature, all males were aged 4-6 d post pupal emergence. When we compared flies with the same genotype raised at different temperatures, 29°C males were aged 4-5 d post emergence and 18°C males were aged 6-7 d post emergence. We aged flies from the colder temperature for more days than flies from the warmer temperature because developmental rate is positively correlated with developmental temperature in flies (Atkinson 1996). The ages we selected ensure that all males were physiologically capable of mating, while also sampling flies at similar physiological ages across experiments. Aging calculations are reported in Supporting Information 1.

The two males in each experiment were labeled using red and blue luminous powder (BioQuip) by shaking the flies in an 8 oz paper cup. The color assigned to males was switched in each successive batch (i.e., blue $Y^M$ and red $III^M$ in one batch, and then red $Y^M$ and blue $III^M$ in the next batch). In addition, we included the genotype or developmental temperature of the blue-colored male as a fixed effect in our statistical analysis (see below), which provides an additional control for color.

For each replicate of the competitive mating assay, we placed the two different males in a 32 oz transparent plastic container, along with a single virgin female. Each plastic container also contained a 1:1 mixture of dry milk:sugar in a 1 oz paper cup and water in a glass scintillation vial plugged with a cotton roll. Virgin females from the LPR strain (Scott et al. 1996) raised at 25°C were used for all combinations of males. The LPR strain has a different genetic background than the males used in the assay, minimizing any effects of co-adaptation between females and a particular subset of males. All flies were transferred into the mating containers using an aspirator and without anesthesia. All matings were performed in a 25°C incubator because copulation latency is too long for experimentally tractable measurement at lower temperatures. The color (i.e., genotype) of the first male to mate was recorded, as well as the time to mate.

We used the glmer() function in the lme4 package in R (Bates et al. 2015) to test for the effects of genotype and temperature on male mating success. First, to test the effect of genotype, we constructed a logistic regression model with developmental temperature, genotype of the blue male, and their interaction as fixed effects. Experimental batch was modeled as a random effect, with the winning male (CS vs IsoCS) as a response variable. We then assessed significance of fixed effects (type II sum of squares) using the Anova() function in the car package in R (Fox et al. 2013). To test for the effect of temperature on mating success, we similarly constructed a logistic regression model with genotype and the developmental temperature of the blue male as fixed effects and experimental batch as a random effect, with the winning male (18°C vs 29°C) as a response variable.

### *Single-choice mating assays*

We performed experiments to measure copulation latency, or the amount of time elapsed before successful mating, according to male type ($Y^M$ or $III^M$). In these experiments, we used the same IsoCS ($Y^M$) and CS ($III^M$) strains as above and in Hamm et al. (2009). We also tested another strain from each genotype. CSrab ($III^M$) was created by backcrossing the $III^M$ chromosome from the rspin strain isolated in New York onto the CS background (Shono and Scott 2003; Son et al. 2019). CSaY ($Y^M$) was created by backcrossing the $Y^M$ chromosome from the aabys genome reference strain onto the CS background (Scott et al. 2014; Meisel et al. 2015). Virgin females used in the assays were all from the LPR strain (Scott et al. 1996), which has a different genetic background than all males tested. In addition, we also assayed LPR males to determine how copulation latencies of $III^M$ and $Y^M$ males compare to those of males from the same genetic background as the females.

We first attempted to test flies reared at the same temperatures as in our competitive mating assays (18°C and 29°C), as well as at an intermediate developmental temperature (22°C). However, we did not generate enough flies at 18°C, and so we only have data for flies raised at 22°C and 29°C. Our results demonstrate that 22°C is a sufficiently low temperature to detect effects of both genotype and developmental temperature on mating success (see below). All larvae from each male strain were reared in 32 oz plastic containers on the same wheat bran diet described above (Hamm et al. 2009). Upon emergence, unmated male and female progeny were separated and fed water and a 1:1 mixture of dry milk:sugar *ad libitum* until assays were conducted. Assays of males raised at 22°C were conducted 10-11 days after eclosion, while those of males raised at 29°C were conducted 6-7 days after eclosion. This ensures that males were assayed at similar physiological ages. Females were all raised at 25°C, and unmated females were aged 8-9 days after eclosion (see Supporting Information 1 for all accumulated degree day calculations).

We followed a similar protocol as in a previous experiment testing copulation latency in *D. melanogaster* (Shorter et al. 2016). Briefly, five males from a single strain were aspirated without anesthesia into an 8 oz container covered with a fine mesh cloth secured by rubber band. Five LPR females were similarly transferred into the container, marking the start of the mating assay. The house flies were then observed every ten minutes over the course of four hours. Copulation latency was determined in two ways. First, we measured the amount of time elapsed between the start of an assay and each successful mating within a container, defined as a male remaining attached to a female for at least 1 minute (Hamm et al. 2009). Male house flies typically remain attached to females for >60 minutes (Bryant 1980), making it unlikely, although possible, for us to miss matings within 10 minute intervals. Individuals who did not mate were excluded from this analysis. Second, we used a binary variable noting whether each male mated during the 4 hour assay. Although we were unable to distinguish between individual males in this

assay, we did not observe any males mate more than once within 4 hours in a pilot study
conducted between one male and five females, suggesting that observed matings were by
different males. All trials were conducted at 22–23°C.

To determine the effects of male type on the amount of time taken to mate, we used the glmer()
function in the lme4 package in R (Bates et al. 2015) to create a mixed effects model, including
male genotype, developmental temperature, and their interaction as fixed effects and batch and
strain as random effects. For the binary measure of copulation latency, we used a binomial
logistic regression of the same model, with whether a fly mated as our dependent variable. We
then assessed significance of fixed effects (type II sum of squares) using the Anova() function in
the car package in R (Fox et al. 2013). Pairwise comparisons between male types (III$^M$, Y$^M$, and
LPR) were conducted using Z-tests of proportions.

## RESULTS AND DISCUSSION

### *Differential expression of odorant binding protein genes between III$^M$ and Y$^M$ males*

We confirmed that the gene expression profiles of III$^M$ and Y$^M$ male heads are minimally
differentiated (Meisel et al. 2015; Son et al. 2019). There are only 40 DE genes between heads of
III$^M$ and Y$^M$ adult males (21 upregulated in Y$^M$ males, 19 upregulated in III$^M$, Table S2). Gene
ontology analysis revealed no significant biological process, molecular function, or cellular
component terms enriched within the list of DE genes.

Within the list of DE genes, we identified one Obp gene (*LOC105261916*) upregulated in Y$^M$
males. House fly Obp genes can be grouped into families corresponding to their *D. melanogaster*
orthologs (Scott et al. 2014). The DE Obp gene in our analysis is orthologous to *Obp56h*. The
*Obp56h* family, as well as other Obp families, was greatly expanded within muscids (house fly
and close relatives, including stable fly and horn fly) compared to *D. melanogaster* (Scott et al.
2014; Olafson and Saski 2020; Olafson et al. 2021). In addition to *LOC105261916*, seven of the
remaining eight house fly *Obp56h* genes for which we obtained RNA-seq count data showed
similar trends of greater expression in Y$^M$ than III$^M$ males, with three of these showing
significant DE ($p<0.05$) before an FDR correction (Fig. 1). All but one of the *Obp56h* genes has
higher expression in Y$^M$ than III$^M$ males (8/9, regardless of significance), which is significantly
greater than the fraction of other genes with higher expression in Y$^M$ males, regardless of
significance, in the rest of the genome (Fisher's exact test, $p = 0.019$). Moreover, the expression
levels of several house fly *Opb56h* genes are sensitive to developmental temperature or the
interaction between temperature and male genotype (Fig S1).

**Figure 1** - Neighbor-joining phylogenetic tree of the *Obp56h* gene family within *M. domestica* and *D. melanogaster* based on protein sequences constructed in MEGA X (Kumar et al. 2018). Amino acid sequences were aligned by MUSCLE (Edgar 2004). *M. domestica Obp56h* genes are identified based on gene IDs. The bootstrap consensus tree was inferred from 10,000 replicates. Branch lengths are scaled according to the number of amino acid substitutions per site. The phylogeny was arbitrarily rooted at *D. melanogaster Obp56h.* Batch-adjusted expression levels for each *M. domestica Obp56h* gene from each replicate are displayed at the branch tips (small circles). Large circles show the average across all replicates, with error bars denoting the standard error (unfilled stars: $p < 0.05$ before FDR correction for multiple comparisons; filled star: $p < 0.05$ after correction).

## III$^M$ confers a mating advantage that is robust to developmental temperature

Knockdown of *Obp56h* in *D. melanogaster* results in decreased male copulation latency, or the time it takes for a male to begin to mate with a female after they are first exposed to one another (Shorter et al. 2016). The *Obp56h* gene family is generally expressed higher in Y$^M$ males relative to III$^M$ males (Fig. 1). A previous study identified a competitive mating advantage of III$^M$ over Y$^M$ male house flies (Hamm et al. 2009), consistent with shorter copulation latency in III$^M$ males because of lower expression of *Obp56h* genes. Segregating variation in such a strongly sexually selected trait likely requires some other factor to maintain the difference in copulation latency between Y$^M$ and III$^M$ males (Kirkpatrick and Ryan 1991; Ingleby et al. 2010). The clinal distribution of Y$^M$ and III$^M$ (Tomita and Wada 1989; Hamm et al. 2005; Kozielska et al. 2008) are suggestive that temperature may differentially affect males carrying these proto-Y chromosomes. In addition, two *Obp56h* genes are only upregulated in Y$^M$ males at 29°C, but not at 18°C (Fig S1), suggesting the effect of Y$^M$ on male mating success may be temperature-dependent. The previous experiment only compared mating performance of Y$^M$ and III$^M$ males at 28°C  (Hamm et al. 2009). We therefore tested if the differences in mating success between Y$^M$ and III$^M$ males are sensitive to temperature and are thus context-dependent.

We performed competitive mating assays in which we allowed males carrying III$^M$ or Y$^M$, reared
at either 18°C or 29°C, to compete for a female of an unrelated strain. We found that III$^M$ males
were more successful at mating than Y$^M$ males regardless of developmental temperature
(ANOVA, $p = 6.53$ x $10^{-6}$, Fig. 2A). This mating advantage of III$^M$ males is consistent with
reduced copulation latency as a result of lower expression of *Obp56h* genes. However, our
results suggest that there is not an effect of developmental temperature on the III$^M$ male mating
advantage.

**Figure 2** - III$^M$ chromosome and developmental temperature affect male mating success. A) Outcomes of
competitive mating assays between III$^M$ and Y$^M$ males reared at 18°C or 29°C. Data points represent
experimental batches. Horizontal lines denote the median across all batches. B) Outcomes of competitive
mating assays conducted between males reared at 29 °C and 18 °C. Trials were conducted between males
of the same proto-Y chromosome genotype (III$^M$ or Y$^M$). Each data point represents ten replicate trials
within a single batch. C) Outcomes of single-choice mating assays in males reared at 22°C. Data points
refer to the percentage of males (five males within one replicate) that mated with females within 4 hours
within each experimental trial. Horizontal lines denote means within male groups. All females used were
from the LPR strain.

To further investigate if there are effects of developmental temperature on mating success, we
tested whether males reared at different developmental temperatures, but with the same
genotype, have a difference in mating success. We found that males reared at 18°C outcompeted
males reared at 29°C (ANOVA, $p = 2.93$ x $10^{-4}$, Fig. 2B), regardless of genotype. This is
consistent with decreased *Obp56h* expression in Y$^M$ males reared at 18°C (Fig S1), which
outcompete Y$^M$ males raised at 29°C. In contrast, III$^M$ males exhibited low *Obp56h* expression,
regardless of developmental temperature (Fig. S1), yet III$^M$ males raised at 18°C outcompeted
III$^M$ males reared at 29°C. Therefore, *Obp56h* expression levels alone cannot explain the effect
of developmental temperature on mating success.

We conclude that there is substantial evidence that both proto-Y chromosome genotype and
developmental temperature affect male mating success. Notably, the effect of temperature on
mating success is consistent across $Y^M$ and $III^M$ male types, suggesting that context-dependence,
in the form of a genotype-by-temperature interaction, cannot explain why genetic variation in
male mating success is maintained in house fly. However, our mating assays were all performed
at the same temperature (25°C), with male house flies that developed at two different
temperatures. We therefore cannot rule out the possibility that differences in mating success are
affected by courtship temperature (as opposed to developmental temperature).

### $III^M$ males have reduced copulation latency

Our competitive mating assays (Fig. 2A), as well as previously published results (Hamm et al.
2009), raise the possibility that $III^M$ males have a reduced copulation latency relative to $Y^M$
males, which could be explained by the lower expression of *Obp56h* genes in $III^M$ males (Fig. 1).
We therefore directly measured copulation latency in single-choice mating assays. To do so, we
combined five males from a single strain raised at a single temperature with five females from
the unrelated strain used in our competitive mating assays. Developmental temperature had a
significant effect on copulation latency (ANOVA, $p = 9.40 \times 10^{-5}$), with males reared at 22°C
mating faster than those reared at 29°C (Fig. S2A). In general, successful matings were rare for
males that developed at 29°C regardless of whether they carry the $Y^M$ or $III^M$ chromosome (15
successful matings out of 75 males tested). These results are consistent with increased mating
success of males raised at 18°C relative to those raised at 29°C in our competitive mating assays
(Fig. 2B).

We found no significant effect of male genotype (ANOVA, $p = 0.89$), or the interaction between
male genotype and developmental temperature (ANOVA, $p = 0.37$), on the time it takes males to
mate. When we consider only males who mated within 4 hours, the time to mate for $Y^M$ and $III^M$
males did not significantly differ at either 22°C (Tukey's post-hoc, $p = 0.99$) or 29°C (Tukey's
post-hoc, $p = 0.78$). However, this analysis is problematic because we have no measure of
copulation latency for males that did not mate within the 4 hour experimental window, which
amounts to >70% of males in three of the four genotype-by-temperature combinations. It is
therefore possible that copulation latency does indeed differ between $Y^M$ and $III^M$ males in a way
that is not detected in our censored data.

To overcome the problems associated with censored data, we next treated copulation latency as a
binary variable by calculating the proportion of the five males per trial that mated within the 4
hour assay. We observed significant effects of male genotype (ANOVA, $p = 6.18 \times 10^{-3}$) and
developmental temperature (ANOVA, $p = 9.04 \times 10^{-4}$) on the proportion of males that mated. The
effect of developmental temperature was largely a result of very few matings for males that
developed at 29°C relative to 22°C (Fig. S2B). The lower copulation latency for flies that

developed at 22°C is consistent with our competitive mating assays that showed males that
developed at a lower temperature have higher mating success (Fig. 2B). In the 22°C treatment, a
significantly greater proportion of III$^M$ males mated within 4 hours than Y$^M$ (61.7% v. 28.3%;
Z-test of proportions, $p = 1.21$ x $10^{-4}$; Fig. 2C). This is evidence that III$^M$ males have reduced
copulation latency, which is consistent with their previously documented competitive mating
advantage (Hamm et al. 2009), the competitive mating advantage that we observe (Fig. 2A), and
the reduced expression of *Obp56h* genes (Fig. 1).

The Y$^M$ and III$^M$ males we and others used in mating experiments all share the CS genetic
background that comes from a III$^M$ strain (Hamm et al. 2009). This raises the possibility that III$^M$
males perform better because they have a proto-Y chromosome that is co-adapted to its genetic
background. To test this hypothesis, we measured copulation latency in Y$^M$ males from the same
strain (LPR) as the females in our experiments. We observed a greater proportion of III$^M$ males
mating within 4 hours when compared to the LPR Y$^M$ males (Z-test of proportions, $p = 0.038$),
although the copulation latency in LPR males was highly variable (Fig. 2C). Therefore, the
reduced copulation latency conferred by the III$^M$ chromosome overwhelms any potential effects
of coadaptation of the proto-Y chromosome to male genetic background or male-female
co-adaptation within strains. The reduced copulation latency of III$^M$ males is only detectable
when house flies develop at 22°C, suggesting that it is either temperature-dependent or we lack
the resolution to detect it when males develop at warmer temperatures (because they take too
long to mate). Our results also provide evidence that the effect of temperature on copulation
latency is independent of genotype, suggesting that context-dependent effects of the proto-Y
chromosomes cannot explain the maintenance of the polymorphism.

***House fly chromosome III genes and* Drosophila *X chromosome genes have correlated***
***expression with* Obp56h *genes***

In order to identify the regulatory architecture underlying the differential expression of *Obp56h*
genes between Y$^M$ and III$^M$ males, we identified 27 co-expression modules across house fly male
heads. One of these modules (containing 122 genes, Table S3) is differentially expressed
between Y$^M$ and III$^M$ males (FDR adjusted $p = 0.001$, Fig. 3). GO analysis revealed significant
enrichment (FDR adjusted $p < 0.05$) of 15 biological process terms including those related to
immune system processes (GO:0032501), responses to stress (GO:0006950), and response to
external stimuli (GO:0009605) within this module (Table S4). This module is also enriched for
house fly chromosome III genes (31 chromosome III genes versus 38 genes assigned to other
chromosomes, Fisher's exact test $p < 1$ x $10^{-5}$, with 53 genes not assigned to a chromosome) and
for DE genes between Y$^M$ and III$^M$ males (16 DE genes in this module versus 24 DE genes
assigned to other modules, Fisher's exact test $p < 1$ x $10^{-5}$). We used the WGCNA measure of
intramodular connectivity, kWithin, to identify hub genes within the module that likely have
important roles in the regulation of gene expression. The top five hub genes are (with

*D. melanogaster* orthologs in parentheses): *LOC101887703 (CG8745), LOC105262120*
(*CG10514*), *LOC101894501 (Gr98c), LOC101893264 (gd)*, and *LOC101893651 (CG2120)* (Fig.
3, S3).

**Figure 3** - Network visualization of the co-expression module that is differentially regulated between III$^M$
and Y$^M$ males. Each circle within the module is a gene, and *Obp56h* genes are indicated with purple fill.
Lines represent edge connections between genes. Genes labeled with "+" are within the top 100 most
strongly connected to *Obp56h* genes. Genes are ordered from top to bottom according to intramodular
connectivity (kWithin), with genes of higher connectivity (i.e., hub genes) on top, and peripheral genes on
the bottom. Borders around genes denote log$_2$ fold-change in expression between Y$^M$ and IIII$^M$ male
heads, with darker blue borders denoting upregulation in Y$^M$, and darker red borders denoting
upregulation in III$^M$. Chromosomal locations in house fly (*Mdom*) and *D. melanogaster* (*Dmel*) are shown
for the 5 hub genes and *Obp56h*.

Three *Obp56h* genes that are DE between III$^M$ and Y$^M$ males (*LOC105261916, LOC101891822,*
and *LOC101891651*) are all assigned to the co-expression module (Fig. 3). The *Obp56h* gene
cluster itself is found on house fly chromosome V, which is unlikely to differ between the Y$^M$ and
III$^M$ males in our experiments—the majority of males compared in the RNA-seq data and mating
experiments have a common genetic background (including chromosome V) and differ only in
whether they carry III$^M$ or Y$^M$. Removing samples with a different background did not affect the
general difference in *Obp56h* expression between III$^M$ and Y$^M$ males (see Supporting Information
2 for a summary of these results). We therefore hypothesized that differential *Obp56h* expression
is at least partly controlled by *trans* regulatory variation that maps to Y$^M$, III$^M$, or both.

If *Obp56h* gene expression is regulated by *trans* factors that map to chromosome III, then we expect the *Obp56h* gene family to be found within, or strongly connected to genes within, the gene module that is differentially regulated between III$^M$ and Y$^M$ males. Indeed, we find that the module is enriched for *Obp56h* genes relative to other Obp genes—three *Obp56h* genes and no other Obp genes were assigned to this module (Fisher's exact test, $p = 5.1 \times 10^{-3}$, Fig. 3). This suggests that *Obp56h* expression is strongly correlated with, and is either regulated by or regulates, genes whose expression is affected by the III$^M$ chromosome. We cannot perform the same analysis for the effect of Y$^M$ because only 51 genes have been assigned to the house fly X/Y$^M$ chromosome (Meisel and Scott 2018), limiting our power to detect an excess of genes.

To test for *trans* regulators of *Obp56h* gene expression, we identified house fly genes whose expression covaries with *Obp56h* genes (Table S5). We found significant enrichment for chromosome III genes within the 100 genes whose expression covaries most with *Obp56h* gene expression (corresponding to the top 0.55% covarying genes); of the 100 genes with the highest *Obp56h* connection scores, 26 are on chromosome III (Fisher's exact test $p = 2.0 \times 10^{-4}$, Fig. 4A). This enrichment is robust to varying the threshold used to classify a gene as in the top covarying; considering genes with the top 1%, 5%, 10% covarying expression also resulted in significant enrichment of chromosome III genes (Fisher's exact test, all $p < 0.05$). This supports the hypothesis that *trans* regulatory variants that differ between III$^M$ and the standard chromosome III are at least partially responsible for DE *Obp56h* genes between III$^M$ and Y$^M$ house fly males.

Our network analysis does not ascribe directions to the edges connecting house fly genes, and it is therefore possible that *Obp56h* DE has *trans* regulatory effects on chromosome III expression. To test this hypothesis, we examined available RNA-seq data from an experiment comparing wild type *D. melanogaster* with flies in which *Obp56h* had been knocked down (Shorter et al. 2016). *Obp56h* is on the right arm of the second chromosome in *D. melanogaster* (2R, or Muller element C), which is homologous to house fly chromosome V (Foster et al. 1981; Weller and Foster 1993). House fly chromosome III is homologous to the *D. melanogaster* X chromosome, which is known as Muller element A (Meisel and Scott 2018; Schaeffer 2018). The *D. melanogaster* males in the RNA-seq experiment all share the same X chromosome, and only differ in one copy of their second chromosome (which either carries a UAS-RNAi knockdown construct or does not). If *Opb56h* genes have *trans* regulatory effects on element A genes in males, we would expect an excess of DE *D. melanogaster* X chromosome genes *Obp56h* knockdown flies. Indeed, we found that *Obp56h* knockdown in *D. melanogaster* resulted in excess DE of X chromosome genes in male head (Fisher's exact test, $p = 0.011$, Fig. 4B) and body ($p = 0.038$, Fig. S4), but not in either tissue sample in females (Fisher's exact test, both $p > 0.49$). These results suggest that there is male-specific *trans* regulatory control of *D. melanogaster* X-linked genes by *Obp56h*. This regulatory architecture associated with *Obp56h* expression may create epistatic or pleiotropic constraints that inhibit selection to reduce

expression of *Obp56h* genes, which should otherwise be favored because reduced expression
shortens copulation latency.

**Figure 4** - Percent of genes on each chromosome within (A) the top 100 genes with the strongest
connections to the *Obp56h* family in house fly, and (B) genes differentially expressed (DE) between
*Obp56h* knockdown and control *D. melanogaster* (black bars: males, grey bars: females). Asterisks
indicate a significant difference between observed (bars) and expected (red lines) counts of genes on each
chromosome compared to all other chromosomes (Fisher's exact test, $p < 0.05$).

Our house fly results suggest that alleles differing between III$^M$ and the standard chromosome III
have *trans* effects on *Obp56h* expression. The *D. melanogaster* data suggest that variation in
*Obp56h* expression affects the expression of genes on the *D. melanogaster* X chromosome
(which is homologous to house fly chromosome III). This raises the possibility that there is *trans*
regulatory feedback from *Obp56h* genes that affects chromosome III gene expression in house
fly. If that regulatory feedback is conserved between house fly and *D. melanogaster*, we expect
that orthologous genes would be DE on Muller element A (house fly chromosome III and the *D.*
*melanogaster* X chromosome) between III$^M$ and Y$^M$ house flies and between *Obp56h* knockdown
and wild-type *D. melanogaster*. Consistent with this expectation, we found that genes that are
downregulated upon knockdown of *Obp56h* in *D. melanogaster* have house fly orthologs that are
more downregulated in III$^M$ male house flies (i.e., lower log$_2$ fold-change) than expected by
chance ($p = 5.60 \times 10^{-3}$, Fig. S5A). In contrast, genes that were upregulated upon *Obp56h*
knockdown in *D. melanogaster* were not significantly differentially regulated between Y$^M$ and
III$^M$ male genotypes, although the observed trend suggests that these genes may be more
downregulated in III$^M$ males than expected ($p = 0.103$, Fig. S5B). Our results therefore suggest
that, genome-wide, the genes which vary in expression along with *Obp56h* (regardless of the
direction of *trans* regulation) are evolutionarily conserved between *M. domestica* and
*D. melanogaster*. Consistent with this hypothesis, we identified the GO term "response to stress"
(GO:0033554) to be significantly enriched amongst genes with strong connection scores with
*Obp56h* expression in *M. domestica* and in the list of DE genes in *D. melanogaster* upon *Obp56h*

knockdown (Table S6). This provides further evidence that there is an evolutionarily conserved
*trans* regulatory feedback loop involving *Obp56h* expression in *Drosophila* and house fly
through similar molecular functions.

*Network analysis reveals candidate regulators of* **Obp56h** *expression*

The house fly co-expression module contains candidate genes and pathways through which
*Obp56h* genes, and likely male copulation latency, are regulated. For example, within the list of
DE genes, we identified one gustatory receptor gene (*LOC101894501*, the ortholog of
*D. melanogaster Gr98c*) upregulated in III$^M$ males ($p_{ADJ}$ = 0.037). Although Obps can have a
variety of functions, they most typically interact with chemosensory receptors (odorant,
ionotropic, and gustatory receptors) in the detection of chemical cues or signals (Zhou 2010;
Benoit et al. 2017; Sun et al. 2018). If *Obp56h* serves a sensory detection role in male house fly
heads, then *Gr98c* is a promising candidate gene with which it interacts. The *M. domestica*
ortholog of *Gr98c* (*LOC101894501*, which we will refer to as *Md-Gr98c*) is a hub gene in the
co-expression module containing *Obp56h* (Fig. 3). *Md-Gr98c* is the only chemosensory receptor
assigned to this module, it is located on chromosome III, and its expression is negatively
correlated with *Obp56h*. A negative correlation between the expression of a chemosensory
receptor and its interacting binding protein has previously been reported in a pair of genes that
modulate male *Drosophila* mating behavior (Park et al. 2006).

**Figure 5** - Allele-specific expression (ASE) in A) *LOC101894501* (*Md-Gr98c*), B) *LOC101893264*
(*Md-gd*), and C) *LOC101893651* (*CG2120* ortholog). The x-axis depicts base pair positions (scaffold
coordinates) of the informative single nucleotide polymorphisms (SNPs) that differ between III$^M$ and
standard chromosome III alleles. The y-axis and data points depict the read depth of a given allele
normalized by the total mapped reads for a given strain-by-experimental batch group combination (FPM
= fragments per million). Lines depict mean read depths at each diagnostic site for III (turquoise) and III$^M$

(salmon) alleles in $III^M$ males, and mean read depths at each site for III alleles in $Y^M$ males (black). Tables
under each graph mark significant differences (*: $p < 0.05$) in normalized read depths at each diagnostic
site for each of three pairwise comparisons: $III^M$ allele vs. III allele in $III^M$ males ($III^M$-III), $III^M$ allele in
$III^M$ males vs both III alleles in $Y^M$ males ($III^M$-$Y^M$), III allele in $III^M$ males vs. both III alleles in $Y^M$ males
($III$-$Y^M$).

We tested if *Md-Gr98c* is differentially regulated between the $III^M$ chromosome and standard
chromosome III by comparing expression in $III^M$ males (i.e., heterozygotes for $III^M$ and a
standard chromosome III) with males homozygous for the standard chromosome III. Differential
expression of the $III^M$ and III chromosome alleles would implicate this gene as having a causal
effect on *Obp56h* expression. *Md-Gr98c* contains 4 exonic SNPs differentiating the $III^M$ and III
chromosomes. Within each $III^M$ strain in each RNA-seq experiment, we observed significantly
greater expression of the $III^M$ allele than the standard chromosome III allele at two of the four
diagnostic SNP sites (Fig. 5A). The other two SNPs showed the same pattern of $III^M$-biased
expression but were not significant (both $p > 0.05$). The $III^M$ allele is also expressed higher than
both III alleles in $Y^M$ males. Higher expression of the $III^M$ allele is consistent with *cis* regulatory
divergence between the $III^M$ and standard chromosome III being partially responsible for
elevated *Md-Gr98c* expression in $III^M$ males. Furthermore, the standard chromosome III allele is
expressed significantly higher in $III^M$ males than $Y^M$ males at two of the four diagnostic SNP
sites (Fig. 5A); we observe the same pattern at the other two sites without significance ($p >$
0.05). Higher expression of the III allele in $III^M$ males than $Y^M$ males suggests that *trans*
regulators further increase the expression of *Md-Gr98c* in $III^M$ males. This combination of *cis*
and *trans* regulatory affects on *Md-Gr98c* expression are consistent with the *trans*-regulatory
loop we hypothesize between *Obp56h* and chromosome III that regulates male mating behavior.
Future experiments could determine whether *Gr98c* and *Obp56h* do indeed interact and, if so,
what pheromonal or other chemical compounds they detect.

A serine protease gene, *LOC101893264*, orthologous to *D. melanogaster gd* (Konrad et al.
1998), is also among the top 5 hub genes within the co-expression module (Fig. 3). This gene is
predicted to encode a positive regulator of the Toll signaling pathway (Valanne et al. 2011),
suggesting that the *M. domestica* ortholog of *gd* (*Md-gd*) could have an important gene
regulatory function within the module via Toll signaling. *Md-gd* is located on chromosome III,
and it is upregulated in $III^M$ males (adj. $p = 0.022$). We identified seven sites where all RNA-seq
reads were mapped to the $III^M$ allele, while no reads were mapped to the standard chromosome
III allele in $III^M$ males (Fig. 5B). At all seven diagnostic SNP sites in this gene, the $III^M$ allele is
significantly more highly expressed than the III allele in $III^M$ males (all $p = 0.021$), and it is more
highly expressed than both III alleles in $Y^M$ males at six of seven sites (all $p = 0.021$). The lack of
expression of the III allele in either $III^M$ or $Y^M$ males is consistent with monoallelic gene

expression of the III$^M$ allele, although further evidence is required to confirm this hypothesis (see
Supporting Information 3 for detailed discussion).

We identified similar evidence of monoallelic gene expression within another hub gene,
*LOC101893651*, which is orthologous to *D. melanogaster CG2120* (Fig. 5C). *LOC101893651* is
among the most central genes within the co-expression module (Fig. 3), and it is strongly
upregulated in III$^M$ males (log$_2$ fold-change: 1.33, adjusted $p$ = 0.033). *LOC101893651* is found
on house fly chromosome III and is predicted to encode a transcription factor. At all four
diagnostic sites within *LOC101893651*, the III$^M$ allele is significantly more highly expressed
than the III allele in III$^M$ males (all $p$ = 0.021), as well as both III alleles in Y$^M$ males (all $p \leq$
0.027). Within the WGCNA module, *Obp56h* expression is most strongly correlated with
*LOC101893651*, suggesting that *LOC101893651* could encode the transcription factor that is
directly responsible for the repression of *Obp56h* expression in III$^M$ males.

Based on their differential expression, allele-specific expression, centrality within the
co-expression module, and their location on chromosome III, *Md-Gr98c*, *Md-gd*, and
*LOC101893651* are all strong candidate genes that may directly or indirectly affect the
expression of house fly *Obp56h* genes. However, the expression of genes on other chromosomes
are also strongly correlated with *Obp56h* expression (Fig. 3), suggesting other chromosomes may
also be involved in the *trans* regulatory loop. In addition, many house fly genes are not yet
mapped to chromosomes (Meisel and Scott 2018), limiting our ability to infer the chromosomes
involved in regulating *Obp56h* expression. We describe two genes (*LOC105262120* and
*LOC101887703*) that may be located on other chromosomes below.

Expression of *LOC101893651* (the ortholog of *CG2120* that is predicted to encode a
transcription factor) is most strongly correlated with *LOC105262120* (an ortholog of *CG10514*),
which is predicted to encode an ecdysteroid kinase. Expression of *Obp56h* and *CG10514* are
correlated with the production of CHCs that are involved in social behavior in *D. melanogaster*
(Shorter et al. 2016) and *Drosophila serrata* (McGraw et al. 2011), respectively. CHCs are often
under strong sexual selection across insect systems, with individual or combinations of CHCs
serving as important mating cues (Thomas and Simmons 2009,Berson et al. 2019a,b). The
correlation of these genes with CHC profiles in *Drosophila* provides additional evidence that
*Obp56h* expression, and the house fly co-expression module more generally, are related to male
mating behavior, and possibly under sexual selection. However, cuticular hydrocarbons also
provide protection against biotic and abiotic stressors (Otte et al. 2018) and play a crucial role in
desiccation resistance (Lockey 1988). This dual role of CHCs in mating and stress resistance
suggests that sexual selection on *Obp56h* expression could be pleiotropically constrained by
trade-offs with stress response. Our GO enrichment analysis on both the house fly and *D.*
*melanogaster* RNA-seq data also revealed that *Obp56h* expression is correlated with the
expression of genes involved in general stress responses, supporting this hypothesis. Pleiotropic

constraints on *Obp56h* expression (because of correlated changes in CHCs) could therefore
reduce the response to selection on male copulation latency, contributing to the maintenance of
genetic variance. Future studies should aim at determining whether III$^M$ and Y$^M$ male house flies
also differ in CHC profiles, as well as their resistance to desiccation and other environmental
stressors.

**Figure 6** - Correlations of gene expression between *Obp56h* (house fly *LOC105261916*) and *CG8745*
(*LOC101887703*) in (A) house fly male head tissue, and (B) *D. melanogaster* male head tissue. Values for
*D. melanogaster* are from count data as reported in Shorter et al. (2016). Linear regression models were
used to determine 95% confidence intervals (shaded in grey) summarizing the effect of *Obp56h*
expression on *CG8745* expression in each species.

*LOC105262120* expression is most strongly correlated with the most central gene within the
co-expression module, *LOC101887703*. *LOC105262120* and *LOC101887703* are both
upregulated in III$^M$ males (*LOC101887703*: log$_2$ fold-change: 2.21, adj. $p = 0.016$;
*LOC105262120*: log$_2$ fold-change: 2.26, adj. $p = 0.007$). *LOC101887703* is orthologous to
*D. melanogaster CG8745*, which is predicted to encode an ethanolamine-phosphate
phospho-lyase and is broadly expressed in many *D. melanogaster* tissues (Chintapalli et al.
2007). In both *D. melanogaster* and house fly, *Obp56h* expression is significantly negatively
correlated with the expression of *CG8745* or *LOC101887703*, respectively (Fig. 6).

*LOC101887703* has a paralog (*LOC101890114*) that is predicted to be on chromosome III. The
two transcripts encoded by these paralogs are <1% diverged in their nucleotide sequences,
suggesting a recent duplication event. Gene duplication is hypothesized to be involved in the
resolution of inter-sexual conflict (Connallon and Clark 2011; Gallach and Betrán 2011; Van
Kuren and Long 2018), raising the possibility that at least one of the two house fly paralogs is
partitioned into a sex-specific function. Moreover, broadly expressed genes often give rise to
paralogs with sex-specific expression (Meisel et al. 2009). Notably, chromosome III is a proto-X
chromosome (III$^M$ is the proto-Y), and there is a general excess of gene duplication from X
chromosomes to the autosomes across flies and other animals, possibly driven by selection on
sex-specific functions (Betrán et al. 2002; Emerson et al. 2004; Meisel et al. 2009; Baker and

Wilkinson 2010). Future work could address a potential sexually dimorphic subfunctionalization
of the two *CG8745* paralogs in the house fly genome.

Together, the hub genes in the co-expression module are likely to be members of an
evolutionarily conserved *trans* regulatory feedback loop that controls and/or is controlled by
*Obp56h* expression (Fig. 7A). Specifically, we infer that *Obp56h* regulates *CG8745*
(*LOC101887703*) because knockdown of *Obp56h* causes an increase in *CG8745* expression in
*D. melanogaster* (Fig. 6B). Based on their locations on house fly chromosome III, positions as
hub genes in the house fly co-expression module (Fig. 3), and divergent expression between the
III$^M$ and standard III chromosomes (Fig. 5), we also hypothesize that *LOC101893651* (*CG2120*),
*Md-gd*, and/or *Md-Gr98c* regulate *Obp56h* in house fly (Fig. 7B). *Md-Gr98c* is a particularly
promising candidate, because chemosensory binding proteins and receptors are known to
co-regulate one another (Park et al. 2006). Future manipulative experiments will help in further
evaluating the direction of regulation of these co-expressed genes.

**Figure 7** - Hypotheses on connections between *Obp56h* expression, proto-Y chromosome genotype, and
male mating behavior based on house fly and *D. melanogaster* gene expression data. A) Summary of
evidence for an evolutionarily conserved *trans* regulatory loop between *Obp56h* and Muller Element A
(house fly chromosome III, and *D. melanogaster* X chromosome). Our hypothesis is based on differential
expression between III$^M$ vs. Y$^M$ male house flies, *Obp56h* knockdown vs. control *D. melanogaster*, and
network connectivity of *Obp56h* family gene expression within house fly. B) Summary of candidate
genes implicated in conserved *trans* regulatory loop. Three of the top five hub genes of module A are
located on house fly chromosome III, are negatively correlated with *Obp56h* expression, and exhibit
either allele-specific expression (ASE) or show signs of monoallelic gene expression biased towards the
III$^M$ allele. Similar correlations between expression measures of *Obp56h* and *CG8745* (*LOC101887703*)
in *D. melanogaster* and house fly male head tissue suggest that *Obp56h* regulates *CG8745*, which is the
primary hub gene in the WGCNA module that is differentially expressed between III$^M$ and Y$^M$ male house

flies. Shared correlations between *Obp56h* expression and copulation latency in both house fly and *D.*
*melanogaster* also suggest that *Obp56h* regulates male fly mating behavior.

## CONCLUSIONS

In this study, we aimed to identify a genetic mechanism explaining phenotypic variation in male mating performance across male proto-Y chromosome genotypes in house fly (Hamm et al. 2009). An *Obp56h* gene is among our small list of genes differentially expressed between III$^M$ and Y$^M$ male heads (Fig. 1). In *D. melanogaster*, *Obp56h* expression affects male copulation latency (Shorter et al. 2016), and we identified a similar difference in copulation latency between male proto-Y genotypes in house fly (Fig. 2). We also identified an excess of house fly chromosome III (Muller element A) genes strongly co-expressed with the *Obp56h* gene family, suggesting *trans* regulation of *Obp56h* by genes from element A (Fig. 3, 4, 5). Similarly, we observe evidence for excess *trans* regulation of element A genes in *D. melanogaster* by *Obp56h* (Fig. 4). The inferred directions of *trans* regulation between *Obp56h* and element A are in opposite directions in house fly and *D. melanogaster* (Fig. 7A), but an excess of the same element A genes are correlated with *Obp56h* expression in both species. This suggests that there is an evolutionarily conserved *trans* regulatory loop affecting and affected by *Obp56h* expression (Fig. 7) between fly species that diverged >50M years ago (Wiegmann et al. 2011). The shared relationship between *Obp56h* expression and copulation latency in both species suggests that this *trans* regulatory loop has conserved effects on male mating behavior across distantly related flies.

House flies and *Drosophila* have independently acquired the same sex chromosome (Muller element A is house fly chromosome III, which is homologous to the *Drosophila* X). This raises the possibility that element A is primed to be recruited as a sex chromosome because of the *trans* regulatory connections with a gene (*Obp56h*) that has important effects on male mating behavior. Convergent sex-linkage of the same chromosomal region has been observed in vertebrates (O'Meally et al. 2012; Furman and Evans 2016; Ezaz et al. 2017), which could be explained by the same gene independently acquiring a sex determining allele in mutiple independent lineages (Takehana et al. 2014). Genes with sex-specific effects (including sexually antagonistic variants) are also expected to be an important selective force in the formation of new sex chromosomes (van Doorn and Kirkpatrick 2007) and the subsequent evolution of X and Y chromosomes (Gibson et al. 2002; Charlesworth et al. 2005; Abbott et al. 2017). Our results suggest that an enrichment of genes that regulate sexually selected behaviors could promote the sex-linkage of the same chromosome in distantly related species without convergent evolution of a master sex determiner.

The apparently simple correlation between *Obp56h* expression and male mating behavior suggests that selection for reduced male copulation latency in house fly could easily be achieved by downregulation of *Obp56h* expression. Our results address three possible mechanisms for the paradoxical maintenance of genetic variation underlying this sexually selected trait: context-dependence, epistasis, and pleiotropy. To address context-dependence, we tested if the effects of $Y^M$ and $III^M$ on copulation latency vary across temperatures. $Y^M$ and $III^M$ are distributed across a latitudinal cline, suggesting that temperature differentially affects the fitness of males according to proto-Y chromosome genotype (Tomita and Wada 1989; Hamm et al. 2005; Feldmeyer et al. 2008; Kozielska et al. 2008). We found that temperature does indeed affect male mating performance, but $III^M$ males outperform $Y^M$ males regardless of temperature (Fig. 2). In addition, the *Obp56h* gene with the strongest signal of differential expression (*LOC105261916*) is only differentially expressed in $Y^M$ males across temperatures (Fig S1), suggesting that *Obp56h* expression levels alone cannot explain the effect of temperature on mating success. Therefore, we hypothesize that proto-Y chromosome genotype and developmental temperature have independent effects on male mating success and copulation latency. The lack of a genotype-by-environment interaction suggests that context-dependent effects cannot explain the maintenance of variation in male copulation latency in house flies.

The interaction between proto-Y chromosome genotype and *Obp56h* expression is suggestive that epistasis may reduce the response to selection on copulation latency in house fly. Similar to our results, Y chromosome genotype in *D. melanogaster* is a major determinant of male fitness, but this effect depends on genetic background (Chippindale and Rice 2001). This suggests that epistatic interactions reduce the efficacy of selection on male-beneficial Y-linked alleles in *D. melanogaster*. Our experiments were not explicitly designed to test for the effect of genetic background. Nonetheless, we found that LPR males exhibit similar delayed copulation latencies as the other $Y^M$ males we tested (Fig. 2C), despite having a different genetic background. This suggests that epistatic interactions do not necessarily modulate the effect of $Y^M$ and/or $III^M$ on male mating. However, we are unable to tease apart the effects of the $Y^M$ chromosome from those of the remaining genetic background, as LPR males also possess a different $Y^M$ chromosome than the other $Y^M$ males we tested. Regardless of this limitation, our results are consistent with the hypothesis that the effect of *Obp56h* expression on male copulation latency depends on male proto-Y chromosome genotype via *trans* regulation, which leaves open the possibility that epistasis constrains the response to selection for this trait.

Our results provide stronger evidence that pleiotropic constraints maintain genetic variation for male mating behavior in house fly. The *trans* regulatory loop between *Obp56h* expression and multiple genes on Muller element A may create pleiotropic constraints that weaken the efficacy of selection on this trait. For example, *trans* regulators are predicted to have pleiotropic effects, which could impede the response to selection on traits they affect (Carroll 2005). In addition, the genes implicated in the *Obp56h trans* regulatory loop may affect other phenotypes, which can

create correlations between traits and weaken the response to selection (Lande and Arnold 1983).
Notably, the expression of both *Obp56h* and *CG10514* are associated with CHC profiles in
*Drosophila* (McGraw et al. 2011; Shorter et al. 2016). CHCs serve dual functions for both
chemical communication and protection against environmental stressors (Blomquist and
Bagnères 2010). Selection on *Obp56h* expression and male mating behavior could thus be
weakened by trade-offs between mating behavior and stress response. This is consistent with the
general prediction that pleiotropy can maintain genetic variance for sexually selected traits in
natural populations (Kirkpatrick and Ryan 1991; Turelli and Barton 2004; Johnston et al. 2013;
Heinen-Kay et al. 2020). Our results therefore provide a potential example of how the regulatory
architecture underlying a sexually selected trait can create pleiotropic constraints that could
impede selection on the trait, maintaining genetic variation in spite of strong selection on the
trait.

## **ACKNOWLEDGEMENTS**

791 Rebecca Presley assisted with house fly mating experiments. This material is based upon work
792 supported by the National Science Foundation under Grant No. DEB-1845686. Any opinions,
793 findings, and conclusions or recommendations expressed in this material are those of the authors
794 and do not necessarily reflect the views of the National Science Foundation.

## REFERENCES

Abbott, J. K., A. K. Nordén, and B. Hansson. 2017. Sex chromosome evolution: historical insights and future perspectives. Proc. Biol. Sci. 284.

Adhikari, K., J. H. Son, A. H. Rensink, J. Jaweria, D. Bopp, L. W. Beukeboom, and R. P. Meisel. 2021. Temperature-dependent effects of house fly proto-Y chromosomes on gene expression.

Allen, S. L., R. Bonduriansky, C. M. Sgro, and S. F. Chenoweth. 2017. Sex-biased transcriptome divergence along a latitudinal gradient. Mol. Ecol. 26:1256–1272.

Arnqvist, G., N. Vellnow, and L. Rowe. 2014. The effect of epistasis on sexually antagonistic genetic variation. Proc. Biol. Sci. 281.

Atkinson, D. 1996. Ectotherm life-history responses to developmental temperature. Pp. 183–204 *in* A. F. B. Ian A. Johnston, ed. Animals and Temperature: Phenotypic and Evolutionary Adaptation.

Baker, R. H., and G. S. Wilkinson. 2010. Comparative genomic hybridization (CGH) reveals a neo-X chromosome and biased gene movement in stalk-eyed flies (genus *Teleopsis*). PLoS Genetics 6:e1001121.

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting linear mixed-effects models using lme4. Journal of Statistical Software 67:1–48.

Benjamini, Y., and Y. Hochberg. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. J. R. Stat. Soc. Series B Stat. Methodol. 57:289–300. [Royal Statistical Society, Wiley].

Benoit, J. B., A. Vigneron, N. A. Broderick, Y. Wu, J. S. Sun, J. R. Carlson, S. Aksoy, and B. L. Weiss. 2017. Symbiont-induced odorant binding proteins mediate insect host hematopoiesis. Elife 6.

Berson, J. D., F. Garcia-Gonzalez, and L. W. Simmons. 2019a. Experimental evidence for the role of sexual selection in the evolution of cuticular hydrocarbons in the dung beetle, *Onthophagus taurus*. J. Evol. Biol. 32:1186–1193.

Berson, J. D., M. Zuk, and L. W. Simmons. 2019b. Natural and sexual selection on cuticular

hydrocarbons: a quantitative genetic analysis. Proc. Biol. Sci. 286:20190677.

Betrán, E., K. Thornton, and M. Long. 2002. Retroposed new genes out of the X in *Drosophila*. Genome

Res. 12:1854–1859.

Bliss, C. I. 1926. Temperature characteristics for prepupal development in *Drosophila melanogaster*. J.

Gen. Physiol. 9:467–495.

Blomquist, G. J., and A.-G. Bagnères. 2010. Insect Hydrocarbons: Biology, Biochemistry, and Chemical

Ecology. Cambridge University Press.

Bray, N. L., H. Pimentel, P. Melsted, and L. Pachter. 2016. Near-optimal probabilistic RNA-seq

quantification. Nat. Biotechnol. 34:525–527.

Brooks, R., and J. A. Endler. 2001. Female guppies agree to differ: phenotypic and genetic variation in

mate-choice behavior and the consequences for sexual selection. Evolution 55:1644–1655. Wiley

Online Library.

Bryant, E. H. 1980. Geographic variation in components of mating success of the housefly, *Musca*

*domestica L.*, in the United States. The American Naturalist 116:655–669.

Carroll, S. B. 2005. Evolution at two levels: on genes and form. PLoS Biol. 3:e245.

Castel, S. E., A. Levy-Moonshine, P. Mohammadi, E. Banks, and T. Lappalainen. 2015. Tools and best

practices for data processing in allelic expression analysis. Genome Biol. 16:195.

Charlesworth, D., B. Charlesworth, and G. Marais. 2005. Steps in the evolution of heteromorphic sex

chromosomes. Heredity 95:118–128.

Chenoweth, S. F., and K. McGuigan. 2010. The genetic basis of sexually selected variation. Annu. Rev.

Ecol. Evol. Syst. 41:81–101. Annual Reviews.

Chintapalli, V. R., J. Wang, and J. A. T. Dow. 2007. Using FlyAtlas to identify better *Drosophila*

*melanogaster* models of human disease. Nat. Genet. 39:715–720.

Chippindale, A. K., and W. R. Rice. 2001. Y chromosome polymorphism is a strong determinant of male

fitness in *Drosophila melanogaster*. Proc. Natl. Acad. Sci. U. S. A. 98:5677–5682.

Chung, H., and S. B. Carroll. 2015. Wax, sex and the origin of species: Dual roles of insect cuticular

hydrocarbons in adaptation and mating. Bioessays 37:822–830.

Connallon, T. 2015. The geography of sex-specific selection, local adaptation, and sexual dimorphism.

Evolution 69:2333–2344.

Connallon, T., and A. G. Clark. 2011. The resolution of sexual antagonism by gene duplication. Genetics

187:919–937.

Connallon, T., S. Sharma, and C. Olito. 2019. Evolutionary consequences of sex-specific selection in

variable environments: four simple models reveal diverse evolutionary outcomes. Am. Nat.

193:93–105.

Delph, L. F., J. Andicoechea, J. C. Steven, C. R. Herlihy, S. V. Scarpino, and D. L. Bell. 2011.

Environment-dependent intralocus sexual conflict in a dioecious plant. New Phytol. 192:542–552.

Dobin, A., C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson, and T. R.

Gingeras. 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29:15–21.

Edgar, R. C. 2004. MUSCLE: a multiple sequence alignment method with reduced time and space

complexity. BMC Bioinformatics 5:113.

Emerson, J. J., H. Kaessmann, E. Betrán, and M. Long. 2004. Extensive gene traffic on the mammalian X

chromosome. Science 303:537–540.

Ezaz, T., K. Srikulnath, and J. A. M. Graves. 2017. Origin of amniote sex chromosomes: an ancestral

super-sex chromosome, or common requirements? J. Hered. 108:94–105.

Feldmeyer, B., M. Kozielska, and B. Kuijper. 2008. Climatic variation and the geographical distribution

of sex-determining mechanisms in the housefly. Evolutionary. evolutionary-ecology.com.

Fenster, C. B., L. F. Galloway, and L. Chao. 1997. Epistasis and its consequences for the evolution of

natural populations. Trends Ecol. Evol. 12:282–286.

Fitzpatrick, M. J. 2004. Pleiotropy and the genomic location of sexually selected genes. Am. Nat.

163:800–808.

Foster, G. G., M. J. Whitten, C. Konovalov, J. T. A. Arnold, and G. Maffi. 1981. Autosomal genetic maps

of the Australian Sheep Blowfly, *Lucilia cuprina dorsalis* R.-D. (Diptera: Calliphoridae), and

possible correlations with the linkage maps of *Musca domestica* L. and *Drosophila melanogaster*

(Mg.). Genet. Res. 37:55–69. Cambridge University Press.

Fox, J., M. Friendly, and S. Weisberg. 2013. Hypothesis tests for multivariate linear models using the car

package. The R Journal 5:39.

Franco, M. G., P. G. Rubini, and M. Vecchi. 1982. Sex-determinants and their distribution in various

populations of *Musca domestica L.* of Western Europe. Genet. Res. 40:279–293.

Furman, B. L. S., and B. J. Evans. 2016. Sequential turnovers of sex chromosomes in African clawed

frogs (*Xenopus*) suggest some genomic regions are good at sex determination. G3 6:3625–3633.

Gallach, M., and E. Betrán. 2011. Intralocus sexual conflict resolved through gene duplication. Trends

Ecol. Evol. 26:222–228.

Gibson, J. R., A. K. Chippindale, and W. R. Rice. 2002. The X chromosome is a hot spot for sexually

antagonistic fitness variation. Proceedings of the Royal Society of London B: Biological Sciences

269:499–505. The Royal Society.

Greene, E., B. E. Lyon, V. R. Muehter, L. Ratcliffe, S. J. Oliver, and P. T. Boag. 2000. Disruptive sexual

selection for plumage coloration in a passerine bird. Nature 407:1000–1003.

Hamm, R. L., J.-R. Gao, G. G.-H. Lin, and J. G. Scott. 2009. Selective advantage for III$^M$ males over Y$^M$

males in cage competition, mating competition, and pupal emergence in *Musca domestica* L.

(Diptera: Muscidae). Environ. Entomol. 38:499–504.

Hamm, R. L., R. P. Meisel, and J. G. Scott. 2015. The evolving puzzle of autosomal versus Y-linked male

determination in *Musca domestica*. G3 5:371–384.

Hamm, R. L., T. Shono, and J. G. Scott. 2005. A cline in frequency of autosomal males is not associated with insecticide resistance in house fly (Diptera: Muscidae). J. Econ. Entomol. 98:171–176.

Hedrick, P. W., M. E. Ginevan, and E. P. Ewing. 1976. Genetic polymorphism in heterogeneous environments. Annu. Rev. Ecol. Syst. 7:1–32. Annual Reviews.

Heinen-Kay, J. L., R. E. Nichols, and M. Zuk. 2020. Sexual signal loss, pleiotropy, and maintenance of a male reproductive polymorphism in crickets. Evolution 74:1002–1009.

Ingleby, F. C., J. Hunt, and D. J. Hosken. 2010. The role of genotype-by-environment interactions in sexual selection. J. Evol. Biol. 23:2031–2045.

Jia, F. Y., M. D. Greenfield, and R. D. Collins. 2000. Genetic variance of sexually selected traits in waxmoths: maintenance by genotype x environment interaction. Evolution 54:953–967.

Johnston, S. E., J. Gratten, C. Berenos, J. G. Pilkington, T. H. Clutton-Brock, J. M. Pemberton, and J. Slate. 2013. Life history trade-offs at a single locus maintain sexually selected genetic variation. Nature 502:93–95.

Jones, A. G. 2016. Theory of sexual selection. Pp. 119–127 *in* R. M. Kliman, ed. Encyclopedia of Evolutionary Biology. Academic Press, Oxford.

Kirkpatrick, M., and M. J. Ryan. 1991. The evolution of mating preferences and the paradox of the lek. Nature 350:33–38.

Kokko, H., and K. Heubel. 2008. Condition-dependence, genotype-by-environment interactions and the lek paradox. Genetica 134:55–62.

Konrad, K. D., T. J. Goralski, A. P. Mahowald, and J. L. Marsh. 1998. The gastrulation defective gene of *Drosophila melanogaster* is a member of the serine protease superfamily. Proc. Natl. Acad. Sci. U. S. A. 95:6819–6824.

Koštál, V., J. Korbelová, T. Štětina, R. Poupardin, H. Colinet, H. Zahradníčková, I. Opekarová, M. Moos,

and P. Šimek. 2016. Physiological basis for low-temperature survival and storage of quiescent larvae

of the fruit fly *Drosophila melanogaster*. Sci. Rep. 6:32346.

Kozielska, M., B. Feldmeyer, I. Pen, F. J. Weissing, and L. W. Beukeboom. 2008. Are autosomal

sex-determining factors of the housefly (*Musca domestica*) spreading north? Genet. Res.

90:157–165. Cambridge University Press.

Kumar, S., G. Stecher, M. Li, C. Knyaz, and K. Tamura. 2018. MEGA X: Molecular evolutionary

genetics analysis across computing platforms. Mol. Biol. Evol. 35:1547–1549.

Lande, R., and S. J. Arnold. 1983. The measurement of selection on correlated characters. Evolution

37:1210–1226.

Langfelder, P., and S. Horvath. 2008. WGCNA: an R package for weighted correlation network analysis.

BMC Bioinformatics 9:559.

Langfelder, P., P. S. Mischel, and S. Horvath. 2013. When is hub gene selection better than standard

meta-analysis? PLoS One 8:e61505.

Lasne, C., S. B. Hangartner, T. Connallon, and C. M. Sgrò. 2018. Cross-sex genetic correlations and the

evolution of sex-specific local adaptation: Insights from classical trait clines in *Drosophila*

*melanogaster*. Evolution 72:1317–1327. Wiley Online Library.

Leek, J. T., W. E. Johnson, H. S. Parker, A. E. Jaffe, and J. D. Storey. 2012. The sva package for

removing batch effects and other unwanted variation in high-throughput experiments. Bioinformatics

28:882–883.

Levene, H. 1953. Genetic equilibrium when more than one ecological niche is available. Am. Nat.

87:331–333. The University of Chicago Press.

Li, A., and S. Horvath. 2007. Network neighborhood analysis with the multi-node topological overlap

measure. Bioinformatics 23:222–231.

Lockey, K. H. 1988. Lipids of the insect cuticle: origin, composition and function. Comparative

939   Biochemistry and Physiology Part B: Comparative Biochemistry 89:595–645.

940 Loeb, J., and J. H. Northrop. 1917. On the influence of food and temperature upon the duration of life.

941   Journal of Biological Chemistry 32:103–121.

942 Love, M. I., W. Huber, and S. Anders. 2014. Moderated estimation of fold change and dispersion for

943   RNA-seq data with DESeq2. Genome Biol. 15:550.

944 Maere, S., K. Heymans, and M. Kuiper. 2005. BiNGO: a Cytoscape plugin to assess overrepresentation of

945   Gene Ontology categories in Biological Networks. Bioinformatics 21:3448–3449.

946 Marcillac, F., Y. Grosjean, and J.-F. Ferveur. 2005. A single mutation alters production and discrimination

947   of *Drosophila* sex pheromones. Proc. Biol. Sci. 272:303–309.

948 McGraw, E. A., Y. H. Ye, B. Foley, S. F. Chenoweth, M. Higgie, E. Hine, and M. W. Blows. 2011.

949   High-dimensional variance partitioning reveals the modular genetic basis of adaptive divergence in

950   gene expression during reproductive character displacement. Evolution 65:3126–3137.

951 McKenna, A., M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis, A. Kernytsky, K. Garimella, D.

952   Altshuler, S. Gabriel, M. Daly, and M. A. DePristo. 2010. The Genome Analysis Toolkit: a

953   MapReduce framework for analyzing next-generation DNA sequencing data. Genome Res.

954   20:1297–1303.

955 Meisel, R. P., T. Davey, J. H. Son, A. C. Gerry, T. Shono, and J. G. Scott. 2016. Is multifactorial sex

956   determination in the house fly, *Musca domestica (L.)*, stable over time? J. Hered. 107:615–625.

957 Meisel, R. P., C. A. Gonzales, and H. Luu. 2017. The house fly Y Chromosome is young and minimally

958   differentiated from its ancient X Chromosome partner. Genome Res. 27:1417–1426.

959 Meisel, R. P., M. V. Han, and M. W. Hahn. 2009. A complex suite of forces drives gene traffic from

960   *Drosophila* X chromosomes. Genome Biol. Evol. 1:176–188.

961 Meisel, R. P., and J. G. Scott. 2018. Using genomic data to study insecticide resistance in the house fly,

962   *Musca domestica*. Pestic. Biochem. Physiol. 151:76–81.

Meisel, R. P., J. G. Scott, and A. G. Clark. 2015. Transcriptome differences between alternative sex

determining genotypes in the house fly, *Musca domestica*. Genome Biol. Evol. 7:2051–2061.

Merilä, J., and B. C. Sheldon. 1999. Genetic architecture of fitness and nonfitness traits: empirical

patterns and development of ideas. Heredity 83 ( Pt 2):103–109.

Olafson, P. U., S. Aksoy, G. M. Attardo, G. Buckmeier, X. Chen, C. J. Coates, M. Davis, J. Dykema, S. J.

Emrich, M. Friedrich, C. J. Holmes, P. Ioannidis, E. N. Jansen, E. C. Jennings, D. Lawson, E. O.

Martinson, G. L. Maslen, R. P. Meisel, T. D. Murphy, D. Nayduch, D. R. Nelson, K. J. Oyen, T. J.

Raszick, J. M. C. Ribeiro, H. M. Robertson, A. J. Rosendale, T. B. Sackton, P. Saelao, S. L. Swiger,

S.-H. Sze, A. M. Tarone, D. B. Taylor, W. C. Warren, R. M. Waterhouse, M. T. Weirauch, J. H.

Werren, R. K. Wilson, E. M. Zdobnov, and J. B. Benoit. 2021. The genome of the stable fly,

*Stomoxys calcitrans*, reveals potential mechanisms underlying reproduction, host interactions, and

novel targets for pest control. BMC Biol. 19. Springer Science and Business Media LLC.

Olafson, P. U., and C. A. Saski. 2020. Chemosensory-Related Gene Family Members of the Horn Fly,

(Diptera: Muscidae), Identified by Transcriptome Analysis. Insects 11.

O'Meally, D., T. Ezaz, A. Georges, S. D. Sarre, and J. A. M. Graves. 2012. Are some chromosomes

particularly good at sex? Insights from amniotes. Chromosome Res. 20:7–19.

Otte, T., M. Hilker, and S. Geiselhardt. 2018. Phenotypic plasticity of cuticular hydrocarbon profiles in

insects. J. Chem. Ecol. 44:235–247.

Park, S. K., K. J. Mann, H. Lin, E. Starostina, A. Kolski-Andreaco, and C. W. Pikielny. 2006. A

*Drosophila* protein specific to pheromone-sensing gustatory hairs delays males' copulation attempts.

Curr. Biol. 16:1154–1159.

Ritchie, M. E., B. Phipson, D. Wu, Y. Hu, C. W. Law, W. Shi, and G. K. Smyth. 2015. limma powers

differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res.

43:e47.

Rueffler, C., T. J. M. Van Dooren, O. Leimar, and P. A. Abrams. 2006. Disruptive selection and then

what? Trends Ecol. Evol. 21:238–245.

Ruzicka, F., L. Dutoit, P. Czuppon, C. Y. Jordan, X. Li, C. Olito, A. Runemark, E. I. Svensson, H. P.

Yazdi, and T. Connallon. 2020. The search for sexually antagonistic genes: Practical insights from

studies of local adaptation and statistical genomics. Evolution Letters 57:289.

Schaeffer, S. W. 2018. Muller "elements" in *Drosophila*: How the search for the genetic basis for

speciation led to the birth of comparative genomics. Genetics 210:3–13.

Scott, J. G., P. Sridhar, and N. Liu. 1996. Adult specific expression and induction of cytochrome P450lpr

in house flies. Arch. Insect Biochem. Physiol. 31:313–323.

Scott, J. G., W. C. Warren, L. W. Beukeboom, D. Bopp, A. G. Clark, S. D. Giers, M. Hediger, A. K.

Jones, S. Kasai, C. A. Leichter, M. Li, R. P. Meisel, P. Minx, T. D. Murphy, D. R. Nelson, W. R.

Reid, F. D. Rinkevich, H. M. Robertson, T. B. Sackton, D. B. Sattelle, F. Thibaud-Nissen, C.

Tomlinson, L. van de Zande, K. K. O. Walden, R. K. Wilson, and N. Liu. 2014. Genome of the house

fly, *Musca domestica L.*, a global vector of diseases with adaptations to a septic environment.

Genome Biol. 15:466.

Seehausen, and van Alphen. 1999. Can sympatric speciation by disruptive sexual selection explain rapid

evolution of cichlid diversity in Lake Victoria? Ecol. Lett. 2:262–271.

Shannon, P., A. Markiel, O. Ozier, N. S. Baliga, J. T. Wang, D. Ramage, N. Amin, B. Schwikowski, and

T. Ideker. 2003. Cytoscape: a software environment for integrated models of biomolecular

interaction networks. Genome Res. 13:2498–2504.

Shono, T., and J. G. Scott. 2003. Spinosad resistance in the housefly, *Musca domestica*, is due to a

recessive factor on autosome 1. Pestic. Biochem. Physiol. 75:1–7.

Shorter, J. R., L. M. Dembeck, L. J. Everett, T. V. Morozova, G. H. Arya, L. Turlapati, G. E. St Armour,

C. Schal, T. F. C. Mackay, and R. R. H. Anholt. 2016. *Obp56h* modulates mating behavior in

*Drosophila melanogaster*. G3 6:3335–3342.

Son, J. H., T. Kohlbrenner, S. Heinze, L. W. Beukeboom, D. Bopp, and R. P. Meisel. 2019. Minimal effects of proto-Y chromosomes on house fly gene expression in spite of evidence that selection maintains stable polygenic sex determination. Genetics 213:313–327.

Son, J. H., and R. P. Meisel. 2021. Gene-level, but not chromosome-wide, divergence between a very young house fly proto-Y chromosome and its homologous proto-X chromosome. Mol. Biol. Evol. 38:606–618.

Sun, J. S., S. Xiao, and J. R. Carlson. 2018. The diverse small proteins called odorant-binding proteins. Open Biol. 8:180208.

Takehana, Y., M. Matsuda, T. Myosho, M. L. Suster, K. Kawakami, T. Shin-I, Y. Kohara, Y. Kuroki, A. Toyoda, A. Fujiyama, S. Hamaguchi, M. Sakaizumi, and K. Naruse. 2014. Co-option of *Sox3* as the male-determining factor on the Y chromosome in the fish *Oryzias dancena*. Nat. Commun. 5:4157.

Thomas, M. L., and L. W. Simmons. 2009. Sexual selection on cuticular hydrocarbons in the Australian field cricket, *Teleogryllus oceanicus*.

Tomita, T., and Y. Wada. 1989. Multifactorial sex determination in natural populations of the housefly (*Musca domestica*) in Japan. The Japanese Journal of Genetics 64:373–382. The Genetics Society of Japan.

Tomkins, J. L., J. Radwan, J. S. Kotiaho, and T. Tregenza. 2004. Genic capture and resolving the lek paradox. Trends Ecol. Evol. 19:323–328.

Turelli, M., and N. H. Barton. 2004. Polygenic variation maintained by balancing selection: pleiotropy, sex-dependent allelic effects and G x E interactions. Genetics 166:1053–1079.

Valanne, S., J.-H. Wang, and M. Rämet. 2011. The *Drosophila Toll* signaling pathway. The Journal of Immunology 186:649–656.

van Doorn, G. S., and M. Kirkpatrick. 2007. Turnover of sex chromosomes induced by sexual conflict.

Nature 449:909–912.

van Doorn, G. S., and F. J. Weissing. 2002. Ecological versus sexual selection models of sympatric

speciation: a synthesis. Selection 2:17–40. Akadémiai Kiadó.

Van Kuren, N. W., and M. Long. 2018. Gene duplicates resolving sexual conflict rapidly evolved essential

gametogenesis functions. Nature Ecology & Evolution 2:705–712.

Weller, G. L., and G. G. Foster. 1993. Genetic maps of the sheep blowfly *Lucilia cuprina*: linkage-group

correlations with other dipteran genera. Genome 36:495–506.

Wiegmann, B. M., M. D. Trautwein, I. S. Winkler, N. B. Barr, J.-W. Kim, C. Lambkin, M. A. Bertone, B.

K. Cassel, K. M. Bayless, A. M. Heimberg, B. M. Wheeler, K. J. Peterson, T. Pape, B. J. Sinclair, J.

H. Skevington, V. Blagoderov, J. Caravas, S. N. Kutty, U. Schmidt-Ott, G. E. Kampmeier, F. C.

Thompson, D. A. Grimaldi, A. T. Beckenbach, G. W. Courtney, M. Friedrich, R. Meier, and D. K.

Yeates. 2011. Episodic radiations in the fly tree of life. Proc. Natl. Acad. Sci. U. S. A.

108:5690–5695.

Zhou, J.-J. 2010. Odorant-binding proteins in insects. Vitamins & Hormones 241–272.