Dynamic Phosphorylation of RGS Provides Spatial Regulation of G-alpha And Promotes Completion of Cytokinesis

William C. Simke[1], Andrew J. Hart[1], Cory P. Johnson[2], Sari Mayhue[1], P. Lucas Craig[1], Savannah Sojka[1], and Joshua B. Kelley[1,2]

[1] Department of Molecular and Biomedical Sciences, University of Maine

[2] Graduate School of Biomedical Science and Engineering, University of Maine

Running Title: Phosphorylated RGS and Cytokinesis

Corresponding Author: Joshua Kelley, joshua.b.kelley@maine.edu

Abstract

Yeast use a G-protein coupled receptor (GPCR) signaling pathway to detect mating pheromone, arrest in G1, and direct polarized growth towards the potential mating partner. The primary negative regulator of this pathway is the regulator of G-protein signaling (RGS), Sst2, which induces Gα GTPase activity and subsequent inactivation of all downstream signaling. MAPK phosphorylates the RGS in response to pheromone, but the role of this modification is unknown. We set out to examine the role of RGS phosphorylation during the pheromone response. We found that phosphorylation of the RGS peaks early in the pheromone response and diminishes RGS localization to the polarization site and focuses Gα/MAPK complexes there. At later time points, RGS is predominantly unphosphorylated, which promotes RGS localization to the polar cap and broadens the distribution of Gα/MAPK complexes relative to the Cdc42 polarity machinery. Surprisingly, we found that phosphorylation of the RGS is required for the completion of cytokinesis prior to pheromone induced growth. The completion of cytokinesis in the presence of pheromone is promoted by the formin Bnr1 and the kelch-repeat protein, Kel1, both proteins previously found to interact with the RGS.

Introduction

When cells receive signals to perform competing processes, they must integrate those two signals into a singular outcome. The budding yeast, *Saccharoymyces cerevisiae*, use a G-protein coupled receptor (GPCR) to detect and grow toward potential mating partners (Alvaro and Thorner, 2016; Arkowitz, 2009; Wang and Dohlman, 2004). However, the cells must complete mitosis and arrest in G1 prior to mating projection morphogenesis (shmoo formation) (Peter et al., 1993). This requires that the cell prioritize the signaling that will drive mitosis and cytokinesis to completion, and only after arrest in G1, allow the pheromone signaling pathway to commandeer the Cdc42 polarity machinery that has shared uses in both mitosis and pheromone-induced morphogenesis (Chiou et al., 2017; Park and Bi, 2007). While the mechanism by which G1 arrest occurs is understood, the mechanism responsible for suppression of receptor driven polarization is unknown.

The pheromone response can be thought of as a response made up primarily of two G-proteins: the receptor-activated large G-protein consisting of the Gα and Gβγ conveying information about where the pheromone receptor is active, and the small G-protein Cdc42 controlling MAPK signaling and actin cytoskeleton polarization. The GPCR Ste2 activates the large G-protein, causing the Gα and Gβγ subunits to dissociate. Gβγ initiates Cdc42 mediated polarization of the actin cytoskeleton to form a mating projection. Gβγ also promotes the activation of the two yeast ERK homologs, Fus3 and Kss1(Wang and Dohlman, 2004). Of these two MAP kinases, Fus3 has pheromone specific roles: it is required for gradient tracking, arrest of the cell cycle in G1, and is scaffolded to the cell periphery by active Gα to regulate actin polymerization (Elion et al., 1993; Hao et al., 2008; Matheos et al., 2004; Metodiev et al., 2002; Pope et al., 2014). For this study we will only be concerned with Fus3 functions, and so all references to MAPK refer to Fus3.

The primary negative regulator of the pheromone pathway is the Regulator of G-protein Signaling (RGS), Sst2, (Chasse et al., 2006) which serves as the GTPase activating protein (GAP) for the Gα subunit (Apanovitch et al., 1998). Upon hydrolyzing GTP, the Gα binds to Gβγ, turning off the pathway. RGS function is required for pathway inactivation and for the ability of the cell to track the pheromone gradient (Dohlman et al., 1995; Segall, 1993). GPCR signaling pathways play a central role in human disease, indicating that understanding the mechanisms by which RGS functions has the potential to inform understanding of human signaling pathways relevant for drug development (Bar-Shavit et al., 2016; Hauser et al., 2017; Lappano and Maggiolini, 2012).

The RGS Sst2 has characterized interactions with the Gα (Gpa1) subunit, the pheromone receptor (Ste2), and the MAPK (Fus3) (Ballon et al., 2006; DiBello et al., 1998; Garrison et al., 1999). RGS serves as a GTPase activating protein for Gα, a function that is enhanced by its binding to the cytoplasmic tail of the receptor (Apanovitch et al., 1998; Ballon et al., 2006; Dixit et al., 2014). The MAPK Fus3 phosphorylates the RGS at serine 539 in a pheromone dependent manner, but this does not impact the sensitivity of the pathway or downstream MAPK or transcriptional outputs (Garrison et al., 1999).

In less well-characterized interactions, the RGS has been found in a yeast two-hybrid screen to interact with the formin Bnr1 and the formin regulatory protein Kel1 (Burchett et al., 2002; Yu et al., 2008). Yeast contain two formins, Bni1 and Bnr1, which drive polymerization of actin in response to Rho family signaling (Breitsprecher and Goode, 2013). Bni1 is activated by Cdc42, and is known to be involved in the pheromone response as its activity is regulated by MAPK bound to active Gα (Evangelista et al., 1997; Matheos et al., 2004; Metodiev et al., 2002). Bnr1 is activated by Rho3 and Rho4, and is involved in cytokinesis, but has no clear role in the pheromone response (Imamura et al., 1997). The formins are partially redundant, as either is sufficient for viability, but the loss of both is lethal (Imamura et al., 1997). Kel1 is a kelch-repeat containing protein which has been shown to act as a negative regulator of Bnr1, is required for efficient mating, and plays a role in the mitotic exit network (Gould et al., 2014; Hofken and Schiebel, 2002; Smith and Rose, 2016). Given the clear roles of Bnr1 and Kel1 in cytokinesis with little known about their roles in the pheromone response, the potential for RGS interactions with these proteins has not been pursued.

In an unsynchronized, logarithmically growing population of yeast, most cells will be between the beginning of S phase and cytokinesis. Because the Cdc42 polarity machinery is shared between both the mitotic machinery and the pheromone machinery, cells must arrest in G1 before responding to pheromone (Bi and Park, 2012; Chiou et al., 2017). This is accomplished by MAPK phosphorylating Far1, which inhibits Cyclin Dependent Kinase (Butty et al., 1998; Peter et al., 1993; Pope et al., 2014). Thus, when cells receive the pheromone signal, they complete their current round of mitosis and cytokinesis, and only then does receptor driven signaling reorient the polarity machinery for pheromone-induced morphogenesis of the mating projection (shmoo) (Chiou et al., 2017).

Here, we set out to determine the role of MAPK phosphorylation of RGS in response to pheromone. We found that RGS phosphorylation is dynamic, with high phosphorylation early in the response, followed by decreased phosphorylation later. Phosphorylation of the RGS decreases its localization to the polar cap, and

reduces the distance between peak active Cdc42 and peak active Gα. Surprisingly, RGS phosphorylation early in the pheromone response is required to ensure that cytokinesis completes prior to the beginning of pheromone induced polarity. We find that cytokinetic completion in the presence of pheromone is also dependent upon the kelch-repeat protein Kel1, a component of the mitotic exit network.

Results

Cells can track a gradient of pheromone independent of phosphorylation at Ser 539.

The ability to track a gradient of pheromone is dependent upon RGS, specifically its GTPase activating protein (GAP) function (Dixit et al., 2014; Segall, 1993). MAPK phosphorylates RGS on serine 539 in response to pheromone, though the nature of this phosphorylation event is poorly understood. While previous studies found that GAP activity was not affected by S539 phosphorylation, it is possible that phosphorylation might change Sst2 function in time or space in a way that affects gradient tracking (Garrison et al., 1999). We tested this hypothesis by using strains expressing an unphosphorylatable RGS mutant ($sst2^{S539A}$, denoted p̶RGS) and phospho-mimetic RGS mutant ($sst2^{S539D}$, denoted p*RGS) from the endogenous SST2 genomic locus, each fused to GFP. As a marker of the polar cap, we used the Cdc42-GTP binding protein Bem1 fused to mRuby2 (Kelley

Figure 1

Figure 1. Phosphorylation state of the RGS does not stop gradient tracking. A) Representative live-cell images of WT, unphosphorylatable (p̶RGS), and phosphomimetic (p*RGS) RGS expressing the polar cap marker (Bem1-mRuby2) and the RGS (Sst2-EGFP) tracking a 0-150nM gradient of pheromone, with pheromone increasing to the right. B) Quantification of gradient tracking cells measured by the cosine of orientation for WT (n =95), p̶RGS ($sst2^{S539A}$, n =39), and p*RGS ($sst2^{S539D}$, n =45) from three experiments. Scale bars represent 5 μm.

Figure 2. Localization of the RGS is dependent on phosphorylation state. A) Confocal images of WT, unphosphorylatable (p̶RGS), and phosphomimetic (p*RGS) RGS fused to EGFP in saturating pheromone. B) Quantification of the average RGS spatial distribution normalized to the polar cap marker (Bem1-mRuby2) in saturating pheromone over a 12hr time-course in a microfluidic gradient chamber, imaged by epifluorescence microscopy. Lines are derived from averaging from 180 min. onward. Bottom graphs display statistical analysis using one-way ANOVA followed by Tukey's HSD, with statistically (p < 0.05) significant differences in localization noted by bars. Data is derived from n = 89 cells (WT), n = 88 p̶RGS (unphosporylatable), and n = 139 p*RGS (phosphomimetic) cells per strain, with 29 time points per cell. C) 3-D kymographs of the spatial distribution of the RGS over 12hrs for WT, p̶RGS, and p*RGS with 37 time points per cell.

et al., 2015).   These strains were examined in a microfluidic gradient chamber by live cell microscopy (Suzuki et al., 2021). We exposed these cells to a 0-150 nM gradient of pheromone and measured their ability to grow toward the source of pheromone (Figure 1). Both phospho-mutant strains were able to track a gradient of pheromone(Figure 1B). Thus, feedback phosphorylation of RGS has no effect on gradient tracking.

RGS localization is regulated by its phosphorylation state

The pheromone driven MAPK Fus3 is central to mating signaling, driving cell cycle arrest, transcriptional changes, and promoting actin cytoskeleton changes (Alvaro and Thorner, 2016; Chiou et al., 2017; Wang and Dohlman, 2004). (Garrison et al., 1999) We hypothesized that phosphorylation of the RGS may regulate its spatial distribution. To test this, we again used the p*RGS (*sst2^{S539D}*) and p̄RGS (*sst2^{S539A}*) mutants tagged with EGFP, as well as Bem1-mRuby2 for polar cap localization. Cells were exposed to saturating pheromone (300 nM) in a microfluidic chamber (Suzuki et al., 2021) for 12 hours and imaged by timelapse epifluorescence microscopy. The signal of RGS along the periphery of the cell is spatially normalized to the center of the polar cap (peak Bem1 signal) and averaged to generate a distribution of the protein during the pheromone response (Kelley et al., 2015). We found that phosphorylation of the RGS diminishes its localization to the polar cap (Figure 2A). The inability to phosphorylate RGS leads to a small but statistically significant increase in association with the polar cap, suggesting that much of the RGS measured in WT cells may be in the unphosphorylated form (Figure 2B). We then examined changes in RGS distribution using an averaged 3D-kymograph (Figure 2C). As expected, RGS fluorescence increases throughout the time course, which is due to persistent pheromone-induced production of RGS (Dohlman et al., 1996).   As the time in pheromone passes 400 minutes, there is broadening of the lower intensity area surrounding the polar cap in both WT and p̄RGS cells, while the p*RGS mutant has a very consistently low polar cap association through time. We conclude that the phospho-form of the RGS is unable to interact with a normal binding partner at the center of the polar cap.

RGS phosphorylation alters the distance between the Polar Cap and the Gα-MAPK complex

While much of the polarity signaling in the pheromone response comes through the Gβγ branch of the pathway, GTP bound Gα is known to recruit active MAPK (Metodiev et al., 2002). The Gα-MAPK complex activates the formin Bni1 and promotes gradient tracking (Errede et al., 2015; Matheos et al., 2004; Metodiev et al., 2002). We hypothesized that the phospho-dependent changes in RGS localization would lead to cor-

Figure 3. The MAPK Fus3 can serve as an indicator of the localization of active Gα. A) MAPK (Fus3-EGFP) localization in WT cells and two Gα mutants, the hyperactive Gα (*gpa1^{G302S}*) and a mutant that cannot bind MAPK (*gpa1^{E21E22}*), show differential recruitment of MAPK to the periphery of the cell under treatment with 300 nM pheromone. B) Quantification of the amount of MAPK on the periphery of the cell spatially normalized to the center of the polar cap as in Figure 1. This analysis is based on the average projection of the stack of images, as that more accurately represents the amount of protein present on the periphery. Shaded areas represent 95% confidence intervals. C) Average distributions of MAPK over time (derived from maximum intensity projections, but scaled to mean intensity). MAPK association with the site of polarity is largely dependent upon the ability of the MAPK to bind to the Gα, as MAPK levels at the polar cap are lower in cells expressing a mutant Gα that cannot bind MAPK (*gpa1^{E21E22}*). In cells with hyperactive Gα (*gpa1^{G302S}*), MAPK association with the periphery of the cell continues to climb throughout the pheromone response and is broadly distributed, as would be expected in when the Gα can be turned on but cannot be turned off. Data is derived from n = 83 (WT), n = 157 (*gpa1^{G302S}*), and n = 86 (*gpa1^{E21E22}*), with 29 time points per cell for B and 37 per cell for C.

responding changes in the localization of active Gα. While Gα is localized across the membrane (Wang et al., 2005), its localization does not indicate whether Gα is activated. However, we hypothesized that we could monitor the localization of the MAPK (Fus3) on the periphery to indicate the location of active Gα.

To test the use of MAPK as a maker for active Gα, we examined a GFP tagged MAPK (Fus3-GFP) in wild type cells, and two mutants with opposing effects: a Gα mutant that does not bind MAPK, $gpa1^{E21\,E22}$ (Metodiev et al., 2002); and a Gα mutant that is hyperactive because it no longer interacts with RGS, $gpa1^{G302S}$ (Figure 3A) (DiBello et al., 1998). We assessed MAPK levels along the periphery of the cell in these strains using the same techniques as above, with the exception that we performed an average projection of un-deconvolved Z-stacks to represent the amount of detected signal more accurately (whereas max intensity projections emphasize shape). We found that hyperactive Gα increased the amount of MAPK at the periphery, and that the Gα MAPK-binding mutant decreased MAPK localization to the periphery (Figure 3B, C). We believe the distribution of signal in the Gα MAPK-binding mutant ($gpa1^{E21E22}$) likely represents the profile of the remaining MAPK binding partners at the polar cap. Fus3 binds to the MAPK scaffold Ste5, and the polar cap is populated with many MAPK substrates (Elion et al., 1993; Wang and Dohlman, 2004). We conclude that MAPK localization can serve as a readout for localization of active Gα.

Having determined that MAPK localization can serve as a proxy for the localization of active Gα and that RGS localization is altered by phosphorylation, we next examined whether RGS phosphorylation affects the distribution of the Gα/MAPK complex. We generated strains expressing fluorescent-protein fusions of a polar cap marker (Bem1-mRuby2) and MAPK (Fus3-EGFP) in the presence of the phospho-mimetic and un-phosphorylatable RGS mutants. The p̄RGS mutant ($sst2^{S539A}$) resulted in a very similar MAPK localization to WT, with some broadening below the peak (Figure 4A/B). The p*RGS ($sst2^{S539D}$) however, induced a large increase in polar cap association (Figure 4A/B).

A potential explanation for this increase in average MAPK signal at the polar cap would be a change in the absolute distribution of Gα/MAPK at individual times, or alternatively, the same distribution has a different relationship with the polar cap. To assess the RGS induced changes in MAPK localization, we examined the distribution of MAPK (Fus3—GFP) spatially normalized to itself (Figure 5A). We see a statistically significant, but small, difference in the distribution of MAPK in both phospho-mutants when compared to WT RGS, so the change in MAPK distribution itself cannot account for the drastic changes we see in the MAPK localization relative to the polar cap seen in Figure 4. We next examined the distribution of the maximal MAPK intensity

Figure 4. RGS Phosphorylation increase Gα/MAPK complex levels at the center of the polar cap. A) Profiles of MAPK (Fus3-GFP) in cells expressing the unphosphorylatable RGS (*sst2^S539A*) or the phosphomimetic RGS (*sst2^S539D*) aligned to the center of the polar cap (Bem1) as described in Figure 2. The phosphomimetic RGS causes a broader distribution of MAPK at the center of the polar cap. Shaded areas represent 95% confidence intervals. Statistical analysis is shown in the graphs below. B) Changes in RGS phosphomutant distribution over time. Data is derived from n = 139 (p̶RGS) and n = 129 (p*RGS) cells and 29 time points per cell in C, and 37 time points per cell in B.

relative to maximal polar cap intensity (Figure 5B). This is effectively measuring the offset along the membrane between active Cdc42 and active Gα. We find that 50% of maxima for WT and p̶RGS fall within ~ 2.8 µm of the polar cap peak, while 50% of MAPK maxima in the p*RGS fall within ~1.4 µm of the polar cap peak. Thus, unphosphorylated RGS appears to drive a greater distance between active Gα and active Cdc42.

RGS phosphorylation peaks early in the pheromone response and diminishes at later time points

Previous characterization of the phosphorylation of the RGS, Sst2, at serine 539 showed Fus3-dependent phosphorylation at one-hour of pheromone treatment (Garrison et al., 1999). By contrast, we examine our cells in microfluidics devices for 12 hours, and we specifically calculate the average distribution of the RGS from 180 minutes onward (approximately the time that the RGS levels reach a new, pheromone-induced steady state, Figure 2C). Our analysis of RGS localization showed that the unphosphorylatable p̶RGS (sst2^S539A) was similar to WT over the 12-hour time course (Figure 2C). Furthermore, the offset of Gα/MAPK from

A)

Figure 5. RGS induced changes in Gα/ MAPK distribution. A) Distribution of MAPK (Fus3-GFP) from Figure 4, spatially normalized to peak MAPK rather than to the polar cap (Bem1). Both RGS phosphomutants result in a mild, but statistically significant sharpening of the MAPK distribution, but the magnitudes of the changes are insufficient to explain the changes seen in Figure 4. Shaded areas represent 95% confidence intervals. Statistical analysis is shown in graphs below. B) We compare the distance between the maxima of the Bem1 and MAPK, effectively the offset between peak Cdc42 activation and peak Gα activation. Graphed is the cumulative sum of MAPK maximum intensities versus distance from the polar cap. Vertical lines show the distance where 50% of maxima have appeared. Statistical significance was evaluated by pairwise Kolomogorov-Smirnov tests. Each pairwise comparison is significant with p-values less than 0.005.

B)

the polar cap was also similar between the two strains. These observations suggest that unphosphorylated

RGS may be the dominant species at later times during the pheromone response. We hypothesized that Sst2

phosphorylation at S539 may be dynamic: peaking earlier in the response and decreasing at later times. To test

the dynamics of phosphorylation, we developed a polyclonal antibody to detect Sst2 phosphorylated on serine

539, LHPH**S**PLSEC, where the underlined serine is phosphorylated. Western blotting of Sst2-GFP versus un-

tagged Sst2 shows a GFP-dependent size shift in the detected band, indicating that the antibody is specific for

Sst2 (Supplemental Figure 1). Western blotting is done in the presence of excess unphosphorylated peptide to ensure that it is specific for the phospho-epitope.

We next tested whether RGS phosphorylation decreased at later times in the pheromone response. We treated two different strains, *Sst2-GFP* and *bar1Δ* with pheromone and took samples every hour for four hours. A complication of examining a potential decrease in pheromone-driven signaling is the desensitizing role of the protease Bar1, which degrades pheromone (Ciejek and Thorner, 1979). By comparing these two strains we can verify that the antibody is detecting Sst2, as the detected band will show a size shift due to the GFP fusion, and we are able to exclude Bar1-mediated degradation of pheromone as driving the decrease in phosphorylation over time. We find that phosphorylation of the RGS peaks between 1 and 2 hours consistent with the literature (Garrison et al., 1999), but phosphorylation levels decrease to lower levels by four hours post pheromone treatment (Figure 6A, B). This decrease is independent of Bar1 activity (Figure 6, Supplemental Figure S1). Thus, phosphorylation likely plays a role early in the pheromone response. Given that 90 minutes after pheromone treatment is roughly the time it would take for the entire population of yeast to complete mitosis and arrest in G1, we hypothesized that the phosphorylation of S539 might have a role in the completion of mitosis.

Figure 6. RGS phosphorylation peaks 1hr in the pheromone response. A) Western blotting of phospho-RGS-GFP responding to saturating pheromone over four hours. G6PDH was probed as a loading control. B) Quantification of Western blotting shown in A, normalized to G6PDH levels. Error bars represent standard error of the mean, n=3.

Phosphorylation of RGS promotes coordination of cytokinesis with the pheromone response.

Cells which have already left G1 must complete mitosis and cytokinesis prior to polarizing and forming a mating projection in response to pheromone. We found that some mother-daughter pairs in our unphosphorylatable pRGS mutant (*sst2^S539A^*) formed mating projections before they had finished cytokinesis (Figure 7A). The frequency of the event was low and we never observed these defects in wild type cells or cells expressing the phosphomimetic mutant during microfluidics experiments. Previous studies found interactions between the RGS and two proteins involved in cytokinesis, Bnr1 and Kel1 (Burchett et al., 2002; Yu et al., 2008). Both of these proteins play a role in mitosis, Bnr1 through the regulation of actin polymerization at the mitotic septin ring and Kel1 through promoting the mitotic exit network (MEN) (Buttery et al., 2007; Gao et al., 2010; Hotz and Barral, 2014; Pruyne et al., 2004; Seshan et al., 2002). Additionally, Kel1 serves as a negative regulator of Bnr1, and may impact cytokinesis through that role as well (Gould et al., 2014). Therefore, we hypothesized the cytokinetic defect may be mediated through interactions with these proteins.

 We examined *bnr1Δ* and *kel1Δ* cells responding to saturating pheromone and found that *kel1Δ* cells also occasionally fail to complete cytokinesis prior to responding to pheromone (Figure 7A, negative data for *bnr1Δ* not shown). These were both rare events and our microfluidic experiments do not contain large numbers of yeast, so we grew wild type, pRGS, and *kel1Δ* yeast in culture, treated with saturating pheromone, and counted the frequency of cells with failed cytokinesis, finding that both mutants lead to rates of failed cytokinesis of ~3-4% (Figure 7B, minimum of 1350 cells per strain). From this, we conclude that both Sst2 and Kel1 are both involved in a mechanism that ensures cytokinesis finishes prior to the pheromone response.

To test the role of phosphorylation of the RGS in promoting cytokinesis, we took advantage of hydroxyurea (HU) to damage DNA and cause stalled cytokinesis (Amaral et al., 2016), followed by pheromone exposure, while maintaining the hydroxyurea stress (Figure 7C). HU causes DNA damage to activate the NoCut pathway, an aurora kinase-dependent checkpoint that stalls cytokinesis if chromatin has not cleared the midbody (Amaral et al., 2016; Norden et al., 2006). We reasoned that any process that delays the polarity machinery from being repurposed by pheromone receptor-driven signaling may exacerbate the defects that spontaneously arise in the unphosphorylatable pRGS or *kel1Δ* mutants. Under these conditions, stalled cytokinesis would be common, enhancing our ability to differentiate the behaviors of cells with altered Sst2 and Kel1 signaling. We pretreated with 100mM HU for two hours and then examined cells four hours post pheromone treatment, while maintaining HU. We found that WT cells frequently stalled cytokinesis with two round cells joined at

A)

DIC | Con A & Calcofluor

ₚRGS (*sst2^{S539A}*)

kel1Δ

5 µm

B) Frequency of Cytokinesis Failure during Pheromone Response

% Cytokinesis Defect

WT | ₚRGS | *kel1Δ*

C) Induce Cytokinetic Defects and Challenge with Pheromone

+ Hydroxyurea → Cytokinesis Arrest → + Pheromone Hydroxyurea → ?

D) Normal Response

DIC | DIC Std. Dev. Proj. | Calcofluor | Characteristics

WT

Single Cell — Unpolarized

Single Cell — Mating Projection

Pair of Cells Both have Mating Projection

Mother/Daughter Pair — No Mating Projection

E) Normal Response to HU & Pheromone

WT | ₚRGS *sst2^{S539A}* | p*RGS *sst2^{S539D}* | *kel1Δ* | *bnr1Δ* | *bni1Δ*

F) Asymmetric Hyperpolarized Growth

kel1Δ

Mother/Daughter Pair — One Hypertrophic Mating Projection

ₚRGS *sst2^{S539A}*

Mother/Daughter Pair — One Hypertrophic Mating Projection

G) Asymmetric Hyperpolarized Growth

WT | ₚRGS *sst2^{S539A}* | p*RGS *sst2^{S539D}* | *kel1Δ* | *bnr1Δ* | *bni1Δ*

Figure 7. Phosphorylated Sst2 and the Kelch-repeat protein Kel1 promote completion of cytokinesis prior to pheromone induced polarization. A) Images of p̄RGS mutant and *kel1Δ* which have failed to complete cytokinesis prior to pheromone induce polarized growth. Cell walls were stained with Calcofluor White and Concanavalin-A 647 to verify the open bud neck. B) Wild type and mutant p̄RGS and *kel1Δ* strains were exposed to pheromone in culture for 90 minutes, fixed, and then failed cytokinetic events were counted. n = 1412 (WT), 1350 (p̄RGS), and 1396 (*kel1Δ*) from two separate experiments. C) In order to drive cytokinetic defects, we pretreated cells with 100mM hydroxyurea, followed by treatment with both hydroxyurea and pheromone to investigate the role of the indicated proteins in delay of pheromone induced polarity until completed mitosis. D) Images of normal phenotypes in response to HU + pheromone. Shown are a single focal plane of DIC, a standard deviation projection of a stack of DIC images to better show the state of the bud neck, and cell wall staining with Calcofluor White. We considered a normal response to hydroxyurea and pheromone to be one of the following: 1) Completion of cytokinesis but arrest as a circular cell, in the event that stress signaling is suppressing the pheromone response (a minority of cells). 2) A lone cell responding to pheromone. 3) Completion of cytokinesis (if the cells had resolved their DNA damage), followed by pheromone induced morphogenesis. These cells may be still associated, but show signs of completed cytokinesis 4) Arrest of cytokinesis yielding a mother daughter pair with no polarized growth. E) Plots of the frequency of normal response to hydroxyurea and pheromone in the indicated strains. Error bars represent 95% confidence intervals. For each strain, n > 640 cells across 3 experiments. F) Examples of the asymmetric hyperpolarized growth phenotype. G) Plots of the frequency of asymmetric hyperpolarized growth in response to hydroxyurea and pheromone.

the bud neck. We scored phenotypes as a normal response if the cells showed evidence of completing cytokinesis prior to undergoing polarized growth, or if they arrested as a mother daughter pair with no evidence of polarized growth (Figure 7D/E). We also found cells that had completed cytokinesis and began mating projection formation. In cells with the unphosphorylatable p̄RGS, we found more cells that had both failed cytokinesis and began polarized growth in one or both the mother and daughter cells. A particularly striking phenotype involves one cell remaining round, while the other shows hyper-polarized growth, which we refer to as asymmetric hyperpolarized growth (Figure 7F). In contrast, cells expressing the phospho-mimetic p*RGS mutant (sst2^{S539D}) suppress polarized growth in the presence of HU-induced DNA damage. When we examine *kel1Δ* cells under these conditions, we find that the asymmetric hyperpolarized growth is a dominant phenotype (Figure 7G). Thus, unphosphorylatable RGS partially phenocopies the loss of Kel1 function. This suggests that phosphorylation of RGS promotes a Kel1 dependent mechanism to prevent the mating machinery from commandeering the polarity machinery prior to the completion of cytokinesis. The asymmetric hyperpolarized growth we see in this experiment is consistent with previously published defects seen with perturbation of the Kel1 binding partner, Lte1 (Geymonat et al., 2010) and suggests a defect in Kel1/Lte1 signaling is brought about by unphosphorylated RGS.

Yeast have two formins: Bni1 is associated with the polar cap and is activated by Cdc42 and the Gα/MAPK complex. Bnr1 is associated with mitotic septin structures and has no known role in the pheromone response but is known to be negatively regulated by Kel1 (Gould et al., 2014). Given the central role that formins play in both mitosis and the pheromone response, we hypothesized that the formins may be playing a role in

the coordination of cytokinesis and the beginning of pheromone induced polarized growth. We performed the same experiment as above, inducing cytokinetic defects with hydroxyurea followed by pheromone treatment, and assessed the ability of cells lacking either Bni1 or Bnr1 to prevent pheromone induced polarization prior to the completion of cytokinesis. Deletion of Bni1 largely stopped polarization of cells prior to completion of cytokinesis, and completely abrogated the asymmetric hyper-elongated phenotype (Figure 7E, G). Deletion of Bnr1 resulted in increased polarization prior to the completion of cytokinesis and increased levels of asymmetric hyper-polarized growth (Figure 7E, G). The asymmetric hyper-elongated growth phenotype is clearly dependent upon Bni1 and inhibited by Bnr1. Thus, coordination of pheromone induced polarity with the completion of cytokinesis is promoted by Bnr1 function and antagonized by Bni1.

Discussion

Here we set out to determine the role of feedback phosphorylation of the RGS, Sst2. We found that the phosphorylation is dynamic through the pheromone response, reaching a maximal level an hour into the response (Figure 6). We found that phosphorylation of the RGS alters the localization of the RGS relative to the polar cap during the pheromone response and leads to a broadened distribution of the Gα/MAPK complex (Gpa1/Fus3) (Figure 4, 5). Strikingly, cells unable to phosphorylate the RGS sometimes began polarized growth in response to pheromone without waiting for the completion of cytokinesis. By inducing cytokinetic defects with hydroxyurea, we were able to determine that phosphorylation of the RGS enhances the ability of the cell to stall cytokinesis without initiating pheromone induce polarity, thereby correctly integrating both internal stress response and an external morphogenesis response (Figure 7). This coordination appears to use the kelch repeat protein Kel1, and the formin Bnr1.

The role of RGS phosphorylation

The RGS Sst2 is the primary negative regulator of the pheromone response, a role dominated by its GTPase activating function to turn off the Gα subunit, Gpa1 (Apanovitch et al., 1998; Chasse et al., 2006). The pheromone-induced feedback phosphorylation of the RGS by the MAPK Fus3 is known to have no effect on the pathway sensitivity (Garrison et al., 1999). These data lead us to pose the question; why is Sst2 phosphorylated? We conclude that the unphosphorylated RGS is important for normal spatial regulation of Gα signaling, but impedes cytokinesis and so the RGS is phosphorylated by MAPK early in the pheromone response to suppress polarity until the completion of cytokinesis (Figure 8).

Figure 8. Proposed Role of RGS phosphorylation. In cells that have detected pheromone during mitosis and prior to cytokinesis, phospho-RGS and Kel1 inhibit the recruitment of the polarity machinery by active pheromone receptor. This may be because inactive RGS inhibits Kel1 function. After cytokinesis while cells are generating a mating projection or tracking a gradient of pheromone, unphosphorylated RGS promotes a greater distance between active Cdc42 and active Gα, potentially enhancing polar cap wandering.

An obvious question arises from this investigation: Does the RGS plays a role in cytokinesis in the absence of pheromone? There are multiple lines of evidence to suggest that RGS has no role in normal cytokinesis. First, In previous studies on cells lacking the RGS, we have not observed any cytokinetic defects (Kelley et al., 2015). Secondly, baseline Sst2 levels are an order of magnitude higher in haploids than in diploids (de Godoy et al., 2008). If the RGS played an important role in cytokinesis in the absence of pheromone, then haploid and diploid cells would need different mechanisms for regulating cytokinesis, an unlikely scenario.

Our data is consistent with unphosphorylated RGS inhibiting a subset of Kel1 function, as the unphosphorylatable pRGS phenocopies the spontaneous failure to complete cytokinesis before mating projection formation that we see in cells lacking Kel1 (Figure 7). We would expect this inhibition of Kel1 to involve stochiometric binding (directly or through an intermediary) and so in the absence of pheromone, where RGS levels are low (Figure 2C and (Dohlman et al., 1996)) there would be very little impact of unphosphorylated RGS on Kel1 activity. We presume that feedback phosphorylation is most important for cells that have just committed to going through the cell cycle when the mating signal starts, as they will have the most accumulation of RGS by the time they reach cytokinesis. In this scenario, a large amount of unphosphorylated RGS present would be able to inhibit Kel1. Phosphorylation of the RGS, however, would prevent its inhibition of Kel1. In the absence of pheromone, the low levels of unphosphorylated RGS would lead to a small impact on Kel1 activity. We conclude that RGS is phosphorylated early in the response to allow normal Kel1 function during the

completion of the cell cycle, and dephosporylated at later times to modulate the location of active Gα, likely through modulation of Kel1 function (Figure 8).

Coordination of the end of cytokinesis with the beginning of receptor mediated morphogenesis

In an unsynchronized population, cells will be evenly distributed through the 90-minute yeast cell cycle. Upon stimulation with pheromone, receptor signaling will immediately begin with subsequent MAPK activation, and downstream phosphorylation of the protein Far1 (Arkowitz, 2009). Far1 serves two purposes in the pheromone response: (1) to inhibit cyclin dependent kinase activity, leading to arrest in G1 (Pope et al., 2014), and (2) to couple the Cdc42 GEF, Cdc24, to free Gβ, thereby promoting polarization to sites of active receptor (Nern and Arkowitz, 1999). The duration of receptor signaling prior to the repurposing of the polar cap will vary depending on where in the cell cycle each cell is when pheromone signaling begins. Thus, some cells may be an hour or more into pheromone signaling prior to completing cytokinesis, while others may be able to immediately start mating projection formation or experience a delay of only a few minutes. A potentially significant difference between these two scenarios is the amount of RGS present in the cell (Figure 2), as SST2 transcription is upregulated by pheromone signaling (Dohlman et al., 1996), and so cells that must delay receptor driven polarity for a long time prior to cytokinesis may be more prone to RGS-induced errors and be more dependent upon MAPK phosphorylation of the RGS.

We have found the delay of receptor driven polarity is promoted by phosphorylation of the RGS and by the kelch-repeat protein, Kel1, which associates with the polar cap, regulates the formin Bnr1, is required for efficient mating, and takes part in the mitotic exit network (MEN) (Gould et al., 2014; Hofken and Schiebel, 2002; Philips and Herskowitz, 1998; Smith and Rose, 2016). Kel1 contributes to the MEN by anchoring the Ras regulator Lte1 to the daughter cell during mitosis (Geymonat et al., 2010; Hotz and Barral, 2014). In addition to promoting mitotic exit, Kel1 and Lte1 have been found to suppress spurious polarization prior to the completion of mitosis, a role that may be separate from their role in MEN (Geymonat et al., 2010). Failure of Lte1 suppression of polarized growth leads to asymmetric hyperpolarized growth very similar to what we see in HU (Figure 7) (Geymonat et al., 2010). Based on our results with HU, it appears that unphosphorylated RGS is disrupting this Lte1 polarity suppression pathway, while phosphorylation of the RGS promotes Lte1 polarity suppression. Geymonat and colleagues described this suppression of polarity in terms of Rsr1/Bud1 driven polarity (Geymonat et al., 2010), which would normally drive the activation of Cdc42 at the insipient bud site for the next round of mitosis (Kozminski et al., 2003). Here we see a block in polarity driven by Gβγ associat-

ed Cdc42 GEF (Cdc24) (Butty et al., 1998), which suggests that the ability of Kel1 and Lte1 to inhibit polarity is broadly applicable and may serve further functions in the pheromone response.

The role of unphosphorylated RGS

At later time points the RGS is dephosphorylated (Figure 6), suggesting a switch in the requirements for RGS as the pheromone response progresses. We found that the localization of the RGS to the center of the polar cap is inhibited by phosphorylation. This phospho-dependent change in localization may seem unintuitive at first glance, as RGS binds to the Receptor at the periphery of the cell, in an interaction required for proper pathway desensitization (Ballon et al., 2006). Disruption of receptor binding with the *sst2^Q304N* mutant causes supersensitivity of the pheromone pathway (Ballon et al., 2006; Dixit et al., 2014), but phosphorylation state of RGS has no effect on pathway sensitivity (Garrison et al., 1999). Additionally, receptor binding is inhibited by receptor phosphorylation (Ballon et al., 2006), and so only a subset of the receptor on the cell surface at any given time may be capable of RGS binding. All of this suggests that the change in polar cap localization of the RGS is due to interactions with a protein other than the receptor, and indeed, we do still see a spike in phosphomimetic p*RGS concentration at the center of the polar cap that may represent the receptor associated RGS. Our data suggest that the functional consequence of RGS phosphorylation is altered spatial regulation of the pathway (Figure 8).

Unphosphorylated RGS (e.g. WT at later time points or the p̶RGS mutant) drives a larger distance between active large G-protein (Gpa1) and active small G-protein (Cdc42) than cells expressing the phospho-mimetic p*RGS (Figure 5). The simplest explanation of this observation is that the presence of RGS at the polar cap locally suppresses Gα activation. This type of negative feedback to the center of active signaling could help drive wandering of the polar cap by promoting large G-protein signaling further from the current site of polarization (Howell et al., 2012). Wandering of the polar cap is important for sensitive gradient tracking (Dyer et al., 2013), and so the small difference in the ability to track the gradient very well that we see in the p*RGS mutant (Figure 1) may be due to its decreased offset between the receptor driven large G-protein and the Cdc42 driven polarity machinery. Additionally, an offset between receptor signaling and the polar cap has recently been proposed to play a role in gradient tracking (Ghose et al., 2021), and while we did not see an effect on gradient tracking here, under more difficult conditions, the observed Gα offset may enhance chemotropic growth.

The phosphorylation of RGS at early time points appears to allow Kel1 to function in delaying early

polarization, while unphosphorylatable RGS inhibits that function. This suggests that the role of unphosphory-lated RGS at later time points likely interferes with Kel1 function. This may be to regulate the formin, Bnr1, or it may be to regulate Lte1. Regulation of a formin might be the expected function, but Bnr1 appears to have little role in the pheromone pathway. Bni1 would seem a more likely candidate, and while there is no evidence that Kel1 can regulate Bni1, neither is there clear evidence that it cannot (Gould et al., 2014). Our hydroxyurea experiments show that the asymmetric hyper-polarized growth requires Bni1 and is inhibited by Kel1, consis-tent with a role for Kel1 negatively regulating Bni1, but as mentioned above, this phenotype is likely dependent upon the polarity suppressing effects of Lte1. An intriguing possibility is that unphosphorylated RGS regulates Lte1 function at later times to promote polarity.

Conclusion

Here we have established a role for the phosphorylation of the RGS, Sst2, in promoting the completion of cytokinesis prior to the pheromone pathway repurposing the mitotic Cdc42 machinery for production of the mating projection, or shmoo. The exacerbation of this defect by the presence of a DNA-damaging agent emphasizes that the cell must integrate the competing signals of a check point instructing the cell to stop mitosis with a GPCR signaling pathway instructing the cell to polarize towards a mating partner. The use of a short-term phosphorylation event to temporarily alter RGS function and thereby allow the mitotic checkpoint to complete seems a likely motif to repeat in other systems.

Acknowledgements

This research was supported by the National Institute of General Medical Sciences of the National Institutes of Health under Award Number R15GM128026. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. The authors declare no conflicts of interest.

Author Contributions

William Simke contributed to conceptualization, investigation, data curation, formal analysis, investigation, visualization, writing -original draft, and writing -review & editing. Andrew Hart contributed to conceptualiza-tion, investigation, formal analysis, and visualization and writing -review and editing. Cory Johnson contrib-uted to investigation, methodology, validation, and writing -review and editing. Sari Mayhue contributed to conceptualization, methodology, investigation, and data analysis. Lucas Craig contributed to investigation and

formal analysis. Savannah Sojka contributed to investigation. Joshua Kelley contributed to funding acquisition, conceptualization, formal analysis, methodology, investigation, supervision, software, visualization, and writing - original draft.

Methods

Yeast Strains. Strains used in this study are shown in Table 2. Strains were constructed in the MATa haploid Sac-charomyces cerevisiae strain, BY4741. Proteins were tagged with EGFP or mRuby2 at its chromosomal locus through oligonucleotide-directed homologous recombination with EGFP-spHIS5 amplified with primers listed in Table 3 from the tagging vectors listed in Table 1 (Lee et al., 2013) or the GFP collection(Huh et al., 2003). Sst2 phosphomutants were made by integrating the codon of interest with a PCR amplified CORE cassette(Storici and Resnick, 2006). Deletions were performed by first amplifying the genomic locus from the MATa haploid de-letion collection (Dharmacon) with primers listed in Table 3 and transformed using the standard high efficiency lithium acetate transformation.

Cells were grown in rich medium (YPD) or synthetic medium (SC) at 30°C unless otherwise indicated. PCR products were transformed into yeast strains using standard lithium acetate transformation procedure. Individual colonies were isolated by growth on standard selective media (SC leu-, SC ura-, SC his-,), selective media with 5-fluoroorotic acid (Zymo Research, Tustin, CA), or YPD selective media (YPD G418+). Transfor-mants were verified using fluorescence microscopy, sequencing, and/or PCR.

Hydroxyurea experiments. Yeast cultures were grown to an OD600 of 0.4-0.6 at 30°C and then pre-treated with 100mM of hydroxyurea (Alfa Aesar, Tewksbury, MA) for 2 hours at 30°C. After 2 hours of pre-treatment with HU, a saturating concentration of α-factor was added and cultures continued to incubate. Cultures were then fixed at 90 minutes and 240 minutes using an overnight ethanol fixation at -20°C. Following ethanol fixation, yeast were resuspended and washed twice in 50nM sodium citrate buffer (pH 7.2). Next, the cultures were incubated with 20mg/mL RNase A (Thermo Scientific) for a minimum of 1 hour at 37°C. Following RNase incubation, proteinase K (Thermo Scientific) was added to the cultures at a final concentration of 0.4mg/mL and incubated at 55°C for a minimum of 1 hour then placed at 4°C overnight. For imaging, cells were pelleted, then washed and mounted in 1X PBS (pH 7.4). Cells were then imaged on the IX83 epiflourescent microscope (Olympus).

Spontaneous cytokinesis defect experiments. Yeast strains were grown in liquid synthetic complete media with 2 % dextrose (SCD) at 30 °C to an OD600 of 0.6 – 0.8. Cells were treated with 30 μM α-factor for 90 minutes. Cells then were fixed in 4 % paraformaldehyde, 2 % glucose, and 30 μM α-factor for 20 minutes. After fixation

and 3 washes with 1x phosphate buffered saline (PBS), the cells were stained with 7 μM Calcofluor White for 20 minutes and 50 μg/mL of Concanavalin A (both obtained from Biotium, Fremont CA) for 30 minutes. The cells were once again washed 3 times with 1x PBS and then imaged. Randomly chosen fields were imaged and then cells were scored for failed cytokinesis.

Antibody production. The following peptides corresponding to the Sst2 amino acid sequence surround Serine 539 were synthesized by Genscript (Piscataway NJ), the phospho-Sst2 S539 peptide LHPH**S**PLSEC, where the **S** was phosphorylated, and the unphosphorylated peptide LHPHSPLSEC. The phospho-peptide was injected into Rabbits by Cocalico Biologicals (Reamstown PA) according to their standard protocol. Antibody was affinity purified on phospho-peptide covalently bound to a SulfoLink column according to the manufacturer's instructions (Thermo Scientific).

Western blotting. Phosphorylation state of Sst2 was assessed by Western blotting. Yeast cultures were grown overnight in 30°C. Cells were lysed with TCA buffer and protein concentrations were determined using DC protein assay kit (BioRad). Protein separation was performed with a 7.5% SDS-PAGE and transferred to nitrocellulose at 100 V for 90 mins. Primary antibody (1:1,000) and non-phosphopepetide (1:10,000) were incubated in 1% PBST blocking solution overnight followed by secondary antibody incubation (1:10,000) in 1% PBST blocking solution for 1 hr. Band intensity was detected via Odyssey CLx imaging system (LI-COR) and then quantified using ImageJ.

Imaging on Agarose Pad. Yeast were imaged on an Olympus IX83 with a 60X-TIRF 1.49 NA objective, a Photometrics Prime95b camera, X-Cite LED 120 Boost fluorescence light source (Excelitas), and filters for DAPI and GFP (Semrock). Cells were grown to mid-log phase (OD600 = 0.1 to 0.8) at 30°C in Synthetic Complete Media with 2% dextrose (SCD) and then imaged on pads made of 2% agarose in SCD, as the use of agarose leads to lower autofluorescence than standard agar pads. Imaging was performed with an objective heater (Bioptechs) set to 30°C. Cells were pelleted and then resuspended in SC with 3uM α-factor and placed on an agarose pad as above.

Microfluidics Experiments. Microfluidic devices were made by using a Silicone polymer poured onto a microfluidics device mold (Suzuki et al., 2021) fabricated by UMaine FIRST. SYLGARD 184 Silicone Polymer was mixed at a ratio of 10:1, part A to part B, using a glass stirring rod to mix (Dow). Mixed polymer was poured onto the device mold and placed in a vacuum chamber for 1hr. After all air bubbles were removed, the mixture was placed in an oven at 80°C for 1hr. After cooling to room temperature, devices were cut out using a razor and ports

were punctured using an 18g Leuer stub. Prepped devices and coverslips were cleaned by spraying with methanol, ethanol, then water, and dried using an air hose. Devices and coverslips were exposed to oxygen plasma for 45 seconds in a Harrick Plasma PDC32G Cleaner followed by fusion of the device to the cover slip.

Cultures were grown in SC to an OD600 between 0.1–0.8 at 30°C. Live-cell microfluidics experiments were performed using an IX83 (Olympus, Waltham MA) microscope with a Prime 95B CMOS Camera (Photometrics) controlled by Cell Sens 1.17 (Olympus). Fluorescence and Differential Interference Contrast (DIC) images were acquired using an Olympus-APON-60X-TIRF objective. Z-stacks of GFP and RFP images were acquired using an X-Cite 120 LEDBoost (Excelitas). Cells were imaged in a microfluidic device based on the Dial-a-wave design that allows for the rapid switching of media while holding the yeast in place (Appendix A) (Bennett 2008, Dixit 2014). Pheromone addition was verified using AlexaFluor 647 dye (Life Technologies) imaged with a single plane image. Cells were imaged at 20 min intervals for 12 hours for 300nM experiments and 5 min intervals for 0-150nM experiments. Confocal microscopy was conducted on a Leica DMi8 (Leica) imaging platform equipped with an automated stage, SP8X white light laser (capped at 70% of total power), an argon laser (Leica Microsystems Buffalo Grove, IL). All imaging was conducted using HyD hybrid detectors. Imaging settings were determined based on experimental needs and were replicated for repeat experiments.

Image Analysis. . Images were deconvolved using Huygens Software (Scientific Volume Imaging, Hilversum, Netherlands) Classic Maximum Likelihood Estimation (CMLE) Deconvolution Algorithm. Masks of cells were made using ImageJ (Schindelin et al., 2012) and data analysis was performed using MATLAB (MathWorks, Natick, MA). To quantify the fraction of protein localization over time, MATLAB was utilized. Masks of cells were loaded into MATLAB and each cell was labeled over time to track individual cells over time. The fluorescent intensity of each fluorescent protein was extracted over time using a line width of 5 pixels as previously described (Kelley et al., 2015). Peak Bem1 was used as a reference to normalize the distribution of proteins of interest in relation to the polar cap. This was done by setting peak Bem1 as the midpoint and shifting the protein of interest in the same manner. For profiles reporting fraction of protein at each position, fluorescence was normalized by subtracting the minimum value from each line-scan, followed by normalization of the subtracted data to sum to one. The normalized fluorescence intensity was plotted at each point along the cell periphery with shaded regions showing 95% confidence intervals derived by bootstrapping with 10000 resamplings. Statistical analysis was performed between profiles using a sliding one-way ANOVA and Tukeys honestly significant difference (HSD) test followed by false discover rate adjustment with the MATLAB *mafdr()*

function with p values <0.05 denoted as significant. Where indicated, a pairwise Kolmogorov-Smirnov test was performed using the MATLAB *kstest2()* function.

References

Alvaro, C.G., and J. Thorner. 2016. Heterotrimeric G Protein-coupled Receptor Signaling in Yeast Mating Pheromone Response. *The Journal of biological chemistry*. 291:7788-7795.

Amaral, N., A. Vendrell, C. Funaya, F.Z. Idrissi, M. Maier, A. Kumar, G. Neurohr, N. Colomina, J. Torres-Rosell, M.I. Geli, and M. Mendoza. 2016. The Aurora-B-dependent NoCut checkpoint prevents damage of anaphase bridges after DNA replication stress. *Nat Cell Biol*. 18:516-526.

Apanovitch, D.M., K.C. Slep, P.B. Sigler, and H.G. Dohlman. 1998. Sst2 is a GTPase-activating protein for Gpa1: purification and characterization of a cognate RGS-Galpha protein pair in yeast. *Biochemistry*. 37:4815-4822.

Arkowitz, R.A. 2009. Chemical gradients and chemotropism in yeast. *Cold Spring Harbor perspectives in biology*. 1:a001958.

Ballon, D.R., P.L. Flanary, D.P. Gladue, J.B. Konopka, H.G. Dohlman, and J. Thorner. 2006. DEP-domain-mediated regulation of GPCR signaling responses. *Cell*. 126:1079-1093.

Bar-Shavit, R., M. Maoz, A. Kancharla, J.K. Nag, D. Agranovich, S. Grisaru-Granovsky, and B. Uziely. 2016. G Protein-Coupled Receptors in Cancer. *Int J Mol Sci*. 17.

Bi, E., and H.O. Park. 2012. Cell polarization and cytokinesis in budding yeast. *Genetics*. 191:347-387.

Breitsprecher, D., and B.L. Goode. 2013. Formins at a glance. *J Cell Sci*. 126:1-7.

Burchett, S., P. Flanary, C. Aston, L. Jiang, K. Young, P. Uetz, S. Fields, and H. Dohlman. 2002. Regulation of stress response signaling by the N-terminal dishevelled/EGL-10/pleckstrin domain of Sst2, a regulator of G protein signaling in Saccharomyces cerevisiae. *The Journal of biological chemistry*. 277:22156-22167.

Buttery, S.M., S. Yoshida, and D. Pellman. 2007. Yeast formins Bni1 and Bnr1 utilize different modes of cortical interaction during the assembly of actin cables. *Mol Biol Cell*. 18:1826-1838.

Butty, A.C., P.M. Pryciak, L.S. Huang, I. Herskowitz, and M. Peter. 1998. The role of Far1p in linking the heterotrimeric G protein to polarity establishment proteins during yeast mating. *Science*. 282:1511-1516.

Chasse, S., P. Flanary, S. Parnell, N. Hao, J. Cha, D. Siderovski, and H. Dohlman. 2006. Genome-scale analysis reveals Sst2 as the principal regulator of mating pheromone signaling in the yeast Saccharomyces cerevisiae. *Eukaryotic cell*. 5:330-346.

Chiou, J.G., M.K. Balasubramanian, and D.J. Lew. 2017. Cell Polarity in Yeast. *Annu Rev Cell Dev Biol*. 33:77-101.

Ciejek, E., and J. Thorner. 1979. Recovery of S. cerevisiae a cells from G1 arrest by alpha factor pheromone requires endo-peptidase action. *Cell*. 18:623-635.

de Godoy, L.M., J.V. Olsen, J. Cox, M.L. Nielsen, N.C. Hubner, F. Frohlich, T.C. Walther, and M. Mann. 2008. Comprehensive mass-spectrometry-based proteome quantification of haploid versus diploid yeast. *Nature*. 455:1251-1254.

DiBello, P.R., T.R. Garrison, D.M. Apanovitch, G. Hoffman, D.J. Shuey, K. Mason, M.I. Cockett, and H.G. Dohlman. 1998. Selective uncoupling of RGS action by a single point mutation in the G protein alpha-subunit. *The Journal of biological chemistry*. 273:5780-5784.

Dixit, G., J.B. Kelley, J.R. Houser, T.C. Elston, and H.G. Dohlman. 2014. Cellular noise suppression by the regulator of G protein signaling Sst2. *Molecular cell*. 55:85-96.

Dohlman, H., D. Apaniesk, Y. Chen, J. Song, and D. Nusskern. 1995. Inhibition of G-protein signaling by dominant gain-of-function mutations in Sst2p, a pheromone desensitization factor in Saccharomyces cerevisiae. *Molecular and cellular biology*. 15:3635-3643.

Dohlman, H.G., J. Song, D. Ma, W.E. Courchesne, and J. Thorner. 1996. Sst2, a negative regulator of pheromone signaling in the yeast Saccharomyces cerevisiae: expression, localization, and genetic interaction and physical association with Gpa1 (the G-protein alpha subunit). *Molecular and cellular biology*. 16:5194-5209.

Dyer, J.M., N.S. Savage, M. Jin, T.R. Zyla, T.C. Elston, and D.J. Lew. 2013. Tracking shallow chemical gradients by actin-driven wandering of the polarization site. *Current biology : CB*. 23:32-41.

Elion, E.A., B. Satterberg, and J.E. Kranz. 1993. FUS3 phosphorylates multiple components of the mating signal transduction cascade: evidence for STE12 and FAR1. *Mol Biol Cell*. 4:495-510.

Errede, B., L. Vered, E. Ford, M.I. Pena, and T.C. Elston. 2015. Pheromone-induced morphogenesis and gradient tracking are dependent on the MAPK Fus3 binding to Galpha. *Mol Biol Cell*. 26:3343-3358.

Evangelista, M., K. Blundell, M.S. Longtine, C.J. Chow, N. Adames, J.R. Pringle, M. Peter, and C. Boone. 1997. Bni1p, a yeast formin linking cdc42p and the actin cytoskeleton during polarized morphogenesis. *Science*. 276:118-122.

Gao, L., W. Liu, and A. Bretscher. 2010. The yeast formin Bnr1p has two localization regions that show spatially and temporally distinct association with septin structures. *Mol Biol Cell*. 21:1253-1262.

Garrison, T., Y. Zhang, M. Pausch, D. Apanovitch, R. Aebersold, and H. Dohlman. 1999. Feedback phosphorylation of an RGS protein by MAP kinase in yeast. *The Journal of biological chemistry*. 274:36387-36391.

Geymonat, M., A. Spanos, S. Jensen, and S.G. Sedgwick. 2010. Phosphorylation of Lte1 by Cdk prevents polarized growth

during mitotic arrest in S. cerevisiae. *The Journal of cell biology*. 191:1097-1112.

Ghose, D., K. Jacobs, S. Ramirez, T. Elston, and D. Lew. 2021. Chemotactic movement of a polarity site enables yeast cells to find their mates. *Proc Natl Acad Sci U S A*. 118.

Gould, C.J., M. Chesarone-Cataldo, S.L. Alioto, B. Salin, I. Sagot, and B.L. Goode. 2014. Saccharomyces cerevisiae Kelch proteins and Bud14 protein form a stable 520-kDa formin regulatory complex that controls actin cable assembly and cell morphogenesis. *The Journal of biological chemistry*. 289:18290-18301.

Hao, N., S. Nayak, M. Behar, R.H. Shanks, M.J. Nagiec, B. Errede, J. Hasty, T.C. Elston, and H.G. Dohlman. 2008. Regulation of cell signaling dynamics by the protein kinase-scaffold Ste5. *Molecular cell*. 30:649-656.

Hauser, A.S., M.M. Attwood, M. Rask-Andersen, H.B. Schioth, and D.E. Gloriam. 2017. Trends in GPCR drug discovery: new agents, targets and indications. *Nat Rev Drug Discov*. 16:829-842.

Hofken, T., and E. Schiebel. 2002. A role for cell polarity proteins in mitotic exit. *The EMBO journal*. 21:4851-4862.

Hotz, M., and Y. Barral. 2014. The Mitotic Exit Network: new turns on old pathways. *Trends Cell Biol*. 24:145-152.

Howell, A., M. Jin, C.-F. Wu, T. Zyla, T. Elston, and D. Lew. 2012. Negative feedback enhances robustness in the yeast polarity establishment circuit. *Cell*. 149:322-333.

Huh, W.K., J.V. Falvo, L.C. Gerke, A.S. Carroll, R.W. Howson, J.S. Weissman, and E.K. O'Shea. 2003. Global analysis of protein localization in budding yeast. *Nature*. 425:686-691.

Imamura, H., K. Tanaka, T. Hihara, M. Umikawa, T. Kamei, K. Takahashi, T. Sasaki, and Y. Takai. 1997. Bni1p and Bnr1p: downstream targets of the Rho family small G-proteins which interact with profilin and regulate actin cytoskeleton in Saccharomyces cerevisiae. *The EMBO journal*. 16:2745-2755.

Kelley, J.B., G. Dixit, J.B. Sheetz, S.P. Venkatapurapu, T.C. Elston, and H.G. Dohlman. 2015. RGS proteins and septins cooperate to promote chemotropism by regulating polar cap mobility. *Current biology : CB*. 25:275-285.

Kozminski, K., L. Beven, E. Angerman, A. Tong, C. Boone, and H. Park. 2003. Interaction between a Ras and a Rho GTPase couples selection of a growth site to the development of cell polarity in yeast. *Molecular biology of the cell*. 14:4958-4970.

Lappano, R., and M. Maggiolini. 2012. GPCRs and cancer. *Acta Pharmacol Sin*. 33:351-362.

Lee, S., W.A. Lim, and K.S. Thorn. 2013. Improved Blue, Green, and Red Fluorescent Protein Tagging Vectors for S. cerevisiae. *PLoS ONE*. 8:e67902.

Matheos, D., M. Metodiev, E. Muller, D. Stone, and M.D. Rose. 2004. Pheromone-induced polarization is dependent on

the Fus3p MAPK acting through the formin Bni1p. *The Journal of cell biology*. 165:99-109.

Metodiev, M.V., D. Matheos, M.D. Rose, and D.E. Stone. 2002. Regulation of MAPK function by direct interaction with the mating-specific Galpha in yeast. *Science*. 296:1483-1486.

Nern, A., and R.A. Arkowitz. 1999. A Cdc24p-Far1p-Gbetagamma protein complex required for yeast orientation during mating. *The Journal of cell biology*. 144:1187-1202.

Norden, C., M. Mendoza, J. Dobbelaere, C.V. Kotwaliwale, S. Biggins, and Y. Barral. 2006. The NoCut pathway links completion of cytokinesis to spindle midzone function to prevent chromosome breakage. *Cell*. 125:85-98.

Park, H.O., and E. Bi. 2007. Central roles of small GTPases in the development of cell polarity in yeast and beyond. *Microbiology and molecular biology reviews : MMBR*. 71:48-96.

Peter, M., A. Gartner, J. Horecka, G. Ammerer, and I. Herskowitz. 1993. FAR1 links the signal transduction pathway to the cell cycle machinery in yeast. *Cell*. 73:747-760.

Philips, J., and I. Herskowitz. 1998. Identification of Kel1p, a kelch domain-containing protein involved in cell fusion and morphology in Saccharomyces cerevisiae. *The Journal of cell biology*. 143:375-389.

Pope, P.A., S. Bhaduri, and P.M. Pryciak. 2014. Regulation of cyclin-substrate docking by a G1 arrest signaling pathway and the Cdk inhibitor Far1. *Current biology : CB*. 24:1390-1396.

Pruyne, D., L. Gao, E. Bi, and A. Bretscher. 2004. Stable and dynamic axes of polarity use distinct formin isoforms in budding yeast. *Mol Biol Cell*. 15:4971-4989.

Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J.Y. Tinevez, D.J. White, V. Hartenstein, K. Eliceiri, P. Tomancak, and A. Cardona. 2012. Fiji: an open-source platform for biological-image analysis. *Nature methods*. 9:676-682.

Segall, J.E. 1993. Polarization of yeast cells in spatial gradients of alpha mating factor. *Proc Natl Acad Sci U S A*. 90:8332-8336.

Seshan, A., A.J. Bardin, and A. Amon. 2002. Control of Lte1 localization by cell polarity determinants and Cdc14. *Current biology : CB*. 12:2098-2110.

Smith, J.A., and M.D. Rose. 2016. Kel1p Mediates Yeast Cell Fusion Through a Fus2p- and Cdc42p-Dependent Mechanism. *Genetics*. 202:1421-1435.

Storici, F., and M.A. Resnick. 2006. The delitto perfetto approach to in vivo site-directed mutagenesis and chromosome rearrangements with synthetic oligonucleotides in yeast. *Methods Enzymol*. 409:329-345.

Suzuki, S.K., J.B. Kelley, T.C. Elston, and H.G. Dohlman. 2021. Gradient Tracking by Yeast GPCRs in a Microfluidics Chamber. *In* G Protein-Coupled Receptor Screening Assays. 275-287.

Wang, Y., and H.G. Dohlman. 2004. Pheromone signaling mechanisms in yeast: a prototypical sex machine. *Science*. 306:1508-1509.

Wang, Y., L.A. Marotti, Jr., M.J. Lee, and H.G. Dohlman. 2005. Differential regulation of G protein alpha subunit trafficking by mono- and polyubiquitination. *The Journal of biological chemistry*. 280:284-291.

Yu, H., P. Braun, M.A. Yildirim, I. Lemmens, K. Venkatesan, J. Sahalie, T. Hirozane-Kishikawa, F. Gebreab, N. Li, N. Simonis, T. Hao, J.F. Rual, A. Dricot, A. Vazquez, R.R. Murray, C. Simon, L. Tardivo, S. Tam, N. Svrzikapa, C. Fan, A.S. De Smet, A. Motyl, M.E. Hudson, J. Park, X. Xin, M.E. Cusick, T. Moore, C. Boone, M. Snyder, F.P. Roth, A.L. Barabasi, J. Tavernier, D.E. Hill, and M. Vidal. 2008. High-Quality Binary Protein Interaction Map of the Yeast Interactome Network. 322:104-110.

A)

B)

Supplemental Figure S1. Characterization of the phospho-Sst2 antibody. A) Dot blot with the indicated quantities of peptide. The antibody is specific for phospho-peptide under the conditions used. All blotting is carried out with excess unphosporylated peptide to stop nonspecific binding. B) Our anti-phospo-Sst2 antibody detects a higher molecular weight band when detecting Sst2-GFP than when detecting native Sst2. Additionally, the time dependent decrease in phospho-Sst2 is not dependent upon Bar1-mediated degradation of pheromone.