Title: Metabolic crossfeeding structures the assembly of polysaccharide degrading communities

Sammy Pontrelli[1], Rachel Szabo[2], Shaul Pollak[2], Julia Schwartzman[2], Daniela Ledezma-Tejeida[1], Otto X. Cordero[2], Uwe Sauer[1]

[1]Institute of Molecular Systems Biology, ETH Zürich, Switzerland
[2]Department of Civil and Environmental Engineering, Massachusetts Institute of Technology, Cambridge

One sentence summary: Specialized biopolymer degraders structure microbial community assembly through interconnected modes of metabolic crossfeeding

Abstract:
Metabolic processes that fuel the growth of heterotrophic microbial communities are initiated by specialized biopolymer degraders that decompose complex forms of organic matter. It is unclear, however, to what extent degraders structure the downstream assembly of the community that follows polymer breakdown. Investigating a model marine microbial community that degrades chitin, we show that chitinases secreted by different degraders produce oligomers of specific chain lengths that not only select for specialized consumers but also influence the metabolites secreted by these consumers into a shared resource pool. Each species participating in the breakdown cascade exhibits unique hierarchical preferences for substrates, which underlies the sequential colonization of metabolically distinct groups as resource availability changes over time. By identifying the metabolic underpinnings of microbial community assembly, we reveal a hierarchical crossfeeding structure that allows biopolymer degraders to shape the dynamics of community assembly.

INTRODUCTION

Microbial communities mediate a staggering number of biological processes that contribute to the health of humans, animals(*1*) and the planet as a whole(*2*). The metabolic activity of co-occurring species results in the formation and depletion of nutrients in shared resource pools, where different modes of competition and cooperation impact community composition and form a network of metabolic crossfeeding(*3*, *4*, *13–22*, *5*, *23*, *6–12*). Individual species often provide specific metabolic functions that influence the growth of a broader community, including use of terminal electron acceptors(*14*, *24*, *25*), degradation of complex carbon sources(*26–29*), removal of toxic byproducts(*30–32*), or assimilation of nitrogen and sulfur(*18*, *33–35*). However, we remain with a limited molecular understanding of the extent to which individual species impact the formation and structure of crossfeeding networks and community assembly. Here, we demonstrate multiple metabolic mechanisms by which specialized biopolymer degraders influence the trajectory of community assembly, exemplified with an 18 member community that thrives on chitin, the second most abundant polysaccharide on the planet(*36*). Similar to other polymer degrading communities(*26–28*), chitin communities require specialized degraders to supply nutrients to nonspecialized downstream consumers. Our community consists of phylogenetically diverse seawater isolates with distinct metabolic capabilities, which become abundant at different points during assembly of chitin communities. This provides a system that allows us to demonstrate how specialized chitin degraders initiate a hierarchal food web that shapes the community during the assembly process.

RESULTS

Functional classification of bacterial isolates

We first classified the 18 isolates, collected from natural polymer degrading communities(*5*, *26*)(Table S1), into three functional guilds based on their phenotypic capabilities for chitin utilization (Fig 1A).

These species were tested for the ability to grow on chitin, the chitin monomer *N*-acetylglucosamine (GlcNAc), and the dimer chitobiose (Fig 1AB, S1, S2): Five degraders used all three substrates as a sole carbon and nitrogen source, and six exploiters could not use chitin but grew on GlcNAc, four of which also grew on chitobiose. Seven scavengers could not use any of these substrates; hence, they must rely on nutrients secreted by other species to thrive in the community. Previously published genome data of these 18 isolates(*5, 26*) revealed a wide range of chitin utilization potential (Fig 1A) that closely matches phenotypes. All five degraders, and two exploiters contain chitinases, uptake components specific for GlcNAc, and all genes required for conversion of GlcNAc into glycolytic intermediates: GlcNAc kinase, GlcNAc-6-phosphate deacetylase, and glucosamine-6-phosphate deaminase. The remaining four exploiters contain all metabolic components required to metabolize and transport GlcNAc (Fig 1A). All seven scavengers lack at least one transporter or catabolic gene required for GlcNAc utilization (Fig 1A). These varying metabolic capabilities prime guilds to become abundant at different points during community succession due to the different nutrients that become available over time(*5*). To visualize the trajectory of each functional guild during community assembly, we mapped 16s sequences of all species to 16s ribosomal gene exact sequence variants whose abundance changes were previously determined on chitin particles colonized from a large species pool in seawater(*26*). The mean trajectories of each functional guild revealed unique dynamics (Fig 1C), whereby degraders are the first to reach their maximum abundance, followed by exploiters and finally scavengers. This succession pattern portrays an order in a trophic cascade; i. e. degraders cause the release of chitin degradation products that are consumed by exploiters, which in turn release broader sets of metabolites that support scavengers.

Downstream metabolic influence of secreted chitinases

It is likely that degraders form publicly available products that can support the growth of exploiters and possibly select for specific species. To test this, extracellular enzymes were concentrated from culture supernatants of each degrader during growth on chitin and used to digest fresh colloidal chitin. Liquid chromatography mass spectrometry (LC-MS) quantification of oligosaccharides with six or fewer residues revealed that each digest resulted in a unique oligomer profile, with the disaccharide chitobiose and the monomer GlcNAc as the most abundant products (Fig 2A). Each of the six exploiters was then grown on the five chitin digests (Fig 2B, Fig S3). *Sil*A3R06 and three other exploiters grew well on all digests. While *Pha*B3M02 and *Alt*A3R04 hardly grew on the *Psy*6C06 digest that contained chitobiose as the major oligomer (Fig 2A), but grew on *Vib*1A01 and *Psy*6C06 digests containing larger quantities of GlcNAc and other oligomers (Fig 2A). Consistently, *Pha*B3M02 and *Alt*A3R04 did not grow on chitobiose as the sole carbon source (Fig 2B, Fig S3). Inspired by this observation, we tested whether the differences in hydrolysis product profiles among degraders could control which exploiters grow in coculture. For this purpose, we grew degraders *Vib*1A01 and *Psy*6C06 in coculture with the four exploiters. In accord with the enzyme digest data (Fig 2A), *Vib*1A01, which produces a range of oligomers supported growth of all species in coculture (Fig 2D). The chitobiose producer *Psy*6C06 was able to support all exploiters except *Pha*B3M02, showing that the composition of degradation products can influence the abundance of specific exploiters. Surprisingly, the exploiter *Alt*A3R04 also grew in coculture with *Psy*6C06, although it grew poorly on the enzyme digest of *Psy*6C06 and showed no growth on chitobiose (Fig 2b). This suggests that additional metabolites were secreted by *Psy*6C06 or another exploiter to support growth of *Alt*A3R04 but not of *Pha*B3M02, and more generally, that multiple resource pools are formed during chitin degradation that allow downstream growth of less specialized species.

To understand the mechanism that facilitated the growth of AltA3R06 in coculture, we investigated whether degraders secrete additional metabolites during growth on chitin to provide a carbon or nitrogen source independent from GlcNAc oligomers. We grew the five degraders on colloidal chitin until early stationary phase and inoculated the cell-free culture supernatants individually with six exploiters and seven scavengers (Fig 2C). While four of the supernatants did not support growth of any species to detectable optical densities, presumably because degraders consumed available limiting nutrients required for growth, *Psy*6C06 supernatant supported the growth of six scavengers and three exploiters. LC-MS targeted metabolomics revealed that GlcNAc and chitobiose concentrations were below the limit of detection in all five supernatants. However, nine metabolites were present in at least one supernatant, including succinate, glutamate, aspartate, and propionic acid (Fig S4, supplemental dataset 1). Since catabolism of GlcNAc into the glycolytic intermediate glucose 6-phosphate entails deacetylation and deamination, acetate or ammonia were quantified with enzyme assays (Fig 2E). Ammonia was likely the nitrogen source provided to other species since it was present at substantial concentrations in all supernatants. Acetate was present only in the *Psy*6C06 supernatant at a concentration 10.9 fold higher than the cumulative molar concentrations of all other detected metabolites. Hence, acetate was presumably the major carbon source in *Psy*6C06 cocultures that fueled growth of *Alt*A3R04 (Fig 2D 2C). Collectively, these results demonstrate that non-degrading community members are selectively supported by a combination of freely available chitin monomers, oligomers and metabolic byproducts secreted during growth of degraders.

We suspected that exploiters also secrete metabolites that less specialized species consume, and wondered if the chitin degradation products influence these secretion profiles. To test this hypothesis, we cultivated each exploiter on colloidal chitin digested with extracellular enzymes obtained from each of the five degraders. After 36 hours, culture supernatants were analyzed with the discovery

metabolomics method flow-injection time-of-flight (FIA-QTOF) MS(37). Overall, we detected 2521 ions, of which 216 could be annotated based on accurate mass. Principle component analysis using the 216 annotated ions showed a clear clustering of the secretion profiles based on the oligomer substrate (Fig 2F). For example, degraders *Vib*1A01 and *Vib*I3M07 produced similar sets of n1-n4 GlcNAc oligomers (Fig 2A). The secretion profiles of exploiters that used these substrates also showed no clear separation, suggesting that *Vib*1A01 and *Vib*I3M07 exert similar effects on exploiters *via* their similar chitinases. Alternatively, degraders *Psy*6C06 and *Vib*6D03 produced primarily chitobiose and GlcNAc, respectively, and secretion profiles of exploiters grown on these digests clustered distinctly. These results demonstrate that the different chitin degradation products play a major role in determining which metabolites exploiters secrete into a shared resource pool.

Identification of strain-specific metabolic secretion and uptake profiles

We designed two experiments to identify which of the many secreted metabolites could potentially serve as primary nutrient sources for scavengers. First, all scavengers were cultivated on cell-free supernatants collected from each of the degraders and exploiters after growth on GlcNAc that was used by all species. All supernatants except *Sil*A3R06 supported growth of at least two, but typically more scavengers (5.27 ± 2.4 with $OD_{600} > 0.1$, Fig 3A), demonstrating secretion of large quantities of metabolites by most species. Measuring 24 absolute metabolite concentrations by targeted LC-MS and enzymatic assays confirmed acetate and ammonia as the major carbon and nitrogen sources, respectively (Fig 3BC, supplemental dataset 1), akin to the results obtained with the *Psy*6C06 chitin supernatant. Yet, additional metabolic niches are manifested by a variety of organic acids, amino acids, and nucleosides that made up as much 31% of the total secreted carbon pool, as in the case of *Col*C2M11. The second experiment was designed to test how these additional metabolites are partitioned among all community members. For this purpose, supernatants of degraders grown on

chitin and degraders and exploiters grown on GlcNAc were pooled into a single mixture containing all secreted metabolites. All 18 strains were inoculated in this supernatant and grew individually to densities between 0.15 and 0.7 OD_{600} units, corresponding to 4-6 doublings (Fig S5). To identify consumed metabolites that support this growth, we determined relative changes by FIA-QTOF-MS and searched for metabolites that were i) produced (log_2 fold change of ion intensity > 1) by degraders or exploiters after growth to late exponential phase in at least one of the cultures, and ii) consumed (log_2 < -0.5) by at least one species in samples taken over the course of growth in the pooled supernatant (Fig 3D, (Supplementary Dataset 2). Each species had a distinct metabolite consumption profile and a unique ordered hierarchy of which metabolites are consumed first (Fig 3D). This was exemplified by acetate, GlcNAc, pyruvate and glutamate (Fig 3E, Fig S6). For example, *Alt*A3R04 co-metabolized all four substrates, while *Sil*A3R06 and *Pha*B3M02 metabolized them with specific ordered preferences or not at all. These results suggest that species may be specialized to consume unique sets of resources with a distinct hierarchy of preferences, thus enhancing the potential for many species to coexist in environments with fluctuating nutrient availability.

Given the sequential colonization of degraders, exploiters, and scavengers (FIG 1C), we wanted to test whether unique metabolic preferences of guilds and species drive this succession. For this purpose, we approximated the availability of resource pools at early timepoints by determining which metabolites are produced by degraders after individual growth on colloidal chitin, since chitin degradation initiates community succession. Conversely, we approximated the resource pools that degraders and exploiters produce at later timepoints by determining metabolites they secreted after growth on GlcNAc. We then calculated the fraction of metabolites that each species consumed from the early or late resource pools (Fig 3F). Of the 32 metabolites present in the chitin-derived resource pool, degraders, exploiters, and scavengers consumed on average 13%, 28%, and 25% of the available metabolites, respectively.

Scavengers greatly increased their consumption to 45% of the 54 metabolites in the GlcNAc-derived resource pool, while degrader and exploiter consumption did not change much. Thus, metabolites produced at later timepoints in community succession are consumed preferentially by scavengers, explaining how this functional guild is specialized to thrive on resources that become available at later timepoints in community succession (Fig 1C).

Inferring metabolite exchange networks that explain coculture outcomes

In light of the unique secretion and uptake profiles of each species, we explored how these traits may contribute to the outcome of communities in coculture. Cocultures were seeded with one of three degraders and one of four exploiters with an additional set of the same four scavengers to create twelve six-member communities (Fig 4A). The three degraders extracellularly produced different sets of chitin degradation products (Fig 2A), which appeared to influence the prevalence of exploiters in several cases. For example, exploiter *Alt*A3R04 used GlcNAc but not chitobiose (Fig 2B) and consistently coexisted well with degrader *Vib*6D03 (Fig. 4A) that produced GlcNAc as a major degradation product (Fig 2A). However, degrader-exploiter coexistence was not entirely explained by their dependence on GlcNAc oligomers. For instance, exploiter *Col*C2M11 did not grow in coculture with degrader *Vib*1A01 (Fig. 4A), despite its ability to grow on chitobiose and GlcNAc formed by *Vib*1A01(Fig 2A, S3). This is likely the result of poor substrate affinity for GlcNAc oligomers and suggests that additional secreted metabolites may fuel *Col*C2M11.

For a more comprehensive and temporal view, we inferred coculture-specific hypothetical nutrient exchange networks by leveraging the sequential production and consumption data. Specifically, we compiled metabolite secretion and uptake data of individual species (Supplementary Dataset 3) to form pools where each metabolite was assumed to either be present or absent (Supplementary Dataset 4).

The first metabolite pool includes only degrader-derived metabolites; i.e. those produced during chitin degradation that then become available to the exploiter that next colonized the community. The second pool consists of exploiter-derived metabolites upon growth on the first metabolite pool. The exploiter-derived metabolite pool is defined as the degrader-derived pool minus metabolites consumed by the exploiter plus those secreted by the exploiter. A hypothetical metabolite exchange occurred between degrader and exploiter if a metabolite in the degrader-derived pool was produced by the degrader and consumed by the exploiter. Metabolite exchanges with scavengers were inferred from metabolites present in the exploiter-derived pool (i.e. produced by degrader or exploiter) and consumed by a scavenger. This resulted in twelve hypothetical exchange networks (Supplementary Dataset 5) where nodes represent a species that survived in coculture and edges represent exchanged metabolites. Figure 4B depicts a simplified illustration of two networks where the edge width is the number of exchanged metabolites between each species. These networks identify and quantify the number of metabolites exchanged between pairs of species, providing a basis to infer which interactions may contribute to coculture outcomes.

To assess how exploiters influenced the flow of metabolites from degraders to scavengers, we evaluated how exploiters may potentially compete with scavengers for degrader-derived metabolites by identifying the percentage of degrader-derived metabolites that are consumed by exploiters in the exchange networks (Table S2). No scavengers coexisted with *Sil*A3R06 in any coculture (Fig 4A), and on average *Sil*A3R06 consumed 78.3% of all degrader-derived metabolites, compared to at most 33.3% by any of the other exploiters. This suggests that *Sil*A3R06 competes with scavengers for degrader-derived metabolites to a much higher degree than any other exploiter. Moreover, to evaluate whether *Sil*A3R06 itself contributed metabolites to scavengers that may support their growth, we calculated the number of metabolites produced by each exploiter that scavengers may consume (Table S3, Fig 4B), revealing

that *Sil*A3R06 did not secrete any such metabolite. On the contrary, our networks predict that in six cases, exploiters secreted a larger variety metabolites than they consumed, suggesting that exploiters have the ability to expand the number of available metabolic niches to support a more diverse population. This is exemplified with *Col*C2M11 in coculture with *Vib*6D03 (Fig 4B), which only consumed GlcNAc, chitobiose and glutamate, yet it engaged in 27 hypothetical exchanges with scavengers that are mediated by ten metabolites. Subsequently, this is the only coculture in which growth of all scavengers was observed. Collectively, these data illustrate how exploiters, after their abundance in the community is influenced by degraders (Fig 2D), have downstream influence on community diversity by changing the resources available to scavengers.

Lastly, we aimed to identify metabolites that accounted for the growth of individual species in coculture (Figure 4A). To this end, we combined the metabolic uptake/secretion profiles of individual species (Supplementary Dataset 3) with the 6-member coculture outcomes (Figure 4A) and their respective metabolite pools (Supplementary Dataset 4). Given that many metabolite abundances have only relative and not absolute quantities, we used a Boolean approach that takes into consideration only the presence or absence of each metabolite. This approach inferred a potential causal relationship between a metabolite and a given species when three conditions were true: (1) the metabolite was shown to be consumed by the species in its uptake profile, (2) the metabolite was present in the inferred metabolite pool available to the species in all of the cocultures where it showed growth, and (3) the metabolite was absent in the inferred metabolite pool available to the species in all cocultures where it did not grow. With this approach we were able to identify potential causal metabolites, primarily organic acids, amino acids, and one vitamin, for 5 species (Table S4). In the case of *Col*C2M11, which grew in presence of two degraders but not in presence of *Vib*1A01, glutamate was the only metabolite that could potentially explain the growth phenotype. In another case, the scavengers *Para*C2R09, *Cit*C3M06 and *Mar*F3R11

coexisted together in eight of the twelve cocultures (Fig 4A), suggesting that there may be an overlap in metabolites that contribute to their growth. Our approach identified 4 potential causal metabolites that overlapped between two or three of these scavengers: isocitrate, alanine, pantothenate, and the isoleucine precursor, citraconate (Table S4). In a more specific example, scavenger *Mar*D2M19 only grew in coculture with exploiter *Col*C2M11 and degrader *Vib*6D03. We inferred seven metabolites in the exploiter-derived resource pool of this coculture that may underlie the growth of *Mar*D2M19. These include malate, fumarate, and glycolate, which are only secreted by *Col*C2M11 in comparison to the other three exploiters. Figure 4C is a graphical summary of metabolite exchanges that we speculate contribute to the coculture outcomes based on our findings. Included are all the inferred causal metabolites (Table S4) and metabolites that were likely to be broadly exchanged between many species. The latter includes metabolites that were commonly produced by degraders and exploiters after growth on GlcNAc (Fig 3B) which were shown to be consumed by scavengers (Fig 3D), such as glutamate, pyruvate, oxoproline, and the growth supporting substrate acetate (Fig 3B). Collectively, these metabolite exchanges illustrate how specific metabolites, in addition to broadly exchanged metabolites, may contribute to the presence of individual species.

DISCUSSION

Resolving microbial community function to the level of contributing species is hampered by the difficulty in monitoring how nutrients are dispersed between species and how individual species can impact the broader community through metabolic interaction networks. Here we demonstrate an approach for how community interaction networks can be learned from the characterization of individual species, and how specialized biopolymer degraders scaffold these networks to shape the population. Degraders secrete chitinases that produce different oligomer profiles that influence the abundance of exploiters, which thrive primarily on oligomers that they cannot generate themselves. Moreover, the composition

of oligomer profiles influences which metabolites exploiters secrete into a shared resource pool that is available to the entire population. As such, degraders initiate a hierarchical cascade of nutrient flow into a population.

An open question regarding the exchange of nutrients in a community is the extent to which it is mediated by broad and nonspecific crossfeeding networks(*6*), or by species-specific interactions with an ordered structure(*38–40*). Our results provide evidence that the exchange of nutrients has an ordered structure that allows individual species to shape the flow of metabolites, and subsequently that nonspecific crossfeeding networks do not capture the complexity of nutrient exchange. We show that a multitude of secreted metabolites provide metabolic niches to support a large population, including major growth supporting substrates such as acetate, ammonium and glutamate whose exchange appears to be nonspecific between most species. However, our results further demonstrate that scavengers, which can grow on neither chitin nor its degradation products, preferentially consume metabolites that are formed at later points in community succession, demonstrating how degraders scaffold downstream colonization of non-specialized species. With our approach of integrating individual species consumption and secretion data, we were able to infer metabolic exchange networks that hypothesize specific metabolites that support growth of different species.

Collectively, these results demonstrate how a hierarchical structure is formed within microbial communities by radiating crossfeeding networks. In principle, this crossfeeding structure can be generalized to the multitude of communities that require specialized biopolymer degraders, including those involved in human health(*1*), biogeochemical transformations(*2*), and biotechnology(*41*). Although hierarchical structures may be convoluted in more complex environments, we outline a concept for

generating hypotheses on crossfeeding networks when constructing smaller consortia that recapitulate key aspect of the entire community.

MATERIALS AND METHODS

Bacterial species and chemicals

All chemicals were purchased from Sigma Aldrich unless otherwise noted. GlcNAc oligonucleotide standards were obtained from Omicron Biochemicals. Media used are either Marine Broth 2216 (Difco #279100, Fisher) or MBL minimal media supplemented with a carbon source. MBL is mixed from several stock components: 4x concentrated seawater salts (NaCl 80g/l, $MgCl_2*6H_2O$ 12g/l, $CaCl_2*2H_2O$ 0.6g/l, KCL 2g/l), 1000x concentrated trace minerals ($FeSO_4*7H_2O$ 2.1g/l, H_3BO_3 30mg/l, $MnCl_2*4H_2O$ 100mg/l, $CoCl_2*6H_2O$ 190mg/l, $NiCl_2*6H_2O$ 24 mg/l, $CuCl_2*2H_2O$ 2mg/l, $ZnSO_4*7H_2O$ 144 mg/l, $Na_2MoO_4*2H_2O$ 36mg/l, $NaVO_3$ 25mg/l, $NaWO_4*2H2O$ 25mg/l, $Na_2SeO_3*5H_2O$ 6mg/l), 1000x concentrated vitamins (riboflavin 100mg/l, D-biotin 30mg/l, thiamine hydrochloride 100mg/l, L-ascorbic acid 100mg/l, Ca-d-pantothenate 100 mg/l, folate 100mg/l, nicotinate 100mg/l, 4-aminobenzoic acid 100mg/l, pyridoxine HCl 100mg/l, lipoic acid 100mg/l, NAD 100mg/l, thiamin pyrophosphate 100mg/l, cyanocobalamin 10mg/l), 1mM phosphate dibasic, 1mM sodium sulfate,50mM HEPES pH 8.2, and a carbon source. All stocks are diluted accordingly and the volume is adjusted to reach 1x concentrations of all components.

Species were received from the lab of Otto Cordero from Massachusetts Institute of Technology (table S1). All species were previously collected from polymer degrading communities of natural seawater communities from Canoe Beach, Nahant, MA, USA; 42°25'11.5'' N, 70°54'26.0'' W(*5, 26*). Species were stored in -80°C in glycerol stocks and streaked onto Marine Broth 2216 plates containing 1.5% agar (BD #214010) prior to use.

Crossfeeding with pooled supernatant

Single colonies were transferred into liquid Marine Broth 2216 and grown overnight at 220RPM at 27°C. Cells were centrifuged and washed in MBL containing no carbon source two times. This was then used to inoculate the respective culture to a final OD_{600} of 0.01. All supernatants were harvested from species grown in 125ml shake flasks containing 20ml of the respective media and were shaken at 100RPM at 27°C. Supernatants were collected by centrifugation and further filtered using a 0.2 µM PES Stericup Filter unit (Millipore). Supernatants were stored at -20°C until further use. Pooled supernatants were prepared by pooling equal volumes of each supernatant. This includes supernatants collected from each degrader grown on chitin, and each degrader and each exploiter grown on GlcNAc. The pooled supernatant was diluted 25% with fresh MBL containing no carbon source prior to inoculation. Species were inoculated into 1.8ml of pooled supernatant contained in 96 deep well plates with a 4mm glass bead. Plates were shaken at 220RPM at 27°C. Samples were taken every 2-4 hours after inoculation, and cells were removed by centrifugation. Supernatant was stored at -20°C prior to measurements.

Preparation of colloidal chitin

10g of chitin powder (Sigma C7170) was added to 100ml of phosphoric acid (85 wt. %) and incubated at 4°C for 48 hours. Roughly 500ml of fresh H_2O was added to this mixture and shaken until the chitin precipitated. This was then filtered using vacuum filter paper (Macherey-Nagel MN615) and fresh water was used to wash the filtrate three times. Chitin precipitate was then added into 3kDa dialysis tubing cellulose membrane (Sigma) and placed in water to remove residual monomers and oligomers. Fresh water was replaced daily three times. Colloidal chitin was then collected, adjusted to pH 7 with NaOH, and homogenized using a Bosch Silent Mixx Pro blender. The colloidal chitin was autoclaved before use.

Assay for acetate quantification

The following master mix was used for assays: 100mM Tris pH 7.4, 5mM $MgCl_2$, 1mM phosphoenolpyruvate, 1mM ATP, 3u/ml pyruvate kinase, 3u/ml acetate kinase. The assay was started by adding 20μl of sample to 200μl of master mix. The assay was monitored by measuring an absorbance at OD_{230}, and concentrations were determined by comparing the rate of PEP consumption over time compared to acetate standards.

Assay for ammonia quantification

The following master mix was used for assays: 100mM Tris pH 8.3, alpha-ketoglutarate 7mM, NADPH 0.25mM, 0.1u/ml glutamate dehydrogenase. The assay was started by adding 20μl of sample to 200μl of master mix. Consumption of NADPH was measured over time by measuring an absorbance at OD_{340}, and concentrations were determined by comparing slopes to an analytical standard of ammonium chloride.

Concentration of extracellular chitinases and enzyme digests

Degraders were inoculated into 50ml of 2g/l colloidal chitin MBL after growing in Marine Broth 2216 as a preculture. Cultures were grown until mid-late log phase (OD_{600} ~1), and all remaining chitin and cells were removed by centrifugation at 4°C. Supernatant was filtered using a 0.2 μM PES Stericup Filter unit (Millipore). Enzymes were concentrated from the supernatant 50 fold using a 10 kDa centrifugation protein concentration filter (Amicon Ultra, Millipore). Following this, the enzyme was added to fresh MBL containing 5g/l of chitin and sampled every 24 hours to measure for degradation products. The resulting digest was further sterile filtered and diluted by a factor of 2.5 and used to support growth of exploiters. Protein concentration was determined using the Bradford reagent.

Chitin coculture experiments

Individual colonies were inoculated into Marine Broth 2216 from solid medium and grown overnight at 220RPM at 27°C. Cells were then washed twice in MBL with no carbon and further inoculated into MBL with 2g/l chitin at $1x10^6$ CFU/ml, except for degraders, which were inoculated at $5x10^6$ CFU/ml for qPCR experiments and $2x10^7$ CFU/ml for 16s coculture experiments. All coculture experiments used 96 deep well plates containing 1.8ml media and a 4mm glass bead. Samples for quantifying species abundance were taken of the planktonic phase. Removal of colloidal chitin from samples was done by centrifuging the cultures at 500rcf for 30 seconds and the top later of culture from each well was sampled. Samples were stored at -20°C. Cell abundance was measured with qPCR or 16s sequencing.

Genomic DNA purification

Genomic DNA purification was done using DNAdvance Kit (Beckman A48705) following the manufacturers protocol with the modification that the input for the kit is 100µL of Lysis Master Mix combined with 100µL of bacterial culture that is diluted to an OD_{600} of less than 0.2.

Cell absolute quantification with qPCR

qPCR was performed using GoTaq qPCR Master Mix (Promega) following the manufacturer's protocol. qPCR reactions were prepared with a final volume of 15µL using prepared genomic DNA as a template. Samples were measured and data analyzed using a QuantStudio 3 Real-Time PCR System with ΔΔCT method. Absolute quantification of species was obtained by comparing C_t values with those obtained from standard cultures that have known CFU/ml values. PCR primers were designed to have an efficiency of 90-105% and sequences are listed in table S5. Primers were tested for specificity to the bacterial species used in each coculture.

16s sequencing

Measuring relative abundance of species in coculture was performed with amplicon 16s sequencing

using the MetaFast protocol (Fasteris, Switzerland) including proprietary MetaFast barcoded primers.

Amplicons were prepared by amplifying purified genomic DNA samples with a unique pair of MetaFast

barcoded primers with Kapa HIFI DNA polymerase. Further library preparation was performed by

Fasteris and samples were sequencing with an Illumina MiSeq platform.

Paired, quality-filtered reads obtained from 16S sequencing were processed using dada2, version 1.16 to

determine exact sequence variants (ESVs). Except where noted, the default parameters were used(*42*).

Briefly, sequences were trimmed to remove adapters and further trimmed from positions 25 to 175,

yielding 150 bp sequences with quality score >38. The maximum expected error rate was set to

maxEE(2,5) to ensure stringent trimming and better differentiate reads belonging to species with very

similar sequences. After learning error rates, samples were pooled to define variants. Species were

matched to ESVs using a custom BLAST built from the full length 16S genes of species used in the

experiment(*43*). The taxonomy of the ESVs was further confirmed by taxonomic identification using

DECIPHER. The SILVA SSU r138 database from 2019 was used for DECIPHER analysis(*44*).

Genome annotation

The species used for this analysis were isolated and sequenced as part of two experiments previously

performed in the Cordero lab for studying the succession of natural microbial communities from

seawater on polysaccharide particles(*5, 26*). For each genome, protein-coding genes were predicted and

translated using Prodigal v2.6.3(*45*). Predicted protein sequences were compared to a custom database

of profile hidden Markov models (HMMs) of proteins involved in growth on chitin using the *hmmsearch*

function of HMMER v3.3 with default parameters(*46*). Publicly-available HMMs were downloaded from

the Pfam v33.1(*47*) or TIGRFAM v15.0 databases(*48*) (see Supplementary dataset 6 for accession

numbers). Custom HMMs were made by identifying experimentally-verified proteins of interest(*49*, *50*),

finding their homologs in the UniProtKB/Swiss-Prot v2020_06 database(*51*), creating a seed alignment

using MAFFT v7 with default parameters(*52*, *53*), and building the profile HMMs using the *hmmbuild*

function of HMMER with default parameters (see Supplementary dataset 6 for details on each custom

HMM). The isolate strain's proteins were annotated based on the *hmmsearch* results if the protein

length was at least 100 amino acids, the independent E-value was less than 1×10^{-9}, and the domain

score was greater than 30. Only most significant annotation was used for each protein sequence. Gene

copy numbers were calculated for each genome by tallying the number of annotations made for each

protein group. Genome sequences for each organism can be found using NCBI accession numbers listed

in table S1.

Metagenomic 16S rRNA dynamics

The 16S rRNA sequence of each genome was determined by first amplifying it using the 27F (5-

AGAGTTTGATCMTGGCTCAG-3) + 1492R (5-GGTTACCTTGTTACGACTT-3) universal bacterial primers(19,

20), followed by sanger sequencing. We mapped the 16S sequences of our isolates back to the

metagenomic 16S trajectories (PRJNA319196, PRJNA478695) using blast(*43*). For each isolate 16S

matches in the metagenome were found using a moving threshold approach – we searched for matches

starting at 100% identity and going down to 97% identity at 1% identity steps, and if a match was found

in any identity threshold the search was ended and all metagnomic hits above that threshold were

considered to belong to that isolate 16S. The read counts were transformed into relative frequencies

and only entries with a relative abundance $>10^{-3}$ were retained for downstream analysis. Each experiment was originally performed in triplicate, and for each isolate we take the mean relative abundance at each time-point. For each isolate the trajectories were normalized by their highest relative abundance (so that the highest point of any trajectory is 1), and the average of these normalized trajectories is what is reported in Fig 1C.

Untargeted metabolite profiling and analysis

Untargeted methods were carried out using Flow Injection Analysis Quadroupole Time of Flight Mass Spectrometry (FIA-QTOF-MS). All supernatants were first diluted 100fold into water before measurements. Metabolomics was carried out with a binary LC pump (Agilent Technologies) and a MPS2 Autosampler (Gerstel) coupled to an Agilent 6520 time-of-flight mass spectrometer (Agilent Technologies). This was operated in negative mode, at 2Ghz for extended dynamic range, with a m/z (mass over charge ratio) range of 50-1000, as described previously(*37*). The mobile phase consisted of isopropanol:water (60:40, v/v) with 5mM ammonium fluoride buffer at pH 9 and the flow rate was 150 μl/minute. Raw data for all measurements was subjected to a spectral processing and alignment pipeline using Matlab (The Mathworks, Natick) as described previously(*37*).

Overall, we detected 2521 ions, of which 216 could be annotated based on accurate mass, corresponding to a maximum of 526 unique metabolites. Ions were annotated with a tolerance of 0.005 Da against a curated compound library that contains all metabolites predicted to be present in at least one of the species used in this study based on BioCyc databases. If a single m/z ion is matched with multiple isomeric or isobaric compounds within the compound library, the compound that participates in the largest number of enzymatic reactions in pathway databases for all 18 organisms used in this work

is chosen as the top annotation. Certain isomers were resolved using LC-MS methods as described

below, in which case the correct annotations were manually edited.

LC-MS exometabolomic compound measurements

Extracellular metabolomics measurements were performed as described previously(*54*). Briefly,

chromatographic separation was performed using a Poroshell 120 EC-CN 2.1 x 150 mm, 2.7 μm column

(Agilent) in an Agilent 1290 infinity stack set to 40°C. The method performed was isocratic, with a mobile

phase consisting of 10mM ammonia acetate pH 5.9 and 5% acetonitrile with a flow rate of 250μL/min

and a 5 μL injection volume. An Agilent 6520 series quadrupole time of flight mass spectrometer was

used for measurements with a m/z range of 20-400 operated in negative mode. Prior to LC-MS

measurements, samples were diluted both 20 and 100 fold in water to maximize the number of

measurable compounds within the limits of quantification. Data analysis was performed using Agilent

Masshunter Quantitative Analysis software. Peaks were integrated using spectral summation integration

with integration windows that are defined based on retention times observed from analytical standards.

The Agilent software was used to determine polynomial calibration curves with a goodness of fit (R^2 >

0.97).

GlcNAC oligomer quantification

GlcNAc oligomers were quantified using LC-MS with an Agilent 1100 series stack. Chromatographic

separation used a Poroshell 120 HILIC-Z 2.1 x 100 mm, 2.7 μm column (Agilent) kept at a temperature of

30°C and a constant flow rate of 1ml/min. 5 μL of sample was injected after five-fold dilution in 80%

acetonitrile. Mobile phase A consists of 10mM ammonia acetate pH 9 in water, and mobile phase B

consists of 10mM ammonia acetate pH 9 in 90% acetonitrile. At injection, the mobile phase contains

90% B phase for 2 minutes, followed by a gradient to 40% B phase until 12 minutes and held for one

more minute before being re-equilibrated at the starting condition. An Agilent 6520 series quadrupole

time of flight mass spectrometer was used for measurements, operated in negative mode at 2GHz for

extended dynamic range, with a m/z range of 100-1500. Calibration curves were measured using

analytical standards and used to calculate concentrations of oligomers in the sample.

Construction of hypothetical metabolite exchange networks

Metabolic exchange networks were constructed based on coculture outcomes (Fig 4A). This includes

secretion profiles of degraders grown on chitin (Supplementary dataset 1 and 2), GlcNAc oligomers that

degraders form extracellularly (Fig 3A), secretion profiles of exploiters after growth on pooled

supernatant (Supplementary dataset 2), and metabolites consumed by exploiters and scavengers in

pooled supernatant (Fig 3D). Combined metabolite production and consumption data used for

construction of metabolic networks is contained in Supplementary dataset 3. The metabolic networks

for each coculture infer two metabolite pools that mimic the sequential colonization of chitin particles

by degraders, exploiters and finally scavengers (Supplementary Dataset 4). In order to incorporate both

targeted datasets, which include absolute quantification, and untargeted metabolomics datasets, which

include a broad coverage of metabolites without absolute quantification, metabolites were noted to

have one of two possible states: present or absent. The first metabolite pool is a degrader-derived

metabolite pool consisting of metabolites produced by the degrader in the culture and represents

resources available to support growth of the exploiter. The second resource pool is an exploiter-derived

metabolite pool that represents metabolites available to support scavengers after the exploiter

consumed or added metabolites to the first, degrader-derived metabolite pool. To construct the

exploiter-derived metabolite pool, metabolites that are consumed by an exploiter are subtracted from

the degrader-derived metabolite pool, and metabolites that are produced an exploiter are added to the

degrader-derived metabolite pool. We next used these pools to infer metabolite exchanges between the

species in coculture to construct 12 coculture specific metabolite exchange networks (Supplementary Dataset 5). Each species that survived in coculture is a node in these networks, exchanged metabolites are represented by edges. To illustrate these networks in figure 4B, the edge width represents the number of exchanged metabolites. Metabolites exchanged between a degrader and an exploiter are metabolites present in the degrader-derived metabolite pool that are consumed by the exploiter. Metabolites exchanged between a scavenger and a degrader or exploiter are metabolites consumed by the scavenger from the exploiter-derived metabolite pool that are secreted by either the degrader or exploiter. In order to identify potentially causal metabolites, we used a Boolean approach where three conditions were needed to be simultaneously true in order to infer a causal relationship between a metabolite and a given strain: (1) the metabolite was shown to be consumed by the species in its uptake profile, (2) the metabolite was present in the degrader-derived or exploiter-derived metabolite pools of all cocultures where the exploiter or scavenger showed growth, respectively, and (3) the metabolite was absent in the degrader-derived or exploiter-derived metabolite pools of all cocultures where the exploiter or scavenger did not show growth, respectively. These criteria were only applicable for species that did not grow in at least one coculture. These inferred causal metabolites are in table S5.

Acknowledgments

Funding: This work was supported by a grant from the Simons Foundation (ID608247, SP and ID542395 OXC), as part of the collaboration on Principles of Microbial Ecosystems (PriME).

Author contributions: S Pontrelli designed and performed all experiments, analyzed data and wrote the manuscript. R Szabo performed genomic analysis of bacterial isolates. S Pollak mapped 16s sequences of bacterial isolates to ESVs of developing chitin communities. J Schwartzman performed 16s amplicon sequencing analysis. D Ledezma performed network analysis of chitin coculture outcomes. O Cordero assisted in data interpretation and writing the manuscript U Sauer designed the experiments and wrote the manuscript.

Competing interests: Authors declare no competing interests

Data and materials availability: All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. Metabolomics spectral files can be accessed from the MassIVE database (ftp://massive.ucsd.edu/MSV000087455/). The short read 16 s rRNA amplicon sequencing data (BioSample: SAMN11023523-SAMN11023755) have been deposited in NCBI with the BioProject identifier PRJNA478695. All deposited genome sequences have been deposited in NCBI with BioProject identifier and BioSample IDs specified in table S1.

Figure 1: Successional dynamics of chitin degrading seawater communities. A) Species used in this study. Heatmap depicting (left to right) final OD_{600} after growth on 2 g/l colloidal chitin, final OD_{600} after growth on 10 mM chitobiose, growth rate on 20 mM GlcNAc, classified functional guild, and copy numbers of genes relevant for chitin degradation. Growth data was obtained from three independent biological replicates. B) Schematic of functional guilds: Degraders can grow on chitin as a sole carbon source through breakdown of chitin using chitin degrading enzymes, exploiters can grow on monomeric or oligomeric N-acetylglucosamine (GlcNAc) as a sole carbon source, and scavengers require metabolites secreted by other species to be sustained in the community. C) 16s sequences of the 18 investigated species are mapped to 16s ribosomal gene exact sequence variants whose abundances were previously determined on chitin particles colonized from a species pool in seawater(*26*). Mean normalized frequencies of species that comprise each functional guild are plotted over time. TBDR, TolB-Dependent receptor

Figure 2: Chitin degradation products influence exploiter population and their secreted resources A) Quantification of GlcNAc oligonucleotide during *in vitro* digestion of 5 g/l chitin by secreted enzymes from each degrader. B) OD_{600} of three exploiters after 36 h growth on 20 mM GlcNAc, 10 mM chitobiose, and three enzyme digests. C) Growth of scavengers and exploiters after 36 hours on cell-free supernatant produced by each of the five degraders upon growth on colloidal chitin. D) Absolute abundance of species in cocultures of one degrader (*Vib*1A01 or *Psy*6C06) and four exploiters during grown on colloidal chitin, as determined by qPCR. E) Enzymatic quantification of ammonia and acetate in

degrader supernatants upon growth on colloidal chitin. F) Principal component analysis exploiter metabolite secretion profiles after growth on each colloidal chitin digest. Colors represent the digest used as a growth substrate. All error bars represent standard deviation from three biological replicates.

Figure 3: Metabolite consumption and production profiles among community members A) Growth of scavengers at 36 hours on cell-free supernatants obtained from degraders and exploiters after growth on GlcNAc. B) Metabolite concentration in cultures of degraders or exploiters after growth on 20 mM GlcNAc. C) Ammonia concentration in cultures of degraders or exploiters after growth on GlcNAc. D) Consumption of crossfed metabolites in pooled supernatant. (left) FIA-QTOF-MS measurements of metabolites that are consumed by degraders, exploiters or scavengers (Ion intensity $Log_2 < -0.5$). (right) Time course measurements of metabolite consumption. Points represent the fraction of the final OD_{600} at which metabolite consumption is detected for each species E) Relative change of acetate, pyruvate, GlcNAc, and glutamate during growth in pooled supernatant by three exploiters. Curves are fit using local polynomial regression. F) Fraction of available metabolites contained in chitin or GlcNAc derived resource pools that is consumed by each functional guild. All experiments were performed in triplicate. Error bars represent standard deviation. DEHEM, 4-deoxy-α-L-erythro-hex-4-enopyranuronate-β-D-mannuronate; OHCU, 2-oxo-4-hydroxy-4-carboxy-5-ureidoimidazoline. Strain names in green are degraders, blue are exploiters, and orange are scavengers.

Figure 4: Outcomes of chitin degrading cocultures and hypothetical metabolic exchange networks A) Each coculture was seeded with six species: one degrader (green), one exploiter (blue), and each of the four scavengers *Para*C2R09 (yellow), *Cit*C3M06 (orange), *Mar*D2M19 (red), *Mar*F3R11 (purple). Cultures were grown for five weeks with four serial dilutions on colloidal chitin and relative abundance is shown for the final culture. Experiments were performed in triplicate. B) Two of twelve hypothetical metabolite exchange networks that illustrate the number of metabolites exchanged between species that coexisted in cocultures outcomes from panel A. Edge width represents the number of exchanged metabolites. Exchange networks based on the remaining ten coculture outcomes are contained in Supplementary Dataset 5. C) A graphical summary of metabolite exchanges that likely contribute to the coculture outcomes. Bolded metabolites are all inferred causal metabolites that may contribute to the growth of individual species. Grey metabolites represent broadly exchanged growth supporting substrates that are secreted by degraders or exploiters grown on GlcNAc and consumed by scavengers. All remaining secreted metabolites and crossfeeding interactions that are not included in this illustration are contained in (Supplementary Dataset 1, Supplementary Dataset 3, and Supplementary Dataset 5). Abbreviation: DHMB, 2,3-dihydroxy-2-methylbutanoate

Figure 1

Figure 2

Figure 3

Figure 4

REFERENCES

1.   P. J. Turnbaugh, R. E. Ley, M. Hamady, C. M. Fraser-Liggett, R. Knight, J. I. Gordon, The Human Microbiome Project. *Nature*. **449**, 804–810 (2007).

2.   A. Konopka, What is microbial community ecology. *ISME J.* **3**, 1223–1230 (2009).

3.   O. Ponomarova, N. Gabrielli, D. C. Sévin, M. Mülleder, K. Zirngibl, K. Bulyha, S. Andrejev, E. Kafkia, A. Typas, U. Sauer, M. Ralser, K. R. Patil, Yeast Creates a Niche for Symbiotic Lactic Acid Bacteria through Nitrogen Overflow. *Cell Syst.* **5**, 345-357.e6 (2017).

4.   R. Baran, E. L. Brodie, J. Mayberry-Lewis, E. Hummel, U. N. Da Rocha, R. Chakraborty, B. P. Bowen, U. Karaoz, H. Cadillo-Quiroz, F. Garcia-Pichel, T. R. Northen, Exometabolite niche partitioning among sympatric soil bacteria. *Nat. Commun.* **6** (2015), doi:10.1038/ncomms9289.

5.   M. S. Datta, E. Sliwerska, J. Gore, M. F. Polz, O. X. Cordero, Microbial interactions lead to rapid micro-scale successions on model marine particles. *Nat. Commun.* **7** (2016), doi:10.1038/ncomms11965.

6.   J. E. Goldford, N. Lu, D. Bajić, S. Estrela, M. Tikhonov, A. Sanchez-Gorostiaga, D. Segrè, P. Mehta, A. Sanchez, Emergent simplicity in microbial community assembly. *Science (80-.).* **361**, 469–474 (2018).

7.   S. Blasche, Y. Kim, R. A. T. Mars, D. Machado, M. Maansson, E. Kafkia, A. Milanese, G. Zeller, B. Teusink, J. Nielsen, V. Benes, R. Neves, U. Sauer, K. R. Patil, Metabolic cooperation and spatiotemporal niche partitioning in a kefir microbial community. *Nat. Microbiol.* **6**, 196–208 (2021).

8.   A. R. Pacheco, M. Moel, D. Segrè, Costless metabolic secretions as drivers of interspecies interactions in microbial ecosystems. *Nat. Commun.* **10** (2019), doi:10.1038/s41467-018-07946-9.

9.   S. Louca, S. M. S. Jacques, A. P. F. Pires, J. S. Leal, D. S. Srivastava, L. W. Parfrey, V. F. Farjalla, M. Doebeli, High taxonomic variability despite stable functional structure across microbial communities. *Nat. Ecol. Evol.* **1**, 1–12 (2017).

10.   O. X. Cordero, L. A. Ventouras, E. F. DeLong, M. F. Polz, Public good dynamics drive evolution of iron acquisition strategies in natural bacterioplankton populations. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 20059–20064 (2012).

11.   J. H. Hehemann, P. Arevalo, M. S. Datta, X. Yu, C. H. Corzett, A. Henschel, S. P. Preheim, S. Timberlake, E. J. Alm, M. F. Polz, Adaptive radiation by waves of gene transfer leads to fine-scale resource partitioning in marine microbes. *Nat. Commun.* **7** (2016), doi:10.1038/ncomms12860.

12.   M. F. Romine, D. A. Rodionov, Y. Maezato, A. L. Osterman, W. C. Nelson, Underlying mechanisms for syntrophic metabolism of essential enzyme cofactors in microbial communities. *Nat. Publ. Gr.* **11**, 1434–1446 (2017).

13.   L. Kešnerová, R. A. T. Mars, K. M. Ellegaard, M. Troilo, U. Sauer, P. Engel, Disentangling metabolic functions of bacteria in the honey bee gut. *PLoS Biol.* **15**, 1–28 (2017).

14.   A. J. M. Stams, C. M. Plugge, Electron transfer in syntrophic communities of anaerobic bacteria

and archaea. *Nat. Rev. Microbiol.* **7**, 568–577 (2009).

15. Z. Wei, T. Yang, V. Friman, Y. Xu, Q. Shen, A. Jousset, Trophic network architecture of root-associated bacterial communities determines pathogen invasion and plant health. *Nat. Commun.* (2015), doi:10.1038/ncomms9413.

16. D. A. Rodionov, A. A. Arzamasov, M. S. Khoroshkin, S. N. Iablokov, S. A. Leyn, S. N. Peterson, P. S. Novichkov, Micronutrient Requirements and Sharing Capabilities of the Human Gut Microbiome. **10**, 1–22 (2019).

17. F. Moens, M. Verce, L. De Vuyst, Lactate- and acetate-based cross-feeding interactions between selected strains of lactobacilli, bifidobacteria and colon bacteria in the presence of inulin-type fructans. *Int. J. Food Microbiol.* **241**, 225–236 (2017).

18. J. R. Gallon, N2 fixation in phototrophs: Adaptation to a specialized way of life. *Plant Soil*. **230**, 39–48 (2001).

19. J. Kramer, Ö. Özkaya, R. Kümmerli, Bacterial siderophores in community and host interactions. *Nat. Rev. Microbiol.* **18**, 152–163 (2020).

20. A. Zelezniak, S. Andrejev, O. Ponomarova, D. R. Mende, P. Bork, K. R. Patil, Metabolic dependencies drive species cooccurrence in diverse microbial communities. *Proc. Natl. Acad. Sci. U. S. A.* **112**, E7156 (2015).

21. O. S. Venturelli, A. V Carr, G. Fisher, R. H. Hsu, R. Lau, B. P. Bowen, S. Hromada, T. Northen, A. P. Arkin, Deciphering microbial interactions in synthetic human gut microbiome communities. *Mol. Syst. Biol.* **14**, 1–19 (2018).

22. D. Bajic, A. Sanchez, The ecology and evolution of microbial metabolic strategies. *Curr. Opin. Biotechnol.* **62**, 123–128 (2020).

23. J. K. Cole, J. R. Hutchison, R. S. Renslow, Y. M. Kim, W. B. Chrisler, H. E. Engelmann, A. C. Dohnalkova, D. Hu, T. O. Metz, J. K. Fredrickson, S. R. Lindemann, Phototrophic biofilm assembly in microbial-mat-derived unicyanobacterial consortia: Model systems for the study of autotroph-heterotroph interactions. *Front. Microbiol.* **5**, 1–18 (2014).

24. A. J. M. Stams, F. A. M. De Bok, C. M. Plugge, M. H. A. Van Eekert, J. Dolfing, G. Schraa, Exocellular electron transfer in anaerobic microbial communities. *Environ. Microbiol.* **8**, 371–382 (2006).

25. P. T. Ha, S. R. Lindemann, L. Shi, A. C. Dohnalkova, J. K. Fredrickson, M. T. Madigan, H. Beyenal, Syntrophic anaerobic photosynthesis via direct interspecies electron transfer. *Nat. Commun.* **8**, 1–7 (2017).

26. T. N. Enke, M. S. Datta, J. Schwartzman, N. Cermak, D. Schmitz, J. Barrere, A. Pascual-García, O. X. Cordero, Modular Assembly of Polysaccharide-Degrading Marine Microbial Communities. *Curr. Biol.* **29**, 1528-1535.e6 (2019).

27. H. J. Flint, K. P. Scott, S. H. Duncan, P. Louis, E. Forano, Microbial degradation of complex carbohydrates in the gut. *Gut Microbes*. **3** (2012), doi:10.4161/gmic.19897.

28. E. A. Bayer, E. Morag, R. Lamed, The cellulosome - a treasure- trove for biotechnology. *Trends Biotechnol.* **12**, 259–265 (1994).

29. C. Belzer, W. Chia, S. Aalvink, B. Chamlagain, V. Piironen, W. M. de Vos, Microbial Metabolic

Networks at the Mucus Layer Lead to Diet-Independent Butyrate and Vitamin B12 Production by Intestinal. *MBio*. **8**, 1–14 (2017).

30. B. Bayer, C. Pelikan, M. J. Bittner, T. Reinthaler, M. Könneke, G. J. Herndl, Proteomic Response of Three Marine Ammonia-Oxidizing Archaea to Hydrogen Peroxide and Their Metabolic Interactions with a Heterotrophic Alphaproteobacterium, 1–15 (2019).

31. J. J. Morris, Z. I. Johnson, M. J. Szul, M. Keller, E. R. Zinser, Dependence of the cyanobacterium Prochlorococcus on hydrogen peroxide scavenging microbes for growth at the ocean's surface. *PLoS One*. **6** (2011), doi:10.1371/journal.pone.0016805.

32. E. E. Lilja, D. R. Johnson, Segregating metabolic processes into different microbial cells accelerates the consumption of inhibitory substrates. *ISME J.* **10**, 1568–1578 (2016).

33. R. P. Kiene, L. J. Linn, J. A. Bruton, New and important roles for DMSP in marine microbial communities. **43**, 209–224 (2000).

34. Y. Morono, T. Terada, M. Nishizawa, M. Ito, F. Hillion, N. Takahata, Y. Sano, Carbon and nitrogen assimilation in deep subsea fl oor microbial cells. **108** (2011), doi:10.1073/pnas.1107763108.

35. Y. Yesiltepe, B. R. Morton, A. B. Cory, H. C. Cardamone, K. S. Hofmockel, M. S. Lipton, J. J. Moran, R. S. Renslow, J. K. Fredrickson, R. Lindemann, Nitrogen Source Governs Community Carbon Metabolism in a Model Hypersaline Benthic Phototrophic Biofil. *Msystems*, 1–12 (2020).

36. M. G. Peter, Applications and Environmental Aspects of Chitin and Chitosan. *J. Macromol. Sci. Part A*. **32**, 629–640 (1995).

37. T. Fuhrer, D. Heer, B. Begemann, N. Zamboni, High-throughput, accurate mass metabolome profiling of cellular extracts by flow injection-time-of-flight mass spectrometry. *Anal. Chem.* **83**, 7074–7080 (2011).

38. S. Blasche, Y. Kim, R. A. T. Mars, D. Machado, M. Maansson, E. Kafkia, A. Milanese, G. Zeller, B. Teusink, J. Nielsen, V. Benes, R. Neves, U. Sauer, K. R. Patil, Metabolic cooperation and spatiotemporal niche partitioning in a kefir microbial community. *Nat. Microbiol.* **6**, 196–208 (2021).

39. A. Goyal, V. Dubinkina, S. Maslov, Multiple stable states in microbial communities explained by the stable marriage problem. *ISME J.* **12**, 2823–2834 (2018).

40. S. Estrela, J. C. C. Vila, N. Lu, D. Bajić, M. Rebolleda-Gómez, C.-Y. Chang, J. E. Goldford, A. Sanchez-Gorostiaga, Á. Sánchez, Functional attractors in microbial community assembly. *Cell Syst.*, 1–14 (2021).

41. J. C. Liao, L. Mi, S. Pontrelli, S. Luo, Fuelling the future: Microbial engineering for the production of sustainable biofuels. *Nat. Rev. Microbiol.* **14**, 288–304 (2016).

42. B. J. Callahan, P. J. Mcmurdie, M. J. Rosen, A. W. Han, A. J. A. Johnson, S. P. Holmes, DADA2 : High-resolution sample inference from Illumina amplicon data. **13** (2016), doi:10.1038/nmeth.3869.

43. T. Madden, The BLAST sequence analysis tool in The NCBI Handbook [Internet]. 2nd edition. *Natl. Cent. Biotechnol. Inf.* (2013).

44. E. S. Wright, Using DECIPHER v2 . 0 to Analyze Big Biological Sequence Data in R. **XX**, 1–8 (2016).

45. D. Hyatt, G. Chen, P. F. Locascio, M. L. Land, F. W. Larimer, L. J. Hauser, Prodigal : prokaryotic gene recognition and translation initiation site identification (2010).

46. S. R. Eddy, Accelerated Profile HMM Searches. **7** (2011), doi:10.1371/journal.pcbi.1002195.

47. J. Mistry, S. Chuguransky, L. Williams, M. Qureshi, G. A. Salazar, E. L. L. Sonnhammer, S. C. E. Tosatto, L. Paladin, S. Raj, L. J. Richardson, R. D. Finn, A. Bateman, Pfam : The protein families database in 2021. **49**, 412–419 (2021).

48. D. H. Haft, B. J. Loftus, D. L. Richardson, F. Yang, J. A. Eisen, I. T. Paulsen, O. White, TIGRFAMs : a protein family resource for the functional identification of proteins. **29**, 41–43 (2001).

49. K. L. Meibom, X. B. Li, A. T. Nielsen, C. Wu, S. Roseman, G. K. Schoolnik, The Vibrio cholerae chitin utilization program. **101**, 2524–2529 (2004).

50. S. Eisenbeis, S. Lohmiller, M. Valdebenito, S. Leicht, V. Braun, NagA-Dependent Uptake of N - Acetyl-Glucosamine and N -Acetyl-Chitin Oligosaccharides across the Outer Membrane of Caulobacter crescentus ▨. **190**, 5230–5238 (2008).

51. T. U. Consortium, UniProt : a worldwide hub of protein knowledge. **47**, 506–515 (2019).

52. K. Katoh, D. M. Standley, MAFFT Multiple Sequence Alignment Software Version 7 : Improvements in Performance and Usability Article Fast Track. **30**, 772–780 (2013).

53. F. Madeira, Y. Park, J. Lee, N. Buso, T. Gur, N. Madhusoodanan, P. Basutkar, A. R. N. Tivey, S. C. Potter, D. Finn, R. Lopez, The EMBL-EBI search and sequence analysis tools APIs in 2019 F abio. **47**, 636–641 (2019).

54. S. Pontrelli, U. Sauer, Salt-Tolerant Metabolomics for Exometabolomic Measurements of Marine Bacterial Isolates. *Anal. Chem.* **93**, 7164–7171 (2021).