Effect of a low water concentration in chloride, sodium and potassium on oocyte maturation, oocyte hydration, ovulation and egg quality in rainbow trout.

Emilien Segret[1,2*], Emilie Cardona[1,3], Sandrine Skiba-Cassy[3], Frédéric Cachelou[2], Julien Bobe[1*].

[1] INRAE, LPGP, Fish Physiology and Genomics, F-35000 Rennes, France.

[2] Viviers de Sarrance, 64490 Sarrance, France.

[3] INRAE, Univ. Pau & Pays Adour, E2S UPPA, NUMEA, 64310, Saint-Pée-sur-Nivelle, France.

*corresponding authors: Emilien Segret : emilien.segret@inrae.fr

                              Julien Bobe : julien.bobe@inrae.fr

Key words: ultrasound staging; non-invasive phenotyping; egg viability; VisEgg; ion and water transport.

Highlights:

-   **Low water salinity impairs final oocyte maturation and egg quality in rainbow trout**
-   **Low water salinity induces delayed ovulation and impaired oocyte hydration**
-   **Low water salinity induces a dysregulation of several key ovulatory genes**
-   **Monitoring of final oocyte maturation can be performed using ultrasound staging**

Abstract

Water salinity is an important environmental factor known to have detrimental effects on salmonid reproduction, mostly when migrating female broodfish are held in sea water. In contrast, data obtained in freshwater are scarce and the impact of low water salinity, and more specifically of low water concentrations in sodium, chloride and potassium, during reproduction in freshwater is currently unknown. For this reason, and because ion and water fluxes are critical for the final steps of the female gamete formation, including oocyte hydration and ovulation, the aim of the present study was to investigate the impact of low salinity water on final oocyte maturation, ovulation and, ultimately, on egg quality, using rainbow trout as a physiological model and relevant aquaculture species.

Fish from the same commercial strain were raised throughout their lifecycle either in a site characterized by low concentrations of Na+, K+, and Cl- ions in the water or in a closely located control site exhibiting higher concentration in these elements. Egg quality and duration of final oocyte maturation were investigated using innovative phenotyping tools such as automatic assessment of egg viability using the VisEgg system and non-invasive echograph-based monitoring of final oocyte maturation duration, respectively. Oocyte hydration during final oocyte maturation and after ovulation was also investigated. Finally, molecular phenotyping was performed using real-time PCR-based monitoring of several key players of final oocyte maturation and ovulation associated with ion and water transport, inflammation, proteolytic activity, and coagulation. Oocyte hydration and gene expression data were analyzed in the light of the duration of final oocyte maturation.

Here we show that low water salinity (i.e., low water concentration in chloride, sodium and potassium) negatively influences final oocyte maturation, ovulation and, ultimately, egg quality. Low water salinity triggered delayed ovulation and lower oocyte viability. When investigating the impact of low water salinity on final oocyte maturation duration, individuals presenting the most severe phenotypes

exhibited impaired oocyte hydration and abnormally reduced gene expression levels of several key players of the ovulatory process. While the under expression of water (e.g., aquaporins) and ion (e.g., solute carriers) transporters is consistent with impaired oocyte hydration, our observations also indicate that the overall ovulatory gene expression program is disrupted. Our results raise the question of the mechanisms underlying the negative influence of low salinity water on the dynamics of the preovulatory phase, on the control of the oocyte homeostasis, including hydration, and on the overall success of the maturation-ovulation process.

1 Introduction

Fish egg quality is defined as the egg ability to be fertilized and to subsequently develop into a normal embryo (Bobe and Labbé, 2010). Egg quality can be highly variable in fish under both natural and aquaculture conditions. In most species, a lack of knowledge of mechanisms influencing egg quality is a limiting factor for the mass production of fry (Migaud et al. 2013). Many external factors such as broodstock nutrition, temperature and photoperiod, to name just a few, are known to have a major impact on egg quality in various fish species (Aegerter and Jalabert, 2004; Bonnet et al., 2007; Carrillo et al., 1989; Izquierdo et al., 2001; Pankhurst et al., 1996). In contrast, other factors such as composition and physico-chemical parameters of the water have received far less attention (Brooks et al., 1997; Bobe and Labbe, 2010). Ionic composition of the water is however likely to influence the final steps of oogenesis (i.e., final oocyte maturation - FOM) given the importance of water exchange during this critical period (Dolomatov, 2012). Among mineral elements, chloride, sodium and potassium are of specific interest because (i) they are the most abundant electrolytes in the body of living organisms and (ii) they serve a vital function in controlling osmotic pressure and acid-base equilibrium (Lall, 2003). It is commonly accepted that deficiencies in chloride, sodium and potassium are difficult to trigger because fish derive these mineral elements from surrounding water (Lall, 2003). To our knowledge, the impact of a low ionic composition in these three key mineral elements on female reproduction was never investigated.

Several concomitant, yet distinct, events take place in the full-grown ovarian follicle during FOM. The follicle-enclosed oocyte progressively acquires the ability to resume meiosis whereas the follicle is preparing the release of oocyte from surrounding somatic layers at ovulation (Grier et al., 2007; Lubzens et al., 2010). In fish, as in other vertebrates, the mechanisms involved in the ovulatory process are described as inflammatory-like (Espey, 1994; Thibault and Levasseur, 1988). In contrast to meiosis resumption and ovulation that are common to all vertebrates, FOM in fish is associated with a significant hydration of the oocyte. This phenomenon results in a dramatic increase of oocyte volume in marine species and is triggered by water entry due to the increase in free amino acid content

resulting from yolk protein processing (Finn et al., 2002). In the saltmarsh species, *Fundulus heteroclitus*, the influx in potassium and sodium is a major cause of the uptake of osmotically obligated water and subsequent volume increase experienced by maturing oocytes (Wallace et al., 1992). In fresh water species, a more limited, yet significant, increase in water content is observed during FOM that could contribute mechanically to the ovulatory process, ultimately leading to oocyte release from surrounding somatic follicular layers (Craik and Harvey, 1984; Milla et al., 2006).

Rainbow trout (*Oncorhynchus mykiss*) is a major aquaculture species that produce eggs every year starting at age 2 (Bromage et al., 1992). In this species oocyte maturation, ovulation (Cerdà et al., 2007; Grier et al., 2007; Jalabert and Fostier, 1984 ; Lubzens et al., 2010), and egg quality (Bobe, 2015; Bobe and Labbé, 2010; Bromage and Jones, 1992; Brooks et al., 1997) have been extensively studied, including at the molecular level (Bobe et al., 2004, 2006, 2009; Yamashita et al., 2000). As previously documented, exposition to stress factors (e.g., temperature) during FOM can alter the quality of unfertilized eggs released from the ovary at ovulation (Aegerter and Jalabert, 2004; Bobe and Labbé, 2010; Colson et al., 2019).

For these reasons, and because water and ion fluxes are important for several key steps of oocyte development during FOM, the aim of the present study was to investigate the impact of water ionic composition, and more specifically low concentrations of sodium, chloride and potassium on FOM, ovulation and, ultimately, on egg quality, using rainbow trout as a physiological model and relevant aquaculture species. To achieve this objective, we used two closely located production sites in the French Pyrenees where the stream water exhibits dramatically different concentrations of chloride, sodium and potassium all year round, including during the reproductive period.

2 Material and methods

2.1 Ethical statement

Investigations were conducted in compliance with EU Directive 2010/63/EU for animal experiments. The experimental protocol registered under the reference #201904021747602 was specifically

approved by INRAE-LPGP Animal Care and Use Committee. For oocyte sampling, echograph-based monitoring and ovulation detection, fish were anaesthetized in tricaine metanesulfonate (MS222) at a concentration of 130 mg/liter. For gonads sampling fish were euthanized in MS222 at a concentration of 300mg/liter, until 10 minutes after cessation of opercular movement.

2.2   Experimental design and water characterization

Two experimental sites located in Rébénacq [43.155948,-0.405067] and Sarrance [43.049999,-0.6014075] (French Pyrenees), and presenting similar water temperature [9°-10°C] and oxygen conditions were used. The two sites are closely located (<20 km) but located in different valleys (i.e., hydrogeologic systems) differing mostly by ionic abundance in Sodium (Na+), Chloride (Cl-) and Potassium (K+) in the water (Fig 1.A , Supplemental datafiles 1 and 2). The site presenting significantly lower abundance of these 3 mineral elements will be referred to hereafter as "Low Salinity Water (LSW)", whereas the other site, for which water ionic composition was significantly higher, will be referred to as "Control". Water composition analyses were performed monthly throughout the year, for both sites. Samples were analyzed by a specialized analytical laboratory (Ultra Traces Analyses Aquitaine, Pau, France). Ionic abundance analysis was performed using inductively coupled plasma atomic emission spectroscopy (ICP-AES) technique (for Na+ and K+) and by ionic chromatography (for Cl-). Data obtained were not normally-distributed (Shapiro-Wilk test, $p > 0.005$). Comparison of two groups were performed using non parametric U-tests. A comprehensive water analysis of both sites is available in supplemental data files 1 and 2. Approximately two weeks before expected ovulation (estimated based on previous ovulation dates), fish were transferred into an 8 m^3 raceway to facilitate manipulations. Experimental raceways on both sites were of similar dimensions (8 m in length, 1 m in width and 1 m in depth) with a water renewal of 200% per hour. Ovulation was monitored three time a week. On the control site, a total of 32 fish were used including 18 for egg quality assessment using the VisEgg system (Cardona et al. 2020) (4 of them being also used for the oocyte hydration study and 11 of them being also used for blood sampling) and 14 for the molecular analysis. On the LSW site, a

total of 46 fish were used including 30 for egg quality assessment using the VisEgg system (4 of them being also used for the oocyte hydration study and 9 of them being also used for blood sampling) and 16 for the molecular analysis.

2.3   Animal husbandry

Three years old female rainbow trout (*Oncorhynchus mykiss*) from the same commercial strain (Viviers SA Company, France) were used in this study. After rearing in a nursery site, fish were transferred to either Rébénacq (control site) or Sarrance (LSW site) locations at 6 months post-fertilization. On both sites, fish were reared under natural photoperiod in $125m^3$ raceways at a $60kg.m^{-3}$ density, with a water renewal of 75% per hour. Similar feeding protocols, including feeding rate and diet composition, were applied for both sites throughout their lifecycle. Fish were fed at an identical daily rate (0.6% of the live weight) with the same commercial diet (46% crude protein, 19% crude fat, 0.9% phosphorus, 11.5% ash, 19 MJ/kg gross energy, Omega-3 4.8%, Omega-6 1.25%) (Le Gouessant, Lamballe, France). Following a first reproduction at age 2, fish were held under natural photoperiod during their second reproductive cycle. Spawning naturally occurred during November-December. Feeding was stopped when first ovulations were detected.

2.4   Egg quality assessment using VisEgg

In both groups, specific fish (control, n=18; LSW, n=30) were individually monitored until ovulation. Ovulations were checked every 2-3 days, under anesthesia, by gentle pressure on the abdomen. Oocyte viability evaluation was systematically performed using the VisEgg protocol when ovulation occurred. This recently developed tool allows a robust assessment of different egg features including egg viability (Cardona et al., 2020). The process uses a standardized image-base batch analysis method of 24h-hydrated unfertilized eggs and an image processing algorithm. Outputs are the presence and percentage of whitening (i.e., non-viable) eggs within the batch. These features can be used to reliably assess egg integrity and viability. Analyze of presence of white eggs depending on rearing sites or the maturation duration yielded contingency tables analyzed using Khi-2 tests. Data obtained from VisEgg

were not normally distributed (Shapiro-Wilk test, p < 0.001). Comparison of two groups were performed using non parametric U-tests.

2.5 Oocyte development staging during FOM

A non-invasive to the female ultrasound-based staging was used to estimate progress into FOM. This approach, previously described for assessing reproductive stages in wild salmons (Nevoux et al., 2019), was used here for the first time to assess progression into FOM in rainbow trout. An ultrasound scanner M-Turbo (Sonosite) with a 5-10MHz linear transducer (multiple frequency) was used. Anesthetized fish were placed in a plastic tray for a short duration (<30 sec) and sagittal images of the abdominal cavity were obtained by aligning the transducer with the lateral line of the fish. Late vitellogenic oocytes appeared white on the screen, while a dark area progressively appeared at the center of the oocyte as the oocytes progressed into FOM (Fig 3.A).

In order to validate echograph-based staging, direct brightfield observation was performed. In rainbow trout, late-vitellogenic (LV) oocytes observed with the naked eye appear opaque and no noticeable changes in yolk structure or any sign of meiosis resumption can be observed. Post-vitellogenesis (PV) oocytes exhibit a central yolk clarification as meiosis progressively resumes. Germinal vesicle is still visible and lipid droplets remain relatively small. During oocyte maturation (Mat), germinal vesicle breakdown (GVBD) progressively occurs and oocytes become totally clear with well-defined lipid droplets. The above description of phenotypic changes occurring in the rainbow trout oocyte during FOM was used to validate the echograph-based staging (Grier et al., 2007; Jalabert and Fostier, 1984; Milla et al., 2006; Patiño and Sullivan, 2002; Hurk and Peute, 1979; Wallace and Selman, 1990). In 100% of the cases (n= 57) the direct observation confirmed the echograph-based staging of the oocyte during FOM. Echograph-based oocyte staging was preferred due to its non-invasiveness nature and was thus subsequently used to measure the duration of the oocyte maturation phase (i.e., between fully transparent oocyte (Mat) stage and detected ovulation) in the present study. The duration of the maturation phase was temperature-normalized using the actual temperature recorded on both sites

during experiments and expressed in hours at 10°C in both experimental groups. Comparisons of the two groups were performed using non-parametric U-tests.

2.6 Plasma sampling and analysis

Blood was sampled during oocyte maturation on both sites (control, n=9; LSW, n=11). Blood samples were centrifuged (1000 g, 10 min, 4°C), and plasma were sampled and stored at −20 °C until further analysis. Plasma samples were analyzed by a specialized analytical laboratory (Ultra Traces Analyses Aquitaine, Pau, France). Ionic abundance analysis was performed using inductively coupled plasma atomic emission spectroscopy (ICP-AES) technique (for Na^+ and K^+) and by ionic chromatography (for Cl^-). Comparison of two groups were performed using non parametric U-tests.

2.7 Hydration measurement

Individual fish originating from the LSW site were separated in two groups, based on the duration of the maturation phase [Mat<64H at 10°C (n=10), Mat⩾64H at 10°C (n=21)]. Oocyte hydration was measured (n=4 individuals of each group) and monitored from late vitellogenesis stage until ovulation. The method used has been previously described (Milla et al., 2006). Briefly, gentle manual striping was performed under anesthesia to collect oocytes, from late vitellogenesis stage until ovulation. As previously described (Finet et al., 1988), non-ovulated oocytes were expelled from the follicle layers using two fine tip tweezers. Oocyte development staging was performed under a binocular microscope. A total of 60 to 80 oocytes per female were obtained and rapidly weighted to obtain wet mass (WM). After heating these oocytes 24 h at 105°C in a drying oven (Memmert, Buechenbach, Germany), dry mass (DM) was measured. The water content (WC) was measured from subtracting DM from WM. For each individual, WC was measured (n=4 individuals of each group) and monitored from late vitellogenesis stage until ovulation. Analysis of the increase in water content (WC) in the two experimental groups (Mat <64H ; Mat ⩾ 64H) was performed using non parametric U-test.

2.8   Molecular analysis

2.8.1   Tissue collection

Ovaries from control (n=4 late-vitellogenesis stage and n=10 oocyte maturation stage) and LSW (n=5 late-vitellogenesis stage and n=11 oocyte maturation stage) groups were sampled for RNA extraction. After euthanasia, ovaries were dissected out of the abdominal cavity. Oocyte staging was performed using ultrasound images prior to euthanasia followed by macroscopic validation of oocyte stage under microscope after dissection. Ovarian follicles were deyolked as previously described (Bobe and Goetz, 2000) by pressing the entire tissue between two stainless steel screens while continuously applying ice-cold Cortland medium with a squirt bottle, then immediately frozen in liquid Nitrogen and stored at -80°C until RNA extraction.

2.8.2   Total RNA extraction and reverse transcription

Ovarian tissue was homogenized in Trizol reagent (Molecular Research Center, Cincinnati, USA) at a ratio of 1ml per 100mg of tissue. Tissue was grinded using bead beating technology (Precellys Evolution, Bertin Technologies, Saint-Quentin en Yvelines, France). Total RNA extraction was performed following the manufacturer's instructions. Briefly, 2 µg of total RNA were reverse transcribed using Maxima First Strand cDNA Synthesis Kit (Maxima Reverse Transcriptase enzyme, derived from Moloney murine Leukemia virus enzyme). Reverse transcription was performed by 10 minutes of incubation at 25°C followed by a 15 minutes step at 60°C and a 5-minute step at 75°C. Control reactions were run without Maxima reverse transcriptase and used as negative controls in the real-time polymerase chain reaction study.

2.8.3   Real-time PCR analysis

Real-time PCR was performed using a LightCycler480 (Roche, Switzerland). Reverse transcription products were diluted to 1/30. Technical replicates of each biological sample were run. Runs were performed in 5µL reactions containing 2µL of RT product, 250 nM of each primer and 1X PowerUp SYBR Green Master Mix (Applied Biosystems). The program used was: 50°C for 2 minutes, then 95°C

for 2 minutes for initial denaturation, 40 cycles of 95°C for 3 seconds and 60°C for 30 seconds. After amplification, a fusion curve was obtained by 15-second step followed by a 0.5°C increase, repeated 70 times, and starting at 60°C

2.8.4 Gene expression analysis

Several concomitant events occur in the ovary during FOM that involve distinct biological processes including ovulation, an inflammatory-like process characterized by major tissue rearrangements, blood coagulation, detachment of somatic follicular cells from the oocyte, and oocyte hydration which involves specific water and ion transports. Genes used for the molecular analysis were thus selected for their putative role in inflammation, proteolytic activity, or water and ion transport and were previously reported to be dramatically differentially up-regulated during FOM in rainbow trout (Bobe et al., 2006). Primers used in this prior study (Bobe et al., 2006) were used here to specifically monitor the expression of the following genes. Aquaporin 4 (*aqp4,* GenBank accession # BX885214), Vasotocin-neurophysin VT 1 (*vt1,* GenBank accession # CA375992) and Solute carrier family 26 (*slc26a4,* GenBank accession # BX873066) are involved in ion and water transport processes whereas A Disintegrin And Metalloproteinases 22 (*adam22*, GenBank accession # CA363158), CXC chemokine L14 (*cxcl14,* GenBank accession # BX868653) and coagulation factor V (*f5*, GenBank accession # BX879767) are inflammation-related genes. 18S RNA level was measured in each sample and used for messenger RNA abundance normalization. Expression levels at the late-vitellogenic (LV) stage are not presented but were not significantly different between groups and were arbitrarily set to 1 for each group. Expression data at the oocyte maturation stage were expressed as a percentage of the transcript abundance at the LV stage. Difference between LSW and control groups at oocyte maturation stage were analyzed using non-parametric U-tests.

3   Results and discussion

3.1   Low water salinity results in lower egg viability

The automatic phenotyping of egg viability using the VisEgg system (Fig 2.A) revealed that 60% of the females exhibited non-viable eggs at the time of ovulation in the LSW group in sharp contrast with the control group in which only 18% of the females exhibited at least one non-viable egg at ovulation. More specifically, the percentage of non-viable eggs was dramatically increased (+378%) in the LSW group in comparison to the control group (Fig 2.B). Salinity is known to have an impact on reproduction in fish, including in salmonids. In Atlantic salmon (*Salmo salar*) and Coho salmon (*Oncorhynchus kisutch*), whole-life breeding in freshwater can result in the production of poor quality gametes including poor semen production for males and presence of atretic follicles in females (Valdebenito et al., 2015). Similarly, other studies reported that rearing of Atlantic salmon in sea water induces delays and variability in the timing of ovulation (Haffray, 1995) and poor gamete quality (Maisse et al., 1998). In Arctic charr (*Salvelinus alpinus*), gamete quality is improved for broodstock transferred in saltwater during their gametogenesis (Atse et al., 2002). Together, these studies indicate that water salinity is a key factor for the reproduction success of salmonids that all reproduce in freshwater even though a part of their life cycle occurs in sea water in some species. Most studies have however focused on the impact of high water salinity during final oocyte maturation. In contrast, data on low salinity water remain scarce. In addition, most studies have focused on migrating species that return to freshwater to reproduce. In the present study, we confirm that water salinity is a key factor of reproductive success and show that, similarly to high salinity, low salinity can also have a detrimental effect on reproductive success.

3.2   Low water salinity triggers delayed ovulation

Together, our results indicate that low water concentration in chloride, sodium and potassium triggers a significant decrease in egg viability at ovulation. In order to shed light on underlying mechanisms, we aimed at investigating the preovulatory period that is known to be critical for egg quality. Given the

favorable temperature range (i.e., 9-10°C) and similar feeding regime throughout the reproductive cycle, we reasoned that LSW could perturbate FOM and investigated the duration of the maturation phase from meiosis resumption to ovulation. Using non-invasive echograph-based analysis we could measure the duration of the oocyte maturation phase. After monitoring the first signs of intra-oocyte clearing (i.e., post-vitellogenic stage) (Fig 3.A and B), oocytes became fully transparent (i.e., maturation stage) until they were released from the ovary at ovulation. The duration of the maturation phase was calculated in each group after normalization by the daily temperature on site. We observed (Fig 3.C) that the maturation phase was significantly longer in the LSW group in comparison to the control group. The duration of oocyte maturation was on average 53 hours (at 10°C) in the LSW group while it was 70 hours (at 10°C) in the control group, which corresponds to a 32% increase in FOM duration in the LSW group.

Several studies described the temporal aspects of final oocyte maturation in rainbow trout. The *in vivo* duration of FOM, including progressive central clearing of the oocyte, ranged from 48 to 72 hours at 13°C (Bry, 1981). The detachment of the mature oocyte from surrounding somatic follicular layers was reported to occur between 72 and 96 hours after the first observable signs of oocyte maturation (i.e., central clarification) at 10°C (Jalabert, 1978; Jalabert and Szöllösi, 1975). Other authors showed that ovulation occurs generally 48-96 hours at 14°C after the peak concentration of maturation-inducing steroid (Fostier, 1981). Another study shown a duration of 48-72 hours at 12°C between maturation and detected ovulation for rainbow trout (Breton et al., 1998).

The duration of oocyte maturation, from full oocyte cytoplasm clearing to ovulation, reported here for the control group, is therefore fully consistent with the duration of FOM reported in these prior studies that usually included the post-vitellogenic phase (i.e., germinal vesicle migration towards the periphery of the oocyte). Our observations however suggest that the duration of FOM in the LSW group is higher or in the upper range of what is classically observed in rainbow trout at 10°C. In addition, the LSW group exhibits a strong interindividual variability with individuals that ovulated 97 hours (2 individuals) and 130 hours (1 individual) after detected oocyte maturation (Fig 3.C). These individuals presented

the usual phenotypic traits of maturation (i.e., color pattern and body shape), but ovulation occurred drastically later. Heterogeneity in FOM duration and, in some cases, absence of ovulation has often been reported in trout and other fish. This heterogeneity has however systematically always been associated with premature induction of FOM and ovulation, typically when FOM was triggered in maturationally incompetent (i.e., vitellogenic) follicles *in vivo* or *in vitro*. This is for instance the case for hormonally injected fish (Bry, 1981; Goetz and Bergman, 1978; Jalabert, 1978). Using non-invasive ultrasound-based FOM monitoring, we were able to demonstrate here that oocyte maturation (i.e., oocyte clearing) occurred but that ovulation was delayed. Previous studies have demonstrated that oocyte maturation and ovulation, while being interconnected, were two independent processes. In rainbow trout it is possible to trigger ovulation without oocyte maturation (Jalabert, 1972). Together, our results show that low salinity (i.e., low water concentration in chloride, sodium and potassium) during FOM triggers delayed ovulation, sometimes by several days. Our observations also indicate that, while oocyte maturation occurs normally, the ovulatory process appears impaired or inhibited, ultimately leading to abnormally long FOM.

3.3 Reduced egg viability in low salinity water is linked with delayed ovulation

Together our results show that low water concentration in chloride, sodium and potassium triggered reduced egg viability and delayed ovulation. We also noticed a high inter-individual variability in both egg viability and maturation phase duration in the LSW group (Figs 2.B & 3.C). We reasoned that delayed ovulation could be at the origin of lower egg viability in the LSW group and investigated further any link between egg viability and maturation phase duration. For data analysis, LSW fish were separated in two groups, based on the duration of the maturation phase. We observed a dramatic and highly significant difference in egg viability in the LSW group depending on the duration of the maturation phase (Fig 4). A very high proportion (85%) of the fish exhibiting a maturation phase duration of 64 hours or more exhibited non-viable eggs at ovulation, while only 10% of the fish that underwent oocyte maturation in less than 64 hours exhibited this feature (Fig 4.A). In addition, the

percentage of non-viable eggs was dramatically increased in the ⩾64h group in comparison to the <64h group (Fig 4.B).

These results strongly support our hypothesis of a significant loss of egg quality after delayed ovulation. A possible explanation for the higher rate of non-viable eggs would be the ageing of the oocytes. This well-known phenomenon has been extensively described (Aegerter and Jalabert, 2004; Billard, 1981; Craik and Harvey, 1984; Escaffre, 1976; Lahnsteiner, 2000; Samarin et al., 2015, 2008; Springate et al., 1984) and is known to trigger major egg quality defects. All existing studies on oocyte ageing prior to fertilization were however conducted after post-ovulatory ageing. To our knowledge, oocyte ageing prior to ovulation (i.e., within the ovarian follicle) is reported here for the first time in a fish species. In salmonids, in contrast to other fish species, post-ovulatory ageing does not induce a rapid decrease of egg quality. In rainbow trout, ovulated eggs can be held in the body cavity for 5 days at 12°C without any significant decrease of their developmental success (Aegerter et al., 2005). It is therefore unlikely that the modest intra follicular post-meiotic ageing observed here in the LSW group, in which ovulation is delayed, on average, by 17 hours, is sufficient to trigger alone the drop of egg quality observed in LSW. The osmotic properties of the rainbow trout egg have been thoroughly investigated in rainbow trout (Gray, 1932) and another explanation for the increase in whitening egg occurrence could be a loss of the oocyte homeostasis in LSW ultimately leading to non-viable eggs. In brook trout, a diminution of ionic concentration (Cl-) in surrounding medium has as direct negative effect of membrane resistance (Marshall et al., 1985). Disruption of the oocyte homeostasis could therefore originate from a fragile oocyte vitelline membrane. In addition, the loss in oocyte homeostasis could originate from difficulties (i.e., high energetic cost) in maintaining the osmotic difference across oocyte membrane in a hypotonic external medium.

Together, our results show that the reduced egg viability observed in LSW is linked to delayed ovulation. This reduced egg viability can be due, at least in part, to intra-follicular oocyte ageing. We cannot, however, rule out the possibility that LSW also affects egg viability through other mechanisms that could disrupt the maintenance of oocyte homeostasis and ultimately decrease egg viability.

Further investigations are needed to unravel the mechanisms triggering decreased egg viability when FOM is occurring in LSW.

3.4 Low water concentration in chloride, sodium and potassium results in reduced oocyte hydration during FOM

The release of the oocyte from surrounding follicular layers at the time of ovulation results from the ovulatory process that has been described initially in mammals as an inflammatory-like reaction. In fish, in which the oocyte is directly in contact with surrounding follicular layers, a prerequisite is the separation of the oocyte from surrounding granulosa layers (Jalabert, 1976; Lubzens et al., 2010). This phenomenon has been thoroughly described in rainbow trout in which suppression of tight junctions between granulosa cells and the oocyte results in the apparition of a thin periplasmic space separating the oocyte from surrounding somatic layers (Lubzens et al., 2010). Many observations, including direct brightfield *in vitro* follicle observation, have shown that preovulatory rainbow trout oocyte exhibits a significant mechanical pression onto surrounding somatic layers (Lubzens et al., 2010). Somatic follicular layers appear to be stretched around the oocyte at that stage. Because of the angiotensin-like nature of the ovulatory process and the contractile actions of theca layers during ovulation (Hajnik and Goetz, 1998; Hsu et al., 1992), postovulatory follicles appear shrinked after ovulation. It was hypothesized that the modest (i.e., 20%), yet significant, increase in water content observed during rainbow trout FOM would contribute to the increasing pressure applied by the oocyte onto surrounding somatic layers and would ultimately mechanically facilitate the exit of the oocyte through a hole in somatic layers (Milla et al., 2006). Mechanic pressure applied by the oocyte swelling is one cause of follicle rupture and the completeness of hydration facilitates the complete liberation of unfertilized eggs into the abdominal cavity. We reasoned that a lower hydration would decrease the efficiency of ovulation and investigated hydration in the two above defined sub-groups exhibiting respectively <64h and ⩾64h maturation phase duration. Oocyte water content (WC) was monitored throughout FOM in both groups. In the <64h group, WC progressively increased throughout FOM in

comparison to the late vitellogenic (LV) stage, with a 15% increase during oocyte maturation and a 27% increase at ovulation to reach 36% during the post-ovulatory period (Fig 5, light blue). In contrast, a much more limited increase in water content was observed in the ≥64h group. In comparison to the LV stage, a 2% increase in water content was observed during oocyte maturation while a 13% increase was monitored at ovulation that did not further increase during the post-ovulatory period. Observations in the <64h group are fully consistent with existing data in rainbow trout that were obtained using the same methodology (Milla et al., 2006). In contrast the hydration in the ≥64h appears to be drastically reduced. At all stages, the water increase was significantly lower in the ≥64h group in comparison to the <64h group. Together these results indicate that oocyte hydration is impaired during FOM occurring in low salinity water.

Oocyte hydration has been described for several species (Finn et al., 2002; LaFleur and Thomas, 1991; Milla et al., 2006; Watanabe and Kuo, 1986). Liberation of free amino acids and an uptake of inorganic ions occur during maturation. These events create a change in osmotic equilibrium, resulting in the passive transport of water molecule into the oocyte. This transport mechanism uses several proteins activity such as solute carriers, Na+/K+ ATPases, Aquaporins, and demonstration was made that protein activity and hydration were closely dependent on K+ concentration in the medium (Ecker and Dennis Smith, 1971; LaFleur and Thomas, 1991). K+ and Na+ accumulation in the oocyte during maturation plays an important role in the oocyte hydration process. For *Fundulus heteroclitus*, oocytes Na+ and K+ uptakes by gap junctions (Lubzens et al., 2010) are main responsible of osmolality increase (Greeley et al., 1991; Wallace et al., 1992). For Atlantic halibut *Hippoglossus hippoglossus*, Cl- uptakes also contributes to the increase of oocyte osmolality (Finn et al., 2002). Together these studies demonstrate the importance of Na+, K+ and Cl- ions for the oocyte hydration process during FOM. In contrast, the impact of low water salinity was never investigated, mostly because these experiments were often conducted in marine species. We hypothesize that FOM in low salinity water is associated in reduced ions uptake during maturation ultimately leading to incomplete oocyte hydration and

thereby causing a large variability in the timing of ovulation. Further investigations are needed to elucidate underlying mechanisms.

3.5 Low water concentration in chloride, sodium and potassium induces dysregulated expression of ovulation genes

The ovulatory process involves the activation of a specific gene expression program that has been extensively documented in mammals (Richards et al., 2002) and in fish (Bobe et al., 2009, 2006; Cerdà et al., 2007; Fabra et al, 2005; Hajnik and Goetz, 1998; Knoll-Gellida, 2006). The inflammatory-like nature of the ovulatory process has been well characterized and the expression of many inflammation-related genes have been reported in various fish species (Hagiwara, 2020; Ohnishi, 2005; Takahashi, 2019) including in rainbow trout (Bobe et al., 2006). A subset of these previously identified genes related to inflammation, proteolytic activity, or ion and water transport was monitored to characterize the ovulatory process in low salinity water.

Solute transporters and hydration-related genes are under-expressed in low salinity water

Aquaporin are molecular channels responsible of passive water transport across membranes. The role of aquaporins during fish oocyte hydration has been described by several studies. The up-regulation of *aquaporin 4* (*aqp4*) was reported during rainbow trout follicular maturation (Bobe et al., 2006), suggesting that oocyte hydration of fresh water fish is aquaporin mediated, in consistency with the demonstrated role of aquaporin1-like in oocyte hydration of saltwater fish (Cerdà, 2009; Fabra, 2005).

Solute carrier family 26 member 4 gene (*slc26a4*), previously known as pendrin, codes for a transmembrane anion exchanger and is involved in osmoregulation mechanisms (Nakada et al., 2005). Identified as dramatically up-regulated in the rainbow trout ovary during maturation, slc26a4 is believed to play an important role in the osmotic regulation, causing water influx into oocyte during meiotic maturation. In the present study, a marked under expression of *slc26a4* and *aqp4* is observed in the ovary when FOM occurs in LSW (Fig 6). Given the role of both genes in ion and water transport,

this under expression is fully consistent with the impaired oocyte hydration observed in LSW, in comparison to the control group.

Vasotocin-neurophysin VT 1 protein (*vt1* gene) has two key roles in FOM-ovulation in catfish (Joy and Chaube, 2015). It is an important link in the gonadotropin cascade controlling FOM and ovulation. By a common action with maturation inductions hormones, VT 1 is a regulator of germinal vesicle breakdown and meiosis resumption. VT 1 also plays a key role in follicular hydration by regulating the aquaporin action (Balment et al., 2006). Here we show a marked under expression of *vt1* observed in the LSW group in comparison to the control group (Fig 6). This profile is fully consistent with both the observed impaired ovulatory process and the under expression of *aquaporin 4*. In addition, the position of *vt1* in the gonadotropin-dependent regulatory cascade controlling oocyte maturation and ovulation suggests that it could be a relay in mediating the inhibitory effect of LSW.

Inflammation-related genes are under-expressed in low salinity water

Ovulation is accompanied by a broad-spectrum proteolysis within follicular layers that implies a large number of proteases. Members of A Disintegrin and Metalloproteinase gene family are dramatically up-regulated during trout FOM-ovulation phases (Berndtson and Goetz, 1988; 1990; Bobe et al., 2006). *Adam22* plays an important role in cell-to-cell and cell-to-matrix interaction regulation (White, 2003). By regulating or processing the deposit of the collagen fibrils, ADAMs proteases cause the extra-cellular matrix proteolysis (Ohnishi et al., 2005).

CXC chemokines stimulate recruitment of leukocytes and play a role of pro-inflammatory mediators. *Cxcl14* implication in immune–related functions has been described in fish (Baoprasertkul et al., 2005). The over-expression of *cxcl14* during FOM-ovulation phases is fully in line with the inflammatory-like nature of the ovulatory process.

Coagulation factor V (*f5* gene) is involved in synthesis of blood coagulant factors and has been characterized in mammals and fish (Hanumanthaiah et al., 2002; Rothberger, 1984). It was speculated

that synthesis of coagulation factor V prevents bleeding caused by rupture of follicle layers during ovulation.

Here we show that *cxcl14*, *adam22*, and *f5* are dramatically under expressed during oocyte maturation in the LSW group in comparison to the control group (Fig 6). These expression profiles are consistent with the delayed ovulation observed in LSW and suggest that the inflammatory nature of ovulation is affected. This is especially true for *cxcl14* for which a 5-fold decrease in gene expression is observed. Together, these results show that several important components of the ovulatory process are toned down when FOM occurs in low salinity water. It is noteworthy that most, if not all, aspects of the ovulatory process including inflammation, coagulation, swelling, water and ion transport as well and proteolysis are under activated. Even if previous studies have shown gene expression changes when fish adapt to water salinity change (e.g. migration, precipitation) for salmonids (Morro, 2020; Norman et al., 2014) and other species (Fiol, 2006; Pujante, 2018), evidence of such an impact of low water salinity on the overall ovulatory process were, to our knowledge, previously unreported. This raises the question of how low water salinity can inhibit the overall ovulatory process and not only osmotic exchanges.

3.6 Lower blood plasma levels of sodium, chloride and potassium are observed in the LSW site

In order to assess the impact of water concentration in chloride, sodium and potassium on fish homeostasis, we measured blood plasma concentration in these three elements in both control and LSW sites. We observed a significantly lower plasmatic concentration of these three mineral elements in fish of the LSW site in comparison to the control site (Fig 1.B). This indicates that the water composition can lead to significant differences in plasma concentration in these elements. This also further supports our hypothesis that reproduction in fresh water exhibiting low sodium, chloride and potassium concentration perturbates the fish oocyte osmolarity and the hydration process during final oocyte maturation and ovulation.

3.7 Role of other water components

While marked and highly significant differences in water concentration of chloride, sodium and potassium are found between the two experimental sites, other differences exist in water composition that could also contribute, at least in part, to the impaired FOM and ovulation and reduced egg quality phenotype observed in the LSW site. A lower concentration in Iron (Fe) and Manganese (Mn) is observed in the LSW site in comparison to the control site. For Iron, the difference is however not significant when only the reproductive period is considered (Supplemental datafile 2). Manganese is known to inhibit ovulation in rainbow trout (Jalabert and Szollosi, 1975) and inhibit meiotic maturation in terrestrial animals (Bilodeau-Goeseels, 2001). It is therefore difficult to link low Mn concentration with impaired FOM and ovulation. In addition, it is commonly accepted that the diet is considered to be the major source of Manganese and Iron (Lall, 2002). It is therefore extremely unlikely that the lower water concentration in Mn and Fe found in the LSW site is responsible for the impaired reproductive phenotype observed in the LSW site. Similarly, a lower nitrate concentration was observed in the LSW site. While we cannot rule out that this could contribute to the abnormal reproductive phenotype observed in the LSW site, this seems however, to our knowledge and in the light of previous studies led on salmonids and others fish species (Dolomatov, 2011; Tripathi and Krishna, 2008), extremely unlikely.

4 Conclusion

Here we show that low water concentration in chloride, sodium and potassium negatively influences final oocyte maturation, ovulation and, ultimately, egg quality. Low salinity water triggers delayed ovulation and lower oocyte viability. When investigating the impact of LSW on FOM duration, individuals presenting the most severe phenotypes exhibited impaired oocyte hydration and abnormally reduced gene expression levels of several key players of the ovulatory process. While the under expression of water (e.g., aquaporins) and ion (e.g., solute carriers) transporters is consistent with impaired oocyte hydration, our observations also indicate that the entire ovulatory gene

expression sequence is disrupted. Our results raise the question of the mechanisms underlying the negative influence of low water concentration in chloride, sodium and potassium on the dynamics of the preovulatory phase, on the control of the oocyte homeostasis, including hydration, and on the overall success of the maturation-ovulatory process. A global analysis of differentially regulated genes in LSW during final oocyte maturation is needed to unravel these mechanisms.

5 Declarations

5.1 Competing interests

The authors declare no competing interest.

5.2 Authors' contributions

ES conceived the study, performed experiments and data analysis and drafted the manuscript. EC participated in experiments. SS-K and FC participated in the design of the study. JB conceived the study and participated in manuscript writing and data analysis.

5.3 Funding

This work was supported by the National Association for Technologic Research (ANRT) [grant number 2018/0457]; the Nouvelle-Aquitaine regional council [grant number 3379120].
Funding sources have no involvement in the conduct of the research.

5.4 Acknowledgements

The authors thank Floriane Colon and Viviers Company staff for their help in fish rearing. The authors thank the Nouvelle-Aquitaine regional council and ANRT for the financial support which allowed the study.

6 Figure legends

Color should be used for all figures in print.

Figure 1: Water and plasma composition on both experimental sites

A: Distribution of Potassium, Chloride and Sodium concentration (mg/L) in Control (in grey, n=12) and Low Salinity Water (in blue, n=12) sites.

B: Distribution of Potassium, Chloride and Sodium concentration (g/L) in fish blood plasma reared in Control (in grey, n=11) and Low Salinity Water (in blue, n=9) sites.

Figure 2: Egg quality in the two experimental sites

A: Occurrence of non-viable eggs in batches evaluated using the VisEgg system in Control (n=18) and LSW (n=30) groups. Khi-2 test (p = 0.007).

B: Rate (percent) of non-viable eggs evaluated using the VisEgg system in Control (n=18) and LSW (n=30) groups.

Figure 3: Ultrasound staging for measure of maturation duration

A: Echograph-based observation of oocyte changes during final oocyte maturation. Changes in oocyte aspect from LV to MAT is illustrated by a progressive central darkening, with brightening of peripheral layer before ovulation.

B: Pictures of direct observation of oocytes classified by stage. Oocyte were obtained by stripping and defolliculation (Greeley et al., 1991). Yolk clarification and lipid droplet formation are observable.

C: Distribution of maturation duration evaluated by ultrasound in Control (n=18) and LSW (n=30) from central oocyte clarification and ovulation.

Figure 4: Egg quality depending on the duration of oocyte maturation

A: Percentage of females exhibiting non-viable eggs in low salinity water, evaluated using the VisEgg system. Individuals are separated in two groups depending on the duration of oocyte maturation (Mat<64H, n=10; Mat⩾64H, n=20). Khi-2 test (p = 1.34E-04).

B: Percentage of non-viable eggs depending on the duration of oocyte maturation (Mat<64H, n=10; Mat⩾64H, n=20).

Figure 5: Hydration of oocytes during final oocyte maturation

Increase of oocyte water content (WC), in comparison to late vitellogenic stage, for oocytes sampled at four ovarian stages: Late-Vitellogenesis (LV), oocyte maturation (Mat), ovulation (Ov) and 3 days post-ovulation (Post-Ov). Individuals are separated in two groups based on the duration of oocyte maturation (Mat<64H, n=4; Mat⩾64H, n=4).

Figure 6: Ovarian expression profiles of genes involved in FOM and ovulation

Ovarian expression profiles of *aqp4*, *slc26*, *cf5*, *adam22*, *cxcl14* and *vt5* during oocyte maturation in control (n = 10) and low salinity water (LSW, n = 11) groups. Messenger RNA abundance was measured using real-time PCR and normalized using the abundance of 18S. Abundance was arbitrarily set to 1 in the LV stage in each group.

7 Additional Files

	CONTROL		LSW		
	Mean	**SD**	**Mean**	**SD**	
Conductivity (µS/cm)	**264,00**	41,64	**248,92**	34,07	
Major ions, in milligrams per liter					
Bicarbonate	**145,50**	17,77	**140,00**	17,04	
Calcium	**44,79**	8,57	**43,60**	5,78	
Chloride	**2,29**	0,84	**1,24**	0,39	*
Magnesium	**4,29**	0,51	**5,16**	2,41	
Potassium	**0,47**	0,13	**0,21**	0,09	*
Sodium	**1,83**	0,28	**1,09**	0,27	*
Sulfur	**3,03**	0,79	**5,08**	3,32	*
Sulfate	**8,33**	2,06	**13,05**	9,54	
Nutrients, in milligrams per liter					
Nitrate	**2,92**	1,28	**1,03**	0,27	*
Trace elements, in micrograms per liter					
Copper	**0,22**	0,09	**0,14**	0,07	
Iron	**14,99**	14,12	**3,00**	4,58	*
Manganese	**1,47**	1,28	**0,15**	0,18	*
Selenium	**0,13**	0,02	**0,15**	0,05	
Iode	**1,35**	0,53	**1,04**	0,18	
Zinc	**1,37**	0,70	**1,35**	0,83	

Supplemental data file 1: Water analysis in both control (Rébénacq) and LSW (Sarrance) sites.

Mean (N= 12) yearly (2020-2021) values and standard deviation are shown. Difference between LSW and control water were analyzed using non-parametric U-tests.

* denotes significant differences.

	CONTROL		LSW		
	Mean	**SD**	**Mean**	**SD**	
Conductivity (µS/cm)	**282.00**	20.785	**242.67**	29.48	
Major ions, in milligrams per liter					
Bicarbonate	**154.50**	25.05	**143.67**	10.97	
Calcium	**48.97**	5.11	**42.43**	4.66	
Chloride	**2.98**	0.45	**1.47**	0.06	*
Magnesium	**4.59**	0.32	**3.51**	2.66	

Potassium	**0.60**	0.19	**0.18**	0.04	*
Sodium	**2.02**	0.17	**0.99**	0.18	*
Sulfur	**3.60**	0.09	**3.10**	2.82	
Sulfate	**9.70**	0.65	**9.5**	8.37	
Nutrients, in milligrams per liter					
Nitrate	**4.09**	1.79	**1.11**	0.09	*
Trace elements, in micrograms per liter					
Copper	**0.26**	0.15	**0.16**	0.13	
Iron	**8.07**	6.72	**2.00**	0.44	
Manganese	**0.70**	0.33	**0.12**	0.07	*
Selenium	**0.13**	0.01	**0.17**	-	
Iode	**1.45**	0.50	**1.15**	0.08	
Zinc	**1.26**	0.56	**1.44**	0.54	

Supplemental data file 2: Water analysis in both control (Rébénacq) and LSW (Sarrance) site during the reproductive period.

Mean (N= 3) values (from 2020-10-15 to 2020-12-15) and standard deviation are shown. Difference between LSW and control water were analyzed using non-parametric U-tests.

* denotes significant differences.

8 References

Aegerter, S., Jalabert, B., 2004. Effects of post-ovulatory oocyte ageing and temperature on egg quality and on the occurrence of triploid fry in rainbow trout, Oncorhynchus mykiss. Aquaculture 231, 59–71. https://doi.org/10.1016/j.aquaculture.2003.08.019

Aegerter, S., Jalabert, B., & Bobe, J., 2005. Large scale real-time PCR analysis of mRNA abundance in rainbow trout eggs in relationship with egg quality and post-ovulatory ageing. Molecular Reproduction and Development: Incorporating Gamete Research, 72(3), 377-385. https://doi.org/10.1002/mrd.20361

Atse, C.B., Audet, C., De La Noüe, J., 2002. Effects of temperature and salinity on the reproductive success of Arctic charr, Salvelinus alpinus (L.): Egg composition, milt characteristics and fry survival. Aquac. Res. 33, 299–309. https://doi.org/10.1046/j.1355-557x.2002.00674.x

Balment, R.J., Lu, W., Weybourne, E., Warne, J.M., 2006. Arginine vasotocin a key hormone in fish physiology and behaviour : A review with insights from mammalian models. General and comparative endocrinology. 147, 9–16. https://doi.org/10.1016/j.ygcen.2005.12.022

Baoprasertkul, P., He, C., Peatman, E., Zhang, S., Li, P., Liu, Z., 2005. Constitutive expression of three novel catfish CXC chemokines: Homeostatic chemokines in teleost fish. Mol. Immunol. 42, 1355–1366. https://doi.org/10.1016/j.molimm.2004.12.012

Berndtson, K., Goetz, W., 1988. Protease activity in brook trout (Salvelinus fontalis) follicle walls demonstrated by substrate-polyacrylamide gel electrophoresis. Biol. Reprod. 38 (2), 511–516. https://doi.org/10.1095/biolreprod38.2.511

Berndtson, A.K., Goetz, F.W., 1990. Metallo-protease activity increases prior to ovulation in brook trout (Salvelinus fontinalis) and yellow perch (Perca flavescens) follicle walls. Biol. Reprod. 42, 391–398

Billard, R., Bry, C., Gillet, C., 1981. Stress, environment and reproduction in teleost fish. Stress and fish, Academic Press, 0-12-554550-9. hal-01600274

Bilodeau-Goeseels, S., 2001. Manganese inhibits spontaneous nuclear maturation in bovine cumulus-enclosed oocytes. Can. J. Anim. Sci. 81, 223–228. https://doi.org/10.4141/A00-074

Bobe, J., 2015. Egg quality in fish: Present and future challenges. Anim. Front. 5, 66–72. https://doi.org/10.2527/af.2015-0010

Bobe, J., Labbé, C., 2010. Egg and sperm quality in fish. Gen. Comp. Endocrinol. 165, 535–548. https://doi.org/10.1016/j.ygcen.2009.02.011

Bobe, J., Montfort, J., Nguyen, T., Fostier, A., 2006. Identification of new participants in the rainbow trout (Oncorhynchus mykiss) oocyte maturation and ovulation processes using cDNA microarrays. Reprod. Biol. Endocrinol. 4, 1–16. https://doi.org/10.1186/1477-7827-4-39

Bobe, J., Nguyen, T., Fostier, A., 2009. Ovarian function of the trout preovulatory ovary: New insights from recent gene expression studies. Comp. Biochem. Physiol. - A Mol. Integr. Physiol. 153, 63–68. https://doi.org/10.1016/j.cbpa.2008.10.021

Bobe, J., Nguyen, T., Jalabert, B., 2004. Targeted gene expression profiling in the rainbow trout (Oncorhynchus mykiss) ovary during maturational competence acquisition and oocyte maturation. Biol. Reprod. 71, 73–82. https://doi.org/10.1095/biolreprod.103.025205

Bobe, J., Goetz, F.W., 2000. A tumor necrosis factor decoy receptor homologue is up-regulated in the brook trout (Salvelinus fontinalis) ovary at the completion of ovulation. Biol Reprod. 62(2):420-6. https://doi.org/10.1095/biolreprod62.2.420

Bonnet, E., Montfort, J., Esquerre, D., Hugot, K., Fostier, A., Bobe, J., 2007. Effect of photoperiod manipulation on rainbow trout (Oncorhynchus mykiss) egg quality: A genomic study. Aquaculture 268, 13–22. https://doi.org/10.1016/j.aquaculture.2007.04.027

Breton, B., Govoroun, M., Mikolajczyk, T., 1998. GTH I and GTH II secretion profiles during the reproductive cycle in female rainbow trout: Relationship with pituitary responsiveness to GnRH-a stimulation. Gen. Comp. Endocrinol. 111, 38–50. https://doi.org/10.1006/gcen.1998.7088

Bromage, N., Jones, J., 1992. The effects of seasonal alterations in ration on fecundity and maturation in rainbow trout (oncorhynchus mykiss). Proc. Br. Soc. Anim. Prod. 1992, 81–81. https://doi.org/10.1017/s03082296000219422

Bromage, N., Jones, J., Randall, C., Thrush, M., Davies, B., Springate, J., Duston, J., Barker, G., 1992. Broodstock management, fecundity, egg quality and the timing of egg production in the rainbow trout (Oncorhynchus mykiss). Aquaculture 100, 141–166. https://doi.org/10.1016/0044-8486(92)90355-O

Brooks, S., Tyler, C.R., Sumpter, J.P., 1997. Egg quality in fish: What makes a good egg? Rev. Fish Biol. Fish. 7, 387–416. https://doi.org/10.1023/A:1018400130692

Bry, C., 1981. Temporal aspects of macroscopic changes in rainbow trout (Salmo gairdneri) oocytes before ovulation and of ova fertility during the post-ovulation period; effect of treatment with 17α-hydroxy-20β-dihydroprogesterone. Aquaculture 24, 153–160. https://doi.org/10.1016/0044-8486(81)90052-1

Cardona E., Bugeon J., Segret E., Bobe J., 2020. VisEgg: a robust phenotyping tool to assess rainbow trout egg features and viability. Fish Physiol Biochem. Epub ahead of print. https://doi.org/10.1007/s10695-020-00844-2

Carrillo, M., Bromage, N., Zanuy, S., Serrano, R., Prat, F., 1989. The effect of modifications in photoperiod on spawning time, ovarian development and egg quality in the sea bass (Dicentrarchus labrax L.). Aquaculture 81, 351–365. https://doi.org/10.1016/0044-8486(89)90159-2

Cerdà, J., Fabra, M., Raldúa, D., 2007. Physiological and molecular basis of fish oocyte hydration. Fish Oocyte From Basic Stud. to Biotechnol. Appl. 349–396. https://doi.org/10.1007/978-1-4020-6235-3-12

Cerdà, J., 2009. Molecular pathways during marine fish egg hydration: The role of aquaporins. J. Fish Biol. 75, 2175–2196. https://doi.org/10.1111/j.1095-8649.2009.02397.x

Colson V, Cousture M, Damasceno D, Valotaire C, Nguyen T, Le Cam A, Bobe J. Maternal temperature exposure impairs emotional and cognitive responses and triggers dysregulation of neurodevelopment genes in fish. PeerJ. 2019 Jan 31;7:e6338. doi: 10.7717/peerj.6338. PMID: 30723624; PMCID: PMC6360074.

Craik, J.C.A., Harvey, S.M., 1984. Biochemical changes occurring during final maturation of eggs of some marine and freshwater teleosts. J. Fish Biol. 24, 599–610. https://doi.org/10.1111/j.1095-8649.1984.tb04830.x

Dolomatov, S.I., Shekk, P. V., Zukow, W., Kryukova, M.I., 2011. Features of nitrogen metabolism in fishes. Rev. Fish Biol. Fish. 21, 733–737. https://doi.org/10.1007/s11160-011-9212-z

Dolomatov, S.I., Zukow, W., Novikov, N.Y., Muszkieta, R., Bulatowicz, I., Dzierzanowski, M., Kazmierczak, U., Strojek, K., 2012. The regulation of osmotic and ionic balance in fish reproduction and in the early stages of ontogeny. Russ. J. Mar. Biol. 38, 365–374. https://doi.org/10.1134/S1063074012050057

Ecker, R.E., Dennis Smith, L., 1971. Influence of exogenous ions on the events of maturation in Rana pipiens oocytes. J. Cell. Physiol. 77, 61–70. https://doi.org/10.1002/jcp.1040770108

Escaffre, A. M., Billard, R., 1979. Évolution de la fécondabilité des ovules de truite arc-en-ciel (Salmo gairdneri) laissés dans la cavité abdominale au cours de la période post-ovulatoire. Bulletin Français de Pisciculture. 272, 56-70. https://doi.org/10.1051/kmae:1979012

Espey, L.L., 1994. Current status of the hypothesis that mammalian ovulation is comparable to an inflammatory reaction. Biol. Reprod. 50, 233–238. https://doi.org/10.1095/biolreprod50.2.233

Fabra, M., Raldúa, D., Power, D. M., Deen, P. M., & Cerda, J., 2005. Marine fish egg hydration is aquaporin-mediated. Science. 307(5709), 545-545. https://doi.org/10.1126/science.1106305

Finn, R.N., Østby, G.C., Norberg, B., Fyhn, H.J., 2002. In vivo oocyte hydration in Atlantic halibut (Hippoglossus hippoglossus); proteolytic liberation of free amino acids, and ion transport, are

driving forces for osmotic water influx. J. Exp. Biol. 205, 211–224. https://doi.org/10.1242/jeb.205.2.211

Fiol, D.F., Chan, S.Y., Kültz, D., 2006. Regulation of osmotic stress transcription factor 1 (Ostf1) in tilapia (Oreochromis mossambicus) gill epithelium during salinity stress. J Exp Biol Aug;209(Pt 16):3257-65. doi: 10.1242/jeb.02352. PMID: 16888073.

Fostier A, Breton B, Jalabert B, Marcuzzi O, 1981. Evolution des niveaux plasmatiques de la gonadotropine glycoprotéique et de la 17 alpha hydroxy-20 beta dihydroprogestérone au cours de la maturation et de l'ovulation chez la Truite arc-en-ciel. Salmo gairdnerii. CR Seances Acad. Sci. III. 293(15):817-20. French. PMID: 6804018.

Goetz, F.W., Bergman, H.L., 1978. The effects of steroids on final maturation and ovulation of oocytes from brook trout (Salvelinus fontinalis) and yellow perch (Perca flavescens). Biol. Reprod. 18, 293–298. https://doi.org/10.1095/biolreprod18.2.293

Gray, J., 1932. The osmotic properties of the eggs of the trout (Salmo Fario). Journal of Experimental Biology. 9(3), 277-299. https://doi.org/10.1242/jeb.9.3.277

Greeley Jr, M. S., Hols, H., Wallace, R. A., 1991. Changes in size, hydration and low molecular weight osmotic effectors during meiotic maturation of Fundulus oocytes in vivo. Comparative Biochemistry and Physiology, Part A: Physiology. 100(3), 639-647. https://doi.org/10.1016/0300-9629(91)90383-N

Grier, H.J., Uribe, M.C., Parenti, L.R., 2007. Germinal epithelium, folliculogenesis, and postovulatory follicles in ovaries of rainbow trout, Oncorhynchus mykiss (Walbaum, 1792). J Morphol. 268(4), 293-310. https://doi.org/10.1002/jmor.10518

Haffray, P., Fostier, A., M., Jalabert, B., 1995. Influence du maintien en mer ou de la période de transfert en eau douce des reproducteurs de saumon atlantique Salmo salar sur la maturation sexuelle et la qualité des gamètes. Aquatic Living Resources. 8(2), 135-145. https://doi.org/10.1051/alr:1995010

Hajnik, C.A., Goetz, F.W., Hsu, S.Y., Sokal, N, 1998. Characterization of a ribonucleic acid transcript from the brook trout (Salvelinus fontinalis) ovary with structural similarities to mammalian adipsin/complement factor D and tissue kallikrein, and the effects of kallikrein-like serine proteases on follicle contraction. Biol Reprod. 58(4):887-97. https://doi.org/10.1095/biolreprod58.4.887

Hagiwara A, Ogiwara K, Sugama N, Yamashita M, Takahashi T, 2020. Inhibition of medaka ovulation by gap junction blockers due to its disrupting effect on the transcriptional process of LH-induced Mmp15 expression. Gen Comp Endocrinol. 1;288:113373. https://doi.org/10.1016/j.ygcen.2019.113373

Hanumanthaiah, R., Day, K., Jagadeeswaran, P., 2002. Comprehensive analysis of blood coagulation pathways in teleostei: Evolution of coagulation factor genes and identification of Zebrafish factor VIIi. Blood Cells, Mol. Dis. 29, 57–68. https://doi.org/10.1006/bcmd.2002.0534

Hurk, R. van den, Peute, J., 1979. Cyclic changes in the ovary of the rainbow trout, Salmo gairdneri, with special reference to sites of steroidogenesis. Cell Tissue Res. 199, 289–306. https://doi.org/10.1007/BF00236140

Hsu, S., Goetz, F.W., Sciences, B., Dame, N., Dame, N., 1992. Angiotensins stimulate in vitro ovulation and contraction of brook trout (Salvelinus fontinalis) follicles. Fish Physiol. Biochem. 10, 277–282. https://doi.org/10.1007/BF00004476

Izquierdo, M.S., Fernández-Palacios, H., Tacon, A.G.J., 2001. Effect of broodstock nutrition on reproductive performance of fish. Aquaculture. 197, 25–42. https://doi.org/10.1016/S0044-8486(01)00581-6

Jalabert B, Breton B, Bry C., 1972. Maturation et ovulation in vitro des ovocytes de la truite arc-en-ciel Salmo gairdneri. CR Acad Hebd Seances Acad Sci D. 11;275(11):1139-42. French. PMID: 4196379.

Jalabert, B., Fostier, A., 1984. The follicular sensitivity in vitro to maturation-inducing hormones in rainbow trout, Salmo gairdnerii: role of oestradiol-17β. Aquaculture. 43(1-3), 1-11. https://doi.org/10.1016/0044-8486(84)90004-8

Jalabert, B., 1976. In vitro oocyte maturation and ovulation in rainbow trout (Salmo gairdnerii), Northern Pike (Esox lucius), and Goldfish (Carassius auratus). J. Fish. Res. Board Canada. 33, 974–988. https://doi.org/10.1139/f76-124

Jalabert, B., Theron, M. C., Heydorff, M., 1978. Production of fertilizable oocytes from follicles of rainbow trout (Salmo gairdnerii) following in vitro maturation and ovulation. Annales de Biologie Animale Biochimie Biophysique (Vol. 18, No. 2B, pp. 461-470). EDP Sciences.

Jalabert, B., Szöllösi, D., 1975. In vitro ovulation of trout oocytes : Effect of prostaglandins on smooth muscle-like cells of the theca. Prostaglandins. 765–778. https://doi.org/10.1016/0090-6980(75)90113-6

Joy, K.P., Chaube, R., 2015. General and Comparative Endocrinology Vasotocin – A new player in the control of oocyte maturation and ovulation in fish. Gen. Comp. Endocrinol. https://doi.org/10.1016/j.ygcen.2015.02.013

Knoll-Gellida, A., André, M., Gattegno, T., Forgue, J., Admon, A., & Babin, P. J., 2006. Molecular phenotype of zebrafish ovarian follicle by serial analysis of gene expression and proteomic profiling, and comparison with the transcriptomes of other animals. BMC genomics. 7(1), 1-28. https://doi.org/10.1186/1471-2164-7-46

LaFleur, G.J., Thomas, P., 1991. Evidence for a role of Na+,K+-ATPase in the hydration of Atlantic croaker and spotted seatrout oocytes during final maturation. J. Exp. Zool. 258, 126–136. https://doi.org/10.1002/jez.1402580114

Lahnsteiner, F., 2000. Morphological, physiological and biochemical parameters characterizing the over-ripening of rainbow trout eggs. Fish physiology and biochemistry. 23(2), 107-118. https://doi.org/10.1023/A:1007839023540

Lall, S.P., 2003. The Minerals. Fish Nutr. 259–308.

    https://doi.org/10.1016/b978-012319652-1/50006-9

Lubzens, E., Young, G., Bobe, J., Cerdà, J., 2010. Oogenesis in teleosts: How fish eggs are formed. Gen.

    Comp. Endocrinol. 165, 367–389. https://doi.org/10.1016/j.ygcen.2009.05.022

Maisse, G., Billard, R., Cosson, J., Labbé, C., Loir, M., Le Gac, F., & Fierville, F., 1998. Influence du

    maintien en mer des mâles de saumon atlantique (Salmo salar), pendant la période de

    reproduction, sur la qualité du sperme. Bulletin Français de la Pêche et de la Pisciculture. (350-

    351), 349-357. https://doi.org/10.1051/kmae:1998010

Marshall, W.S., Habibi, H.R., Lessman, C.A., 1985. Electrophysiology of oocytes during meiotic

    maturation and after ovulation in brook trout (Salvelinus fontinalis). Can. J. Zool. 63, 1904–1908.

    https://doi.org/10.1139/z85-283

Milla, S., Jalabert, B., Rime, H., Prunet, P., Bobe, J., 2006. Hydration of rainbow trout oocyte during

    meiotic maturation and in vitro regulation by 17,20β-dihydroxy-4-pregnen-3-one and cortisol. J.

    Exp. Biol. 209, 1147–1156. https://doi.org/10.1242/jeb.02094

Morro, B., Doherty, M.K., Balseiro, P., Handeland, S.O., Albalat, A., 2020. Plasma proteome profiling of

    freshwater and seawater life stages of rainbow trout (Oncorhynchus mykiss). PLoS One. 2020 Jan

    3;15(1):e0227003. https://doi.org/10.1371/journal.pone.0227003

Nakada, T., Zandi-Nejad, K., D.B., Hirose, S., 2005. Roles of Slc13a1 and Slc26a1 sulfate transporters of

    eel kidney in sulfate homeostasis and osmoregulation in freshwater. Am. J. Physiol. - Regul.

    Integr. Comp. Physiol. 289. https://doi.org/10.1152/ajpregu.00725.2004

Nevoux, M., Marchand, F., Forget, G., Huteau, D., Tremblay, J., & Destouches, J. P., 2019. Field

    assessment of precocious maturation in salmon parr using ultrasound imaging. Peer Community

    in Ecology, 1101/425561.

Norman, J.D., Ferguson, M.M., Danzmann, R.G., 2014. Transcriptomics of salinity tolerance capacity in

Arctic charr (Salvelinus alpinus): a comparison of gene expression profiles between divergent QTL genotypes. Physiol Genomics. 2014 Feb 15;46(4):123-37. https://doi.org/10.1152/physiolgenomics.00105.2013

Ohnishi, J., Ohnishi, E., Shibuya, H., Takahashi, T., 2005. Functions for proteinases in the ovulatory process. Biochimica et Biophysica Acta (BBA)-Proteins and Proteomics. 1751(1), 95-109. https://doi.org/10.1016/j.bbapap.2005.05.002

Pankhurst, N.W., Purser, G.J., Van Der Kraak, G., Thomas, P.M., Forteath, G.N.R., 1996. Effect of holding temperature on ovulation, egg fertility, plasma levels of reproductive hormones and in vitro ovarian steroidogenesis in the rainbow trout Oncorhynchus mykiss. Aquaculture. 146, 277–290. https://doi.org/10.1016/S0044-8486(96)01374-9

Patiño, R., Sullivan, C. V., 2002. Ovarian follicle growth, maturation, and ovulation in teleost fish. Fish Physiology and Biochemistry. 26(1), 57-70. https://doi.org/10.1023/A:1023311613987

Pujante, I. M., Moyano, F. J., Martos-Sitcha, J. A., Mancera, J. M., & Martínez-Rodríguez, G., 2018. Effect of different salinities on gene expression and activity of digestive enzymes in the thick-lipped grey mullet (Chelon labrosus). Fish physiology and biochemistry, 44(1), 349–373. https://doi.org/10.1007/s10695-017-0440-6

Richards, J.A.S., Russell, D.L., Ochsner, S., Espey, L.L., 2002. Ovulation: New dimensions and new regulators of the inflammatory-like response. Annu. Rev. Physiol. 64, 69–92. https://doi.org/10.1146/annurev.physiol.64.081501.131029

Rothberger, H., McGee, M. P., 1984. Generation of coagulation factor V activity by cultured rabbit alveolar macrophages. The Journal of experimental medicine. 160(6), 1880-1890. https://doi.org/10.1084/jem.160.6.1880

Samarin, A.M., Ahmadi, M.R., Azuma, T., Rafiee, G.R., Amiri, B.M., Naghavi, M.R., 2008. Influence of the time to egg stripping on eyeing and hatching rates in rainbow trout Oncorhynchus mykiss under cold temperatures. Aquaculture. 278, 195–198.

https://doi.org/10.1016/j.aquaculture.2008.03.034

Samarin, A.M., Policar, T., Lahnsteiner, F., 2015. Fish oocyte ageing and its effect on egg quality. Rev. Fish. Sci. Aquac. 23, 302–314. https://doi.org/10.1080/23308249.2015.1053560

Springate, J.R.C., Bromage, N.R., Elliott, J.A.K., Hudson, D.L., 1984. The timing of ovulation and stripping and their effects on the rates of fertilization and survival to eying, hatch and swim-up in the rainbow trout (Salmo gairdneri R.). Aquaculture 43, 313–322. https://doi.org/10.1016/0044-8486(84)90032-2

Takahashi, T., Hagiwara, A., Ogiwara, K., 2019. Follicle rupture during ovulation with an emphasis on recent progress in fish models. Reproduction. 157(1):R1-R13. https://doi.org/10.1530/REP-18-0251

Thibault, C., Levasseur, M.C., 1988. Ovulation. Hum Reprod. 3(4):513-23. https://doi.org/10.1093/oxfordjournals.humrep.a136737

Tripathi, V., Krishna, A., 2008. Changes in nitric oxide (NO) synthase isoforms and NO in the ovary of Heteropneustes fossilis (Bloch.) during the reproductive cycle. J. Endocrinol. 199, 307–316. https://doi.org/10.1677/JOE-07-0509

Valdebenito, I.I., Gallegos, P.C., Effer, B.R., 2015. Gamete quality in fish: Evaluation parameters and determining factors. Zygote 23, 177–197. https://doi.org/10.1017/S0967199413000506

Wallace, R. A., & Selman, K., 1981. Cellular and dynamic aspects of oocyte growth in teleosts. American zoologist. 21(2), 325-343. https://doi.org/10.1093/icb/21.2.325

Wallace, R. A., Greeley, M. S., McPherson, R., 1992. Analytical and experimental studies on the relationship between Na+, K+, and water uptake during volume increases associated with Fundulus oocyte maturation in vitro. Journal of Comparative Physiology B. 162(3), 241-248. https://doi.org/10.1007/BF00357530

Watanabe, W. O., & Kuo, C. M., 1986. Water and ion balance in hydrating oocytes of the grey mullet,

Mugil cephalus (L.), during hormone-induced final maturation. Journal of fish biology. 28(4), 425-437. https://doi.org/10.1111/j.1095-8649.1986.tb05180.x

White, J.M., 2003. ADAMs : modulators of cell – cell and cell – matrix interactions. Current opinion in cell biology. 15(5), 598-606. https://doi.org/10.1016/j.ceb.2003.08.001

Yamashita M, Mita K, Yoshida N, Kondo T., 2000. Molecular mechanisms of the initiation of oocyte maturation: general and species-specific aspects. Prog Cell Cycle Res. 4:115-29. https://doi.org/10.1007/978-1-4615-4253-7_11

Figure 1 : Water and plasma composition classified by site

Figure 2 : Egg quality classified by site

A Late vitellogenesis = LV

Post-vitellogenesis = PV

Maturation = MAT

Ultrasound oocyte staging

B LV

PV

MAT

Direct oocyte staging

C

p=0.002*

Hours (10°C)

Control   LSW

Duration of FOM

Figure 3: Ultrasound staging for measure of maturation duration

Figure 4: Egg quality depending on the duration of oocyte maturation

Figure 5: Hydration of oocyte during final oocyte maturation

Figure 6: **Ovarian expression profiles of genes involved in FOM and ovulation mechanisms**