Atypical integrative element with strand-biased circularization activity assists interspecies

antimicrobial resistance gene transfer from *Vibrio alfacsensis*

Lisa Nonaka,[1,2*] Michiaki Masuda,[1] Hirokazu Yano[3,4*]

[1]Department of Microbiology, Dokkyo Medical University School of Medicine, Mibu,

Tochigi, Japan

[2]Faculty of Human Life Sciences, Shokei University, Kumamoto, Japan

[3]Faculty of Life and Environmental Sciences, University of Tsukuba, Tsukuba, Japan

[4]Graduate School of Life Sciences, Tohoku University, Sendai, Japan

Short title: Integrative elements with strand-biased circularization activity

*Corresponding authors

Email: nonaka20@shokei-gakuen.ac.jp (LN); yano.hirokazu@ige.tohoku.ac.jp (HY)

Abstract

The exchange of antimicrobial resistance (AMR) genes between aquaculture and terrestrial microbial populations has emerged as a serious public health concern. However, the nature of the mobile genetic elements in marine bacteria is poorly documented. To gain insight into the genetic mechanisms underlying AMR gene transfer from marine bacteria, we mated a multidrug-resistant *Vibrio alfacsensis* strain with an *Escherichia coli* strain, and then determined the complete genome sequences of the donor and transconjugant strains. Sequence analysis revealed a multidrug resistance conjugative plasmid in the donor strain, which was integrated into the chromosome of the recipient. The plasmid backbone in the transconjugant chromosome was flanked by two copies of a 7.1 kb unclassifiable integrative element harboring a β-lactamase gene. The 7.1 kb element (named SE-6945) and previously reported Tn*6283* share four coding sequences, two of which encode the catalytic R-H-R-Y motif of tyrosine recombinases. Polymerase chain reaction and sequencing experiments revealed that SE-6945 generates a circular copy of its one specific strand without generating an empty donor site, in contrast to the movement of integron gene cassette or ICE/IMEs discovered to date. Copy numbers and location of SE-6945 in the chromosome affected antibiotic resistance levels of the transconjugants. SE-6945/Tn*6283*-related integrative elements are collectively termed SEs (s̲trand-biased circularizing integrative e̲lements). SEs were identified in the genomes of other *Vibrio* species. Overall, these results suggest that in addition to conjugative plasmids, SEs are involved in the spread of AMR genes among marine bacteria.

Introduction

Antimicrobials have been used globally in aquaculture to control fish diseases. Although this approach helps to maintain a stable supply of aquacultural products, misuse of antimicrobials has led to the emergence of antimicrobial-resistant microbes and the accumulation of resistance genes at aquaculture sites [1,2]. As the spread of multidrug-resistant (MDR) bacteria and pan-drug-resistant bacteria threatens human life [3,4], it is important to obtain clues about whether and how aquatic and terrestrial microbial populations exchange genetic materials.

Plasmids [5] and integrative and conjugative elements (ICE) [6] are DNA units that can move from one cell to another through the conjugation machinery. They can directly move between species [7,8], and indirectly mobilize genes on specific mobile elements [9–11]. Plasmids can be classified into families according to replicon type or mobilization machinery type for epidemiological purposes [12,13].

Transposons have been defined as specific DNA segments that can repeatedly insert into one or more sites in one or more genomes [14]. Several distinct strand exchange mechanisms and the routes used for their location changes, i.e., *transposition*, have been discovered to date [15]. In the comprehensive review by Curcio and Derbyshire [15], transposons were classified into five families based on the DNA strand exchange enzymes used. In prokaryotes, the mode of transposition was largely classified into 'cut-out paste-in' (often called 'excision integration' for the ICEs [6]), 'copy-out paste-in,' or 'copy-in' for which replication is necessary to complete transposition; e.g., Tn*3*, IS*91*. DNA insertion involving DNA replication can also be seen in the insertion of a folded single-strand DNA of Integron gene cassette into *attI* [16] and of phage CTX genome into *dif* [17]; both use tyrosine

recombinase(s) as the strand exchange enzyme. In transposon classification, ICEs are

classified as transposons that use Y-transposase (tyrosine recombinase) or S-transposase

(serine recombinase) [15]. Other than ICEs, DNA segments that can move to a location in

other cells via the excision – transfer (rolling-circle replication) – integration route have

been termed IME [18], MGI [10], CIME [19], or MTn [20] (hereafter collectively referred to

as IME). IMEs encode site-specific recombinase or DDE transposase as strand exchange

enzyme [21]. They have been recognized as degraded forms of ICEs [19]. ICEs/IMEs so far

studied in the laboratory seem to change their location via the 'excision integration' route

only [22–25]. Genes embedded in these mobile genetic elements are readily shared among

bacteria and provide genetic resources for microbial adaptation in changing environments.

To better understand the genetic mechanisms underlying the spread of antibiotic

resistance genes at aquaculture sites, we previously collected resistant bacteria from

sediments at an aquaculture site in Kagawa, Japan [26–28]. We identified a self-

transmissible MDR plasmid of the MOB$_H$-family, named pAQU1, that can replicate in both

the original host *Photobacterium damselae* subsp. *damselae* and in *Escherichia coli* [27].

Subsequently, another MOB$_H$-family MDR plasmid, named pSEA1, was identified in a *Vibrio*

alfacsensis isolate (previously identified as *V. ponticus*) [29]. pSEA1 carries a 12 kb

unclassifiable integrative element, Tn*6283* (renamed as SE-6283 in this study), in addition to

the antimicrobial resistance (AMR) genes. Although pSEA1 could not replicate in *E. coli* at

42 °C, following conjugation, it could integrate into the *E. coli* chromosome by homologous

recombination between two SE-6283 copies: one on pSEA1 and another that moved from

pSEA1 into the chromosome [29]. Further, SE-6283 was suggested to move via a circular

copy intermediate of one specific strand without generating an empty donor site,

presumably using tyrosine recombinases. However, it is not known whether interspecies

gene transfer assisted by SE-6283-like elements is common in nature.

To obtain further insights into the mechanisms behind genetic exchange among

aquaculture-associated bacteria, in this study we mated another MDR *Vibrio* isolate, *V.*

alfacsensis 04Ya249, with *E. coli* strains in the laboratory and then determined the genome

sequences of both the donor and transconjugant strains. We identified a SE-6283-like

element involved in the integration of the multidrug resistance plasmid into *E. coli*

chromosome. Based on the findings in this study, we have termed the currently

unclassifiable integrative element, SE-6283, and its related elements as a new class of

mobile elements: <u>s</u>trand-biased circularizing integrative <u>e</u>lement (SE).

In this study, several types of intracellular DNA movements need to be discussed. For

clarity, terms *transposition*, *integration*, and *insertion* are used here based on the following

rules. We call the movement of a DNA segment from one location to the other location in

one or two genomes, mediated by any DNA strand exchange enzymes, as transposition. The

physical incorporation of a plasmid into a chromosome is called integration regardless of the

molecular mechanisms and genomic locations involved. The physical incorporation of a DNA

molecule into a specific site in another molecule mediated by a site-specific recombinase

(so-called integrase) has traditionally been called integration, but we refer to it as an

insertion to distinguish this event from the plasmid integration defined above. The term

translocation has frequently been used when referring to rarely occurring location change

of a large DNA segment between chromosomes in organisms carrying multiple

chromosomes. We did not use translocation in this study as it is not relevant to this study.

Results

Identification of SE carrying a β-lactamase gene

We previously isolated *V. alfacsensis* strain 04Ya249 from sea sediment at an aquaculture

site [26]. 04Ya249 showed resistance to erythromycin, tetracycline, and ampicillin [30]. To

identify active mobile elements in this strain that are relevant to its AMR, the strain was

mated with macrolide-sensitive *E. coli* JW0452, and the transconjugant was first selected in

the presence of erythromycin at 42 °C. One transconjugant was named strain TJ249.

The complete genome sequences of strains 04Ya249 and TJ249 were determined

using the PacBio RS II platform. The genome of strain 04Ya249 consists of four replicons:

two chromosomes, a 306 kb putative conjugative plasmid pSEA2, and an 82 kb putative

conjugative plasmid pVA249. Strain 04Ya249 has a very similar genome architecture to the

database strain *V. alfacsensis* CAIM 1831 based on gene synteny (S1 Fig A) and average

nucleotide identity (S1 Fig B). Seven AMR genes were identified on plasmid pSEA2 (S2 Fig A).

One β-lactamase gene (*CARB-19* allele in the CARD database [31]) was located within the

7.1-kb repeat region found in both chromosome 1 and plasmid pSEA2 (S2 Fig B), which

contained four protein-coding sequences in addition to a β-lactamase gene. These four

coding sequences showed homology to the four coding sequences clustered at one end of

SE-6283 from strain 04Ya108 (Fig 1A), although gene product identities between SE-6945

and SE-6283 counterparts were very low: 22.5 % for IntA, 20.5 % for CDS2, 23.2 % for IntB,

and 26.6 % for CDS4 in blastp.

This 7.1-kb region was confirmed to be active in generating its circular copy of top

strand in *E. coli* and *V. alfacsensis* and inserting it into new target locations in *E. coli* in this

study. The 7.1kb region was designated as a new member of SE, SE-6945. The four

conserved coding sequences (SE core genes) were tentatively called *intA*, CDS2, *intB*, and

CDS4. The biochemical functions of their products were estimated using Pfam and

InterProScan in EMBL. The *intA* and *intB* products generated a hit on phage_integrase

(CL0382) in Pfam and were further aligned with known tyrosine recombinases using

structure-aware alignment program PROMAL3D [32] to confirm the presence of conserved

motifs. The *intA* and *intB* products contained the catalytic R-H-R-Y motif (G-<u>R</u> motif in box I

and <u>H</u>-<u>R</u>-H-<u>Y</u> motif in box II (Fig 1B)) conserved in tyrosine recombinases [33], though the N-

terminal region of IntB is extraordinarily long among tyrosine recombinases. CDS2 and CDS4

products did not generate hits in the Pfam or InterProScan databases. The SEs carry

imperfect 19 bp inverted repeat motifs termed C and C' at two ends (Fig. 1 A), which might

be equivalent to core-type sites [34] used in strand change mediated by tyrosine

recombinase(s).

Fig 1. Genetic features of SEs. (A) Upper panel: Genetic organization of SE-6283 and SE-

6945. Lower panel: Nucleotide sequences of imperfect inverted repeat motifs C and C'. The

nicking sites used for circularization were six bases upstream of the 5⬛-end of motif C and

the 3⬛-end of motif C' (black arrowheads in the upper panel). No strand exchange was

detected on the bottom strand (white arrowheads). (B) Presence of R-H-R-Y motif in IntA

and IntB. IntA and IntB sequences are aligned with sequences of known tyrosine

recombinases based on secondary structure prediction using PROMAL 3D [32]. The aligned

regions corresponding to box I and box II in [33] are shown. (C) Schematic representation of

SE-6945 and pSEA2 insertion locations in *E. coli* TJ249.

As pSEA2 showed > 99.9 % nucleotide identity on the aligned region to pSEA1

(accession no. LC081338.1) from strain 04Ya108 [29], we also determined the complete

genome of strain 04Ya108 for comparison with strain 04Ya249. As predicted from Southern

hybridization [29], strain 04Ya108 possessed two copies of SE-6283, one in chromosome 1

and the other in plasmid pSEA1. Strain 04Ya108 also carries SE-6945 in the same two

locations as strain 04Ya249 (S1 Fig A, S2 Fig B).

The chromosome of *E. coli* TJ249 contains two notable insertions (Fig 1C), a smaller

insertion of SE-6945 alone immediately downstream of the putative transcription

termination site for *yjjN* (*yjjNt*) located between *yjjN* and *yjjO* (in GenBank accession no.

U00096.3), and a larger insertion into *insJ*, containing a nearly complete copy of pSEA2 and

an additional copy of SE-6945. The 14.2 kb *mphA-cat* gene region flanked by repeats in

pSEA2 (contig 1 in S2 Fig A) was detected as a separate circular contig in the TJ249 genome

assembly. These indicate the movement between different genomic locations (pSEA2, *yjjNt*,

InsJ), thereby transposition, of SE-6945 in *E. coli*.

The border regions between SE-6945 and its target locations were termed *attL* (*intA*-

proximal side) and *attR* (*intA*-distal side) (Fig 2 A). The nucleotide sequence alignments of

attL and *attR* from two *E. coli* chromosomal locations and two locations in 04Ya249

genomes (pSEA2 and chr 1) are shown in S3 Fig and S4 Fig, respectively. Sequence

comparison of the *attL*, *attR*, and unoccupied target site in the host (*attB*) in *E. coli* (Fig 2B)

revealed that the 6 bp next to the 3⊡ end of C' in *attR* in *E. coli* (5'-TTTTCT-3') was identical

to the 6 bp next to the 5⊡ end of C at *attL* on pSEA2, while the 6 bp next to the 5⊡ end of C

in *attL* (5⊡-TTCTTT-3⊡ in *insJ*, 5⊡-TTTTTT-3⊡ in *yjjNt*) was from *attB* (S3 Fig, S4 Fig, Fig 2B).

Therefore, SE-6945 inserts itself, ending with 5⊡-GTA-3⊡ (Fig 1 A) along with an additional 6

bp next to the 5☐ end of C at *attL* on the donor molecule into the target site. The 6 bp from

the donor molecule is placed next to the 3☐ end of C' in the newly formed *attR.*

Target site preference of SE-6945

After the isolation and genome sequence determination of TJ249, additional mating

experiments between *E. coli* JW0452 and *V. alfacsensis* 04Ya249 were performed to test

whether pSEA2 integration into a specific location of *E. coli* chromosome accompanying SE-

6945 transposition is reproducible. Independent mating was performed more than 17 times,

and transconjugant was screened in the presence of Tc alone, Ap alone, or both Ap and Tc.

SE-6945 insertion pattern was investigated by Southern hybridization for 16 randomly

chosen Tc resistant transconjugants, and one Tc sensitive but Ap-resistant transconjugant

obtained from Ap alone selection (S5 Fig). Among 16 Tc resistant transconjugant strains, 12

strains generated *intA*-containing fragments originating from the *yjjNt*-SE-6945 junction or

the *insJ*-SE-6945 junction. These 16 strains also generated pSEA2-derived fragments (S5 Fig).

The remaining four strains generated pSEA2-derived fragments and contained only one

copy of *intA*, suggesting SE-6945-independent pSEA2 integration into chromosome;

autonomous replication of pSEA2 seems unlikely given the high frequency of pSEA2

integration into the chromosome in other transconjugant strains. Strain LN52, obtained

from Ap alone selection, carried only one SE-6945 copy, and it was inserted into *yjjNt* (S5

Fig). Taken together, these observations suggest that pSEA2 integration into *E. coli*

chromosome generally accompanies the SE-6945 transposition, and that SE-6945 prefers

yjjNt or *insJ*, but not random sites in the *E. coli* genome.

To obtain further insights into the target site of SE-6945, bacterial genomes carrying

SE-6945 were searched based on blastn in NCBI nucleotide collection (nr/nt) database using

the SE-6945 core gene region (*intA*-CDS2-*intB*-CDS4) as the query. The genome of *Vibrio*

harveyi strain WXL538 contained a segment showing 88.9% nucleotide sequence identity to

the SE-6945 core genes. Genome structure comparisons between WXL538 and 04Ya249

revealed that WXL538 carries a 16.4 kb SE-6945-like SE (tentatively termed SE-VhaWXL538)

containing a different set of cargo genes (Fig 2 A). SE-VhaWXL538 is inserted into the

promoter region of a *yaaA* gene encoding an H2O2_YaaD_Superfamily protein (NCBI

conserved domain accession no. cl01143) (locus_tag F9277_14400 in CP045070.1). This

location corresponds to the SE-6945 insertion location in *V. alfacsensis* chromosome 1 (Fig 2

A). Sequences of unoccupied target sites (*attB*) in *E. coli*, *V. alfacsensis*, and *V. harveyi* are

shown in Fig 2B. The conservation level of nucleotide in *attB* is presented in Fig 2C as

Sequence logos information contents [35]. The central 6 bp putative crossover region in *attB*

was T rich. However, other motifs, such as imperfect inverted repeat structure observed in

attC/attI of integron (IntI target) [36,37], or *dif* of CTX phage integration site (XerC/XerD

target) [17], could not be detected in *attB, attL* (S3 Fig), or *attR* (S4 Fig) for SE-6945 in the

current data set.

Fig 2. Locations of SE-6945 and its related elements. (A) Comparisons of genetic

organization around SE insertion locations. GenBank accession numbers (*Vibrio* strains only)

and the coordinate of genomic segments compared are shown below the strain name. (B)

Alignment of *attB* sequences. The sequences are retrieved from SE-free genomes. Since the

SE-6945-free pSEA2 ancestor could not be found in the database, the ancestral *attB*

sequence on pSEA2 in the putative crossover region shown as x is unpredictable. Nicking is

expected to take place next to the central box: one site on the left side on the top strand.

(C) Sequence logos representation of conservation levels of *attB* sequence.

Transposition of SE-6945 and plasmid integration in mating

The direct repeat structure of SE-6945 in the TJ249 chromosome suggested that

pSEA2 was integrated into the recipient chromosome through the transposition of SE-6945,

with subsequent integration of pSEA2 into the chromosome by homologous recombination

between SE-6945 copies: one on the chromosome and the other on pSEA2 (referred to as a

two-step gene transfer mechanism in the previous study [29]). If this scenario holds,

insertion of SE-6945 alone should occur more frequently than pSEA2 integration upon

mating, and pSEA2 integration would be reduced in frequency or abolished altogether in the

chromosome of a *recA*-null mutant recipient. To test these possibilities, we conducted

additional mating assays.

Quantitative mating assays were performed using a rifampicin-resistant recipient

strain, JW0452rif, and its isogenic strain carrying *recA* deletion JW0452Δ*recA*rif. The transfer

of Tc Ap resistance from *V. alfacsensis* 04Ya249 to *E. coli* JW0452rif was detected only in 2

out of 4 replicated mating experiments, and the transfer frequency was determined to be

approximately 10^{-9} per donor, indicating a very rare event, generating 0–3 transconjugant

colonies per mating (Fig 3). The transfer of Ap resistance alone was observed at >300-fold

higher frequency than the transfer of Tc Ap-resistant transconjugants (compare the first and

the second groups from left in Fig 3), indicating that insertion of SE-6945 alone occurs more

frequently than pSEA2 integration during mating.

The transfer of Tc Ap resistance to JW0452Δ*recA*rif could not be detected in all four

independent mating assays (below detection limit of 1.02×10^{-9} per donor). Unexpectedly

transfer frequency of Ap resistance was also reduced to the 1/17 level of JW0452rif in

JW0452Δ*recA*rif (compare the second and the fourth groups from the left in Fig 3). Thus, the

direct effect of *recA* knockout on pSEA2 integration could not be evaluated due to the

reduced detection sensitivity of gene transfer. However, when LN52rif (which already

carries one copy of SE-6945 and wild type *recA*) was used as the recipient, Tc Ap-resistant

transconjugants were obtained at a 56-fold higher frequency (in mean) than that observed

when using JW0452rif as the recipient (compare the first and fifth groups in Fig 3). This

indicates that the presence of the 7.1 kb long homology stretches in both target DNA and

pSEA2 promotes pSEA2 integration in the recipient.

The Tn*3*-like cointegrate formation [15] might happen if SE-6945 could

simultaneously introduce nick at one site on the top strand at *attL* (or *attR*) and one site on

the bottom strand at *attR* (or *attL*), and DNA-protein crosslink (DPC) repair [38] could clean

the linked tyrosine recombinase to produce a 3′-OH priming site. If SE-mediated

cointegrate formation occurs, then Tc Ap-resistant transconjugant should emerge at a

comparable frequency to Ap-resistant transconjugant, which was not the case in our

observation.

Collectively, these results and the nature of tyrosine recombinases support the

model that pSEA2 integration mainly occurs through a two-step gene transfer mechanism;

transposition followed by homologous recombination than through a one-step Tn*3*-like

cointegrate formation process.

Fig 3. Resistance gene transfer from *V. alfacsensis*. Transfer frequency (Y-axis) is the $\log_{10}$-

transformed value of the transconjugant colony-forming units (CFU) divided by the donor

CFU. Four replicate mating experiments were performed. No transconjugant was detected

in any of the four replicate experiments using JW0452Δ*recA*rif with Tc Ap selection or in the

two replicate experiments using JW0452rif with Tc Ap selection (indicated by open circles

placed at the bottom). In one experiment on LN52rif, the donor CFU was not evaluated, but

the transconjugant CFU was obtained at a comparable frequency to that observed in the

other three replicate experiments. The detection limit of this experiment was around 1.02 ×

10^{-9} per donor.

Strand-biased circularization of SE-6945

We previously [29] revealed the following unique features of SE-6283: (i) it does not

generate an unoccupied donor site upon its circularization *in vivo*, and (ii) the circular form

of SE-6283 is generated using only one strand as a template at least in *E. col*. We note that

this assertion is based on the notion that tyrosine recombinases have a nature to not

produce blunt ends at *att* sites on double-strand DNA during strand exchange. Although SE-

6283 encodes tyrosine recombinases and not a single DDE transposase, the outcomes

described above, which result only from a figure-eight structure containing a single-strand

DNA bridge (Path 1 in Fig 4A), is reminiscent of the first step of the 'copy-out paste-in'

transposition of ISs [39] (IS*3* route), rather than the excision of ICEs (ICE/IME route) or the

single-strand circle generation of Integron gene cassette by *attC* × *attC* recombination (gene

cassette route), both of which generate empty donor sites [24,40,41]. If the figure-eight

structure containing a single-strand DNA bridge is subject to replication without DPC repair,

the products would be a single-strand SE circle, the original copy of donor molecule, and the

unjoined chromosome, which is eventually lost (Fig 4A bottom left). To test whether circular

copy production of SE-6945 also follows Path 1, we created a pSEA2-free 04Ya249 derivative

strain LN95 carrying only a single copy of SE-6945 in the chromosome and analyzed the

production of its circular molecule in this strain and the sequence of strand exchange

products.

Pairs of primers were designed such that one primer anneals to the inside of the

integrative element and the other anneals to the outside of the element (Fig 4A), amplifying

attL (intA side, Fig 4B, product size 304 bp), attR (bla side, 248 bp), empty donor site (attB,

345 bp), occupied attB (7.4 kb), and attS (joint of motif C and C' formed on the circular SE-

6945, 206 bp). Primer set 1-4 in Fig 4 A were used to detect both empty attB and occupied

attB.

The circular form of SE-6945 was detected in LN95 (Fig 4B) as well as the parent

strain 04Ya249 (S6 Fig A). This suggests that the cis-elements (attL/attR) on chromosome 1

and strand exchange enzymes encoded by the chromosomal copy of SE-6945 are functional

in V. alfacsensis. The empty attB (345 bp) could not be detected under PCR conditions with

short extension times (Fig 4 B(i)), while a larger product of occupied attB (7.4 kb) was

detected when the Long-PCR condition was used (Fig 4 B(ii)).

Fig 4. PCR and sequencing analyses of SE-6945 circle. (A) Design of PCR experiment to

detect SE-6945 circle. Upper panel: primer annealing locations (triangles in gray) on SE-6945

and the attB region. The number corresponds to the primer set numbers in B. Primer

number indicates the following primers in Table 2: 1, LN_181_Valc_chr1_attB_L; 2,

LN_142_junction2; 3, LN_143_junction2; 4, LN183_Valc_chr1_attB_R2. Lower panel:

possible routes of SE movement. Two strands of DNA are shown in separate lines. Lines in

orange are SEs. In Path 1 (solid arrows), one strand is nicked, and the SE side is circularized

while the host DNA side of the strand remains unjoined, leading to a figure-eight structure

containing a single-strand DNA bridge. The figure-eight form can be subject to replication,

giving rise to a double-strand SE circle (if the recombinase bound 3▯ end is cleaned, for

example, by DPC repair [38]) (copy-out route), or single-strand DNA circle (if the 3▯ end is

not cleaned). In Path 2, strand exchange generates a figure-eight containing a double-strand

DNA bridge. After going through Path 2, there are two possible outcomes: (i) another round

of strand exchange at bottom strand leading to circular a double-strand SE as well as a

circular donor molecule carrying empty donor site (ICE/IME route), (ii) replication on the

figure-eight leading to single-strand SE circle and donor molecule with empty *attB*, and the

original donor molecule (gene cassette route). (B) The result of PCR. Numbers in the primer

set correspond to primer numbers in panel A. PCR products were electrophoresed in 2%

agarose gel (i) or 0.8% agarose gel (ii). (C) Putative nicking sites in *attL* and *attR*. The SE DNA

is shown in orange. Nicking and ligation at positions shown in arrowheads in the same color

can generate *attS* with a 6 bp spacer. (D) Observed *attS* sequences. *attS* type 1 can be

generated from strand exchange at the top strand on chromosome 1 or bottom strand on

pSEA2, while *attS* type 1 can be generated from strand exchange at the top strand on pSEA2

or bottom strand on chromosome 1.

To estimate the copy number of the circular form of SE-6945 in the cell population,

we first searched for Illumina reads of strain LN95 that spanned *attS* or the "hypothetical"

attB. Although next-generation sequencing reads were obtained at 229× chromosome

coverage, no reads spanning *attS* or *attB* were detected [42]. Quantitative PCR further

revealed that the mean *attS* to *gyrB* ratio was 0.0012 (S6 Fig B). This relative copy number

was consistent with the result from a previous study detecting the circular form of SE-6283

in *E. coli*, which showed an *attS* to chromosome ratio of 0.001 [29].

To determine the nicking site on *attL* and *attR* (Fig 4D), we cloned the PCR products

of *attS* and then sequenced ten cloned molecules. Based on the *attL*, *attR*, and the original

attB sequences in the TJ249 chromosome (S3 Fig, S4 Fig), we hypothesized four nicking sites

(arrowheads in Fig 4D) in *attL* and *attR* on pSEA2 and chromosome 1 in strain 04Ya249. The

cloned *attS* PCR products of 04Ya249 contained a 6 bp spacer sequence between C and C'

(Fig 4E). This observation supports the hypothesis that nicking occurs 6 bp upstream of the

5⊠ end of C side and 3⊠ end of C' side and/or 6 bp upstream of the 5⊠ end of C' side and 3⊠

end of C side. Two sequences, 5⊠-TTTTTT-3⊠ (*attS* type 1) and 5⊠-TTTTCT-3⊠ (*attS* type 2)

were detected at a 1:9 ratio in the spacer region between C and C' in strain 04Ya249

(examples of the Sanger sequencing trace files have been posted in figshare [42]). These

two types of spacer sequences can arise from both pSEA2 and chromosome 1 (Fig 4D).

However, only one sequence (*attS* type 1) was detected in the pSEA2-free strain LN95 (Fig

4E), indicating that strand exchange occurs only on the top strand in *attL* × *attR*

recombination on chromosome 1 (Fig 4C). The occurrence of *attS* type 1 and *attS* type 2 in

04Ya249 (Fig 4E) agrees with the top strand exchange in *attL* × *attR* recombinations on

chromosome 1 and pSEA2, respectively (Fig 4D). Collectively, these observations are

consistent with the model that SE-6945 follows Path 1 (Fig 4 A) without undergoing strand

exchange of the bottom strand in *V. alfacsensis*.

Nicking at one strand on the double-strand DNA and its circularization is often seen in

the reactions mediated by HUH endonucleases (relaxase/rolling-circle replication initiator)

[43]. To confirm that observed nicking at *attL* and *attR* is independent of HUH

endonucleases produced from plasmids or other mobile elements in *V. alfaccensis*, the

production of *attS* was additionally investigated using *E. coli* strain LN52 that carries a single

copy of SE-6945 but lacks pSEA2. *attS* was produced even in LN52 (S6 Fig C). Therefore,

strand-biased (top strand only) *attL* × *attR* recombination is likely mediated by DNA strand

exchange enzymes encoded by SE-6945.

The insertion copy number and location of SE-6945 affects β-lactam

resistance levels of the host cell

The naturally occurring strains 04Ya249 and 04Ya108 carry two copies of SE-6945 in

genomes. Furthermore, 04Ya108 also carries two copies of SE-6283. However, it remains

unclear whether increased SE copies in a genome confer an advantage to the host cell.

Three copies of SE-6945 in the transconjugant TJ249 were initially detected by genome

sequencing. Southern hybridization analysis of other JW0452 transconjugants revealed one

to two copies of SE-6945 in their genomes (S5 Fig). Location of SE-6945 insertion became

evident based on Southern hybridization for 15 among 19 transconjugants. Therefore, we

used these transconjugant strains to test whether multiple insertions of SE-6945 or its

insertion location can increase the antibiotic resistance phenotype of the transconjugants.

Sixteen *E. coli* strains with or without SE-6945 were grown in Luria-Bertani broth

with ampicillin, and then the minimum inhibitory concentrations (MICs) of ampicillin were

determined using the broth dilution method following the Clinical Laboratory Standards

Institute (CLSI) guidelines. The MIC was highest in the clone carrying three copies of SE-6945,

followed by the group of transconjugants carrying two copies and the group carrying one

copy, and the MIC was the lowest in JW0452 (Fig 5). Although within-group variation was

also observed, the increased copy number of SE-6945 in the genome generally resulted in

the increased β-lactam resistance of the host cell.

MIC of individual strain is shown in Table S1. When focused on a class of MIC 500,

there was a trend that strains with SE-6945 insertion into *insJ* show a higher MIC than

strains with insertion into *yjjNt*. Thus, the location of SEs can also affect phenotype variation,

possibly due to the differences in the expression levels of their cargo genes.

Fig 5. The copy number of SE-6945 affects the resistance level. Sixteen transconjugants and

a control strain (JW0452) were divided into four groups shown in distinct colors based on

the SE-6945 copy number in the genome. MIC of ampicillin is shown according to copy

number.

Target site preference of SE-6283

We previously reported SE-6283 insertion into the coding region of the *bcp* gene encoding

peroxiredoxin (locus_tag b2480) in *E. coli.* SE-6283 also carries a *bcp* homolog (locus_tag

VYA_19800 in AP024165.1) as if it complements the disrupted *bcp* gene in the host genome

upon insertion [29]. Genome sequencing of strain 04Ya108 in this study revealed that one

SE-6283 copy was inserted into a *bcp* gene homolog in chromosome 1 (Fig 6 A). To obtain

further insight into the target site preference of SE-6283, SE-6283-like elements in public

databases were screened using blastn. SE-6283-like SEs were identified in three other *Vibrio*

species (Fig 6 A). The sizes of SE-6283 and its related SEs ranged from 13.8 kb (SE-6283) to

41.3 kb (SE-ValK09K1 in Fig 6 A). Although the motifs C and C' of the three SE-6283-like

elements were slightly different from those of SE-6283 (see *attL* in S7 Fig and *attR* in S8 Fig),

all are inserted into *bcp* orthologs in chromosome 1: chromosome 1 (or I) of *V. alginolyticus*

strain K09K1 is incorrectly named as chromosome II in CP017919. Comparison of *attL*, *attR*,

and unoccupied *attB* found in the SEs-free genome of each *Vibrio* species suggests the

insertion of SE-6283-like elements and the 6 bp sequence into the central 6 bp putative

crossover region in *attB* (Fig 6 B). Nucleotide sequences around putative nicking sites in *attB*

for SE-6283 were G and A-rich and did not show inverted repeat structure like the case of

attB for SE-6945.

Fig 6. Location of SE-6283 and its related SEs. (A) Comparisons of genetic organization

around SE elements in six genomic locations. GenBank accession numbers and the

coordinates of genomic segments compared are shown below the strain name. Except SE-

6283 on pSEA1, all elements are inserted into a *bcp* gene, while they all carry a *bcp* gene. (B)

Alignment of *attB* sequences from different bacterial species. (C) Sequence logos showing

conservation levels of *attB*. The arrowhead indicates the putative nicking site on the top

strand.

Discussion

Vibrio is one of the major bacterial genera found in marine sediments [44], and is among the

most common microbiota of wild and farmed shrimp [45] and fish [46,47]. Several members

of the genus *Vibrio* are pathogens of fishes reared in aquaculture [48], while other subsets

of *Vibrio* species, including *V. cholerae*, *V. parahaemolyticus,* and *V. vulnificus*, which are

ubiquitous in relatively low-salinity sea water, are seafood-borne human pathogens [48,49].

Thus, *Vibrio* can be considered as a key genus linking AMR genes between aquatic

environments and human-associated environments. Indeed, the accumulation of AMR

genes in this genus has attracted increased research attention, particularly in *V. cholerae*

[50,51]. However, direct experimental evidence for AMR gene transfer from *Vibrio* species

to other human-relevant bacteria is limited.

Known genetic mechanisms of AMR gene transfer from *Vibrio* include A/C plasmids

[52,53], unclassified conjugative plasmids [54], pAQU1-type MOB$_H$-family conjugative

plasmids [55], pSEA1-type MOB$_H$-family conjugative plasmids [29], ICEs [56], IMEs [57], and

a combination of a sedentary chromosomal integron and a conjugative plasmid carrying an

integron [58]. The host range of pSEA1-type MOB$_H$ plasmids, discovered in *V. alfacsensis* (for

which only one complete genome was available until the present study), is unknown. In this

study, we investigated how a pSEA1-like plasmid can contribute to AMR gene transfer in a

laboratory setting.

In contrast to A/C plasmids, autonomous replication of pSEA2 and pSEA1 was

difficult to achieve in *E. coli,* since transconjugant selection generally yielded *E. coli* clones

carrying plasmid DNA integrated into the chromosome. The current most likely mechanism

behind this phenomenon is transposition of SE followed by homologous recombination,

which enabled AMR gene transfer beyond the plasmid's replication host range. This mode of

horizontal gene transfer may be important among marine bacteria, since the plasmid

conjugation host range is expected to be wider than the replication host range [59,60].

pSEA1 was found to carry two SEs (SE-6283 and SE-6945). However, in the previous

study [29], only transposition of SE-6283 was observed, likely because we did not recognize

SE-6945 on pSEA1 and did not intend to detect its transposition. The newly identified SE-

6945 is the smallest SE identified to date (Fig 2 and Fig 6), and the only known SE harboring

an AMR gene. As SEs can have more than one target site in a single genome, they may

contribute to microbial adaptation to the antimicrobials used at aquaculture sites, with

diverse mechanisms beyond mediating the horizontal transmission of AMR genes, such as

increasing the resistance level or gene redundancy preceding evolutionary innovation of an

AMR gene [61].

The discovery of SE-6945 highlights four potential core genes (*intA*, CDS2, *intB*, and

CDS4) present in SEs. However, the specific roles of these gene products in their

transposition steps remain to be determined. Among the transposon moves classified by

Curcio and Derbyshire [15], 'copy-out' is currently the most consistent with our

observations of SEs regarding the first step of its location change. One model for strand

exchange on single-strand DNA mediated by tyrosine recombinase, leading to a single-

strand circle, has been proposed for Integron gene cassette excision (replicative resolution

model in [41]). In the replicative resolution model, strand exchange between two folded

attC bottom strands and the following replication generates a gene cassette circle and an

empty *attC* [40]. SE circle formation barely accompanies the generation of empty *attB* (Fig 4

and [29]). Thus the rejoining of the host DNA side at nicking sites is somehow prevented,

possibly by the action of multiple proteins involved: i.e., CDS2, CDS4 products, unlike the

movement of the gene cassette. Currently, we do not have information on the fate of the

unjointed donor molecule. Unlike DDE transposase, tyrosine recombinases usually do not

generate a free 3′ OH end and are covalently linked to the phosphate of substrate DNA (Fig

4A) [62]. Therefore, DPC repair-equivalent process [38], which might require both host and

SE-encoding factors, is necessary to achieve the IS*3*-like copy-out process. Nonetheless, we

speculate that the alternate single-strand DNA circle generation route, accompanying

chromosome loss (bottom left in Fig 4B), is a less likely route as it does not give any

advantage to target site-selective and putatively non-transferrable SEs left in dying cells.

While strand-biased *attL* × *attR* recombination is repeatedly observed for the two

SEs, not much information has been obtained regarding their integration step. While 6 bp

from the 5' end of C at *attL* on the donor molecule is incorporated into a circular SE copy,

the incorporated 6 bp tends to be placed next to the 3' end of C' at *attR* after insertion into

the target site (see *attL* on pSEA2 and *attRs* on TJ249, and previously detected SE-6283

transposition from pSEA1 to *bcp* in *E. coli*). Thus, there seems to be a strict rule in the order

of strand exchange at the integration step, but determining the precise order of nicking and

nicking sites on the two strands during the integration step requires further investigations.

A notable difference between IS and SE is the strong strand bias unique to SE upon

strand exchange: IS can generate a single-strand bridge using either of the two strands (but

not both simultaneously) [63,64], while SEs use only top strand. The dissimilarity between

SE and ICE/IMEs is that ICE/IMEs encode the recombination directionality factor Xis [24],

while SEs do not encode Xis homologs. CDS2 or CDS4 products may function exclusively in

either the initial circularization step or the integration as the recombination directionality

factor. Another dissimilarity between SE and ICE/IME is that the former does not encode

HUH endonuclease. There is still a possibility that SEs carry *oriT* recognized by a family of

relaxase and are mobilized like CIME [20], though we could not identify *oriT* in SE-6945 and

SE-6283 by local similarity search alone.

We propose that interplay between SEs and MDR conjugative plasmids contribute to

AMR gene transmission in marine bacteria. Further biochemical and bioinformatic studies

on SEs are needed to reveal the mechanisms of gene transfer among the genus *Vibrio* and

other aquaculture-associated bacteria.

To refer to individual SE members, we used digits that are compatible with the

existing transposon database [65]. However, further debate is needed regarding handling

SEs in mobile DNA databases.

Materials and methods

Strains, plasmids, and culture media

Strains and plasmids used in the study are listed in Table 1. Strain 04Ya108 was previously

identified as *Vibrio ponticus* based on 16S rRNA gene sequence similarity. However,

determination of the complete sequence in this study revealed that strain 04Ya108 shows

>96% average nucleotide identity to *V. alfacsensis* strain CAIM 1831 (DSM 24595) (S1 Fig)

[66]. Therefore, strain 04Ya108 was newly classified as *V. alfacsensis*. Strain LN95 is a pSEA2-

free tetracycline-susceptible derivative of strain 04Ya249. This strain was generated through

a repeated batch culture of 04Ya249, and subsequent single-colony isolation. The absence

of pSEA2 in LN95 was confirmed by next-generation sequencing.

Table 1. Strains and plasmids used.

Strains or plasmids	Relevant genotype, phenotype, descriptions[a c]	Reference
E. coli		
DH5α	F⁻, Φ80d*lacZ*ΔM15, Δ(*lacZYA-argF*)U169, *deoR*, *recA*1, *endA*1, *hsdR*17(r$_K^-$, m$_K^+$), *phoA*, *supE*44, λ⁻, *thi*-1, *gyrA*96, *relA*1	Takara Bio Inc., Shiga, Japan
JW0452	λ- *rph-1* Δ(*rhaD-rhaB*)568 *hsdR*514 Δ*lacZ*4787(::*rrnB-3*) Δ(*araD-araB*)567 Δ*acrA*748::*kan*; Km^r	[67]
JW0452rif	Spontaneous rifampicin-resistant mutant of JW0452	This study
W0452Δ*recA*	JW0452Δ*recA*::*cat*; Km^r, Cm^r	This study
JW0452Δ*recA*rif	Spontaneous rif resistant mutant of JW0452Δ*recA*; Km^r, Cm^r, Rif^r	This study

TJ249	JW0452 *insJ*::SE-6945::pSEA2 *yjjNt*::SE-6945; Ap^r Tc^r Ery^r Cm^r	This study
LN28	JW0452 chr::pSEA2^b; Ap^r Tc^r Ery^r Cm^r	This study
LN29	JW0452 *yjjNt*::SE-6945::pSEA2; Ap^r Tc^r Ery^r Cm^r	This study
LN30	JW0452 *yjjNt*::SE-6945::pSEA2; Ap^r Tc^r Ery^r Cm^r	This study
LN32	JW0452 chr::pSEA2; Ap^r Tc^r Ery^r Cm^r	This study
LN33	JW0452 *insJ*::SE-6945::pSEA2; Ap^r Tc^r Ery^r Cm^r	This study
LN34	JW0452 *insJ*::SE-6945::pSEA2; Ap^r Tc^r Ery^r Cm^r	This study
LN35	JW0452 *insJ*::SE-6945::pSEA2; Ap^r Tc^r Ery^r Cm^r	This study
LN36	JW0452 chr::pSEA2^b; Ap^r Tc^r Ery^r Cm^r	This study
LN37	JW0452 *insJ*::SE-6945::pSEA2; Ap^r Tc^r Ery^r Cm^r	This study
LN52	JW0452 *yjjNt*::SE-6945; Ap^r	This study
LN52rif	Spontaneous rifampicin-resistant mutant of LN52; Rif^r, Ap^r	This study
LN58	JW0452 *insJ*::SE-6945::pSEA2; Ap^r Tc^r Ery^r Cm^r	This study
LN82	JW0452 *yjjNt*::SE-6945::pSEA2; Ap^r Tc^r Ery^r Cm^r	This study
LN84	JW0452 *insJ*::SE-6945::pSEA2; Ap^r Tc^r Ery^r Cm^r	This study
LN86	JW0452 chr::pSEA2; Ap^r Tc^r Ery^r Cm^r	This study
LN88	JW0452 *yjjNt*::SE-6945::pSEA2; Ap^r Tc^r Ery^r Cm^r	This study
LN89	JW0452 *insJ*::SE-6945::pSEA2; Ap^r Tc^r Ery^r Cm^r	This study
LN90	JW0452 *insJ*::SE-6945::pSEA2; Ap^r Tc^r Ery^r Cm^r	This study
LN91	JW0452 *yjjNt*::SE-6945::pSEA2; Ap^r Tc^r Ery^r Cm^r	This study
Vibrio alfacsensis		
04Ya249	Aquaculture isolate harboring conjugative plasmid pSEA2; Tc^r, Ap^r, Ery^r Cm^r	[26]
04Ya108	Aquaculture isolate harboring conjugative plasmid pSEA1; Tc^r, Ap^r, Ery^r Cm^r	[26]
LN95	pSEA2 free derivative of strain 04Ya249; Ap^r	This study
Plasmids		
pGEM-T	Cloning vector; pUC replicon, Ap^r	Promega, Madison, WI, USA
pGEM-attS	pGEM-T carrying the 206 bp *attS* fragment; pUC replicon, Ap^r	This study
pGEM-int	pGEM-T carrying the 103 bp *intA* fragment; pUC replicon, Ap^r	This study
pGEM-gyrB	pGEM-T carrying the 110 bp *gyrB* fragment from strain 04Ya249; pUC replicon, Ap^r	This study
pKD46	Lambda-Red recombinase expression plasmid; pSC101ts replicon, Ap^r	[68]
pKD3	Donor of the *cat* gene for the Lambda-Red recombination experiment; R6K replicon, Cm^r	[68]

a. Antimicrobial resistance phenotypes (ʳ) listed were confirmed by experiments.

Abbreviations are as follows: Ap, ampicillin; Tc, tetracycline: Ery, erythromycin; Cm,

chloramphenicol; Rif, rifampicin; *kan*, kanamycin resistance gene. *yjjNt* denotes the putative

transcription termination region for the *yjjN* gene.

b. Integration of pSEA2 without generating a new junction of SE and chromosome.

Integration location of pSEA2 is unknown.

c. 19 transconjugants listed (TJ249, LN28, LN29, LN30, LN32, LN33, LN34, LN35, LN36, LN37,

LN52, LN58, LN82, LN84, LN86, LN88, LN89, LN90, LN91) were obtained from 19

independent mating experiments. Selection conditions used in transconjugant screening are

as follows: Tc 42 °C for LN28, LN29 LN30, LN32, LN33, LN34, LN35, LN36, and LN37; Ap 42 °C

for LN52; Ap Tc 42 °C for LN58 LN82, LN84, LN86, LN88, LN89, LN90, and LN91; Ery 42 °C for

TJ249.

532   *E. coli* strain LN52, which carries a single copy of SE-6945, is a JW0452

transconjugant obtained by ampicillin selection at 42 °C. TJ249 is also a JW0452

transconjugant obtained by erythromycin selection at 42 °C. The presence of a single SE-

6945 copy in the LN52 genome was confirmed by Southern hybridization (S5 Fig). The *recA*-

null mutant of JW0452, JW0452Δ*recA*, was constructed using the lambda-Red method [68].

A DNA fragment containing the 5 and 3 sequences of the *recA* gene and a

chloramphenicol resistance gene was amplified by PCR using primers YO-175 and

RecA_stop_primingsite_2 (Table 2) and plasmid pKD3 [68] as a template. The PCR products

(700 ng) were introduced into electrocompetent cells of JW0452 carrying pKD46 by

electroporation using a Gene Pulser Xcell™ (BioRad, Hercules, CA, USA). The occurrence of

recombination at the expected site was confirmed by PCR using primers CAT-584 and

BW25113_2815723f (Table 2). The absence of the *recA* gene was also confirmed by PCR

using primers LN192_recA1 and LN193_recA2.

Vibrio alfacsensis strains were cultured in BD Bacto™ brain heart infusion medium

(BD237500; Becton, Dickinson, and Company, Franklin Lakes, NJ, USA) supplemented with

up to 2% NaCl at 25 °C. *E. coli* strains were cultured in BD Difco™ LB Broth, Miller

(BD244520; Becton, Dickinson, and Company) at 37 °C or 42 °C. BD Difco™ Mueller Hinton

Broth (BD 275730; Becton, Dickinson, and Company) was used for antibiotic susceptibility

testing of *E. coli*. BD Difco™ Marine Broth 2216 (BD279110; Becton, Dickinson, and

Company) was used for filter mating. Solid media were prepared by adding 1.5% agar to the

broth. Antibiotics were added to the medium at the following concentrations when

required: erythromycin (Nacalai Tesque, Kyoto, Japan), 100 µg/ml; tetracycline (Nacalai

Tesque), 10 µg/ml; rifampicin (Sigma-Aldrich, St. Louis, MO, USA), 50 µg/ml; ampicillin

(Nacalai Tesque), 100 µg/ml.

Table 2. Oligonucleotides used.

Name	Sequence (5′ to 3′)	Purpose
LN112	GGGTTACCTTCCCAATGCGT	Southern hybridization probe for SE-6945 *intA*
LN113	CGACTGTTGGTAGCGACTGT	Southern hybridization probe for SE-6945 *intA*
LN_142_junction2	AAGATGGTAAAAGTGTTCCA	Detection of *attS* (206 bp) by qPCR and *attL* (304 bp) by standard PCR
LN_143_junction2	TTTGTGTGTAGCCCTTGTG	Detection of *attS* (206 bp) by

		qPCR and *attR* (248 bp) by standard PCR
LN_150_intA2	GGTTATGTGGAGAAGTTGCC	Detection of *intA* (103 bp) by qPCR
LN_151_intA2	TGAGTTCGGTTTCTTGCTTC	Detection of *intA* (103 bp) by qPCR
LN_181_Valc_chr1_attB_L	CGAGGGTAAAGTGCCAACAT	Detection of chromosomal *attB* (345 bp if empty, 7.4 kb if occupied) and *attL* (304 bp) by standard PCR
LN183_Valc_chr1_attB_R2	ACATCAGCAGGAGTTAGTTG	Detection of chromosomal *attB* (345 bp, 7.4 kb) and *attR* (248 bp) by standard PCR
LN184_04Ya249_gyrBf1	AACAGAATTGCACCCAGAAG	Detection of *gyrB* (110 bp) by qPCR
LN185_04Ya249_gyrBr1	GAAGACCGCCTGATACTTTG	Detection of *gyrB* (110 bp) by qPCR
YO-175	CAGAACATATTGACTATCCGGTATTACCCGGCATGACAGGAGTAAAAATGTGTAGGCTGGAGCTGCTTCG	Amplification of *cat* gene for Lambda-Red recombination
RecA_stop_primingsite_2	ATGCGACCCTTGTGTATCAAACAAGACGATTAAAAATCTTCGTTAGTTTCCATATGAATATCCTCCTTA	Amplification of *cat* gene for Lambda-Red recombination
CAT-584	AAGCCATCACAAACGGCATG	Confirmation of *cat* gene insertion

BW25113_2815723f	AATACGCGCAGGTCCATAAC	Confirmation of *cat* gene insertion
LN192_recA1	GTTCCATGGATGTGGAAACC	Confirmation of *recA* deletion
LN193_recA2	ATATCGACGCCCAGTTTACG	Confirmation of *recA* deletion

Conjugation

The donor *Vibrio* strain and *E. coli* recipient strains were grown overnight at 25 °C and 37 °C,

respectively. A 500 µl aliquot of each culture was mixed, centrifuged, and resuspended in 50

µl of Luria-Bertani broth. The cell mixture was spotted on a 0.45 µm pore-size nitrocellulose

filter (Merck, Millipore Ltd., Tullagreen, Ireland) placed on marine broth agar, and allowed

to mate for 24 h at 25 °C. After incubation, the cell mixture on the filter was serially diluted

in 1× phosphate-buffered saline, and then 100 µl of the mixture was plated on an

appropriate agar medium to measure the CFU. The transconjugant strain TJ249 was

obtained by the first mating experiment of strain 04Ya249 with *E. coli* strain JW0452 as the

donor and recipient, respectively, followed by Ery selection at 42 °C. A high temperature

(42 °C) was used to suppress the growth of the *Vibrio alfacsensis* strain on LB agar. After

genome sequence determination of TJ249, more than 17 independent mating experiments

were performed to obtain insights into the variation of pSEA2 integration patterns using

JW0452 as the recipient, and transconjugants were screened at 42 °C in the presence of Tc

alone, Ap alone, or both Ap and Tc. The AMR phenotype of each transconjugant was further

determined after single-colony isolation using Ery, Cm, Ap, and Tc (Table 1).

Quantitative mating assays (Fig 3) were performed to address the involvement of

homologous recombination in pSEA2 integration using strains JW0452rif, JW0452Δ*recA*rif,

and LN95rif as recipients. Selection conditions for transconjugants were Rif Ap Tc 42 °C or

Rif Ap 42 °C. The *Vibrio* donor strain was selected on brain heart infusion agar with 2% NaCl

supplemented with Tc at 25 °C. CFUs of donor, recipient, and transconjugant were

determined after 24 hours of incubation.

PCR experiments

The PCR detection of *attL*, *attR,* and *attS* was performed using TaKaRa Ex Taq (TaKaRa Bio

Inc., Kusatsu, Japan), and genomic DNA of 04Ya249, LN95, and LN52 prepared using QIAGEN

DNeasy Blood & Tissue kit (QIAGEN GmbH, Hilden, Germany). PCR premix was prepared in a

50 µl scale, and 1 µl of the template (10 ng/µl) was added to the reaction mix throughout

the study. The PCR cycle used was 94 °C for 1 min, followed by 35 cycles of 94 °C for 30 s

and 60 °C for 30 s. As the empty *attB* was not amplified using TaKaRa Ex Taq, we conducted

additional Long-PCR using LA Taq polymerase (TaKaRa Bio Inc.) to detect occupied *attB* and

confirm the functionality of primers used for detection of empty *attB*. The PCR cycles used

for Long-PCR were 1 cycle of 94 °C for 1 min, followed by 35 cycles of 95 °C for 1 min, 65 °C

for 1 min, and 72 °C for 1 min. PCR products of *attS* were further cloned using the pGEM-T

easy vector system (Promega, Madison, WI, USA) and DH5α competent cells to analyze the

sequence variation of PCR products.

Quantitative PCR was performed using THUNDERBIRD® SYBR qPCR Mix (Toyobo,

Osaka, Japan) and a CFX connect Real-Time system (BioRad, Hercules, CA, USA) and a two-

step PCR protocol of 40 cycles of 95 °C for 5 s and 60 °C for 30 s. Target quantity was

estimated based on a standard curve of the control plasmid DNA (pGEM-*gyrB*, pGEM-*intA*,

pGEM-*attS*) constructed by TA cloning. Primers used and their purposes are listed in Table 2

and Fig 4 legend.

Southern hybridization

The insertion copy number and insertion location of SE-6945 in transconjugants (JW0452

derivatives) were analyzed by Southern hybridization using the 5ʹ end of *intA*, amplified

using primers LN112 and LN113 (Table 2) and the PCR DIG Synthesis Kit (Roche, Basel,

Switzerland) as a probe. Genomic DNA (2.5 µg) was double digested with either *Nde*I and

*Sph*I or *Nde*I and *Hin*dIII (New England Biolabs, Ipswich, MA, USA), then electrophoresed in

0.8% agarose gel. Separated DNA fragments were transferred to the nitrocellulose

membranes (Amersham Hybond-N+; GE Healthcare UK Ltd., Buckinghamshire, UK) by

capillary transfer. The probe was detected using CDT-star® (Roche). Restriction map for the

DNA segments around SE-6945 insertion locations and the expected size of DNA segments

with probe signal are shown in S5 Fig A.

Antimicrobial susceptibility testing

615   To examine the antimicrobial susceptibility of transconjugants harboring one to three SE-

616   6945 copies, the MIC of ampicillin was determined using the broth dilution method in 96-

617   well microtiter plate format according to standard M07 of the CLSI [69]. The antibiotic

618   concentrations tested were 1000, 500, 250, 125, 62.5, 31.25, 16, 8, 4, and 2 µg/ml. The test

619   plates were incubated at 35 °C for 24 h.

Genome sequencing

Genomic DNA was extracted from 250–500 µl of *V. alfacsensis* strains 04Ya249, 04Ya108, or

E. coli strain TJ249 culture using the QIAGEN DNeasy Blood & Tissue kit (QIAGEN GmbH).

The extracted genomic DNA was sequenced on the PacBio RS II platform using P6C4

chemistry at Macrogen (Tokyo, Japan) in 2016. Genome assembly was conducted using

HGAP v.3 [70] for *Vibrio* strains and Flye v 2.8.3-b1695 for *E. coli* TJ249 [71]. Reads were

obtained at >120× coverage for the chromosomes of each strain. Reads and the genome

sequence of TJ249 have been posted to figshare [72]. Illumina reads of the pSEA2-free strain

LN95 were obtained using the TruSeq PCR-free library and NovaSeq 6000 platform at

NovogeneAIT Genomics Singapore Pte., Ltd. (Singapore) to confirm the loss of pSEA2 and to

identify the circular form of SE-6945. The mapping results have been posted to figshare [42].

Genomes were compared using MUMmer3.23 [73] and GenomeMatcher [74]. AMR genes in

the assembly were searched using AMRFinderPlus [75]. The average nucleotide identity was

determined using fastANI [76].

Searches for SE-6945 and SE-6283-related SEs

Nucleotide sequences of the SE core gene region (*intA*, CDS2, *intB*, CDS4) (6432 bp for SE-

6283, 5541 bp for SE-6945) were used as queries in blastn searches against the NCBI

nucleotide collection (nr/nt) database (August 3, 2021). Hit subjects showing > 85 %

nucleotide identity with > 99% query coverage were regarded as close relatives of SE-6945

or SE-6283, and were further investigated as to their insertion locations.

Data availability

Sequence data generated in this study and their availability are summarized in Table S2. The

complete genome sequences of strains 04Ya108 and 04Ya249 were submitted to

GenBank/DDBJ/EMBL under accession numbers AP024165 (chr1 of 04Ya108), AP024166

(chr2 of 04Ya108), AP024167 (pSEA1 of 04Ya108), AP024168 (pYa108 of 04Ya108),

AP019849 (chr1 of 04Ya249), AP019850 (chr2 of 04Ya249), AP019851 (pSEA2 of 04Ya249),

and AP019852 (pVA249 of 04Ya249). The raw reads for strains 04Ya108, 04Ya249, and LN95

are available from Sequence Read Archive under accession numbers DRA011098,

DRA008632, and DRA011762, respectively.

Acknowledgments

We thank the National Bioresource Project of the National Institute of Genetics, Japan, for

providing the *E. coli* strains BW25113, JW2669, and JW0452. We thank Atsushi Ota and

Fumito Maruyama at Hiroshima University for their support for the annotation on the TJ249

assembly. We thank Yuichi Otsuka at Saitama University for helpful discussions and Yuta

Sugimoto at Ehime University for experimental support. We thank Kenji K Kojima at Genetic

Information Research Institute for the discussion on transposon classification, and Yasukazu

Daigaku at Tohoku University for the discussion on DNA protein cross link repair. We thank

Editage for language editing (job number LINON_4).

References

664   1.   Cabello FC, Godfrey HP, Tomova A, Ivanova L, Dölz H, Millanao A et al. (2013)

665   Antimicrobial use in aquaculture re-examined: its relevance to antimicrobial resistance

666   and to animal and human health. Environ Microbiol 15: 1917-1942.

2.   Cabello FC, Godfrey HP, Buschmann AH, Dölz HJ (2016) Aquaculture as yet another

environmental gateway to the development and globalisation of antimicrobial

resistance. Lancet Infect Dis 16: e127-e133.

3.   Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG et al. (2012)

Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an

international expert proposal for interim standard definitions for acquired resistance.

Clin Microbiol Infect 18: 268-281.

4.   Karakonstantis S, Kritsotakis EI, Gikas A (2020) Pandrug-resistant Gram-negative

bacteria: a systematic review of current epidemiology, prognosis and treatment

options. J Antimicrob Chemother 75: 271-282.

5.   Rodríguez-Beltrán J, DelaFuente J, León-Sampedro R, MacLean RC, San Millán Á (2021)

Beyond horizontal gene transfer: the role of plasmids in bacterial evolution. Nat Rev

Microbiol 19: 347-359.

6.   Johnson CM, Grossman AD (2015) Integrative and Conjugative Elements (ICEs): What

They Do and How They Work. Annu Rev Genet 49: 577-601.

7.   Yano H, Rogers LM, Knox MG, Heuer H, Smalla K, Brown CJ et al. (2013) Host range

diversification within the IncP-1 plasmid group. Microbiology (Reading) 159: 2303-

2315.

8.   Dahmane N, Libante V, Charron-Bourgoin F, Guédon E, Guédon G, Leblond-Bourget N

et al. (2017) Diversity of integrative and conjugative elements of *Streptococcus*

salivarius and their intra- and interspecies Transfer. Appl Environ Microbiol 83:

e00337-17.

9.   Cabezón E, Lanka E, de la Cruz F (1994) Requirements for mobilization of plasmids

RSF1010 and ColE1 by the IncW plasmid R388: *trwB* and RP4 *traG* are interchangeable.

J Bacteriol 176: 4455-4458.

10. Carraro N, Matteau D, Luo P, Rodrigue S, Burrus V (2014) The master activator of

IncA/C conjugative plasmids stimulates genomic islands and multidrug resistance

dissemination. PLoS Genet 10: e1004714.

11. Daccord A, Ceccarelli D, Burrus V (2010) Integrating conjugative elements of the

SXT/R391 family trigger the excision and drive the mobilization of a new class of *Vibrio*

genomic islands. Mol Microbiol 78: 576-588.

12. Garcillán-Barcia MP, Francia MV, de la Cruz F (2009) The diversity of conjugative

relaxases and its application in plasmid classification. FEMS Microbiol Rev 33: 657-687.

13. Villa L, Carattoli A (2020) Plasmid Typing and Classification. Methods Mol Biol 2075:

309-321.

14. Roberts AP, Chandler M, Courvalin P, Guédon G, Mullany P, Pembroke T et al. (2008)

Revised nomenclature for transposable genetic elements. Plasmid 60: 167-173.

15. Curcio MJ, Derbyshire KM (2003) The outs and ins of transposition: from mu to

kangaroo. Nat Rev Mol Cell Biol 4: 865-877.

16. Bouvier M, Demarre G, Mazel D (2005) Integron cassette insertion: a recombination

process involving a folded single strand substrate. EMBO J 24: 4356-4367.

17. Val ME, Bouvier M, Campos J, Sherratt D, Cornet F, Mazel D et al. (2005) The single-

stranded genome of phage CTX is the form used for integration into the genome of

Vibrio cholerae. Mol Cell 19: 559-566.

18. Doublet B, Boyd D, Mulvey MR, Cloeckaert A (2005) The Salmonella genomic island 1 is

an integrative mobilizable element. Mol Microbiol 55: 1911-1924.

19.   Pavlovic G, Burrus V, Gintz B, Decaris B, Guédon G (2004) Evolution of genomic islands

by deletion and tandem accretion by site-specific recombination: ICE*St1*-related

elements from *Streptococcus thermophilus*. Microbiology (Reading) 150: 759-774.

20.   Achard A, Leclercq R (2007) Characterization of a small mobilizable transposon,

MTnSag1, in Streptococcus agalactiae. J Bacteriol 189: 4328-4331.

21.   Guédon G, Libante V, Coluzzi C, Payot S, Leblond-Bourget N (2017) The obscure world

of integrative and mobilizable elements, highly widespread elements that pirate

bacterial conjugative systems. Genes (Basel) 8: E337.

22.   Caparon MG, Scott JR (1989) Excision and insertion of the conjugative transposon

Tn*916* involves a novel recombination mechanism. Cell 59: 1027-1034.

23.   Auchtung JM, Lee CA, Monson RE, Lehman AP, Grossman AD (2005) Regulation of a

Bacillus subtilis mobile genetic element by intercellular signaling and the global DNA

damage response. Proc Natl Acad Sci U S A 102: 12554-12559.

24.   Burrus V, Waldor MK (2003) Control of SXT integration and excision. J Bacteriol 185:

5045-5054.

25.   Naito M, Ogura Y, Itoh T, Shoji M, Okamoto M, Hayashi T et al. (2016) The complete

genome sequencing of *Prevotella intermedia* strain OMA14 and a subsequent fine-

scale, intra-species genomic comparison reveal an unusual amplification of conjugative

and mobile transposons and identify a novel *Prevotella*-lineage-specific repeat. DNA

Res 23: 11-19.

26.   Nonaka L, Ikeno K, Suzuki S (2007) Distribution of tetracycline resistance gene, *tet(M)*,

in Gram-positive and Gram-negative bacteria isolated from sediment and seawater at

a coastal aquaculture site in Japan. Microbes Environ 22: 355-364.

27.   Nonaka L, Maruyama F, Miyamoto M, Miyakoshi M, Kurokawa K, Masuda M (2012)

737   Novel conjugative transferable multiple drug resistance plasmid pAQU1 from

738   *Photobacterium damselae* subsp. *damselae* isolated from marine aquaculture

739   environment. Microbes Environ 27: 263-272.

740   28.   Nonaka L, Maruyama F, Suzuki S, Masuda M (2015) Novel macrolide-resistance genes,

741   *mef*(C) and *mph*(G), carried by plasmids from *Vibrio* and *Photobacterium* isolated from

742   sediment and seawater of a coastal aquaculture site. Lett Appl Microbiol 61: 1-6.

743   29.   Nonaka L, Yamamoto T, Maruyama F, Hirose Y, Onishi Y, Kobayashi T et al. (2018)

744   Interplay of a non-conjugative integrative element and a conjugative plasmid in the

745   spread of antibiotic resistance via suicidal plasmid transfer from an aquaculture *Vibrio*

746   isolate. PLoS One 13: e0198613.

747   30.   Neela FA, Nonaka L, Suzuki S (2007) The diversity of multi-drug resistance profiles in

748   tetracycline-resistant Vibrio species isolated from coastal sediments and seawater. J

749   Microbiol 45: 64-68.

750   31.   Alcock BP, Raphenya AR, Lau TTY, Tsang KK, Bouchard M, Edalatmand A et al. (2020)

751   CARD 2020: antibiotic resistome surveillance with the comprehensive antibiotic

752   resistance database. Nucleic Acids Res 48: D517-D525.

753   32.   Pei J, Kim BH, Grishin NV (2008) PROMALS3D: a tool for multiple protein sequence and

754   structure alignments. Nucleic Acids Res 36: 2295-2300.

755   33.   Nunes-Düby SE, Kwon HJ, Tirumalai RS, Ellenberger T, Landy A (1998) Similarities and

756   differences among 105 members of the Int family of site-specific recombinases.

757   Nucleic Acids Res 26: 391-406.

758   34.   Landy A (1989) Dynamic, structural, and regulatory aspects of lambda site-specific

759   recombination. Annu Rev Biochem 58: 913-949.

760   35.   Schneider TD, Stephens RM (1990) Sequence logos: a new way to display consensus

761   sequences. Nucleic Acids Res 18: 6097-6100.

762   36.   Stokes HW, O'Gorman DB, Recchia GD, Parsekhian M, Hall RM (1997) Structure and

763   function of 59-base element recombination sites associated with mobile gene

764   cassettes. Mol Microbiol 26: 731-745.

765   37.   Stokes HW, Hall RM (1989) A novel family of potentially mobile DNA elements

766   encoding site-specific gene-integration functions: integrons. Mol Microbiol 3: 1669-

767   1683.

768   38.   Zhang H, Xiong Y, Chen J (2020) DNA-protein cross-link repair: what do we know now.

769   Cell Biosci 10: 3.

770   39.   Chandler M, Fayet O, Rousseau P, Ton Hoang B, Duval-Valentin G (2015) Copy-out-

771   paste-in transposition of IS*911*: a major transposition pathway. Microbiol Spectr 3:

772   40.   Collis CM, Hall RM (1992) Site-specific deletion and rearrangement of integron insert

773   genes catalyzed by the integron DNA integrase. J Bacteriol 174: 1574-1585.

774   41.   Escudero JA, Loot C, Nivina A, Mazel D (2015) The integron: adaptation on demand.

775   Microbiol Spectr 3: MDNA3-0019.

776   42.   Yano H, Nonaka L (2021) Illumina reads analysis of *Vibrio alfacsensis* pSEA2-free strain

777   LN95. URL: https://doiorg/106084/m9figshare14338883

778   43.   Chandler M, de la Cruz F, Dyda F, Hickman AB, Moncalian G, Ton-Hoang B (2013)

779   Breaking and joining single-stranded DNA: the HUH endonuclease superfamily. Nat

780   Rev Microbiol 11: 525-538.

781   44.   Chen J, McIlroy SE, Archana A, Baker DM, Panagiotou G (2019) A pollution gradient

782   contributes to the taxonomic, functional, and resistome diversity of microbial

783   communities in marine sediments. Microbiome 7: 104.

784   45.   Cornejo-Granados F, Lopez-Zavala AA, Gallardo-Becerra L, Mendoza-Vargas A, Sánchez

785   F, Vichido R et al. (2017) Microbiome of Pacific Whiteleg shrimp reveals differential

786   bacterial community composition between Wild, Aquacultured and AHPND/EMS

787   outbreak conditions. Sci Rep 7: 11783.

788   46.   Llewellyn MS, McGinnity P, Dionne M, Letourneau J, Thonier F, Carvalho GR et al.

789   (2016) The biogeography of the atlantic salmon (*Salmo salar*) gut microbiome. ISME J

790   10: 1280-1284.

791   47.   Egerton S, Culloty S, Whooley J, Stanton C, Ross RP (2018) The Gut Microbiota of

792   Marine Fish. Front Microbiol 9: 873.

793   48.   Thompson FL, Iida T, Swings J (2004) Biodiversity of Vibrios. Microbiol Mol Biol Rev 68:

794   403-431.

795   49.   Takemura AF, Chien DM, Polz MF (2014) Associations and dynamics of *Vibrionaceae* in

796   the environment, from the genus to the population level. Front Microbiol 5: 38.

797   50.   Verma J, Bag S, Saha B, Kumar P, Ghosh TS, Dayal M et al. (2019) Genomic plasticity

798   associated with antimicrobial resistance in *Vibrio cholerae*. Proc Natl Acad Sci U S A

799   116: 6226-6231.

800   51.   Osei Sekyere J, Reta MA (2020) Genomic and resistance epidemiology of Gram-

801   negative bacteria in Africa: a systematic review and phylogenomic analyses from a one

802   health perspective. mSystems 5: e00897-20.

803   52.   Carraro N, Sauvé M, Matteau D, Lauzon G, Rodrigue S, Burrus V (2014) Development

804   of pVCR94ΔX from *Vibrio cholerae*, a prototype for studying multidrug resistant IncA/C

805   conjugative plasmids. Front Microbiol 5: 44.

806   53.   Folster JP, Katz L, McCullough A, Parsons MB, Knipe K, Sammons SA et al. (2014)

807   Multidrug-resistant IncA/C plasmid in *Vibrio cholerae* from Haiti. Emerg Infect Dis 20:

808   1951-1953.

809   54.   Liu M, Wong MH, Chen S (2013) Molecular characterisation of a multidrug resistance

810   conjugative plasmid from *Vibrio parahaemolyticus*. Int J Antimicrob Agents 42: 575-

811   579.

812   55.   Nonaka L, Maruyama F, Onishi Y, Kobayashi T, Ogura Y, Hayashi T et al. (2014) Various

813   pAQU plasmids possibly contribute to disseminate tetracycline resistance gene *tet*(M)

814   among marine bacterial community. Front Microbiol 5: 152.

815   56.   Beaber JW, Burrus V, Hochhut B, Waldor MK (2002) Comparison of SXT and R391, two

816   conjugative integrating elements: definition of a genetic backbone for the mobilization

817   of resistance determinants. Cell Mol Life Sci 59: 2065-2070.

818   57.   Carraro N, Rivard N, Ceccarelli D, Colwell RR, Burrus V (2016) IncA/C conjugative

819   plasmids mobilize a new family of multidrug resistance islands in clinical *Vibrio*

820   *cholerae* non-O1/non-O139 isolates from Haiti. mBio 7: e00509-16.

821   58.   Rowe-Magnus DA, Guerout AM, Mazel D (2002) Bacterial resistance evolution by

822   recruitment of super-integron gene cassettes. Mol Microbiol 43: 1657-1669.

823   59.   Encinas D, Garcillán-Barcia MP, Santos-Merino M, Delaye L, Moya A, de la Cruz F

824   (2014) Plasmid conjugation from proteobacteria as evidence for the origin of

825   xenologous genes in cyanobacteria. J Bacteriol 196: 1551-1559.

826   60.   Kishida K, Inoue K, Ohtsubo Y, Nagata Y, Tsuda M (2017) Host range of the conjugative

827   transfer system of IncP-9 naphthalene-catabolic plasmid NAH7 and characterization of

828   its *oriT* region and relaxase. Appl Environ Microbiol 83: e02359-16.

829   61.   Elliott KT, Cuff LE, Neidle EL (2013) Copy number change: evolving views on gene

830   amplification. Future Microbiol 8: 887-899.

831   62.   Jayaram M, Ma CH, Kachroo AH, Rowley PA, Guga P, Fan HF et al. (2015) An overview

832   of tyrosine site-specific recombination: from an Flp perspective. Microbiol Spectr 3:

833   63.   Polard P, Chandler M (1995) An *in vivo* transposase-catalyzed single-stranded DNA

834   circularization reaction. Genes Dev 9: 2846-2858.

835   64.   Rousseau P, Loot C, Turlan C, Nolivos S, Chandler M (2008) Bias between the left and

836   right inverted repeats during IS*911* targeted insertion. J Bacteriol 190: 6111-6118.

837   65.   Tansirichaiya S, Rahman MA, Roberts AP (2019) The Transposon Registry. Mob DNA

838   10: 40.

839   66.   Gomez-Gil B, Roque A, Chimetto L, Moreira APB, Lang E, Thompson FL (2012) *Vibrio*

840   *alfacsensis* sp. nov., isolated from marine organisms. Int J Syst Evol Microbiol 62: 2955-

841   2961.

842   67.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M et al. (2006) Construction of

843   *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. Mol

844   Syst Biol 2: 2006.0008.

845   68.   Datsenko KA, Wanner BL (2000) One-step inactivation of chromosomal genes in

846   *Escherichia coli* K-12 using PCR products. Proc Natl Acad Sci U S A 97: 6640-6645.

847   69.   Wayne PA (2018) Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria

848   That Grow Aerobically. 11th ed. CLSI standard M07. Clinical and Laboratory Standards

849   Institute.

850   70.   Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C et al. (2013)

851   Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing

852   data. Nat Methods 10: 563-569.

853   71.   Kolmogorov M, Yuan J, Lin Y, Pevzner PA (2019) Assembly of long, error-prone reads

854   using repeat graphs. Nat Biotechnol 37: 540-546.

855   72.   Yano H, Nonaka L (2021) Complete genome of transconjugant TJ249. URL:

856   https://doiorg/106084/m9figshare13332467

857   73.   Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C et al. (2004)

858   Versatile and open software for comparing large genomes. Genome Biol 5: R12.

859   74.   Ohtsubo Y, Ikeda-Ohtsubo W, Nagata Y, Tsuda M (2008) GenomeMatcher: a graphical

860   user interface for DNA sequence comparison. BMC Bioinformatics 9: 376.

861   75.   Feldgarden M, Brover V, Haft DH, Prasad AB, Slotta DJ, Tolstoy I et al. (2019) Validating

862   the AMRFinder tool and resistance gene database by using antimicrobial resistance

863   genotype-phenotype correlations in a collection of isolates. Antimicrob Agents

864   Chemother 63: e00483-19.

865   76.   Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S (2018) High

866   throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries.

867   Nat Commun 9: 5114.

Supporting information

S1 Fig. Comparison of the genome structure between strain 04Ya249 with 04Ya108, CAIM

1831 or TJ249. Structure comparison was performed using nucmer in MUMmer3 [73].

Purple dots indicate a match on the Watson strand (5ᐧ to 3ᐧ on the top strand in GenBank

file), and light blue indicates a match on the Crick strand (5' to 3' on the bottom strand). (B)

Average nucleotide identity (ANI) between two strains as determined by fastANI [76]. The

commands used were as follows: (A) $nucmer -minmatch

60../../data/04Ya249_submission.fas../../data/reference.fas

$mummerplot -x "[0,6000000]" -y "[0,22000000]" -postscript -p test out.delta; (B) $fastANI -

q../../data/04Ya249_submission.fas -r../../data/CAIM1831_Refseq.fas -o

04Ya249vsCAIM1831.txt

$fastANI -q../../data/04Ya249_submission.fas -r../../data/04Ya108_submission.fas -o

04Ya249vs04Ya108.txt

$fastANI -q../../data/04Ya249_submission.fas -r../../data/04Ya108_submission.fas -o

04Ya249vs04Ya108.txt

S2 Fig. Genetic map of pSEA2. (A) Locations of antimicrobial resistance (AMR) genes and SE-

6945 in pSEA2. AMR genes were inferred using AMRFinderPlus [75]. Genes are visualized

using CLC Sequence Viewer (Qiagen, Hilden Germany). Green arc denotes the region

detected as a circular contig (contig1) in Pacbio RSII reads assembly of transconjugant strain

TJ249. (B) Location of SEs in pSEA1 and pSEA2. Four SE core genes are indicated by four

distinct colors. Red pentagons are the β-lactamase gene.

S3 Fig. *attL* sequences formed in the SE-6945 insertion regions. Sequences are derived

from 04Ya249 genomes (AP019851.1, AP019849.1), TJ239 genome (*insJ*, *yjjNt*), and *Vibrio*

harveri strain WXL538 (CP045070.1) carrying a SE-6945-like element carrying a different set

of cargo genes. Sequences in red are a part of the mobile DNA unit of the SE element. 6-bp

underlined sequence is incorporated into *attS* during SE circle formation. 19 bp motif C

highlighted in yellow form an imperfect inverted repeat with the motif C' located the other

terminus of SE. Sequence highlighted in cyan is a putative transcription termination motif.

Asterisks indicate conserved nucleotides.

S4 Fig. *attR* sequences formed in the SE-6945 insertion regions. Sequences are derived

from 04Ya249 genomes (AP019851.1, AP019849.1), TJ239 genome (*insJ*, *yjjNt*), and *Vibrio*

harveri strain WXL538 (CP045070.1). Sequences in red are a part of the mobile DNA unit of

the SE element. The 6 bp underlined sequences originate from the original *attB* or SE circle,

and they are not incorporated into the *attS* of *the* SE circle.

S5 Fig. Inference of SE-6945 insertion number and its insertion location by Southern

hybridization. (A) *in silico* restriction map of DNA segments around SE-6945 insertion

locations in strain 04Ya249, TJ249, and a hypothetical *E. coli* strain carrying a pSEA2

insertion into SE-6945 integrated into *yjjNt*. Locations of SE-6945 and *intA* probe

hybridization region are indicated by the yellow square and black filled square, respectively,

above the restriction map. Horizontal black lines indicate the fragments detected by the

probe and their sizes. (B) pSEA2-free *Vibrio* strain LN95 and parental strain 04Ya249. (C)

Nineteen *E. coli* transconjugants obtained from 19 independent mating assays. Upper panel:

digestion with *Nde*I and *Sph*I. Lower panel: digestion with *Nde*I and *Hin*dIII. The color of the

strain name indicates the pattern of SE/pSEA2 insertion: blue, three SE copies (*yjjNt*::SE-

6945, *insJ*::SE-6945::pSEA2); red, two SE copies (*insJ*::SE-6945::pSEA2); purple, (*yjjNt*::SE-

6945::pSEA2); lime, one SE copy (*yjjNt*::SE-6945 or chr::pSEA2); Strains with asterisk (*) are

expected to carry pSEA2 in an unknown chromosomal location without generating a SE-

6945-chromosome junction.

S6 Fig. PCR detection of SE-6945 circle in *V. alfacsensis* strain 04Ya249 and *E. coli* strain

LN52. (A) PCR detection of chromosomal *attL*, *attR*, *attS* in 04Ya249. (B) Copy number of

attS in pSEA2 free *V. alfacsensis* strain LN95 by qPCR. Copy number is represented as

relative copy number. Primers used are shown in Table 2. (C) PCR detection of *attS* in pSEA2

free *E. coli* strain LN52.

S7 Fig. *attL* sequences formed in the SE-6283 insertion regions. Sequences are derived

from strain 04Ya108 (AP024165.1, AP024167.1), *E. coli* transconjugant TJ108W0 sequenced

in our previous study [29], and *Vibrio* strains carrying SE-6283-like elements listed in Fig 5.

Sequences in red are a part of the mobile DNA unit of SE. Incorporation of the 6 bp

underlined sequence of TJ108W0 and pSEA1 into *attS* was demonstrated in the previous

study.

S8 Fig. *attR* sequences formed in the SE-6283 insertion regions. Sequences are derived

from genomes listed in S7 Fig. Regarding SE-6283, the 6 bp underlined sequences in *attR*

originate from inserting SE circle, and they are not incorporated into *attS* upon SE

circularization.

S1 Table. MIC of transconjugant strains

S2 Table. Data availability

A

Vibrio alfacsensis strain 04Ya249 plasmid pSEA2
AP019851.1
113,882 - 139,270

attL(pSEA2)   *attR* (pSEA2)

SE-6945

E. coli strain TJ249
153,608 - 175,527

attL(insJ)

pSEA2

E. coli strain TJ249
444,587 - 470,661

attR(insJ)

pSEA2

intA CDS2 *intB* CDS4   *bla*

70 75 80 85 90 95 100
% identity in blastn

E. coli strain TJ249
4,214,970 - 4,238,688 (complementary)

attL(yjjNt)   *attR*(yjjNt)

Vibrio harveyi strain WXL538 chr 1
CP045070.1
2,941,000 - 2,967,316 (complementary)

attL(yaaAp)   *attR*(yaaAp)

SE-VhaWXL538

cl01143: H2O2_YaaD Superfamily

Vibrio alfacsensis strain 04Ya249 chr 1
AP019849.1
2,684,000 - 2,700,426 (complementary)

attL(yaaAp)   *attR*(yaaAp)

SE-6945

B

```
attB_pSEA2_AP019851.1        GCGCCTTTTCATATTGGCATATCGCCATGTxxxxxxTGTTTTCGATGTCACTATTGAGCTTACGAA
attB_Eco(insJ)_U00096.3      CTTTTTCTGGATCAACTGATATTGCAATGTTCTTTGGTGCCAATCTTGAGCGCGCGTAAACCAG
attB_Eco(yjjNt)_U00096.3     TGCAGCGTCCGGGGAAGTGTTGGGCGCTGTTTTTTTGTTTCTTAATAATGTGTTGTAAGCCGTAG
attB_Val_CAIM1831_CP032093.1 GATACAAAAGCCATGCAGACAACGCATGGCTTTTTTGTACTCTAA-TGAAAAATAGAATTAGGGA
attB_Vha_ATCC33843_CP009467.2 ATTCAAAAAGCCATGCAGACAATGCATGGCTTTTTTGTACTCTAAATGGAAAAATAGAATTAGGGA
```

C

A

B

attB_Vro_B64D1_Chr2_CP018312.1 TGCTGACCAGTTCGGTGTGTGGGGCGAGAAAAAGTTCATGGGTAAAGTTTACGATGGTCTTCACCG
attB_Vsc_FP3289_MDCJ01000002   GGCAGAACAGTTTGGCGTTTGGGGCGAGAAGAAATTTATGGGTAAAGTGTATGATGGCTTGCACCG
attB_Valg_ATCC17749_CP006718.1 TGCAGACCAGTTCGGCGTTTGGGGTGAAAGAAATTCATGGGCAAAGTATACGACGGTCTTCACCG
attB_Val_CAIM1831_CP032093.1   TGCAGAACAGTTTGGTGTTTGGGGTGAAAGAAGTTCATGGGTAAAGTGTACGACGGACTTCACCG
attB_Eco_W3110_AP009048.1   GTGCGAACAATTCGGCGTCTGGGGTGAAAGTCCTTCATGGGCAAAACCTACGATGGCATTCATCG
attB_Val_pSEA2_AP019851.1   CGTCTATGCTATTGATTCTCTTTATGAAGAAGAGGGCAAAGGCGTCAAATATGAGTTCGTATACAA

C