Atypical integrative element with strand-biased circularization activity assists interspecies

antimicrobial resistance gene transfer from *Vibrio alfacsensis*

Lisa Nonaka,[1,2,*] Michiaki Masuda,[1] Hirokazu Yano[3,4,†*]

[1]Department of Microbiology, Dokkyo Medical University School of Medicine, Mibu,

Tochigi, Japan

[2]Faculty of Human Life Sciences, Shokei University, Kumamoto, Japan

[3]Faculty of Life and Environmental Sciences, University of Tsukuba, Tsukuba, Japan

[4]Graduate School of Life Sciences, Tohoku University, Sendai, Japan

† Current address: Antimicrobial Resistance Research Center, National Institute of

Infectious Diseases, Higashimurayama, Japan

Short title: Integrative elements with strand-biased circularization activity

*Corresponding authors

Email: nonaka20@shokei-gakuen.ac.jp (LN); h-yano@niid.go.jp (HY)

Abstract

The exchange of antimicrobial resistance (AMR) genes between aquaculture and terrestrial microbial populations has emerged as a serious public health concern. However, the nature of the mobile genetic elements in marine bacteria is poorly documented. To gain insight into the genetic mechanisms underlying AMR gene transfer from marine bacteria, we mated a multidrug-resistant *Vibrio alfacsensis* strain with an *Escherichia coli* strain, and then determined the complete genome sequences of the donor and the transconjugant strains. Sequence analysis revealed a conjugative multidrug resistance plasmid in the donor strain, which was integrated into the chromosome of the recipient. The plasmid backbone in the transconjugant chromosome was flanked by two copies of a 7.1 kb unclassifiable integrative element harboring a β-lactamase gene. The 7.1 kb element and the previously reported element Tn*6283* share four coding sequences, two of which encode the catalytic R-H-R-Y motif of tyrosine recombinases. Polymerase chain reaction and sequencing experiments revealed that these elements generate a circular copy of one specific strand without leaving an empty site on the donor molecule, in contrast to the movement of integron gene cassettes or ICE/IMEs discovered to date. These elements are termed SEs (strand-biased circularizing integrative elements): SE-6945 (the 7.1 kb element) and SE-6283 (Tn*6283*). The copy number and location of SE-6945 in the chromosome affected the antibiotic resistance levels of the transconjugants. SEs were identified in the genomes of other *Vibrio* species. Overall, these results suggest that SEs are involved in the spread of AMR genes among marine bacteria.

Introduction

Antimicrobials have been used globally in aquaculture to control fish diseases. Although this

approach helps to maintain a stable supply of aquacultural products, misuse of

antimicrobials has led to the emergence of antimicrobial-resistant microbes and the

accumulation of resistance genes at aquaculture sites [1,2]. As the spread of multidrug-

resistant (MDR) bacteria and pan-drug-resistant bacteria threatens human life [3,4], it is

important to obtain clues about whether and how aquatic and terrestrial microbial

populations exchange genetic materials.

Conjugative plasmids [5] and integrative and conjugative elements (ICE) [6] are DNA

units that can move from one cell to another through the conjugation machinery that they

encode. They can directly move between species [7,8], and indirectly mobilize genes on

specific mobile elements [9–11]. Plasmids can be classified into families according to the

replicon type or the mobilization machinery type for epidemiological purposes [12,13].

Transposons have been defined as specific DNA segments that can repeatedly insert

into one or more sites in one or more genomes [14]. Several distinct strand exchange

mechanisms and the routes used for their location changes, i.e., *transposition*, have been

discovered to date [15]. In a comprehensive review by Curcio and Derbyshire [15],

transposons were classified into five families based on the DNA strand exchange enzymes

used. In prokaryotes, the mode of transposition was largely classified into 'cut-out paste-in'

(often called 'excision integration' for the ICEs [6]), 'copy-out paste-in,' or 'copy-in' for which

replication is necessary to complete transposition; e.g., Tn*3*, IS*91*. DNA insertion involving

DNA replication can also be seen in the integration of integron gene cassettes into *attI* [16]

and the phage CTX into *dif* [17]; both use tyrosine recombinase(s) as the strand exchange

enzyme. In transposon classification, ICEs are classified as transposons that use a Y-

transposase (tyrosine recombinase) or an S-transposase (serine recombinase) [15]. Other

than ICEs, DNA segments that can move to a location in other cells via the excision –

transfer (rolling-circle replication) – integration route have been termed IME (integrative

mobilizable element) [18], MGI (mobilizable genomic island) [19], CIME (*cis*-mobilizable

element) [20], or MTn (mobilizable transposon) [21] (hereafter collectively referred to as

IME). IMEs encode site-specific recombinase or DDE transposase as strand exchange

enzyme [22]. They have been recognized as degraded forms of ICEs [20]. ICEs/IMEs so far

studied in the laboratory seem to change their location via the 'excision integration' route

only [23–26]. Genes embedded in these mobile genetic elements are readily shared among

bacteria and provide genetic resources for microbial adaptation in changing environments.

To better understand the genetic mechanisms underlying the spread of antimicrobial

resistance (AMR) genes at aquaculture sites, we previously collected resistant bacteria from

sediments at an aquaculture site in Kagawa, Japan [27–29]. We identified a self-

transmissible MDR plasmid of the MOB_H-family, named pAQU1, that can replicate in both

the original host *Photobacterium damselae* subsp. *damselae* and in *Escherichia coli* [28].

Subsequently, another MOB_H-family MDR plasmid, named pSEA1, was identified in a *Vibrio*

alfacsensis isolate (previously identified as *V. ponticus*) [30]. pSEA1 carries a 12 kb

unclassifiable integrative element, Tn*6283*, in addition to the AMR genes. Although pSEA1

could not replicate in *E. coli* at 42 °C, following conjugation, it could cointegrate with the *E.*

coli chromosome by homologous recombination between two Tn*6283* copies: one on pSEA1

and another that moved from pSEA1 into the chromosome [30]. Further, Tn*6283* was

suggested to circularize one specific strand without producing an empty site, presumably

using tyrosine recombinases. However, it is not known whether interspecies gene transfer

assisted by Tn*6283*-like elements is common in nature.

To obtain further insights into the mechanisms behind genetic exchange among

aquaculture-associated bacteria, in this study we mated another MDR *Vibrio* isolate, *V.*

alfacsensis 04Ya249, with *E. coli* strains in the laboratory and then determined the genome

sequences of both the donor and transconjugant strains. We identified a Tn*6283*-like

element involved in the cointegration of the multidrug resistance plasmid and the *E. coli*

chromosome. Based on the findings in the previous study and this study, we have termed

Tn*6283* and a related element as a new class of mobile elements: s̲trand-biased circularizing

integrative e̲lement (SE). Tn*6283* is hereafter referred to as SE-6283.

In this study, several types of intracellular DNA movement need to be discussed. For

clarity, the terms *transposition*, *cointegration*, and *integration* are used here based on the

following rules. We call the movement of a DNA segment from one location to the other

location in one or two genomes, mediated by any DNA strand exchange enzymes,

transposition. The physical incorporation of a plasmid into a chromosome is called

cointegration regardless of the molecular mechanisms and genomic locations involved. The

physical incorporation of a DNA molecule into a specific site in another molecule mediated

by a site-specific recombinase (so-called integrase) is called integration.

Results

Identification of SE carrying a β-lactamase gene

We previously isolated *V. alfacsensis* strain 04Ya249 from sea sediment at an aquaculture

site [27]. 04Ya249 showed resistance to erythromycin, tetracycline, and ampicillin [31]. To

identify active mobile elements in this strain that are relevant to its AMR, the strain was

mated with macrolide-sensitive *E. coli* JW0452, and transconjugants were selected on

erythromycin at 42 °C. One transconjugant was named strain TJ249.

The complete genome sequences of strains 04Ya249 and TJ249 were determined

using the PacBio RS II platform. The genome of strain 04Ya249 consists of four replicons:

two chromosomes, a 306 kb putative conjugative plasmid pSEA2, and an 82 kb putative

conjugative plasmid pVA249. Strain 04Ya249 has a very similar genome architecture to the

database strain *V. alfacsensis* CAIM 1831 based on gene synteny (S1 Fig A) and average

nucleotide identity (S1 Fig B). Seven AMR genes were identified on pSEA2 (S2 Fig A). One β-

lactamase gene (60 % product identity to bla_{VHH-1} in the CARD database [32]) was located

within a 7.1-kb repeat region found in both chromosome 1 and pSEA2 (S2 Fig B). This β-

lactamase gene has been assigned a new family name, bla_{GMA} (Gammaproteobacterial

Mobile Class A β-lactamase), and allele name, bla_{GMA-1}, in NCBI.

The chromosome of *E. coli* TJ249 contains two notable insertions (Fig 1 A), a smaller

insertion of the 7.1-kb element alone immediately downstream of the putative transcription

termination site for *yjjN* (*yjjNt*) located between *yjjN* and *yjjO* (in GenBank accession no.

U00096.3), and a larger insertion into *insJ* of IS*3*, containing a nearly complete copy of pSEA2

and an additional copy of the 7.1 kb element. These indicate the movement between

different genomic locations (pSEA2, *yjjNt*, *insJ*), thereby transposition, of the 7.1 kb element

in *E. coli*. Thus, the 7.1 kb element was labelled a transposon and assigned a four digit

identifier, 6945, by the transposon registry [33] . As the 7.1-kb element was confirmed to

generate a circular copy of one specific strand without leaving an empty site like SE-6283 in

this study (see below), the 7.1 kb element was regarded as a new SE and designated SE-

6945.

As pSEA2 showed > 99.9 % nucleotide identity in the aligned region to pSEA1

(accession no. LC081338.1) from strain 04Ya108 [30], we also determined the complete

genome of strain 04Ya108 for comparison with strain 04Ya249. As predicted from Southern

hybridization [30], strain 04Ya108 possessed two copies of SE-6283, one in chromosome 1

and the other in plasmid pSEA1. Strain 04Ya108 also carries two copies of SE-6945, in the

same locations on the chromosome and plasmid as strain 04Ya249 (S1 Fig A, S2 Fig B). The

four coding sequences in SE-6945 showed homology to the four coding sequences clustered

at one end of SE-6283 (Fig 1 B). The biochemical functions of the products of the four genes

were predicted using Pfam and InterProScan in EMBL. The *intA* and *intB* products generated

a hit on phage_integrase (CL0382) in Pfam and were further aligned with known tyrosine

recombinases using structure-aware alignment program PROMAL3D [34] to confirm the

presence of conserved motifs. The *intA* and *intB* products contained the catalytic R-H-R-Y

motif (G-R motif in box I and H-R-H-Y motif in box II (Fig 1C)) conserved in tyrosine

recombinases [35], though the N-terminal region of IntB is extraordinarily long among

tyrosine recombinases. CDS2 and CDS4 products did not generate hits in the Pfam or

InterProScan databases. SE-6283 and SE-6945 carry imperfect 19 bp inverted repeat motifs

at their ends (Fig 1 B), termed C and C' based apparent on functional equivalence to the

core-type sites of phages [36]. Gene product identities between the SE-6945 and SE-6283

counterparts were 22.5 % for IntA, 20.5 % for CDS2, 23.2 % for IntB, and 26.6 % for CDS4 in

blastp.

Fig 1. Genetic features of SEs.

(A) Schematic representation of the 7.1 kb element and pSEA2 insertion locations in *E. coli*

TJ249. Yellow pentagons are the 7.1 kb element. The sequence from pSEA2 is indicated by a

horizontal line. (B) Upper panel: Genetic organization of SE-6283 and SE-6945. Lower panel:

Nucleotide sequences of imperfect inverted repeat motifs C and C'. Dots indicate that bases

in C are identical to those in C'. (C) Presence of R-H-R-Y motif in IntA and IntB. IntA and IntB

sequences are aligned with sequences of known tyrosine recombinases based on secondary

structure prediction using PROMAL 3D [34]. The aligned regions corresponding to box I and

box II in [35] are shown.

The border regions between SE-6945 and its target locations were termed *attL* (*intA*-

proximal side) and *attR* (*intA*-distal side) (Fig 2 A). The nucleotide sequence alignments of

attL and *attR* from the two *E. coli* chromosomal locations and the two locations in 04Ya249

genomes (pSEA2 and chr 1) are shown in S3 Fig and S4 Fig, respectively. Sequence

comparison of the *attL*, *attR*, and unoccupied target site in the host (*attB*) in *E. coli* (Fig 2 B)

revealed that the 6 bp next to the 3' end of C' in *attR* in *E. coli* (5'-TTTTCT-3') was identical to

the 6 bp next to the 5' end of C at *attL* on pSEA2, while the 6 bp next to the 5' end of C in

attL (5'-TTCTTT-3' in *insJ*, 5'-TTTTTT-3' in *yjjNt*) was from *attB* (S3 Fig, S4 Fig, Fig 2 B).

Therefore, SE-6945 inserts itself, ending with 5'-GTA-3' (Fig 1 B) along with an additional 6

bp next to the 5' end of C at *attL* on the donor molecule into the target site. The 6 bp from

the donor molecule is placed next to the 3' end of C' in the newly formed *attR.*

Target site preference of SE-6945 and SE-6283

After the isolation and genome sequence determination of TJ249, additional mating

experiments between *E. coli* JW0452 and *V. alfacsensis* 04Ya249 were performed to test

whether pSEA2 insertion into a specific location in the *E. coli* chromosome accompanying

SE-6945 transposition is reproducible. Independent mating was performed more than 17

times, and transconjugants were screened on tetracycline (Tc) alone, ampicillin (Ap) alone,

or both Ap and Tc. The SE-6945 integration pattern was investigated by Southern

hybridization for 16 randomly chosen Tc resistant transconjugants, and one Tc sensitive but

Ap-resistant transconjugant obtained from selection with Ap alone (S5 Fig C). Among 16 Tc

resistant transconjugants, 12 strains had *intA* at the *yjjNt*-SE-6945 junction or the *insJ*-SE-

6945 junction. These 16 strains also generated pSEA2-derived fragments (S5 Fig C). The

remaining four strains generated pSEA2-derived fragments and contained only one copy of

intA, suggesting SE-6945-independent pSEA2 cointegration with the chromosome;

autonomous replication of pSEA2 seems unlikely given the high frequency of pSEA2

cointegration with the chromosome in other transconjugants. Strain LN52, obtained from

selection with Ap alone, carried only one SE-6945 copy, and it was integrated into *yjjNt* (S5

Fig A, C). Taken together, these observations suggest that pSEA2 integration relies on SE-

6945 transposition, and that SE-6945 prefers *yjjNt* or *insJ* of IS*3*, rather than random sites in

the *E. coli* genome.

To obtain further insights into the target site of SE-6945, bacterial genomes carrying

SE-6945 were searched for based on blastn in the NCBI nucleotide collection (nr/nt)

database using the SE-6945 core gene region (*intA*-CDS2-*intB*-CDS4) as the query. The

genome of *Vibrio harveyi* strain WXL538 contains a segment showing 88.9% nucleotide

sequence identity to SE-6945 core genes. Genome structure comparisons between WXL538

and 04Ya249 revealed that WXL538 carries a 16.4 kb SE-6945-like SE (provisionally termed

SE-VhaWXL538) containing a different set of cargo genes (Fig 2 A). SE-VhaWXL538 is

integrated into the promoter region of a *yaaA* gene encoding an H_2O_2_YaaD_Superfamily

protein (NCBI conserved domain accession no. cl01143) (locus_tag F9277_14400 in

CP045070.1). This location corresponds to the SE-6945 insertion location in *V. alfacsensis*

chromosome 1 (Fig 2 A). Sequences of target sites (*attB*) in *E. coli*, *V. alfacsensis*, and *V.*

harveyi are shown in Fig 2 B. The conservation level of nucleotides in *attB* is presented in Fig

2 C as sequence logo information contents [37]. The central 6 bp putative crossover region

in *attB* ~~was~~ is T rich. However, other motifs, such as imperfect inverted repeat structures

observed in *attC/attI* of integrons (IntI target) [38,39], or *dif* of CTX phage integration site

(XerC/XerD target) [17], could not be detected in *attB, attL* (S3 Fig), or *attR* (S4 Fig) for SE-

6945 in the current data set.

Fig 2. Locations of SE-6945 and its related elements.

(A) Comparisons of genetic organization around SE insertion locations. GenBank accession

numbers (*Vibrio* strains only) and the coordinates of genomic segments compared are

shown below the strain name. (B) Alignment of *attB* sequences. The sequences are

retrieved from SE-free genomes. Since the SE-6945-free pSEA2 ancestor could not be found

in the database, the bases in the putative crossover region could not be defined and are

shown as x. Nicking is expected to take place next to the central box: position is shown by

black arrowhead. (C) Sequence logo representation of conservation levels of *attB* sequence.

We previously reported SE-6283 integration into the *bcp* gene encoding

peroxiredoxin (locus_tag b2480) in *E. coli.* SE-6283 also carries a *bcp* homolog (locus_tag

VYA_19800 in AP024165.1) as if it complements the disrupted *bcp* gene in the host genome

upon insertion [30]. Genome sequencing of strain 04Ya108 in this study revealed that one

SE-6283 copy was integrated into a *bcp* gene homolog in chromosome 1 (Fig 3 A). To obtain

further insight into the target site preference of SE-6283, SE-6283-like elements in public

databases were screened for using blastn. SE-6283-like SEs were identified in three other

Vibrio species (Fig 3 A). The sizes of SE-6283 and related SEs ranged from 13.8 kb (SE-6283)

to 41.3 kb (SE-ValK09K1 in Fig 3 A). Although the motifs C and C' of the three SE-6283-like

elements were slightly different from those of SE-6283 (see *attL* in S6 Fig and *attR* in S7 Fig),

all are integrated into *bcp* orthologs in chromosome 1: chromosome 1 (or I) of *V.*

alginolyticus strain K09K1 is incorrectly named as chromosome II in CP017919. Comparison

of *attL*, *attR*, and unoccupied *attB* found in the SEs-free genome of each *Vibrio* species

suggests the integration of SE-6283-like elements and the 6 bp sequence into the central 6

bp putative crossover region in *attB* (Fig 3 B). Nucleotide sequences around putative nicking

sites in *attB* for SE-6283 are G and A-rich and, like *attB* of SE-6945 did not possess an

inverted repeat structure (Fig 3 C). These observations collectively suggest that, like ICEs,

SEs target a few selected locations.

Fig 3. Location of SE-6283 and its related SEs.

(A) Comparisons of genetic organization around SE elements in five genomic locations.

GenBank accession numbers and the coordinates of genomic segments compared are

shown below the strain name. Except SE-6283 on pSEA1, all elements are integrated into a

bcp gene and carry a *bcp* gene. (B) Alignment of *attB* sequences from different bacterial

species. (C) Sequence logo showing conservation levels of *attB*. The arrowhead indicates the

putative nicking site on the top strand.

Transposition of SE-6945 aids pSEA2-chromosome cointegration

The direct repeat structure of SE-6945 in the TJ249 chromosome suggested that

pSEA2 was integrated into the recipient chromosome through the transposition of SE-6945,

with subsequent cointegration of pSEA2 with the chromosome by homologous

recombination between SE-6945 copies: one on the chromosome and the other on pSEA2

(referred to as a two-step gene transfer mechanism in the previous study [30]). If this

scenario holds, integration of SE-6945 alone should occur more frequently than pSEA2-

chromosome cointegration upon mating, and the cointegration would be reduced in

frequency or abolished altogether in a *recA*-null mutant recipient. To test these possibilities,

we conducted additional mating assays.

Quantitative mating assays were performed using a rifampicin-resistant recipient

strain, JW0452rif, and its isogenic strain carrying *recA* deletion JW0452Δ*recA*rif. Note that Tc

resistance is on pSEA2, while Ap resistance is on SE-6945. The transfer of Tc Ap resistance

from *V. alfacsensis* 04Ya249 to *E. coli* JW0452rif was detected only in 2 out of 4 replicated

mating experiments, and the transfer frequency was determined to be approximately 10^{-9}

per donor, indicating a very rare event, generating 0–3 transconjugant colonies per mating

(Fig 4). The transfer of Ap resistance alone was observed at >300-fold higher frequency than

the transfer of both Tc and Ap resistance (compare the first and the second groups from left

in Fig 4), indicating that integration of SE-6945 alone occurs more frequently than pSEA2-

chromosome cointegration during mating.

The transfer of Tc Ap resistance to JW0452Δ*recA*rif could not be detected in any of

the four independent mating assays (below detection limit of 1.02×10^{-9} per donor).

Unexpectedly the mean transfer frequency of Ap resistance to JW0452Δ*recA*rif was about

17-fold lower than that to JW0452rif (compare the second and the fourth groups from the

left in Fig 4). Thus, the direct effect of *recA* knockout on the cointegration could not be

evaluated due to the reduced detection sensitivity of gene transfer. However, the mean

transfer frequency of Tc Ap resistance to LN52rif (which already carries one copy of SE-6945

and wild type *recA*) was about 56-fold higher than that to JW0452rif (compare the first and

fifth groups in Fig 4). This indicates that the presence of the 7.1 kb long identical sequence

in both target DNA and pSEA2 promotes cointegration of pSEA2 and the recipient

chromosome.

Tn*3*-like cointegrate formation [15] might happen if SE-6945 could simultaneously

introduce nicks at one site on the top strand at *attL* (or *attR*) and one site on the bottom

strand at *attR* (or *attL*), and DNA-protein crosslink (DPC) repair [40] could clean the linked

tyrosine recombinase to produce a 3'-OH priming site. If SE-mediated cointegrate formation

occurs, then Tc Ap-resistant transconjugants should emerge at a comparable frequency to

Ap-resistant transconjugant, which was not the case in our observation.

Collectively, these results and the nature of tyrosine recombinases support the

model that pSEA2-chromosome cointegration mainly occurs through a two-step gene

transfer mechanism, transposition followed by homologous recombination, rather than

through a one-step Tn*3*-like cointegrate formation process.

Fig 4. Resistance gene transfer from *V. alfacsensis*.

Transfer frequency (Y-axis) is the $\log_{10}$-transformed value of the transconjugant colony-

forming units (CFU) divided by the donor CFU. Four replicate mating experiments were

performed. No transconjugants were detected in any of the four replicate experiments using

JW0452Δ*recA*rif with Tc Ap selection or in the two replicate experiments using JW0452rif

with Tc Ap selection (indicated by open circles placed at the bottom). In one experiment on

LN52rif, the donor CFU was not evaluated, but the transconjugant CFU was obtained at a

comparable frequency to that observed in the other three replicate experiments. The

detection limit of this experiment was around 1.02×10^{-9} per donor.

Strand-biased circularization of SE-6945

We previously [30] revealed the following unique features of SE-6283: (i) it does not

generate an unoccupied donor site upon circularization *in vivo*, and (ii) the circular form of

SE-6283 is generated using only one strand as a template at least in *E. coli*. We note that

this assertion is based on the notion that tyrosine recombinases generally do not produce

blunt ends at *att* sites on double stranded DNA during strand exchange. Although SE-6283

encodes tyrosine recombinases and not a single DDE transposase, the outcomes described

above, which result only from a figure-eight structure containing a single stranded DNA

bridge (Path 1 in Fig 5 A), is reminiscent of the first step of the 'copy-out paste-in'

transposition of some ISs [41] (IS*3* route), rather than the excision of ICEs (ICE/IME route) or

the generation of circular integron gene cassettes by *attC* × *attC* recombination (gene

cassette route) [25,42,43]. If the figure-eight structure containing a single stranded DNA

bridge is subject to replication without DPC repair, the products would be a single stranded

SE circle, the original copy of donor molecule, and the nicked chromosome, which is

eventually lost (Fig 5 A bottom left). To test whether circular copy production of SE-6945

also follows Path 1, we created a pSEA2-free 04Ya249 derivative strain LN95 carrying only a

single copy of SE-6945 in the chromosome (S5 Fig B) and analyzed the production of its

circular molecule in this strain and the sequence of strand exchange products.

Pairs of primers were designed such that one primer anneals inside the integrative

element and the other outside it (Fig 5 A), amplifying *attL* (*intA* side, Fig 5 B, product size

304 bp), *attR* (*bla* side, 248 bp), empty donor site (*attB,* 345 bp), occupied *attB* (7.4 kb), and

attS (joint of motif C and C' formed on the circular SE-6945, 206 bp). Primer set '1-4' in Fig 5

A was used to detect both empty *attB* and occupied *attB*.

The circular form of SE-6945 was detected in LN95 (Fig 5 B(i)) as well as the parent

strain 04Ya249 (S8 Fig A). This suggests that the chromosomal copy of SE-6945 in *V.*

alfacsensis is functional. The empty *attB* (345 bp) could not be detected under PCR

conditions with short extension times (Fig 5 B(i)), while a larger product of occupied *attB*

(7.4 kb) was detected when long PCR conditions were used (Fig 5 B(ii)).

Fig 5. PCR and sequencing analyses of SE-6945 circle.

(A)   PCR experiment to detect SE-6945 circle. Upper panel: primer annealing locations

(triangles in gray) on SE-6945 and the *attB* region. The number corresponds to the primer

set numbers in B. Primer number indicates the following primers in Table 2: 1,

LN_181_attB_L; 2, LN_142_junction2; 3, LN_143_junction2; 4, LN183_attB_R2. Lower panel:

possible routes of SE movement. Two strands of DNA are shown as separate lines, with SE in

orange. In Path 1 (solid arrows), one strand is nicked, and the SE side is circularized while

the host DNA side of the strand remains not joined, leading to a figure-eight structure

containing a single stranded DNA bridge. The figure-eight form can be subject to replication,

giving rise to a double stranded SE circle (if the recombinase bound 3' end is cleaned, for

example, by DPC repair [40], referred to as 'IS*3* (copy-out) route'), or single stranded DNA

circle (if the 3' end is not cleaned). In Path 2, strand exchange generates a figure-eight

containing a double stranded DNA bridge. After going through Path 2, there are two

possible outcomes: (i) another round of strand exchange at bottom strand leading to a

circular double stranded SE as well as a circular donor molecule carrying empty donor site

('ICE/IME route'), (ii) replication on the figure-eight leading to single stranded SE circle and

donor molecule with empty *attB*, and the original donor molecule ('gene cassette route').

(B) The result of PCR. PCR products were electrophoresed in 2% agarose gel (i) or 0.8%

agarose gel (ii). (C) Putative nicking sites in *attL* and *attR*. The SE DNA is shown in orange.

Nicking and ligation at positions shown as arrowheads in the same color can generate *attS*

with a 6 bp spacer. (D) Observed *attS* sequences. *attS* type 1 can be generated from strand

exchange at the top strand on chromosome 1 or bottom strand on pSEA2, while *attS* type 1

can be generated from strand exchange at the top strand on pSEA2 or bottom strand on

chromosome 1.

To estimate the copy number of the circular form of SE-6945 in the cell population,

we first searched for Illumina reads of strain LN95 that spanned *attS* or the "hypothetical"

attB. Although next-generation sequencing reads were obtained at 229× chromosome

coverage, no reads spanning *attS* or *attB* were detected [44]. Quantitative PCR further

revealed that the mean *attS* to *gyrB* ratio was 0.0012 (S8 Fig B). This relative copy number

was consistent with the result from a previous study detecting the circular form of SE-6283

in *E. coli*, which showed an *attS* to chromosome ratio of 0.001 [30].

To determine the nicking site on *attL* and *attR* (Fig 5 C), we cloned the PCR products

of *attS* and then sequenced ten cloned molecules. Based on the *attL*, *attR*, and the original

attB sequences in the TJ249 chromosome (S3 Fig, S4 Fig), we hypothesized four nicking sites

(arrowheads in Fig 5 C) in *attL* and *attR* on pSEA2 and chromosome 1 in strain 04Ya249. The

cloned *attS* PCR products of 04Ya249 contained a 6 bp spacer sequence between C and C'

(Fig 5 D). This observation supports the hypothesis that nicking occurs 6 bp upstream at the

C end and the C' end and/or 6 bp upstream at the C' end and the C end. Two sequences, 5'-

TTTTTT-3' (*attS* type 1) and 5'-TTTTCT-3' (*attS* type 2) were detected at a 1:9 ratio in the

spacer region between C and C' in strain 04Ya249 (examples of the Sanger sequencing trace

files have been posted in figshare [44]). These two types of spacer sequences can arise from

both pSEA2 and chromosome 1 (Fig 5 D). However, only one sequence (*attS* type 1) was

detected in the pSEA2-free strain LN95 (Fig 5 D), indicating that strand exchange occurs only

on the top strand in *attL* × *attR* recombination on chromosome 1 (Fig 5 C). The occurrence

of *attS* type 1 and *attS* type 2 in 04Ya249 (Fig 5 D) agrees with the top strand exchange in

attL × *attR* recombinations on chromosome 1 and pSEA2, respectively (Fig 5 C). Collectively,

these observations are consistent with the model that SE-6945 follows Path 1 (Fig 5 A)

without undergoing strand exchange of the bottom strand in *V. alfacsensis*.

Nicking of one strand of the double stranded DNA and its circularization is often seen in

the reactions mediated by HUH endonucleases (relaxase/rolling-circle replication initiator)

[45]. To confirm that observed nicking at *attL* and *attR* is independent of HUH

endonucleases produced from plasmids or other mobile elements in *V. alfaccensis*, the

production of *attS* was additionally investigated using *E. coli* strain LN52 that carries a single

copy of SE-6945 but lacks pSEA2. *attS* was produced even in LN52 (S8 Fig C). Therefore,

strand-biased (top strand only) *attL* × *attR* recombination is likely mediated by DNA strand

exchange enzymes encoded by SE-6945.

The insertion copy number and location of SE-6945 affects Ap resistance levels of the host cell

The naturally occurring strains 04Ya249 and 04Ya108 carry two copies of SE-6945 in their genomes. However, it remains unclear whether increased SE copies in a genome confer an advantage to the host cell. Three copies of SE-6945 in the transconjugant TJ249 were initially detected by genome sequencing. Southern hybridization analysis of other JW0452 transconjugants revealed one to two copies of SE-6945 in their genomes (S5 Fig). The location of SE-6945 became evident based on Southern hybridization for 15 among 19 transconjugants. Therefore, we used these transconjugant strains to test whether multiple integrations of SE-6945 or its integration location can increase the Ap resistance phenotype of the transconjugants.

Sixteen *E. coli* strains with or without SE-6945 were grown in Luria-Bertani broth with Ap, and then the minimum inhibitory concentrations (MICs) against individual strains were determined using the broth dilution method following the Clinical Laboratory Standards Institute (CLSI) guidelines. The MIC was highest in the clone carrying three copies of SE-6945, followed by the group of transconjugants carrying two copies and the group carrying one copy, and the MIC was the lowest in JW0452 (Fig 6). Although within-group variation was also observed, the increased copy number of SE-6945 in the genome generally resulted in the increased Ap resistance of the host cell.

The MICs of strains are shown in Table S1. There was a trend that strains with SE-6945 integrated into *insJ* showed a higher MIC than strains with integration into *yjjNt*. Thus, the location of SEs can also affect phenotype variation, possibly due to the differences in the expression levels of their cargo genes.

Fig 6. The copy number of SE-6945 affects the Ap resistance level. Sixteen transconjugants and a control strain (JW0452) were divided into four groups shown in distinct colors based on the SE-6945 copy number in the genome. MIC of ampicillin is shown according to copy number.

Discussion

Vibrio is one of the major bacterial genera found in marine sediments [46], and is among the most common microbiota of wild and farmed shrimp [47] and fish [48,49]. Several members of the genus *Vibrio* are pathogens of fishes reared in aquaculture [50], while other subsets of *Vibrio* species, including *V. cholerae*, *V. parahaemolyticus,* and *V. vulnificus*, which are ubiquitous in relatively low-salinity sea water, are human pathogens from seafood and/or water. [50,51]. Thus, *Vibrio* can be considered as a key genus aiding movement of AMR genes between aquatic environments and human-associated environments. Indeed, the accumulation of AMR genes in this genus has attracted increased research attention, particularly in *V. cholerae* [52,53]. However, direct experimental evidence for AMR gene transfer from *Vibrio* species to other human-relevant bacteria is limited.

Known genetic mechanisms of AMR gene transfer from *Vibrio* include A/C plasmids [54,55], unclassified conjugative plasmids [56], pAQU1-type MOB$_H$-family conjugative plasmids [57], pSEA1-type MOB$_H$-family conjugative plasmids [30], ICEs [58], IMEs [19], and a combination of a sedentary chromosomal integron and a conjugative plasmid carrying an integron [59]. The host range of pSEA1-type MOB$_H$ plasmids, discovered in *V. alfacsensis* (for which only one complete genome was available until the present study), is unknown. In this

study, we investigated how a pSEA1-like plasmid can contribute to AMR gene transfer in a

laboratory setting.

In contrast to A/C plasmids, autonomous replication of pSEA2 and pSEA1 was

difficult to achieve in *E. coli,* since transconjugant selection generally yielded *E. coli* clones

carrying plasmid DNA integrated into the chromosome. The current most likely mechanism

behind this phenomenon is transposition of SE followed by homologous recombination,

which enabled AMR gene transfer beyond the plasmid's replication host range. This mode of

horizontal gene transfer may be important among marine bacteria, since the plasmid

conjugation host range is expected to be wider than the replication host range [60,61].

pSEA1 was found to carry two SEs (SE-6283 and SE-6945). However, in the previous

study [30], only transposition of SE-6283 was observed, likely because we did not recognize

SE-6945 on pSEA1 and did not intend to detect its transposition. The newly identified SE-

6945 is the smallest SE identified to date (Fig 2 and Fig 3), and the only known SE harboring

an AMR gene. As SEs can have more than one target site in a single genome, they may

contribute to microbial adaptation to the antimicrobials used at aquaculture sites, with

diverse mechanisms beyond mediating the horizontal transmission of AMR genes, such as

increasing the resistance level or gene redundancy preceding evolutionary innovation of an

AMR gene [62].

The discovery of SE-6945 highlights four potential core genes (*intA*, CDS2, *intB*, and

CDS4) present in SEs. However, the specific roles of these gene products in their

transposition steps remain to be determined. Among the types of transposon movement

classified by Curcio and Derbyshire [15], 'copy-out' is currently the most consistent with our

observations of SEs regarding the first step of its location change. One model for strand

exchange on single stranded DNA mediated by tyrosine recombinase, leading to a single

stranded circle, has been proposed for integron gene cassette excision (replicative

resolution model in [43]). In the replicative resolution model, strand exchange between two

folded *attC* bottom strands and the following replication generates a gene cassette circle

with *attC* and results in the removal of one gene cassette with *attC* from the gene cassette

array on the donor molecule. This can cause formation of integron variants with fewer gene

cassettes and even integrons with an empty *attI,* as observed in *in vivo* experiments [42]. SE

circle formation rarely leaves the empty *attB* (Fig 5 B (i) and [30]). Thus the rejoining of the

host DNA side at nicking sites is somehow prevented, possibly by the action of multiple

proteins involved: possibly the CDS2, CDS4 products, unlike the movement of gene

cassettes. Currently, we do not have information on the fate of the nicked donor molecule.

Unlike DDE transposases, tyrosine recombinases usually do not generate a free 3' OH end

and are covalently linked to the phosphate of substrate DNA (Fig 5 A) [63]. Therefore, a

process equivalent to DPC repair [40], which might require both host and SE-encoding

factors, is necessary to achieve the IS*3*-like copy-out process. Nonetheless, we speculate

that the alternate single stranded DNA circle generation route, accompanying chromosome

loss (bottom left in Fig 5 A), is a less likely route as it does not give any advantage to target

site-selective and putatively non-transferrable SEs left in dying cells.

While strand-biased *attL* × *attR* recombination was repeatedly observed for the two

SEs, not much information has been obtained regarding their integration ~~step~~. While 6 bp

from the 5' end of C at *attL* on the donor molecule is incorporated into a circular SE copy,

the incorporated 6 bp tends to be placed next to the 3' end of C' at *attR* after integration

into the target site (see *attL* on pSEA2 and *attRs* on TJ249, and previously detected SE-6283

in transposition from pSEA1 to *bcp* in *E. coli*). Thus, there seems to be a strict rule in the

order of strand exchange at the integration step, but determining the precise order of

nicking and nicking sites on the two strands during the integration step requires further

investigations.

A notable difference between IS and SE is the strong strand bias unique to SE upon

strand exchange: some ISs can generate a single stranded bridge using either of the two

strands (but not both simultaneously) [64,65], while SEs use only top strand. The

dissimilarity between SE and ICE/IMEs is that ICE/IMEs encode the recombination

directionality factor Xis [25], while SEs do not encode Xis homologs. CDS2 or CDS4 products

may function exclusively in either the initial circularization step or the integration as the

recombination directionality factor. Another dissimilarity between SE and ICE/IME is that

the former does not encode an HUH endonuclease. There is still a possibility that SEs carry

an *oriT* recognized by a family of relaxases and are mobilized like CIME [21], though we

could not identify an *oriT* in SE-6945 or SE-6283 by local similarity search alone. It may also

be possible that IntB, a large tyrosine recombinase, functions as a relaxase, since a

conjugative plasmid from *Clostridium* has been shown to use a tyrosine recombinase

homolog as a conjugative relaxase [66].

We propose that interplay between SEs and conjugative MDR plasmids contribute to

AMR gene transmission in marine bacteria. Further biochemical and bioinformatic studies

on SEs are needed to reveal the mechanisms of gene transfer among the genus *Vibrio* and

other aquaculture-associated bacteria.

To refer to transposable individual SE members, we used numbers that are

compatible with the existing transposon database [33]. However, further debate is needed

regarding handling SEs in mobile DNA databases.

Materials and methods

Strains, plasmids, and culture media

Strains and plasmids used in the study are listed in Table 1. Strain 04Ya108 was previously

identified as *Vibrio ponticus* based on 16S rRNA gene sequence similarity. However,

determination of the complete sequence in this study revealed that it shows >96% average

nucleotide identity to *V. alfacsensis* strain CAIM 1831 (DSM 24595) (S1 Fig B) [67].

Therefore, strain 04Ya108 was re-classified as *V. alfacsensis*. Strain LN95 is a pSEA2-free

tetracycline-susceptible derivative of strain 04Ya249. This strain was generated through a

repeated batch culture of 04Ya249, and subsequent single-colony isolation. The absence of

pSEA2 from LN95 was confirmed by next-generation sequencing.

Table 1. Strains and plasmids used.

Strains or plasmids	Relevant genotype, phenotype, descriptions[a]	Reference
Vibrio alfacsensis		
04Ya249	Aquaculture isolate harboring conjugative plasmid pSEA2; Tc^r, Ap^r, Ery^r Cm^r	[27]
04Ya108	Aquaculture isolate harboring conjugative plasmid pSEA1; Tc^r, Ap^r, Ery^r Cm^r	[27]
LN95	pSEA2 free derivative of strain 04Ya249; Ap^r	This study
E. coli		
DH5α	F$^-$, Φ80d*lacZ*ΔM15, Δ(*lacZYA-argF*)U169, *deoR*, *recA*1, *endA*1, *hsdR*17(r$_K^-$, m$_K^+$), *phoA*, *supE*44, λ$^-$, *thi*-1, *gyrA*96, *relA*1	Takara Bio Inc., Shiga, Japan
JW0452	λ- *rph-1* Δ(*rhaD-rhaB*)568 *hsdR514* Δ*lacZ4787*(::*rrnB-3*) Δ(*araD-araB*)567 Δ*acrA748*::*kan*; Km^r	[68]
JW0452rif	Spontaneous rifampicin-resistant mutant of JW0452	This study
JW0452Δ *recA*	JW0452Δ*recA::cat*; Km^r, Cm^r	This study
JW0452Δ *recA*rif	Spontaneous rif resistant mutant of JW0452Δ*recA*; Km^r, Cm^r, Rif^r	This study

Transconjugants[c]		
TJ249	JW0452 *insJ*::SE-6945::pSEA2 *yjjNt*::SE-6945; Ap^r Tc^r Ery^r Cm^r	This study
LN28, LN32, LN36, LN86	JW0452 chr::pSEA2[b]; Ap^r Tc^r Ery^r Cm^r	This study
LN29, LN30, LN82, LN88, LN91	JW0452 *yjjNt*::SE-6945::pSEA2; Ap^r Tc^r Ery^r Cm^r	This study
LN33, LN34, LN35, LN37, LN58, LN84, LN89, LN90	JW0452 *insJ*::SE-6945::pSEA2; Ap^r Tc^r Ery^r Cm^r	This study
LN52	JW0452 *yjjNt*::SE-6945; Ap^r	This study
LN52rif	Spontaneous rifampicin-resistant mutant of LN52; Rif^r, Ap^r	This study
Plasmids		
pGEM-T	Cloning vector; pUC replicon, Ap^r	Promega, Madison, WI, USA
pGEM-attS	pGEM-T carrying the 206 bp *attS* fragment; pUC replicon, Ap^r	This study
pGEM-int	pGEM-T carrying the 103 bp *intA* fragment; pUC replicon, Ap^r	This study
pGEM-gyrB	pGEM-T carrying the 110 bp *gyrB* fragment from strain 04Ya249; pUC replicon, Ap^r	This study
pKD46	Lambda-Red recombinase expression plasmid; pSC101ts replicon, Ap^r	[69]
pKD3	Donor of the *cat* gene for the Lambda-Red recombination experiment; R6K replicon, Cm^r	[69]

a. Antimicrobial resistance phenotypes (^{r}) listed were confirmed by experiments.

Abbreviations are as follows: Ap, ampicillin; Tc, tetracycline: Ery, erythromycin; Cm,

chloramphenicol; Rif, rifampicin; *kan*, kanamycin resistance gene. *yjjNt* denotes the putative

transcription termination region for the *yjjN* gene. It is unclear whether the resistance genes

on pSEA2 are still in the original locations for other transconjugants besides TJ249.

b. Cointegration of plasmid pSEA2 with chromosome, independent of SE-6945 transposition.

Location of pSEA2 in the transconjugant genome could not be defined based on Southern

hybridization results.

c. The 19 transconjugants listed (TJ249, LN28, LN29, LN30, LN32, LN33, LN34, LN35, LN36,

LN37, LN52, LN58, LN82, LN84, LN86, LN88, LN89, LN90, LN91) were obtained from 19

independent mating experiments. Selection was done at 42 °C. Antibiotic selection used was

as follows: Tc for LN28, LN29 LN30, LN32, LN33, LN34, LN35, LN36, and LN37; Ap for LN52;

Ap Tc for LN58 LN82, LN84, LN86, LN88, LN89, LN90, and LN91; Ery for TJ249.

The presence of a single SE-6945 copy in the LN52 genome was confirmed by

Southern hybridization (S5 Fig). The *recA*-null mutant of JW0452, JW0452Δ*recA*, was

constructed using the lambda-Red method [69]. A DNA fragment containing the 5' and 3'

sequences of the *recA* gene and a chloramphenicol resistance gene was amplified by PCR

using primers YO-175 and RecA_stop_primingsite_2 (Table 2) and plasmid pKD3 [69] as a

template. The PCR products (700 ng) were introduced into electrocompetent cells of

JW0452 carrying pKD46 by electroporation using a Gene Pulser Xcell™ (BioRad, Hercules,

CA, USA). The occurrence of recombination at the expected site was confirmed by PCR using

primers CAT-584 and BW25113_2815723f (Table 2). The absence of the *recA* gene was also

confirmed by PCR using primers LN192_recA1 and LN193_recA2.

Vibrio alfacsensis strains were cultured in BD Bacto™ brain heart infusion medium

(BD237500; Becton, Dickinson, and Company, Franklin Lakes, NJ, USA) supplemented with

up to 2% NaCl at 25 °C. *E. coli* strains were cultured in BD Difco™ LB Broth, Miller

(BD244520; Becton, Dickinson, and Company) at 37 °C or 42 °C. BD Difco™ Mueller Hinton

Broth (BD 275730; Becton, Dickinson, and Company) was used for antibiotic susceptibility

testing of *E. coli*. BD Difco™ Marine Broth 2216 (BD279110; Becton, Dickinson, and

Company) was used for filter mating. Solid media were prepared by adding 1.5% agar to the

broth. Antibiotics were added to the medium at the following concentrations when

required: erythromycin (Nacalai Tesque, Kyoto, Japan), 100 μg/ml; tetracycline (Nacalai

Tesque), 10 μg/ml; rifampicin (Sigma-Aldrich, St. Louis, MO, USA), 50 μg/ml; ampicillin

(Nacalai Tesque), 100 μg/ml.

Table 2. Oligonucleotides used.

Name	Sequence (5' to 3')	Purpose
LN112	GGGTTACCTTCCCAATGCGT	Southern hybridization probe for SE-6945 *intA*
LN113	CGACTGTTGGTAGCGACTGT	Southern hybridization probe for SE-6945 *intA*
LN_142_junction2	AAGATGGTAAAAGTGTTCCA	Detection of *attS* (206 bp) by qPCR and *attL* (304 bp) by standard PCR
LN_143_junction2	TTTGTGTGTAGCCCTTGTG	Detection of *attS* (206 bp) by qPCR and *attR* (248 bp) by standard PCR
LN_150_intA2	GGTTATGTGGAGAAGTTGCC	Detection of *intA* (103 bp) by qPCR
LN_151_intA2	TGAGTTCGGTTTCTTGCTTC	Detection of *intA* (103 bp) by qPCR
LN_181_attB_L	CGAGGGTAAAGTGCCAACAT	Detection of chromosomal *attB* (345 bp if empty, 7.4 kb if occupied) and *attL* (304 bp) by standard PCR (04Ya249)

LN183_attB_R2	ACATCAGCAGGAGTTAGTTG	Detection of chromosomal *attB* (345 bp, 7.4 kb) and *attR* (248 bp) by standard PCR (04Ya249)
LN184_gyrBf1	AACAGAATTGCACCCAGAAG	Detection of *gyrB* (110 bp) by qPCR (04Ya249)
LN185_gyrBr1	GAAGACCGCCTGATACTTTG	Detection of *gyrB* (110 bp) by qPCR (04Ya249)
YO-175	CAGAACATATTGACTATCCGGTATT ACCCGGCATGACAGGAGTAAAAAT GTGTAGGCTGGAGCTGCTTCG	Amplification of *cat* gene for Lambda-Red recombination
RecA_stop_primingsite_2	ATGCGACCCTTGTGTATCAAACAA GACGATTAAAAATCTTCGTTAGTTT CCATATGAATATCCTCCTTA	Amplification of *cat* gene for Lambda-Red recombination
CAT-584	AAGCCATCACAAACGGCATG	Confirmation of *cat* gene insertion
BW25113_2815723f	AATACGCGCAGGTCCATAAC	Confirmation of *cat* gene insertion
LN192_recA1	GTTCCATGGATGTGGAAACC	Confirmation of *recA* deletion
LN193_recA2	ATATCGACGCCCAGTTTACG	Confirmation of *recA* deletion

Conjugation

The donor *Vibrio* strain and *E. coli* recipient strains were grown overnight at 25 °C and 37 °C, respectively. A 500 µl aliquot of each culture was mixed, centrifuged, and resuspended in 50 µl of Luria-Bertani broth. The cell mixture was spotted on a 0.45 µm pore-size nitrocellulose filter (Merck, Millipore Ltd., Tullagreen, Ireland) placed on marine broth agar, and allowed to mate for 24 h at 25 °C. After incubation, the cell mixture on the filter was serially diluted in 1× phosphate-buffered saline, and then 100 µl of the mixture was plated on an appropriate agar medium to measure the CFU and incubated at 42 °C to suppress growth of *Vibrio* strains. After genome sequence determination of TJ249, more than 17 independent mating experiments were performed to obtain insights into the variation of pSEA2 integration patterns using JW0452 as the recipient, and transconjugants were screened at 42 °C in the presence of Tc alone, Ap alone, or both Ap and Tc. The AMR phenotype of each transconjugant was further determined after single-colony isolation using Ery, Cm, Ap, and Tc (Table 1).

Quantitative mating assays (Fig 4) were performed to address the involvement of homologous recombination in pSEA2 integration using strains JW0452rif, JW0452Δ*recA*rif, and LN95rif as recipients. Selection conditions for transconjugants were Rif Ap Tc 42 °C or Rif Ap 42 °C. The *Vibrio* donor strain was selected on brain heart infusion agar with 2% NaCl supplemented with Tc at 25 °C. CFUs of donor, recipient, and transconjugant were determined after 24 hours of incubation.

PCR experiments

The PCR detection of *attL*, *attR,* and *attS* was performed using TaKaRa Ex Taq (TaKaRa Bio

Inc., Kusatsu, Japan), and genomic DNA of 04Ya249, LN95, and LN52 prepared using QIAGEN

DNeasy Blood & Tissue kit (QIAGEN GmbH, Hilden, Germany). PCR premix was prepared in a

50 μl scale, and 1 μl of the template (10 ng/μl) was added to the reaction mix throughout

the study. The PCR cycle used was 94 °C for 1 min, followed by 35 cycles of 94 °C for 30 s

and 60 °C for 30 s. As the empty *attB* was not amplified using TaKaRa Ex Taq, we conducted

additional long PCR using LA Taq polymerase (TaKaRa Bio Inc.) to detect occupied *attB* and

confirm the functionality of primers used for detection of empty *attB*. The PCR cycles used

for long PCR were 1 cycle of 94 °C for 1 min, followed by 35 cycles of 95 °C for 1 min, 65 °C

for 1 min, and 72 °C for 1 min. PCR products of *attS* were further cloned using the pGEM-T

easy vector system (Promega, Madison, WI, USA) and DH5α competent cells to analyze the

sequence variation of PCR products.

Quantitative PCR was performed using THUNDERBIRD® SYBR qPCR Mix (Toyobo,

Osaka, Japan) and a CFX connect Real-Time system (BioRad, Hercules, CA, USA) and a two-

step PCR protocol of 40 cycles of 95 °C for 5 s and 60 °C for 30 s. Target quantity was

estimated based on a standard curve of the control plasmid DNA (pGEM-*gyrB*, pGEM-*intA*,

pGEM-*attS*) constructed by TA cloning. Primers used and their purposes are listed in Table 2

and Fig 5 legend.

Southern hybridization

The copy number and integration location of SE-6945 in transconjugants (JW0452

derivatives) were analyzed by Southern hybridization using the 5' end of *intA*, amplified

using primers LN112 and LN113 (Table 2) and the PCR DIG Synthesis Kit (Roche, Basel,

Switzerland) as a probe. Genomic DNA (2.5 μg) was double digested with either *Nde*I and

*Sph*I or *Nde*I and *Hin*dIII (New England Biolabs, Ipswich, MA, USA), then electrophoresed in

0.8% agarose gel. Separated DNA fragments were transferred to the nitrocellulose

membranes (Amersham Hybond-N+; GE Healthcare UK Ltd., Buckinghamshire, UK) by

capillary transfer. The probe was detected using CDT-star® (Roche). Restriction map for the

DNA segments around SE-6945 integration locations and the expected size of DNA segments

with probe signal are shown in S5 Fig A.

Antimicrobial susceptibility testing

611   To examine the antimicrobial susceptibility of transconjugants harboring one to three SE-

612   6945 copies, the MIC of ampicillin was determined using the broth dilution method in 96-

613   well microtiter plate format according to standard M07 of the CLSI [70]. The Ap

614   concentrations tested were 1000, 500, 250, 125, 62.5, 31.25, 16, 8, 4, and 2 μg/ml. The test

615   plates were incubated at 35 °C for 24 h.

Genome sequencing

Genomic DNA was extracted from 250–500 μl of *V. alfacsensis* strains 04Ya249, 04Ya108, or

E. coli strain TJ249 culture using the QIAGEN DNeasy Blood & Tissue kit (QIAGEN GmbH).

The extracted genomic DNA was sequenced on the PacBio RS II platform using P6C4

chemistry at Macrogen (Tokyo, Japan) in 2016. Genome assembly was conducted using

HGAP v.3 [71] for *Vibrio* strains and Flye v 2.8.3-b1695 for *E. coli* TJ249 [72]. Reads were

obtained at >120× coverage for the chromosomes of each strain. Reads and the genome

sequence of TJ249 have been posted to figshare [73]. Illumina reads of the pSEA2-free strain

LN95 were obtained using the TruSeq PCR-free library and NovaSeq 6000 platform at

NovogeneAIT Genomics Singapore Pte., Ltd. (Singapore) to confirm the loss of pSEA2 and to

identify the circular form of SE-6945. The mapping results have been posted to figshare [44].

Genomes were compared using MUMmer3.23 [74] and GenomeMatcher [75]. AMR genes in

the assembly were searched for using AMRFinderPlus [76]. The average nucleotide identity

was determined using fastANI [77].

Searches for SE-6945 and SE-6283-related SEs

Nucleotide sequences of the SE core gene region (*intA*, CDS2, *intB*, CDS4) (6432 bp for SE-

6283, 5541 bp for SE-6945) were used as queries in blastn searches against the NCBI

nucleotide collection (nr/nt) database (August 3, 2021). Hit subjects showing > 85 %

nucleotide identity with > 99% query coverage were regarded as close relatives of SE-6945

or SE-6283, and were further investigated as to their integration locations.

Data availability

Sequence data generated in this study and their availability are summarized in Table S2. The

complete genome sequences of strains 04Ya108 and 04Ya249 were submitted to

GenBank/DDBJ/EMBL under accession numbers AP024165 (chr1 of 04Ya108), AP024166

(chr2 of 04Ya108), AP024167 (pSEA1 of 04Ya108), AP024168 (pYa108 of 04Ya108),

AP019849 (chr1 of 04Ya249), AP019850 (chr2 of 04Ya249), AP019851 (pSEA2 of 04Ya249),

and AP019852 (pVA249 of 04Ya249). The raw reads for strains 04Ya108, 04Ya249, LN95 and

TJ249 are available from Sequence Read Archive under accession numbers DRA011098,

DRA008632, DRA011762, and DRA013462, respectively.

Acknowledgments

We thank the National Bioresource Project of the National Institute of Genetics, Japan, for

providing the *E. coli* strains BW25113, JW2669, and JW0452. We thank Atsushi Ota and

Fumito Maruyama at Hiroshima University for their support for the annotation on the TJ249

assembly. We thank Yuichi Otsuka at Saitama University for helpful discussions and Yuta

Sugimoto at Ehime University for experimental support. We thank Kenji K Kojima at Genetic

Information Research Institute for the discussion on transposon classification, and Yasukazu

Daigaku at Tohoku University for the discussion on DNA protein cross link repair. We thank

Editage for language editing (job number LINON_4_4).

References

1. Cabello FC, Godfrey HP, Tomova A, Ivanova L, Dölz H, Millanao A et al. (2013)

Antimicrobial use in aquaculture re-examined: its relevance to antimicrobial resistance

and to animal and human health. Environ Microbiol 15: 1917-1942.

2. Cabello FC, Godfrey HP, Buschmann AH, Dölz HJ (2016) Aquaculture as yet another

environmental gateway to the development and globalisation of antimicrobial

resistance. Lancet Infect Dis 16: e127-e133.

3. Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG et al. (2012)

Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an

international expert proposal for interim standard definitions for acquired resistance.

Clin Microbiol Infect 18: 268-281.

4. Karakonstantis S, Kritsotakis EI, Gikas A (2020) Pandrug-resistant Gram-negative

bacteria: a systematic review of current epidemiology, prognosis and treatment

options. J Antimicrob Chemother 75: 271-282.

5. Rodríguez-Beltrán J, DelaFuente J, León-Sampedro R, MacLean RC, San Millán Á (2021)

Beyond horizontal gene transfer: the role of plasmids in bacterial evolution. Nat Rev

Microbiol 19: 347-359.

6. Johnson CM, Grossman AD (2015) Integrative and Conjugative Elements (ICEs): What

They Do and How They Work. Annu Rev Genet 49: 577-601.

7. Yano H, Rogers LM, Knox MG, Heuer H, Smalla K, Brown CJ et al. (2013) Host range

diversification within the IncP-1 plasmid group. Microbiology (Reading) 159: 2303-

2315.

8. Dahmane N, Libante V, Charron-Bourgoin F, Guédon E, Guédon G, Leblond-Bourget N

et al. (2017) Diversity of integrative and conjugative elements of *Streptococcus*

salivarius and their intra- and interspecies Transfer. Appl Environ Microbiol 83:

e00337-17.

9. Cabezón E, Lanka E, de la Cruz F (1994) Requirements for mobilization of plasmids

RSF1010 and ColE1 by the IncW plasmid R388: *trwB* and RP4 *traG* are interchangeable.

J Bacteriol 176: 4455-4458.

10. Carraro N, Matteau D, Luo P, Rodrigue S, Burrus V (2014) The master activator of

IncA/C conjugative plasmids stimulates genomic islands and multidrug resistance

dissemination. PLoS Genet 10: e1004714.

11.   Daccord A, Ceccarelli D, Burrus V (2010) Integrating conjugative elements of the

692   SXT/R391 family trigger the excision and drive the mobilization of a new class of *Vibrio*

693   genomic islands. Mol Microbiol 78: 576-588.

694   12.   Garcillán-Barcia MP, Francia MV, de la Cruz F (2009) The diversity of conjugative

695   relaxases and its application in plasmid classification. FEMS Microbiol Rev 33: 657-687.

696   13.   Villa L, Carattoli A (2020) Plasmid typing and classification. Methods Mol Biol 2075:

697   309-321.

698   14.   Roberts AP, Chandler M, Courvalin P, Guédon G, Mullany P, Pembroke T et al. (2008)

699   Revised nomenclature for transposable genetic elements. Plasmid 60: 167-173.

700   15.   Curcio MJ, Derbyshire KM (2003) The outs and ins of transposition: from mu to

701   kangaroo. Nat Rev Mol Cell Biol 4: 865-877.

702   16.   Bouvier M, Demarre G, Mazel D (2005) Integron cassette insertion: a recombination

703   process involving a folded single strand substrate. EMBO J 24: 4356-4367.

704   17.   Val ME, Bouvier M, Campos J, Sherratt D, Cornet F, Mazel D et al. (2005) The single-

705   stranded genome of phage CTX is the form used for integration into the genome of

706   Vibrio cholerae. Mol Cell 19: 559-566.

707   18.   Doublet B, Boyd D, Mulvey MR, Cloeckaert A (2005) The *Salmonella* genomic island 1 is

708   an integrative mobilizable element. Mol Microbiol 55: 1911-1924.

709   19.   Carraro N, Rivard N, Ceccarelli D, Colwell RR, Burrus V (2016) IncA/C conjugative

710   plasmids mobilize a new family of multidrug resistance islands in clinical *Vibrio*

711   *cholerae* non-O1/non-O139 isolates from Haiti. mBio 7: e00509-16.

712   20.   Pavlovic G, Burrus V, Gintz B, Decaris B, Guédon G (2004) Evolution of genomic islands

713   by deletion and tandem accretion by site-specific recombination: ICE*St1*-related

714   elements from *Streptococcus thermophilus*. Microbiology (Reading) 150: 759-774.

21.   Achard A, Leclercq R (2007) Characterization of a small mobilizable transposon,

MTn*Sag1*, in *Streptococcus agalactiae*. J Bacteriol 189: 4328-4331.

22.   Guédon G, Libante V, Coluzzi C, Payot S, Leblond-Bourget N (2017) The obscure world

of integrative and mobilizable elements, highly widespread elements that pirate

bacterial conjugative systems. Genes (Basel) 8: E337.

23.   Caparon MG, Scott JR (1989) Excision and insertion of the conjugative transposon

Tn*916* involves a novel recombination mechanism. Cell 59: 1027-1034.

24.   Auchtung JM, Lee CA, Monson RE, Lehman AP, Grossman AD (2005) Regulation of a

Bacillus subtilis mobile genetic element by intercellular signaling and the global DNA

damage response. Proc Natl Acad Sci U S A 102: 12554-12559.

25.   Burrus V, Waldor MK (2003) Control of SXT integration and excision. J Bacteriol 185:

5045-5054.

26.   Naito M, Ogura Y, Itoh T, Shoji M, Okamoto M, Hayashi T et al. (2016) The complete

genome sequencing of *Prevotella intermedia* strain OMA14 and a subsequent fine-

scale, intra-species genomic comparison reveal an unusual amplification of conjugative

and mobile transposons and identify a novel *Prevotella*-lineage-specific repeat. DNA

Res 23: 11-19.

27.   Nonaka L, Ikeno K, Suzuki S (2007) Distribution of tetracycline resistance gene, *tet(M)*,

in Gram-positive and Gram-negative bacteria isolated from sediment and seawater at

a coastal aquaculture site in Japan. Microbes Environ 22: 355-364.

28.   Nonaka L, Maruyama F, Miyamoto M, Miyakoshi M, Kurokawa K, Masuda M (2012)

Novel conjugative transferable multiple drug resistance plasmid pAQU1 from

Photobacterium damselae subsp. *damselae* isolated from marine aquaculture

environment. Microbes Environ 27: 263-272.

29.   Nonaka L, Maruyama F, Suzuki S, Masuda M (2015) Novel macrolide-resistance genes,

740   *mef*(C) and *mph*(G), carried by plasmids from *Vibrio* and *Photobacterium* isolated from

741   sediment and seawater of a coastal aquaculture site. Lett Appl Microbiol 61: 1-6.

742   30.   Nonaka L, Yamamoto T, Maruyama F, Hirose Y, Onishi Y, Kobayashi T et al. (2018)

743   Interplay of a non-conjugative integrative element and a conjugative plasmid in the

744   spread of antibiotic resistance via suicidal plasmid transfer from an aquaculture *Vibrio*

745   isolate. PLoS One 13: e0198613.

746   31.   Neela FA, Nonaka L, Suzuki S (2007) The diversity of multi-drug resistance profiles in

747   tetracycline-resistant *Vibrio* species isolated from coastal sediments and seawater. J

748   Microbiol 45: 64-68.

749   32.   Alcock BP, Raphenya AR, Lau TTY, Tsang KK, Bouchard M, Edalatmand A et al. (2020)

750   CARD 2020: antibiotic resistome surveillance with the comprehensive antibiotic

751   resistance database. Nucleic Acids Res 48: D517-D525.

752   33.   Tansirichaiya S, Rahman MA, Roberts AP (2019) The transposon registry. Mob DNA 10:

753   40.

754   34.   Pei J, Kim BH, Grishin NV (2008) PROMALS3D: a tool for multiple protein sequence and

755   structure alignments. Nucleic Acids Res 36: 2295-2300.

756   35.   Nunes-Düby SE, Kwon HJ, Tirumalai RS, Ellenberger T, Landy A (1998) Similarities and

757   differences among 105 members of the Int family of site-specific recombinases.

758   Nucleic Acids Res 26: 391-406.

759   36.   Landy A (1989) Dynamic, structural, and regulatory aspects of lambda site-specific

760   recombination. Annu Rev Biochem 58: 913-949.

761   37.   Schneider TD, Stephens RM (1990) Sequence logos: a new way to display consensus

762   sequences. Nucleic Acids Res 18: 6097-6100.

38.   Stokes HW, O'Gorman DB, Recchia GD, Parsekhian M, Hall RM (1997) Structure and

function of 59-base element recombination sites associated with mobile gene

cassettes. Mol Microbiol 26: 731-745.

39.   Stokes HW, Hall RM (1989) A novel family of potentially mobile DNA elements

encoding site-specific gene-integration functions: integrons. Mol Microbiol 3: 1669-

1683.

40.   Zhang H, Xiong Y, Chen J (2020) DNA-protein cross-link repair: what do we know now.

Cell Biosci 10: 3.

41.   Chandler M, Fayet O, Rousseau P, Ton Hoang B, Duval-Valentin G (2015) Copy-out-

paste-in transposition of IS*911*: a major transposition pathway. Microbiol Spectr 3:

42.   Collis CM, Hall RM (1992) Site-specific deletion and rearrangement of integron insert

genes catalyzed by the integron DNA integrase. J Bacteriol 174: 1574-1585.

43.   Escudero JA, Loot C, Nivina A, Mazel D (2015) The integron: adaptation on demand.

Microbiol Spectr 3: MDNA3-0019.

44.   Yano H, Nonaka L (2021) Illumina reads analysis of *Vibrio alfacsensis* pSEA2-free strain

LN95. URL: https://doiorg/106084/m9figshare14338883

45.   Chandler M, de la Cruz F, Dyda F, Hickman AB, Moncalian G, Ton-Hoang B (2013)

Breaking and joining single-stranded DNA: the HUH endonuclease superfamily. Nat

Rev Microbiol 11: 525-538.

46.   Chen J, McIlroy SE, Archana A, Baker DM, Panagiotou G (2019) A pollution gradient

contributes to the taxonomic, functional, and resistome diversity of microbial

communities in marine sediments. Microbiome 7: 104.

47.   Cornejo-Granados F, Lopez-Zavala AA, Gallardo-Becerra L, Mendoza-Vargas A, Sánchez

F, Vichido R et al. (2017) Microbiome of Pacific Whiteleg shrimp reveals differential

bacterial community composition between Wild, Aquacultured and AHPND/EMS

outbreak conditions. Sci Rep 7: 11783.

48.   Llewellyn MS, McGinnity P, Dionne M, Letourneau J, Thonier F, Carvalho GR et al.

(2016) The biogeography of the atlantic salmon (*Salmo salar*) gut microbiome. ISME J

10: 1280-1284.

49.   Egerton S, Culloty S, Whooley J, Stanton C, Ross RP (2018) The gut microbiota of

marine fish. Front Microbiol 9: 873.

50.   Thompson FL, Iida T, Swings J (2004) Biodiversity of Vibrios. Microbiol Mol Biol Rev 68:

403-431.

51.   Takemura AF, Chien DM, Polz MF (2014) Associations and dynamics of *Vibrionaceae* in

the environment, from the genus to the population level. Front Microbiol 5: 38.

52.   Verma J, Bag S, Saha B, Kumar P, Ghosh TS, Dayal M et al. (2019) Genomic plasticity

associated with antimicrobial resistance in *Vibrio cholerae*. Proc Natl Acad Sci U S A

116: 6226-6231.

53.   Osei Sekyere J, Reta MA (2020) Genomic and resistance epidemiology of Gram-

negative bacteria in Africa: a systematic review and phylogenomic analyses from a one

health perspective. mSystems 5: e00897-20.

54.   Carraro N, Sauvé M, Matteau D, Lauzon G, Rodrigue S, Burrus V (2014) Development

of pVCR94ΔX from *Vibrio cholerae*, a prototype for studying multidrug resistant IncA/C

conjugative plasmids. Front Microbiol 5: 44.

55.   Folster JP, Katz L, McCullough A, Parsons MB, Knipe K, Sammons SA et al. (2014)

Multidrug-resistant IncA/C plasmid in *Vibrio cholerae* from Haiti. Emerg Infect Dis 20:

1951-1953.

56.   Liu M, Wong MH, Chen S (2013) Molecular characterisation of a multidrug resistance

conjugative plasmid from *Vibrio parahaemolyticus*. Int J Antimicrob Agents 42: 575-

579.

57.   Nonaka L, Maruyama F, Onishi Y, Kobayashi T, Ogura Y, Hayashi T et al. (2014) Various

pAQU plasmids possibly contribute to disseminate tetracycline resistance gene *tet*(M)

among marine bacterial community. Front Microbiol 5: 152.

58.   Beaber JW, Burrus V, Hochhut B, Waldor MK (2002) Comparison of SXT and R391, two

conjugative integrating elements: definition of a genetic backbone for the mobilization

of resistance determinants. Cell Mol Life Sci 59: 2065-2070.

59.   Rowe-Magnus DA, Guerout AM, Mazel D (2002) Bacterial resistance evolution by

recruitment of super-integron gene cassettes. Mol Microbiol 43: 1657-1669.

60.   Encinas D, Garcillán-Barcia MP, Santos-Merino M, Delaye L, Moya A, de la Cruz F

(2014) Plasmid conjugation from proteobacteria as evidence for the origin of

xenologous genes in cyanobacteria. J Bacteriol 196: 1551-1559.

61.   Kishida K, Inoue K, Ohtsubo Y, Nagata Y, Tsuda M (2017) Host range of the conjugative

transfer system of IncP-9 naphthalene-catabolic plasmid NAH7 and characterization of

its *oriT* region and relaxase. Appl Environ Microbiol 83: e02359-16.

62.   Elliott KT, Cuff LE, Neidle EL (2013) Copy number change: evolving views on gene

amplification. Future Microbiol 8: 887-899.

63.   Jayaram M, Ma CH, Kachroo AH, Rowley PA, Guga P, Fan HF et al. (2015) An overview

of tyrosine site-specific recombination: from an Flp perspective. Microbiol Spectr 3:

64.   Polard P, Chandler M (1995) An in vivo transposase-catalyzed single-stranded DNA

circularization reaction. Genes Dev 9: 2846-2858.

65.   Rousseau P, Loot C, Turlan C, Nolivos S, Chandler M (2008) Bias between the left and

834   right inverted repeats during IS911 targeted insertion. J Bacteriol 190: 6111-6118.

835   66.   Wisniewski JA, Traore DA, Bannam TL, Lyras D, Whisstock JC, Rood JI (2016) TcpM: a

836   novel relaxase that mediates transfer of large conjugative plasmids from *Clostridium*

837   *perfringens*. Mol Microbiol 99: 884-896.

838   67.   Gomez-Gil B, Roque A, Chimetto L, Moreira APB, Lang E, Thompson FL (2012) *Vibrio*

839   *alfacsensis* sp. nov., isolated from marine organisms. Int J Syst Evol Microbiol 62: 2955-

840   2961.

841   68.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M et al. (2006) Construction of

842   *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. Mol

843   Syst Biol 2: 2006.0008.

844   69.   Datsenko KA, Wanner BL (2000) One-step inactivation of chromosomal genes in

845   *Escherichia coli* K-12 using PCR products. Proc Natl Acad Sci U S A 97: 6640-6645.

846   70.   Wayne PA (2018) Methods for dilution antimicrobial susceptibility tests for bacteria

847   that grow aerobically. 11th ed. CLSI standard M07. Clinical and Laboratory Standards

848   Institute.

849   71.   Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C et al. (2013)

850   Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing

851   data. Nat Methods 10: 563-569.

852   72.   Kolmogorov M, Yuan J, Lin Y, Pevzner PA (2019) Assembly of long, error-prone reads

853   using repeat graphs. Nat Biotechnol 37: 540-546.

854   73.   Yano H, Nonaka L (2021) Complete genome of transconjugant TJ249. URL:

855   https://doiorg/106084/m9figshare13332467

74.   Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C et al. (2004)

Versatile and open software for comparing large genomes. Genome Biol 5: R12.

75.   Ohtsubo Y, Ikeda-Ohtsubo W, Nagata Y, Tsuda M (2008) GenomeMatcher: a graphical

user interface for DNA sequence comparison. BMC Bioinformatics 9: 376.

76.   Feldgarden M, Brover V, Haft DH, Prasad AB, Slotta DJ, Tolstoy I et al. (2019) Validating

the AMRFinder tool and resistance gene database by using antimicrobial resistance

genotype-phenotype correlations in a collection of isolates. Antimicrob Agents

Chemother 63: e00483-19.

77.   Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S (2018) High

throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries.

Nat Commun 9: 5114.

Supporting information

S1 Fig. Comparison of the genome structure between strain 04Ya249 with 04Ya108, CAIM

1831 and TJ249. Structure comparison was performed using nucmer in MUMmer3 [74].

Purple dots indicate a match on the Watson strand (5' to 3' on the top strand in GenBank

file), and light blue indicates a match on the Crick strand (5' to 3' on the bottom strand). (B)

Average nucleotide identity (ANI) between two strains as determined by fastANI [77]. The

commands used were as follows: (A) $nucmer -minmatch

60../../data/04Ya249_submission.fas../../data/reference.fas

$mummerplot -x "[0,6000000]" -y "[0,22000000]" -postscript -p test out.delta; (B) $fastANI -

q../../data/04Ya249_submission.fas -r../../data/CAIM1831_Refseq.fas -o

04Ya249vsCAIM1831.txt

$fastANI -q../../data/04Ya249_submission.fas -r../../data/04Ya108_submission.fas -o

04Ya249vs04Ya108.txt

$fastANI -q../../data/04Ya249_submission.fas -r../../data/04Ya108_submission.fas -o

04Ya249vs04Ya108.txt

S2 Fig. Genetic map of pSEA2. (A) Locations of antimicrobial resistance (AMR) genes and SE-

6945 in pSEA2. AMR genes were inferred using AMRFinderPlus [76]. Genes were visualized

using CLC Sequence Viewer (Qiagen, Hilden Germany). (B) Location of SEs in pSEA1 and

pSEA2. Four SE core genes are indicated by four distinct colors. Red pentagons are the β-

lactamase gene.

S3 Fig. *attL* sequences formed in the SE-6945 integration regions. Sequences are derived

from 04Ya249 genomes (AP019851.1, AP019849.1), TJ239 genome (*insJ*, *yjjNt*), and *Vibrio*

harveri strain WXL538 (CP045070.1) carrying a SE-6945-like element carrying a different set

of cargo genes. Sequences in red are incorporated into circular copy of SE. 6-bp underlined

sequence is incorporated into *attS* during SE circle formation. 19 bp motif C highlighted in

yellow form an imperfect inverted repeat with the motif C' located the other terminus of SE.

Sequence highlighted in cyan is a putative transcription termination motif. Asterisks indicate

conserved nucleotides.

S4 Fig. *attR* sequences formed in the SE-6945 integration regions. Sequences are derived

from 04Ya249 genomes (AP019851.1, AP019849.1), TJ239 genome (*insJ*, *yjjNt*), and *Vibrio*

harveri strain WXL538 (CP045070.1). Sequences in red are incorporated into circular copy of

SE. The 6 bp underlined sequences originate from the original *attB* or SE circle, and they are

not incorporated into the *attS* of *the* SE circle.

S5 Fig. Inference of SE-6945 insertion number and its integration location by Southern

hybridization. (A) *in silico* restriction map of DNA segments around SE-6945 integration

locations in strain 04Ya249, TJ249, and a hypothetical *E. coli* strain carrying a pSEA2

insertion into SE-6945 integrated into *yjjNt*. Locations of SE-6945 and *intA* probe

hybridization region are indicated by the yellow square and black filled square, respectively,

above the restriction map. Horizontal black lines indicate the fragments detected by the

probe and their sizes. (B) Southern blots of the pSEA2-free Vibrio strain LN95 and the

parental strain 04Ya249. Genomic DNA was double digested with *NdeI* and *HindIII* (the left

two lanes) or *NdeI* and *SphI* (the right two lanes). The probe used was 5' end of *intA*. Four

unique bands originate from fragments shown in the first two rows (04Ya249 pSEA2,

04Ya249 chr1) in panel A. (C) Southern blots of 19 *E. coli* transconjugants obtained from 19

independent mating assays. Upper panel shows digestion with *NdeI* and *SphI*. Lower panel

shows digestion with *NdeI* and *HindIII*. The color of the strain name indicates the pattern of

SE/pSEA2 insertion deduced from restriction maps in panel A: blue, three SE copies

(*yjjNt*::SE-6945, *insJ*::SE-6945::pSEA2); red, two SE copies (*insJ*::SE-6945::pSEA2); purple,

(*yjjNt*::SE-6945::pSEA2); lime, one SE copy (*yjjNt*::SE-6945 or chr::pSEA2); Strains with

asterisk (*) are expected to carry pSEA2 in an unknown chromosomal location without

generating a SE-6945-chromosome junction.

S6 Fig. *attL* sequences formed in the SE-6283 insertion regions. Sequences are derived

from strain 04Ya108 (AP024165.1, AP024167.1), *E. coli* transconjugant TJ108W0 sequenced

in our previous study [30], and *Vibrio* strains carrying SE-6283-like elements listed in Fig 3.

Sequences in red are incorporated into circular copy of SE. Incorporation of the 6 bp

underlined sequence of TJ108W0 and pSEA1 into *attS* was demonstrated in the previous

study.

S7 Fig. *attR* sequences formed in the SE-6283 insertion regions. Sequences are derived

from genomes listed in Fig 3. Regarding SE-6283, the 6 bp underlined sequences in *attR*

originate from inserting SE circle, and they are not incorporated into *attS* upon SE

circularization.

S8 Fig. PCR detection of SE-6945 circle in *V. alfacsensis* strain 04Ya249 and *E. coli* strain

LN52. (A) PCR detection of chromosomal *attL*, *attR*, *attS* in 04Ya249*.* (B) Relative copy

number of *attS* in *V. alfacsensis* LN95 (no pSEA2). Copy number is represented as relative

copy number to *gyrB*. Primers used are shown in Table 2. (C) PCR detection of *attS* in pSEA2

free *E. coli* strain LN52.

S1 Table. MIC of transconjugant strains

S2 Table. Data availability

A

IS3
insJ

yjjN-yjjO
intergenic (*yjjNt*)

E.coli MG1655 (U00096.3)

3,721,004

4,597,068

Transconjugant TJ249

intA

ARGs

7.1 kb

pSEA2

B

SE-6945
7.1 kb

C   C′

intA   CDS2   *intB*   CDS4   *bla*_{GMA-1}

408 aa   581aa   720 aa   135 aa   cargo

SE-6283
(Tn*6283*)

C   C′

intA   CDS2   *intB*   CDS4   cargo

13.8 kb   467 aa   773 aa   731 aa   165 aa

5′   3′   Terminus

SE-6945   C′   TAACTTTATGTTAAGGGTA
            C   .TTAGA......GGCT...

SE-6283   C′   TAAAGAATTTCCAAGGGTA
            C   ....A......T......G

C

                                G R   Box I
λ-Int   (NP_040609.1)   196 --PCWLRLAMELAVVTGQRVGDLCEMKWSDIV---DGY-------- 228
XerD_Eco   (AAC75932.1)   130 PLELRDKAMLEVLYATGLRVSELVGLTMSDIS---LRQG------ 165
Int_SXT   (AAL59748.1)   225 -ITEKNRIYIKLLLLLGGRKGELIQAEKHHFD---LQSA------ 259
IntI1_R388 (HXF36_RS00185)   131 ---GEHRLFAQLLYGTGMRISEGLQLRVKDLD---FDHGT------ 164
IntA_6945   (BBM67680.1)   196 -ISGEFRLLVATVINSGLRAIEVADIKPRHFK---VPKGFKGKTYT 237
IntA_6283   (BCN27155.1)   264 --PKAANLIFRLCLEAGLRIDEATHFPSSDIG---IADCSDLDVVP 304
IntB_6945   (BBM67682.1)   343 ----EGACFGACSAFTGMRISELTQIEGDSYKEVDIDGVK------ 378
IntB_6283   (BCN27153.1)   349 -----GGCLWGLMARTGMRGDEMYSLNTVQGCTKDIIDRQN----- 385

                                H R   Box II   H   Y
λ-Int   (NP_040609.1)   299 ---SFEGDPPTFHELR-SLSARLYEKQ---------ISDKFAQHLLGHKSDTMASQYRDD 345
XerD_Eco   (AAC75932.1)   237 ----DSEKLSPHVLRHAFATHLLNHG---------ADLRVVQMLLGHSDLSTTQIYTHV 282
Int_SXT   (AAL59748.1)   331 SLS-VEMEHWSMHDLRRTMRTRMSAIT----------TQEVAELMIGHSKKGLDAIYNQY 379
IntI1_R388 (HXF36_RS00185)   271 -----TKPATPHTLRHSFATALLRSG---------YDIRTVQDLLGHSDVSTTMIYTHV 315
IntA_6945   (BBM67680.1)   328 ---SNPHFDHDFHDTRATYAANKLDLLLNIPDLSTTQALKLLKDELGHKDLSVTMRYLTH 384
IntA_6283   (BCN27155.1)   389 ----DRSWYYRIHDLRSTFATHWLWNESKERDVGYDYLMDELKELMGHSSTSITEKYIKY 444
IntB_6945   (BBM67682.1)   500 NERGELYWRFNTHSLRRTFAHFVVGHGL--------VSLASLKHQFKHIHLAMTAIYASH 551
IntB_6283   (BCN27153.1)   487 SFELGSNYEFTPHQLRRSFAYYLIGYEL--------CNFPQLKQQFSHVSMAMTRHYAKN 538

A

B

attB_pSEA2_AP019851.1	GCGCCTTTTCATATTGGCATATCGCCATGT**xxxxxx**TGTTTTCGATGTCACTATTGAGCTTACGAA
attB_Eco(insJ)_U00096.3	CTTTTTCTGGATCAACTGATATTGCAATG**TTCTTT**TGGTGCCAATCTTGAGCGCGCGTAAACCAG
attB_Eco(yjjNt)_U00096.3	TGCAGCGTCCGGGGAAGTGTTGGGCGCTG**TTTTTT**TGTTTCTTAATAATGTGTTGTAAGCCGTAG
attB_Val_CAIM1831_CP032093.1	GATACAAAAGCCATGCAGACAACGCATGG**CTTTTTT**TGTACTCTAA-TGAAAAATAGAATTAGGGA
attB_Vha_ATCC33843_CP009467.2	ATTCAAAAGCCATGCAGACAATGCATGG**CTTTTTT**TGTACTCTAAATGGAAAATAGAATTAGGGA

C

A

B

```
attB_Vro_B64D1_Chr2_CP018312.1    TGCTGACCAGTTCGGTGTGTGGGGCGAGAAAAAGTTCATGGGTAAAGTTTACGATGGTCTTCACCG
attB_Vsc_FP3289_MDCJ01000002      GGCAGAACAGTTTGGCGTTTGGGGCGAGAAAGAAATTTATGGGTAAAGTGTATGATGGCTTGCACCG
attB_Valg_ATCC17749_CP006718.1    TGCAGACCAGTTCGGCGTTTGGGGCGAAAAGAAATTCATGGGCAAAGTATACGACGGTCTTCACCG
attB_Val_CAIM1831_CP032093.1      TGCAGAACAGTTTGGTGTTTGGGGTGAAAAGAAGTTCATGGGTAAAGTGTACGACGGACTTCACCG
attB_Eco_W3110_AP009048.1         GTGCGAACAATTCGGCGTCTGGGGTGAAAAGTCCTTCATGGGCAAAACCTACGATGGCATTCATCG
attB_Val_pSEA2_AP019851.1         CGTCTATGCTATTGATTCTCTTTATGAAGAAGAGGGCAAAGGCGTCAAATATGAGTTCGTATACAA
```

C