1 Title Page

2 Uncovering the Genetic Profiles Underlying the Intrinsic Organization of

3 the Human Cerebellum

Running Title: **Genetic Basis of Cerebellar Functional Organization**

Yaping Wang[1,2,3,4,15], Lin Chai[2,3,4,15], Congying Chu[2,3,4*], Deying Li[2,3,4], Chaohong Gao[1,2,3,4],

Xia Wu[2,3,4], Zhengyi Yang[2,3,4], Yu Zhang[5], Junhai Xu[6], Jens Randel Nyengaard[1,7,8], Simon

B. Eickhoff[9,10], Bing Liu[11], Kristoffer Hougaard Madsen[1,12,13], Tianzi Jiang[1,2,3,4,14],

Lingzhong Fan[1,2,3,4,14*]

[1]Sino-Danish Center, [2]University of Chinese Academy of Sciences, 100190, Beijing, China

[3]Brainnetome Center, [4]National Laboratory of Pattern Recognition, Institute of Automation,

Chinese Academy of Sciences, 100190, Beijing, China

[5]Research Center for Healthcare Data Science, Zhejiang Lab, 311100, Hangzhou, China

[6]School of Computer Science and Technology, Tianjin Key Laboratory of Cognitive

Computing and Application, Tianjin University, 300350, Tianjin, China

[7]Core Centre for Molecular Morphology, Section for Stereology and Microscopy,

Department of Clinical Medicine, Aarhus University, 8000, Aarhus, Denmark

[8]Department of Pathology, Aarhus University Hospital, 8200, Aarhus, Denmark

[9]Institute of Neuroscience and Medicine (INM-7: Brain and Behaviour), Research Centre

Jülich, 52425, Jülich, Germany

[10]Institute of Systems Neuroscience, Heinrich Heine University Düsseldorf, 40225,

Düsseldorf, Germany

[11]State Key Laboratory of Cognitive Neuroscience and Learning, Beijing Normal

University, 100875, Beijing. China

[12]Informatics and Mathematical Modelling, Technical University of Denmark, 2800,

Kongens Lyngby, Denmark

[13]Danish Research Centre for Magnetic Resonance, Centre for Functional and Diagnostic

Imaging and Research, Copenhagen University Hospital-Amager and Hvidovre, 2650,

Hvidovre, Denmark

[14]CAS Center for Excellence in Brain Science and Intelligence Technology, Institute of

Automation, Chinese Academy of Sciences, 100190, Beijing, China

[15]These authors contributed equally: Yaping Wang, Lin Chai.

* To whom correspondence should be addressed:

Email: lingzhong.fan@ia.ac.cn or chucongying@gmail.com

Abstract

The functional diversity of the human cerebellum is largely believed to be derived more from its extensive connections rather than being limited to its mostly invariant architecture. However, whether and how the determination of cerebellar connections in its intrinsic organization interact with microscale gene expression is still unknown. Here we decode the genetic profiles of the cerebellar functional organization by investigating the genetic substrates simultaneously linking cerebellar functional heterogeneity and its drivers, i.e., the connections. We not only identified 443 network-specific genes but also discovered that their co-expression pattern correlated strongly with intra-cerebellar functional connectivity (FC). Ninety of these genes were also linked to the FC of cortico-cerebellar cognitive-limbic networks. To further discover the biological functions of these genes, we performed a "virtual gene knock-out" by observing the change in the coupling between gene co-expression and FC and divided the genes into two subsets, i.e., a positive gene contribution indicator (GCI^+) involved in cerebellar neurodevelopment and a negative gene set (GCI^-) related to neurotransmission. A more interesting finding is that GCI^- is significantly linked with the cerebellar connectivity-behavior association and many recognized brain diseases that are closely linked with the cerebellar functional abnormalities. Our results could collectively help to rethink the genetic substrates underlying the cerebellar functional organization and offer possible micro-macro interacted mechanistic interpretations of the cerebellum-involved high order functions and dysfunctions in neuropsychiatric disorders.

Introduction

Converging evidence from animal and human studies is advancing our understanding of the human cerebellum, which is engaged in motor, complex cognitive, and emotional behaviors [1-4]. While such functional diversity of the cerebellum was believed to derive more from its extensive afferent and efferent connections to extra-cerebellar structures, rather than being limited to a regular lattice-like anatomic feature of cerebellar-cortical cytoarchitecture [1, 5-8]. Considering the widely accepted understanding that the macroscale functional organization of the human nervous system is ultimately regulated by the underlying microscale gene expression [9-12], it is thus intriguing to unravel the genetic profiles underlying the cerebellar functional organization. As yet, what remains unclear is whether and how the hypothesized determination of cerebellar connections in its intrinsic functional organization interact with microscale gene expression.

To date, the genetic mechanism supporting the functional organization of the human cerebellum is mainly unknown. Only a few studies have attempted to investigate the gene expression pattern of the human cerebellum, but they provided inconsistent results in gene expression variability. For instance, Hawrylycz et al. [13] and Negi and Guda [14] both found that gene expression is highly homogeneous across the anatomical regions of the healthy adult cerebellum. In contrast, Aldinger et al. [15] and Wang and Zoghbi [12] found that cerebellar development and function are governed by the precise regulation of molecular and cellular programs and that the gene expression pattern is heterogeneous across spatial and temporal scales. In addition, differences in gene expression patterns between the cerebellar gyri and sulci [16], and considerable cerebellar regional specializations containing specific cell types, as revealed by high-throughput single-

nucleus RNA-seq [17] have been found in the mouse cerebellum. Because relevant studies

that showed homogeneity [13, 14] explored the overall cerebellar genetic expression

pattern across its gross macro-anatomical boundaries (e.g., cerebellar lobules) and might

have failed to fully reflect the functional organization of the human cerebellum [18, 19].

This inconsistency in the genetic variability of the cerebellum needs to be further explored.

In the past decade, functional topological maps describing the organization of the

human cerebellum using task [20] and task-free functional magnetic resonance imaging

(fMRI) [21, 22], specifically, separate cerebellar functional networks [21, 22] and intra-

cerebellar functional gradients [23], have been proposed. In particular, Buckner et al. [21]

employed resting-state functional connectivity (FC) of the cerebello-cortical circuit as a

tool to map the intrinsic functional architecture of the human cerebellum and proposed a

possible functional parcellation into 7 networks and 17 networks. It is thus possible to

decode the genetic profiles of the cerebellar functional organization by investigating the

molecular genetic substrates linking cerebellar functional heterogeneity and its drivers, i.e.,

the connections. One promising approach is imaging-transcriptomics analysis [24-26],

which allows the brain-wide spatial analysis of microscopic transcriptome data combined

with macroscopic neuroimaging phenotypes [9]. Moreover, it also offers the opportunity

to link the transcriptome data with the behavior variations via the neuroimaging [27]. These

cross-scale analyses could provide a better understanding of the interaction between

microscale gene expression and macroscale functional network, and ultimately involved in

the individual behaviors, and importantly, their putative multiscale interactions in

cerebellar related diseases [28].

Thus, our goal was to investigate the neurobiological genetic substrates underlying the

functional organization of the human cerebellum. More specifically, we will seek to

address the following three progressive questions (Fig. S1). Are there differentially

expressed genes in the diverse intra-cerebellar functional networks with largely invariant

cytoarchitecture? How do gene expression and cerebellar connectivity relate to each other,

and is there any link between this association and human behavior, as well as brain disease?

In the current study, we first examined network-specific genes by differential gene

expression analysis across diverse cerebellar functional networks characterized by largely

invariant cytoarchitecture. This analysis allowed us to further investigate the genetic

explanations to the cerebellar inconsistency between functional heterogeneity and

cytoarchitecture near-homogeneity. Then to explore the relationship between these

network-specific genes with the cerebellar connection, we constructed the gene co-

expression matrix using the network-specific genes and found it is highly correlated with

both intra-cerebellar and cerebello-cortical cognitive-limbic FC. Furthermore, we tested

the contribution of each network-specific gene to this correlation by virtual gene knock-

out (KO) and divided the network-specific genes into two subsets, i.e., a positive gene

contribution indicator (GCI^+) and a negative gene set (GCI^-). We found that the GCI^+

appears to be mainly involved in cerebellar neurodevelopment. Whereas GCI^- seems to be

related to neurotransmission, emotion-cognitional behaviors and is significantly enriched

in various neuropsychiatric disorders that are closely linked with cerebellar functional

abnormalities. Together, the current exploration provides a starting point for associating

the genetic and behavior markers of the functional network to cerebellar involvements in

higher order non-motor functions and dysfunctions in various neuropsychiatric disorders

[1, 29, 30].

Materials and Methods

The schematic of the experimental design is depicted in Fig. 1 and includes three steps. Step 1 (Fig. 1A): to investigate whether network-specific genes occurred across cerebellar functional networks, we performed the differential gene expression analysis based on the combination of the Allen Human Brain Atlas (AHBA) transcriptome data [9] with a cerebellar functional parcellation atlas [21]. Step 2 (Fig. 1B): then, the co-expression matrix was constructed using the network-specific genes from step 1 and compared with FC to explore their overall correlation (referred to as the Gene-FC correlation for simplicity). Meanwhile, the relationship between gene and cerebello-cortical FC also has been explored. Step 3 (Fig. 1C): the "virtual gene knock-out" was leveraged to examine the direction of each gene's contribution to this Gene-FC correlation from step 2 and used to separate the network-specific genes into two subsets. Furthermore, we applied a series of functional annotation tools to explore the role of these genes, including gene enrichment analysis, Behavior-FC-Gene mapping analysis, disease enrichment analysis, and integrative temporal specificity analysis.

A. Step 1: Differential gene expression analysis

Sample assignment

Example visual network
Vs. other 6 networks by limma

443 network-specific genes

Network	Specific genes
Control	0
Default	0
DorsAtten	51
Limbic	170
VentAtten	0
Som/Motor	3
Visual	221

B. Step 2: Correlation between gene co-expression and FC

C. Step3: Functional annotation

Virtual knock-out
(virtual KO)

Gene contribution indicator
$$GCI_X = R_{X-} - R_{All}$$

GCI+ gene set   GCI- gene set

Behavior-FC-Gene
mapping analysis

Gene enrichment
analysis

Integrative temporal
specificity analysis

Fig. 1. Analysis pipeline. A Step 1: differential gene expression analysis. We assigned the AHBA cerebellar samples into seven cerebellar functional networks (left) [21] and averaged each gene's expression within the same network individually. Then we compared the gene expression in each network with all the other networks by limma [31] (middle) with a fold change > 1 and $p < 0.05$ (FDR corrected) as an indicator (Red indicates that the genes were found significantly positively expressed in the visual network.). Thus, we obtained the network-specific genes for the seven networks (right). **B Step 2:** correlations between the gene co-expression and the FC included intra-cerebellar and cerebello-cortical circuit. Intra-cerebellum: for each pair of networks, we calculated the gene expression similarity between them using the cerebellar network-specific genes and then constructed the gene co-expression matrix. The FC matrix was constructed by correlating the BOLD signal for all pairs. Then the relationship between the genetic correlation and functional correlation was evaluated. Cerebello-cortical circuit: we first defined the cortical network-specific genes as we did for the cerebellum and tested whether any convergently differentially expressed genes occurred. Then we used the overlapping genes to obtain the cortical genetic correlation for each cerebellar network and evaluated the relationship between the cortical genetic and functional correlation for each cerebellar network. **C Step 3:** functional annotation includes virtual gene knock-out (KO), Behavior-FC-Gene mapping, gene enrichment analysis, and integrative temporal specificity analysis.

AHBA preprocessing

The AHBA [9] is a publicly available transcriptome dataset (http://www.brain-map.org), which provides normalized microarray gene expression data from six adult donors (ages 24, 31, 34, 49, 55, and 57 years old; $n = 4$ left-hemisphere only, $n = 2$ both left and right

hemispheres). Table S1 shows the demographic information. Tissue collection was

approved by Institutional Review Boards of the Maryland Department of Health and

Hygiene, University of Maryland Baltimore and University of California Irvine, and the

informed consents were obtained from decedent next-of-kin [9]. Although AHBA provides

gene expression from only six adult donors, it still has many unprecedented advantages.

While some existing human gene expression atlases just cover multiple brain regions, only

the AHBA delivers high-resolution coverage of nearly the entire brain [32]. It includes the

expression of more than 20,000 genes taken from 3,702 spatially distinct tissue samples [9]

ranging from the cerebral cortex to the cerebellum.

Referred to Anderson et al. [26], the current preprocessing pipeline included data

filtering, probe selection, sample selection, and assignment. We first filtered the probes

with the AHBA binary indicator to mitigate the background noise and excluded probes

without an Entrez ID. Then for the genes that corresponded to two or more probes, we

chose the probe with the maximum summed adjacency to represent the corresponding gene

expression; otherwise, we retained the probe with the highest mean expression, using the

CollapseRows function [33] in R. The first two steps generated 20,738 unique mRNA

probes, which provided expression data for 20,738 genes. As suggested by Arnatkeviciute

et al. [32] and given the known transcriptional differences [13] between the cortical and

sub-cortical regions and the cerebellum, we separated the cortical and cerebellar samples a

priori based on the slab type and structure name provided by AHBA and processed them

separately later. In the end, 337 samples were retained for the cerebellar cortex and 1,701

samples for the cortical cortex.

Finally, we respectively assigned these 337 cerebellum samples and 1,701 cortical

samples into the cerebellar functional networks atlas [21] and cortical functional networks

atlas [34], both of which have 7- and 17-network parcellation strategies. For each cerebellar

sample, we first generated a single $1 \times 1 \times 1$ mm^3 region of interest (ROI) at the MNI

coordinate for each sample using the AFNI 3dmaskdump -nbox function [35]. The network

label from either 7- or 17-network parcellation was assigned if the ROI fell within a

cerebellar network of the Buckner atlas. Considering the uneven and discrete sampling of

the AHBA data [9], if the $1 \times 1 \times 1$ mm^3 ROI did not overlap with any network, the

associated ROI was expanded to $3 \times 3 \times 3$ mm^3. And if the $3 \times 3 \times 3$ mm^3 ROI overlapped

with the functional atlas, the network that had the maximum number of shared voxels with

the ROI was assigned. Otherwise, the steps above were repeated for a $5 \times 5 \times 5$ mm^3 ROI.

The cerebellar samples were excluded if the $5 \times 5 \times 5$ mm^3 ROI did not overlap with any

cerebellar networks. Tables S2 and S3 show the distributions of the cerebellar sample

assignment for the 7-network and 17-network atlases. The assignment of the AHBA

cortical samples into the cortical functional network atlas was consistent with the method

used for the cerebellum, and the cortical sample distributions are shown in Tables S4 and

S5.

Differential gene expression analysis across functional networks

The gene expressions of the cerebellar samples within the same network were averaged for

each gene across the samples individually, resulting in 20,738 genes $\times$ 7 or 17 networks

matrices for each donor. Here we used as many of the most original overall genes as

possible, because some existing methods [13, 32] for screening genes do not consider the

specificity of gene expression within the human cerebellum. Then we calculated the

differentially expressed genes across the 7 networks using the R limma package [31] by

comparing the gene expression in one network (e.g., control) with the remaining 6

networks (e.g., default, limbic, visual, etc.). Since the gene expression vectors of the same

network derived from different donors can be treated as biological replicates [26], the

limma's duplicatedCorrelation tool [36] was leveraged to evaluate the independence

between replicates. Specifically, the correlations between replicates were estimated using

the duplicateCorrelation function, which individually fits a mixed linear model by

restricted maximum likelihood for each gene and returns a consensus correlation [36]. This

consensus correlation between replicates would then be incorporated into the limma's

linear model to preserve more information than simply averaging the biological replicates

[36]. The traditional minimum fold change threshold was unsuitable for determining

biologically meaningful but subtly different expressions [37], in particular for the

cerebellum, due to the highly homogenous cytoarchitecture. Instead, we applied the

Benjamini-Hochberg (BH) method to control the false discovery rate (FDR), and the

statistical threshold $p \leq 0.05$ (FDR corrected) combined with a fold change > 1 was used

as the key indicator for differentially expressed genes. For simplicity, the differentially

expressed genes across cerebellar networks are referred to as cerebellar network-specific

genes throughout this paper. The cortical network-specific genes were identified in the

same way. The only difference was that the gene expression of the cortical samples was

first averaged within each parcel (51 and 114 parcels, which corresponded to the 7- and

17-networks, respectively) [34] and then averaged within each network.

To test the specificity of these network-specific genes derived based on task-free 7

functional networks, we also applied the differential gene expression analysis with the

same procedure in other different cerebellar atlases. There are the different resolution of

the same task-free functional atlas [21], i.e., 17-network parcellation, independent task-

based multi-domain task battery (MDTB) [20] functional atlas and different resolutions,

i.e., 11-, 28-lobular parcellations, of the anatomical atlas [38]. Then we estimated the

overlapping genes between the network-specific genes derived based on the task-free 7

functional networks with other atlases. Moreover, we applied the differential gene

expression analysis in only 4 left-hemisphere donors with the same procedure to test the

sensibility of these network-specific genes.

Cerebellar resting-state functional connectivity (FC)

The minimally preprocessed [39, 40] Human Connectome Project (HCP) S1200 release

dataset [41], which has 1,018 subjects (aged from 22 to 37 years old) with both structural

MRI and resting-state functional MRI (rs-fMRI, HCP S1200 manual), was used. The

preprocessing pipeline includes artifact correction (correction of gradient nonlinearity

distortion, realignment for head motion, registration of fMRI data using structural data,

reduction of geometric distortions due to B0 field inhomogeneity, etc.) as well as denoising

by ICA-FIX [42, 43]. Time courses were extracted from these CIFTI grayordinate-format

preprocessed rs-fMRI images, and the global signal was regressed as well. The resting-

state blood oxygen level-dependent time series were averaged within each cortical parcel

of the 7- or 17-network cortical atlases and within each cerebellar network of the 7- or 17-

network cerebellar atlases [21], separately. The FC within the cerebellum was computed

using Pearson's correlation for the averaged time courses for each interest ROI. Because

four runs were performed for each subject, the correlation values were separately calculated

for each run, Fisher's z-transformed, and averaged across the runs, resulting in a 17 $\times$ 17

cerebellar networks matrix. The same process was used to calculate the correlations

between each functional cerebellar network and each cortical parcel, resulting in a 114

cortical parcels $\times$ 7 cerebellar networks functional correlation matrix, which represents

the FC across the cerebello-cortical circuit. Regardless of whether the FC was within the

cerebellum or across the cerebello-cortical circuit, both categories of FC were defined

using the more fine-grained 17-network parcellation to increase the spatial resolution. The

only exception was that the cerebellar 7-network was applied while calculating the FC

across the cerebello-cortical circuit to compare each cerebellar network directly.

Correlation between gene co-expression with intra-cerebellar FC

To fully capture the genetic correlation with the FC within the cerebellum, we leveraged

the genetic samples of the two bi-hemisphere donors when constructing the gene co-

expression matrix since the FC of the cerebellum is bilateral. Therefore, the gene co-

expression was analyzed for the two bi-hemisphere donors using the network-specific

genes derived from all six donors across 7 networks, using a finer 17-network parcellation

to increase the spatial resolution. Ten networks that contained samples from both bi-

hemisphere donors were retained (Table S3). For each bi-hemispheric donor, the log2 gene

expression of the cerebellar samples was mean-normalized and then averaged within each

network. The cerebellar 10 $\times$ 10 networks correlation matrix was calculated using

Spearman's correlations individually (since the non-normality of gene expression data),

then Fisher's z-transformed, and finally averaged to construct the final 10 networks gene

co-expression matrix. The correlation significance level of the gene co-expression was

evaluated using the overlap between the correlation significance matrix for these two

individuals after being adjusted by Bonferroni correction. Meanwhile, we transformed the

17×17 networks FC matrix into a 10×10 networks size to be consistent with the gene

co-expression matrix. Finally, the relationship between the 10×10 networks gene co-

expression and the 10×10 networks FC matrix was computed using Pearson's

correlation. The correlation between the gene co-expression and FC is referred to as the

Gene-FC correlation throughout the present paper for simplicity.

To test the significance of the Gene-FC relationships while taking into account the

spatial autocorrelation (SA), we leveraged Brain Surrogate Maps with Autocorrelated

Spatial Heterogeneity (BrainSMASH) [44] to generate 10,000 surrogate maps which

persevered the genetic SA of the 443 genes $\times$ 106 samples matrix. The 106 samples were

equivalent to the amount of samples for the 10 networks that were used to construct the

gene co-expression matrix since they contained samples from both bi-hemisphere donors.

The parameter "knn" was set to 100 to keep it roughly consistent with the number of

samples. These 10,000 surrogate maps were used to construct the gene co-expression

matrix and thus generate an empirical null distribution of the Gene-FC correlation while

taking SA into account. The p value (p_{SA}) was defined as the proportion of correlation

values produced by the surrogate maps that exceeded the correlation coefficient for the real

data.

In addition, to evaluate the robustness of the verified Gene-FC correlation within the

cerebellum, we also recalculated it using several different parcellations, i.e., task-free 7-

network parcellation, and independent task-based MDTB functional parcellation [20]. The

criteria for each step were consistent with our primary method. Moreover, we employed a

control test to learn whether the Gene-FC correlation could be obtained using only the

network-specific genes. That is, no Gene-FC correlation while using other genes. We

randomly selected 443 genes from the full gene set without the network-specific genes and

referred to them as non-network-specific genes. Then we calculated the Gene-FC

correlation using the non-network-specific genes and ran this step randomly 10,000 times.

In addition to these thresholdless non-network-specific genes, we applied a set of

thresholds to the averaged original log2 gene expression data to confirm that these non-

network-specific genes were expressed in the cerebellum. Doing this can also help us test

whether the gene co-expression pattern constructed using these threshold non-network-

specific genes was correlated with FC.

Lastly, we also verified the robustness of this Gene-FC correlation using the 218

unrelated participants from the HCP S1200 release [41]. Besides, to constitute an

independent neuroimaging validation dataset, we leveraged 296 participants with four runs

(aged 38−58 years old) from preprocessed HCP-Aging Lifespan 2.0 Release [45]. This

dataset, together with the 1,018 participants from the HCP S1200 release of our primary

strategy, neatly covered the age ranges of AHBA (extends from 24 to 57 years old). The

analysis strategy for these two additional datasets is similar to our primary approach.

Correlation between gene co-expression and FC across the cerebello-cortical circuit

To thoroughly investigate the cerebellar functional organization, we also explored the

relationship between the cerebello-cortical FC and the genetic correlation based on the

strategy used in Anderson et al. [26]. First, we defined the network-specific genes in the

cortex using the same procedure as we had for the cerebellum and examined the genes that

overlapped within the same network of the cerebellum and the cortex. Then the gene co-

expression matrix was constructed between 6 cerebellar networks and 59 cortical parcels

from the two bi-hemisphere donors, using the 90 unique genes derived from the overlap

between the cortical network-specific genes and the cerebellar network-specific genes.

Here, the cerebellar 7-network parcellation was selected to compare the different cerebellar

networks directly. The visual network was excluded because it only had two samples from

one of the 2 bi-hemisphere donors. For the cerebral cortex, 59 cortical parcels that

contained samples from both bi-hemisphere donors were estimated. The log2 mean-

normalized expression within each cerebellar network and each cortical parcel was

estimated individually and correlated using Spearman's ρ, Fisher's z-transformed, and

averaged. We transformed the 114 cortical parcels $\times$ 7 cerebellar networks FC matrix into

59 cortical parcels $\times$ 6 cerebellar networks size to be consistent with the gene co-

expression matrix. Finally, the relationship between the cortical genetic correlation and the

cerebello-cortical FC matrix was computed using Pearson's correlation across six

cerebellar networks and adjusted by the BH method to correct for multiple comparisons.

Gene functional annotation

Virtual gene knock-out (KO)

To extend our investigation of the overall relationship between gene co-expression and FC

within the cerebellum, we referred to a similar previous approach [46, 47] and termed it

the "Virtual Gene Knock-out (KO)" to evaluate each gene's contribution to the Gene-FC

correlation. In brief, we deleted each of the 443 cerebellar network-specific genes one-by-

one to simulate the gene KO, then constructed the gene co-expression matrix without that

gene, analyzed the correlation between the FC and the gene co-expression, and finally

calculated the difference in the correlation coefficient between before and after the

simulated deletion, with the result being defined as the gene contribution indicator (GCI)

[46]. Based on the GCI, we identified two different gene sets that had opposite effects on

the Gene-FC correlation: a GCI positive gene set (GCI$^+$) and a GCI negative gene set

(GCI$^-$). The virtual KO of GCI$^+$ increased the Gene-FC correlation, and, accordingly, its

expression decreased the Gene-FC correlation; in contrast, the virtual KO of GCI$^-$

decreased the correlation, and, accordingly, its expression increased the Gene-FC

correlation.

Behavior-FC-Gene mapping analysis

To evaluate the relationship between these genes and human behaviors, we leveraged the

stable brain-behavior space strategy proposed by Ji et al. [27], which could be briefly

divided into the following three sections. These procedures were called Behavioral-FC-

Gene mapping analysis to facilitate a better understanding and fluency of the paper.

(1). Principal component analysis (PCA) of HCP behavioral data: 59 behavioral

measures [48] (Table S6) covering six behavior categories that represent the general

domains of human behavior were used, including alertness, cognition, emotion, motor,

personality, and sensory. The 218 HCP unrelated participants were selected to avoid

violation of the exchangeability assumption of permutation tests, and subjects with missing

data were excluded, resulting in 211 participants. The significant principal components

(PCs) were calculated by a permutation test, which randomly and independently shuffled

subjects' order for each behavior measure to re-run PCA 10,000 times to establish the null

model. The PCs that explained variance exceeded chance ($p < 0.05$ across permutation test)

were considered significant and retained for the next step. The nomination of PCs was

based on the pattern of loadings on the original 59 behavior measures.

(2). Mass univariate Behavior-FC map: the relationship between significant PCs scores

and individual intra-cerebellar FC between 17 networks was quantified by a mass

univariate regression procedure called PALM tool [49] across 211 unrelated participants.

For each significant PC, the derived regression coefficients on the total 136 functional

connections were then Z-scored and referred to as the Behavior-FC map for this PC. The

significances of the regression coefficients were assessed by a permutation test, which

randomly shuffled the order of the subjects for each significant PC score 10,000 times.

(3). Behavior-FC-Gene co-expression mapping analysis: to evaluate the relationship

between the Behavior-FC map and the co-expression map of GCI^+ and GCI^-, we first

constructed these maps across the ten cerebellar networks that corresponded to the

networks containing genetic samples from both AHBA bi-hemisphere donors. The

relationship between the Behavior-FC maps of eight significant PCs and the co-expression

patterns of GCI^+ and GCI^- was calculated by the Pearson's correlation respectively, and the

significance was assessed by a permutation test which randomly shuffled the gene co-

expression pattern 10,000 times.

GO, pathway, and disorder enrichment analysis (ToppGene portal)

To characterize the biological role of GCI^+ and GCI^-, we applied the *ToppGene portal* [50]

to conduct a gene ontology (GO), pathway, and disorder enrichment analysis. The GO [51]

enrichment analysis provides ontologies to describe accumulated knowledge of genes in

three biological domains: biological process, cellular component, and molecular function.

The BH method for FDR (FDR-BH correction) ($p < 0.05$) was chosen to correct for

multiple comparisons.

Integrative temporal specificity analysis

To investigate the overall temporal expression features of these genes, we applied an online

cell type-specific expression analysis (CSEA) tool [52] to do the enrichment analysis of

the genes within the cerebellum during different lifespan windows. Here, a specificity

index probability (pSI = 0.05, 0.01, 0.001, and 0.0001, permutation corrected) was used to

define the probability of a gene being expressed in each time window relative to all other

time windows to represent the varying stringencies for enrichment. The significance of the

overlap between the interest gene set and those enriched in a specific time window was

evaluated by Fisher's exact test, and the BH method for FDR (FDR-BH correction) was

chosen to correct for multiple comparisons.

Data and availability

R 3.6.1 and custom scripts were used to perform statistical analysis. All R packages were

mentioned explicitly in the text where the package was used. The code is freely available

at https://github.com/FANLabCASIA/CerebellarGeneFCCorrelation. The ToppGene

(https://toppgene.cchmc.org) and CSEA tool (http://genetics.wustl.edu/jdlab/csea-tool-2/)

which used to do the functional annotation of genes were all freely accessible. All data

needed to evaluate the conclusions in this study are present in the article and the

Supplementary Materials.

Results

The cerebellar network-specific genes derived based on the functional segregation within the cerebellum

The genes that were expressed much more in one network than in all the other six networks in the cerebellum and cerebral cortex were identified based on the differential gene expression analysis and are referred to as cerebellar network-specific genes and cortical network-specific genes, respectively. We identified 443 cerebellar network-specific genes (Supplementary Sheets 1 and 3) using all samples from six donors across 7 networks. The distribution of these network-specific genes is shown in Table 1 (Supplementary Sheet 2), which shows that these were mainly expressed in the limbic ($n = 170$), dorsal attention ($n = 51$), somatomotor ($n = 3$), and visual ($n = 221$) networks.

Table 1. Counts of significant differentially expressed genes within each network compared to other networks, referred to as the network-specific genes.

Networks	Cerebellum	Cortex	Overlapping genes
Control	0	33	0
Default	0	25	0
Dorsal Attention	51	80	0
Limbic	170	2920	56
Ventral Attention	0	103	0
SomatoMotor	3	960	2
Visual	221	3586	33
Total (unique)	443	6987	90

The cerebellar ($n = 443$, left column, Supplementary Sheets 2 and 3) and cortical network-specific genes ($n = 6987$, middle column, Supplementary Sheets 7 and 8) were defined across the cerebellar [21] and cortical [34] 7-network strategies, respectively. The rightmost column measures the overlap between the cerebellar and cortical network-specific genes for each network (Supplementary Sheets 9 and 10).

To test the specificity of these 443 network-specific genes in reflecting the difference

of gene expression across task-free cerebellar functional atlas, we applied the differential

expression analysis with the same procedure in other different cerebellar atlases (see more

details in "Methods"). More overlaps between these 443 network-specific genes with the

differentially expressed genes were observed in the different resolution of the same task-

free functional atlas, whereas less in the independent task-based functional atlas and

different resolutions of the lobular atlas (Supplementary Sheet 4). We also tested the

sensitivity of network-specific genes in only four left-hemisphere donors with the same

procedure and observed 230 cerebellar network-specific genes (Supplementary Sheet 5)

with preferentially expressed in the limbic ($n = 49$, overlap $= 44$) and visual ($n = 181$,

overlap $= 97$) networks.

Meanwhile, we obtained 6,987 cortical network-specific genes (Supplementary Sheets

6−8 and Table 1) using the same strategy and found that the cerebellar and cortical

network-specific genes distribution patterns across 7-network parcellation were highly

correlated ($r = 0.95$, $p = 0.001$). Moreover, we found that 90 of these 443 cerebellar

network-specific genes (~20%) (Supplementary Sheets 9 and 10 and Table 1) were

convergently expressed in the cerebral cortex (overlap in limbic $= 56$, somatomotor $= 26$,

visual $= 33$). These results mean that the 56 limbic genes were differentially expressed in

the limbic cortex and the limbic cerebellum and that the two somatomotor genes were

differentially expressed in the somatomotor cortex and somatomotor cerebellum, as well

as the 33 overlap genes in the visual network.

The co-expression of the cerebellar network-specific genes highly correlated with

intra-cerebellar FC

Using the 443 cerebellar network-specific genes, we constructed the gene co-expression

matrix for the two bi-hemisphere donors and explored the relationship between gene

correlation and FC within the cerebellum. Noted, the 443 network-specific genes were

defined based on 7-nework parcellation, while the genetic co-expression was constructed

using 17-network parcellation to increase the spatial precision. Across all the available

network-network pairs, the genetic co-expression correlates with the FC within the

cerebellum ($r = 0.48$, $p_{SA} = 0.008$, Fig. 2), which remains significant when assessed by the

spatial autocorrelation (SA) preserving strategy. This correlation between gene co-

expression and FC was referred to as Gene-FC correlation throughout the present paper for

simplicity. To validate the Gene-FC correlation within the cerebellum, we also leveraged

the different resolution of the same task-free functional atlas and the independent task-

based MDTB functional parcellation [20] to re-perform the aforementioned steps (Fig. 2D).

The gene co-expression and FC within the cerebellum also correlated when analyzed based

on the 7-network parcellation (Figs. 2D and S2): $r = 0.76$, $p_{SA} = 0.006$, and the MDTB

functional parcellation (Figs. 2D and S3): $r = 0.42$, $p_{SA} < 0.001$. Moreover, this Gene-FC

correlation was also obtained using unrelated participants from HCP S1200 release

(Supplementary Sheet 17): $r = 0.49$, $p_{SA} = 0.010$, and another independent HCP-Aging

neuroimaging dataset (Supplementary Sheet 18): $r = 0.45$, $p_{SA} = 0.013$.

Fig. 2. Network-specific gene co-expression correlates with functional connectivity (FC) within the cerebellum. A Genetic correlation was shown by the co-expression matrix (Supplementary Sheet 11) constructed for two bi-hemisphere donors across ten cerebellar networks using 443 cerebellar network-specific genes derived from all six donors. The 10 cerebellar networks corresponded to the networks containing samples from both bi-hemisphere donors (Table S3). *$p \leq 0.05$ (Bonferroni corrected). **B** The FC matrix (Supplementary Sheet 12) shows the functional correlation for the ten cerebellar networks using 1,018 subjects from the HCP S1200 release [41]. All passed the significant threshold

$p \leq 0.01$ (Bonferroni corrected). **C** The probability distribution of null models preserving
genetic spatial autocorrelation generated by BrainSMASH. The vertical black dashed lines
correspond to the p values of 0.05 and 0.95; the blue vertical line showed the practical
observed Gene-FC correlation ($r = 0.48$) and the corresponding $p_{SA} = 0.008$. **D** The overall
intra-cerebellar Gene-FC correlation using different atlases: task-free 7-network (orange),
17-network (red) parcellation of the cerebellar functional atlas based on the cerebello-
cortical FC, and task-based MDTB functional parcellation (blue) based on the task
activation pattern. The Pearson's correlation r and p_{SA} values are shown by the
corresponding colors. **E** The GCI$^+$ ($n = 246$, left, red) and GCI$^-$ ($n = 197$, right, blue) gene
lists were displayed on the flattened shape of the cerebellum. The word size and color
intensity both indicate the GCI value.

Therefore, the 443 cerebellar network-specific genes that we derived based on the

functional segregation of the cerebellum also correlated with the functional integration of

the cerebellum. This Gene-FC correlation was not generated by the genetic SA, so it was

consistently significant whenever using unrelated participants from HCP, independent

HCP-Aging dataset, different parcellation resolution, or independent cerebellar functional

atlas for the calculation. Moreover, the control test exhibited no Gene-FC correlation when

the gene co-expression was constructed using non-network-specific genes (Supplementary

Sheet 19) regardless of whether using a threshold or not. These findings confirmed that

these 443 network-specific genes play a crucial role in intra-cerebellar functional

organization.

Convergently expressed genes among the cerebellar and cortical network-specific

genes correlated with the FC across the cerebello-cortical cognitive-limbic networks

Since 90 of the 443 cerebellar network-specific genes were convergently expressed across

the cerebello-cortical circuit, we wanted to know whether these ~20% genes correlated

with the FC across the cerebello-cortical circuit. A correspondence between the genetic

and functional correlations was identified for the limbic (Fig. 3A): $r = 0.36, p = 0.030$ (FDR

corrected), and control networks (Fig. 3B): $r = -0.33, p = 0.034$ (FDR corrected), but was

not significant for the somatomotor: $r = -0.15$, $p = 0.394$ (FDR corrected), dorsal attention:

$r = -0.19$, $p = 0.281$ (FDR corrected), ventral attention: $r = -0.04$, $p = 0.779$ (FDR corrected),

or default: $r = 0.10$, $p = 0.544$ (FDR corrected) networks. The high cortical genetic

similarity between the limbic system and the adjacent control network: $r = -0.90$, $p < 0.001$

(FDR corrected), somatomotor network: $r = -0.55$, $p < 0.001$ (FDR corrected), and ventral

attention network: $r = -0.72$, $p < 0.001$ (FDR corrected) indicates that the gene co-

expression between the cerebellar limbic network and the cortex reflects a gradual genetic

gradient rather than genetic dissimilarity between the cerebellar limbic network and the

other cerebellar networks. In addition, while controlling the effect of the cortical genetic

similarity between the limbic and control networks, the partial correlation showed no

cortical Gene-FC correlation for the control network: $r = -0.13$, $p = 0.316$. It implies that

the significant cortical Gene-FC correlation for the control network could be explained by

the high cortical genetic similarity between the cerebellar limbic and control networks.

This observation is also consistent with the finding that convergently expressed genes were

only observed in the limbic network but not in the control network (Table 1). Overall, these

443 cerebellar network-specific genes not only correlated with the intra-cerebellar FC, but

~20% of them were also linked with the cerebello-cortical cognitive-limbic networks.

Fig. 3. Cortical genetic and functional correlation of cerebellar limbic and control networks seeds. Both were calculated for two bi-hemisphere donors across six cerebellar networks and 59 cortical parcels that contained samples from both bi-hemisphere donors. **A** Limbic: the cortical gene co-expression (Supplementary Sheet 20) was calculated using the 90 overlapping genes between the cerebellar and cortical network-specific genes by Spearman's correlation. The FC across each cerebellar network with each cortical parcel was calculated using Pearson's correlation (Supplementary Sheet 21). The cortical limbic genetic and functional correlations were correlated with each other: $r = 0.36$, $p = 0.030$ (FDR corrected). **B** Control: the cortical gene co-expression and the FC for the control network were correlated with each other: r = -0.33, p = 0.034 (FDR corrected). Notice that the color bar of gene co-expression was inverted considering the negative Gene-FC correlation for the control network.

Functional annotation revealed distinct biological properties of GCI⁺ and GCI⁻

separated by virtual KO

In addition to the overall correlation between gene co-expression and the functional integration of the human cerebellum, we investigated each gene's importance to the intra-cerebellar Gene-FC correlation by scoring the 443 cerebellar network-specific genes based on the GCI. Using the virtual gene KO procedure (see more details in "Methods"), we were able to classify the 443 network-specific genes that linked cerebellar functional segregation and integration into two groups: a 246 GCI positive gene set (GCI⁺, Fig. 2E left and

Supplementary Sheet 22) and a 197 GCI negative gene set (GCI⁻, Fig. 2E right and

Supplementary Sheet 23). The distinction between the two sets is that the virtual KO of

GCI⁺ genes increased the Gene-FC correlation, whereas the virtual KO of GCI⁻ genes

decreased the Gene-FC correlation. Based on the winner-take-all principle, GCI⁻ genes

may have a critical impact on the functional organization of the cerebellum. An example

is that the top genes, LCP1 and TESC, enable GTPase binding and calcium binding,

respectively [53], which are key functions within signaling transduction and consequently

brain functions. Therefore, we applied a range of functional annotation tools to further

explore the underlying roles of the GCI⁺ and GCI⁻.

Behavior-FC-Gene mapping analysis

Since these genes largely engaged in the cerebellar functional network which could predict

human behaviors, whether they also involved in human behaviors with the FC acting as an

intermediary role? We leveraged the Behavior-FC-Gene mapping analysis (see more

details in methods) to explore the relationship between the co-expression of GCI⁺ and GCI⁻

with the human behaviors. Based on the PCA of 59 behavior measures (Table S6) across

211 HCP unrelated participants, 8 significant PCs were derived (Supplementary Sheet 24).

Then the Behavior-FC map of each significant PC calculated by mass univariate regression

procedure was compared with the co-expression pattern of GCI⁺ and GCI⁻ respectively

(Supplementary Sheet 25). Among the eight significant PCs, we observed correlations of

the Behavior-FC maps of PC1, PC3, and PC4 (Supplementary Sheet 25) with the GCI⁻ co-

expression, rather than the GCI⁺.

But the PC1 and PC4 had few regression coefficients (1.47% and 0.74%, respectively)

in the Behavior-FC map that survived the permutation test (Supplementary Sheet 25)

compared with PC3 (Figs. 4B and S4). These significant Behavior-FC regression

coefficients mainly occurred in the connections between default, control, and somatomotor

networks with others (Fig. 4C), such as the connection between default B network with

somatomotor A, somatomotor B, ventral attention A, limbic pole, and default A networks.

Taking the intra-cerebellar FC as an intermediate role, PC3 had a significant correlation

with the GCI$^-$ co-expression (Fig. 4E, F): $r = 0.67$, permutation test $p < 0.001$. In contrast,

no correlation between GCI$^+$ co-expression and PC3 Behavior-FC map: $r = 0.27$, $p > 0.05$.

Based on the loading pattern of PC3 for the original 59 behavior measures, the PC3 reflects

the variation in the emotion-cognitional behavior measures (Fig. 4A). Specifically, PC3

has a large positive loading for the number of correct responses in emotion recognition,

executive function, and fluid ability of cognition tasks, whereas it has a large negative

loading for the response time in emotion recognition and fluid intelligence tasks. It suggests

a high PC3 score might reflect high emotion and cognition ability. This Behavior-FC-Gene

mapping analysis revealed that the GCI$^-$ not only plays an important profiling role in the

functional organization of the cerebellum but also implicates emotion-cognitional behavior

vicariously through the intra-cerebellar FC.

A

Behavior: PC3

D

Gene-set: GCl⁻

B   **E**

R = 0.67

C   **F**

Fig. 4. The Behavior-FC-Gene mapping analysis. A The loading pattern of principal component 3 (PC3) for the original 59 behavior measures (Table S6) was obtained from principal component analysis (PCA). The word sizes represent the absolute values of the loading coefficients, and the different colors indicate the six categories of the behaviors (Table S6). **B** Behavior-FC map for PC3, which are the z-scored regression coefficients of PC3 in each intra-cerebellar functional connection via mass univariate regression procedure. **C** The chord plot shows the significant regression coefficients after the permutation test consisting of 10,000 permutations. The circularly arranged gray histogram shows the absolute magnitudes of the regression coefficients, which correspond to the values of the regression coefficients shown in **B**. The color intensity of chords is the same as the color bar of **B**. **D** The GCI⁻ gene set which used to construct the GCI⁻ co-expression map **E**. **F** The chord plot shows the significant correlations: $p \leq 0.05$ (Bonferroni corrected). The circularly arranged grey histogram shows the absolute magnitudes of the correlation coefficients, which correspond to the values of the correlation coefficients shown in **E**. The color intensity of chords is the same as the color bar of **E**.

GO, pathway, and disorder enrichment analysis

The GO enrichment analysis of the GCI⁺ and GCI⁻ is shown in Fig. 5A. The GCI⁺ was mainly enriched in microtubule-related terms, including the microtubule associated complex (ID: 0005875), motile cilium (ID: 0031514), and dynein complex (ID: 0030286). Compared with GCI⁺, the GCI⁻ was not only enriched in microtubule-related terms but was also significantly enriched in terms related to neurotransmitter transport, such as calcium ion binding (ID: 0005509), regulation of hormone levels (ID: 0010817), response to catecholamine (ID: 0071869), response to monoamine (ID: 0071867), and regulation of neurotransmitter receptor activity (ID: 0099601). These findings are consistent with their different pathway enrichment results (Supplementary Sheets 26 and 27) in that the GCI⁺ was primarily enriched in some basic biological pathways: proximal tubule bicarbonate reclamation (ID: M4361) and glycolysis/gluconeogenesis (ID: M39474), which provides the energy needed during microtube-related processes. In contrast, the GCI⁻ was primarily involved in signaling transduction, especially in some neurotransmission pathways, such as the neuroactive ligand-receptor interaction (ID: M13380).

A Gene ontology enrichment

B Disease enrichment

Fig. 5. The gene ontology (GO) and disease enrichment analysis for GCI⁺ and GCI⁻.
A Bubble plots showing the GO enrichment top 20 terms for GCI⁺ (left) and GCI⁻ (right)
(Complete results are shown in Supplementary Sheets 26 and 27, respectively). The

biological process (BP), cellular component (CC), and molecular function (MF) are
displayed together. The dot size (count) represents the number of genes that are within the
interest GCI$^+$ or GCI$^-$ gene panels as well as a specific GO term (*y*-axis). The different
color intensities indicate the *p* value (FDR corrected). **B** Gradient bar plots showing the
disease enrichment for all representative results for GCI$^+$ and top 15 representative terms
for GCI$^-$. The different color intensities indicate the *p* value (FDR corrected).

Since the GCI$^+$ and GCI$^-$ are involved in different biological processes, we

hypothesized that they also play different roles in brain disease or are related to different

brain diseases. Unexpectedly, we found no link between GCI$^+$ and any brain-related

illnesses (Fig. 5B, left) but observed involvement of GCI$^-$ in various neuropsychiatric

disorders (Fig. 5B, right), including autistic disorder (ID: C0004325), alcoholic

intoxication (ID: C0001973), mental depression (ID: C0011570), pain (ID: C0030193),

learning disorders (ID: C0023186) and others (Supplementary Sheet 27). Many of these,

especially mental depression and autistic disorder, have a close relationship with the human

cerebellum, in which patients have shown FC abnormalities [54, 55]. The mental

depression- and autistic disorder-associated genes were TRH, PENK, TTR, ADCY5,

NRXN1, HTR1A, HTR2C, NTS, PEX5L (*n* = 9, Supplementary Sheet 27), and DLGAP2,

TRH, PENK, RYR3, SEMA3A, NRXN1, TESC, ABCG2, PCDH10, CNTN4, HTR1A,

CALB2, HTR2C, DNAAF4, FOLR1, NTS, GRM8, UPP2 (*n* = 18, Supplementary Sheet

27), respectively, and the overlapping genes were TRH, PENK, NRXN1, HTR1A, HTR2C,

NTS (*n* = 6).

Integrative temporal specificity analysis

In light of the distinct properties of GCI$^+$ and GCI$^-$, we wanted to know whether the roles

played by these two gene sets showed variable prevalence at different ages. By leveraging

the BrainSpan dataset [56] and applying the analysis strategy of the CSEA tool [52], we

found that GCI+ showed significant overexpression in early middle fetal, late middle fetal,

late fetal, neonatal early infancy, and adolescence compared with GCI− (Fig. 6A). These

stages neatly correspond to the timeline of the protracted development of the human

cerebellum [57], which extends from the early embryonic period until the end of the first

postnatal year. This result appears to be consistent with the observation that the GCI+ is

involved in some fundamental biological processes, especially microtubule-related activity,

whose dynamics play a key role in cerebellar neurodevelopment [58]. In contrast,

compared with the GCI+, the GCI− was significantly expressed in early mid fetal, neonatal

early infancy, late infancy, early childhood, adolescence, and young adulthood (Fig. 6B).

These periods include the highest neurodevelopmental risk windows for autism spectrum

disorder (ASD) [59] and major depression disorder (MDD) [60], both of which also existed

in the disease enrichment analysis of GCI−.

Fig. 6. Integrative temporal specificity analysis within the human cerebellum for
GCI$^+$ and GCI$^-$. A Bar plot showing the temporal specificity results for GCI$^+$. The
specificity index probability (pSI = 0.05, 0.01, 0.001, and 0.0001, permutation corrected,
shown as different color intensities) was used to determine how likely a gene was to be
expressed in a given time window relative to all other time windows [52]. The *x*-axis
corresponds to the -$\log_{10}p$ (FDR corrected), and for esthetics if -$\log_{10}p$ (FDR corrected) >
2.5, -$\log_{10}p$ (FDR corrected) = 2.5; the y-axis represents the 10 development windows
collected by BrainSpan [56]. The vertical dark dashed line corresponds to the *p* (FDR
corrected) = 0.05. Complete results are shown in Supplementary Sheet 28. **B** Bar plot
showing the temporal specificity results for GCI$^-$.

Discussion

The current study provided a tentative exploration of the genetic differential and co-

expression linked with the functional organization of the human cerebellum and has the

potential for elaborating and rethinking the neurobiological underpinnings of the cerebellar

functional organization. Furthermore, we identified two gene sets involved in cerebellar

neurodevelopment and neurotransmission and found indirect but interesting genetic

evidence supporting a key role played by the cerebellar functional network in emotion-

cognitional behaviors, as well as many neuropsychiatric disorders. These findings hint at a

micro-macro interacted mechanistic possibility for the cerebellar contributions to emotion-

cognitional behaviors, and consequently to related neuropsychiatric disorders.

The genetic profiles underlying cerebellar functional segregation correlate with intra-

cerebellar and cerebello-cerebral connections

In this study, we found correlations between the identified cerebellar network-specific

genes with the intra-cerebellar connection and cerebello-cerebral FC. These findings could

provide possible empirical genetic support for the hypothesized decisive role of cerebellar

connectivity in the functional heterogeneity of the cerebellum. First, while obtaining the

network-specific genes, we found significant differences in the number of identified genes

between the functional specificity (i.e., limbic, visual networks) and functional diversity

networks (i.e., the control, default networks); specifically, more differentially expressed

genes were in the former and vice versa in the latter [61]. This observation was also found

in a previous cortical gene expression homogeneity analysis [9] that showed that a

relatively high differential expression pattern was observed in the primary sensory cortex,

area 38, and the primary visual cortex. But the findings related to the inconsistency in the

amount of somatomotor cerebellar ($n = 3$) and somatomotor cortical network-specific

genes ($n = 960$) were unclear. One possible explanation may be that the preferential links

between the cerebellar representations of body space and the motor, somatosensory, and

premotor cortices are challenging to distinguish [21]. The finding of cerebellar network-

specific genes is consistent with the elaborate regional difference in cerebellar-cortical

cytoarchitecture [2, 62] in addition to the uniform cell types and their connectivity, such as

regional variation in the cell size and packing density of Purkinje, granule, and Golgi cells.

Second, the overall distribution patterns of the cerebellar and cortical network-specific

genes were highly correlated, a finding in line with a similar macroscale principle identified

in the cerebellar and cortical functional organization [23, 63]. These correlated patterns

may be related to defining the cerebellar networks, which were by projecting the cerebral

cortical networks onto the cerebellum by computing the functional connections between

the two structures [21]. More interestingly, the molecular genetic substrates simultaneously

linking functional heterogeneity and integration could be observed across different

functional subdivisions, regardless of whether the parcellation was based on the task-free

cerebello-cortical FC [21] or the intra-cerebellar task-based activation pattern [20]. These

interpretations are further supported by the widely accepted notion about the human

cerebellum that its functional specialization is dominated more by its connection with

extra-cerebellar structures than within its generally homogeneous cytoarchitecture [8].

Although no intra-cerebellar anatomical fiber connections linking adjacent or distant

cerebellar regions have been observed [64, 65], it is well recognized that the intra-

cerebellar functional map is a consequence of the topological arrangement of its extra-

cerebellar anatomical connections [8]. This fact is especially important to note when

explaining the Behavior-FC results that although our results were obtained through intra-

cerebellar connections, the cerebellar-cortical connections must also be involved actually.

The proposed relationship between extra- and intra-cerebellar connectivity, in turn, can be

expected to affect the resting-state activity between cerebellar regions [23]. Furthermore,

for the interaction between gene and FC, they influence one another bidirectional [66]

through gene-environment (G-E) interplay [67] with the development as an essential

element [68], instead of a unidirectional determination role of the genes in the brain

connection. Here we observed the Gene-FC interaction from the adult perspective (AHBA

donors aged from 24 to 57 years old), which lacks exploration in the G-E interplay

compared with the early development [68].

Third, in addition to the intra-cerebellar Gene-FC correlation, we observed a direct

correlation between genes underlying the cerebellar functional specialization and

cerebello-cerebral FC of the limbic and control networks. The Gene-FC correlation in the

control network was mainly caused by the genetic similarity between these two networks;

this interaction between limbic-emotion and control-cognition has been confirmed

anatomically and behaviorally [69]. For instance, the integrated processing by the emotion

and cognition areas has been identified solely based on their anatomical connections [70].

This relationship can also be observed in that when looking at the top of a hill, a sad mood

induces a steeper perception of the hill than a happy one [71]. One possible reason why we

only obtained this correspondence in the limbic network may be the low functional

heterogeneity [61] and inter-individual functional variability [72] of the limbic network

compared with others as well as the complexity of gene expression; i.e., the Gene-FC

correlation is not fully portrayed by the differentially expressed genes [26]. Considering

the indirect connection between cerebellum and cortex, and large differences between the

cerebellum and cortex in terms of their gene expression patterns [13] and structure-function

relationships [73], as well as the individual variability of their functional networks [75],

the convergently expressed genes associated with the cerebello-cortical cognitive-limbic

networks were evident through simple analysis of co-expression, demonstrating these

genes are very significant and may hold clues to the molecular underpinnings of the

cognitive-emotion roles played by the cerebello-cortical circuit. For example, the HTR1A

and HTR2C, both preferentially expressed in the cerebellar and cortical limbic network,

are pivotal genes in serotonin transmission and play a modulation role in the limbic system

act as important therapeutic targets in limbic system-related disorders [74]. This

observation was further confirmed by the intra-cerebellar Behavior-FC-Gene mapping

analysis, which showed that some of the genes were strongly associated with the emotion-

cognitional behaviors.

Cerebellar neurodevelopment feature of GCI$^+$, cerebellar neurotransmission,

emotion-cognitional behavior, and neuropsychiatric related features of GCI$^-$

Interestingly, we identified two gene subsets with pronouncedly different characteristics

based solely on the direction in which each gene influenced the intra-cerebellar Gene-FC

correlation by applying a simple virtual KO approach on the 443 cerebellar network-

specific genes. By using a series of bioinformatic tools, we found converging evidence for

GCI$^+$ and GCI$^-$ involvement in cerebellar neurodevelopment and cerebellar

neurotransmission, respectively. It is also interesting to speculate that these 443 network-

specific genes that link both cerebellar functional segregation and integration have a

relationship with some brain disorders since prior evidence showed that the cerebellar

functional organization plays a key role in various neurological [30, 75] and psychiatric

disorders [29], most of which possess common underlying genetic risks [76]. But a tricky

problem emerged in that the genes we are interested in were derived from healthy

individuals. It could be tackled to some extent by using the virtual KO method, which can

simulate the different expression levels of each gene and thus coarsely corresponds to a

fraction of the expression level under normal health and disease situations. This is why we

thought that we might be able to see whether the GCI^+ and GCI^- are related to a specific

disease even though the genes were derived from healthy individuals.

The GCI^+ is involved in many microtubule-related terms and is overexpressed

throughout the protracted development of the cerebellum. The dynamics and flexibility of

microtubules were found to be essential throughout cerebellar development via leading the

morphological alterations of Purkinje cells [58]. In addition, some genes of the GCI^+, such

as GTPBP2 [77] and Lin28b [78], were found to play a key role in neurodevelopment;

overexpression of the Lin28b gene can induce the development of pathological lobulation

in the cerebellum [78]. This converging evidence prompts our speculation that the GCI^+ is

engaged in cerebellar neurodevelopment. Unexpectedly, the GCI^+ showed no link to brain-

related diseases, which appears to be consistent with its primary involvement in many

fundamental biological functions. However, this lack of disease linkage is inconsistent with

the significant overexpression of GCI^+ genes during the protracted development of the

cerebellum. Many researchers pointed out that this protracted development increased the

susceptibility of the cerebellum to many psychiatric disorders [57]. It is likely

complemented by the overexpression of GCI^- in the early middle fetal and neonatal early

infancy periods. Other possible explanations include few genetic studies of the cerebellum

compared with the cerebral cortex, and large gene expression differences between the

cerebellum and extra-cerebellar structures [13], so the related datasets may lack sufficient

information specific to the cerebellum. It calls for future studies to provide a complete

explanation by considering multiple perspectives.

The GCI^- was found to be related to emotion-cognitional behaviors, involved in many

neurotransmission processes, enriched in various neurological and psychiatric disorders,

and significantly overexpressed in late infancy, early childhood, adolescence, and young

adulthood compared with GCI^+. These results are mutually supportive. The

neurotransmission has long been believed to play a crucial role in emotional [79] and

cognitional behaviors [80], and its abnormality has been extensively associated with

various neurological [81] and psychiatric disorders [82, 83] with the characteristic of

cognitive and emotional impairments. For example, the abnormal transmission of

monamines and catecholamines, such as serotonin and dopamine, has been widely linked

with many psychiatric disorders, and these transmitters have thus become potential

treatment targets [84]. The period through which the GCI^- genes are expressed includes

the high-risk time windows for GCI^- enriched disorders, such as mental depression (aged

18–29) [60] and autistic disorder (from infancy to childhood) [59]. And the high expression

of GCI^- in early middle fetal life might be associated with the prenatal risk factors

associated with depression [85] and autism [86].

A possible micro-macro interacted mechanistic explanation for the

function/dysfunction of the human cerebellum

The cerebellum regulates motor, emotion, and cognition by functioning as an oscillation

dampener that combines multiple internal representations with external stimuli and

appropriate responses to maintain the behavior around a homeostatic baseline

automatically to optimize performance according to the cortex [87]. Interestingly, we found

that the GCI$^-$ associated with the emotion-cognitional behaviors via the intra-cerebellar FC,

mainly includes the connections between control, default, and somatomotor networks

which have been widely confirmed to be involved in these behaviors via human clinical

data or functional imaging studies [4, 88-90]. This result suggests that the interaction

between genes and FC is also involved in various higher functions of the cerebellum,

providing new evidence and molecular substrates for the progressively complete notion

that the cerebellum modulates cognition and emotion as it does to the motor control [1].

Beyond that, the specifics of how the cerebellum is involved in emotional and cognitive

functions are explained by many compatible theories [87]. Such as the internal model [91],

in which the cerebellum shows increased activations for negative feedback compared to

the positive feedback in the reversal-learning task [92], which is maybe consistent with the

negative Behavior-FC map found in the present study.

Meanwhile, the regions that showed a significant correlation between emotion-

cognitional behaviors and GCI$^-$ via connections, correspond exactly to the cerebellar

posterior lobe whose lesions produce the dysmetria of thought and emotion, i.e., the

cerebellar cognitive affective syndrome, which is characterized by the deficits in executive

function, visual-spatial processing, linguistic skills and affect regulation [93]. And no

significant difference in the performance on newly established objective cognitive tests

battery between patients with isolated cerebellar lesions versus complex cerebello-cortical

connection disorders, such as schizophrenia, mental depression, and autistic disorder [94].

It is thus reasonable and mutually validating that we found that the GCI$^-$ enriched in many

neuropsychiatric disorders involving alternation in these two categories of behaviors,

including mental depression, autistic disorder, pain, alcoholic intoxication, learning

disorder, and others. Moreover, these disorders are all closely related to the alterations of

the cerebellar FC [29, 54, 95-97].

Therefore, the GCI^- provides a possible micro-macro interacted mechanistic

explanation for the functions and dysfunctions of the human cerebellum. The genes

underlying the functional organization of the human cerebellum are likely involved in

emotion-cognitional behaviors through their interaction with the cerebellar connection.

One of the possible ways the risk genes act in the pathogenesis of corresponding diseases

maybe through their interactions with the cerebellar FC, which results in dysregulation of

cerebellar FC and thus pathologically manifested as its functional connection abnormalities

and consequently transient or long-term impairment at the cognitive and emotional

behavioral levels of these diseases. For instance, the fluctuation in the correspondence of

the Gene-FC relationship found in the present study and widespread altered cerebellar FC

excavated previously in diverse neuropsychiatric disorders [29]. The GCI^- also provides a

promising genetic resource for investigating the cerebellar involvements in emotion-

cognitional behaviors and related brain diseases. For example, the observed overlapping

genes, i.e., NRXN1, are associated with mental depression and autistic disorder supported

previous clinical studies showing that rare and common variants in NRXN1 carried risks

for MDD [98], ASD, and schizophrenia [99]. And HTR1A, which has a high expression in

the cerebellum [14] and has been linked to the cognitive process in both animal models and

human [79], was found to be involved in pain, mental depression, autistic disorder,

alcoholic intoxication, learning disorder, and other conditions (Supplementary Sheet 27).

Methodological considerations

The interpretation of our findings is not without limitations. First, we validated our results

across independent imaging datasets that cover the age range of AHBA, and various

cerebellar atlases with different parcellation criteria and resolutions. But we are unable to

test our results in an independent gene dataset as there is no other publicly available dataset

with a detailed sampling of subregions of the human cerebellum. Second, the gene co-

expression we constructed only considered one small part of the relationship between the

genes and FC. Thereby it did not fully recapitulate the complexity of the brain

transcriptome, such as gene-gene interactions [100]. That is one possible reason we only

found a cerebello-cortical Gene-FC correlation for the cognitive-limbic networks. Third,

the simple correlation approach [25, 61] used in this study and other linear regression

models like partial least squares analysis [101], can only prioritize genes for further

investigation and cannot fully explore the causal relationship between genes and functional

organization. As a result, further exploration is hindered by the intricacies of genetic and

epigenetic regulation. It makes the discussion and explanation of the different directions of

this correlation challenging. For example, why the direction of influence on the Gene-FC

correlation could separate these 443 genes into two distinct gene sets with different

functions remains unclear. Hence further related exploration is necessary but very

challenging. Fourth, since the 443 genes were not derived based on the correlation with a

spatially-defined phenotype, the newly proposed strategy [102] that mitigates the bias of

leveraging gene enrichment approach in the spatial transcriptomic data cannot be directly

utilized. Nonetheless, it cannot be determined whether this type of bias affects the present

results, which calls for future efforts to develop methods to answer this question.

Briefly, more donors, the overall pattern of gene expression, gene regulation,

epigenomics, and improved cellular resolution are needed and imperative for developing

more appropriate and ingenious approaches to understand the bidirectional relationship

between genes and functional organization, which is a greater challenge for neuroscience

than just identifying a link between genetic and imaging data. Nevertheless, in light of the

currently limited understanding of how microscale genes contribute to macroscale brain

functional organization, the prioritization of genes and the related functional annotations

presented here are still necessary [24, 26].

Conclusions

In summary, we found that the network-specific genes underlying cerebellar functional

heterogeneity correlated with the intra-cerebellar and cerebello-cerebral FC. It indicates

that the genetic infrastructure associated with functional segregation coalesces to form a

collective system, which closely relates to the functional integration of these functional

subregions. The current study has thus unveiled part of the neurobiological genetic

substrates underlying the cerebellar functional organization. We also identified important

indirect genetic markers that support the key role played by the cerebellar functional

network in emotion-cognitional behaviors and many brain disorders. These findings hint

at the possibility of establishing a cerebellar "gene—connection—function/dysfunction"

chain, as well as of helping to bridge the knowledge gap between the genetic mechanisms

driving the cerebellar functional organization and the predictive markers of behaviors and

heritable risks of disorders, especially major depression and autistic disorder.

Acknowledgements

This work was partially supported by the Science and Technology Innovation 2030—Brain Science and Brain-Inspired Intelligence Project (Grant No. 2021ZD0200203), the Natural Science Foundation of China (Grant Nos. 82072099 and 91432302), the Strategic Priority Research Program of the Chinese Academy of Sciences (XDB32030214), the Youth Innovation Promotion Association, the Beijing Advanced Discipline Fund and the Major Scientific Project of Zhejiang Lab (Grant No. 2021ND0PI01). Data were provided by the Human Connectome Project, WU-Minn Consortium (Principal Investigators: David Van Essen, and Kamil Ugurbil; 1U54MH091657) funded by the 16 NIH Institutes and Centers that support the NIH Blueprint for Neuroscience Research; and by the McDonnell Center for Systems Neuroscience at Washington University. The authors appreciate the English language and editing assistance of Rhoda E. and Edmund F. Perozzi, PhDs.

Author Contributions

YW and LF designed the research; YW and LC performed the experiments; YW, LF, CC, TJ, BL, and ZY contributed new analytic tools; YW, LC, DL, XW and CG analyzed data; YW and LF wrote the paper; and LF, CC, KHM, JRN, YZ, ZY, JX, SBE, BL, and TJ contributed analytic expertise, theoretical guidance, paper revisions, and informed interpretation of the results; and LF and CC supervised the project.

Conflict of Interest

All authors report no biomedical financial interests or potential conflicts of interest.

References

1. Schmahmann JD, Guell X, Stoodley CJ, Halko MA. The Theory and Neuroscience of Cerebellar Cognition. *Annu Rev Neurosci* 2019; **42:** 337-364.

2. De Zeeuw CI, Lisberger SG, Raymond JL. Diversity and dynamism in the cerebellum. *Nat Neurosci* 2021; **24**(2)**:** 160-167.

3. Strick PL, Dum RP, Fiez JA. Cerebellum and nonmotor function. *Annu Rev Neurosci* 2009; **32:** 413-434.

4. Buckner RL. The cerebellum and cognitive function: 25 years of insight from anatomy and neuroimaging. *Neuron* 2013; **80**(3)**:** 807-815.

5. Sathyanesan A, Zhou J, Scafidi J, Heck DH, Sillitoe RV, Gallo V. Emerging connections between cerebellar development, behaviour and complex brain disorders. *Nat Rev Neurosci* 2019; **20**(5)**:** 298-313.

6. Guell X, Schmahmann JD, Gabrieli JDE. Functional Specialization is Independent of Microstructural Variation in Cerebellum but Not in Cerebral Cortex. *bioRxiv* 2018**:** 424176.

7. Voogd J, Glickstein M. The anatomy of the cerebellum. *Trends Cognit Sci* 1998; **2**(9)**:** 307-313.

8. Schmahmann JD. The role of the cerebellum in affect and psychosis. *J Neurolinguistics* 2000; **13**(2)**:** 189-214.

9. Hawrylycz MJ, Lein ES, Guillozet-Bongaarts AL, Shen EH, Ng L, Miller JA *et al.* An anatomically comprehensive atlas of the adult human brain transcriptome. *Nature* 2012; **489**(7416)**:** 391-399.

10. Johnson MB, Kawasawa YI, Mason CE, Krsnik Z, Coppola G, Bogdanovic D *et al.* Functional and evolutionary insights into human brain development through global transcriptome analysis. *Neuron* 2009; **62**(4)**:** 494-509.

11. Guevara EE, Hopkins WD, Hof PR, Ely JJ, Bradley BJ, Sherwood CC.
Comparative analysis reveals distinctive epigenetic features of the human
cerebellum. *PLoS Genet* 2021; **17**(5)**:** e1009506.

12. Wang VY, Zoghbi HY. Genetic regulation of cerebellar development. *Nat Rev*
Neurosci 2001; **2**(7)**:** 484-491.

13. Hawrylycz M, Miller JA, Menon V, Feng D, Dolbeare T, Guillozet-Bongaarts AL
et al. Canonical genetic signatures of the adult human brain. *Nat Neurosci* 2015;
18(12)**:** 1832-1844.

14. Negi SK, Guda C. Global gene expression profiling of healthy human brain and its
application in studying neurological disorders. *Sci Rep* 2017; **7**(1)**:** 897.

15. Aldinger KA, Thomson Z, Phelps IG, Haldipur P, Deng M, Timms AE *et al.* Spatial
and cell type transcriptional landscape of human cerebellar development. *Nat*
Neurosci 2021; **24**(8)**:** 1163-1175.

16. Zeng T, Chen H, Fakhry A, Hu X, Liu T, Ji S. Allen mouse brain atlases reveal
different neural connection and gene expression patterns in cerebellum gyri and
sulci. *Brain Struct Funct* 2015; **220**(5)**:** 2691-2703.

17. Kozareva V, Martin C, Osorno T, Rudolph S, Guo C, Vanderburg C *et al.* A
transcriptomic atlas of the mouse cerebellum reveals regional specializations and
novel cell types. *bioRxiv* 2020**:** 2020.2003.2004.976407.

18. Ren Y, Guo L, Guo CC. A connectivity-based parcellation improved functional
representation of the human cerebellum. *Sci Rep* 2019; **9**(1)**:** 9115.

19. Bernard JA, Seidler RD, Hassevoort KM, Benson BL, Welsh RC, Wiggins JL *et al.*
Resting state cortico-cerebellar functional connectivity networks: a comparison of
anatomical and self-organizing map approaches. *Front Neuroanat* 2012; **6:** 31.

20.   King M, Hernandez-Castillo CR, Poldrack RA, Ivry RB, Diedrichsen J. Functional
boundaries in the human cerebellum revealed by a multi-domain task battery. *Nat*
Neurosci 2019; **22**(8)**:** 1371-1378.

21.   Buckner RL, Krienen FM, Castellanos A, Diaz JC, Yeo BT. The organization of
the human cerebellum estimated by intrinsic functional connectivity. *J*
Neurophysiol 2011; **106**(5)**:** 2322-2345.

22.   Ji JL, Spronk M, Kulkarni K, Repovs G, Anticevic A, Cole MW. Mapping the
human brain's cortical-subcortical functional network organization. *Neuroimage*
2019; **185:** 35-57.

23.   Guell X, Schmahmann JD, Gabrieli JDE, Ghosh SS. Functional gradients of the
cerebellum. *eLife* 2018; **7:** e36652.

24.   Fornito A, Arnatkeviciute A, Fulcher BD. Bridging the Gap between Connectome
and Transcriptome. *Trends Cogn Sci* 2019; **23**(1)**:** 34-50.

25.   Richiardi J, Altmann A, Milazzo AC, Chang C, Chakravarty MM, Banaschewski
T *et al.* BRAIN NETWORKS. Correlated gene expression supports synchronous
activity in brain networks. *Science* 2015; **348**(6240)**:** 1241-1244.

26.   Anderson KM, Krienen FM, Choi EY, Reinen JM, Yeo BTT, Holmes AJ. Gene
expression links functional networks across cortex and striatum. *Nat Commun* 2018;
9(1)**:** 1428.

27.   Ji JL, Helmer M, Fonteneau C, Burt JB, Tamayo Z, Demšar J *et al.* Mapping brain-
behavior space relationships along the psychosis spectrum. *eLife* 2021; **10:** e66968.

28.   van den Heuvel MP, Scholtens LH, Kahn RS. Multiscale Neuroscience of
Psychiatric Disorders. *Biol Psychiatry* 2019; **86**(7)**:** 512-522.

29.   Phillips JR, Hewedi DH, Eissa AM, Moustafa AA. The cerebellum and psychiatric
disorders. *Front Public Health* 2015; **3:** 66.

30.   Rajendran AG, Nutakki C, Sasidharakurup H, Nair eBB, Diwakar S. Cerebellum
in Neurological Disorders: A Review on the Role of Inter-Connected Neural
Circuits. *J neurol stroke* 2017; **6**(2)**:** 00196.

31.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W *et al.* limma powers
differential expression analyses for RNA-sequencing and microarray studies.
Nucleic Acids Res 2015; **43**(7)**:** e47.

32.   Arnatkeviciute A, Fulcher BD, Fornito A. A practical guide to linking brain-wide
gene expression and neuroimaging data. *Neuroimage* 2019; **189:** 353-367.

33.   Miller JA, Cai C, Langfelder P, Geschwind DH, Kurian SM, Salomon DR *et al.*
Strategies for aggregating gene expression data: the collapseRows R function. *BMC*
Bioinformatics 2011; **12:** 322.

34.   Yeo BT, Krienen FM, Sepulcre J, Sabuncu MR, Lashkari D, Hollinshead M *et al.*
The organization of the human cerebral cortex estimated by intrinsic functional
connectivity. *J Neurophysiol* 2011; **106**(3)**:** 1125-1165.

35.   Cox RW. AFNI: software for analysis and visualization of functional magnetic
resonance neuroimages. *Comput Biomed Res* 1996; **29**(3)**:** 162-173.

36.   Smyth GK, Michaud J, Scott HS. Use of within-array replicate spots for assessing
differential expression in microarray experiments. *Bioinformatics* 2005; **21**(9)**:**
2067-2075.

37.   Dalman MR, Deeter A, Nimishakavi G, Duan Z-H. Fold change and p-value cutoffs
significantly alter microarray interpretations. *BMC Bioinformatics* 2012; **13**(2)**:**
S11.

38.   Diedrichsen J, Balsters JH, Flavell J, Cussans E, Ramnani N. A probabilistic MR
atlas of the human cerebellum. *Neuroimage* 2009; **46**(1)**:** 39-46.

39. Smith SM, Beckmann CF, Andersson J, Auerbach EJ, Bijsterbosch J, Douaud G *et*
al. Resting-state fMRI in the Human Connectome Project. *Neuroimage* 2013; **80:**
144-168.

40. Glasser MF, Sotiropoulos SN, Wilson JA, Coalson TS, Fischl B, Andersson JL *et*
al. The minimal preprocessing pipelines for the Human Connectome Project.
Neuroimage 2013; **80:** 105-124.

41. Van Essen DC, Ugurbil K, Auerbach E, Barch D, Behrens TE, Bucholz R *et al.*
The Human Connectome Project: a data acquisition perspective. *Neuroimage* 2012;
62(4)**:** 2222-2231.

42. Salimi-Khorshidi G, Douaud G, Beckmann CF, Glasser MF, Griffanti L, Smith SM.
Automatic denoising of functional MRI data: combining independent component
analysis and hierarchical fusion of classifiers. *Neuroimage* 2014; **90:** 449-468.

43. Griffanti L, Salimi-Khorshidi G, Beckmann CF, Auerbach EJ, Douaud G, Sexton
CE *et al.* ICA-based artefact removal and accelerated fMRI acquisition for
improved resting state network imaging. *Neuroimage* 2014; **95:** 232-247.

44. Burt JB, Helmer M, Shinn M, Anticevic A, Murray JD. Generative modeling of
brain maps with spatial autocorrelation. *Neuroimage* 2020; **220:** 117038.

45. Bookheimer SY, Salat DH, Terpstra M, Ances BM, Barch DM, Buckner RL *et al.*
The Lifespan Human Connectome Project in Aging: An overview. *Neuroimage*
2019; **185:** 335-348.

46. Seidlitz J, Vasa F, Shinn M, Romero-Garcia R, Whitaker KJ, Vertes PE *et al.*
Morphometric Similarity Networks Detect Microscale Cortical Organization and
Predict Inter-Individual Cognitive Variation. *Neuron* 2018; **97**(1)**:** 231-247 e237.

47. Kong X-Z, Tzourio-Mazoyer N, Joliot M, Fedorenko E, Liu J, Fisher SE *et al.* Gene
Expression Correlates of the Cortical Network Underlying Sentence Processing.
Neurobiology of Language 2020; **1**(1)**:** 77-103.

48.   Tian Y, Zalesky A. Characterizing the functional connectivity diversity of the
insula cortex: Subregions, diversity curves and behavior. *Neuroimage* 2018; **183:**
716-733.

49.   Winkler AM, Ridgway GR, Webster MA, Smith SM, Nichols TE. Permutation
inference for the general linear model. *Neuroimage* 2014; **92:** 381-397.

50.   Chen J, Bardes EE, Aronow BJ, Jegga AG. ToppGene Suite for gene list
enrichment analysis and candidate gene prioritization. *Nucleic Acids Res* 2009;
37(Web Server issue)**:** W305-311.

51.   Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM *et al.* Gene
ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat*
Genet 2000; **25**(1)**:** 25-29.

52.   Dougherty JD, Schmidt EF, Nakajima M, Heintz N. Analytical approaches to RNA
profiling data for the identification of genes enriched in specific cells. *Nucleic Acids*
Res 2010; **38**(13)**:** 4218-4230.

53.   Huntley RP, Sawford T, Mutowo-Meullenet P, Shypitsyna A, Bonilla C, Martin
MJ *et al.* The GOA database: gene Ontology annotation updates for 2015. *Nucleic*
Acids Res 2015; **43**(Database issue)**:** D1057-1063.

54.   Zhu DM, Yang Y, Zhang Y, Wang C, Wang Y, Zhang C *et al.* Cerebellar-cerebral
dynamic functional connectivity alterations in major depressive disorder. *J Affect*
Disord 2020; **275:** 319-328.

55.   Verly M, Verhoeven J, Zink I, Mantini D, Peeters R, Deprez S *et al.* Altered
functional connectivity of the language network in ASD: role of classical language
areas and cerebellum. *Neuroimage Clin* 2014; **4:** 374-382.

56.   Miller JA, Ding SL, Sunkin SM, Smith KA, Ng L, Szafer A *et al.* Transcriptional
landscape of the prenatal human brain. *Nature* 2014; **508**(7495)**:** 199-206.

57.   ten Donkelaar HJ, Lammens M, Wesseling P, Thijssen HO, Renier WO.
1062   Development and developmental disorders of the human cerebellum. *J Neurol* 2003;
1063   **250**(9)**:** 1025-1036.

1064   58.   Munoz-Castaneda R, Diaz D, Peris L, Andrieux A, Bosc C, Munoz-Castaneda JM
1065   *et al.* Cytoskeleton stability is essential for the integrity of the cerebellum and its
1066   motor- and affective-related behaviors. *Sci Rep* 2018; **8**(1)**:** 3072.

1067   59.   Beyreli I, Karakahya O, Cicek AE. Deep multitask learning of gene risk for
1068   comorbid neurodevelopmental disorders. *bioRxiv* 2020**:** 2020.2006.2013.150201.

1069   60.   Villarroel MA, Terlizzi EP. Symptoms of Depression Among Adults: United States,
1070   2019. *NCHS Data Brief* 2020; (379)**:** 1-8.

1071   61.   Anderson ML, Kinnison J, Pessoa L. Describing functional diversity of brain
1072   regions and brain networks. *Neuroimage* 2013; **73:** 50-58.

1073   62.   Cerminara NL, Lang EJ, Sillitoe RV, Apps R. Redefining the cerebellar cortex as
1074   an assembly of non-uniform Purkinje cell microcircuits. *Nat Rev Neurosci* 2015;
1075   **16**(2)**:** 79-93.

1076   63.   Margulies DS, Ghosh SS, Goulas A, Falkiewicz M, Huntenburg JM, Langs G *et al.*
1077   Situating the default-mode network along a principal gradient of macroscale
1078   cortical organization. *Proc Natl Acad Sci U S A* 2016; **113**(44)**:** 12574-12579.

1079   64.   Schmahmann JD, Pandya DN. Disconnection syndromes of basal ganglia, thalamus,
1080   and cerebrocerebellar systems. *Cortex* 2008; **44**(8)**:** 1037-1066.

1081   65.   Schmahmann JD. From movement to thought: anatomic substrates of the cerebellar
1082   contribution to cognitive processing. *Hum Brain Mapp* 1996; **4**(3)**:** 174-198.

1083   66.   van den Heuvel MP, Yeo BTT. A Spotlight on Bridging Microscale and Macroscale
1084   Human Brain Architecture. *Neuron* 2017; **93**(6)**:** 1248-1251.

67. Krubitzer L, Kaas J. The evolution of the neocortex in mammals: how is phenotypic
diversity generated? *Curr Opin Neurobiol* 2005; **15**(4)**:** 444-453.

68. Boyce WT, Sokolowski MB, Robinson GE. Genes and environments, development
and time. *Proc Natl Acad Sci U S A* 2020; **117**(38)**:** 23235-23241.

69. Storbeck J, Clore GL. On the interdependence of cognition and emotion. *Cogn
Emot* 2007; **21**(6)**:** 1212-1237.

70. Ghashghaei HT, Barbas H. Pathways for emotion: interactions of prefrontal and
anterior temporal pathways in the amygdala of the rhesus monkey. *Neuroscience*
2002; **115**(4)**:** 1261-1279.

71. Riener CR, Stefanucci JK, Proffitt DR, Clore G. An effect of mood on the
perception of geographical slant. *Cogn Emot* 2011; **25**(1)**:** 174-182.

72. Mueller S, Wang D, Fox MD, Yeo BT, Sepulcre J, Sabuncu MR *et al.* Individual
variability in functional connectivity architecture of the human brain. *Neuron* 2013;
77(3)**:** 586-595.

73. Guell X, Schmahmann JD, Gabrieli JD. Functional Specialization is Independent
of Microstructural Variation in Cerebellum but Not in Cerebral Cortex. *bioRxiv*
2018**:** 424176.

74. Hensler JG. Serotonergic modulation of the limbic system. *Neurosci Biobehav Rev*
2006; **30**(2)**:** 203-214.

75. Schmahmann JD. Disorders of the cerebellum: ataxia, dysmetria of thought, and
the cerebellar cognitive affective syndrome. *J Neuropsychiatry Clin Neurosci* 2004;
16(3)**:** 367-378.

76. Anttila V, Bulik-Sullivan B, Finucane HK, Walters RK, Bras J, Duncan L *et al.*
Analysis of shared heritability in common disorders of the brain. *Science* 2018;
360(6395)**:** eaap8757.

77.   Bertoli-Avella AM, Garcia-Aznar JM, Brandau O, Al-Hakami F, Yuksel Z, Marais
A *et al.* Biallelic inactivating variants in the GTPBP2 gene cause a
neurodevelopmental disorder with severe intellectual disability. *Eur J Hum Genet*
2018; **26**(4)**:** 592-598.

78.   Wefers AK, Lindner S, Schulte JH, Schuller U. Overexpression of Lin28b in Neural
Stem Cells is Insufficient for Brain Tumor Formation, but Induces Pathological
Lobulation of the Developing Cerebellum. *Cerebellum* 2017; **16**(1)**:** 122-131.

79.   Wang F, Yang J, Pan F, Ho RC, Huang JH. Editorial: Neurotransmitters and
Emotions. *Front Psychol* 2020; **11: 21.**

80.   Svob Strac D, Pivac N, Muck-Seler D. The serotonergic system and cognitive
function. *Transl Neurosci* 2016; **7**(1)**:** 35-49.

81.   Swoboda KJ, Hyland K. Diagnosis and treatment of neurotransmitter-related
disorders. *Neurol Clin* 2002; **20**(4)**:** 1143-1161, viii.

82.   Brown RP, Mann JJ. A clinical perspective on the role of neurotransmitters in
mental disorders. *Hosp Community Psychiatry* 1985; **36**(2)**:** 141-150.

83.   Kato TA, Yamauchi Y, Horikawa H, Monji A, Mizoguchi Y, Seki Y *et al.*
Neurotransmitters, psychotropic drugs and microglia: clinical implications for
psychiatry. *Curr Med Chem* 2013; **20**(3)**:** 331-344.

84.   Seo D, Patrick CJ, Kennealy PJ. Role of Serotonin and Dopamine System
Interactions in the Neurobiology of Impulsive Aggression and its Comorbidity with
other Clinical Disorders. *Aggress Violent Behav* 2008; **13**(5)**:** 383-395.

85.   Su Y, D'Arcy C, Meng X. Research Review: Developmental origins of depression
– a systematic review and meta-analysis. *J Child Psychol Psychiatry* 2021; **62**(9)**:**
1050-1066.

86.   Newschaffer CJ, Fallin D, Lee NL. Heritable and nonheritable risk factors for
autism spectrum disorders. *Epidemiol Rev* 2002; **24**(2)**:** 137-153.

87. Guell X, Gabrieli JDE, Schmahmann JD. Embodied cognition and the cerebellum:
Perspectives from the Dysmetria of Thought and the Universal Cerebellar
Transform theories. *Cortex* 2018; **100:** 140-148.

88. Habas C. Functional Connectivity of the Cognitive Cerebellum. *Front Syst*
Neurosci 2021; **15:** 642225.

89. Stoodley CJ, MacMore JP, Makris N, Sherman JC, Schmahmann JD. Location of
lesion determines motor vs. cognitive consequences in patients with cerebellar
stroke. *Neuroimage Clin* 2016; **12:** 765-775.

90. Saarimaki H, Ejtehadian LF, Glerean E, Jaaskelainen IP, Vuilleumier P, Sams M
et al. Distributed affective space represents multiple emotion categories across the
human brain. *Soc Cogn Affect Neurosci* 2018; **13**(5)**:** 471-482.

91. Ito M. Control of mental activities by internal models in the cerebellum. *Nat Rev*
Neurosci 2008; **9**(4)**:** 304-313.

92. Peterburs J, Hofmann D, Becker MPI, Nitsch AM, Miltner WHR, Straube T. The
role of the cerebellum for feedback processing and behavioral switching in a
reversal-learning task. *Brain Cogn* 2018; **125:** 142-148.

93. Schmahmann JD. The role of the cerebellum in cognition and emotion: personal
reflections since 1982 on the dysmetria of thought hypothesis, and its historical
evolution from theory to therapy. *Neuropsychol Rev* 2010; **20**(3)**:** 236-260.

94. Hoche F, Guell X, Vangel MG, Sherman JC, Schmahmann JD. The cerebellar
cognitive affective/Schmahmann syndrome scale. *Brain* 2018; **141**(1)**:** 248-270.

95. Ramos TC, Balardin JB, Sato JR, Fujita A. Abnormal Cortico-Cerebellar
Functional Connectivity in Autism Spectrum Disorder. *Front Syst Neurosci* 2018;
12: 74.

96. Baker KG, Harding AJ, Halliday GM, Kril JJ, Harper CG. Neuronal loss in
functional zones of the cerebellum of chronic alcoholics with and without
Wernicke's encephalopathy. *Neuroscience* 1999; **91**(2)**:** 429-438.

97. Abdallah M, Zahr NM, Saranathan M, Honnorat N, Farrugia N, Pfefferbaum A *et*
al. Altered Cerebro-Cerebellar Dynamic Functional Connectivity in Alcohol Use
Disorder: a Resting-State fMRI Study. *The Cerebellum* 2021; **20**(6)**:** 823-835.

98. Freire-Cobo C, Wang J. Dietary phytochemicals modulate experience-dependent
changes in Neurexin gene expression and alternative splicing in mice after chronic
variable stress exposure. *Eur J Pharmacol* 2020; **883:** 173362.

99. Hu Z, Xiao X, Zhang Z, Li M. Genetic insights and neurobiological implications
from NRXN1 in neuropsychiatric disorders. *Mol Psychiatry* 2019; **24**(10)**:** 1400-
1414.

100. Hua J, Yang Z, Jiang T, Yu S. Pairwise interactions in gene expression determine
a hierarchical transcriptional profile in the human brain. *Sci Bull* 2021; **66**(14)**:**
1437-1447.

101. Ball G, Seidlitz J, Beare R, Seal ML. Cortical remodelling in childhood is
associated with genes enriched for neurodevelopmental disorders. *Neuroimage*
2020; **215:** 116803.

102. Fulcher BD, Arnatkeviciute A, Fornito A. Overcoming false-positive gene-
category enrichment in the analysis of spatially resolved transcriptomic brain atlas
data. *Nat Commun* 2021; **12**(1)**:** 2669.

1181