Intracranial Human Recordings Reveal Intensity Coding for the Pain of Others in the Insula

Efe Soyman[1,2,†], Rune Bruls[1,†], Kalliopi Ioumpa[1,†], Laura Müller-Pinzler[3], Selene Gallo[1,4], Elisabeth C.W. van Straaten[5], Matthew W. Self[6], Judith C. Peters[6,7,8], Jessy K. Possel[6], Yoshiyuki Onuki[9], Johannes C. Baayen[10], Sander Idema[10], Christian Keysers[1,11,‡], Valeria Gazzola[1,11,‡]

[†,‡] **These authors contributed equally to this work.**

Abstract

Based on neuroimaging data, the insula is considered important for people to empathize with the pain of others, whether that pain is perceived through facial expressions or the sight of limbs in painful situations. Here we present the first report of intracranial electroencephalographic (iEEG) recordings from the insulae collected while 7 presurgical epilepsy patients rated the intensity of a woman's painful experiences viewed in movies. In two separate conditions, pain was deduced from seeing facial expressions or a hand being slapped by a belt. We found that broadband activity in the 20-190 Hz range correlated with the trial-by-trial perceived intensity in the insula for both types of stimuli. Using microwires at the tip of a selection of the electrodes, we additionally isolated 8 insular neurons with spiking that correlated with perceived intensity. Within the insula, we found a patchwork of locations with differing selectivities within our stimulus set, some representing intensity only for facial expressions, others only for the hand being hit, and others for both. That we found some locations with intensity coding only for faces, and others only for hand across simultaneously recorded locations suggests that insular activity while witnessing the pain of others cannot be entirely reduced to a univariate salience representation. Psychophysics and the temporal properties of our signals indicate that the timing of responses encoding intensity for the sight of the hand being hit are best explained by kinematic information; the timing of those encoding intensity for the facial expressions are best explained by shape information in the face. In particular, the furrowing of the eyebrows and the narrowing of the eyes of the protagonist in the movies suffice to predict both the rating of and the timing of the neuronal response to the facial expressions. Comparing the broadband activity in the iEEG signal with spiking activity and an fMRI experiment with similar stimuli revealed a consistent spatial organization for the representation of intensity from our hand stimuli, with stronger intensity representation more anteriorly and around neurons with intensity coding. In contrast, for the facial expressions, we found that the activity at the three levels of measurement do not coincide, suggesting a more disorganized representation. Together, our intracranial recordings indicate that the insula encodes, in a partially intermixed layout, both static and dynamic cues from different body parts that reflect the intensity of pain experienced by others.

Author affiliations:

Social Brain Lab, Netherlands Institute for Neuroscience, Royal Netherlands Academy of Art and Sciences, 1105 BA, Amsterdam, the Netherlands

Social Cognitive and Affective Neuroscience Lab, Kadir Has University, Cibali 34083, Fatih, Istanbul, Turkey

Social Neuroscience Lab, Department of Psychiatry and Psychotherapy, University of Lübeck, 23562, Lübeck, Germany

Current Address: Department of Psychiatry, Amsterdam University Medical Center, 1105 AZ, Amsterdam, the Netherlands

Department of Neurology and Clinical Neurophysiology, Amsterdam UMC, Vrije Universiteit Amsterdam, 1117 HV, Amsterdam, the Netherlands

Department of Vision and Cognition, Netherlands Institute of Neuroscience, Royal Netherlands Academy of Art and Sciences, 1105 BA, Amsterdam, the Netherlands

Department of Cognitive Neuroscience, Faculty of Psychology and Neuroscience, Maastricht University, 6229 EV, Maastricht, the Netherlands

Maastricht Brain Imaging Center (M-BIC), Maastricht University, 6229 EV, Maastricht, the Netherlands

Department of Neurosurgery, Jichi Medical University, 329-0498, Tochigi, Japan

Department of Neurosurgery, location VUmc, Amsterdam University Medical Center, 1117 HV, Amsterdam, The Netherlands

Brain and Cognition, Department of Psychology, University of Amsterdam, 1018 WS, Amsterdam, The Netherlands

Correspondence to: Valeria Gazzola or Christian Keysers

Full address: Netherlands Institute for Neuroscience, Meibergdreef 47, 1105 BA Amsterdam, NL

E-mail: v.gazzola@nin.knaw.nl or c.keysers@nin.knaw.nl

Running title: Perceived intensity coding in the insula

Keywords: Insula; Pain; Empathy; intracranial EEG; Broadband Gamma

Abbreviations: BA44/45=Brodmann Area 44/45; BBP=Broad-Band Power 20-190 Hz; BF_{10}=Bayes Factor in favour of a two-tailed alternative hypothesis; BF_{01}=Bayes Factor in favour of the null hypothesis relative to the two-tailed alternative hypothesis; BF_{+0}=Bayes Factor in favour of a one-tailed alternative hypothesis; Fp1=Frontal pole region 1; iEEG=intracranial EEG; MNI=Montreal Neurological Institute coordinate system; OP8/9=Frontal Opercular regions 8 and 9; p_1=one-tailed p-value; p_2=two-tailed p-value; r_P=Pearson's correlation coefficient; r_S=Spearman's rank correlation coefficient; r_K=Kendall's Tau correlation coefficient.

Introduction

Sharing the distress of others is central to empathy. fMRI shows a number of brain regions involved in the experience of pain also increase their activity while participants perceive the pain of others, including the cingulate cortex, the insula, and the somatosensory cortices.[1–4] Across humans, primates, and rodents, lesions in these regions impair the perception and the sharing of others' emotions.[5] Directly recording electrical signals from these regions in humans would complement the more indirect fMRI measurements and sharpen our understanding of how these regions represent the intensity of other people's pain, for at least two reasons. First, fMRI records a mixed signal that includes synaptic input and local neural processing. Localizing BOLD activity that encodes a particular task or stimulus property in a particular brain region thus cannot ensure that neurons in that region actually have spiking activity that encodes that property.[6] For instance, BOLD signals in V1 fluctuate based on whether a stimulus is perceived or not in binocular rivalry.[6,7] In contrast, simultaneous electrical recordings show that broadband gamma activity, which is tightly coupled to spiking[8–10] in the same region responds to a stimulus equally well whether it is perceived or suppressed - only the slower components, <20Hz, that are known to carry feedback synaptic input, fluctuate with perception.[7] Being able to record electrical activity, particularly in the broadband gamma range, is thus critical to localize where in this circuitry neuronal spiking indeed represents the pain of others. Second, fMRI's low temporal resolution makes it difficult to characterize the time-course of responses.

For the anterior cingulate we have intracranial recordings: Hutchison[11] documented a single neuron in epileptic patients that responded to the sight of a finger being pin-pricked with increased firing rate, and a recent rodent study revealed that cingulate neurons responding to pain experience have responses that increase with the intensity of the pain experienced by another rat.[12] In contrast, although the insula is central in the neuroimaging literature on empathy, and shows increases of BOLD signal for watching painful compared to non-painful social stimuli,[1,3,4,13–16] we still lack such intracranial recordings while individuals witness the pain of others. Intracortical EEG (iEEG) has been recorded in the insula during the self-experience of pain,[12] and the insula and adjacent SII are the only cortical regions where iEEG

electrode stimulation can induce painful sensations.[18,19] However, to our knowledge, there are no published studies recording from insular electrodes while patients witness the pain of others. The degree to which neuronal activity local to the insula, as opposed to feedback synaptic input from other regions such as the cingulate gyrus, encodes the intensity of other people's pain therefore remains unclear and the time-course of such neural activity remains undercharacterized.

To fill this gap and characterize the insula's electrophysiological responses to the pain of others, we recorded iEEG in 7 epileptic patients during pre-surgical exploration, while participants rated the pain they perceived another person in a video to experience (Fig. 1). All these patients had macro-contacts in their insulae that yielded local field potentials (LFP, Fig. 1c) capable of measuring broad-band gamma activity. Three, additionally, had micro-wires at the tip of some electrodes to record from isolated insular neurons. Based on fMRI studies showing increased activity in the insula for more painful stimuli, we expected increases in power in higher LFP frequencies and spike counts for more intense stimuli, and used one-tailed testing unless specified otherwise. Our stimuli included two ways in which pain is perceived in others (Fig. 1a). Half the stimuli (Face videos), showed a female receiving electroshocks on the hand and expressing pain through facial expressions (furrowing eyebrows and tightening eyes). The other half (Hand videos), showed the protagonist's hand slapped by a leather belt, and pain intensity had to be deduced from the movements of the belt and hand. In both cases, movies, rather than static images, were chosen to provide richer and more ecologic stimuli and provide information about the temporal dynamics with which such movies are represented in a field still dominated by the presentation of static images.[20,21] We used these two classes of stimuli, because both tap into the visual perception of other people's pain, and we start to understand that they do so through dissociable routes.[2–4] For instance, the hand stimuli depend on the hand region of the somatosensory cortex,[22] while facial expressions are depend on the ventral somatosensory cortex and the insula.[23–25]

The insula is thought to contribute to many aspects of our mental life, including affect and our awareness of our emotional states, sensorimotor functions, nociception, decision-making under uncertainty, empathy, and saliency[26–29]. Here we focus on its encoding of the pain of other individuals not because we think this is the insula's primary role, but because within the literature on empathy, the insula plays a key role, and the pain of others is a salient stimuli that can affect nociception[30], affect[31] and decision-making[22,32], and thereby engages many of the functions associated with the insula.

Figure 1. Experimental design and recording site locations. (a) Frames extracted from a Hand and Face movie. For the Face, the first second of each movie showed a neutral facial expression, the second, the facial reaction to the shock. For the Hand, the movie started with the belt resting on the hand. The first second showed the belt lifting and coming down again, to hit the hand at the 1 s mark exactly. The hand then reacted to the force of the belt. Both the slap and the shock delivery happened in the middle of the movies, splitting them into a 1 s neutral and a 1 s pain period. (b) Single trial structure diagram. After the presentation of each video, patients expressed their choice at their pace using the keyboard keys f, g, j, k for pain intensities 1-2, 3-4, 5-6, 7-8, respectively. ITI started with participant's response. (c) Position (i.e. midpoint between two adjacent electrodes) of the 85 bipolar recording sites shown as dots and of the microwire locations shown as plusses, color coded by patient. Data from the two hemispheres and all latero-medial coordinates are projected here onto a single sagittal slice of the insula taken at X=38 from the brain of one of the patients. For a list of all MNI coordinates, see Supplementary Table 3.

Materials and methods

Patients and electrodes

Depth electrode recordings were collected from 9 epileptic volunteers, admitted at the Amsterdam UMC to localize seizure origin. Patients received initial study information from the neurosurgeon, and provided informed consent to the experimenter before the surgery occurred. Our single session experiment started on average 4 days after surgery (std = 1.89 days). Two patients were excluded from the analyses due to poor behavioral performance (Supplementary Materials Section 1), and seven were included (4 females, 34.3y±9std). The study was approved by the medical ethical committee of the Vrije University Medical Center

(protocol 2016/037) and each patient signed a written informed consent according to the Declaration of Helsinki. There was no indication of epileptic onset in electrode locations used in our analyses (see Supplementary Materials Section 2). Patients were implanted with Behnke-Fried depth electrodes[33] (Ad-Tech Medical Instrument Corporation) targeted at the right or left, anterior or posterior insula.

Video pain rating task

The 2 s videos were generated as in Gallo et al.[22] and showed a Caucasian female receiving either electrical shocks to the hand (reaction conveyed by the facial expression only; Face condition) or a slap with a belt to the hand (reaction conveyed by the hand only; Hand condition). Hence the location of the noxious stimulation was maintained across conditions (dorsum of the left hand), but the cues through which participants could deduce the painfulness differed. All videos started with 1 s of baseline: neutral facial expression for Face and static hand for Hand stimuli. Movies were cut, so that evidence of pain started at 1 s (Fig. 1a). Before the experiment, participants were instructed to rate pain intensity ("How much pain do you think the person felt?") on a scale from 0 (no pain at all) to 10 (the worst imaginable pain). To reassure patients that no real harm was never inflicted to the actor in the movie, they were informed that during video recording stimulations in the 9-10 range were never used. Participants had to rate pain intensity after each video at their own pace, using 4 keyboards-keys (Fig. 1b). Only the relevant keys were presented on the screen, intensities were not indicated. Patients watched each of the 60 videos (30 Hand, 30 Face) twice in fully randomized fashion with a random interval of 1.5 s±0.5. The videos were matched in terms of intensity and standard deviation based on a validation in Gallo et al.[22]

Preprocessing of LFP signals

To reduce artefacts and extract local signals, iEEG macro contact recordings were digitally re-referenced in a bipolar layout (Supplementary Fig. 2). This generated 85 bipolar recordings from 102 contacts in the insula, with patients having between 5 and 19 bipolar channels (Fig. 1c, Supplementary Table 3). Re-referencing attenuated 50 Hz noise sufficiently to omit digital filters that distort data. Continuous recordings were separated into trials of 4 s: 1 s premovie baseline, 2 s video, and 1 s postmovie. Trials were visually checked for ground failure and amplitude saturation (none was detected), downsampled to 400 Hz, and detrended.

Time-frequency decomposition of LFP signals

A sliding window Hanning taper based approach was used for each trial with the following parameters: frequencies from 1 to 200 Hz in steps of 1 Hz; time points from -1 s (relative to movie onset) to 3 s in steps of 0.0025 s; and for each frequency, a single sliding Hanning taper window with the duration of 8 cycles (maximum=1 s; minimum=0.1 s). Trials were expressed as % power change relative to baseline (-1 s to 0 s) separately for each frequency: $y(t)=(P(t)-P_0)/P_0)$, with P_0=average of baseline. Points with $y(t)$ ±10 standard deviations from the mean of the other trials were excluded to not reject entire trials, but only

outlier time-frequency points in some trials (rejections were rare, mean rejected time-frequency points=0.0032%±0.0035std).

General statistical approach and intensity coding

Our core question is how the insula codes perceived intensity. Supplementary Materials Section 4 describes our statistical approach in detail. Briefly, we consider a bipolar channel to show *intensity coding*, if its trial-by-trial power variations correlate positively with the variation in pain intensity reported by the patient. We always recoded the 1-2, 3-4, 5-6, 7-8 rating options as 1, 2, 3, 4. For each bipolar recording, we then calculated the Spearman's rank correlations (due to the ordinal nature of intensity ratings) between the patient's intensity rating and power estimate over all trials for each time-frequency intersection separately, or within a certain power-band. A one-sample t-test was used to test whether the average correlation over the 85 bipolar recordings differed from 0. Correlations were not Fisher r->z transformed, because r and z values remain virtually identical for $-0.5 < r < 0.5$, which is the range in which the correlations we examined remain. Where Shapiro-Wilk tests suggest deviations from normality, non-parametric tests are used, and indicated by W (Wilcoxon signed rank) or U (Mann-Whitney U). Additionally, where evidence for the absence of an effect is relevant, we provide Bayes-factors that quantify the relative evidence for H_1 and H_0 (BF_{10}), or for a one-tailed H_+ vs H_0 (BF_{+0}), and use $BF < \frac{1}{3}$ as the critical bound for evidence of absence as detailed in ref [34]. For p-values, we specify one-tailed vs two-tailed tests for t-tests using the index 1 or 2 (p_1=one-tailed *p*-value, p_2=two-tailed *p*-value). For F-tests, two-tailed tests are reported.

Spike sorting and selection of responsive neurons in microwires

Three patients had microwires (Behnke-Fried electrodes[33], Ad-Tech Medical) in the insula protruding from the electrode tip (Fig. 1c '+'). Spikes were detected and sorted using Wave_Clus2.[35] In short, raw data was filtered between 300-3000 Hz. As per default settings, spike waveforms were extracted from 0.625 ms before to 1.375 ms after the signal exceeded a 5*noise threshold, where noise was the unbiased estimate of the median absolute deviation. Wave_Clus2 sorted and clustered the waveforms automatically and were manually checked by author RB. Clusters were excluded in which >2% of spikes were observed with an inter-spike interval <2 ms or with firing rate <1 Hz. To identify cells that responded to our stimuli, we used a Wilcoxon signed rank test comparing spike counts during baseline (-1 s to 0 s) against that during the pain period (1 s to 2 s) for Hand and Face trials together. Only cells that showed a response to the stimuli ($p_1 < 0.05$), irrespective of pain intensity, were considered for further analysis.

Intensity coding in spiking

Similar to LFP analyses, a cell was said to show *intensity coding*, if spike counts rank correlated positively with reported intensity. Because JASP includes Bayesian statistics using Kendall's Tau but not Spearman r, we used the former to quantify evidence for or against intensity coding.

Broadband power analyses in microwires

To explore whether intensity coding in cells and the broadband power (BBP, 20-190 Hz, Fig. 2b) from the same microwire were related, for the 10 microwires that yielded responsive neurons (whether these neurons showed intensity coding or not) we quantified the association between BBP averaged over the pain period (1-2 s) and intensity ratings (1-2, 3-4, 5-6, 7-8) using rank correlation coefficients separately for face and hand videos (again using Kendall's Tau to provide BF_{+0} estimates). All 8 microwires protruding from the same electrode were first re-referenced to the microwire with the least spiking and lowest artefacts, yielding seven microwire recordings for each of the 4 electrode-tips with wires in the insula. Data were filtered to remove 50 Hz noise and harmonics at 100 and 150 Hz. Subsequently, they were separated into trials of 4 s (-1 s to 3 s relative to video onset), downsampled to 400 Hz and visually checked for artifacts. The time-frequency decomposition of power followed the same procedure as for the macro contact recordings. Finally, intensity coding at the level of spikes (i.e. r_K(spikes,rating)) and BBP (r_K(BBP,rating)) from the same wire were compared using a Kendall's Tau coefficient.

fMRI experiment

Participants. Twenty-five healthy volunteers participated in the study. The full dataset of two participants was excluded from the analyses because head motions were above 3mm. Analyses were performed on the remaining twenty-three participants (13 females; mean age =28.76 years old +-SD=6.16). The study was approved by the local ethics committee of the University of Amsterdam (project number: 2017-EXT-8542).

Stimuli and rating task. The video pain rating task was performed as described above with the following differences. Each trial started with a grey fixation cross lasting 7-10 s, followed by a red fixation cross for 1 s, followed by the 2 s video, followed by a red fixation cross lasting 2-8 s, followed by the rating scale ranging from 'not painful at all' ('0') to 'most intense imaginable pain' ('10'). The design also includes another condition we will not analyse here, in which participants viewed videos varying in color saturation and had to report on a scale from 'not a change' ('0') to 'a very big change' ('10'). Participants were asked to provide a rating by moving the bar along the scale using two buttons for right and left (index and middle finger) and a third one for confirming their response (ring finger) using their left hand. The direction of the scale and the initial position of the bar was randomized in each trial. The videos used for the Face and Hand conditions for the electrophysiology and fMRI experiment were generated in the same way but were not identical.The task was split up into 46 blocks of 30 trials each, 2 blocks of electrical pain stimulations and, 2 blocks of mechanical slaps by a belt and 2 blocks of videos with changes in color saturation (presented in 46 separate fMRI acquisition runs). The last condition will not be discussed in this manuscript. Anatomical images were recorded between the fourth and fifth run of fMRI acquisition.

Data acquisition and processing. MRI images were acquired with a 3-Tesla Philips Ingenia CX system using a 32-channel head coil. One T1-weighted structural image (matrix = 240x222; 170 slices; voxel size = 1x1x1mm) was collected per participant together with EPI (echo-planar imaging) volumes (matrix M x P: 80 x 78; 32 transversal slices acquired in

ascending order; TR = 1.7 s; TE = 27.6 ms; flip angle: 72.90°; voxel size = 3x3x3 mm, including a .349 mm slice gap). MRI data were processed in SPM12. EPI images were slice-time corrected to the middle slice and realigned to the mean EPI. High quality T1 images were coregistered to the mean EPI image and segmented. The normalization parameters computed during the segmentation were used to normalize the gray matter segment (1x1x1 mm) and the EPI images (2x2x2 mm) to the MNI templates. Finally, EPIs images were smoothed with a 6 mm kernel.

At the first level, we defined separate regressors for the Hand videos, Face videos, rating-scale and button-presses. Analyses focused on the 2 s that the videos were presented. The trial-by-trial ratings given by the participants were used as a parametric modulator, one modulator for the face and one for the Hand trials, on the respective video regressor. The rating-scale regressor started from the moment the scale appeared and ended with participants' confirmation button press. The button-press regressor, finally, had zero duration and was aligned to the moment of all the button presses. Six additional regressors of no interest were included to model head movements. To quantify the degree to which each voxel in the insula had BOLD (blood-oxygen-level-dependent imaging) activity associated with trial-by-trial ratings, we then brought the parameter estimate for the parametric modulator obtained for Hand and Face trials separately to a second level t-test with 23 participants, and then used the resulting t-value as a measure of the random effect effect size of the association. We used t-values rather than the average value of the parametric modulator, because these values are to be compared against out-of-sample values of patients, and the topography of t-values is a better predictor for out-of-sample generalizations. However, the average parameter value correlated above 0.9 with the t-value.

To investigate whether the pattern of BOLD activity across all voxels in the insula encodes intensity, we additionally performed a multivariate regression analysis akin to the PLSR for the BBP described in Supplementary Materials Section 8. For each participant, we performed a general linear model that estimated a separate parameter estimate for the video epoch of trials in which participants gave a rating of 0-2, 3-4, 5-6 or 7-8 respectively, separately for Hand and Face trials. In matlab we then loaded for each subject the parameter estimate images for each level of rating, and only included voxels that fell within our insula mask (Faillenot *et al.*, 2017). We then trained a weighted partial least-square regression using the matlab function plsregress and the data from all but one participant to predict rating based on a linear combination of the parameter estimates in each voxel and then used this linear combination to predict the rating of the left-out participant, then repeated the procedure for each participant. We weighted the regression by replicating each parameter estimate image in the training and testing set by the number of trials that went into it. Then we quantified how accurately the regression predicted the rating of the left-out participants using Kendall's tau. We then tested whether the performance was above chance by comparing the 23 prediction accuracies (Kendall's Tau) against zero in a one-tailed test. Based on the analysis on the BBP we performed this analysis with procedure using 2 or 3 components.

Data Availability

The data presented in this work are available upon request

Results

Our patients provided ratings with similar standard deviation for the Hand and Face stimuli ($SD_{Hand}=0.82$, $SD_{Face}=0.91$), although they rated the Hand stimuli as slightly higher in pain intensity than the Face stimuli ($Mean_{Hand}=2.7\pm0.82sd$, $Mean_{Face}:1.95\pm0.91$; Supplementary Table 1). Comparing the ratings of the 7 patients that were included in our final analysis with those of an age and gender matched control group revealed that the ratings of the patients were in the normal range (Supplementary Materials Section 1).

iEEG activity in the insula correlates with the perceived intensity of the pain of others

Figure 2. Intensity coding in the local field potentials from macro-contacts in the insula. (a) For each frequency and time relative to stimulus onset, the average r_S-value over all insular bipolar recordings between iEEG power and rating for Face and Hand trials together. (b) The same as (a), but cluster corrected for multiple comparison as described in Supplementary Materials Section 5.1.[36] BBP: Broad-Band Power, the cluster of significant positive intensity coding frequencies (i.e. $r_S>0$, 20-190 Hz) used throughout the paper. (c) Mean±sem time course of intensity coding in BBP over the 85 recordings when Face and Hand trials are combined. Above the x-axis, orange bars show periods of significant intensity coding after circular shift correction as defined in Supplementary Materials Section 5.1 during the pain period, the brief black bar indicates negative correlation significance during the neutral period. Below the x-axis, the black block marks the neutral and the warm block the pain period. (d) The t-value of a t-test comparing the intensity coding of all 85 bipolar recordings combining Hand and Face trials within the pain period (1–2 s post-stimulus onset) in the insula against zero (red) was higher than the distribution of corresponding t-values obtained when performing the same test using 85 bipolar recordings randomly selected 100,000 times from the macro-contacts of our 7 patients anywhere in the brain (Supplementary Materials Section 5.4). (e) Mean±sem % BBP changes relative to baseline as a function of reported intensity separately for the neutral (black) and pain (yellow-red) period when combining Hand and Face trials. BF_{10} values:

Bayes-factor quantifying evidence for H_1 relative to H_0 from a non-parametric t-test comparing BBP power during the pain period against that during the neutral period. ***: $p<0.001$ relative to the preceding reported intensity. See Supplementary Materials Section 5.3 for ANOVA details. (f) Time course of intensity coding in BBP for Hands and Faces separately. $r_S>0$ indicated with green bars for Hands and purple for Faces. Black bars indicate $r_{S\_Hand}>r_{S\_Face}$. The early and late periods that result for Hands and Faces, respectively, will be used throughout the paper. (g) Mean±sem BBP as a function of rating for Hands and Faces in the early and late periods. Green ***: $p<0.001$ relative to the preceding intensity for the Hand. Black *: $p<0.001$ main effect of rating (i.e. combining Hand and Face) similar to panel (e). Purple BF_{10}: evidence for a lack of difference across ratings for the Hand. Black BF_{10}: lack of difference between rating 1-2 and 3-4 with Hand and Face combined.

Plotting the power over the bipolar electrodes as a function of perceived intensity, irrespectively of whether Hand or Face movies were shown (Supplementary Fig. 3a-d) suggests an increase in power for the highest ratings. Correlating power with reported intensity and applying cluster corrections (Supplementary Materials Section 5.1) revealed: a cluster of positive correlations ranging from 20-190 Hz and 1.12-1.62 s ($p_1<0.001$; p_1=one-tailed p value); one at very low frequencies (1-6 Hz, 0.02-2.055 s; $p_1<0.001$), and a small cluster of negative correlations (13-17 Hz, 1.295-1.83 s; $p_1=0.004$; not further discussed, Fig. 2a-b). Intensity coding was apparent in all traditional frequency ranges, except alpha (Supplementary Materials Section 5.2 and Supplementary Fig. 3f-j) and, as expected, was significant in the pain period. With no obvious differences among frequency-bands above alpha, we henceforth used the frequency band 20-190 Hz for all analyses and refer to it as broadband power (BBP). We concentrate on BBP rather than oscillatory signals in lower frequencies because BBP is more closely linked to neural spiking[8–10], cannot be explored in non-invasive EEG recordings, and is the frequency range that can supplement the information available for the substantial fMRI literature.[6,7] The temporal profile of the BBP-rating association revealed two periods with significant positive correlations: 1.1375–1.54 s and 1.7375–1.9575 s (Fig. 2c). Averaging BBP power over the entire pain period revealed that out of 85 macro contacts, 27 (32%) showed a significant positive correlation (assessed as $p_1<0.05$, Supplementary Fig. 3k) between perceived intensity and BBP (n=120 trials, all $r_{S[118]}>0.156$, $p_1<0.045$), which was extremely unlikely to occur by chance (Binomial $p_1=5\times10^{-15}$, $BF_{+0}=3\times10^{12}$). Also, randomly picking 85 electrodes anywhere in the brain yielded BBP-rating associations that were significantly lower than those we find in the insula ($p_1=4\times10^{-5}$, Fig. 2d and Supplementary Materials Section 5.4), confirming that the BBP in the insula has enriched intensity coding. The effect of reported intensity depends mainly on BBP power increases for the two highest intensity ratings (Fig. 2e and Supplementary Fig. 3a-d).

Intensity coding arises earlier for the Hand than Face stimuli

To investigate how intensity coding depends on the stimulus, we then separated Face and Hand trials. The time-frequency analysis revealed regions of broadband intensity coding earlier for Hand than Face, as reflected in the direct contrast (Fig. 2f, and Supplementary Materials Section 5.5). Focusing on the BBP frequency range identified independently of stimulus type, significant intensity coding for the Hand from 1.0075 to 1.4375 s (hereafter 'early period') and for the Face from 1.75 to 1.8625 s and from 1.905 to 1.975 s, which jointly (1.75-1.975 s) will be called 'late period' (Figure 2f). The insula thus reflects in broadband

activity the perceived intensity with differential time courses for hand and face videos in the current study.

　　　To explore the shape of the BBP-rating relation, we averaged BBP over time for the early and the late periods for each pain rating separately (Fig. 2g). For the early period, a stimulus (Hand, Face) x rating repeated-measures ANOVA revealed a significant interaction (Greenhouse-Geisser-corrected $F_{[2.183,102.621]}$=13.55, p=3x10^{-6}, BF$_{incl}$=2x10^6). Planned comparisons provided evidence that BBP for Faces in the early period was similar for consecutively increasing painfulness level pairs (n=48 since all possible rating options were used by 4 patients with a total of 48 electrodes, all $t_{[47]}$<0.252, p_2>0.802, BF$_{10}$<0.163). Note that throughout the text, for t-tests, p_1=one-tailed p-value, and p_2=two-tailed p-value.), whereas, there was an orderly increase in broadband power for increasing pain ratings for Hands from 3-4 onwards (3-4 vs 5-6: $t_{[47]}$= 5.97, p_2=3x10^{-7}, BF$_{10}$= 51110; 5-6 vs 7-8: W= 188, p_2=2x10^{-5}, BF$_{10}$= 764.63). However, BBP for ratings of 1-2 was unexpectedly higher compared to the ratings of 3-4 (W=1014, p_2=3x10^{-6}, BF$_{10}$= 847.14). A similar ANOVA for the late period, revealed evidence for the absence of an interaction ($F_{[3,141]}$=0.55, p=0.650, BF$_{incl}$=0.034). There was only a significant main effect of rating ($F_{[3,141]}$=16.54, p=3x10^{-9}, BF$_{incl}$=2x10^7), indicating that BBP in the late period of the Hand and Face videos together was the same for ratings 1-2 and 3-4 (W= 597, p_2=0.931, BF$_{10}$= 0.163), but thereafter showed significant increases with each consecutive increase in pain ratings (3-4 vs 5-6: $t_{[47]}$= 3.46, p_2=0.001, BF$_{10}$= 25.147; 5-6 vs 7-8: $t_{[47]}$= 2.90, p_2= 0.006, BF$_{10}$= 6.292). Taken together, these analyses indicate BBP in the insula reflects perceived intensity only for the Hand stimuli in the early, and for both stimulus types in the late window.

The timing of shape information matches that of Face intensity coding

Figure 3. Temporal dynamics of pain perception. (a) Mean±sem of Spearman correlation between each participant in the frame rating task and the average ratings of the other participants in the video rating task (r_S(frame,average_video), green and purple) compared against that between the participants rating of the video and the average rating of the other participants in the video rating task (r(video,average_video), gray) for Hands and Faces. Black statistics above the bars compare the respective frame and video rating, the colored statistics compare the frame rating against zero. The black statistic under the bars compare the frame rating across Hand and Face. See Supplementary Materials Section 6.1 for additional methods and analyses. (b) An example frame from the face videos that was used in the frame rating task. The average activation of the Action Units (AU) 4 and 7 was used to estimate the intensity of the shape information in Faces. Mean±sem shape estimation as a function of time for Face videos rated as 1-2, 3-4, 5-6, 7-8. Purple bars indicate the period with significant BBP-rating correlations for Faces, as seen in Fig. 2f. (c) The same as (b), but for motion information in Face videos. (d) The same as (c), but for motion information in Hand videos. Green bar shows the period with significant BBP-rating correlations for Hands, as in Fig. 2f. (e) r_s between motion energy and BBP (blue) or between subjective rating and BBP (orange) for the 6 consecutive bins of 333 ms during the movie. Error bars=sem. All statistics are two-tailed parametric t-tests against zero, because r_s values were normally distributed (All Shapiro-Wilk $p>0.05$). Values are indicated in the table above each panel for each time-bin of 1/3s. FDR correction is over the 6 bins. No r_s-to-z transform was used because the r_s-values were in the range $-0.5<r_S<0.5$ for which r and z value are extremely similar. (f) same for the partial correlations: r_S(BBP,motion|rating) in blue and r_S(BBP,rating|motion) in orange.

Our facial stimuli have both shape and kinematic information. Leveraging the high temporal resolution of our iEEG recordings, we can ask which of these two sources of information best match the timing of our intensity coding in the insula. We first subjected our stimuli to more detailed, time-resolved analyses (see Supplementary Materials Section 6 for detailed methods). Psychophysics on a subset of 40 participants from our original online video rating revealed that participants can recognize the pain intensity of Face stimuli from static frames taken at key moments in the stimuli with accuracy even better than that for the entire video (W=193, p_2=0.003, BF_{10}=31.293, Fig. 3a). The same is not true for the Hand stimuli, for which accuracy dropped significantly when using still frames (n=39 participants who successfully rated both the dynamic videos and still frames, $t_{[38]}$=11.959, p_2=2x10^{-14}, BF_{10}=4x10^{11}). This suggests that static shape information could suffice to explain ratings for Face stimuli. Using the automated face analysis software FaceReader to extract the two most reliable shape features of painful facial expressions (facial Action Units AU4 and AU7[37], Fig. 3b), we find that how lowered the eye-brows (AU4) and how tightened the eye-lids are (AU7) in key frames suffices to predict participants' perceptions of the pain intensity in Face stimuli with r_p=95% accuracy (Supplementary Materials Section 6.2), and we thus use the average time-course of these two AUs to track shape information in our Face stimuli (Supplementary Fig. 5a). Figure 3c shows shape information increases towards the end of the movies with rating intensity. Comparing the timing of intensity coding for the Face in the insula BBP (purple bar in Fig. 2f and 3b) with the timing of the shape information for the Face (separation between the curves in Fig. 3b) shows a nice correspondence, with both highest late in the movie. Regarding kinematics, we calculated the changes in pixel-values across consecutive frames to track the timing of motion (Fig. 3c,d), and this information could also predict the rating of our patients with high accuracy for face stimuli (Supplementary Fig. 5c). Comparing the timing of intensity coding in the insula for Face stimuli (Fig. 3c purple bar) with the timing of motion and kinematic information (Fig. 3b-c) shows that maximizes when motion information has already declined. This suggests that shape information could dominate Face intensity coding. Using static facial expressions in the future experiments will however be essential to determine if static information alone can suffice. A more quantitative approach, that estimates how much motion and shape information predict our patients' ratings on a frame-by-frame basis, and how the neural intensity coding lags behind this information (Supplementary Materials Section 6 and Supplementary Fig. 5g) confirms this analysis, and suggests that neural responses lags behind the shape information by 40-320 ms.

Specific kinematic information could drive Hand intensity coding

Motion energy is also a reliable predictor of pain intensity rating for the Hand stimuli (Supplementary Fig. 5c). In our movies, motion occurs at two time points: early, when the belt is lifted up, and then again, while the belt hits the hand (Figure 3d). Pain however, occurs only at the second point in time. This allows us to explore whether the insula is coding movement in general, or movement that is associated with pain more specifically. We thus divided the two seconds of the Hand movies in 6 segments, and asked, for each segment, how well BBP relates to motion energy in the same segment (Figure 3e). Over the 85 channels, we had

evidence of absence for a relationship during the neutral period that contained the period during which the belt was seen to move upwards (all $BF_{10}<1/3$), and evidence for a relationship during the first 666ms of the pain period when the belt is seen to slap the hand (both $p_{unc}<0.003$, or $p_{bonf}<0.018$ corrected for 6 bins, both $BF_{10}>8.8$). Indeed a repeated measurement ANOVA comparing the correlation values across the 6 bins confirms that the relationship between motion energy and BBP changes as a function of time ($F_{[5,420]}=2.9$, $p=0.014$, $BF_{incl}=1.52$). This shows that the BBP response in the insula does not code motion in general, but motion at a time when it is relevant, here to assess the pain intensity. Next, we asked whether subjective rating or motion energy was the best predictor of BBP across the 6 bins (Figure 3e). Rating per se was an even better predictor of BBP than motion energy (rmANOVA, 2 predictor x 6 bin, main effect of predictor: $F_{[1,84]}=23$, $p=7\times10^{-6}$, $BF_{incl}=13473$). Interestingly, using partial correlations, we see that the correlation between rating and BBP remains highly significant when seeing the belt hit the hand even after removing what can be explained by motion energy, but we have evidence for the absence of a correlation between motion energy and BBP if removing the variance explained by rating (Fig. 3f). Together, this data supports the idea that the insula could employ motion to encode the painfulness in our Hand stimuli, but does not respond to simply seeing the motion of the belt, and that subjective rating of intensity appears to mediate the relationship between motion and insular response.

A quantitative approach that estimates how much the neural intensity coding lags behind motion information (Supplementary Materials Section 6 and Supplementary Fig. 5e) suggests that for the Hand stimuli, the insula response lags behind the motion information by 0-80 ms.

The insula contains a surprising number of stimulus-specific intensity coding locations

Figure 4. Relation between hand and face intensity coding in the insula. (a) Topographical maps of BBP-rating correlation coefficients for Hands and Faces in the early and late periods. Each circle is one of the recording sites (as in Fig 1c), with filled circles indicating locations with significant correlation coefficients ($p_1<0.05$). (b) Classification of recording locations based on their Hand (early period) and Face (late period) intensity coding. Bipolar recordings in the gray zone (n=5) significantly co-represent intensity for Hands and Faces (both $p_1<0.05$, i.e. beyond dashed line). Recordings in the purple (n=6) and in the green (n=10) zone represent intensity coding selective for Faces or Hands, respectively, i.e. one $p_1<0.05$ and one $BF_{+0}<⅓$. (c) Location of all 85 bipolar recordings color-coded by selectivity as described in (b). Note that locations Hand and Face without further specification are those with r_S values for the other stimulus type falling between the dashed and dotted line, that provides inconclusive evidence, and thus show neither significant dual coding, nor evidence of absence. Open circles represent all locations with $p_1>0.05$ for both Hand and Face. (d) Correlation coefficients for Hands and Faces separated by coding characteristics in (b) for all patients together and for an exemplary patient. p_{bino} refers to the likelihood to find the observed number of locations in that quadrant using a binomial distribution as explained in Supplementary Materials Section 7. (e) The left two panels depict the average pearson correlation

coefficients, together with corresponding resampling null distributions, as a measure of the accuracy of decoding intensity ratings using the partial least square regression (PLSR) beta coefficients of BBP in the early period for Hands and in the late period for Faces as described in Supplementary Materials Section 8. The right panels are similar to left panels, but show the accuracies of cross-decoding, predicting Hand ratings from the Face BBP and vice versa. The dotted lines indicate 95[th] percentiles of the resampling null distributions.

We next focused on how individual recording sites reflected perceived intensity. In the early period, for Hand videos, 21/85 (25%) showed significant intensity coding (rating-BBP correlations, n=60 trials, all $r_{S[58]}$>0.219, p_1<0.046), which was above chance (Binomial, 21/85 at α=0.05, p_1=9x10^{-10}, BF$_{+0}$=2x10^7). In contrast, for the Face, only 3/85 showed intensity coding (4%), which is expected by chance (Binomial p_1=0.804, BF$_{+0}$=0.025). During the late period, above chance numbers of recordings showed intensity coding for Hand (14/85, 17%, p_1=8x10^{-5}, BF$_{+0}$=201.41), and the same was true for Face videos (15/85, 18%, Binomial p_1=2x10^{-5}, BF$_{+0}$=808.49; Fig. 4a).

If the insula simply represents salience, one might expect a tight association between intensity coding for the Hand and Face, and an above chance number of locations showing intensity coding for Face and Hand. In contrast, if the insula also represents more specific information, we would expect above-chance numbers of locations with intensity coding for the Face but not the Hand and vice versa. Statistically, we infer the *presence* of intensity coding based on r_S>0, p_1<0.05, like elsewhere in the manuscript, and its *absence* using Bayesian statistics [34], with BF$_{+0}$<⅓. Plotting each bipolar recording's r_S values on an x-y plot, with x representing r_S for the Hand and y for the Face, with dashed and dotted lines at critical r_S values corresponding to p_1<0.05 and BF$_{+0}$<⅓, we define 9 quadrants, three of which are of conceptual importance (Fig. 4b): those of locations with dual intensity coding (i.e. p_1<0.05 for Face and Hand), those with Face-selective intensity coding (i.e. Face p_1<0.05 but Hand BF$_{+0}$<⅓) and those for Hand-selective intensity coding (i.e. Hand p_1<0.05 but Face BF$_{+0}$<⅓). We then used binomial tests to compare the proportion of locations falling in these three quadrants against chance, and found that all three quadrants contain more locations than expected by chance (Fig. 4d and Supplementary Materials Section 7 for details of the statistics and why for BF$_{+0}$<⅓, false negative p≈0.05 based on [34,38]). Indeed even within a single patient, amongst simultaneously recorded channels, we find above chance numbers of Face and Hand-selective channels (Fig. 4d). Also, calculating the association between intensity coding across Hand and Face through a simple correlation of the respective r-values, confirms the presence of a significant but weak, and barely worth mentioning (in a Bayesian sense) association (r_K=0.131, p_1=0.038, BF$_{+0}$=1.27). Together, this shows the insula is a patchwork, with some locations representing the Hand but not the Face, others the Face but not the Hand, and a small number finally representing both in terms of intensity coding. The spatial distribution of these locations is shown in Fig. 4c.

In addition, we used a multivariate partial least square regression (PLSR) approach to assess how well the pattern of BBP across the insula can predict participants' pain ratings (see Supplementary Materials Section 8). BBP across the 85 sites in the early period for Hand videos can be used to predict the patients average rating of the stimulus with reasonably high accuracy (n=10 trials since 1/3 of the 30 unique videos were used for testing decoding performance for each randomization, $r_{P[8]}$=0.575, p_1=9x10^{-4} based on reshuffled distribution), and BBP in the

late period for Face videos with almost significant accuracy ($r_{P[8]}$=0.331, p_1=0.058, Fig. 5e). A direct comparison of the performance of the two PLSR indicates that the performance was higher for the Hand than the Face (Non-parametric test across the decoding performances, W=944605, p_2=9x10^{-260}, BF$_{10}$=7x10^{39}). To test if intensity was encoded through similar patterns for the two stimulus types, we repeated the analyses training the PLSR on one stimulus type and testing it on the other. We found above-chance cross-decoding in both cases (Hand->Face: $r_{P[8]}$=0.343, p_1=0.029; Face->Hand: $r_{P[8]}$=0.389, p_1=0.023; Fig. 4e). However, when the 5 contacts that significantly co-represented pain intensity for both hand and face videos (the black dots in Figure 4c) were excluded from the analyses, the cross-decoding accuracy fell to insignificant levels (Hand->Face: $r_{P[8]}$=0.175, p_1=0.153; Face->Hand: $r_{P[8]}$=0.185, p_1=0.149). These findings corroborate the above results, indicating that perceived pain intensity is reflected in the insula as a mixture of hand-specific, face-specific, and hand-face common representations.

Hand intensity coding increases anteriorly as in a similar fMRI task

Figure 5. **Relationship between BBP and BOLD activity and intensity coding.** (**a**) Correlation (r_K) between MNI coordinates and BBP intensity coding, separately for Hand (green) and Face (purple). Bold numbers mark evidence for (BF$_{10}$>3) or against (BF$_{10}$<⅓) a correlation. Statistical values were obtained by correlating separately the x, y or z coordinate of each bipolar recording with the r_S(BBP,rating) of each recording over all 85 recordings using JASP. Tau refers to Kendall's Tau, p_2 and BF$_{10}$ the two-tailed probability and BF based on H$_0$:Tau=0. (**b**) Results of the regression analysis between resting state connectivity and intensity coding for the 85 bipolar recording coordinates for the Hand. Warm colors indicate significant positive, and cold, negative regression values. (**c**) Predictive performance of a PLSR trained to predict ratings based on the pattern of BOLD activity across all voxels in the insula for different ratings. A leave-one-out cross-validation was used, and each circle represents the r_K between the predicted and actual rating for each left out participant, and the p_1 and BF$_{+0}$ values then test these n=23 correlation values against zero using a non-parametric association test (Kendall's Tau). Results are shown as mean±sem separately for Hand and Face trials, and using 2 or 3 components. (**d**) Topography of intensity coding for the Hand (left) and Face (right), as assessed at the group level, by the parametric modulator capturing changes in BOLD activity that correlate with trial-by-trial differences in participant's rating. t-values testing the parametric modulator >0 at the group level are shown as a function of y and z coordinate in the insula

mask. For each coordinate, the maximum value across all x-coordinates within the two insulae is indicated. **(e)** Relation (r_P because of normality) between the t-value of the parametric modulator for the rating in the fMRI bold data (x-axis) and the BBP intensity coding (computed in the early period for the Hand, green, and late period for the Face, purple) in the iEEG signal (y-axis) for each of the 85 contact locations. Note that for the fMRI signal, the value is taken from the voxel closest to the MNI coordinates of the corresponding contact in the iEEG signal.

To examine the spatial distribution of intensity coding, we examined the relationship between MNI coordinates of the bipolar recordings and intensity coding (i.e. r_S(BBP,rating), Fig. 5a). The only significant association was that more anterior recordings (i.e. more positive y-coordinates) have higher Hand intensity coding. Interestingly, we found evidence against a right-left difference (i.e. $BF_{10}<\frac{1}{3}$ for x-coordinates) for the Face and Hand, providing moderate evidence against a left-right lateralization. To exclude that this finding could be driven by differences across patients, we also performed a random intercept mixed linear model using x, y and z coordinates as predictors of Hand intensity coding (without interactions) with patients as random nesting variables. This analysis confirmed the y coordinates predict hand intensity coding (X: $F_{[1,79.23]}=0.022$, $p_2=0.881$; Y: $F_{[1,80.97]}=13.23$, $p_2=0.0005$; Z: $F_{[1,73.95]}=0.166$, $p_2=0.685$).

To better understand the origin of the anterior gradient for Hand intensity coding, we performed a regression analysis between intensity coding of the 85 insular recording locations (for Hand and Face separately), and resting state connectivity seeded at corresponding MNI locations in Neurosynth (see Supplementary Materials Section 9). Insular locations with higher Hand intensity coding had higher resting state connectivity with the left anterior insula and ventral prefrontal cortex (including BA44/45, OP8/9, Fp1), with the right frontal orbital cortex; with the bilateral cingulate (incl. BA24/33); and the right cerebellum (Crus I and lobules VI, VII and VIII) (clusterwise FWE corrected at the cluster-level at p<0.05 using initial cluster-cutting at $p_{unc}<0.001$, $t_{[82]}=3.19$, and then setting the minimum cluster-size to $FWEc=772$ as determined by the random field theoretical calculation in SPM; Supplementary Table 4). In line with the lack of spatial gradients for the Face stimuli in the insula of our patients, examining which voxels had higher resting state connectivity with insular locations with higher Face intensity coding did not yield any significant voxels even at $p_{unc}<0.001$ (all $p_{unc}>0.001$).

Finally, to compare the spatial gradient we find using iEEG with that using fMRI, we leveraged existing data from an unpublished study in our lab using a similar design to measure brain activity using fMRI in healthy participants. BOLD activity in the insula also contained significant information about the perceived intensity for Hand and Face stimuli (Fig. 5c), and performance did not differ across Hand and Face (t-test comparing the leave-one-subject out performance for Hand and Face, with 2 components, $t_{[22]}=0.675$, $p=0.5$, $BF_{10}=0.27$; 3 components: $t_{[22]}=-0.39$, $p=0.7$, $BF_{10}=0.23$). For both the Hand and the Face stimuli, we found a gradient along the y axis with more anterior locations showing a stronger, and more positive association between BOLD activity and rating (Fig. 5d). For the Hand, across our 85 bipolar recordings in the patients, locations with higher BBP intensity coding in iEEG also show higher t-values in the BOLD signal (Fig. 5e). For the Face, we have evidence of absence for an association of the two measures (Fig. 5e).

The insula contains neurons with intensity coding for the Hand and/or Face

The insula thus displays intensity coding in a broad frequency range, including locations with Hand- or Face-specific intensity coding, as well as locations showing intensity coding for both stimulus types. To explore this representation at the level of single neurons, we analyzed the microwire data from the 3 patients (patients C, D, and E) that had microwires in the ventral anterior insula (see '+' symbols in Fig. 1c for locations and Supplementary Table 3). Spike sorting resulted in a total of 28 candidate neurons. From these, 13 showed more spikes during the pain period than the pre-stimulus baseline. Amongst those, 8 show intensity coding for the Face and/or Hand (Fig. 6), with significant Kendall's Tau correlations between perceived intensity (1-2, 3-4, 5-6, 7-8) and spike count during the pain period (1–2 s post-stimulus onset) for at least one stimulus type: 4/8 for the Face and 5/8 for Hand (Binomial, Face: p_1=0.003, BF_{+0}=27; Hands: p_1=3x10^{-4}, BF_{+0}=282). Considering the p_1-value for the intensity coding, two cells (a,b) showed intensity coding for both hand and face trials, 3 (c-e) only for hand and 3 (f-h) only for face trials. If we additionally consider the BF_{+0} values below ⅓ as evidence for the absence of coding in the other stimulus type, we find 3 Hand-specific cells (c,d,e) and 2 Face-specific cells (g,h). Importantly, within patient D, we observe the co-existence of Hand-selective (c,d) and Face-selective intensity coding (g).

Figure 6. Intensity coding in single cells and corresponding local field potentials. (a-h) Left (Face) and right (Hand) columns display, for each cell, the rastergrams and peri-stimulus time histogram (PSTH) for 8 cells that showed intensity coding for at least one stimulus type. For the PSTH, each curve represents the mean±sem of the firing rate in each bin for trials with the corresponding rating. Not all patients gave all possible ratings in each

condition. For the rastergram, trials are sorted in order of rating, with the highest ratings upmost. The colorbar next to the rastergram indicates the rows corresponding to each rating. p_1 and BF_{+0} values result from a one-tailed test of the Kandell's Tau between rating and spike-count in the pain period (marked by the dashed lines). *= significant intensity coding (p_1<0.05), t= trend (p_1<0.1), X= evidence of absence for a positive intensity coding (BF_{+0}<⅓). The x-axis (time) is relative to movie onset. The color bar on the leftmost side indicates from which patient the data is taken. Middle columns show the BBP averaged over the pain period for the corresponding microwire as a function of rating as a boxplot showing the variance across trials, with the box and whiskers representing the quartiles across trials, and the p_1 and BF_{+0}, the Kendall's Tau test of the association of rating and BBP. Note that cells c and d were taken from the same microwire, and therefore have only one BBP graph.

To explore how spiking relates to BBP, we analysed the BBP from the 10 wires that yielded the 13 cells showing stimulus triggered responses. Using Kendall's Tau correlations between BBP (20-190 Hz) and the patient's intensity ratings (1-2, 3-4, 5-6, 7-8), and comparing these results with the coding of the cells on the same wire reveals some relationship between the two. For the Hand videos, two out of three wires that yielded cells with intensity coding also showed significant association of ratings and BBP (Fig. 6a,c,d). Indeed, intensity coding (i.e. correlation between intensity rating and spiking/BBP) were significantly correlated across the 10 microwires (r_K=0.57, p_1=0.012, BF_{+0}=7.69). For the Face videos, only 1/5 wires with spike intensity coding cells showed significant intensity coding in the BBP, and ⅖ showed a trend. Across the wires, there was a trend towards an association between the intensity coding in the spikes and BBP (r_K=0.34, p_1=0.088, BF_{+0}=1.63).

Discussion

Here we characterize how the insula's iEEG activity encodes the intensity of other people's emotions, using pain as an important category. LFPs indicate that neural activity in the insula within a broad range of frequencies, including the conventional theta, beta and gamma frequency bands, scales with the perceived intensity of pain expressed by others. Interestingly, the insula only appeared to be recruited once the perceived pain level was at least moderate: activity was increased for moderate (5-6) compared to mild (3-4) and for severe (7-8) compared to moderate. However, activity for mild pain (3-4) was not significantly increased compared to minimal pain (1-2), or baseline activity. This echoes a recent finding that BBP activity in the insula is selectively increased only once thermal stimulation is consistently painful.[17] Furthermore, we isolate a small number of insular neurons increasing their firing with increases in the intensity of pain experienced by others.

As mentioned in the introduction, BOLD signals can dissociate from neural spiking.[6,7] Just as V1 BOLD signals fluctuate with perception while spiking in V1 does not,[7] the observation that BOLD signals in the insula fluctuate with perceived pain intensity alone cannot guarantee that neuronal spiking in the insula does. The insula's BOLD signal could instead fluctuate with perceived intensity simply as a result of feedback synaptic input from other brain regions that encode perceived intensity (e.g., Area 24 in the cingulate gyrus[12]). The foremost impact of our broad-band gamma and spiking data is thus to provide what is arguably the first evidence that the intensity of other people's pain is indeed locally encoded in the spiking of neurons in the insula.

The human insula has been in the focus of pain neuroscience as part of the pain matrix recruited by first-hand experience of pain.[39] In this tradition, neuroimaging evidence for activation of the insula while witnessing pain experienced by others has lead many to suggest it may represent a neural basis for empathy for pain.[1,3,4,40] However, the insula is also recruited by a variety of tasks beyond nociception and pain empathy, including other affective states, sensorimotor functions and decision-making under uncertainty.[26–29] With so many tasks recruiting the insula, the idea has become popular that it may play a key role in attaching salience to behaviorally important stimuli.[26,27,41,42] In the present study, we do not intend to (and cannot) address the selectivity of the insula for the pain of others over other emotions, and we do not claim the neural responses we report are pain-specific. Instead we characterize how the insula's iEEG activity encodes the intensity of other people's emotions, using pain as an important category, and use the agnostic terminology of 'intensity coding', rather than 'pain intensity coding' throughout the paper. That the insula's broadband activity correlated with motion energy in our Hand stimuli only while the motion was associated with pain (during the slapping) but not while motion was innocuous (during the initial belt-lifting) shows that the effects we measured cannot be reduced to visual motion detection. That perceived pain intensity was mediating the association between motion-energy and broadband activity further speaks against an interpretation of our results as reflecting unspecific motion processing. Future experiments using a wider gamut of control stimuli, that are matched in motion but that differ in emotion will be critical to address the specificity of the responses we describe, and hence, what state they could reliably signal.[41] For the hand stimuli, this could include seeing hands experience a range of different salient affective experiences, such as a hand being hurt, being caressed and being rejected in addition to a neutral hand-shake.[16] For the face, this could include disgusted, angry, fearful and happy facial expressions matched for motion.

An important, and somewhat related, question has been whether pain or salience cues are represented in a modality-specific or modality-general manner in the insula. fMRI studies have shown that the anterior insula is coactivated with different brain regions depending on whether the pain of others is conveyed via indirect cues or via the actual body part, such as the hand, that directly receives the noxious stimulation.[1–4,22] Here, we provide electrophysiological measures of neural activity that speak to that issue. We focus on the broad-band gamma signal known to have comparatively high spatial specificity and be closest to neural spiking,[8–10] and find a mixed organization, consisting of modality-specific and -general locations in a partially intermixed layout: we found locations with broadband activity and spiking associated with perceived intensity for the Hand, but not the Face; others associated with the Face, but not the Hand; and others still associated with perceived intensity for both.

Leveraging our high temporal and spatial resolution, we found that locations that showed intensity-coding for the Hand stimuli have activity timing echoing the timing of pain-related motion cues with relatively short latencies <100 ms, and that the association of motion energy and broadband activity is mediated by the perceived intensity. Locations that show intensity coding for the Face appear to have activity echoing the timing of shape information with latencies no longer than 320 ms. These latencies are in a range similar to that found following nociceptive stimulation[17] or static disgusted facial expressions.[43,44] Using automated software to detect the level of activation of the facial action units 4 and 7 (i.e. lowering eye-brows and tightening eye-lids), we find that this information suffices to predict participants'

rating of the stimuli with high accuracy, and followed the time course of the neural activity in the Face intensity encoding locations well enough to suggest that it provides traction on the analyses of dynamic pain-related facial expressions.

An important consideration is whether this selectivity for Face or Hand stimuli could simply originate from some participants finding our facial stimuli more salient, and others the hand stimuli, particularly, given that our patients rated the Hand stimuli with slightly higher pain intensity than our Face stimuli. That we find face and hand selectivity side-by-side in simultaneously recorded locations and neurons in single patients suggests that this cannot suffice to explain our data. This is because if a patient were to find Hand stimuli more salient than Face stimuli, and the insula simply codes saliency, we would expect to find Hand-specific intensity coding in that patient's insula, but we would not expect to find side by side locations with Hand-specific and Face-specific coding. Our findings instead suggest that different locations might encode behaviorally relevant (and hence salient) information about the intensity of the pain of others with some degree of specialization for a particular type of information (hand vs face). The insula intensity coding we measure could have a dual function: help perceive the intensity of other people's emotions and tag intense emotions as salient, thereby reconciling the notion that it contributes to empathy and to saliency. One peculiar observation in our data, is that broadband power for the Hand stimuli did not increase monotonically as a function of reported pain intensity, but in a J shape, with higher power for the lowest than second lowest rating. Some have argued that the insula may be part of an active inference circuitry in the brain that learns to predict emotionally relevant states and stimuli[45]. In such a perspective, a J shaped response curve could reflect a combination of underlying neurons representing the intensity of other people's emotions (with a monotonic increase in activity) and others representing the prediction error (which for randomly presented intensities would have a U shape centered on the average intensity). Future experiments could contrast responses to a given pain intensity in blocks of high and blocks of low average presented intensity to disentangle the effect of observed and expected intensity on neural responses in the insula and shed further light on this predictive component.

On the other hand, we also found evidence that some locations and cells represent intensity coding for both the Hand and the Face stimuli, and overall, if we train a partial least square regression to predict perceived intensity from the activity pattern across all recorded locations, we find that training the decoder on Hand activity pattern and testing it on Face activity patterns (or vice versa) leads to above chance decoding. This confirms that the insular representation of intensity can support stimulus independent decoding - despite our partial least square regression not being biased to focus on signals that do generalize well across stimulus types. This provides an electrophysiological basis for recent fMRI studies that show that stimuli depicting situations in which others' experience of pain or not can be discriminated using the same pattern across hand and face stimuli.[46]

In addition to the broadband results we report in detail, we find that theta power increases with perceived intensity. Given a growing animal literature establishing that interareal theta synchronization promotes learning about threats,[47–50] examining the coherence in the theta range across iEEG electrodes in different brain regions during pain observation may in the future shed light on how humans learn about safety through others.

Spatially, finally, we found that Hand intensity coding was enriched in the anterior dorsal insula, where we also found the largest proportion of locations encoding both Hand and Face intensity. This anterior bias was also observed in our BOLD signal for similar Hand stimuli. A recent meta-analysis identified that the most consistent BOLD activations when observing limbs in painful situations within the insula occur bilaterally around MNI coordinates y=13 and z=10,[4] which closely matches where we find the highest density of Hand intensity coding (Fig. 5a). Interestingly, locations with higher Hand intensity coding have increased connectivity at rest with extra-insular regions involved in processing two relevant stimulus dimensions. Connectivity was higher with cerebellar lobules VI, VII, and VIII, and with the inferior frontal gyrus, all of which are recruited by[51–53] and necessary for[53–55] perceiving the very kinematics of hand actions we find to be good predictors of BBP activity in the current study. Connectivity is also higher with the mid- and anterior cingulate cortex associated with pain witnessing in humans[1,3,4] and that contain neurons with pain intensity coding in rats witnessing the pain of other rats.[12]

With respect to Faces, somewhat surprisingly, we did not find a clear spatial clustering of intensity coding in our electrophysiological data. Overall, our ability to decode perceived intensity from our Face stimuli was also lower than that from the Hand stimuli. In that context, it is important to reflect on the fact that, despite our efforts to match the perceived intensity based on previous data[22], patients (and to a lesser extent an age and gender matched control group) perceived the Hand stimuli as more intense than the face stimuli. Given that responses were strongest for rating 4, and that this rating was given more rarely for Faces than Hands, this difference could have contributed to making our Face results less consistent. At the same time, the variance in rating, a critical factor for determining the efficiency with which a regression can detect the presence of a rating-BBP relationship, did not differ across Hand and Face. Together, this difference in perceived intensity cautions us against overinterpreting the lack of detectability of a topographic organization of responses to faces. Indeed, in our BOLD data, where decoding performance was similar for Hand and Face, face intensity coding also had a clear spatial organization, being stronger more anteriorly, and meta-analyses show the left anterior insula to be reliably recruited by the observation of painful facial expressions.[4] However, that we find fewer locations and less reliable spatial organization for Face than Hand intensity coding does dovetail with recent meta-analyses of the fMRI literature show that, when comparing studies showing limbs in painful situations with those showing painful facial expressions, the insula is more reliably recruited by the sight of limbs.[3,4] Indeed, that we find a macroscopic organization for the Hand, but not the Face, intensity coding is echoed at the mesoscale: microwires with cells with Hand intensity coding also tend to show Hand intensity coding in the BBP signal that is thought to pool the spiking of many neighbouring neurons, but the same is not true for the Face. In terms of lateralization, we find that our data is more likely if one assumes that both hemispheres have similar intensity coding, than if one hemisphere were dominant. This echoes the fact that during noxious stimulation on the right hand, both insulae show significant iEEG responses (although slightly stronger in the left insula[17]) and that fMRI fails to find robust lateralization of responses to empathy for pain.[3,4]

Acknowledgments

We thank Pieter Roelfsema for enabling the collaboration that led to the access to the patients, Eline Ramaaker for her assistance in electrode localization, Agneta Fischer and George Bulte at UvA for advice and help with the use of FaceReader, and Tess den Uyl for advice on how to use FaceReader specifically to analyse facial expressions of pain.

Funding

This work was supported by Dutch Research Council (NWO) VIDI grant (452-14-015) to VG and VICI grant (453-15-009) to CK.

Competing interests

The authors report no competing interests

References

1. Lamm C, Decety J, Singer T. Meta-analytic evidence for common and distinct neural networks associated with directly experienced pain and empathy for pain. *NeuroImage*. 2011;54(3):2492-2502. doi:10.1016/j.neuroimage.2010.10.014
2. Keysers C, Kaas JH, Gazzola V. Somatosensation in social perception. *Nat Rev Neurosci*. 2010;11(6):417-428. doi:10.1038/nrn2833
3. Timmers I, Park AL, Fischer MD, et al. Is Empathy for Pain Unique in Its Neural Correlates? A Meta-Analysis of Neuroimaging Studies of Empathy. *Front Behav Neurosci*. 2018;12. doi:10.3389/fnbeh.2018.00289
4. Jauniaux J, Khatibi A, Rainville P, Jackson PL. A meta-analysis of neuroimaging studies on pain empathy: investigating the role of visual information and observers' perspective. *Soc Cogn Affect Neurosci*. 2019;14(8):789-813. doi:10.1093/scan/nsz055
5. Paradiso E, Gazzola V, Keysers C. Neural mechanisms necessary for empathy-related phenomena across species. *Curr Opin Neurobiol*. Published online 2021. doi:10.1016/j.conb.2021.02.005
6. Boynton GM. Spikes, BOLD, attention, and awareness: a comparison of electrophysiological and fMRI signals in V1. *J Vis*. 2011;11(5):12. doi:10.1167/11.5.12
7. Maier A, Wilke M, Aura C, Zhu C, Ye FQ, Leopold DA. Divergence of fMRI and neural signals in V1 during perceptual suppression in the awake monkey. *Nat Neurosci*. 2008;11(10):1193-1200. doi:10.1038/nn.2173
8. Bartoli E, Bosking W, Chen Y, et al. Functionally Distinct Gamma Range Activity Revealed by Stimulus Tuning in Human Visual Cortex. *Curr Biol CB*. 2019;29(20):3345-3358.e7. doi:10.1016/j.cub.2019.08.004
9. Buzsáki G, Anastassiou CA, Koch C. The origin of extracellular fields and currents-- EEG, ECoG, LFP and spikes. *Nat Rev Neurosci*. 2012;13(6):407-420. doi:10.1038/nrn3241
10. Miller KJ, Honey CJ, Hermes D, Rao RPN, denNijs M, Ojemann JG. Broadband changes in the cortical surface potential track activation of functionally diverse neuronal populations. *NeuroImage*. 2014;85 Pt 2:711-720. doi:10.1016/j.neuroimage.2013.08.070
11. Hutchison WD, Davis KD, Lozano AM, Tasker RR, Dostrovsky JO. Pain-related

neurons in the human cingulate cortex. *Nat Neurosci*. 1999;2(5):403-405. doi:10.1038/8065

12. Carrillo M, Han Y, Migliorati F, Liu M, Gazzola V, Keysers C. Emotional Mirror Neurons in the Rat's Anterior Cingulate Cortex. *Curr Biol*. 2019;29(8):1301-1312.e6. doi:10.1016/j.cub.2019.03.024

13. Wicker B, Keysers C, Plailly J, Royet JP, Gallese V, Rizzolatti G. Both of us disgusted in My insula: the common neural basis of seeing and feeling disgust. *Neuron*. 2003;40(3):655-664.

14. Singer T, Seymour B, O'Doherty J, Kaube H, Dolan RJ, Frith CD. Empathy for pain involves the affective but not sensory components of pain. *Science*. 2004;303(5661):1157-1162. doi:10.1126/science.1093535

15. Jabbi M, Swart M, Keysers C. Empathy for positive and negative emotions in the gustatory cortex. *NeuroImage*. 2007;34(4):1744-1753. doi:10.1016/j.neuroimage.2006.10.032

16. Meffert H, Gazzola V, den Boer JA, Bartels AAJ, Keysers C. Reduced spontaneous but relatively normal deliberate vicarious representations in psychopathy. *Brain J Neurol*. 2013;136(Pt 8):2550-2562. doi:10.1093/brain/awt190

17. Liberati G, Mulders D, Algoet M, et al. Insular responses to transient painful and non-painful thermal and mechanical spinothalamic stimuli recorded using intracerebral EEG. *Sci Rep*. 2020;10(1):22319. doi:10.1038/s41598-020-79371-2

18. Jobst BC, Gonzalez-Martinez J, Isnard J, et al. The Insula and Its Epilepsies. *Epilepsy Curr*. 2019;19(1):11-21. doi:10.1177/1535759718822847

19. Mazzola L, Isnard J, Peyron R, Mauguière F. Stimulation of the human cortex and the experience of pain: Wilder Penfield's observations revisited. *Brain J Neurol*. 2012;135(Pt 2):631-640. doi:10.1093/brain/awr265

20. Adolphs R, Tranel D, Damasio AR. Dissociable neural systems for recognizing emotions. *Brain Cogn*. 2003;52(1):61-69. doi:10.1016/S0278-2626(03)00009-5

21. Zinchenko O, Yaple ZA, Arsalidou M. Brain Responses to Dynamic Facial Expressions: A Normative Meta-Analysis. *Front Hum Neurosci*. 2018;12:227. doi:10.3389/fnhum.2018.00227

22. Gallo S, Paracampo R, Müller-Pinzler L, et al. The causal role of the somatosensory cortex in prosocial behaviour. *eLife*. 2018;7. doi:10.7554/eLife.32740

23. Adolphs R, Damasio H, Tranel D, Cooper G, Damasio AR. A role for somatosensory cortices in the visual recognition of emotion as revealed by three-dimensional lesion mapping. *J Neurosci*. 2000;20(7):2683-2690. doi:10.1523/jneurosci.20-07-02683.2000

24. Dal Monte O, Krueger F, Solomon JM, et al. A voxel-based lesion study on facial emotion recognition after penetrating brain injury. *Soc Cogn Affect Neurosci*. 2013;8(6):632-639. doi:10.1093/scan/nss041

25. Mattavelli G, Pisoni A, Casarotti A, et al. Consequences of brain tumour resection on emotion recognition. *J Neuropsychol*. 2019;13(1):1-21. doi:https://doi.org/10.1111/jnp.12130

26. Uddin LQ. Salience processing and insular cortical function and dysfunction. *Nat Rev Neurosci*. 2015;16(1):55-61. doi:10.1038/nrn3857

27. Uddin LQ, Nomi JS, Hébert-Seropian B, Ghaziri J, Boucher O. Structure and Function of the Human Insula. *J Clin Neurophysiol Off Publ Am Electroencephalogr Soc*. 2017;34(4):300-306. doi:10.1097/WNP.0000000000000377

28. Craig AD. How do you feel? Interoception: The sense of the physiological condition of the body. *Nat Rev Neurosci*. 2002;3(8):655-666. doi:10.1038/nrn894

29. Craig ADB. How do you feel--now? The anterior insula and human awareness. *Nat Rev Neurosci*. 2009;10(1):59-70. doi:10.1038/nrn2555

30. Martin LJ, Hathaway G, Isbester K, et al. Reducing social stress elicits emotional contagion of pain in mouse and human strangers. *Curr Biol*. 2015;25(3):326-332. doi:10.1016/j.cub.2014.11.028

31. Batson CD, et al, O'Quin K, Fultz J, Vanderplas M. Influence of self-reported distress and empathy on egoistic versus altruistic motivation to help. *J Pers Soc Psychol*. 1983;45(3):706-718. doi:10.1037/0022-3514.45.3.706

32. Lengersdorff LL, Wagner IC, Lockwood PL, Lamm C. When Implicit Prosociality Trumps Selfishness: The Neural Valuation System Underpins More Optimal Choices When Learning to Avoid Harm to Others Than to Oneself. *J Neurosci Off J Soc Neurosci*. 2020;40(38):7286-7299. doi:10.1523/JNEUROSCI.0842-20.2020

33. Fried I, Wilson CL, Maidment NT, et al. Cerebral microdialysis combined with single-neuron and electroencephalographic recording in neurosurgical patients. Technical note. *J Neurosurg*. 1999;91(4):697-705. doi:10.3171/jns.1999.91.4.0697

34. Keysers C, Gazzola V, Wagenmakers E-J. Using Bayes factor hypothesis testing in neuroscience to establish evidence of absence. *Nat Neurosci*. 2020;23(7):788-799. doi:10.1038/s41593-020-0660-4

35. Quiroga RQ, Nadasdy Z, Ben-Shaul Y. Unsupervised Spike Detection and Sorting with Wavelets and Superparamagnetic Clustering. *Neural Comput*. 2004;16(8):1661-1687. doi:10.1162/089976604774201631

36. Maris E, Oostenveld R. Nonparametric statistical testing of EEG- and MEG-data. *J Neurosci Methods*. 2007;164(1):177-190. doi:10.1016/j.jneumeth.2007.03.024

37. Kunz M, Meixner D, Lautenbacher S. Facial muscle movements encoding pain-a systematic review. *Pain*. 2019;160(3):535-549. doi:10.1097/j.pain.0000000000001424

38. Jeffreys H. *Theory of Probability*. 2nd ed. Oxford University Press; 1939.

39. Ingvar M. Pain and functional imaging. Howseman A, Zeki S, eds. *Philos Trans R Soc Lond B Biol Sci*. 1999;354(1387):1347-1358. doi:10.1098/rstb.1999.0483

40. Bernhardt BC, Singer T. The Neural Basis of Empathy. *Annu Rev Neurosci*. 2012;35(1):1-23. doi:10.1146/annurev-neuro-062111-150536

41. Zaki J, Wager TD, Singer T, Keysers C, Gazzola V. The Anatomy of Suffering: Understanding the Relationship between Nociceptive and Empathic Pain. *Trends Cogn Sci*. 2016;20(4):249-259. doi:10.1016/j.tics.2016.02.003

42. Legrain V, Iannetti GD, Plaghki L, Mouraux A. The pain matrix reloaded: A salience detection system for the body. *Prog Neurobiol*. 2011;93(1):111-124. doi:10.1016/j.pneurobio.2010.10.005

43. Krolak-Salmon P, Hénaff M-A, Isnard J, et al. An attention modulated response to disgust in human ventral anterior insula. *Ann Neurol*. 2003;53(4):446-453. doi:10.1002/ana.10502

44. Chen Y-H, Dammers J, Boers F, et al. The temporal dynamics of insula activity to disgust and happy facial expressions: a magnetoencephalography study. *NeuroImage*. 2009;47(4):1921-1928. doi:10.1016/j.neuroimage.2009.04.093

45. Seth AK, Friston KJ. Active interoceptive inference and the emotional brain. *Philos Trans R Soc Lond B Biol Sci*. 2016;371(1708):20160007. doi:10.1098/rstb.2016.0007

46. Zhou F, Li J, Zhao W, et al. Empathic pain evoked by sensory and emotional-communicative cues share common and process-specific neural representations. Shackman A, Büchel C, Mehta M, Vatansever D, eds. *eLife*. 2020;9:e56929. doi:10.7554/eLife.56929

47. Tovote P, Fadok JP, Lüthi A. Neuronal circuits for fear and anxiety. *Nat Rev Neurosci*. 2015;16(6):317-331. doi:10.1038/nrn3945

48. Likhtik E, Gordon JA. Circuits in sync: decoding theta communication in fear and safety. *Neuropsychopharmacol Off Publ Am Coll Neuropsychopharmacol*.

2014;39(1):235-236. doi:10.1038/npp.2013.228

49. Likhtik E, Stujenske JM, Topiwala MA, Harris AZ, Gordon JA. Prefrontal entrainment of amygdala activity signals safety in learned fear and innate anxiety. *Nat Neurosci*. 2014;17(1):106-113. doi:10.1038/nn.3582

50. Taub AH, Perets R, Kahana E, Paz R. Oscillations Synchronize Amygdala-to-Prefrontal Primate Circuits during Aversive Learning. *Neuron*. 2018;97(2):291-298.e3. doi:10.1016/j.neuron.2017.11.042

51. Gazzola V, Keysers C. The Observation and Execution of Actions Share Motor and Somatosensory Voxels in all Tested Subjects: Single-Subject Analyses of Unsmoothed fMRI Data. *Cereb Cortex*. 2009;19(6):1239-1255. doi:10.1093/cercor/bhn181

52. Caspers S, Zilles K, Laird AR, Eickhoff SB. ALE meta-analysis of action observation and imitation in the human brain. *NeuroImage*. 2010;50(3):1148-1167. doi:10.1016/j.neuroimage.2009.12.112

53. Abdelgabar AR, Suttrup J, Broersen R, et al. Action perception recruits the cerebellum and is impaired in patients with spinocerebellar ataxia. *Brain*. 2019;142(12):3791-3805. doi:10.1093/brain/awz337

54. Pobric G, de C. Hamilton AF. Action Understanding Requires the Left Inferior Frontal Cortex. *Curr Biol*. 2006;16(5):524-529. doi:10.1016/j.cub.2006.01.033

55. Keysers C, Paracampo R, Gazzola V. What neuromodulation and lesion studies tell us about the function of the mirror neuron system and embodied cognition. *Curr Opin Psychol*. 2018;24:35-40. doi:10.1016/j.copsyc.2018.04.001