Moving Beyond the 'CAP' of the Iceberg: Intrinsic Connectivity Networks in fMRI are Continuously Engaging and Overlapping

A. Iraji[1,*], A. Faghiri[1], Z. Fu[1], P. Kochunov[2], B. M. Adhikari[2], A. Belger[3], J.M. Ford[4,5], S. McEwen[6], D.H. Mathalon[4,5], G.D. Pearlson[7], S.G. Potkin[8], A. Preda[8], J.A. Turner[9], T.G.M. van Erp[10], C. Chang[11], and V.D. Calhoun[1,*]

[1]Tri-Institutional Center for Translational Research in Neuroimaging and Data Science (TReNDS), Georgia State University, Georgia Institute of Technology, and Emory University, Atlanta, GA, USA

[2]Maryland Psychiatric Research Center, Department of Psychiatry, School of Medicine, University of Maryland, Baltimore, MD, USA

[3]Department of Psychiatry, University of North Carolina, Chapel Hill, NC, USA

[4]Department of Psychiatry, University of California San Francisco, San Francisco, CA, USA

[5]San Francisco VA Medical Center, San Francisco, CA, USA

[6]Department of Psychiatry and Biobehavioral Sciences, University of California Los Angeles, Los Angeles, CA, USA

[7]Departments of Psychiatry and Neuroscience, Yale University, School of Medicine, New Haven, CT, USA

[8]Department of Psychiatry and Human Behavior, University of California Irvine, Irvine, CA, USA

[9]Department of Psychology, Georgia State University, Atlanta, GA, USA

[10]Clinical Translational Neuroscience Laboratory, Department of Psychiatry and Human Behavior, University of California Irvine, Irvine, CA, USA

[11]Department of Electrical Engineering and Computer Science, Vanderbilt University, Nashville, TN, USA

1   * Correspondence address to:

2   armin.iraji@gmail.com (Armin Iraji)

3   vcalhoun@gsu.edu (Vince Calhoun)

Abstract

Resting-state functional magnetic resonance imaging is currently the mainstay of functional

neuroimaging and has allowed researchers to identify intrinsic connectivity networks (aka functional

networks) at different spatial scales. However, little is known about the temporal profiles of these

networks and whether it is best to model them as continuous phenomena in both space and time or,

rather, as a set of temporally discrete events. Both categories have been supported by series of studies

with promising findings. However, a critical question is whether focusing only on time points presumed

to contain isolated neural events and disregarding the rest of the data is missing important information,

or worse, producing misleading conclusions. In this work, we argue that brain networks identified within

the spontaneous blood oxygenation level-dependent (BOLD) signal are not limited to temporally sparse

burst moments and that these event present time points (EPTs) contain valuable but incomplete

information about the underlying functional patterns.

We show that brain networks are continuously present in BOLD signal, including during their event

absent time points (EATs), i.e., time points that exhibit minimum activity and are the least likely to

contain an event. Moreover, our findings suggest that EPTs do not contain all the available information

about their corresponding networks. We observe distinct default mode connectivity patterns obtained

from all time points (AllTPs), EPTs, and EATs. We show evidence of robust relationships with

schizophrenia symptoms that are both common and unique to each of the sets of time points (AllTPs,

EPTs, EATs), likely related to transient patterns of connectivity. Together, these findings indicate the

importance of leveraging the full temporal data in functional studies, including those using event-detection approaches.

Keywords

functional connectivity (FC), event present time points (EPTs), event absent time points (EATs), activation spatial map (ASM), independent component analysis (ICA), intrinsic connectivity networks (ICNs), co-activation patterns (CAPs)

Introduction

Brain function results from the interaction among local and distributed brain areas. Thus, the temporal dependency among brain regions, commonly known as functional connectivity, is a widely-used tool for studying brain function (Friston, 2011). The advent of resting-state functional magnetic resonance imaging (rsfMRI) revealed the presence of strong temporal dependencies between functionally related brain regions (e.g., bilateral motor cortices) even without external stimulation (Biswal *et al.*, 1995). This led to a rapid growth in rsfMRI research and identification of spatial patterns of functionally connected regions termed intrinsic connectivity networks (ICNs), or functional networks, at different spatial scales from large-scale distributed networks to fine-grained, spatially local ones (Damoiseaux *et al.*, 2006; Allen *et al.*, 2011; Iraji *et al.*, 2019b). Interestingly, spatial patterns obtained by applying independent component analysis (ICA) to spatial maps of thousands of different activation conditions derived from the BrainMap meta-analytic tool (https://www.brainmap.org/) closely resemble large-scale networks (Smith *et al.*, 2009), as do data-driven analyses of activation maps from task data (Calhoun and Allen, 2013), further supporting the functional relevance of functional networks. These approaches are implicitly built upon the notion of continuous information processing and interaction among brain regions.

In parallel to this work, another family of approaches, that we collectively called event detection approaches, was developed originally based upon a hypothesis that functional patterns and brain networks emerge from discrete, neural events (Chialvo, 2010; Tagliazucchi *et al.*, 2012). Instead of using all available data, event detection approaches first select a set of extreme time points (e.g., top 10%) as event present time points (EPTs) and use only these EPTs to obtain corresponding brain functional patterns (Tagliazucchi *et al.*, 2012; Liu *et al.*, 2013). These approaches assume that these time points alone are sufficient to estimate the temporal dependency and to derive the functional connectivity patterns observed in functional connectivity studies (Tagliazucchi *et al.*, 2011; Tagliazucchi *et al.*, 2016; Cifre *et al.*, 2020). As such, event-based studies shift away from typical functional connectivity and mainly focus on capturing functional patterns as (co-)activation patterns of sparse EPTs using first-order statistics and signal amplitude rather than statistical dependence between them (Tagliazucchi *et al.*, 2011; Tagliazucchi *et al.*, 2012; Petridou *et al.*, 2013). Event detection approaches have shown direct correspondence between such spatial co-activation patterns and large-scale brain networks obtained from functional connectivity analysis (Tagliazucchi *et al.*, 2012; Liu *et al.*, 2013). They also demonstrated that different EPTs of a given node represent different co-activation patterns reflecting dynamic information of rsfMRI data (Liu and Duyn, 2013; Karahanoğlu and Van De Ville, 2015). However, these findings were limited to large-scale distributed brain networks with no established observations of fine-grained networks (ICNs) obtained from high-order ICA. This, in turn, may indicate that event detection approaches are limited to dominant large covarying functional patterns.

After years of using first-order statistics in event-based studies, which followed early works utilizing the temporal dependency between extreme time points to derive the functional connectivity (Tagliazucchi *et al.*, 2011; Tagliazucchi *et al.*, 2012), recent works revisited the idea of using extreme time points to study (dynamic) functional connectivity in the context of second and higher-order statistics and

demonstrated the potential to provide additional insights into cognition and behavior (Tagliazucchi *et al.*, 2016; Zamani Esfahlani *et al.*, 2020; Sporns *et al.*, 2021).

These intriguing findings, along with the potential clinical and behavioral relevance of EPTs (Chen *et al.*, 2015; Bolton *et al.*, 2020b; Gaviria *et al.*, 2021) and a number of supporting observations in animal studies (Liang *et al.*, 2015; Matsui *et al.*, 2016, 2019), further buttress the assumption that rsfMRI signal may originate from brief, sporadically events rather than continuous dynamic functional interactions. This has led to a growing interest in studying brain function using momentarily sparse time points (Bolton *et al.*, 2020a; Zhang *et al.*, 2020).

However, an important question that has not yet been addressed is whether estimating results from only EPTs omits important information, or worse, provides misleading conclusions. In this work, we argue against the proposition that functional patterns estimated from the spontaneous blood oxygenation level-dependent (BOLD) signal exist only in temporally sparse burst moments and these discrete EPTs contain incomplete information about the underlying functional patterns. This position can be supported by two specific observations. First, it has been shown that spiking activity propagates using both asynchronous and synchronous spiking (Kumar *et al.*, 2010). It is unlikely that sparse events alone can completely capture both of these modes. Second, the fact that connectivity estimators based on both amplitude (e.g., Pearson correlation) and phase (e.g., instantaneous phase synchrony) modulation that have been shown to contain meaningful information (Allen *et al.*, 2011; Omidvarnia *et al.*, 2016) suggests that sparse events based only on amplitude in the space-time domain may give incomplete results.

We evaluated our premise by concentrating on the default mode as it is arguably the most detected/studied functional pattern in event-based studies and is commonly used to support the discrete

nature of functional patterns in the spontaneous BOLD signal. We first asked whether the default mode pattern is present only in its so-called EPTs or whether it can also be identified from the event absent time points (EATs), i.e., time points with the least (almost zero) probability of being EPTs. Additionally, we explored whether the default mode EPTs contain similar functional connectivity information as the full data in the context of their associations with a diagnosis of schizophrenia and with schizophrenia symptoms. Finally, we asked whether the (usually ignored) EATs of the default mode could provide additional unique information about schizophrenia.

It should be mentioned that while the term "co-activation pattern/map" is commonly used to describe spatial patterns identified by event-detection approaches, the term co-activation pattern (CAP) also refers to a specific category of event detection approaches (Liu *et al.*, 2018), and the term co-activation has also been used in other contexts, such as describing task co-activation patterns across fMRI studies (Toro *et al.*, 2008; Eickhoff *et al.*, 2011). Therefore, we opted for the term activation spatial maps (ASMs) to describe the first-order activation patterns that are being evaluated in this study to avoid confusion. Furthermore, to simplify notation in the remainder of this paper, unless we explicitly indicate otherwise, the terms EATs and EPTs refer to the EATs and EPTs of the default mode.

Materials and Methods

Dataset and Inclusion Criteria: We used 3-Tesla resting-state functional magnetic resonance imaging (rsfMRI) data, comprising 744 subjects selected from three datasets with different data acquisition parameters (detailed in (Iraji *et al.*, 2021)), including Functional Imaging Biomedical Informatics Research Network (FBIRN), Center for Biomedical Research Excellence (COBRE), and Maryland Psychiatric Research Center (MPRC). The inclusion criteria include 1) the number of time points (volumes) of the BOLD time series larger than 100 (the minimum number of time points is 135), 2) head motion transition less than 3º rotations and 3 mm translations in every direction, 3) mean framewise

displacement (mFD) less than 0.25, 4) high-quality registration to an echo-planar imaging (EPI)

template, and 5) spatial overlap between individual mask and group mask above 80%.

Preprocessing: The rsfMRI data were preprocessed primarily using the statistical parametric mapping

(SPM12, http://www.fil.ion.ucl.ac.uk/spm/) toolbox. The first five volumes were discarded for

magnetization equilibrium purposes. Rigid body motion correction was performed to correct subject

head motion during the scan. Slice-timing correction was applied to correct timing differences (temporal

misalignment) in slice acquisition. The data of each subject were warped into a Montreal Neurological

Institute (MNI) echo-planar imaging (EPI) template and resampled to 3 mm^3 isotropic voxels. Next, the

data were spatially smoothed using a Gaussian kernel with a 6 mm full width at half-maximum

(FWHM).

Because the fMRI signal has a low signal-to-noise ratio (SNR), the detection of EPTs can be sensitive to

noise. For example, noise can change the time points that survive thresholding and alter the time points

that manifest as local maxima. To address this, we performed the following cleaning steps on the time

course of each voxel. Voxel time courses were detrended by removing linear, quadratic, and cubic

trends. We regressed out six motion realignment parameters and their derivatives. Outliers were detected

based on the median absolute deviation, spike threshold $(c_1) = 2.5$ (Jo *et al.*, 2013), and replaced with the

best estimate using a third-order spline fit to the clean portions of the time courses. Bandpass filtering

was applied using a fifth-order Butterworth filter with a cutoff frequency of 0.01 Hz-0.15 Hz. We also

evaluated if the cleaning procedure drives the findings by repeating the analysis using uncleaned data.

Extracting Default Mode Time Course: We used several procedures to obtain the time series associated

with the default mode network to illustrate that our findings are not the result of a selected node or time

courses. Similar to most event-based studies, we first used a region of interest (ROI) in the posterior

cingulate cortex (PCC) area as a node for the default mode. For this purpose, we used the two most

commonly used ROIs including a $6 \times 6 \times 6$ mm^3 cube centered at (x = 0, y = −53, z = 26) (Liu and

Duyn, 2013) and a sphere with 3mm-radius, centered at (x = −6, y = −58, z = 28)) (Chang and Glover,

2010) in MNI coordinates. We called these two $Node_{Seed1}$ and $Node_{Seed2}$. We defined an additional node

($Node_{Meta}$) using a term-based meta-analysis for the term "default mode" in Neurosynth

(https://www.neurosynth.org/) (Yarkoni *et al.*, 2011). The default mode associated map was registered

to the study common space, and the top 27 voxels were selected as $Node_{Meta}$ for the default mode. The

$Node_{Meta}$ includes a cluster of size 20 voxel in the PCC and a cluster of size seven voxel in the angular

gyrus. For these three nodes, the average time course within each node was used as the representation of

the default mode time course to identify time points with the highest and lowest contributions of the

default mode.

While existing event-based studies use anatomically fixed nodes to estimate EPTs, one may argue our

findings are the result of using predefined anatomically fixed nodes for all individuals, which could

result in inaccurate estimation of the default mode time course. As such, we also utilized ICA to obtain

subject-specific default mode maps and associated time courses ($Node_{ICA}$) (Calhoun *et al.*, 2001;

Calhoun and Adalı, 2012). ICA was performed using the GIFT v4.0c software package

(https://trendscenter.org/software/gift/) (Iraji *et al.*, 2020a). We applied a low model order (model order

= 20) group-level spatial ICA to obtain large-scale brain networks (Iraji *et al.*, 2016), including the

default mode. We applied variance normalization (z-score) on voxel time courses and computed subject-

level spatial principal components analysis (PCA) to retain maximum subject-level variance (greater

than 99%). Subject-level principal components were then concatenated together across the time

dimension, and group-level spatial PCA was applied to the concatenated subject-level principal

components. The 20 group-level principal components that explained the maximum variance were

selected as the input for the Infomax algorithm to calculate 20 group independent components. The Infomax ICA algorithm was run 20 times, and only components with (ICASSO cluster quality index > 0.8) were used to identify networks. Next, we used the group component as a reference in a spatially constrained ICA (called group information guided ICA or GIG-ICA) to calculate subject-specific default mode and its associated time courses (Du *et al.*, 2015).

Detecting Event Present and Absent Time Points (EPTs and EATs) of the Default Mode: The main idea behind event-based studies is that functional connectivity patterns, such as the default mode, are derived from sparse, abrupt neuronal events which manifest as momentarily changes in the amplitude (or other characteristics) of the BOLD signal of associated regions. Thus, the key step for all event-based studies is to identify a set of time points with the largest association to a predefined node (i.e., EPTs). Here, to evaluate our hypothesis, we also focus on the set of time points with the weakest association to the predefined node (i.e., EATs), which are at the opposite extreme, the time points with the minimum chance of event occurrence. Event-based studies use different techniques to detect EPTs from the time series of a given node, which are suggested to lead to similar results (Cifre *et al.*, 2020). The most common techniques include 1) choosing the time points with maximum amplitudes (e.g., top 10 percent or top 20 percent time points), 2) choosing time points that pass a threshold value (e.g., above one standard deviation of the time series), 3) selecting time points which are the local maxima/minima of the time series, and 4) deconvolving the BOLD signal using hemodynamic models, which resolve into a set of sparse EPTs.

Focusing on the default mode, our main objective here is not to identify the default mode and its EPTs, but rather to assess the possibility of estimating the default mode from its assumed EATs, and secondarily whether we are losing information if we focus only on the EPTs. In other words, we first aim to evaluate if the default mode can be retrieved by only using EATs. We can use any event detection

techniques and assign time points not selected as EPTs as EATs, but here we chose a more extreme

approach and defined EATs as time points that are the least likely EPTs of the default mode. For this

purpose, we selected the 20 time points (equal to the ICA model order) with the minimum amplitude of

a given node time series as EATs (**Figure 1**). In the same manner, the 20 time points with the largest

amplitude are considered to be the EPTs (**Figure 1**). We also evaluate the finding by choosing the 20

time points with the highest absolute value.

7
Figure 1. Schematic of the analysis pipeline. First, we select one of the four default mode nodes and use its time course to
identify event present time points (EPTs) and event absent time points (EATs) for the default mode. The 20 time points with
the largest amplitude are labeled as the EPTs (blue dots), while EATs are the 20 time points with minimum amplitudes (red
dots). We selected this time point selection technique to minimize the likelihood of the presence of an event in EATs. We
also repeated the analysis by using the amplitude of absolute value (as some studies use both positive and negative extremes
as EPTs). We compared EPTs and EATs with the functional patterns obtain using all time points (AllTPs, green dots). We
evaluated the presence of functional patterns in EATs, EPTs, and AllTPs of default mode (DM) using different tools,
including high-order statistics using independent component analysis (ICA), presented in the figure, second-order statistics
using pair-wise Pearson correlation, and first-order statistics using (weighted-) activation spatial maps (ASMs). We were
specifically interested in whether (1) the DM is present within the EATs (magenta box) and (2) EPTs of the DM also include
information about the non-DM networks (cyan box). aDM: anterior default mode network (as opposed to classical (posterior)
DM), ATN: attention network, LFP/RFP: left/right frontoparietal network, SUB: subcortical network, MTR: Somatomotor
network, AUD: auditory network, CER: cerebellar network, and VIS: visual network.

In addition, we used EPTs to investigate if other large-scale networks can be obtained by applying ICA

to the EPTs of the default mode (**Figure 1**). In other words, we evaluated if the default mode EPTs are

only occupied or mainly dominated by default mode or other networks similarly present and contribute

to these time points. While some recent event-based works consider the possibility of the coexistence of

a few functional patterns and allow the temporal overlap between their events (Karahanoğlu and Van De

Ville, 2015; Zoller et al., 2019), event-based studies commonly assume the activation peaks of different

networks do not coincide together. Here, we argue for the continuous presence of all networks including

EPTs of other networks and assess this by focusing on EPTs of the default mode.

Evaluating Event Present and Absent Time Points (EPTs and EATs): We used the following

procedures to evaluate the presence of the large-scale networks in the EATs (EPTs) of the default mode

estimated using the various nodes mentioned earlier. First, we applied ICA to the EATs (EPTs) of the

default mode to evaluate if we can obtain brain networks from these time points. We used ICA with the

same parameters as explained above. However, instead of using all time points, we only used the EATs

(EPTs) of each subject for this procedure. We also calculated whole-brain functional connectivity maps

by calculating Pearson correlation between a node time series and all brain voxels using only EATs

(EPTs). The main objective of this procedure was to determine if the default mode can be obtained from

EATs using linear second-order statistics or if high-order statistics (for example, using ICA) are required

to extract the network.

Finally, we calculated whole-brain ASMs to evaluate if the co-activation pattern of the default mode can

be estimated from EATs in addition to EPTs. First, we calculated the weighted-ASM (wASM) (Eq. 1),

where weight at each time point, $w(t)$, is node signal amplitude and $X(t)$ is the preprocessed BOLD

signal at a given time point.

$$wASM_{voxel \times 1} = \frac{1}{T} \sum_{t=1}^{T} w(t) * X_{voxel \times 1}(t) \quad Eq. 1$$

We also calculated ASM by averaging EATs (EPTs) while correcting the sign of node signal at a given

time point (Eq. 2). The sign correction is needed to prevent positive and negative values from canceling

each other. When EPTs are obtained using the amplitude of a given node time course, EPTs are

expected to have positive values, and Eq. 2 is equal to the simple averaging. However, many studies use

the maximum absolute values of a node time course or L^1-norm (or L^2-norm) of time series obtained

from several node amplitudes at any given time, which requires sign correction.

$$ASM_{voxel \times 1} = \frac{1}{T} \sum_{t=1}^{T} sign\big(w(t)\big) * X_{voxel \times 1}(t) \quad Eq.2$$

Recursive ICA: When we use GIG-ICA to obtain a subject-specific node for the default mode

(Node$_{ICA}$), EATs are estimated from subject-specific default mode time courses. Thereby, the EATs are

largely dominated by other networks, and typically it is assumed there would be no significant presence

of default mode within these timepoints. However, we hypothesized that even in this scenario, the

default mode network is present and can be recovered. To evaluate our hypothesis, we introduce an

approach called recursive ICA, which is simply performing ICA on the residual of the previous ICA. For

this purpose, we first estimate ICA components from the EATs and obtained the stable components.

Then, we regress out these components from each subject EATs and estimate components from the

residual.

Event Present and Event Absent Time Point (EPT and EAT) Signatures in Schizophrenia: We next

investigated whether default mode EPTs carry all information about the default mode or if instead

different temporal portions, EPTs, EATs, and all time points (AllTPs), contain distinct and potentially

complementary information about the default mode. We used Node$_{ICA}$ as it provides the best estimation

of the default mode EPTs and EATs from data.

We first evaluated the association between a schizophrenia symptom score and the default mode

functional network connectivity (DM-FNC). Because different symptom scores, the Positive and

Negative Syndrome Scale (PANSS) and the Brief Psychiatric Rating Scale (BPRS), were recorded for different individuals with schizophrenia, we used a prior conversion algorithm obtained from 3767 individuals to convert BPRS total scores to PANSS total scores (Leucht *et al.*, 2013) and used PANSS total score in our analysis. We calculated FNC between the default mode network and nine other large-scale networks for EPTs, EATs, and AllTPs (aka, static DM-FNC). We regressed out age, gender, site, and mean framewise displacement (mFD) from each FNC. For each approach, i.e., AllTPs, EPTs, and EATs, we used nine DM-FNCs as the independent variables and the total PANSS score as the dependent variable. We applied the least absolute shrinkage and selection operator (LASSO) with ten-fold cross-validations (CV = 10) and 50 Monte Carlo repetitions for cross-validation to identify the symptom-related DM-FNC features. We chose the model and the regularization coefficient (lambda) that resulted in the minimum cross-validated mean squared error (MSE). The DM-FNCs with nonzero coefficients are the most relevant attributes, among the independent variables in the model, to best describe the PANSS score in our dataset. If all coefficients are zero for a model, it means none of the FNC pairs significantly (i.e., beyond constant value) relate to the symptom severity score. Next, we evaluated if symptom-related DM-FNCs (DM-FNCs with nonzero coefficients, if there are any), obtained using individuals with schizophrenia, can also differentiate between individuals with schizophrenia and typical controls using a logistic regression model. Similar to the previous analysis, the logistic regression was applied after regressing out age, gender, site, and mFD covariates.

Results

The Default Mode Never Rests: The results for using the amplitude of Node$_{Seed1}$ time series to obtain the default mode from EATs and EPTs are presented in **Figure 2**. The ICA results show that the default mode network can be estimated well from just the EATs. We also performed an additional test by removing local peaks from EATs and applying ICA on the remaining EATs, which shows the presence

of the default mode network. The spatial similarity between the default mode obtained from EATs (EPTs) and that using all time points is $\rho = 0.84$ (0.97). As expected, the spatial patterns of the default mode in EATs demonstrated meaningful differences. For example, the location of $Node_{Seed1}$ itself does not strongly contribute to the default mode network in these EATs, which explains the lower spatial similarity in EATs. In other words, the default mode exists strongly in $Node_{Seed1}$ EATs, but the contribution of $Node_{Seed1}$ to the default mode in EATs is not as strong as the rest of the default mode regions. The fact that the default mode can still be estimated from the EATs, where it should be least likely to be detected, provides evidence that intrinsic networks are best modeled as continuous in both space and time, rather than as discrete events.

We next studied the connectivity profile of $Node_{Seed1}$ in the EATs to evaluate if $Node_{Seed1}$ is completely dissociated from the default mode or, despite not being a part of the default mode network dominant pattern, $Node_{Seed1}$ maintains its connection to the default mode regions. Interestingly, even in the EATs, $Node_{Seed1}$ is significantly connected to the default mode regions, and the default mode connections are the dominant functional connectivity of $Node_{Seed1}$ even in its EATs (**Figure 2**). The activity patterns further supported the functional connectivity results in that both ASMs and weighted ASMs (wASMs) show co-activation patterns resembling the default mode network for AllTPs, ETPs, and EATs. The connectivity and co-activity patterns of $Node_{Seed1}$ in EATs are somewhat noisier because of the small amplitude and fluctuations of $Node_{Seed1}$ time series in these time points; however, the default mode pattern is still clearly present.

Performing analysis using the absolute value of $Node_{Seed1}$ time series resulted in similar findings as using the signed amplitude of $Node_{Seed1}$ time series (Supplementary Materials). The findings were replicated when other nodes were used to identify EPTs and EATs. **Figure 3** (and Supplementary Materials) shows the results for $Node_{Seed2}$ and $Node_{Meta}$ when the absolute value (amplitude) of their

time courses were used to extract EATs and EPTs. These results are similar to those for $Node_{Seed1}$,

suggesting our findings are not the results of specific node or event selection procedures.

ICA analysis also reveals an interesting finding regarding the dynamic interplay between networks,

consistent across experiments (i.e., using different nodes and event detection techniques). In EATs,

many regions of the executive control network (identified as the cyan arrows in **Figure 2**) strongly

contribute to the default mode component, which appears as control and default mode networks to be

emerging as one component. These two networks have been repeatedly reported as two separate

networks (Beckmann *et al.*, 2005; Smith *et al.*, 2009), as we also observed when running the same

decomposition (ICA with model order 20) for EPTs and also for using all time points.

10
Figure 2. $Node_{Seed1}$ default mode functional patterns. The plots are thresholded at *z*-value = 1. The first column shows the
functional patterns obtained using all time points (AllTPs) of individuals. The second column represents the results obtained
by applying analysis on the event present time points (EPTs), and the third column is the functional patterns obtained using
only event absent time points (EATs). The first row is showing the ICA results. The top left shows the default mode network
calculated by applying ICA on the whole dataset (using all time points). The default mode network was successfully obtained
for both EPTs and EATs. For EATs, $Node_{Seed1}$, which is in the posterior cingulate area (PCC), does not strongly contribute to
the default mode network (see the 'hole' in seed location in the top right panel. However, even in EATs, the $Node_{Seed1}$
connectivity with the default mode regions is stronger than its connectivity with the rest of the brain as shown in the
functional connectivity map (FCM) of $Node_{Seed1}$ obtained using pair-wise Pearson correlation (the second row). In other
words, this result suggests the default mode network exists strongly at EAT time points, and $Node_{Seed1}$ is functionally
connected to it. Activation spatial maps (ASMs) and weighted ASMs (wASMs) show very similar co-activation patterns for
AllTPs, ETPs, and EATs consist of default mode regions. Indeed, the EATs maps are noisier because the small amplitude of

Node$_{Seed1}$ time course in EATs making it more susceptible to noise. The arrows in cyan color show regions commonly
contribute to the executive control network.

3
Figure 3. The default mode functional patterns were obtained using the absolute value of Node$_{Seed2}$ (a) and Node$_{Meta}$ (b) time
courses. The plots are thresholded at z-value = 1. The first, second, and third columns illustrate the results obtained using all
time points (AllTPs), using only the event present time points (EPTs), and using only event absent time points (EATs),
respectively. Node$_{Seed2}$ and Node$_{Meta}$ results reproduce the findings obtained from Node$_{Seed1}$ (see Figure 1).

Next, we evaluated the presence of the default mode in EATs (and EPTs) identified from the time course

of the subject-specific default mode network. As we predicted, because these EATs were obtained from

the same subject's data, the default mode network activity was weak and concealed by the strong

activity of other sources (such as other networks and artifacts). By using recursive ICA and removing

the contribution of dominant components from EATs first, we successfully estimated the default mode

network from the EATs alone (**Figure 4**). We also observed significant functional connectivity and co-

activation in EATs between the time course of the default mode network and other regions involved in

the default mode (**Figure 4**). The result confirmed the previous findings using other nodes, suggesting

the default mode and functional connectivity between associated regions consistently present in

spontaneous BOLD signals. Results were also consistent when we did not perform cleaning steps on the

voxel time course.

8
Figure 4. The default mode functional patterns were obtained using the amplitude of $Node_{ICA}$. $Node_{ICA}$ is subject-specific
default mode network obtain using group-information guided independent component analysis (GIG-ICA) (Du *et al.*, 2015).
The plots are thresholded at z-value = 1. The first, second, and third columns illustrate the results obtained using all time
points (AllTPs), using only the event present time points (EPTs), and using only event absent time points (EATs),
respectively. * indicates the default mode network obtain using recursive ICA.

The Default Mode Event Time Points are also Occupied by Other Networks: Next, we evaluated the

presence of large-scale networks obtained in static analysis using all time points in EPTs and EATs. For

all investigated nodes (i.e., $Node_{Seed1}$, $Node_{Seed2}$, $Node_{Meta}$, and $Node_{ICA}$), we successfully obtained the

large-scale networks for both EPTs and EATs, suggesting that the large-scale networks present in the

data regardless of the strength of the default mode and associated regions. **Figure 5** shows the large-

scale networks obtained using the amplitude value of $Node_{Meta}$ (i.e., Neurosynth default mode associated map) time course. The large-scale networks for other nodes and conditions can be found in Supplementary Materials.

Figure 5. The large-scale networks calculate using the amplitude value of Node$_{Meta}$ time courses. The plots are thresholded at
z-value = 1. The first, second, and third columns illustrate the results obtained using all time points (AllTPs), using only the
event present time points (EPTs), and using only event absent time points (EATs), respectively. This result shows EPTs are
not solely dominated by the default mode network, and the large-scale networks are equally present in EPTs. We also retrieve
large-scale networks from EATs, suggesting the networks consistently present in the BOLD signal. DM: default mode
network (also known as posterior DM). aDM: anterior default mode network (as opposed to classical (posterior) DM), ATN:
attention network, LFP/RFP: left/right frontoparietal network, SUB: subcortical network, MTR: Somatomotor network,
AUD: auditory network, CER: cerebellar network, and VIS: visual network.

Event Present and Event Absent Time Point (EPT and EAT) Unique Signatures in Schizophrenia:

We next studied the DM-FNC obtained from different portions (EPTs, EATs, and AllTPs) of data in the

context of their associations with schizophrenia and schizophrenia symptoms. **Figure 6** shows the

minimum cross-validated mean squared error results for generalized linear models obtained using

LASSO with ten-fold cross-validations and 50 repetitions, where the response variable (dependent

variable) is the total PANSS score and the predictor variables (independent variables) are nine DM-

FNCs obtained using AllTPs, EPTs, or EATs. The lines in **Figure 6** show the symptom-related DM-

FNCs for AllTPs model in red, EPTs model in blue, and EATs model in green, and the width of each

line represents the magnitude of the coefficients. The connection between the cerebellum and the default

mode (CER-DM) contributes to all three models, suggesting its steady effect over time. However, it has

much smaller coefficient in EPTs (|b|= 0.61) compared to AllTPs (|b| = 2.14) and EATs (|b| = 2.81).

We also observed unique symptom-related patterns across models. For example, the AUD-DM

connection contributes only to AllTPs model (|b| = 0.58), the RPF-DM connection contributes only to

EPTs model (|b| = 0.28), and the LPF-DM connection contributes only to EATs model (|b| = 4.54). Post-

hoc analyses were applied to explore the findings. We observed that the weights of symptom-related

DM-FNCs (1) are almost identical across LASSO repetitions and (2) did not show significant

differences for regression model applied across different subsets of data. Moreover, we calculate the

strength of association with the PANSS total score by carrying out the final regressions on different

subsets of data using predictor variables identified from each subset. Among three models, EATs,

despite having fewer predictor variables (two versus three), showed the highest correlation value (|r| =

0.191) with PANSS score compared to EPTs (|r| = 0.143) and AllTPs (|r| = 0.125) for their own

predictors (the blue cells in Table 1). Moreover, EATs also show the highest correlation value for using

the predictor variables obtained from AllTPs and EPTs (the last row in Table 1).

5
6   **Table 1.** Applying regression on different subsets (EATs, EPTs, AllTPs), using the predictor variables identified for each subset. AllTPs: all time points, EPTs: event present time points, and EATs: event absent time points.

The subset of data used to obtain predictor variables	AllTPs	EPTs	EATs
Number of predictors	3	3	2
AllTPs: correlation value (|r|) /p-value	0.125/0.35	0.121/0.38	0.100/0.35
EPTs: correlation value (|r|) /p-value	0.130/0.32)	0.143/0.23	0.086/0.47
EATs: correlation value (|r|) /p-value	**0.131/0.31)**	**0.156/0.17**	**0.191/0.02**

7   Additionally, we found the symptom-related DM-FNCs differentiate (FDR < 0.05) between

8   schizophrenia and typical controls in all three AllTPs, EPTs, and EATs models, further supporting their

9   biological relevance. These results suggest that different portions of data, including EATs, carry

10   unique/additional information about the default mode, and a focus only on EPTs may miss this

11   important information.

Figure 6. The association between schizophrenia symptoms and the default mode functional network connectivity (DM-FNC). The lines represent the coefficients of the generalized linear model obtained using the least absolute shrinkage and selection operator (LASSO) with ten cross-validations (CV = 10) and 50 Monte Carlo repetitions for the model with the minimum cross-validated mean squared error (MSE), where the response variable is the total of positive and negative syndrome scale (PANSS) scores and the predictor variables are nine default mode-FNC pairs estimated using all time points (AllTPs) in red, using the default mode event present time points (EPTs) in blue, and using the default mode event absent time points (EATs) in green. The width of each line represents the magnitude of the coefficients. aDM: anterior default mode network (as opposed to classical (posterior) DM), ATN: attention network, LFP/RFP: left/right frontoparietal network, SUB: subcortical network, MTR: Somatomotor network, AUD: auditory network, CER: cerebellar network, and VIS: visual network.

Discussion and Perspective

The main objective of functional neuroimaging is to relate the measurements of imaging modalities to underlying neural activity and biological variability. Among different imaging modalities, fMRI, which measures the BOLD signal as a proxy for average neural activity, has advantages over others because of its ability to record data from the whole brain, arguably the best trade-off between spatial and temporal resolutions. However, there is still little known about the underlying functional architecture and how neural activity contributes to the spontaneous BOLD signal. To decipher this mystery, the neuroimaging

community has dedicated great effort to develop analytical approaches to model spontaneous BOLD signal and interpret the underlying functional patterns.

Building upon the notion of dynamic, ongoing functional interactions across the brain, the dominant category of approaches model intrinsic functional patterns (i.e., functional networks) as continuous entities in time. A major departure from this category seeks to model brain function as a set of discrete (i.e., sparse in time) functional patterns, initially developed based on the hypothesis that spontaneous BOLD signal results from brief and isolated epochs of neural events (Chialvo, 2010). Because detecting event present time points is the core component and the real differentiator of this category, we used the term "event detection approaches" as the general term to describe this category of approaches. Similar to the main category, event detection studies have presented intriguing findings, suggesting their potential to facilitate understanding of underlying neural activities with great opportunities for both clinical and research settings. These results in a rapidly growing interest in event detection approaches and using only a subset of data (EPTs), instead of all available time points to study brain function.

This study argues that intrinsic networks have an ongoing, continuous presence in BOLD signal, and EPTs do not contain all the information of their corresponding functional patterns. As such, using only EPTs may result in missing important information, or worse, providing misleading conclusions. The findings of this study support this. Thus, while we emphasize the potential of event detection approaches and advocate for further studies of this category, we call for caution on the assumptions and interpretation of the findings.

We summarize the findings of this study in three main analyses. The first analysis stemmed from the premise that network patterns derived from discrete, temporally sparse events, compared to continuous ongoing interactions between associated regions. We focused on the default mode pattern because it has

been repeatedly detected as a dominant global co-activation pattern and arguably most studied in ROI-based event-driven studies. We evaluated the continuous presence of the default mode in spontaneous BOLD signal by investigating if the default mode significantly exists in the time points with the least probability of being the default mode event (EATs), i.e., the time points with amplitudes near to baseline (zero). We tested our proposition using two commonly used default mode nodes in event-based studies, using a node obtained from meta-analysis for the term "default mode" in Neurosynth, and even using the time course of subject-specific default mode obtained using group ICA followed by multi-objective optimization (Du *et al.*, 2015). We observed that regardless of choice of node, preprocessing steps (cleaning), analysis (first, second, and higher-order statistics), the default mode patterns strongly present in both activity and connectivity spaces. These findings suggest that the default mode is a continuous phenomenon or at least has a continuous footprint in spontaneous BOLD signal. Therefore, EPTs do not contain the complete information of functional patterns and are insufficient to model their properties. Considering that the true nature of the underlying neuronal activity is still unknown, focusing only on EPTs may obscure our understanding of brain function. The success of event detection approaches in identifying brain networks and capturing useful information using only EPTs should not be seen as evidence that networks only present or originated from discrete infrequent (Liu *et al.*, 2018).

While the main objective of the first analysis was to evaluate the presence of the default mode at EATs of the default mode, the second analysis targeted its EPTs. We were particularly interested in whether spontaneous BOLD signal in these time points is induced by default mode events and merely reflects the default mode or subsists on other functional patterns, obtained from the whole data. ICA analysis shows that large-scale brain networks can be equally obtained from EPTs as well as EATs. The results of these analyses provide evidence that intrinsic networks have an ongoing, continuous presence in the BOLD signal, and they are best modeled as continuous in both space and time, rather than as discrete events.

The third analysis was designed to answer whether EPTs can capture a full picture of functional

properties or if different portions of time carry unique/complementary information. First, we studied the

association between a schizophrenia symptom severity score and the DM-FNC calculated using only

EPTs, only EATs, and AllTPs. We observed common patterns such as the contribution of CER-DM

connection across all three models; in addition, robust and unique symptom-related links were found for

all three scenarios. For instance, unlike the EPT model, the AllTPs model identified the AUD-DM

connection to be related to the symptom score suggesting the EPT approach may be missing some

information present in the full-time series. Further supporting this, the LFP-DM connection from the

EATs shows a robust association with symptoms. At the same time, we advocate for using EPTs to

study brain function as they can contain unique information about brain states. We observed that the

RFP-DM connection contributes only to the EPT model. These findings are aligned with previous

studies showing relationships between symptom severity and functional connectivity of regions from

cerebellar, default mode, and left/right frontoparietal networks (Whitfield-Gabrieli *et al.*, 2009;

Woodward *et al.*, 2011; Wang *et al.*, 2015; Chahine *et al.*, 2017; Brady *et al.*, 2019). For instance, Brady

et al. identified the functional connectivity of the bilateral dorsolateral prefrontal regions (regions in LFP

and RFP) with the default mode regions covaried significantly with symptom severity (Brady *et al.*,

2019). In particular, the functional connectivity of the right dorsolateral prefrontal was associated with

negative symptom severity (Brady *et al.*, 2019). We observed DM-RFP FNC is associated with PANSS

total score in the EPTs. The functional connectivity between a default mode node in PCC and left

middle frontal gyrus (regions involved in LFP) and PANSS general symptom is shown to be correlated

(Woodward *et al.*, 2011). Similarly, a DM-LFP FNC association with the PANSS total score was found

in the EAT analysis. LFP dysfunction has been frequently associated with a range of abnormalities in

schizophrenia, correlated with PANSS score, and suggested as a potential endophenotypic marker of

schizophrenia (Chahine *et al.*, 2017). The relationship between the PANSS and the default mode has

also been reported for within-DM functional connectivity (Garrity *et al.*, 2007). Strikingly for all three

models (i.e., AllTPs, EPTs, and EATs), the symptom-related DM-FNCs also differentiated between

schizophrenia and typical controls, supporting their potential biological relevance. Building on these

findings, we propose different portions of time, to be exact EPTs versus EATs versus AllTPs, carry

unique/complementary information, and using EPTs alone is not sufficient to study brain function.

Furthermore, because symptom scales like PANSS are the primary clinical tools to assess psychotic

behavioral disorders, their functional fingerprints can provide a better understanding of schizophrenia-

related brain functional changes. Despite promising findings, the symptom severity scores on the

schizophrenia subjects are unfortunately differ between the two samples (PANSS versus BPRS), which

we attempt to harmonized them using established a prior conversion algorithm obtained to convert

BPRS total scores to PANSS total scores. This, on the other hand, results in very blunt instrument

(PANSS total score) compared to the finely tuned tools. We propose that future studies should dedicate

more efforts to deciphering behavior-functional imaging relationships and delineating the symptom

scale brain functional fingerprint, particularly by leveraging a finer breakdown of severity symptom

scores into positive and negative and disorganized symptoms.

In addition to differences in the context of schizophrenia, our analyses illustrated differences in the

functional patterns of EPTs and EATs, suggesting they may reflect different states of brain functional

architecture. One intriguing finding was observed in the executive control network. The executive

control network has consistently been identified as a separate functional network with anticorrelative

patterns with the default mode in both time and space. The executive control and default mode were also

reported to have opposite responses (activation versus deactivation) during cognitively demanding tasks

(Sridharan *et al.*, 2008). Similar to previous work, these two appeared as two independent components

in our ICA analyses using AllTPs and EPTs (see **Figure 2** and **Figure 3**). However, for EATs obtained

using voxels mainly in the PCC (i.e., $Node_{Seed1}$, $Node_{Seed2}$, and $Node_{Meta}$), we observed these two

networks seem to emerge into one component which contains the same default mode regions as EPTs

and AllTPs and many regions of the executive control network.

Considering the premise about the central role of the PCC in coordination between these intrinsic

connectivity networks and supporting internally-directed cognition (Leech and Sharp, 2014), which is

important during goal-directed tasks, these moments may reflect information integration between these

two networks and modulation of top-down processing (Wen et al., 2013). This finding is also supported

by the recent spatial dynamic observation of dynamic integration and segregation between brain

networks (Iraji et al., 2019a). Iraji et al. show that intrinsic networks, commonly considered separate

entities in previous spatial static studies, transiently merge and separate, reflecting their dynamic

segregation and integration (Iraji et al., 2019a; Iraji et al., 2020b). Our findings suggest that the default

mode and executive networks are fully segregated at EPTs, which is expected as the default mode

activity is maximum and main default mode regions are expected to strongly connect to each other

compared to their functional connections to other regions. However, in the EATs, the functional

connectivity among default mode regions, particularly the PCC, is lower relative to their connections

with other regions and potentially reflects the transfer of information and modulatory interaction

between these two momentarily integrated networks. As such, we argue that focusing only on EPTs

would be limited to dominant within network dynamics and unveil dynamic segregation patterns, while

including EATs would allow us to better capture dynamic integration and spatial fluidity between

networks (Iraji *et al.*, 2019a).

In addition, existing event-based studies are mainly limited to identifying spatial patterns that resemble

large-scale distributed networks obtained in functional connectivity studies. Early event-based studies

have used this similarity to support their hypothesis and assert that functional connectivity results from coactivation in EPTs. We argue that while the activation maps of EPTs successfully identify gross dominant large-scale covarying functional patterns, more spatially local functional connectivity patterns in the data will be missed. In other words, the activity patterns cannot explain the functional patterns obtained across multiple spatial scales, such as fine-grained ICNs obtained in higher-order ICA analysis (Iraji *et al.*, 2019b; Iraji *et al.*, 2021). Moreover, studies have shown that functional connectivity can occur in cross-frequency and at different frequencies within a given network (Chang and Glover, 2010; Allen *et al.*, 2018; Yaesoubi *et al.*, 2020). As a matter of fact, various types of neural activities with distinct spatial and temporal representation may contribute to the BOLD signal (Logothetis, 2002, 2008).

Concluding Remarks and Future Considerations

We propose that the BOLD signal can be best modeled as a combination of processes that occur at different spatial and temporal scales. Leveraging different existing models and analytical approaches and developing new ones can help better understand underlying neuronal processes that contribute to the BOLD signal, leading to better insights into brain function and its alterations in various brain conditions. This includes event detection approaches and other single frame-based techniques (use single time points as the elements of analysis). However, one should consider the impact of noise on these approaches considering the signal-to-noise ratio (SNR) of the BOLD signal.

We also recommend leveraging data-driven approaches instead of predefined anatomical nodes to detect EPTs. When a region is a central part of a functional pattern (e.g., the PCC for the default mode), its high amplitude time points can effectively depict the functional pattern. However, as our results show, the amplitude of a given fixed node does not well represent the activity of the corresponding functional pattern, rather the activity of a given node and its contribution to functional patterns. For instance, while

EPTs obtained from a PCC node extract the default mode, the default mode is also strongly present in its

EATs, but the node's contribution to the default mode is not as strong as the rest of the default mode

regions. This, for example, can be seen in the top right panel of **Figure 2**, which shows the default mode

network obtained by applying ICA on the EDAs of Node$_{Seed1}$, where the node itself appears as a 'hole'

and does not have a strong contribution to the default mode. This is expected given the EATs were

selected to have minimal relationship to Node$_{Seed1}$, but it is striking the even in this extreme case the rest

of the default mode can still be well-estimated.

Related to this, different methods capture different information regarding the association of a region to a

functional pattern. Both ASM and FCM (first and second-order statistics) calculate the contributions of

regions/voxels to their dominated functional patterns without considering their contribution to other

functional patterns. On the other hand, multivariate analyses like ICA calculate the degree of

associations while controlling for the effect of other functional patterns in a model. This can result in

differences in spatial maps of different functional patterns (e.g., **Figure 2** and **Figure 3**). We suggest

that using multivariate data-driven approaches to extract the time course of a functional pattern might

better detect its EPTs. This can be done using fully blind approaches like clustering or ICA (Allen *et al.*,

2011), or hybrid approaches that use spatial constraints to facilitate comparability across analyses while

also adapting to the data to ensure functional coherence (Du *et al.*, 2020).

Future studies should assess our findings using other functional patterns and use other event detection

approaches such as using local maxima and minima, deconvolving the hemodynamic response, or using

other approaches such as instantaneous functional connectivity (second or higher-order statistics) instead

of activity (first-order statistic). Finally, future single frame-based studies, including event detection

approaches, should also explore the benefits of incorporating spatial dynamics (Iraji *et al.*, 2019a; Iraji *et*

al., 2019c; Iraji *et al.*, 2020b).

In sum, functional networks have continuous, evolving temporal patterns with different portions carrying unique/complementary information highlights the necessity of utilizing all time points and opens new possibilities to study brain function.

Acknowledgements

This work was supported by grants from the National Institutes of Health grant numbers 1U24RR021992, 1U24RR025736, R01EB020407, R01MH118695, and P20GM103472 to Dr. Vince D. Calhoun and by grants from the VA Merit I01CX000497 program and VA Senior Research Career Award to Dr. Judith M Ford.

References

Allen, E. A., Damaraju, E., Eichele, T., Wu, L. & Calhoun, V. D. (2018) 'EEG Signatures of Dynamic Functional Network Connectivity States', *Brain Topogr*, **31**(1), pp. 101-116.

Allen, E. A., Erhardt, E. B., Damaraju, E., Gruner, W., Segall, J. M., Silva, R. F., Havlicek, M., Rachakonda, S., Fries, J., Kalyanam, R., Michael, A. M., Caprihan, A., Turner, J. A., Eichele, T., Adelsheim, S., Bryan, A. D., Bustillo, J., Clark, V. P., Feldstein Ewing, S. W., Filbey, F., Ford, C. C., Hutchison, K., Jung, R. E., Kiehl, K. A., Kodituwakku, P., Komesu, Y. M., Mayer, A. R., Pearlson, G. D., Phillips, J. P., Sadek, J. R., Stevens, M., Teuscher, U., Thoma, R. J. & Calhoun, V. D. (2011) 'A baseline for the multivariate comparison of resting-state networks', *Front Syst Neurosci*, **5**, p. 2.

Beckmann, C. F., DeLuca, M., Devlin, J. T. & Smith, S. M. (2005) 'Investigations into resting-state connectivity using independent component analysis', *Philos Trans R Soc Lond B Biol Sci*, **360**(1457), pp. 1001-1013.

Biswal, B., Yetkin, F. Z., Haughton, V. M. & Hyde, J. S. (1995) 'Functional connectivity in the motor cortex of resting human brain using echo-planar MRI', *Magn Reson Med*, **34**(4), pp. 537-541.

Bolton, T. A. W., Tuleasca, C., Wotruba, D., Rey, G., Dhanis, H., Gauthier, B., Delavari, F., Morgenroth, E., Gaviria, J., Blondiaux, E., Smigielski, L. & Van De Ville, D. (2020a) 'TbCAPs: A toolbox for co-activation pattern analysis', *Neuroimage*, **211**, p. 116621.

Bolton, T. A. W., Wotruba, D., Buechler, R., Theodoridou, A., Michels, L., Kollias, S., Rössler, W., Heekeren, K. & Van De Ville, D. (2020b) 'Triple Network Model Dynamically Revisited: Lower Salience Network State Switching in Pre-psychosis', *Front Physiol*, **11**, p. 66.

Brady, R. O., Jr., Gonsalvez, I., Lee, I., Öngür, D., Seidman, L. J., Schmahmann, J. D., Eack, S. M., Keshavan, M. S., Pascual-Leone, A. & Halko, M. A. (2019) 'Cerebellar-Prefrontal Network Connectivity and Negative Symptoms in Schizophrenia', *Am J Psychiatry*, **176**(7), pp. 512-520.

Calhoun, V. D. & Adalı, T. (2012) 'Multisubject independent component analysis of fMRI: a decade of intrinsic networks, default mode, and neurodiagnostic discovery', *IEEE Rev Biomed Eng*, **5**, pp. 60-73.

Calhoun, V. D., Adali, T., Pearlson, G. D. & Pekar, J. J. (2001) 'A method for making group inferences from functional MRI data using independent component analysis', *Hum Brain Mapp*, **14**(3), pp. 140-151.

Calhoun, V. D. & Allen, E. (2013) 'Extracting intrinsic functional networks with feature-based group independent component analysis', *Psychometrika*, **78**(2), pp. 243-259.

Chahine, G., Richter, A., Wolter, S., Goya-Maldonado, R. & Gruber, O. (2017) 'Disruptions in the left frontoparietal network underlie resting state endophenotypic markers in schizophrenia', *Hum Brain Mapp*, **38**(4), pp. 1741-1750.

Chang, C. & Glover, G. H. (2010) 'Time-frequency dynamics of resting-state brain connectivity measured with fMRI', *Neuroimage*, **50**(1), pp. 81-98.

Chen, J. E., Chang, C., Greicius, M. D. & Glover, G. H. (2015) 'Introducing co-activation pattern metrics to quantify spontaneous brain network dynamics', *Neuroimage*, **111**, pp. 476-488.

Chialvo, D. R. (2010) 'Emergent complex neural dynamics', *Nature Physics*, **6**(10), pp. 744-750.

Cifre, I., Zarepour, M., Horovitz, S. G., Cannas, S. A. & Chialvo, D. R. (2020) 'Further results on why a

point process is effective for estimating correlation between brain regions', *Papers in Physics*, **12**, p.

120003.

Damoiseaux, J. S., Rombouts, S. A., Barkhof, F., Scheltens, P., Stam, C. J., Smith, S. M. & Beckmann,

C. F. (2006) 'Consistent resting-state networks across healthy subjects', *Proc Natl Acad Sci U S A*,

103(37), pp. 13848-13853.

Du, Y., Fu, Z., Sui, J., Gao, S., Xing, Y., Lin, D., Salman, M., Abrol, A., Rahaman, M. A., Chen, J.,

Hong, L. E., Kochunov, P., Osuch, E. A. & Calhoun, V. D. (2020) 'NeuroMark: An automated and

adaptive ICA based pipeline to identify reproducible fMRI markers of brain disorders', *Neuroimage*

Clin, **28**, p. 102375.

Du, Y., Pearlson, G. D., Liu, J., Sui, J., Yu, Q., He, H., Castro, E. & Calhoun, V. D. (2015) 'A group

ICA based framework for evaluating resting fMRI markers when disease categories are unclear:

application to schizophrenia, bipolar, and schizoaffective disorders', *Neuroimage*, **122**, pp. 272-280.

Eickhoff, S. B., Bzdok, D., Laird, A. R., Roski, C., Caspers, S., Zilles, K. & Fox, P. T. (2011) 'Co-

activation patterns distinguish cortical modules, their connectivity and functional differentiation',

Neuroimage, **57**(3), pp. 938-949.

Friston, K. J. (2011) 'Functional and effective connectivity: a review', *Brain Connect*, **1**(1), pp. 13-36.

Garrity, A. G., Pearlson, G. D., McKiernan, K., Lloyd, D., Kiehl, K. A. & Calhoun, V. D. (2007)

'Aberrant "default mode" functional connectivity in schizophrenia', *Am J Psychiatry*, **164**(3), pp. 450-

457.

Gaviria, J., Rey, G., Bolton, T., Delgado, J., Van De Ville, D. & Vuilleumier, P. (2021) 'Brain functional

connectivity dynamics at rest in the aftermath of affective and cognitive challenges', *Hum Brain Mapp*,

42(4), pp. 1054-1069.

Iraji, A., Calhoun, V. D., Wiseman, N. M., Davoodi-Bojd, E., Avanaki, M. R. N., Haacke, E. M. & Kou,

Z. (2016) 'The connectivity domain: Analyzing resting state fMRI data using feature-based data-driven

and model-based methods', *Neuroimage*, **134**, pp. 494-507.

Iraji, A., Deramus, T. P., Lewis, N., Yaesoubi, M., Stephen, J. M., Erhardt, E., Belger, A., Ford, J. M.,

McEwen, S., Mathalon, D. H., Mueller, B. A., Pearlson, G. D., Potkin, S. G., Preda, A., Turner, J. A.,

Vaidya, J. G., van Erp, T. G. M. & Calhoun, V. D. (2019a) 'The spatial chronnectome reveals a dynamic

interplay between functional segregation and integration', *Hum Brain Mapp*, **40**(10), pp. 3058-3077.

Iraji, A., Faghiri, A., Fu, Z., Rachakonda, S., Kochunov, P., Belger, A., Ford, J. M., McEwen, S.,

Mathalon, D. H., Mueller, B. A., Pearlson, G. D., Potkin, S. G., Preda, A., Turner, J. A., van Erp, T. G.

M. & Calhoun, V. D. (2021) 'Multi-spatial scale dynamic interactions between functional sources reveal

sex-specific changes in schizophrenia', *Network Neuroscience*, pp. 1-48.

Iraji, A., Faghiri, A., Lewis, N., Fu, Z., DeRamus, T., Qi, S., Rachakonda, S., Du, Y. & Calhoun, V.

(2019b) 'Ultra-high-order ICA: an exploration of highly resolved data-driven representation of intrinsic

connectivity networks (sparse ICNs)', International Society for Optics and Photonics.

Iraji, A., Faghiri, A., Lewis, N., Fu, Z., Rachakonda, S. & Calhoun, V. D. (2020a) 'Tools of the trade:

Estimating time-varying connectivity patterns from fMRI data', *Soc Cogn Affect Neurosci*.

Iraji, A., Fu, Z., Damaraju, E., DeRamus, T. P., Lewis, N., Bustillo, J. R., Lenroot, R. K., Belger, A.,

Ford, J. M., McEwen, S., Mathalon, D. H., Mueller, B. A., Pearlson, G. D., Potkin, S. G., Preda, A.,

Turner, J. A., Vaidya, J. G., van Erp, T. G. M. & Calhoun, V. D. (2019c) 'Spatial dynamics within and

between brain functional domains: A hierarchical approach to study time-varying brain function', *Hum Brain Mapp*, **40**(6), pp. 1969-1986.

Iraji, A., Miller, R., Adali, T. & Calhoun, V. D. (2020b) 'Space: A Missing Piece of the Dynamic Puzzle', *Trends Cogn Sci*, **24**(2), pp. 135-149.

Jo, H. J., Gotts, S. J., Reynolds, R. C., Bandettini, P. A., Martin, A., Cox, R. W. & Saad, Z. S. (2013) 'Effective Preprocessing Procedures Virtually Eliminate Distance-Dependent Motion Artifacts in Resting State FMRI', *Journal of Applied Mathematics*, **2013**, p. 935154.

Karahanoğlu, F. I. & Van De Ville, D. (2015) 'Transient brain activity disentangles fMRI resting-state dynamics in terms of spatially and temporally overlapping networks', *Nat Commun*, **6**, p. 7751.

Kumar, A., Rotter, S. & Aertsen, A. (2010) 'Spiking activity propagation in neuronal networks: reconciling different perspectives on neural coding', *Nat Rev Neurosci*, **11**(9), pp. 615-627.

Leech, R. & Sharp, D. J. (2014) 'The role of the posterior cingulate cortex in cognition and disease', *Brain*, **137**(Pt 1), pp. 12-32.

Leucht, S., Rothe, P., Davis, J. M. & Engel, R. R. (2013) 'Equipercentile linking of the BPRS and the PANSS', *Eur Neuropsychopharmacol*, **23**(8), pp. 956-959.

Liang, Z., Liu, X. & Zhang, N. (2015) 'Dynamic resting state functional connectivity in awake and anesthetized rodents', *Neuroimage*, **104**, pp. 89-99.

Liu, X., Chang, C. & Duyn, J. H. (2013) 'Decomposition of spontaneous brain activity into distinct fMRI co-activation patterns', *Front Syst Neurosci*, **7**, p. 101.

Liu, X. & Duyn, J. H. (2013) 'Time-varying functional network information extracted from brief instances of spontaneous brain activity', *Proc Natl Acad Sci U S A*, **110**(11), pp. 4392-4397.

Liu, X., Zhang, N., Chang, C. & Duyn, J. H. (2018) 'Co-activation patterns in resting-state fMRI signals', *Neuroimage*, **180**(Pt B), pp. 485-494.

Logothetis, N. K. (2002) 'The neural basis of the blood-oxygen-level-dependent functional magnetic resonance imaging signal', *Philos Trans R Soc Lond B Biol Sci*, **357**(1424), pp. 1003-1037.

Logothetis, N. K. (2008) 'What we can do and what we cannot do with fMRI', *Nature*, **453**(7197), pp. 869-878.

Matsui, T., Murakami, T. & Ohki, K. (2016) 'Transient neuronal coactivations embedded in globally propagating waves underlie resting-state functional connectivity', *Proc Natl Acad Sci U S A*, **113**(23), pp. 6556-6561.

Matsui, T., Murakami, T. & Ohki, K. (2019) 'Neuronal Origin of the Temporal Dynamics of Spontaneous BOLD Activity Correlation', *Cereb Cortex*, **29**(4), pp. 1496-1508.

Omidvarnia, A., Pedersen, M., Walz, J. M., Vaughan, D. N., Abbott, D. F. & Jackson, G. D. (2016) 'Dynamic regional phase synchrony (DRePS): An Instantaneous Measure of Local fMRI Connectivity Within Spatially Clustered Brain Areas', *Hum Brain Mapp*, **37**(5), pp. 1970-1985.

Petridou, N., Gaudes, C. C., Dryden, I. L., Francis, S. T. & Gowland, P. A. (2013) 'Periods of rest in fMRI contain individual spontaneous events which are related to slowly fluctuating spontaneous activity', *Hum Brain Mapp*, **34**(6), pp. 1319-1329.

Smith, S. M., Fox, P. T., Miller, K. L., Glahn, D. C., Fox, P. M., Mackay, C. E., Filippini, N., Watkins, K. E., Toro, R., Laird, A. R. & Beckmann, C. F. (2009) 'Correspondence of the brain's functional architecture during activation and rest', *Proc Natl Acad Sci U S A*, **106**(31), pp. 13040-13045.

Sporns, O., Faskowitz, J., Teixeira, A. S., Cutts, S. A. & Betzel, R. F. (2021) 'Dynamic expression of brain functional systems disclosed by fine-scale analysis of edge time series', *Network Neuroscience*, **5**(2), pp. 405-433.

Sridharan, D., Levitin, D. J. & Menon, V. (2008) 'A critical role for the right fronto-insular cortex in

switching between central-executive and default-mode networks', *Proc Natl Acad Sci U S A*, **105**(34),

pp. 12569-12574.

Tagliazucchi, E., Balenzuela, P., Fraiman, D. & Chialvo, D. R. (2012) 'Criticality in large-scale brain

FMRI dynamics unveiled by a novel point process analysis', *Front Physiol*, **3**, p. 15.

Tagliazucchi, E., Balenzuela, P., Fraiman, D., Montoya, P. & Chialvo, D. R. (2011) 'Spontaneous

BOLD event triggered averages for estimating functional connectivity at resting state', *Neuroscience*

Letters, **488**(2), pp. 158-163.

Tagliazucchi, E., Siniatchkin, M., Laufs, H. & Chialvo, D. R. (2016) 'The Voxel-Wise Functional

Connectome Can Be Efficiently Derived from Co-activations in a Sparse Spatio-Temporal Point-

Process', *Front Neurosci*, **10**, p. 381.

Toro, R., Fox, P. T. & Paus, T. (2008) 'Functional coactivation map of the human brain', *Cereb Cortex*,

18(11), pp. 2553-2559.

Wang, H., Zeng, L. L., Chen, Y., Yin, H., Tan, Q. & Hu, D. (2015) 'Evidence of a dissociation pattern in

default mode subnetwork functional connectivity in schizophrenia', *Sci Rep*, **5**, p. 14655.

Wen, X., Liu, Y., Yao, L. & Ding, M. (2013) 'Top-down regulation of default mode activity in spatial

visual attention', *J Neurosci*, **33**(15), pp. 6444-6453.

Whitfield-Gabrieli, S., Thermenos, H. W., Milanovic, S., Tsuang, M. T., Faraone, S. V., McCarley, R.

W., Shenton, M. E., Green, A. I., Nieto-Castanon, A., LaViolette, P., Wojcik, J., Gabrieli, J. D. &

Seidman, L. J. (2009) 'Hyperactivity and hyperconnectivity of the default network in schizophrenia and

in first-degree relatives of persons with schizophrenia', *Proc Natl Acad Sci U S A*, **106**(4), pp. 1279-

1284.

Woodward, N. D., Rogers, B. & Heckers, S. (2011) 'Functional resting-state networks are differentially

affected in schizophrenia', *Schizophrenia Research*, **130**(1-3), pp. 86-93.

Yaesoubi, M., Silva, R. F., Iraji, A. & Calhoun, V. D. (2020) 'Frequency-Aware Summarization of

Resting-State fMRI Data', *Front Syst Neurosci*, **14**, p. 16.

Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C. & Wager, T. D. (2011) 'Large-scale

automated synthesis of human functional neuroimaging data', *Nat Methods*, **8**(8), pp. 665-670.

Zamani Esfahlani, F., Jo, Y., Faskowitz, J., Byrge, L., Kennedy, D. P., Sporns, O. & Betzel, R. F. (2020)

'High-amplitude cofluctuations in cortical activity drive functional connectivity', *Proc Natl Acad Sci U S*

A, **117**(45), pp. 28393-28401.

Zhang, X., Pan, W. J. & Keilholz, S. D. (2020) 'The relationship between BOLD and neural activity

arises from temporally sparse events', *Neuroimage*, **207**, p. 116390.

Zoller, D. M., Bolton, T. A. W., Karahanoglu, F. I., Eliez, S., Schaer, M. & Van De Ville, D. (2019)

'Robust Recovery of Temporal Overlap Between Network Activity Using Transient-Informed Spatio-

Temporal Regression', *IEEE Trans Med Imaging*, **38**(1), pp. 291-302.

1 Supplementary 1. Node$_{Seed1}$ default mode functional patterns. The spatial maps are displayed at their
2 peak activation coordinates. AllTPs: all time points, ETPs: the event present time points, EATs: event
3 absent time points (EATs). ICA: independent component analysis, FCM: functional connectivity map,
4 ASM: Activation spatial map, wASMs: weighted ASMs

5   AllTPs   EPTs   EATs

7 Supplementary 2. Node$_{Seed2}$ default mode functional patterns. The spatial maps are displayed at their
8 peak activation coordinates. AllTPs: all time points, ETPs: the event present time points, EATs: event
9 absent time points (EATs). ICA: independent component analysis, FCM: functional connectivity map,
10 ASM: Activation spatial map, wASMs: weighted ASMs

11   AllTPs   EPTs   EATs

Supplementary 3. Node_{Meta} default mode functional patterns. The spatial maps are displayed at their
peak activation coordinates. AllTPs: all time points, ETPs: the event present time points, EATs: event
absent time points (EATs). ICA: independent component analysis, FCM: functional connectivity map,
ASM: Activation spatial map, wASMs: weighted ASMs

Supplementary 4. The large-scale networks calculate using the absolute value of Node_{Seed1} time courses.
The spatial maps are displayed at their peak activation coordinates. AllTPs: all time points, ETPs: the
event present time points, EATs: event absent time points (EATs). ICA: independent component
analysis, FCM: functional connectivity map, ASM: Activation spatial map, wASMs: weighted ASMs

Supplementary 5. The large-scale networks calculate using the absolute value of $Node_{Seed2}$ time courses. The spatial maps are displayed at their peak activation coordinates. AllTPs: all time points, ETPs: the event present time points, EATs: event absent time points (EATs). ICA: independent component analysis, FCM: functional connectivity map, ASM: Activation spatial map, wASMs: weighted ASMs