INTRINSIC MOTOR NEURONE EXCITABILITY IS REDUCED IN SOLEUS AND TIBIALIS ANTERIOR OF OLDER ADULTS

Running title: Intrinsic motor neurone excitability is reduced with ageing.

Lucas B. R. Orssatto [1*], David N. Borg [2], Anthony J. Blazevich [3], Raphael L. Sakugawa [4], Anthony J. Shield [1], Gabriel S. Trajano [1]

[1] School of Exercise and Nutrition Sciences, Faculty of Health, Queensland University of Technology (QUT), Brisbane, Australia.

[2] Griffith University, Menzies Health Institute Queensland, The Hopkins Centre, Brisbane, Australia.

[3] Centre for Exercise and Sports Science Research (CESSR), School of Medical and Health Sciences, Edith Cowan University, Joondalup, Australia.

[4] Biomechanics Laboratory, Department of Physical Education, Federal University of Santa Catarina, Florianopolis, Brazil.

*Corresponding author: Lucas Bet da Rosa Orssatto; email: l.betdarosaorssatto@qut.edu.au; https://orcid.org/0000-0003-3788-3700. School of Exercise and Nutrition Sciences, Faculty of Health, Queensland University of Technology (QUT), Brisbane, Australia.

ABSTRACT

Age-related deterioration within both motor neurones and monoaminergic systems
should theoretically reduce neuromodulation by weakening motor neuronal persistent inward
current (PIC) strength. However, this assumption remains untested. Surface
electromyographic signals were collected using two 32-channel electrode matrices placed on
soleus and *tibialis anterior* of 25 older adults (70±4years) and 17 young adults (29±5 years)
to investigate motor unit discharge behaviours. Participants performed triangular-shaped
plantar and dorsiflexion contractions to 20% of maximum torque at a rise-decline rate of
2%/s of each participant's maximal torque. Pairwise and composite paired-motor unit
analyses were adopted to calculate delta frequency (ΔF) and estimate PIC amplitudes. ΔF has
been used to differentiate between the effects of synaptic excitation and intrinsic motor
neuronal properties and is assumed to be proportional to PIC amplitude. The results show that
soleus and *tibialis anterior* motor units in older adults had lower ΔFs when calculated with
the pairwise (-0.99 and -1.29 pps, respectively) or composite (-1.65 and -2.26 pps,
respectively) methods. Older adults' motor units discharged at lower rates (-2.14 and -2.03
pps, respectively) and were recruited at lower torque levels (-1.50 and -2.06% of maximum,
respectively) than young adults. These results demonstrate reduced intrinsic motor neurone
excitability during low-force contractions in older adults, likely mediated by decreases in the
strength of persistent inward currents. Our findings might be explained by deterioration in the
motor neurones or monoaminergic systems, and could contribute to the decline in motor
function during ageing; these assumptions should be explicitly tested in future investigations.

Keywords: Persistent inward current; Ageing; Motor unit; HD-EMG; Motoneuron.

KEY POINTS

- The persistent inward current (PIC) is an intrinsic motor neurone excitability property responsible for amplifying excitatory synaptic input into an appropriate motor output. This study compared estimates of PICs in *soleus* and *tibialis anterior* between young and older adults.

- Motor units of older adults presented lower estimates of PICs for both *soleus* and *tibialis anterior*. Older adults also presented lower motor unit discharge rates and they were recruited at lower torque levels than young adults.

- These results indicate a reduced intrinsic motor neurone excitability in soleus and tibialis anterior motor units, likely mediated by decreases in the strength of persistent inward currents.

- The reduced PIC strength in older adults may result from deterioration within both motor neurones and monoaminergic systems.

- These novel findings contribute to our understanding of the factors underpinning the declines in force production and motor function during ageing.

INTRODUCTION

The age-related loss of force production has been comprehensively described in the

literature (Vandervoort, 2002; Orssatto *et al.*, 2018; Suetta *et al.*, 2019), with the

physiological alterations affecting force production including changes in several pathways

within the nervous system (Aagaard *et al.*, 2010; Manini *et al.*, 2013; Orssatto *et al.*, 2018;

Larsson *et al.*, 2019). The motor neurone is an important component of the nervous system

affected by ageing as it is responsible for integrating and amplifying excitatory synaptic input

into an appropriate motor output (Heckman & Enoka, 2012). An essential intrinsic property

of the motor neurone is its capacity to set up persistent inward currents (PICs), which are

depolarising currents generated by voltage-sensitive sodium and calcium channels that

increase cell excitability by amplifying and prolonging synaptic input (Gorassini *et al.*, 2002;

Heckman *et al.*, 2005). Importantly, increases in the concentration of the monoamines

serotonin and noradrenaline facilitate PIC development. Under conditions of high

monoaminergic drive, synaptic input can be amplified by at least five-fold, suggesting that

this amplification is a critical determinant of the motor neurone's ability to achieve the

discharge rates observed during normal motor behaviour (Hounsgaard & Kiehn, 1993; Lee &

Heckman, 1999, 2000). Thus, potential physiological alterations in motor neurone intrinsic

properties, or in the monoaminergic input to the motor neurone, might reduce the motor

neurone's ability to discharge at higher rates, thus reducing the ability to produce high muscle

forces.

During ageing, several changes are observed in both the motor neurones and the

monoaminergic system that might potentially reduce PIC strength in older adults, including

lower discharge rates (Orssatto *et al.*, 2020; Kirk *et al.*, 2021), reduced incidence of doublet

discharges (Christie & Kamen, 2006), and an increased afterhyperpolarisation duration

(Piotrkiewicz *et al.*, 2007). These changes are consistent with the lower motor neurone

excitability that is also observed in aged rat models (Morales *et al.*, 1987). With respect to the

monoaminergic system, research using both human and animal models suggests that ageing is

associated with reduced noradrenaline and serotonin secretions and thus input onto the motor

neurones (Johnson *et al.*, 1993; Ko *et al.*, 1997; Míguez *et al.*, 1999; Shibata *et al.*, 2006; Liu

et al., 2020), which might theoretically underpin PIC reduction with ageing. These findings

indicate the possibility that PIC strength might be reduced in older adults; however, this

hypothesis remains to be tested.

The strength of PICs that are initiated close to the recruitment threshold of motor

neurones can be estimated in humans using the paired motor unit technique (Gorassini *et al.*,

2002; Powers *et al.*, 2008; Stephenson & Maluf, 2011), with data obtained using high-density surface electromyography (Holobar & Zazula, 2007; Enoka, 2019). This technique requires the pairing of the discharge rates of a low-threshold (control unit) to a higher-threshold (test unit) motor unit, obtained during a slowly-increasing and decreasing triangular-shaped contraction (Gorassini *et al.*, 2002; Powers *et al.*, 2008; Orssatto *et al.*, 2021). Subsequently, the difference in discharge rate of the control unit between the time of recruitment and de-recruitment of the test unit is computed as the change in frequency (ΔF). ΔF has been used to differentiate between the effects of synaptic excitation and motor neuronal intrinsic properties and is assumed to be proportional to PIC strength (Afsharipour *et al.*, 2020). With these techniques, it is now possible to estimate and compare the strengths of these PICs in motor units of young and older adults.

The present study compared ΔFs to estimate differences in PIC strength of *soleus* and *tibialis anterior* motor units between young and older adults. Additionally, we explored the relationship between ΔF and the peak discharge rates. We hypothesised that there would be a reduction in PIC in both *soleus* and *tibialis anterior*, and that ΔF would be strongly associated with motor unit peak discharge rate. *Soleus* and *tibialis anterior* were selected for study because the control and timing of their activation are both critical to the performance of daily activities such as standing and walking in older adults (Polcyn *et al.*, 1998; Laughton *et al.*, 2003).

METHODS
Participants and ethical procedures

Forty-four participants volunteered for the study, including 18 young adults and 26 older adults (participant characteristics are documented in Table 1). More older adults were recruited because it was expected that fewer motor units would be identified during decomposition and that some participants may not be able to perform the triangle-shaped contractions with the necessary torque rise and fall accuracy. One participant per group was excluded from the study because no motor units were identified in either *soleus* or *tibialis anterior*. To participate, volunteers had to be free from neurological and lower limb musculoskeletal injuries and disorders. Participants were excluded if they were on medications that could influence the monoaminergic system, such as serotonin or noradrenaline (e.g., beta-blockers and serotonin reuptake inhibitors) or consumed caffeinated foods (e.g., coffee). They were also excluded from the analyses if: a) no usable motor units were identified by the decomposition algorithm, or b) if it was not possible to achieve all the

assumptions required in the paired motor unit analysis (as described below) for either *soleus*

or *tibialis anterior*. The study was approved by the University Human Research Ethics

Committee, and all participants gave written informed consent before participating. Data

collection was conducted during the COVID-19 pandemic and all safety procedures followed

the local state government policies.

Table 1. Participant characteristcs.

Variable	Young Adults (n = 17)	Older Adults (n = 25)
Age (years)	29 ± 5	70 ± 4
Sex (n, %)		
Men	9 (53%)	11 (44%)
Women	8 (47%)	14 (56%)
Body mass (kg)	71.9 ± 13.6	77.8 ± 19.3
Height (cm)	174 ± 11	163 ± 17
Peak torque (N·m)		
Plantar flexion	156 ± 47	85 ± 32
Dorsiflexion	41 ± 14	29 ± 7
Physical activity level (MET min/week)	2541 (1902–3039)	2994 (1386–5139)

Note: Data are presented as mean ± standard deviation, except for Physical activity level,

which is presented as median with interquartile range.

Study design and neuromuscular testing procedures

Participants visited the laboratory on a single occasion in which they were

familiarised with the testing procedures and data were collected. Initially, participants signed

the informed consent and completed the International Physical Activity Questionnaire

(Hagströmer *et al.*, 2006). After electrode placement on *soleus* and *tibialis anterior*, the

participants were seated upright in the chair of an isokinetic dynamometer (Biodex System 4,

Biodex Medical system, Shirley, NY) with the knee fully extended (0°) and ankle in the

anatomical position (0°). A warm-up consisting of six 5-s submaximal voluntary isometric

plantar and dorsiflexion contractions ($2 \times 30\%$, $2 \times 60\%$, and $2 \times 80\%$ of perceived maximal

effort) was performed, followed by three maximal voluntary contractions of ~3-s with 30-s

rest intervals. The maximum torque achieved was recorded as the maximal voluntary

contraction peak torque. Subsequently, participants were familiarised with the triangular-shaped contractions to 20% of their maximum voluntary torque level. Triangular contractions to 20% of maximal torque have been extensively used for ΔF calculations using the paired motor unit technique (Gorassini *et al.*, 2002; Afsharipour *et al.*, 2020; Hassan *et al.*, 2020; Kim *et al.*, 2020; Trajano *et al.*, 2020), and this force was considered similar to the maximum torques developed during daily activities such as standing and walking. All contractions had a duration of 20 s (10-s up and 10-s down) and were performed at a rate of torque increase and decrease of ~2%/s. Participants were instructed to follow the torque path provided in real time on a 58-cm computer monitor during each contraction. Data collection commenced 5 min after the end of familiarisation (usually requiring ~3-10 × 20% triangular contractions with 30-s rest), during which the participants then performed four triangular contractions with 60-s rest intervals. When an abrupt increase or decrease in torque was observed (i.e. the torque trajectory was not closely followed), the trial was excluded and repeated. The maximum voluntary isometric torque and order of triangular contraction completion was randomised between *soleus* and *tibialis anterior*.

Surface electromyography

Surface electromyograms (sEMG) were recorded during the 20% triangular contractions using four semi-disposable 32-channel electrode grids with a 10-mm interelectrode distance (ELSCH032NM6, OTBioelettronica, Torino, Italy). After skin shaving, abrasion, and cleansing with 70% isopropyl alcohol, two electrode grids were placed over the medial and lateral portions of *soleus* (either side of the Achilles tendon) and another two electrode grids were placed over the superior and inferior aspect of *tibialis anterior* using a bi-adhesive foam layer and conductive paste (Ten20, Weaver and Company, Colorado, USA). A strap electrode (WS2, OTBioelettronica, Torino, Italy) was dampened and positioned around the ankle joint as a ground electrode. The sEMG signals were acquired in monopolar mode, amplified (256×), band-pass filtered (10–500 Hz), and converted to a digital signal at 2048 Hz by a 16-bit wireless amplifier (Sessantaquattro, OTBioelettronica, Torino, Italy) using OTBioLab+ software (version 1.3.0., OTBioelettronica, Torino, Italy) before being stored for offline analysis.

Motor unit analyses

Motor unit identification

The recorded data were processed offline using the DEMUSE software (Holobar & Zazula, 2007). For each contraction intensity, only the triangular contraction yielding the lowest deviation from the torque trajectory was analysed. If both contractions presented a similar torque trajectory, the contraction with the highest number of identified motor units was analysed. High-density sEMG signals were band-pass filtered (20-500 Hz) with a second-order, zero-lag Butterworth filter. Thereafter, a blind source separation method, the convolutive kernel compensation (CKC) method, was used for signal decomposition (Holobar & Zazula, 2007; Holobar *et al.*, 2014) from each triangular contraction. CKC yields the filters of individual motor units (so-called motor unit filters) that, when applied to high-density sEMG signals, estimate the motor unit spike trains (Holobar & Zazula, 2007; Holobar *et al.*, 2014). After decomposition, a trained investigator manually inspected motor unit spike trains and edited the discharge patterns of the motor units. Only the motor units with a pulse-to-noise ratio equal to or greater than 30 dB were kept for further analysis (Holobar *et al.*, 2014).

Estimation of PIC amplitude (ΔF) and peak discharge rate

The observed discharge events for each motor unit were converted into instantaneous discharge rates and fitted into a 5^{th}-order polynomial function. The maximum value obtained from the polynomial curve was considered the peak discharge rate. Thereafter, PIC amplitude was estimated using the paired motor unit analysis (Gorassini *et al.*, 2002), referred through the manuscript as pairwise method. Motor units with a low recruitment threshold (i.e., control units) were paired with higher recruitment threshold motor units (i.e., test units). ΔF was calculated as the change in discharge rates of the control motor unit from the moment of recruitment to the moment of de-recruitment of the test unit (Gorassini *et al.*, 2002; Heckman *et al.*, 2005). In order to produce motor unit pairs, the following criteria were adopted: 1) rate-to-rate correlations between the smoothed discharge rate polynomials of the test and control units was $r \geq 0.7$; 2) test units were recruited at least 1.0 s after the control units; and 3) the control unit did not show discharge rate saturation after the moment of test unit recruitment (>0.5 pps) (Gorassini *et al.*, 2002; Udina *et al.*, 2010; Vandenberk & Kalmar, 2014; Binder *et al.*, 2020; Hassan *et al.*, 2020). ΔFs obtained for each pair were averaged to obtain a single ΔF for each motor unit.

We also conducted an additional analysis using the composite paired motor unit method to calculate ΔF values for each motor unit (Afsharipour *et al.*, 2020). This method is characterised by the overlay of 3 lower-threshold motor units to construct a single composite

control unit profile to be paired with the test units. The composite method has been suggested to address some of the limitations observed with the pairwise method, such as underestimation and overestimation of ΔF values, reducing its variability. However, strict assumptions are made for eligible motor units to be included in the analysis, which allows its use only in muscles in which it is possible to identify many motor units (e.g., *tibialis anterior*) but not in those in which fewer motor units are identified using the decomposition method (e.g., *soleus*). Moreover, this method does not allow calculation of ΔF values for lower-threshold motor units since they are used to construct the composite control unit. In summary, the control unit is the overlay of at least three motor units recruited at <3% of the maximum voluntary torque and presenting a similar discharge profile. It was assumed that lower-threshold motor units had their discharge rate profile more linearly related to the synaptic input profile because PICs were almost fully activated at the time of recruitment; therefore, it would be more appropriate to use the composite unit as control. Also, this method requires the removal of the acceleration phase of the discharge rates (i.e., secondary range), making the polynomial ascending-to-descending slope ratio near 1, which is important when measuring PIC amplitudes to avoid any rate-dependent effects on motor unit recruitment or derecruitment (Desmedt & Godaux, 1977). The test units should start discharging in the tertiary range (i.e., after the secondary range and before the descending phase). This method provides a single ΔF for each motor unit, which was then used in the data analysis. Figure 1 illustrates the pairwise and composited paired motor unit analysis methods on *tibialis anterior* motor units for one participant per group. Panels C and D display the test units used for both methods. Panels E and F display the control units for the pairwise method and panels G and H display the control units for the composite method.

3   **Figure 1.** Data from a single participant for each group showing torque during triangle-

4   shaped contractions and delta frequency (ΔF) calculation in *tibialis anterior* for both the

5   pairwise and composite paired motor unit analyses. Data from a young adult is displayed in

6   the left panels and from an older adult in the right panels. Panels A and B on the first row

7   show the torque traces for contractions with 20% of the participant's maximal voluntary

torque. The participants' test units are displayed on panels C and D (purple motor unit), control units for the pairwise method on panels E and F (red motor unit), and control units for the composite method on panels G and H (red, green, and blue motor units). The black continuous lines are the 5^{th}-order polynomial fits for the control units. Note that for the composite method the polynomial curve starts from the tertiary range. The gray-shaded areas represent the ΔF amplitude for each participant and analysis.

Data analysis

All analyses were undertaken in R (version 4.0.3) using RStudio environment (version 1.3.1093). Models were fitted using the *lmerTest* package (Kuznetsova *et al.*, 2017). A linear mixed-effects model was used to compare estimates of ΔF of *soleus* and *tibialis anterior* motor units between young and older adults. The model included: age group, muscle type, recruitment threshold, age group by muscle type, and muscle type by recruitment threshold as fixed effects. A random intercept and slope (recruitment threshold by muscle type) was included for each participant in the study, to account for the correlation between repeated observations on each individual. This model was selected from a series of candidate models (Supplement 1), based on the smallest Bayesian Information Criteria value. The recruitment threshold was standardised (mean = 0, SD = 1) before analysis.

Separate linear mixed-effects models were used to analyse peak discharge rate and recruitment threshold data. These models included: age group, muscle type and age group by muscle type as fixed factors; and a random intercept and slope (muscle group) for each participant. The estimated marginal mean difference and 90% and 95% confidence intervals (CI) in ΔF, peak discharge rate, and recruitment threshold between young and older adults, were determined using the *emmeans* package (Lenth *et al.*, 2021). The standardised difference, denoted *d*, was also calculated using the population SD from each respective linear mixed-effects model as the denominator (Lenth *et al.*, 2021).

The association between ΔF and peak discharge rate was determined by fitting two models: (1) a base model including: age group and muscle type as fixed factors, and a random intercept and slope (muscle type) for each participant, and (2) a second model including all covariates from the base model, in addition to ΔF (standardised; mean 0, SD 1). The difference in the variance explained by the fixed effects (R^2) between the models was calculated (Nakagawa *et al.*, 2017) to isolate the contribution of ΔF. The models were also compared to determine whether there was a reduction in the residual sum of squares with the

1 inclusion of ΔF, which is an indication of a better fitting model (Chambers, 1992).
2 Differences between young and older adults in peak plantar flexion and dorsiflexion torque,
3 and physical activity levels, were determined using independent t-tests. The α level for all
4 tests was 5%. The dataset and R code can be found at https://github.com/orssatto/PICs-
5 ageing.
6
RESULTS
Effects of age and muscle group on ΔF, peak discharge rate, and recruitment threshold
Motor units of older adults had lower ΔFs, discharged at lower rates, and were
recruited at lower torque (muscle force) levels than young adults in both *soleus* and *tibialis*
anterior. Also, ΔF levels and peak discharge rates were lower in *soleus* than *tibialis anterior*,
independent of age.
There were effects of age (β = -0.99, 95% CI = -1.49, -0.50; p < .001) and muscle (β
= 1.29, 95% CI = 0.65, 1.91; p < .001) but no age group by muscle effect (β = -0.29, 95% CI
= -1.08, 0.55; p = .45), on ΔF when calculated using the pairwise paired motor unit method.
ΔF was lower in older adults (Figure 2A) in both *soleus* (d = -1.50; Figure 3A) and *tibialis*
anterior (d = -1.42; Figure 3A). There were effects of age (β = -2.14, 95% CI = -2.98, -1.33;
p < .001) and muscle (β = 3.71, 95% CI = 2.64, 4.83; p < .001) on peak discharge rate, but
there was no age by muscle effect on peak discharge rate (β = 0.11, 95% CI = -1.37, 1.56; p =
.88). Peak discharge rate was lower in older adults (Figure 2B) in both *soleus* (d = -0.98;
Figure 3B) and *tibialis anterior* (d = -1.27; Figure 3B). There was an age effect on
recruitment threshold, with thresholds lower in older adults (β = -1.50, 95% CI = -2.89, -0.12;
p = .040; Figure 3C). There was no evidence of muscle (β = 1.12, 95% CI = -0.35, 2.65; p =
.15) or age by muscle (β = 1.12, 95% CI = -2.59, 1.38; p = .58) effects on recruitment
threshold.
The results of the additional, composite method, analysis revealed there was effects of
age (β = 1.65, 95% CI = -2.54, 0.77; p < .001) and muscle (β = 0.82, 95% CI = 0.22, 1.42; p
= .008) on ΔF, but no age by muscle effect on ΔF (β = 0.61, 95% CI = -1.43, 0.21; p = .150).
ΔFs were lower in older adults in both *soleus* (d = -2.14) and *tibialis anterior* (d = -2.93), as
shown in Figure 4.

Figure 2. ΔF calculated with the pairwise paired motor unit method (A), peak discharge rate

(B), and recruitment threshold (C) in *soleus* and *tibialis anterior* in young and older adults.

1   The mean (circle) and 95% confidence interval are offset to the right, with individual data

2   points coloured by participant. pps = peaks per second.

Figure 3. The marginal mean difference with 90% (thick line) and 95% (thick line) confidence intervals between young and older adults for ΔF calculated with the pairwise paired motor unit method (A), peak discharge rate (B), and recruitment threshold (RT) (C) in *soleus* and *tibialis anterior*. Negative values on all panels indicate the lower responses in older adults. There was an effect of age on ΔF, peak discharge rate, and recruitment threshold, with all variables lower in older adults. There was no age by muscle effect on ΔF, peak discharge rate, or recruitment threshold—indicating that age-dependent differences in these variables were similar for both muscles.

Figure 4. ΔF calculated with the composite paired motor unit method (A), and the marginal mean difference with 90% (thick line) and 95% (thick line) confidence intervals between young and older adults for ΔF (B). In panel A, the mean (circle) and 95% confidence interval are offset to the right, with individual data points coloured by participant. In panel B, negative values indicate ΔF was lower responses in older adults. pps = peaks per second.

Contribution of ΔF to peak discharge rate

There was a small, significant improvement in model fit when ΔF was included as a covariate in the prediction of peak discharge rate (X^2 = 10.444, p = .001). The inclusion of ΔF increased the R^2 by 0.017, from 0.445 (i.e., base model without ΔF) to 0.462. There was an statistical effect of ΔF on peak discharge rate (β = 0.28, 95% CI = 0.11, 0.45, p = .001), in addition to effects of age (β = -1.90, 95% CI = -2.68, -1.14, p < .001) and muscle (β = 3.54, 95% CI = 2.83, 4.28, p < .001).

Peak torque and physical activity levels

Peak torques and physical activity levels are shown in Table 1. Peak torque was lower in older adults for both plantar flexion (mean difference = -70 N·m, 95% CI = -95, -46; p < .001; d = -1.83) and dorsiflexion (mean difference = -12 N·m, 95% CI = -18, -5; p < .001; d = -1.18). Physical activity levels were not statistically different between young and older adults (mean difference = 594 MET min/week, 95% CI = -1269, 2457; p = .52; d = 0.20).

Motor unit identification

The number of motor units identified in *soleus* was 113 for the young group and 211for the older group. The median (IQR) number per younger participant was 7 (5–8) and per older participant 8 (6–9). For *tibialis anterior,* 293 motor units were identified in the young group and 411 in the older group. The median number per younger participant was 19 (16–23) and per older participant 20 (16–27). For the pairwise paired motor unit analysis, it was possible to obtain ΔF values from 16 young adults and 23 older adults in *soleus,* and from 16 young adults and 19 older adults in *tibialis anterior*. The number of test unit ΔFs for *soleus* in the young group was 70 and in the older group was 117. The median number per young participant was 5 (3–6) and per older participant was 4 (3–6). The number of test unit ΔFs for *tibialis anterior* in the young group was 185 and in the older group was 257. The median number per younger participant was 11 (8–14) and per older participant was 14 (9–18).

For the composite paired motor unit analysis, it was possible to obtain ΔF values from 4 young adults and 7 older adults for *soleus* and from 15 young adults and 18 older adults for *tibialis anterior*. The number of test unit ΔFs for *soleus* in the young group was 12 and in the older group was 17, with the median number per younger participant being 3 (3–4) and per older participant being 2 (1–4). The number of test unit ΔFs for *tibialis anterior* in the young

group was 93 and in the older group was 111, with the median number per young participant being 5 (4–8) and per older participant being 6 (3–8). No motor units were identified in one young participant and one older participant in *soleus*, and in one young participant and six older participants in *tibialis anterior*, respectively. These participants were not included in the analyses for the respective muscles.

DISCUSSION

The present study compared strengths of motor neuronal PICs, estimated using the paired motor unit technique (ΔF), between young and older adults. The main findings indicate that ΔF values are considerably lower in both *soleus* and *tibialis anterior* in older adults. These reductions are accompanied by lower peak discharge rates and the recruitment of motor units at lower torque levels in the older than younger adults in both muscles. As an exploratory analysis, a model was developed that showed a small contribution of ΔF to the between-subject variance in peak discharge rates after accounting for the effects of age, muscle. These findings point to a meaningful reduction in the intrinsic motor neurone excitability in older adults that is at least partly associated with a reduced amplitude of persistent inward currents triggered by muscle recruitment.

Estimates of persistent inward currents (ΔF)

The ΔF values obtained from *soleus* and *tibialis anterior* in young adults are similar to those obtained in previous studies using triangular-shaped contractions with a 2%/s force increase-decrease rate (Afsharipour *et al.*, 2020; Kim *et al.*, 2020; Trajano *et al.*, 2020) and indicate that ΔFs from older adults are substantially lower than in young adults in both *soleus* and *tibialis anterior*. These differences can be a result of the detrimental changes within the monoaminergic systems, which indicate lower release of serotonin and noradrenaline with ageing. Locus coeruleus and the dorsal raphe nucleus are the major central sources of noradrenaline and serotonin, respectively (Cassano *et al.*, 2009). Diminution of structural integrity (Liu *et al.*, 2019, 2020) and in neuromelanin (a product of noradrenaline synthesis) content in noradrenergic neurones emanating from locus coeruleus (Shibata *et al.*, 2006) indicate impaired noradrenaline secretion in older adults. Moreover, noradrenaline and serotonin concentration decrements have been observed in the brains of aged rats (Ko *et al.*, 1997; Míguez *et al.*, 1999), and degeneration of serotonergic axons projecting to the ventral horn of the lumbar segment of the spinal cord (where motor neurones innervating lower limb muscles emanate) have been detected (Johnson *et al.*, 1993). Also, serotonin receptors are

affected by expanded circulation of cytokines, resulting in increased re-uptake of serotonin (Michaud *et al.*, 2013). Thus, older adults might speculatively present reduced noradrenaline and serotonin secretions, and hence input onto motor neurones, which would then impair the initiation and modulation of PICs in this population. Changes in motor neurone integrity might also partly underpin reductions in intrinsic excitability. During ageing, axonal demyelination due to reduced expression of proteins responsible for myelination (Pannese, 2011) as well as axonal atrophy and degeneration have been observed, possibly subsequent to deregulated Ca^{2+} homeostasis (Nikoletopoulou & Tavernarakis, 2012) and to toxic, metabolic, or infectious injury sustained throughout the lifespan, or due to high levels of chronic inflammation and oxidative stress (Misgeld, 2011; Selman *et al.*, 2012). Motor neuronal death, especially in higher-threshold motor axons, leading to denervation of motor units has also been documented (Gordon *et al.*, 2004). In these cases, denervated motor units may remodel through reinnervation by nearby lower-threshold motor neurones (Manini *et al.*, 2013; Piasecki *et al.*, 2016), which may explain the reduced recruitment threshold observed in the older adults (as discussed below). These detrimental alterations in motor neurone structure are associated with reduced Ca^{2+}-mediated plateau potential durations in striatal neurones (from aged rats) (Dunia *et al.*, 1996), slower conduction velocity of efferent motor axons (Di Iorio *et al.*, 2006), lower incidence of doublet discharges, slower maximum discharge rates (Klass *et al.*, 2008; Orssatto *et al.*, 2020; Kirk *et al.*, 2021) alongside an increased afterhyperpolarisation duration (Piotrkiewicz *et al.*, 2007), and, as evidenced in the present study, lower PIC strength in the motor neurones. It is a logical hypothesis that the changes within the monoaminergic system and motor neurone structural integrity would possibly explain the reduced ΔFs observed in the present study. However, our findings indicate the need for further study of the cause-effect relationship between these mechanisms and reduced PICs in humans.

PICs are highly sensitive to synaptic inhibition, and both reciprocal and recurrent inhibition directly influence intrinsic motor neurone excitability, being effective PIC deactivators by opposing the facilitatory effects on PICs of descending brainstem neuromodulatory systems (Kuo *et al.*, 2003; Hyngstrom *et al.*, 2007, 2008; Bui *et al.*, 2008*a*, 2008*b*). It would therefore be expected that the reduced PIC strength estimates observed in the older adults in the present study might result from an age-dependent increase in reciprocal and/or recurrent inhibition. However, older adults present reduced reciprocal inhibition from the common peroneal nerve onto *soleus* and from the tibial nerve onto the *tibialis anterior* (Kido *et al.*, 2004). Also, recurrent inhibition onto the *soleus* motor neurone pool is suspected

to be relatively unaffected by ageing (Chalmers & Knutzen, 2004). Therefore, a reasonable supposition is that the behaviour of spinal inhibitory/excitatory systems (i.e., reduced reciprocal inhibition and preserved recurrent inhibition) of older adults are unlikely to be responsible for the decreases in PIC strength.

Recruitment threshold

Motor units that were identified in the older adults showed a lower recruitment threshold and were thus recruited at a lower torque level than in young adults. This behaviour may result from a compression in the range of firing rates observed previously in older adults motor units (Barry *et al.*, 2007; Girts *et al.*, 2020). Since motor unit discharge rate modulation in response to force changes is impaired in older adults, additional motor units must be recruited earlier in a triangular-shaped contraction to continually increase muscle force (Barry *et al.*, 2007; Enoka & Duchateau, 2017). Additionally, the observed lower recruitment thresholds might reflect the recruitment of a greater relative number of smaller motor neurones during the task. Older adults have a greater proportion of lower-threshold motor units as a result of motor unit remodelling subsequent to motor neuronal denervation, so motor units previously innervated by higher-threshold motor units become reinnervated by lower-threshold motor units (Piasecki *et al.*, 2016), as described above. In summary, our data indicate that aged motor neurones have a constrained capacity to amplify the excitatory synaptic input, consequently demanding an earlier recruitment of additional motor units to achieve the required motor output.

It is important to note that Ca^{2+} PIC channels can be activated below the action potential threshold (i.e., subthreshold PICs), strongly influencing motor unit recruitment (Li *et al.*, 2004). The possibility exists that the lower recruitment threshold of motor units of the older adults observed in the present study reflects a higher motor neurone excitability (i.e. enhanced motor neurone recruitment for a given input) because of stronger subthreshold PIC activation. An increase in the proportion of subthreshold PICs would reduce the recruitment threshold while also decreasing ΔF values. Subthreshold PICs may be stronger in smaller motor neurones, resulting in a larger overall (sub- plus supra-threshold) PIC strength (Afsharipour *et al.*, 2020). However, the ΔF method only estimates the suprathreshold contribution of the PICs to the discharge behaviour of motor units. Consequently, it is not possible to ascertain the behaviour of subthreshold PICs using the ΔF method. Our analytical approach accounted for any effect of recruitment threshold (and possibly motor neurone size)

on ΔF, that may otherwise confound any age-related effect, by including recruitment threshold in our modelling of ΔF. Further, the reanalysis of ΔF using the composite paired motor unit method appeared to remove the effect of recruitment threshold on ΔF values (from $\beta = 0.27$ to $\beta = -0.05$), which is an expected characteristic of the method (Afsharipour *et al.*, 2020). Therefore, the reduced ΔFs observed in older adults in the present study is not likely to have been an artefact of differences in motor neurone recruitment thresholds.

Peak discharge rates

The lower ΔF values in the older adults were also accompanied by reduced motor neurone discharge rates. The PIC is an important modulator of discharge rate output (Gorassini *et al.*, 2002; Orssatto *et al.*, 2021) and the available monoamines, serotonin and noradrenaline, facilitate PICs to increase motor neuronal gain and alter the input-output relationship according to the required output (Heckman & Binder, 1991; Heckman, 1994; Powers *et al.*, 2008; Powers & Heckman, 2015; Huh *et al.*, 2017; Orssatto *et al.*, 2021). PICs can amplify synaptic input by more than five-fold, and are thus a determinant mechanism influencing the capacity for motor neurones to achieve the necessary discharge rates to obtain very high muscle activation levels (Hounsgaard & Kiehn, 1993; Lee & Heckman, 1999, 2000). We have recently shown that increases in discharge rate during triangular-shaped contractions at different force levels were strongly associated with ΔF increases, and thus PIC amplitudes, using a within-subject design (Orssatto *et al.*, 2021). Nonetheless, data from the current study showed that ΔF explained only 2% of the between-subject variance in discharge rate output. Therefore, it may be case that ΔF explains more of the within-subject than between-subject differences in motor unit discharge rates. It is important to mention that motor unit discharge rates are not only modulated by PICs. Evidence in both animal (Chase *et al.*, 1985; Maxwell *et al.*, 2018) and human models (Rowe *et al.*, 2006) suggests that the synaptic input into the motor neurone decreases with ageing, which may result from increases in intracortical inhibition or reduced intracortical facilitation in older adults (Todd *et al.*, 2003; McGinley *et al.*, 2010; Opie *et al.*, 2020).

Strengths and limitations

The main strength of our study was the use of two validated (Gorassini *et al.*, 2002; Powers *et al.*, 2008; Afsharipour *et al.*, 2020) and a widely used methods to estimate PIC

strength in humans (Gorassini *et al.*, 2002; Powers *et al.*, 2008; Udina *et al.*, 2010; Hassan *et al.*, 2019, 2020; Trajano *et al.*, 2020; Orssatto *et al.*, 2021); however, both methods have limitations that should be pointed out. The pairwise method (Gorassini *et al.*, 2002) allowed us to obtain several pairs of motor units, having a larger amount of test units per participant. On the other hand, this method present a higher variance as a result of under and overestimation of ΔFs as a consequence of adopting control units with varied recruitment thresholds (Afsharipour *et al.*, 2020). Recently, Afsharipour et al., (2020) proposed the use of a composite control unit to reduce the ΔF variance present in the conventional paired motor unit analysis. However, this method requires some additional assumptions, such as the overlay of three lower threshold motor units with a similar discharge rate profile and with recruitment threshold below 3%. When following these assumptions, there was an important reduction of motor unit pairs, particularly for *soleus.* Therefore, we initially adopted the pairwise method as it permitted comparison between *soleus* and *tibialis anterior* ΔFs. We also ran the composite method as an additional analysis to examine whether reducing ΔF variability and removing the influence of the recruitment threshold on ΔF obtained with the pairwise method would affect our main outcomes. Using both methods allowed us to identify a large difference in ΔF between young and older adults, minimising the methodological limitations. Furthermore, ΔF values obtained in our study are derived from lower threshold *soleus* and *tibialis anterior* motor units recruited at a low force level (20% of peak torque); this is a commonly-used force target and is also similar to forces that might be expected in daily activities such as standing and walking. Motor neurones from distinct muscles depict different discharge behaviours during ageing (Kirk *et al.*, 2021); thus, it is possible that ΔF behaviour might also differ. Furthermore, there is evidence that the function and structure of higher threshold motor neurones are more affected than lower threshold motor neurones (Orssatto *et al.*, 2018). Therefore, we recommend that future studies investigate the effect of ageing on ΔF values from different muscles and at different contraction intensities.

Conclusions

The present study provides novel evidence of reduced intrinsic motor neurone excitability in humans by estimating PIC strengths using the paired-motor unit analyses. Older adults had substantially lower ΔFs, and presumably PIC strengths, in both *soleus* and *tibialis anterior* than young adults with matched physical activity level. This would likely influence the capacity of older individuals to activate the muscles, thus requiring a greater descending drive from cortical areas and hence level of volitional effort, and greater number

of recruited motor units to achieve the same force level (relative to maximum). We also identified a small contribution of ΔF to the between-subject variability in peak discharge rates. The present findings contribute to our understanding of the effects of ageing on motor neurone excitability, which is a potential mechanism underpinning motor functional loss during ageing; this hypothesis should be explicitly tested in future studies. A logical next step is to examine the effect of ageing on monoaminergic projections onto the motor neurones and their relationship with the reduced PIC strength observed in the present study.

ADDITIONAL INFORMATION

Data availability

The dataset and R code are available at https://github.com/orssatto/PICs-ageing.

Competing interests

The authors declare no competing interest related to this manuscript.

Author contributions

LBRO, AJS, AJB, and GST contributed with the conception and design of the work. LBRO acquired data. RLS developed the MATLAB script for ΔF, peak discharge rate, and recruitment threshold calculation. LBRO conducted the biological signals data analyses, and DNB developed the R script and conducted statistical analyses. All authors interpreted and discussed the data, drafted the manuscript, and revised it critically providing important intellectual content.

All authors approved the final version of the manuscript; agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved; and qualify for authorship, and all those who qualify for authorship are listed.

Funding

There is no specific funding related to this manuscript.

REFERENCES

Aagaard P, Suetta C, Caserotti P, Magnusson SP & Kjær M (2010). Role of the nervous system in sarcopenia and muscle atrophy with aging: Strength training as a countermeasure. *Scand J Med Sci Sport* **20,** 49–64.

Afsharipour B, Manzur N, Duchcherer J, Fenrich KF, Thompson CK, Negro F, Quinlan KA, Bennett DJ & Gorassini MA (2020). Estimation of self-sustained activity produced by persistent inward currents using firing rate profiles of multiple motor units in humans. *J Neurophysiol* **124,** 63–85.

Barry BK, Pascoe MA, Jesunathadas M & Enoka RM (2007). Rate coding is compressed but variability is unaltered for motor units in a hand muscle of old adults. *J Neurophysiol* **97,** 3206–3218.

Binder MD, Powers RK & Heckman CJ (2020). Nonlinear Input-Output Functions of Motoneurons. *Physiology (Bethesda)* **35,** 31–39.

Bui T V., Grande G & Rose PK (2008*a*). Relative location of inhibitory synapses and persistent inward currents determines the magnitude and mode of synaptic amplification in motoneurons. *J Neurophysiol* **99,** 583–594.

Bui T V., Grande G & Rose PK (2008*b*). Multiple modes of amplification of synaptic inhibition to motoneurons by persistent inward currents. *J Neurophysiol* **99,** 571–582.

Cassano T, Gaetani S, Morgese MG, Macheda T, Laconca L, Dipasquale P, Taltavull J, Shippenberg TS, Cuomo V & Gobi G (2009). Monoaminergic Changes in Locus Coeruleus and Dorsal Raphe Nucleus Following Noradrenaline Depletion. *Neurochem Res* **34,** 1417–1426.

Chalmers GR & Knutzen KM (2004). Recurrent inhibition in the soleus motor pool of elderly and young adults. *Electromyogr Clin Neurophysiol* **44,** 413–421.

Chambers JM (1992). Linear models. In *Statistical Models*, ed. Chambers JM & Hastie TJ. Wadsworth & Brooks/Cole.

Chase MH, Morales FR, Boxer PA & Fung SJ (1985). Aging of motoneurons and synaptic processes in the cat. *Exp Neurol* **90,** 471–478.

Christie A & Kamen G (2006). Doublet discharges in motoneurons of young and older adults. *J Neurophysiol* **95,** 2787–2795.

Desmedt BJE & Godaux E (1977). Ballistic contractions in man: Characteristic recruitment pattern of single motor units of the tibialis anterior muscle. *J Physiol* **264,** 673–693.

Dunia R, Buckwalter G, Defazio T, Dos Santos Villar F, Mcneill TH & Walsh JP (1996). Decreased duration of Ca2+-mediated plateau potentials in striatal neurons from aged rats. *J Neurophysiol* **76,** 2353–2363.

Enoka RM (2019). Physiological validation of the decomposition of surface EMG signals. *J Electromyogr Kinesiol* **46,** 70–83.

Enoka RM & Duchateau J (2017). Rate coding and the control of muscle force. *Cold Spring Harb Perspect Med*; DOI: 10.1101/cshperspect.a029702.

Girts RM, Mota JA, Harmon KK, MacLennan RJ & Stock MS (2020). Vastus Lateralis Motor Unit Recruitment Thresholds Are Compressed towards Lower Forces in Older Men. *J frailty aging* **9,** 191–196.

Gorassini M, Yang JF, Siu M & Bennett DJ (2002). Intrinsic activation of human

motoneurons: Possible contribution to motor unit excitation. *J Neurophysiol* **87,** 1850–1858.

Gordon T, Hegedus J & Tam SL (2004). Adaptive and Maladaptive Motor Axonal Sprouting in Aging and Motoneuron Disease. *Neurol Res* **26,** 174–185.

Hagströmer M, Oja P & Sjöström M (2006). The International Physical Activity Questionnaire (IPAQ): a study of concurrent and construct validity. *Public Health Nutr* **9,** 755–762.

Hassan AS, Kim EH, Khurram OU, Cummings M, Thompson CK, Miller McPherson L, Heckman CJ, Dewald JPA & Negro F (2019). Properties of Motor Units of Elbow and Ankle Muscles Decomposed Using High-Density Surface EMG. *Proc Annu Int Conf IEEE Eng Med Biol Soc EMBS*3874–3878.

Hassan AS, Thompson CK, Negro F, Cummings MQ, Powers RK, Heckman CJ, Dewald J & McPherson LM (2020). Impact of parameter selection on estimates of motoneuron excitability using paired motor unit analysis. *J Neural Eng*; DOI: 10.1088/1741-2552/ab5eda.

Heckman CJ (1994). Computer simulations of the effects of different synaptic input systems on the steady-state input-output structure of the motoneuron pool. *J Neurophysiol* **71,** 1727–1739.

Heckman CJ & Binder MD (1991). Computer simulation of the steady-state input-output function of the cat medial gastrocnemius motoneuron pool. *J Neurophysiol* **65,** 952–967.

Heckman CJ & Enoka RM (2012). Motor unit. *Compr Physiol* **2,** 2629–2682.

Heckman CJ, Gorassini MA & Bennett DJ (2005). Persistent inward currents in motoneuron dendrites: Implications for motor output. *Muscle and Nerve* **31,** 135–156.

Holobar A, Minetto MA & Farina D (2014). Accurate identification of motor unit discharge patterns from high-density surface EMG and validation with a novel signal-based performance metric. *J Neural Eng*; DOI: 10.1088/1741-2560/11/1/016008.

Holobar A & Zazula D (2007). Multichannel blind source separation using convolution Kernel compensation. *IEEE Trans Signal Process* **55,** 4487–4496.

Hounsgaard J & Kiehn O (1993). Calcium spikes and calcium plateaux evoked by differential polarization in dendrites of turtle motoneurones in vitro. *J Physiol* **468,** 245–259.

Huh S, Siripuram R, Lee RH, Turkin V V., O'Neill D, Hamm TM, Heckman CJ & Manuel M (2017). PICs in motoneurons do not scale with the size of the animal: A possible mechanism for faster speed of muscle contraction in smaller species. *J Neurophysiol* **118,** 93–102.

Hyngstrom AS, Johnson MD & Heckman CJ (2008). Summation of excitatory and inhibitory synaptic inputs by motoneurons with highly active dendrites. *J Neurophysiol* **99,** 1643–1652.

Hyngstrom AS, Johnson MD, Miller JF & Heckman CJ (2007). Intrinsic electrical properties of spinal motoneurons vary with joint angle. *Nat Neurosci* **10,** 363–369.

Di Iorio A, Cherubini A, Volpato S, Sparvieri E, Lauretani F, Franceschi C, Senin U, Abate G, Paganelli R, Martin A, Andres-Lacueva C & Ferrucci L (2006). Markers of inflammation, Vitamin E and peripheral nervous system function. The InCHIANTI study. *Neurobiol Aging* **27,** 1280–1288.

Johnson H, Ulfhake B, Dagerlind Å, Bennett GW, Fone KCF & Hökfelt T (1993). The

serotoninergic bulbospinal system and brainstem□spinal cord content of serotonin□, TRH□, and substance P□like immunoreactivity in the aged rat with special reference to the spinal cord motor nucleus. *Synapse* **15,** 63–89.

Kido A, Tanaka N & Stein RB (2004). Spinal excitation and inhibition decrease as humans age. *Can J Physiol Pharmacol* **82,** 238–248.

Kim EH, Wilson JM, Thompson CK & Heckman CJ (2020). Differences in Estimated Persistent Inward Currents Between Ankle Flexors and 2 Extensors in Humans. *J Neurophysiol* **21,** 1–9.

Kirk EA, Christie AD, Knight CA & Rice CL (2021). Motor unit firing rates during constant isometric contraction□: establishing and comparing an age-related pattern among muscles. *J Appl Physiol*; DOI: https://doi.org/10.1152/japplphysiol.01047.2020.

Klass M, Baudry S & Duchateau J (2008). Age-related decline in rate of torque development is accompanied by lower maximal motor unit discharge frequency during fast contractions. *J Appl Physiol* **104,** 739–746.

Ko ML, King MA, Gordon TL & Crisp T (1997). The effects of aging on spinal neurochemistry in the rat. *Brain Res Bull* **42,** 95–98.

Kuo JJ, Lee RH, Johnson MD, Heckman HM & Heckman CJ (2003). Active Dendritic Integration of Inhibitory Synaptic Inputs in Vivo. *J Neurophysiol* **90,** 3617–3624.

Kuznetsova A, Brockhoff PB & Christensen RHB (2017). lmerTest Package: Tests in Linear Mixed Effects Models . *J Stat Softw*; DOI: 10.18637/jss.v082.i13.

Larsson L, Degens H, Li M, Salviati L, Lee Y il, Thompson W, Kirkland JL & Sandri M (2019). Sarcopenia: Aging-Related Loss of Muscle Mass and Function. *Physiol Rev* **99,** 427–511.

Laughton CA, Slavin M, Katdare K, Nolan L, Bean JF, Kerrigan DC, Phillips E, Lipsitz LA & Collins JJ (2003). Aging, muscle activity, and balance control: Physiologic changes associated with balance impairment. *Gait Posture* **18,** 101–108.

Lee RH & Heckman CJ (1999). Enhancement of bistability in spinal motoneurons in vivo by the noradrenergic α1 agonist methoxamine. *J Neurophysiol* **81,** 2164–2174.

Lee RH & Heckman CJ (2000). Adjustable amplification of synaptic input in the dendrites of spinal motoneurons in vivo. *J Neurosci* **20,** 6734–6740.

Lenth R V., Buerkner P, Herve M, Love J, Riebl H & Singmann H (2021). emmeans: Estimated Marginal Means, aka Least-Squares Means - R package. Available at: https://github.com/rvlenth/emmeans.

Li Y, Gorassini MA & Bennett DJ (2004). Role of Persistent Sodium and Calcium Currents in Motoneuron Firing and Spasticity in Chronic Spinal Rats. *J Neurophysiol* **91,** 767–783.

Liu KY et al. (2020). Noradrenergic-dependent functions are associated with age-related locus coeruleus signal intensity differences. *Nat Commun* **11,** 1–9.

Liu KY, Acosta-Cabronero J, Cardenas-Blanco A, Loane C, Berry AJ, Betts MJ, Kievit RA, Henson RN, Düzel E, Howard R & Hämmerer D (2019). In vivo visualization of age-related differences in the locus coeruleus. *Neurobiol Aging* **74,** 101–111.

Manini TM, Hong SL & Clark BC (2013). Aging and muscle: a neuron's perspective. *Curr Opin Clin Nutr Metab Care* **16,** 1–10.

Maxwell N, Castro RW, Sutherland NM, Vaughan KL, Szarowicz MD, de Cabo R, Mattison

JA & Valdez G (2018). α-Motor neurons are spared from aging while their synaptic inputs degenerate in monkeys and mice. *Aging Cell*; DOI: 10.1111/acel.12726.

McGinley M, Hoffman RL, Russ DW, Thomas JS & Clark BC (2010). Older adults exhibit more intracortical inhibition and less intracortical facilitation than young adults. *Exp Gerontol* **45,** 671–678.

Michaud M, Balardy L, Moulis G, Gaudin C, Peyrot C, Vellas B, Cesari M & Nourhashemi F (2013). Proinflammatory cytokines, aging, and age-related diseases. *J Am Med Dir Assoc* **14,** 877–882.

Míguez JM, Aldegunde M, Paz-Valiñas L, Recio J & Sánchez-Barceló E (1999). Selective changes in the contents of noradrenaline, dopamine and serotonin in rat brain areas during aging. *J Neural Transm* **106,** 1089–1098.

Misgeld T (2011). Lost in elimination: mechanisms of axonal loss. *e-Neuroforum* **2,** 21–34.

Morales FR, Boxer PA, Fung SJ & Chase MH (1987). Basic electrophysiological properties of spinal cord motoneurons during old age in the cat. *J Neurophysiol* **58,** 180–194.

Nakagawa S, Johnson PCD & Schielzeth H (2017). The coefficient of determination R2 and intra-class correlation coefficient from generalized linear mixed-effects models revisited and expanded. *J R Soc Interface*; DOI: 10.1098/rsif.2017.0213.

Nikoletopoulou V & Tavernarakis N (2012). Calcium homeostasis in aging neurons. *Front Genet* **3,** 1–17.

Opie GM, Hand BJ & Semmler JG (2020). Age-related changes in late synaptic inputs to corticospinal neurons and their functional significance: A paired-pulse TMS study. *Brain Stimul* **13,** 239–246.

Orssatto LB da R, Wiest MJ & Diefenthaeler F (2018). Neural and musculotendinous mechanisms underpinning age-related force reductions. *Mech Ageing Dev* **175,** 17–23.

Orssatto LBR, Borg D, Pendrith L, Shield AJ, Blazevich AJ & Trajano GS (2020). Do motoneurons slow with aging? A systematic review and meta-analysis with meta-regression. *bioRxiv*; DOI: 10.1101/2020.12.15.422812.

Orssatto LBR, Mackay K, Shield AJ, Sakugawa RL, Blazevich AJ & Trajano GS (2021). Estimates of persistent inward currents increase with the level of voluntary drive in low-threshold motor units of plantar flexor muscles. *J Neurophysiol* 6.

Pannese E (2011). Morphological changes in nerve cells during normal aging. *Brain Struct Funct* **216,** 85–89.

Piasecki M, Ireland A, Jones DA & McPhee JS (2016). Age-dependent motor unit remodelling in human limb muscles. *Biogerontology* **17,** 485–496.

Piotrkiewicz M, Kudina L, Mierzejewska J, Jakubiec M & Hausmanowa-petrusewicz I (2007). Age-related change in duration of afterhyperpolarization of human motoneurones. *J Physiol* **585,** 483–490.

Polcyn AE, Kerrigan DC & Collins JJ (1998). Age-Related Changes in the Initiation of Gait: Degradation of Central Mechanisms for Momentum Generation. *Arch Phys Med Rehabil* **79,** 1582–1589.

Powers RK & Heckman CJ (2015). Contribution of intrinsic motoneuron properties to discharge hysteresis and its estimation based on paired motor unit recordings: A simulation study. *J Neurophysiol* **114,** 184–198.

Powers RK, Nardelli P & Cope TC (2008). Estimation of the contribution of intrinsic currents

to motoneuron firing based on paired motoneuron discharge records in the decerebrate cat. *J Neurophysiol* **100,** 292–303.

Rowe JB, Siebner H, Filipovic SR, Cordivari C, Gerschlager W, Rothwell J & Frackowiak R (2006). Aging is associated with contrasting changes in local and distant cortical connectivity in the human motor system. *Neuroimage* **32,** 747–760.

Selman C, Blount JD, Nussey DH & Speakman JR (2012). Oxidative damage, ageing, and life-history evolution: Where now? *Trends Ecol Evol* **27,** 570–577.

Shibata E, Sasaki M, Tohyama K, Kanbara Y, Otsuka K, Ehara S & Sakai A (2006). Age-related changes in locus ceruleus on neuromelanin magnetic resonance imaging at 3 Tesla. *Magn Reson Med Sci* **5,** 197–200.

Stephenson JL & Maluf KS (2011). Dependence of the paired motor unit analysis on motor unit discharge characteristics in the human tibialis anterior muscle. *J Neurosci Methods* **198,** 84–92.

Suetta C, Haddock B, Alcazar J, Noerst T, Hansen OM, Ludvig H, Kamper RS, Schnohr P, Prescott E, Andersen LL, Frandsen U, Aagaard P, Bülow J, Hovind P & Simonsen L (2019). The Copenhagen Sarcopenia Study: lean mass, strength, power, and physical function in a Danish cohort aged 20–93 years. *J Cachexia Sarcopenia Muscle* 1–13.

Todd G, Taylor JL & Gandevia SC (2003). Measurement of voluntary activation of fresh and fatigued human muscles using transcranial magnetic stimulation. *J Physiol* **551,** 661–671.

Trajano GS, Taylor JL, Orssatto LBR, McNulty CR & Blazevich AJ (2020). Passive muscle stretching reduces estimates of persistent inward current strength in soleus motor units. *J Exp Biol* **223,** jeb.229922.

Udina E, D'Amico J, Bergquist AJ & Gorassini MA (2010). Amphetamine increases persistent inward currents in human motoneurons estimated from paired motor-unit activity. *J Neurophysiol* **103,** 1295–1303.

Vandenberk MS & Kalmar JM (2014). An evaluation of paired motor unit estimates of persistent inward current in human motoneurons. *J Neurophysiol* **111,** 1877–1884.

Vandervoort A a. (2002). Aging of the human neuromuscular system. *Muscle and Nerve* **25,** 17–25.