Genetic Loci and Metabolic States Associated With Murine Epigenetic Aging

Khyobeni Mozhui [1, 2*], Ake T. Lu[3], Caesar Z Li [3], Amin Haghani [4], Jose Vladimir Sandoval-Sierra [1], Yibo Wu[5,6], Robert W. Williams [2], Steve Horvath[3, 4]

[1] Department of Preventive Medicine, University of Tennessee Health Science Center, College of Medicine, Memphis, Tennessee, USA
[2] Department of Genetics, Genomics and Informatics, University of Tennessee Health Science Center, College of Medicine, Memphis, Tennessee, USA
[3] Department of Human Genetics, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, CA, USA
[4] Department of Biostatistics, Fielding School of Public Health, University of California Los Angeles, Los Angeles, CA, USA
[5]YCI Laboratory for Next-Generation Proteomics, RIKEN Center for Integrative Medical Sciences, Yokohama City, Kanagawa, Japan
[6]University of Geneva, Geneva, Switzerland

*Correspondence:
Khyobeni Mozhui
kmozhui@uthsc.edu

Abstract

Changes in DNA methylation (DNAm) are linked to aging. Here, we profile highly conserved CpGs in 339 predominantly female mice belonging to the BXD family for which we have deep longevity and genomic data. We use a 'pan-mammalian' microarray that provides a common platform for assaying the methylome across mammalian clades. We computed epigenetic clocks and tested associations with DNAm entropy, diet, weight, metabolic traits, and genetic variation. We describe the multifactorial variance of methylation at these CpGs, and show that high fat diet augments the age-associated changes. Entropy increases with age. The progression to disorder, particularly at CpGs that gain methylation over time, was predictive of genotype-dependent life expectancy. The longer-lived BXD strains had comparatively lower entropy at a given age. We identified two genetic loci that modulate rates of epigenetic age acceleration (EAA): one on chromosome (Chr) 11 that encompasses the *Erbb2/Her2* oncogenic region, and a second on Chr19 that contains a cytochrome P450 cluster. Both loci harbor genes associated with EAA in humans including *STXBP4*, *NKX2*-3, and *CUTC*. Transcriptome and proteome analyses revealed associations with oxidation-reduction, metabolic, and immune response pathways. Our results highlight concordant loci for EAA in humans and mice, and demonstrate a tight coupling between the metabolic state and epigenetic aging.

Keywords: epigenetic clock, lifespan, entropy, DNA methylation, genetic mapping, QTL, weight, diet

Introduction

Epigenetic clocks are widely used molecular biomarkers of aging.[1] These DNA methylation (DNAm) age predictors are based on the methylation levels of select CpGs that are distributed across the genome. Each CpG that is used in a clock model is assigned a specific weight, typically derived from supervised training algorithms,[2-4] and collectively, the methylation status across this ensemble of "clock CpGs" are used to estimate the epigenetic age (DNAmAge). This estimate tracks closely, but not perfectly, with an individual's chronological age. How much the DNAmAge deviates from the known chronological age can be a measure of the rate of biological aging. Denoted as "epigenetic age acceleration" (or EAA), a more accelerated clock (positive EAA) suggests an older biological age, and a decelerated clock (negative EAA) suggests a younger biological age. While DNAmAge predicts age, its age-adjusted counterpart, EAA, is associated with variation in health, fitness, exposure to stressors, body mass index (BMI), and even life expectancy.[5-9]

These DNAm clocks were initially reported for humans.[10-12] Since then, many different models of human DNAm clock have been develop, and this rapid expansion was made possible by reliable DNAm microarrays that provide a fixed CpG content—starting with the Illumina Infinium 27K to the current 850K EPIC array.[11,13-15] These clock variants differ in the subset of CpGs that go into the age estimation model. Some clock models are specific to cells or tissues, others are multi-tissue. Some clocks perform better at predicting chronological age, others better capture biological aging and predict health and life expectancy. [8,16-18] The performance of these clocks depend heavily on the training models, and the size and tissue types of the training set.[13]

This age biomarker has also been extended to model organisms, and this has opened up the possibility of directly testing the effects of different interventions such as calorie restriction, rapamycin, and genetic manipulation.[3,19-23] However, one point to note is that model organisms have not benefitted from a microarray platform comparable to that of the human methylation Infinium arrays. Most rodent studies have used enrichment-based DNAm sequencing, and this limits the transferability and reproducibility of clocks between datasets since the same CpGs are not always covered.[21] Moreover, these studies are usually performed in a single inbred strain (for mouse, the canonical C57BL/6), or at most, a few genetic backgrounds, and this makes it impossible to carry out genetic mapping studies that can complement the human genome-wide association studies (GWAS) of epigenetic aging.[24-28]

A new microarray was recently developed to profile CpGs that have high conservation in mammals. This pan-mammalian DNAm array (HorvathMammalMethylChip40) surveys over 37K CpGs and provides a unifying platform to study epigenetic aging in mammals.[29] This array has been used to build multi-tissue universal clocks and lifespan predictors that are applicable to a variety of mammalian species.[30,31] Here, we use this array to examine the dynamism and variability of the conserved CpGs in a genetically diverse cohort of mice belonging to the BXD family.[32,33]

The BXDs are one of the pre-eminent murine genetic reference panels used as the experimental paradigm of precision medicine.[34] They are a large family of recombinant inbred (RI) strains

made by crossing the C57BL/6J (B6) and DBA/2J (D2) parental strains. The family has been
expanded to 150 fully sequenced progeny strains.[34,35] The individual members of the BXD family
(e.g., BXD1, BXD27, BXD102), each represents a replicable isogenic cohort. The family
segregates for a high level of genetic variation, and likewise, family members have high
variation in their metabolic profiles, responses to diet, aging rates, and life expectancies.[32-34,36-
[38] The availability of deep sequence data, and unrivaled multi-omic and phenomic data make
the BXDs a powerful tool with which to evaluate the causal linkage between genome,
epigenome, and aging rates.

In our previous work, we used an enrichment-based sequencing to assay the methylome in a
modest number of BXD mice, and reported rapid age-dependent methylation changes in mice
on high fat diet, and in mice with higher body weight.[39] In the present work, we start by testing
the performance of new pan-tissue and liver-specific epigenetic mouse clocks, and evaluate
how these relate to metabolic states, genotype-dependent life expectancy, and methylome
entropy. We also apply a multi-factor analysis of site-specific CpG methylation to understand
association among four key variables—chronological age, diet, weight, and lifespan—and the
liver methylome. We perform quantitative trait locus (QTL) mapping, along with multi-omic
gene expression analyses, and identify upstream gene loci that modulate the DNAm clocks in
mice.

Our results are consistent with a faster clock for cases on HFD, and with higher body weight.
This may be partly because exposure to HFD augmented the age-dependent gains in
methylation at specific CpGs. We also observed that BXD genotypes with longer life expectancy
tend to have lower methylation at CpGs that undergo age-dependent methylation gains, and
the entropy computed from this set of CpGs have a significant inverse correlation with strain
lifespan. QTL mapping uncovered loci on chromosomes (Chrs) 11 and 19 that are associated
with EAA. A strong candidate gene in the chromosome (Chr) 11 interval (referred to as *Eaa11*) is
Stxbp4, a gene that has been consistently associated with EAA by human genome-wide
association studies (GWAS).[24,26,27] The Chr19 QTL (*Eaa19*) also harbors strong contenders
including *Cyp26a1*, *Myof*, *Cutc*, and *Nkx2–3*, and the conserved genes in humans have been
associated with longevity and EAA.[27,40,41] We performed gene expression analyses using
transcriptomic and proteomic data to clarify the molecular pathways associated with epigenetic
aging, and this highlighted metabolic networks, and also apolipoproteins (including APOE) as
strong expression correlates.

Results

Description of samples

118 Liver DNAm data was from 321 female and 18 male belonging to 45 members of the BXD
119 family, including both parental strains and F1 hybrids. Age ranged from 5.6 to 33.4 months.
120 Mice were all weaned onto a normal chow (control diet; CD) and a balanced subset of cases
121 were then randomly assigned to HFD (see Roy et al. for details[33]). Tissues were collected at
122 approximately six months intervals (see Williams et al.[32]). Individual-level data are in **Data S1**.

DNAm clocks, entropy, and chronological age prediction

We built three different mouse clocks, and each was developed as a pair depending on whether
the training set used all tissues (pan-tissue) or a specific tissue (in this case, liver). These are: (1)
a general DNAm clock (referred to simply as DNAmAge): clock trained without pre-selecting for
any specific CpGs; (2) developmental clock (dev.DNAmAge): built from CpGs that change during
development (defined as the period from prenatal to 1.6 months); and (3) interventional clock
(int.DNAmAge): built from CpGs that change in response to aging related interventions (calorie
restriction and growth hormone receptor knockout). The clocks we report here were trained in
a larger mouse dataset that excluded the BXDs and are therefore unbiased to the
characteristics of the BXD Family.[30,31,42] The specific clock CpGs and coefficients for DNAmAge
computation are in **Data S2**. All the mouse clocks performed well in age estimation and had an
average r of 0.89 with chronological age. However, the interventional clocks had higher
deviation from chronological age and higher median predictive error (**Table 1; Fig 1a**). The age-
adjusted EAA derived from these clocks showed wide individual variation (**Fig 1b**).

We next estimated the methylome-wide entropy as a measure of randomness and information
loss. This was computed from 27966 probes that provide high-quality data and have been
validated to perform well in mice.[29] Consistent with previous reports,[10,43] this property
increased with chronological age, and age accounted for about 6% (in CD) to 28% (in HFD) of
the variance in entropy (**Fig 1c**). As direct correlates of chronological age, all the DNAmAge
estimates also had significant positive correlations with entropy (**Table 1**). We hypothesized
that higher entropy levels will be associated with higher EAA, and based on this bivariate
comparison, most of the EAA showed a significant positive correlation with entropy (**Data S3;**
Fig 1d).

Table 1. Chronological age prediction and correlation with methylome-wide entropy

Clock type	DNAmAge name	Tissue	r with age (n=339) [1]	Age prediction median error	r with entropy (n=339) [1, 2]
Standard clocks	DNAmAge	pan	0.89	0.12	0.43
		liver	0.92	0.10	0.40
Developmental clocks	dev.DNAmAge	pan	0.87	0.14	0.39
		liver	0.91	0.12	0.37
Interventional clocks	int.DNAmAge	pan	0.85	0.17	0.29
		liver	0.86	0.15	0.47

147 [1]p < .0001; [2]p < .0001 Methylome-wide entropy calculated from ~28K CpGs

How the epigenetic readouts relate to diet, sex, and metabolic traits

Diet. HFD was associated with higher EAA for four of the clocks (**Table 2**). For instance, the
liver-specific interventional clock diverged between the diets (**Fig 1a**), and CD mice had an
average of –0.04 years of age deceleration, and HFD mice had an average of +0.11 years of age
acceleration **(Table 2)**. The two clocks that were not affected by diet were the liver general and
developmental clocks. Methylome-wide entropy was not different between the diets.

Body weight. Body weight was first measured when mice were at an average age of 4.5 ± 2.7
months. We refer to this initial weight as baseline body weight (BW0). For mice on HFD, this

Fig 1. Correlates and modifiers of epigenetic clocks and methylome-wide entropy
(a) Correlation between chronological age and predicted age (shown for the liver intervention clock or int.DNAmAge). Black circles are control diet (CD, n = 210); red crosses are high fat diet mice (HFD, n = 128) **(b)** Violin plots of age-adjusted epigenetic age acceleration (EAA) ("int" stands for interventional, "dev" stands for developmental). **(c)** Shannon entropy, calculated from the full set of high quality CpGs, increases with age. **(d)** Methylome entropy has a direct correlation with EAA (shown for the liver int.EAA). **(e)** For 48 mice, initial body weight (BW0) was measured 1 or 3 days after introduction to HFD, and these showed significant correlation with EAA. **(f)** Weight was first measured at mean age of 4.5 ± 2.7 months (BW0), and then at 6.3 ± 2.8 months (BW1). Weight gains during this interval (deltaBW = BW1 − BW0) is a direct correlate of EAA. **(g)** For BXD genotypes with males and females samples, males have higher age acceleration. Bars represent mean ± standard error; 40 females (26 CD, 14 HFD) and 18 males (10 CD, 8 HFD). **(h)** Relative effects of different predictor variables on EAA shown as logworth scores ($-\log_{10}p$). The dashed lines correspond to p = 0.01. Positive logworth values indicate positive regression estimates (for diet, positive means higher in high fat diet compared to control diet). BWF is final weight; Chol is serum total cholesterol; Gluc is fasted glucose levels. **(g)** The residual plot display association between methylome-wide entropy and the pan-tissue int.EAA after adjustment for diet, age, weight, glucose, cholesterol, and batch.

157 **Table 2. Association with diet and weight, and heritability of the epigenetic readouts**

Type	EAA	Diet	Mean (SD)	Diet (p)	r BW0 [a]	p BW0	r BWF [a]	p BWF	H²	Strain r [b]
Mouse clocks	EAA, pan	CD	-0.05 ± 0.21	<.0001	0.19	0.006	0.29	<.0001	0.49	0.54
		HFD	0.07 ± 0.21		0.21	0.01	0.42	<.0001	0.50	
	EAA, liver	CD	0 ± 0.17	ns	0.09	ns	0.20	0.003	0.40	0.73
		HFD	0.03 ± 0.14		0.22	0.01	0.49	<.0001	0.52	
	dev.EAA, pan	CD	-0.04 ± 0.23	0.004	0.09	ns	0.22	0.001	0.53	0.76
		HFD	0.03 ± 0.22		0.27	0.002	0.45	<.0001	0.61	
	dev.EAA, liver	CD	0 ± 0.2	ns	0.19	0.002	0.29	<.0001	0.46	0.78
		HFD	0 ± 0.16		0.29	0.0007	0.47	<.0001	0.60	
	int.EAA, pan	CD	-0.05 ± 0.25	0.0003	0.03	ns	0.21	0.002	0.27	0.66
		HFD	0.06 ± 0.33		0.22	0.01	0.46	<.0001	0.45	
	int.EAA, liver	CD	-0.04 ± 0.22	<.0001	0.05	ns	0.18	0.01	0.59	0.80
		HFD	0.11 ± 0.25		0.27	0.002	0.58	<.0001	0.54	
Entropy	-	CD	2.67 ± 0.02	ns	0.09	ns	0.05	ns	0.31	0.24 (ns)
		HFD	2.67 ± 0.02		0.15	0.09	0.15	0.09	0.32	

[a] BW0 is body weight at about 4.5 months of age (n = 339; 210 CD and 129 HFD); BWF is final weight at tissue collection (1 HFD case missing data; n = 338; 210
CD and 128 HFD)
[b] Pearson correlation between strain means for n = 29 BXD genotypes kept on CD and HFD

was usually before introduction to the diet, except for 48 cases that were first weighed 1 or 3
days after HFD (**Data S1**). In the CD group, only the EAA from the pan-tissue standard and liver
developmental clocks showed modest but significant positive correlations with BW0 (**Table 2**).
In the HFD group, the positive correlation with BW0 was more robust and consistent across all
the clocks, and this may have been due to the inclusion of the 48 cases that had been on HFD
for 1 or 3 days. Taking only these 48 cases, we found that higher weight even after 1 day of HFD
had an age-accelerating effect on all the clocks (**Data S3**), and this was particularly strong for
the interventional clocks (r = 0.45, p = 0.001 for the pan-tissue int.EAA; r = 0.58, p < 0.0001 for
the liver int.EAA) (**Fig 1e**). Second weight was measured 7.4 ± 5.2 weeks after BW0 (mean age
6.3 ± 2.8 months). We refer to this as BW1 and we estimated the weight change as deltaBW =
BW1 – BW0. DeltaBW was a positive correlate of EAA on both diets, albeit more pronounce in
the HFD group (**Fig 1f**; **Data S3**). The final body weight (BWF) was measured at the time of
tissue harvest, and EAA from all the mouse clocks were significant correlates of BWF on both
diets **(Table 2)**. In contrast, entropy did not show an association with either BW0 or BWF. We
do note that when stratified by diet, the entropy level had a slight positive correlation with
BW1 in the HFD group (r = 0.23, p = 0.008), but not in the CD group (**Data S3**).

Sex. Four BXD genotypes (B6D2F1, D2B6F1, BXD102, B6) had cases from both males and
females. We used these to test for sex effects. All the clocks showed significant age acceleration
in male mice, and this effect was particularly strong for the both dev.EAA, and the pan-tissue
int.EAA (**Fig 1g; Data S3**). This effect was independent of the higher BWF of males, and the
higher age-acceleration in males was detected after adjusting for BWF (**Table S1**). There was no
significant difference in entropy between the sexes.

Metabolic measures. 278 cases with DNAm data also had fasted serum glucose and total
cholesterol,[32,33] and we examined whether these metabolic traits were associated with either
the EAA measures or methylome entropy. Since these are highly dependent on diet, weight,
and age, we applied a multivariable model to jointly examine how the different metabolic
variables (cholesterol and glucose, as well as diet and weight) and entropy relate to EAA after
adjusting for age. To test the robustness of associations, we also include the methylation array
batch as another covariate (**Data S1** has batch information; **Data S4** has the full statistics). **Fig**
1h shows the relative strengths and directions of associations between these variables and the
EAA traits. Except for the pan-tissue interventional clock, entropy had a strong positive
association with EAA. For example, a plot of residuals between entropy and the liver int.EAA
indicates that after adjusting for all the other covariates, the methylome-wide entropy explains
17% of the variance in int.EAA (**Fig 1i**). Since diet strongly influences BWF, the inclusion of BWF
in the regression diminished the effect of diet. For the two clocks that were not influenced by
diet (the liver EAA and liver dev.EAA), adjusting for the effect of BWF resulted in an inverse
association with diet (i.e., the residual EAA values after accounting for BWF were slightly lower
in the HFD group). Fasted glucose did not have a significant effect on EAA. Cholesterol had a
positive association with the interventional clocks but the effects were modest (residual R^2 =
0.02 and p = 0.02 for cholesterol and the pan-tissue int.EAA) (**Data S4**).

We also performed a similar analysis with BW0 instead of BWF (**Fig S1**), and here, HFD
remained as an accelerator of the pan-tissue EAA and liver int.EAA. Cholesterol also became a
significant positive correlate of EAA for the interventional clocks (**Fig S1**). This would suggest

that the effect of diet on EAA is mostly mediated by its impact on physiological and metabolic
traits, and BWF becomes a prominent predictor of EAA.

To summarize, our results indicate that the degree of disorder in the methylome increases with
age, and may partly contribute to the epigenetic clocks as higher entropy is associated with
higher EAA. The EAA traits were also associated with biological variables (i.e., body weight, diet,
and sex). Of these, BWF was the strongest modulator of EAA.

How the epigenetic readouts relate to strain longevity

We next obtained longevity data from a parallel cohort of female BXD mice that were allowed
to age on CD or HFD.[33] Since the strain lifespan was determined from female BXDs, we
restricted this to only the female cases. For strains with natural death data from n ≥ 5, we
computed the minimum (minLS), 25th quartile (25Q-LS), mean, median lifespan, 75th quartile
(75Q-LS), and maximum lifespan (maxLS) (**Data S1**). Specifically, we postulated an accelerated
clock for strains with shorter lifespan (i.e., inverse correlation). Overall, the EAA measures
showed the expected inverse correlation trend with the lifespan statistics. However, these
correlations were weak. The correlations were significant only for the pan-tissue general clock
(**Fig S2a**) and the liver intervention clock, with explained variance in lifespan of only ~3% (**Table
S2; Fig S2b, S2c**). When separated by diet, these correlations became weaker indicating that
while we see the expected inverse relationship, the EAA is only weakly predictive of strain
longevity. Entropy estimated from the full set of CpGs was unrelated to strain longevity.

Multifactor variance of the conserved CpGs

Both the entropy and clock readouts capture the overall variation across multiple CpGs, and to
gain insights into the underlying variance patterns, we performed a multivariable epigenome-
wide association study (EWAS). For this, we applied a site-by-site regression on the 27966
validated CpGs,[29] and tested for association with age, BWF, diet, and genotype-dependent
strain median lifespan (full set of probes, annotations, and EWAS results in **Data S5**).

Age was clearly the most influential variable, and this is apparent from the volcano plots (**Fig
2a–d**). We used a cutoff of Bonferroni $p \leq 0.05$ to define differentially methylated CpGs (DMCs),
and 6553 CpGs were associated with age (referred to as age-DMCs), 733 with weight (weight-
DMCs), 321 with diet (diet-DMCs), and 236 with genotype-dependent lifespan (LS-DMCs). We
note extensive overlap among the lists of DMCs that shows that variation at these CpGs are
multifactorial in nature (**1e**). Majority of the age-DMCs (77%) gained methylation (or age-gain),
and consistent with previous observations, age-gain CpGs tended to be in regions with low
methylation, whereas age-DMCs that declined in methylation (age-loss) were in regions with
high methylation (**Fig 2f**).[39,43,44] By overlaying the volcano plots with the age-gain and age-loss
information, we see distinct patterns in how these age-DMCs vary with weight (**Fig 2b**), diet (**Fig
2c**), and genotype lifespan (**Fig 2d**). While the majority of CpGs, including several age-loss CpGs,
had negative regression estimates for weight (i.e., decrease in DNAm with unit increase in
weight), HFD was associated with higher methylation levels (positive regression estimates)
including at several age-DMCs (**Fig 2c**). This pattern of inverse correlation with weight but
heightened methylation due to HFD is illustrated by the CpG in the 3'UTR of *Mettl23*
(cg10587537) (**Fig 2g**). Taking the 6553 age-DMCs, a comparison of the regression estimates for

Fig 2. Multivariable analysis of site-specific methylation
(a) Volcano plot comparing regression estimates (change in methylation beta-value per day of age) versus the statistical significance for age effect. Dashed line denotes the Bonferroni p = 0.05 for ~28K tests). Similar volcano plots for predictor variables: **(b)** final body weight (regression estimates are change per gram of weight), **(c)** diet (change in high fat compared to control diet), and **(d)** strain median lifespan (per day increase in median longevity). CpGs that were significantly associated with age are denoted by colored markers (red circles: age-gain; yellow triangles: age-loss). **(e)** Overlap among the lists of differentially methylated CpGs. **(f)** Each dot represents the mean methylation beta-values for the 5030 age-gain, and 1523 age-loss CpGs. **(g)** Correlation between body weight and methylation beta-values for the CpG (cg10587537) located in the 3'UTR of *Mettl23*. Mice on high fat diet (HFD) have higher methylation than mice on control diet (CD), but the inverse correlation with weight is consistent for both groups (r = –0.45, p < .0001 for CD; r = –0.15, p = 0.08 for HFD). **(h)** Contour density plot for the 6553 CpGs that are significantly associated with age (age-DMCs). This relates the pattern of change with age (x-axis) with change on HFD (y-axis). CpGs that gain methylation with age are also increased in methylation by HFD. **(i)** Correlation between age and methylation at the *Mettl23* 3'UTR CpG (r = 0.35 for CD; r = 0.46). **(j)** For the 6553 age-DMCs, the contour density plot relates the pattern of change with age (x-axis) vs. change with median longevity (y-axis). CpGs that gain methylation with age have lower methylation with higher lifespan.

age (i.e., the change in methylation per day of aging) versus diet (difference in HFD relative to CD) shows that the age-gains were augmented in methylation by HFD (**Fig 2h**), and again, this is illustrated by the CpG in the *Mettl23* 3'UTR (**Fig 2i**). For the LS-DMCs, sites that had negative regression estimates for lifespan (i.e., lower DNAm per day increase in strain median longevity) had higher proportion of age-gain CpGs (**Fig 2d**). A comparison between the regression

250 estimates for age versus the regression estimates for lifespan shows that CpGs that gain
251 methylation with age tended to have lower methylation in strains with longer lifespan (**Fig 2j**).

252 As in Sziráki et al.,[43] we divided the CpGs by age effect: age-gain, age-loss, and those that do not
253 change strongly with age (age-ns; i.e., the remaining 21413 CpGs that were not classified as

Fig 3. Entropy at age-associated CpGs
Entropy values were calculated for the 5020 age-gain and 1523 age-loss CpGs separately.
For both control diet (CD) and high fat diet (HFD), there is significant increase in entropy
with age at the **(a)** age-gain and **(b)** age-loss CpGs. **(c)** The HFD mice also showed a slight
increase in entropy at CpGs that were not strongly associated with age (age-ns). **(d)** The
methylome-wide distribution of beta-values in a young adult mouse (0.6 year old; black
dashed line), and an older mouse (2.3 year old; red line); both CD mice. The young mouse
has higher peaks at the hypo-methylated (closer to 0.1) and hyper-methylated (around 0.9)
beta-values compared to the older mouse. **(e)** The HFD group has higher entropy at the age-
gain CpGs compared to the CD group. **(f)** Entropy at age-loss CpGs is higher with higher
baseline weight (BW0). **(g)** Relative effects of predictor variables on entropy shown as
logworth scores (-$\log_{10}$p). The dashed lines correspond to p = 0.01. Positive values indicate
positive regression estimates (for diet, positive value means higher in HFD). BWF is final
weight; Chol is serum total cholesterol; Gluc is fasted glucose levels; LS is the strain median
lifespan. **(h)** The residual plot (adjusted for age, diet, BWF, glucose, cholesterol, and batch)
shows the inverse association between entropy at age-gain sites, and lifespan. Similar
residual plots show the association between **(i)** BWF and age-loss entropy, and **(j)** between
fasted serum glucose and age-gain entropy.

age-DMCs). For these conserved CpGs, both sets of age-DMCs had significant increases in
entropy with age regardless of diet (**Fig 3a**, **3b**), and even the age-ns showed a modest entropy
gain with age in the HFD group (**Fig 3c**). The reason for this increase in disorder becomes
evident when we compare the density plots using the full set of CpGs for one of the younger
mice (UT319; 0.56 years old) and one of the older mice (UT573; 2.3 years) (**Fig 3d**). Concordant
with previous reports,[43,45] the older sample showed a subtle flattening of the bimodal peaks
towards a slightly more hemi-methylated state. The entropy of the age-gain CpGs was modestly
but significantly higher in the HFD group (p = 0.001; **Fig 3e**). Entropy of the age-loss and age-ns
CpGs were not different between the diets. Body weight on the other hand, was associated
specifically with the entropy score of the age-loss CpGs, and both higher BW0 (**Fig 3f**) and BWF
predicted higher entropy for age-loss CpGs.

We applied a multivariable regression to compare the relative effects of age, diet, BWF,
glucose, cholesterol, and strain median lifespan (**Fig 3g**; full statistics in **Data S6**). Entropy of
age-gain CpGs was increased by HFD but was not associated with BWF. Strain median lifespan
showed a significant inverse correlation with the entropy of age-gain CpGs with an explained
variance of 6% (**Fig 3h**). Entropy of the age-loss CpGs had a significant positive correlation with
BWF (**Fig 3i**), but was not associated with diet, and also had a modestly significant inverse
correlation with median lifespan. Cholesterol was unrelated to the entropy values. Glucose on
the other hand, showed a significant inverse association with entropy of both the age-gain (**Fig
3j**) and age-loss CpGs, and this suggests slightly lower entropy with higher fasted glucose.

Taken together, our results show that the conserved CpGs are influenced by multiple
predictors. HFD augmented the age-dependent changes with a prominent effect on age-gain
CpGs. Body weight showed a strong association with the age-loss CpGs. Additionally, strains
with longer life expectancy tended to have lower methylation levels at age-gain CpGs with an
overall lower entropy state at these CpGs that suggests a more "youthful" methylome for
longer lived genotypes.

Functional and genomic context of DMCs

To uncover the potential biological pathways represented by the DMCs, we performed genomic
regions enrichment analyses for the CpGs.[46] The age-gain CpGs were highly enriched in
transcription factors, regulators of development and growth, menarche and menstrual phases,
energy metabolism, and transcription factor networks such as HNF1 and HNF3B pathways (**Data
S7**). The age-loss CpGs had somewhat modest enrichment, and represented cell adhesion and
cytoskeletal processes, endothelial cell proliferation, and p38 signaling (**Data S7**). The BW-DMCs
were enriched in actin and protein metabolism, and WNT, and platelet-derived growth factor
(PDGF) and ErbB signaling. Similarly, the diet-DMCs were highly enriched in PDGF, epidermal
growth factor (EGFR) and ErbB signaling, as well as the mTOR signaling pathway, and regulation
of energy homeostasis (**Data S7**). Seeming to converge on common pathways, the LS-CpGs that
were negatively correlated with lifespan had modest enrichment in cell signaling pathways such
as EGFR, PDGF, and ErbB signaling. The LS-CpGs with positive correlation with lifespan were
highly enriched in lipid metabolic genes, and also included pathways related to chromosome
maintenance and telomere expansion (**Data S7**).

We next examined the genomic annotations and chromatin states of the DMCs (**Data S8**).
Consistent with previous reports,[39,43] age-gain CpGs were enriched in promoter and 5'UTR
CpGs, but depleted in 3'UTR, exon, and intergenic CpGs (**Data S8; Fig S3a**). Diet- and weight-
DMCs were depleted in promoter regions, and enriched in exons and 3'UTR, and along with the
age-loss CpGs, enriched in introns. For chromatin states, we annotated the CpG regions using
the 15-states chromatin data for neonatal (P0) mouse liver (**Data S8** has annotations for each
site).[47,48] Also included were regions labelled as No Reproducible State or NRS; i.e., regions that
were not replicated.[48] Compared to the array content as background, the age-gain CpGs were
selectively enrich in polycomb associated heterochromatin (Hc-P) and bivalent promoters (Pr-
Bi), chromatin states that were highly depleted among the other DMCs (**Fig S3b**; **Data S8**). In
contrast, strong and permissive transcription sites (Tr-S and Tr-P, respectively) were depleted
among the age-gain CpGs, and enriched among the BW- and diet-DMCs (**Fig S3b**). Age-loss CpGs
were enriched in Tr-P and Tr-I (transcription initiation). Distal enhancers (strong distal or En-Sd,
and poised distal or En-Pd) were also highly enriched among the BW- and diet-DMCs, and also
showed some enrichment among the age-DMCs (**Fig S3b**).

For an overview of the general methylation and variance patterns by chromatin annotations,
we used the full set of 27966 CpGs and computed the average methylation beta-values, and
average regression coefficients (i.e, change in beta-value per unit change in the respective
predictor variable, or contrast between diets). As expected, promoter CpGs and Hc-P were sites
with the lowest methylation. Hc-H, Tr-S, and Tr-P had higher methylation, and many of the
enhancer sites were in the hemi-methylated zone (**Fig S3c, d, e**). For age effect, mean
regression estimates had a significant inverse linear fit with mean methylation ($r = -0.63$, $p =$
0.009; **Fig S3c**) and this is consistent with the greater age-loss at hypermethylated CpGs, and
greater age-gains at hypomethylated CpGs (**Fig 2f**). The effects of diet and weight were not
linearly related to the mean methylation of the chromatin states. Instead, both showed a U-
shaped fit with a significant negative quadratic effect for diet ($R^2 = 0.69$, $p = 0.0005$, quadratic
estimate = -0.05; **Fig S3d**), and a positive quadratic effect for weight ($R^2 = 0.50$, $p = 0.01$,
quadratic estimate = 0.001; **Fig S3e**). Methylation variation as a function on strain longevity did
not relate to mean methylation with either a linear or polynomial fit, and indicates that
variance due to background genotype is less dependent on the chromatin and mean
methylation status. While this is a very low-resolution and broad view of methylation levels and
methylation variation, the observations show that while aging results in erosion of the hypo-
and hypermethylated peaks, diet and body weight appear to have generally stronger
associations with hemi-methylated sites.

Genetic analysis of epigenetic age acceleration

The EAA traits had moderate heritability at an averaged H^2 of 0.50 (**Table 2**).[34] Another way to
gauge level of genetic correlation is to compare between members of strains maintained on
different diets. All the EAA traits shared high strain-level correlations between diets, indicating
an effect of background genotype that is robust to dietary differences (**Table 2**). The

methylome-wide entropy had a heritability of ~0.30, and had no strain-level correlation
between diets.

Fig 4. QTL maps for the DNAm readouts
The Manhattan plots represent the location of genotyped markers (x-axis), and linkage –
log₁₀p (y-axis). **(a)** The peak quantitative trait locus (QTL) for age acceleration from the
pan-tissue interventional clock (int.EAA) is on chromosome (Chr) 11 at ~93 Mb. The inset
shows the mean (± standard error) trait values for BXDs the are homozygous for the
C57BL/6J allele (*BB*; grey) versus BXDs homozygous for the DBA/2J allele (*DD*; black) on
control diet (CD) and high fat diet (HFD). **(b)** The liver-specific int.EAA has a peak QTL on
Chr19 (~38 Mb). Trait means by genotype at this locus are shown in inset; *BB* has higher
age acceleration. **(c)** Linkage statistics are weaker for the methylome-wide entropy.
However, there is a nominally significant linkage on the Chr19 loci, but the peak markers
are at ~47.5 Mb. Here the *BB* genotype has higher entropy.

To uncover genetic loci, we applied QTL mapping using mixed linear modeling that corrects for
the BXD kinship structure.[49] First, we performed the QTL mapping for each EAA traits with
adjustment for diet and body weight. EAA from the two interventional clocks had the strongest
QTLs (**Data S9**). The pan-tissue int.EAA had a significant QTL on Chr11 (90–99 Mb) with the
highest linkage at ~93 Mb (p = 3.5E-06; equivalent to a LOD score of 4.7) (**Fig 4a**). Taking a
genotype marker at the peak interval (BXD variant ID DA0014408.4 at Chr11, 92.750 Mb)[34], we

segregated the BXDs homozygous for either the D2 (*DD*) or the B6 (*BB*) alleles. Strains with *DD*
genotype at this locus had significantly higher int.EAA (**Fig 4a** inset). The liver int.EAA had the
highest QTL on Chr19 (35–45 Mb) with the most significant linkage at markers between 38–42
Mb (p = 9E-07; LOD score of 5.2) (**Fig 4b**). We selected a marker at the peak interval
(rs48062674 at Chr19, 38.650 Mb), and the *BB* genotype had significantly higher liver int.EAA
compared to *DD* (**Fig 4b** inset).

We performed a similar QTL mapping for methylome-wide entropy with adjustment for major
covariates (diet, chronological age, and body weight). There were no genome-wide significant
QTLs. A region on Chr19 that overlapped the liver int.EAA showed a modest peak (**Fig 4c**; **Data**
S9). However, the peak markers for entropy were located slightly distal to the peak EAA QTL
(~47.5 Mb at rs30567369, minimum p = 0.0005). At this locus, the *BB* genotype had higher
average entropy.

To identify regulatory loci that are consistent across the different EAA measures, we applied a
multi-trait analysis and derived the linkage meta-p-value using a p-value combination for the six
EAA traits.[50] The peaks on Chrs 11 and 19 attained the highest consensus p-values (**Fig S4a;**
Data S9). There was another potential consensus peak at combined $-\log_{10}p > 6$ on Chr3 (~54
Mb). We focus on the Chrs 11 and 19 QTLs and refer to these as *EAA QTL on Chr 11* (*Eaa11*),
and *EAA QTL on Chr 19* (*Eaa19*). *Eaa11* extends from 90–99 Mb. For *Eaa19*, we delineated a
broader interval from 35–48 Mb that also encompasses the peak markers for entropy.

We performed marker-specific linkage analyses for each of the clocks using a regression model
that adjusted for diet. With the exception of the liver int.EAA, all the EAA traits had nominal to
highly significant associations with the representative *Eaa11* marker (DA0014408.4), and the
DD genotype had higher age acceleration (**Table 3**). Mean plots by genotype and diet shows
that this effect was primarily in the CD mice (**Fig S4b**). The effect of this locus appeared to be
higher for the pan-tissue clocks compared to the corresponding liver-specific clocks. For
proximal *Eaa19*, the representative marker (rs48062674) was associated with all the EAA traits
and the *BB* mice had higher age acceleration on both diets (**Fig S4c**). We also tested if these
peak markers were associated with the recorded lifespan phenotype and we found no
significant association with the observed lifespan of the BXDs.

Table 3: Marker specific linkage analyses for epigenetic age acceleration and body weight
trajectory

Predictor	Outcome	Linear regression[1]			
		Estimate	Std Error	t Ratio	p
Eaa11 DA0014408.4[*DD*] Chr11, 92.750 Mb (133 *BB* cases, and 173 *DD* cases)	EAA, pan	0.096	0.023	4.184	3.8E-05
	EAA, liver	0.067	0.017	3.880	0.0001
	dev.EAA, pan	0.077	0.025	3.041	0.003
	dev.EAA, liver	0.037	0.020	1.878	0.06
	int.EAA, pan	0.153	0.029	5.278	2.5E-07
	int.EAA, liver	-0.033	0.025	-1.284	0.20
Eaa19 rs48062674[*DD*] Chr19, 38.650 Mb	EAA, pan	-0.083	0.028	-2.954	0.003
	EAA, liver	-0.137	0.020	-6.972	2.0E-11
	dev.EAA, pan	-0.206	0.029	-7.218	4.3E-12

		Estimate	Std Error	t Ratio	p
(238 *BB* cases, and 67 *DD* cases)	dev.EAA, liver	-0.124	0.023	-5.461	9.9E-08
	int.EAA, pan	-0.143	0.035	-4.028	7.1E-05
	int.EAA, liver	-0.250	0.027	-9.238	4.6E-18
Mixed model for longitudinal change in body weight[2]					
Predictor	**Outcome**	**Estimate**	**Std Error**	**t Ratio**	**p**
Eaa11 DA0014408.4[DD] Number of observations = 6885; number of individuals = 2112	Body weight	0.619	0.345	1.794	0.07
Eaa19 rs48062674[DD] Number of observations = 6132; number of individuals = 1852	Body weight	-1.847	0.374	-4.945	7.6E-07

373 [1]Regression model: lm(EAA ~ genotype + diet); [2]lmer(weight ~ age + diet + genotype + (1|mouseID)

Association of EAA QTLs with body weight trajectory

Since gain in body weight with age was an accelerator of the clocks, we examined whether the
selected markers in *Eaa11* and *Eaa19* were also related to body weight change. We retrieved
longitudinal weight data from a larger cohort of the aging BXD mice that were weighed at
regular intervals. After excluding heterozygotes, we tested the effect of genotype. Concordant
with the higher EAA for the *DD* genotype at *Eaa11* in the CD group, the *DD* genotype in the CD
group also had slightly higher mean
weight at older adulthood (12 and 18
months; **Fig 5a**). However, this marker
had no significant association with
body weight when tested using a
mixed effects model (p = 0.07; **Table**
3). In *Eaa19*, it was the *BB* genotype
that consistently exhibited an
accelerated clock on both diets, and
also higher entropy, and the *BB*
genotype had higher average body
weight by 6 months of age (**Fig 5b**),
and this locus had a significant
influence on the body weight
trajectory (p = 7.6E-07; **Table 3**).

Candidate genes for epigenetic age acceleration

397 There are several positional candidate
398 genes in *Eaa11* and *Eaa19*. To narrow
399 the list, we applied two selection

Fig 5. Body weight trajectory by diet and genotype
Body weight was measured at regular age intervals
(x-axis) from **(a)** 2112 BXD mice that were
homozygous at the *Eaa11* marker (DA0014408.4;
842 *BB*, 1279 *DD*), and **(b)** 1852 BXD mice that were
homozygous at the proximal *Eaa19* marker
(rs48062674; 1252 *BB*, 600 *DD*). Mice were
maintained on either control diet (CD) or high fat
diet (HFD). The graphs show the segregation of
body weight over time by diet and genotype. Mean
± standard error; heterozygotes were excluded.

criteria: genes that (1) contain missense and/or stop variants, and/or (2) contain non-coding variants and regulated by cis-acting expression QTLs (eQTL). For the eQTL analysis, we utilized an existing liver transcriptome data from the same aging cohort.[32] We identified 24 positional candidates in *Eaa11* that includes *Stxbp4*, *Erbb2* (*Her-2* oncogenic gene), and *Grb7* (growth factor receptor binding) (**Data S10; Fig 4a**). *Eaa19* has 81 such candidates that includes a cluster of cytochrome P450 genes, and *Chuk* (inhibitor of NF-kB) in the proximal region, and *Pcgf6* (epigenetic regulator) and *Elovl3* (lipid metabolic gene) in the distal region (**Data S10; Fig 4b, 4c**).

For further prioritization, we converted the mouse QTL regions to the corresponding syntenic regions in the human genome, and retrieved GWAS annotations for these intervals.[51] We specifically searched for the traits: epigenetic aging, longevity, age of menarche/menopause/puberty, Alzheimer's disease, and age-related cognitive decline and dementia. This highlighted 5 genes in *Eaa11*, and 3 genes in *Eaa19* (**Table S3**). We also identified a GWAS study that found associations between variants near *Myof-Cyp26a1* and human longevity,[41] and a meta-GWAS that found gene-level associations between *Nkx2–3* and *Cutc*, and epigenetic aging (**Table S3**).[27]

Gene expression correlates of EAA

A subset of the BXD cases had liver RNA-seq data (94 CD, and 59 HFD).[32] Using this set, we performed transcriptome-wide correlation analysis for the general pan-tissue EAA, and the more specific liver int.EAA. To gain insights into biological pathways, we selected the top 2000 transcriptome correlates for functional enrichment analysis (**Data S11**). The common themes for both clocks were: (1) there were far fewer negative correlates (223 out of 2000 for pan-tissue EAA, and 337 out of 2000 transcripts for liver int.EAA) than positive correlates, (2) the negative correlates were highly enriched (Bonferroni correct $p < 0.05$) in oxidation-reduction and mitochondrial genes (**Data S12, Data S13**). The pan-tissue general clock was also highly enriched in pathways related to steroid metabolism, epoxygenase p450 pathway, and xenobiotics, which are pathways that are particularly relevant to liver. The p450 genes included candidates that are in *Eaa19* (e.g., *Cyp2c29*, *Cyp2c37*). The positive correlates were enriched in a variety of gene functions including mitosis for both clocks, and immune and inflammatory response for the general pan-tissue clock (functions that are not specific to liver). 563 transcripts (315 unique genes) were correlated with both the pan-tissue EAA, and the liver int.EAA. Based on hierarchical clustering (HC) of these common mRNA correlates of EAA, the transcripts could be clustered into 3 groups (**Fig 6a;** heatmap in **Fig S5a**). While none of these were significantly enriched in any particular gene ontology (GO), cluster 3 included several oxidation-reduction genes including the *Eaa11* candidate, *Cyp2c29*, and cluster 2 included several cell cycle genes (**Fig 6a**). To verify that these transcriptomic associations are robust to the effect of diet, we repeated the correlation and enrichment analysis in the CD group only for the pan-tissue general clock (n = 94). Again, taking the top 2000 correlates ($|r| \le 0.22$; $p \le 0.03$), we found the same enrichment profiles for the positive correlates (immune, cell cycle) and the negative correlates (oxidation-reduction and mitochondrial) (**Data S12**).

Liver proteome was also available for 164 of the BXDs, and 53 also had adipose proteome. The liver proteome data quantifies over 32000 protein variants from 3940 unique genes and has

Fig 6. Gene expression correlates of epigenetic age acceleration
(a) mRNAs that were correlated with the acceleration of both the pan-tissue general clock (pan EAA), and the liver interventional clock (liver int.EAA) were grouped based on unsupervised hierarchical clustering (HC). Few representative genes and gene ontologies are highlighted. For liver proteome, the level of APOE was the strongest correlate for both the **(b)** liver int.EAA, and **(c)** the pan-tissue EAA. **(d)** For liver proteins that were correlated with both pan-tissue EAA and liver int.EAA, HC grouped the proteins into clusters that were enriched in oxidation-reduction and lipid metabolism, and a cluster enriched in glycogen metabolism. In adipose tissue, the expression level of the APOE protein was higher with higher age acceleration for both the **(e)** liver int.EAA, and **(f)** the pan-tissue EAA.

been reported in Williams et al.[32] Similar to the transcriptome-wide analysis, we extracted the
top 2000 protein correlates of EAA (**Data S14**), and performed functional enrichment analysis
(**Data S12, S13**). For both the liver int.EAA and the pan-tissue EAA, the top liver protein
correlate was APOE, and higher expression of APOE was associated with higher age acceleration
(**Fig 6b, c**). Similar to the transcriptome, the negative correlates of EAA were highly enriched in
oxidation-reduction (several cytochrome proteins), steroid metabolism, and epoxygenase 450
pathway. The positive correlates were also highly enriched in oxidation-reduction (several
hydroxy-delta-5 steroid dehydrogenases proteins), lipid and carbohydrate metabolism, as well
as phospholipid efflux (particularly enriched for the liver int.EAA) (**Data S13**). There was a high
degree of overlap at the proteomic level for the two clocks and 1241 proteins variants (332
unique genes) were correlated with both the pan-tissue EAA and the liver int.EAA (**Data S14**).
For these common protein correlates, the HC divided the proteins into clusters that

represented metabolic pathways mainly related to steroid metabolism, but also glycolysis and
gluconeogenesis (**Fig 6d**; heatmap in **Fig S5b**).

Finally, we used the adipose proteome data for a proteome-wide correlational analysis for the
pan-tissue EAA and liver int.EAA. We took only the top 1000 correlates (due to the small sample
size), and a functional enrichment analysis showed consistent enrichment in metabolic
pathways related to fatty acids and also carbohydrates, and cell proliferation genes for the pan-
tissue EAA (**Data S12, S13**). For the adipose proteome, the cytochrome p450 genes were no
longer enriched. However, the overall functional profile highlighted metabolic pathways as
important gene expression correlates of EAA. Furthermore, for both the liver and adipose
proteomes, APOE levels were highly correlated with EAA that indicates a higher level of this
apolipoprotein in both tissues is associated with higher age acceleration (**Fig 6e, 6f**).

Discussion

Here we have tested the performance of DNAm clocks derived from highly conserved CpGs, and
described the dynamism and variability of site-specific methylation. While age is a major source
of variance, we detected joint modulation by diet, body weight, and genotype-by-diet life
expectancy. HFD had an age accelerating effect on the clocks, and this is concordant with our
previous report where we found more rapid age-associated changes in methylation.[39] This also
concurs with studies in humans that have found that obesity accelerates epigenetic aging.[52,53]
However, when BWF was included in the regression term, the effect of diet became
inconsistent. This suggests that the effect of diet on EAA is mediated by the changes in weight
and metabolic traits such as total cholesterol. Body weight in particular, had a strong age-
accelerating effect. The effect of weight may manifest early on, and even in the CD group,
higher weight gains at younger age (between 4–6 months) was associated with higher EAA later
in life.

We tested different mouse DNAm clocks, and the main difference between these clocks was
the subsets of CpGs that were used for training. It is well-known that DNAm clocks have high
level of degeneracy.[3,14] In other words, highly accurate predictors of chronological age can be
built from entirely different sets of CpGs and different weight coefficients. This is likely because
a large proportion of CpGs undergo some degree of change with age, and combinatorial
information from any subset of this is informative of age. For instance, even at a very stringent
cutoff of Bonferroni 0.05 that treated the 27966 CpGs as "independent", we still detected 6553
CpGs as age-DMC, i.e., close to a quarter of the CpGs we tested. Clocks built from pre-selected
CpGs that are at conserved sequences are known to be sensitive to the effects of pro-longevity
interventions such as calorie restriction and growth hormone receptor deletion.[3,54] And while
all these DNAm clocks achieve reasonably high prediction of chronological age, the age
divergence derived from these different clocks (EAA) can capture slightly different facets of
biological aging, and the better a clock is at predicting chronological age, the lower its
association with mortality risk.[13,14] In the present study, we find that the interventional clocks
deviated most from chronological age, and this is expected as these were built from a much
smaller set of CpGs (see Methods). The interventional clocks were also associated with BWF
and cholesterol, but had weaker associations with BW0. The liver int.EAA had the highest
positive correlation with methylome-wide entropy, and was the clock that had the strongest

inverse correlation with strain longevity. In contrast, the developmental clocks, which were
based on CpGs that change early in life, showed a stronger association with BW0. The contrast
between the interventional and developmental clocks suggests that while one is more
modifiable, the other is more informative of baseline characteristics that influence aging later in
life. The pan-tissue clock, which was not constrained to any preselected set of CpGs or tissue,
also performed well in capturing biological aging and was accelerated by both BW0 and BWF,
diet (when BW0 was the weight term in the regression model), higher entropy, and had a
modest but significant inverse correlation with strain lifespan.

Entropy, a measure of noise and information loss, increases as a function of time and age.[10,55-57]
In the context of the methylome, the shift to higher entropy represents a tendency for the
highly organized hypo- and hypermethylated landscape to erode towards a more hemi-
methylated state.[10,43,45] This increase in disorder, particularly across CpGs that are highly
conserved, could have important functional consequences. The entropy of age-gain CpGs
predicted strain lifespan, and was increase by HFD. Overall, we find that mice belonging to
longer-lived BXD strains had a more "youthful" methylome with lower entropy at the age-gain
CpGs. The entropy of age-loss CpGs on the other hand, was related to the body weight of mice,
and both higher BW0 and BWF were associated with higher entropy. This leads us to suggest
that the rate of noise accumulation, an aspect of epigenomic aging, can vary between
individuals, and the resilience or susceptibility to this shift towards higher noise may be partly
modulate by diet as well as genetic factors.

Somewhat surprising was the inverse correlation between the entropy of age-DMCs and fasted
glucose. This lower entropy of age-gain CpGs with higher glucose is somewhat counter to the
general tendency for strains with shorter lifespan to have higher glucose.[33] In biological
systems, entropy is kept at bay by the uptake of chemical energy, and investment in
maintenance and repair,[57] and we can only speculate that at least in mice, the higher amount of
glucose after overnight fast may be associated with a more ordered methylome. The centrality
of bioenergetics for biological systems may explain why we detect this coupling between the
DNAm readouts (i.e., the clocks, and entropy), and indices of metabolism including weight, diet,
levels of macronutrients, and even expression of metabolic genes. As cogently highlighted by
Donohoe and Bultman,[58] many metabolites (e.g., SAM, NAD$^+$, ATP) are essential co-factors for
enzymes that shape the epigenome, and these could serve as nutrient sensors and mechanistic
intermediaries that regulate how the epigenome is organized in response to metabolic
conditions. Close interactions between macro- and micronutrients, and DNAm is a conserved
process and plays a critical role in defining both physiology and body morphology.[59,60] Overall,
our results suggests that a higher metabolic state is associated with higher entropy and EAA,
and potentially, lower lifespan.

For the BXDs, life expectancy is highly dependent on the background genotype, and mean
lifespan varies from under 16 months for strains such as BXD8 and BXD13, to over 28 months in
strains such as BXD91 and BXD175.[33,36,38] The EAA showed the expected inverse correlation
with lifespan, but the effect was modest and only significant for the pan-tissue EAA and the
liver int.EAA. The association of lifespan with the entropy of age-gain CpGs was slightly
stronger. We must point out that the analysis between the epigenetic readouts and lifespan
was an indirect comparison. Unlike the comparison with body weight and metabolic traits,

which were traits measured from the same individual, the lifespan data are strain
characteristics computed from a parallel cohort of mice that were allowed to survive till natural
mortality, and this may partly explain the weaker associations with EAA. Nonetheless, our
observations indicate that genotypes with higher life-expectancy have generally lower entropy,
and lower methylation levels at the age-gain CpGs, and these properties of the methylome are
likely to be partly under genetic modulation.

Our goal was to take these different clocks and identify regulatory loci that were the most
stable and robust to the slight algorithmic differences in building the clocks. A notable
candidate in *Eaa11* is Syntaxin binding protein 4 (*Stxbp4*, aka, *Synip*), located at 90.5 Mb. *Stxbp4*
is a high-priority candidate due to the concordant evidence from human genetic studies. The
conserved gene in humans is a replicated GWAS hit for the intrinsic rate of epigenetic
aging.[24,26,27] In the BXDs, *Stxbp4* contains several non-coding variants, and a missense mutation
(rs3668623), and the expression of *Stxbp4* in liver is modulated by a *cis*-eQTL. *Stxbp4* plays a
key role in insulin signaling,[61] and has oncogenic activity and implicated in different cancers.[62,63]
Furthermore, GWAS have also associated *STXBP4* with age of menarche.[64,65] *Eaa11* corresponds
to the 17q12-21 region in humans, and the location of additional oncogenic genes, e.g.,
ERBB2/HER2, *GRB7*, and *BRCA1*.[66] The mouse *Brca1* gene is a little distal to the peak QTL region
and is not considered a candidate here, although it does segregate for two missense variants in
the BXDs. *Erbb2* and *Grb7* are in the QTL region, and *Erbb2* contains a missense variant
(rs29390172), and *Grb7* is modulated by a *cis*-eQTL. *Nr1d1* is another candidate in *Eaa11*, and
the co-activation of *Erbb1*, *Grb7*, and *Nr1d1* has been linked to breast and other cancers.[67,68]

Eaa19 was consistently associated with EAA from all the clocks we evaluated, and also with
body weight gains, irrespective of diet. DNAm entropy may also have a weak association with
markers at this interval. The EAA traits have peak markers in the proximal part of *Eaa19*
(around the cytochrome cluster), and the methylome-wide entropy had a weak peak that was
in the distal portion (over candidates like *Elovl3*, *Pcgf3*). Two candidates in *Eaa19* have been
implicated in epigenetic aging in humans based on gene-level meta-GWAS: NK homeobox 3
(*Nkx2-3*, a developmental gene), and CutC copper transporter (*Cutc*).[27] *Eaa19* is also the
location of the *Cyp26a1-Myof* genes, and the human syntenic region is associated with
longevity, metabolic traits, and lipid profiles.[41,69,70] Another noteworthy candidate in *Eaa19* is
Chuk, a regulator of mTORC2, that has been associated with age at menopause.[64,71] *Eaa19*
presents a complex and intriguing QTL related to the DNAm readouts that may also influence
body weight gains over the course of life. Both *Eaa19* and *Eaa11* exemplify the major challenge
that follows when a genetic mapping approach leads to gene- and variant-dense regions.[72,73]
Both loci have several biologically relevant genes, and identifying the causal gene (or genes) will
require a more fine-scaled functional genomic dissection.

The gene expression analyses highlighted metabolic pathways. At the mRNA level, the negative
correlates of EAA were highly enriched in metabolic genes related to oxidation-reduction and
steroid metabolism, while the positive correlates were enriched in pathways related to mitosis,
and immune response for the pan-tissue general EAA. This convergence on metabolic, immune
and cell division genes is very consistent with previous reports.[14,28,44] Here we should note that
depending on the tissue(s) in which the clocks are trains, and the tissue from which the
DNAmAge is estimated, the EAA derivative may put an emphasis on biological pathways or

genes that are most relevant to that tissue. For instance, clocks optimized for neural tissue are
more closely related to neurodegeneration and neuropathologies.[18,74] With the liver clocks,
expression correlates highlighted aspects of metabolism that are relevant to liver function (e.g.,
the cytochrome p450 epoxygenase genes), and this is detected both at the transcriptomic, and
proteomic levels. For the adipose tissue proteome, the cytochrome genes become less
prominent, but the enriched pathways still remained consistent (i.e., oxidation-reduction, lipid
and carbohydrate metabolism, and cell proliferation for the positive correlates of the pan-tissue
EAA). At the proteome level, we also find several phospholipid efflux genes (APOC1, APOA2,
APOC3, APOA1, APOA4, APOE) that are positive correlates of EAA. For both the liver and
adipose proteomes, APOE stands out as the top protein correlate of EAA. A recent human study
has also identified the APOE locus as the strongest GWAS hit for two measures of biological age
acceleration (the phenoAge, and the bioAge).[28] While more specific to liver, the cytochrome
P450 genes presents as both positional candidates, and expression correlates of EAA. These
genes have high expression in liver, and have major downstream impact on metabolism.[75-77]
One caveat is that these CYP genes are part of a gene cluster in *Eaa19* that includes transcripts
with *cis*-eQTLs (e.g., *Cyp2c66*, *Cyp2c39*, *Cyp2c68*), and the tight clustering of the genes, and
proximity of trait QTL and eQTLs may result in tight co-expression due to linkage
disequilibrium.[78] Nonetheless, the cytochrome genes in *Eaa19* are strong candidate modulators
of EAA derived from liver tissue that calls for further investigation.

Aside from *Eaa11* and *Eaa19*, another locus with evidence for consensus QTL was detected on
Chr3. We do not delve into this in the present work, but the Chr3 interval is near genes
associated with human epigenetic aging (*Ift80*, *Trim59*, *Kpna4*).[24,27] However, this QTL is
dispersed across a large interval, and the peak markers do not exactly overlap these human EAA
GWAS hits. While we have focused on *Eaa11* and *Eaa19*, the Chr3 locus presents a potentially
important region for EAA.

In summary, we have identified two main QTLs—*Eaa11* and *Eaa19*—that contribute to variation
in EAA. *Eaa11* contains several genes with oncogenic properties (e.g., *Stxbp4*, *Erbb2*), while
Eaa19 contains a dense cluster of metabolic genes (e.g., *Elovl3*, *Chuk,* the cytochrome genes).
We demonstrate that metabolic profile and body weight are closely related to epigenetic aging
and methylome entropy. The convergence of evidence from genetic and gene expression
analyses indicates that genes involved in metabolism and energy balance contribute to the age-
dependent restructuring of the methylome, which in turn forms the basis of the epigenetic
clocks.

Materials and Methods

Biospecimen collection and processing

Samples for this study were selected from a larger colony of BXD mice that were housed in a
specific pathogen-free (SPF) facility at the University of Tennessee Health Science Center
(UTHSC). All animal procedures were in accordance with a protocol approved by the
Institutional Animal Care and Use Committee (IACUC) at the UTHSC. Detailed description of
housing conditions and diet can be found in.[32,33] Mice were given *ad libitum* access to water,
and either standard laboratory chow (Harlan Teklad; 2018, 18.6% protein, 6.2% fat, 75.2%

carbohydrates), or high-fat chow (Harlan Teklad 06414; 18.4% protein, 60.3% fat, 21.3%
carbohydrate). Animals were first weighed within the first few days of assignment to either
diets, and this was mostly but not always prior to introduction to HFD. Following this, animals
were weighed periodically, and a final time (BWF) when animals were humanely euthanized
(anesthetized with avertin at 0.02 ml per g of weight, followed by perfusion with phosphate-
buffered saline) at specific ages for tissue collection. The present work utilizes the biobanked
liver specimens that were pulverized and stored in -80 °C, and overlaps samples described in
Williams et al.[32] DNA was extracted using the DNeasy Blood & Tissue Kit from Qiagen. Nucleic
acid purity was inspected with a NanoDrop spectrophotometer, and quantified using a Qubit
fluorometer dsDNA BR Assay.

Methylation array and quality checks

DNA samples from ~350 BXD mice were profiled on the Illumina HorvathHumanMethylChip40
array. Samples were in 96-well plate format (**Data S1**), and the plates were randomized for
major covariates such as age and diet. Details of this array are described in Arneson et al.[29] The
array contains probes that target ~36K highly conserved CpGs in mammals. Over 33K probes
map to homologous regions in the mouse genome. For downstream statistical tests, we further
filtered the probes and used only 27966 probes that have been validated for the mouse
genome using calibration data generated from synthetic mouse DNA.[29] Data was normalized
using the SeSame method.[79] Unsupervised HC was performed to identify outliers and failed
arrays, and those were excluded. We also performed strain verification as an additional quality
check. While majority of the probes were free of DNA sequence variants, we found 45 probes
that overlapped variants in the BXD family. We leveraged these as proxies for genotypes, and
performed a principal component analysis (PCA). The top genotype principal components
(genoPC1 and genoPC2; **Data S1**) segregated the samples by strain identity, and samples that
did not cluster close to the reported strains were removed. After excluding outliers, failed
arrays, and samples that failed strain verification, the final liver DNAm data consisted of 339
samples. The beta-values for these ~28K probes in the 339 samples show the expected bimodal
distribution (**Fig S6a**), but for these highly conserved CpGs, we note a much higher
representation of hypermethylated CpGs instead of the slightly hypomethylated state of the
methylome when a wider spectrum of CpGs is assayed.[43]

BXD-unbiased mouse clock estimation

Three different mouse clocks are reported here, and all three are based on penalized regression
modeling using glmnet.[80] Training was done in a larger mouse dataset that excluded the
BXDs.[30,31,42] The clocks are therefore unbiased to the characteristics of the BXDs. For pan-tissue
clocks, all mouse samples were used for training. For the liver specific clocks, the training was
limited to data from liver samples.

The general DNAmAge clock did not preselect for any CpGs and the full set of CpGs that map to
Mus musculus was used. First, a log-linear transformation was applied to the chronological age
using the function:

$$f(Age) = \begin{cases} \dfrac{Age}{1.2 + 0.06} + \log(1.2 + 0.06) - \dfrac{1.2}{1.2 + 0.06}, Age > 1.2 \\ \log(Age + 0.06), Age \leq 1.2 \end{cases}$$

This is similar to the age transformation described in the original Horvath pan-tissue human
clock, but with offset at 0.06, and adult mouse age at 1.2.[11] Following this transformation, an
elastic net regression was implemented to regress the transformed chronological age on the
CpG beta-values in the training data. The alpha was set at 0.5, and the optimal lamda
parameter was determined by 10-fold cross-validation (function cv.glmnet). This selected
subsets of clock CpGs and coefficients (see **Data S2** for the lists of clock CpG, intercepts, and
coefficients). DNAmAge was then calculated as:

$$DNAmAge = f^{-1}\left(\frac{b_0 + b_1 CpG_1 + b_2 CpG_2 + \cdots + b_i CpG_i}{b_0 + b_1 + b_2 + \cdots + b_i}\right)$$

where b_0 is the intercept, and b_1 to b_i are the coefficients, and CpG_1 to CpG_i denote the beta-
values for the respective clock CpGs, and $f^{-1}()$ denotes the inverse function of $f()$.

A similar method was used to build the developmental and interventional clocks, but for these,
the CpGs were pre-selected. For the liver-specific developmental clock, CpGs that change
during mouse development was selected in liver samples based on Pearson correlation with
age in mice that were <1.6 months old. The top 1000 negative and top 1000 positive correlates
were then classified as "developmental CpGs", and the training was done using only this subset
of CpGs. For the pan-tissue dev.DNAmAge, the top 1000 positive and top 1000 negative
developmental CpGs were based on a multi-tissues EWAS, also using Pearson correlation with
age for mice <1.6 months old, and these are CpGs that are strongly correlated with age during
the mouse developmental period when all available tissues are considered.

Training for the interventional clock started with 537 CpGs that relate to gold-standard anti-
aging interventions (calorie restriction, growth hormone receptor knockout).[42,81] These
"interventional CpGs" were identified from an independent mouse liver calorie restriction (n =
95), and one growth hormone receptor knockout (n = 71) data that were not included in the
clock estimation.[42] Top CpGs associated with these interventions were identified and the 537
CpGs are the sites that are consistently associated with these anti-aging interventions. Of the
537, 121 CpGs increased in methylation, and 417 decreased in methylation with application of
the pro-longevity interventions. Given the small number of CpGs that went into training for the
int.DNAmAge, we expected this clock to be less correlated with chronological age, and possibly
more responsive variables such as diet.

Entropy calculation

Methylome-wide entropy was calculated from the 27966 probes. The beta-values were
discretized into 20 bins, and the Shannon entropy for each sample was estimated using the R
package, "entropy" (v1.2.1) with method = "ML": maximum likelihood.[82] The optimal number of
bins was determined using the Freedman-Diaconis rule (breaks = "FD" for the hist() function in
R). We also estimated the methylome-wide entropy after discretizing into 100 and 2000 bins
(values provided in **Data S1**), and the results we report are consistent and robust to the number

of bins. For the age-gain, age-loss, and age-ns CpGs, entropy for each set was estimated, also
following discretization into 20 bins.

Statistics

Statistical analyses were done using R or the JMP Pro software (version 15). Association
between the epigenetic predictors and continuous variables (body weight, strain lifespan,
fasted serum glucose, and total cholesterol) were based on Pearson correlations, and t-test was
used to evaluate the effect of categorical predictors (sex, diet). Multivariable regression models
were also used to control for covariates (R regression equations provided with **Table S1**, **Data**
S4, **S6**, and **Table 3**). All these traits are directly accessible from GeneNetwork 2 (GN2; more
information on how to retrieve these data from GN2 are provided in **Data S15**).[83,84] Longevity
data was obtained from a parallel cohort of BXD mice housed in the same UTHSC colony, and
members of this "longevity cohort" were allowed to age until natural death (more detail on the
longevity cohort can be found in [33]). Males were excluded and strain-by-diet lifespan summary
statistics were derived. Only strain-by-diet groups with 5 or more observations were included in
the correlational analyses with the epigenetic predictors.

Multivariable EWAS

Site-by-site differential methylation analysis (EWAS) was performed on the 27966 CpGs using a
multivariable regression model. As such genome-wide explorations are vulnerable to
unmeasured confounders, we included the top PC derived from a PCA of the 27966 probes.[85]
The top 10 principal components PCs cumulatively accounted for ~62% of the variance (**Fig**
S6b). A plot of PC1 (19% of variance) and PC2 (14% of variance) showed that PC1 captured
some noise due to batch (**Fig S6c**). The remaining top PCs (PC2 onwards) were strongly
associated with biological variables, particular age, and also weight and diet (top 10 PCs
provided in **Data S1**). For this reason, we included PC1 as a correction factor in the EWAS. The
regression model we used was: lm(CpG$_i$~ age + median lifespan + diet + BWF+ PC1), where CpG$_i$
is the ith CpG from 1 to 27966. As lifespan was from female mice, this EWAS excluded the few
male samples.

CpG annotation and enrichment

Functional annotation and enrichment analyses for the DMCs were done using the genomic
region enrichment R package, rGREAT (version 3.0.0)[46] with the array content (i.e., the 27966
CpGs) as background. Enrichment p-values are based on hypergeometric tests, and categories
with Benjamini-Hochberg adjusted p-values ≤ 0.05 are reported. Annotations were for the
GRCm38/mm10 reference genome.

For chromatin state annotation, we used bedtools to annotate the 27966 CpGs coordinates
using chromatin annotation .bed files for neonatal (P0) mouse liver tissue created by Gorkin et
al.[48,86] This provides the 15-states model using ChromHMM,[47] and we downloaded the file for
the "replicated set" (here, the regions annotated as NRS are sites that did not produce
replicable signal). Enrichment and depletion analyses for genomic annotations, and chromatin
annotations were based on the hypergeometric test (phyper R function). The R codes are
provided with the results data (**Data S8**).

Genetic analyses

The broad sense heritability within diet was estimated as the fraction of variability that was explained by background genotype.[34,87,88] For this, we applied an anova: aov(EAA ~ strain), and heritability was computed as: $H^2 = SSq_{strain}/(SSq_{strain} + SSq_{residual})$, where SSq_{strain} is the strain sum of squares, and $SSq_{residual}$ is the residual sum of squares.

All QTL mapping was done on the GN2 platform (trait accession IDs provided in **Data S15**).[83] In the GN2 home page, the present set of BXD mice belongs to the **Group: BXD NIA Longevity Study**, and GN2 provides a direct interface to the genotype data. All QTL mapping was done for genotypes with minor allele frequency ≥ 0.05 using the genome-wide efficient mixed model association (GEMMA) algorithm,[49] which corrects for the BXD kinship matrix. For the EAA traits, diet, weight at 6 months, and final weight were fitted as cofactor. Chronological age had not correlation with EAA and this was not included as a cofactor (including age does not change the results). Genome-wide linkage statistics were downloaded for the full set of markers that were available from GN2 (7320 markers in **Data S9**). For the combined p-values, QTL mapping was done separately using GEMMA for each EAA traits, then the Fisher's p-value combination was applied to get the meta-p-value.[50] We used this method to simply highlight loci that had consistent linkage across the different EAA measures. QTL mapping for methylome-wide entropy was done using GEMMA with adjustment for chronological age, diet, weight at 6 months, and final weight.

For marker specific linkage, we selected SNPs located at the peak QTL regions (DA0014408, rs48062674), and grouped the BXDs by their genotypes (F1 hybrids and other heterozygotes were excluded from this), and marker specific linkage was tested using ANOVA and linear regression (R regression equation given in **Table 3**). rs48062674 is a reference variant that is already catalogued in dbSNP,[89] and is used as a marker in the QTL mapping. DA0014408.4 is an updated variant at a recombinant region in the Chr11 interval and within the peak QTL interval.[34] Genotypes at these markers for individual BXD samples are in **Data S1**.

To test the effect of genotype on body weight change, body weight data measured at approximately 4 (baseline), 6, 12, 18, and 24 months were downloaded from GN2 (**Data S15**). Detailed description of these weight data are in Roy et al.[33] We then applied a mixed effects regression model using the lme4 R package[90]: lmer(weight ~ age + diet + genotype + (1|ID)), where ID is the identifier for individual mouse.

Bioinformatic tools for candidate genes selection

Sequence variation between B6 and D2 in the QTL intervals (Chr11:90–99 Mb, and Chr19:35–48 Mb) were retrieved from the Wellcome Sanger Institute Mouse Genomes Project database (release 1505 for GRCm38/mm10).[91-93] Positional candidates were required to contain at least one coding variant (missense and/or nonsense variants), or have non-coding variants with evidence of *cis*-regulation in liver tissue of the BXDs. *Cis*-eQTLs for the candidate genes were obtained from the liver RNA-seq data described in [32]. An interface to search and analyze this transcriptome data is available from GN2, and is catalogued under *Group: BXD NIA Longevity Study*; *Type: Liver mRNA*; and *Dataset: UTHSC BXD Liver RNA-seq (Oct 19) TMP Log2*.

For human GWAS annotations, we navigated to the corresponding syntenic regions on the
human genome by using the coordinate conversion tool in the UCSC Genome Browser. The
Chr11 90–95 Mb interval on the mouse reference genome (GRCm38/mm10) corresponds to
human Chr17:50.14–55.75 Mb (GRCh38/hg38) (40.7% of bases; 100% span). The Chr11 95–99
Mb interval in the mouse corresponds to human Chr17:47.49–50.14 Mb (29.3% of bases, 57.9%
span), and Chr17:38.19–40.39 Mb (20.7% of bases, 44.1% span). Likewise, for the Chr19 QTL,
the mm10 35–40 Mb corresponds to hg38 Chr10:89.80–95.06 Mb (32.2% of bases, 89.2% span),
40–45 Mb corresponds to hg38 Chr10:95.23–100.98 Mb (46.6% of bases, 95.6% span), and 45–
48 Mb corresponds to hg38 Chr10:100.98–104.41 Mb (46.5% of bases, 100% span). We then
downloaded the GWAS data for these regions from the NHGRI-EBI GWAS catalogue,[51] and
retained the GWAS hits that were related to aging.

Transcriptome and proteome analyses

The liver RNA-seq data mentioned above was also used for the transcriptome-wide
correlational analysis for EAA in the 153 cases that had both DNAm and RNA-seq data. We
considered the top 2000 highest mRNA correlates ($|r|$ = 0.24, p = 0.003 for the pan-tissue EAA;
$|r|$ = 0.3, p = 0.0002 for the liver int.EAA), and the list of transcripts were collapsed to a non-
redundant list of gene symbols, and this was uploaded to the DAVID Bioinformatics Database
(version 2021 update) for GO enrichment analysis.[94,95] Proteome correlational analysis was
carried out using the data: *Group: BXD NIA Longevity Study*; *Type: Liver Proteome*; and *Dataset:*
EPFL/ETHZ BXD Liver Proteome CD-HFD (Nov19). Detailed description of this data is in Williams
et al.[32] 164 BXD cases had both DNAm and liver proteomics, and similar to the RNA-seq, we
selected the top 2000 correlates (($|r|$ = 0.24, p = 0.002 for both the pan-tissue EAA and liver
int.EAA) for enrichment analysis.

59 of the BXD cases also have proteome data from adipose tissue (*Group: BXD NIA Longevity*
Study; *Type: Adipose Proteome*; and *Dataset: Riken-Wu BXD Liver Proteome CD-HFD (Sep20)*).
While small in sample number, we used this data to test whether we could recapitulate the
same functional enrichment profiles in a different tissue. Details on sample preparation and
processing steps for the adipose proteome is provided in the dataset's "Info" page on GN2. In
brief, protein was extracted from the adipose samples by first lysis in a buffer with protease
inhibitor, followed by homogenization with a glass dounce and sonication. The protein fraction
was isolated from the homogenate by centrifugation, and processed for assay on a liquid
chromatography tandem mass spectrometry (LC-M/MS) using a modified Phase Transfer
Surfactant Method as described in Mostafa et al.[96,97] Samples were measured using a Q
Exactive Plus Orbitrap LC–MS/MS System (Thermo Fisher). For each sample, 600 ng was
injected and the samples were measured with data-independent acquisition (DIA). A portion of
the peptides from the samples were pooled and fractionated using a Pierce High pH Reversed-
Phase (HPRP) Peptide Fractionation Kit (Thermo Fisher Scientific) to generate a spectral library.
For the HPRP fractions, 450 ng was injected and the samples were measured with data-
dependent acquisition (DDA). For protein identification, the raw measurement files were
searched against a mouse database using the (uniprot-reviewed_Mus_musculus_10090_.fasta)
using Proteome Discoverer v2.4 software (Thermo Fisher Scientific). Filtered output was used to
generate a sample-specific spectral library using the Spectronaut software (Biognosys,

Switzerland). Raw files from DIA measurements were used for quantitative data extraction with
the generated spectral library, as previously described.[97] The false discovery rate was estimated
with the mProphet approach and set to 0.01 at both peptide precursor level and protein
level.[98,99] Due to the small sample size, for this dataset, we considered the top 1000 protein
correlates of EAA ($|r|$ = 0.25, p = 0.06 for the pan-tissue EAA; $|r|$ = 0.31, p = 0.02 for the liver
int.EAA).

Data availability

The full microarray data will be released via NCBI's Gene Expression Omnibus upon official
publication. Genome annotations of the CpGs can be found on Github
https://github.com/shorvath/MammalianMethylationConsortium. Individual level BXD data,
including the processed microarray data are available on www.genenetwork.org on FAIR+
compliant format; data identifiers, and way to retrieve data are described in **Data S15**.

Acknowledgement. We thank the entire UTHSC BXD Aging Colony team, particularly
Casey J Chapman, Melinda S McCarty, Jesse Ingles, and everyone else who contributed to the
tissue harvest. We thank Evan G Williams for making the gene expression data readily available,
and to David Ashbrook for providing the BXD genotypes. We thank the GeneNetwork team,
especially Zach Sloan and Arthur Centeno, who have been extremely prompt and effective at
assisting with the GeneNetwork interface. We thanks Dr. Garrett Jenkinson for the invaluable
guidance he provided for entropy estimation.

Author contributions. KM contributed to the data, conceived portion of the study, and
performed statistical analysis and drafted the article. ATL, CZL, AH contributed to the data
analysis and in computing the epigenetic clocks. JVS contributed to the lab work. YW
contributed to data generation. RWW conceived of the BXD Aging Colony, and provided access
to the BXD biospecimen and data. SH developed the array platform, and built the epigenetic
clocks and predictor. All authors contributed to, and approved the manuscript.

Research funding. This study was funded by the NIH NIA grants R21AG055841 and
R01AG043930

Competing interests. SH is a founder of the non-profit Epigenetic Clock Development
Foundation, which plans to license several of his patents from his employer UC Regents. The
other authors declare no conflicts of interest.

Ethics approval. All animal procedures were in accordance to protocol approved by the
Institutional Animal Care and Use Committee (IACUC) at the University of Tennessee Health
Science Center.

Reference:

855 1   Horvath, S. & Raj, K. DNA methylation-based biomarkers and the epigenetic clock theory
856   of ageing. *Nature reviews. Genetics* **19**, 371-384, doi:10.1038/s41576-018-0004-3
857   (2018).

858	2	Bell, C. G. *et al.* DNA methylation aging clocks: challenges and recommendations.
859		*Genome biology* **20**, 249, doi:10.1186/s13059-019-1824-y (2019).
860	3	Thompson, M. J. *et al.* A multi-tissue full lifespan epigenetic clock for mice. *Aging* **10**,
861		2832-2854, doi:10.18632/aging.101590 (2018).
862	4	Porter, H. L. *et al.* Many chronological aging clocks can be found throughout the
863		epigenome: Implications for quantifying biological aging. *Aging cell* **20**, e13492,
864		doi:10.1111/acel.13492 (2021).
865	5	Zannas, A. S. *et al.* Lifetime stress accelerates epigenetic aging in an urban, African
866		American cohort: relevance of glucocorticoid signaling. *Genome biology* **16**, 266,
867		doi:10.1186/s13059-015-0828-5 (2015).
868	6	Marioni, R. E. *et al.* The epigenetic clock is correlated with physical and cognitive fitness
869		in the Lothian Birth Cohort 1936. *International journal of epidemiology* **44**, 1388-1396,
870		doi:10.1093/ije/dyu277 (2015).
871	7	Dugue, P. A. *et al.* DNA methylation-based biological aging and cancer risk and survival:
872		Pooled analysis of seven prospective studies. *Int J Cancer* **142**, 1611-1619,
873		doi:10.1002/ijc.31189 (2018).
874	8	Lu, A. T. *et al.* DNA methylation GrimAge strongly predicts lifespan and healthspan.
875		*Aging* **11**, 303-327, doi:10.18632/aging.101684 (2019).
876	9	Ryan, J., Wrigglesworth, J., Loong, J., Fransquet, P. D. & Woods, R. L. A systematic review
877		and meta-analysis of environmental, lifestyle and health factors associated with DNA
878		methylation age. *The journals of gerontology. Series A, Biological sciences and medical*
879		*sciences*, doi:10.1093/gerona/glz099 (2019).
880	10	Hannum, G. *et al.* Genome-wide methylation profiles reveal quantitative views of
881		human aging rates. *Molecular cell* **49**, 359-367, doi:10.1016/j.molcel.2012.10.016
882		(2013).
883	11	Horvath, S. DNA methylation age of human tissues and cell types. *Genome biology* **14**,
884		R115, doi:10.1186/gb-2013-14-10-r115 (2013).
885	12	Bocklandt, S. *et al.* Epigenetic predictor of age. *PloS one* **6**, e14821,
886		doi:10.1371/journal.pone.0014821 (2011).
887	13	Zhang, Q. *et al.* Improved precision of epigenetic clock estimates across tissues and its
888		implication for biological ageing. *Genome Med* **11**, 54, doi:10.1186/s13073-019-0667-1
889		(2019).
890	14	Liu, Z. *et al.* Underlying features of epigenetic aging clocks in vivo and in vitro. *Aging cell*
891		**19**, e13229, doi:10.1111/acel.13229 (2020).
892	15	Lee, Y. *et al.* Blood-based epigenetic estimators of chronological age in human adults
893		using DNA methylation data from the Illumina MethylationEPIC array. *BMC genomics* **21**,
894		747, doi:10.1186/s12864-020-07168-8 (2020).
895	16	Horvath, S. *et al.* Epigenetic clock for skin and blood cells applied to Hutchinson Gilford
896		Progeria Syndrome and ex vivo studies. *Aging* **10**, 1758-1775,
897		doi:10.18632/aging.101508 (2018).
898	17	Levine, M. E. *et al.* An epigenetic biomarker of aging for lifespan and healthspan. *Aging*
899		**10**, 573-591, doi:10.18632/aging.101414 (2018).

18   Shireby, G. L. *et al.* Recalibrating the epigenetic clock: implications for assessing
biological age in the human cortex. *Brain* **143**, 3763-3775, doi:10.1093/brain/awaa334
(2020).
19   Petkovich, D. A. *et al.* Using DNA Methylation Profiling to Evaluate Biological Age and
Longevity Interventions. *Cell metabolism* **25**, 954-960 e956,
doi:10.1016/j.cmet.2017.03.016 (2017).
20   Stubbs, T. M. *et al.* Multi-tissue DNA methylation age predictor in mouse. *Genome*
biology **18**, 68, doi:10.1186/s13059-017-1203-5 (2017).
21   Wang, T. *et al.* Epigenetic aging signatures in mice livers are slowed by dwarfism, calorie
restriction and rapamycin treatment. *Genome biology* **18**, 57, doi:10.1186/s13059-017-
1186-2 (2017).
22   Levine, M. *et al.* A rat epigenetic clock recapitulates phenotypic aging and co-localizes
with heterochromatin. *Elife* **9**, doi:10.7554/eLife.59201 (2020).
23   Han, Y. *et al.* Epigenetic age-predictor for mice based on three CpG sites. *Elife* **7**,
doi:10.7554/eLife.37462 (2018).
24   Gibson, J. *et al.* A meta-analysis of genome-wide association studies of epigenetic age
acceleration. *PLoS genetics* **15**, e1008104, doi:10.1371/journal.pgen.1008104 (2019).
25   Lin, W. Y. Genome-wide association study for four measures of epigenetic age
acceleration and two epigenetic surrogate markers using DNA methylation data from
Taiwan biobank. *Human molecular genetics*, doi:10.1093/hmg/ddab369 (2021).
26   Lu, A. T. *et al.* GWAS of epigenetic aging rates in blood reveals a critical role for TERT.
Nature communications **9**, 387, doi:10.1038/s41467-017-02697-5 (2018).
27   McCartney, D. L. *et al.* Genome-wide association studies identify 137 genetic loci for
DNA methylation biomarkers of aging. *Genome biology* **22**, 194, doi:10.1186/s13059-
021-02398-9 (2021).
28   Kuo, C. L., Pilling, L. C., Liu, Z., Atkins, J. L. & Levine, M. E. Genetic associations for two
biological age measures point to distinct aging phenotypes. *Aging cell* **20**, e13376,
doi:10.1111/acel.13376 (2021).
29   Arneson, A. *et al.* A mammalian methylation array for profiling methylation levels at
conserved sequences. *Nature communications* **13**, 783, doi:10.1038/s41467-022-28355-
z (2022).
30   Lu, A. T. *et al.* Universal DNA methylation age across mammalian tissues. *bioRxiv*,
2021.2001.2018.426733, doi:10.1101/2021.01.18.426733 (2021).
31   Li, C. Z. *et al.* Epigenetic predictors of maximum lifespan and other life history traits in
mammals. *bioRxiv*, 2021.2005.2016.444078, doi:10.1101/2021.05.16.444078 (2021).
32   Williams, E. G. *et al.* Multiomic profiling of the liver across diets and age in a diverse
mouse population. *Cell Syst*, doi:10.1016/j.cels.2021.09.005 (2021).
33   Roy, S. *et al.* Gene-by-environment modulation of lifespan and weight gain in the murine
BXD family. *Nat Metab* **3**, 1217-1227, doi:10.1038/s42255-021-00449-w (2021).
34   Ashbrook, D. G. *et al.* A Platform for Experimental Precision Medicine: The Extended
BXD Mouse Family. *Cell Syst*, doi:10.1016/j.cels.2020.12.002 (2021).
35   Peirce, J. L., Lu, L., Gu, J., Silver, L. M. & Williams, R. W. A new set of BXD recombinant
inbred lines from advanced intercross populations in mice. *BMC genetics* **5**, 7,
doi:10.1186/1471-2156-5-7 (2004).

944 36 de Haan, G. & Williams, R. W. A genetic and genomic approach to identify longevity
945   genes in mice. *Mechanisms of ageing and development* **126**, 133-138,
946   doi:10.1016/j.mad.2004.09.012 (2005).
947 37 Hsu, H. C. *et al.* Age-related thymic involution in C57BL/6J x DBA/2J recombinant-inbred
948   mice maps to mouse chromosomes 9 and 10. *Genes and immunity* **4**, 402-410,
949   doi:10.1038/sj.gene.6363982 (2003).
950 38 Lang, D. H. *et al.* Quantitative trait loci (QTL) analysis of longevity in C57BL/6J by DBA/2J
951   (BXD) recombinant inbred mice. *Aging clinical and experimental research* **22**, 8-19
952   (2010).
953 39 Sandoval-Sierra, J. V. *et al.* Body weight and high-fat diet are associated with epigenetic
954   aging in female members of the BXD murine family. *Aging cell*, e13207,
955   doi:10.1111/acel.13207 (2020).
956 40 Benton, M. C. *et al.* Methylome-wide association study of whole blood DNA in the
957   Norfolk Island isolate identifies robust loci associated with age. *Aging* **9**, 753-768,
958   doi:10.18632/aging.101187 (2017).
959 41 Yashin, A. I. *et al.* Genetics of Human Longevity From Incomplete Data: New Findings
960   From the Long Life Family Study. *The journals of gerontology. Series A, Biological*
961   *sciences and medical sciences* **73**, 1472-1481, doi:10.1093/gerona/gly057 (2018).
962 42 Haghani, A. *et al.* Divergent age-related methylation patterns in long and short-lived
963   mammals. *bioRxiv*, 2022.2001.2016.476530, doi:10.1101/2022.01.16.476530 (2022).
964 43 Sziraki, A., Tyshkovskiy, A. & Gladyshev, V. N. Global remodeling of the mouse DNA
965   methylome during aging and in response to calorie restriction. *Aging cell* **17**, e12738,
966   doi:10.1111/acel.12738 (2018).
967 44 Slieker, R. C. *et al.* Age-related accrual of methylomic variability is linked to fundamental
968   ageing mechanisms. *Genome biology* **17**, 191, doi:10.1186/s13059-016-1053-6 (2016).
969 45 Kerepesi, C. *et al.* Epigenetic aging of the demographically non-aging naked mole-rat.
970   *Nature communications* **13**, 355, doi:10.1038/s41467-022-27959-9 (2022).
971 46 McLean, C. Y. *et al.* GREAT improves functional interpretation of cis-regulatory regions.
972   *Nature biotechnology* **28**, 495-501, doi:10.1038/nbt.1630 (2010).
973 47 Ernst, J. & Kellis, M. ChromHMM: automating chromatin-state discovery and
974   characterization. *Nat Methods* **9**, 215-216, doi:10.1038/nmeth.1906 (2012).
975 48 Gorkin, D. U. *et al.* An atlas of dynamic chromatin landscapes in mouse fetal
976   development. *Nature* **583**, 744-751, doi:10.1038/s41586-020-2093-3 (2020).
977 49 Zhou, X. & Stephens, M. Efficient multivariate linear mixed model algorithms for
978   genome-wide association studies. *Nature methods* **11**, 407-409,
979   doi:10.1038/nmeth.2848 (2014).
980 50 Peirce, J. L., Broman, K. W., Lu, L. & Williams, R. W. A simple method for combining
981   genetic mapping data from multiple crosses and experimental designs. *PloS one* **2**,
982   e1036, doi:10.1371/journal.pone.0001036 (2007).
983 51 Catalog, G. <https://www.ebi.ac.uk/gwas/> (
984 52 Horvath, S. *et al.* Obesity accelerates epigenetic aging of human liver. *Proceedings of the*
985   *National Academy of Sciences of the United States of America* **111**, 15538-15543,
986   doi:10.1073/pnas.1412759111 (2014).

53   Nevalainen, T. *et al.* Obesity accelerates epigenetic aging in middle-aged but not in
elderly individuals. *Clinical epigenetics* **9**, 20, doi:10.1186/s13148-016-0301-7 (2017).
54   Wang, M. & Lemos, B. Ribosomal DNA harbors an evolutionarily conserved clock of
biological aging. *Genome research* **29**, 325-333, doi:10.1101/gr.241745.118 (2019).
55   Xie, H. *et al.* Genome-wide quantitative assessment of variation in DNA methylation
patterns. *Nucleic acids research* **39**, 4099-4108, doi:10.1093/nar/gkr017 (2011).
56   Jenkinson, G., Pujadas, E., Goutsias, J. & Feinberg, A. P. Potential energy landscapes
identify the information-theoretic nature of the epigenome. *Nature genetics* **49**, 719-
729, doi:10.1038/ng.3811 (2017).
57   Hayflick, L. Entropy explains aging, genetic determinism explains longevity, and
undefined terminology explains misunderstanding both. *PLoS genetics* **3**, e220,
doi:10.1371/journal.pgen.0030220 (2007).
58   Donohoe, D. R. & Bultman, S. J. Metaboloepigenetics: interrelationships between energy
metabolism and epigenetic control of gene expression. *J Cell Physiol* **227**, 3169-3177,
doi:10.1002/jcp.24054 (2012).
59   Kucharski, R., Maleszka, J., Foret, S. & Maleszka, R. Nutritional control of reproductive
status in honeybees via DNA methylation. *Science* **319**, 1827-1830,
doi:10.1126/science.1153069 (2008).
60   Dolinoy, D. C., Huang, D. & Jirtle, R. L. Maternal nutrient supplementation counteracts
bisphenol A-induced DNA hypomethylation in early development. *Proceedings of the*
National Academy of Sciences of the United States of America **104**, 13056-13061,
doi:10.1073/pnas.0703739104 (2007).
61   Holman, G. D. A new deadly Syn? *Curr Biol* **9**, R735-737, doi:10.1016/s0960-
9822(99)80471-0 (1999).
62   Michailidou, K. *et al.* Association analysis identifies 65 new breast cancer risk loci.
Nature **551**, 92-94, doi:10.1038/nature24284 (2017).
63   Rokudai, S. *et al.* STXBP4 regulates APC/C-mediated p63 turnover and drives squamous
cell carcinogenesis. *Proceedings of the National Academy of Sciences of the United*
States of America **115**, E4806-E4814, doi:10.1073/pnas.1718546115 (2018).
64   Kichaev, G. *et al.* Leveraging Polygenic Functional Enrichment to Improve GWAS Power.
American journal of human genetics **104**, 65-75, doi:10.1016/j.ajhg.2018.11.008 (2019).
65   Perry, J. R. *et al.* Parent-of-origin-specific allelic associations among 106 genomic loci for
age at menarche. *Nature* **514**, 92-97, doi:10.1038/nature13545 (2014).
66   Albertsen, H. M. *et al.* A physical map and candidate genes in the BRCA1 region on
chromosome 17q12-21. *Nature genetics* **7**, 472-479, doi:10.1038/ng0894-472 (1994).
67   Kauraniemi, P. & Kallioniemi, A. Activation of multiple cancer-associated genes at the
ERBB2 amplicon in breast cancer. *Endocr Relat Cancer* **13**, 39-49,
doi:10.1677/erc.1.01147 (2006).
68   Tanaka, S. *et al.* Coexpression of Grb7 with epidermal growth factor receptor or
Her2/erbB2 in human advanced esophageal carcinoma. *Cancer research* **57**, 28-31
(1997).
69   de Vries, P. S. *et al.* Multiancestry Genome-Wide Association Study of Lipid Levels
Incorporating Gene-Alcohol Interactions. *Am J Epidemiol* **188**, 1033-1054,
doi:10.1093/aje/kwz005 (2019).

70 Richardson, T. G. *et al.* Evaluating the relationship between circulating lipoprotein lipids
and apolipoproteins with risk of coronary heart disease: A multivariable Mendelian
randomisation analysis. *PLoS Med* **17**, e1003062, doi:10.1371/journal.pmed.1003062
(2020).
71 Xu, Y. *et al.* IKK interacts with rictor and regulates mTORC2. *Cell Signal* **25**, 2239-2245,
doi:10.1016/j.cellsig.2013.07.008 (2013).
72 Gallagher, M. D. & Chen-Plotkin, A. S. The Post-GWAS Era: From Association to Function.
American journal of human genetics **102**, 717-730, doi:10.1016/j.ajhg.2018.04.002
(2018).
73 Lappalainen, T. Functional genomics bridges the gap between quantitative genetics and
molecular biology. *Genome research* **25**, 1427-1431, doi:10.1101/gr.190983.115 (2015).
74 Grodstein, F. *et al.* The association of epigenetic clocks in brain tissue with brain
pathologies and common aging phenotypes. *Neurobiol Dis* **157**, 105428,
doi:10.1016/j.nbd.2021.105428 (2021).
75 Olona, A. *et al.* Epoxygenase inactivation exacerbates diet and aging-associated
metabolic dysfunction resulting from impaired adipogenesis. *Mol Metab* **11**, 18-32,
doi:10.1016/j.molmet.2018.03.003 (2018).
76 Schuck, R. N. *et al.* The cytochrome P450 epoxygenase pathway regulates the hepatic
inflammatory response in fatty liver disease. *PloS one* **9**, e110162,
doi:10.1371/journal.pone.0110162 (2014).
77 Wang, Q. *et al.* Time serial transcriptome reveals Cyp2c29 as a key gene in
hepatocellular carcinoma development. *Cancer Biol Med* **17**, 401-417,
doi:10.20892/j.issn.2095-3941.2019.0335 (2020).
78 Mozhui, K. *et al.* Dissection of a QTL hotspot on mouse distal chromosome 1 that
modulates neurobehavioral phenotypes and gene expression. *PLoS genetics* **4**,
e1000260, doi:10.1371/journal.pgen.1000260 (2008).
79 Zhou, W., Triche, T. J., Jr., Laird, P. W. & Shen, H. SeSAMe: reducing artifactual detection
of DNA methylation by Infinium BeadChips in genomic deletions. *Nucleic acids research*
46, e123, doi:10.1093/nar/gky691 (2018).
80 Friedman, J., Hastie, T. & Tibshirani, R. Regularization Paths for Generalized Linear
Models via Coordinate Descent. *J Stat Softw* **33**, 1-22 (2010).
81 Coschigano, K. T. *et al.* Deletion, but not antagonism, of the mouse growth hormone
receptor results in severely decreased body weights, insulin, and insulin-like growth
factor I levels and increased life span. *Endocrinology* **144**, 3799-3810,
doi:10.1210/en.2003-0374 (2003).
82 Hausser, J. & Strimmer, K. Entropy Inference and the James-Stein Estimator, with
Application to Nonlinear Gene Association Networks. *J. Mach. Learn. Res.* **10**, 1469–
1484 (2009).
83 *GeneNetwork 2*, <http://genenetwork.org> (
84 Mulligan, M. K., Mozhui, K., Prins, P. & Williams, R. W. GeneNetwork: A Toolbox for
Systems Genetics. *Methods in molecular biology* **1488**, 75-120, doi:10.1007/978-1-4939-
6427-7_4 (2017).

85   McClay, J. L. *et al.* A methylome-wide study of aging using massively parallel sequencing
1074   of the methyl-CpG-enriched genomic fraction from blood in over 700 subjects. *Human*
1075   *molecular genetics* **23**, 1175-1185, doi:10.1093/hmg/ddt511 (2014).
1076   86   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic
1077   features. *Bioinformatics* **26**, 841-842, doi:10.1093/bioinformatics/btq033 (2010).
1078   87   Ashbrook, D. G. *et al.* Born to Cry: A Genetic Dissection of Infant Vocalization. *Front*
1079   *Behav Neurosci* **12**, 250, doi:10.3389/fnbeh.2018.00250 (2018).
1080   88   Belknap, J. K. Effect of within-strain sample size on QTL detection and mapping using
1081   recombinant inbred mouse strains. *Behavior genetics* **28**, 29-38 (1998).
1082   89   Sherry, S. T. *et al.* dbSNP: the NCBI database of genetic variation. *Nucleic acids research*
1083   **29**, 308-311, doi:10.1093/nar/29.1.308 (2001).
1084   90   Bates, D., Maechler, M., Bolker, B. & Walker, S. lme4: Linear mixed-effects models using
1085   Eigen and S4. . *R package version 1.1-7,* http://CRAN.R-project.org/package=lme4.
1086   91   *Wellcome Sanger Institute Mouse Genome Project*,
1087   <https://www.sanger.ac.uk/sanger/Mouse_SnpViewer/rel-1505> (
1088   92   Keane, T. M. *et al.* Mouse genomic variation and its effect on phenotypes and gene
1089   regulation. *Nature* **477**, 289-294, doi:10.1038/nature10413 (2011).
1090   93   Yalcin, B. *et al.* Sequence-based characterization of structural variation in the mouse
1091   genome. *Nature* **477**, 326-329, doi:10.1038/nature10432 (2011).
1092   94   Huang da, W., Sherman, B. T. & Lempicki, R. A. Bioinformatics enrichment tools: paths
1093   toward the comprehensive functional analysis of large gene lists. *Nucleic acids research*
1094   **37**, 1-13, doi:10.1093/nar/gkn923 (2009).
1095   95   Huang da, W., Sherman, B. T. & Lempicki, R. A. Systematic and integrative analysis of
1096   large gene lists using DAVID bioinformatics resources. *Nature protocols* **4**, 44-57,
1097   doi:10.1038/nprot.2008.211 (2009).
1098   96   Masuda, T., Tomita, M. & Ishihama, Y. Phase transfer surfactant-aided trypsin digestion
1099   for membrane proteome analysis. *J Proteome Res* **7**, 731-740, doi:10.1021/pr700658q
1100   (2008).
1101   97   Mostafa, D. *et al.* Loss of beta-cell identity and diabetic phenotype in mice caused by
1102   disruption of CNOT3-dependent mRNA deadenylation. *Commun Biol* **3**, 476,
1103   doi:10.1038/s42003-020-01201-y (2020).
1104   98   Reiter, L. *et al.* mProphet: automated data processing and statistical validation for large-
1105   scale SRM experiments. *Nature methods* **8**, 430-435, doi:10.1038/nmeth.1584 (2011).
1106   99   Rosenberger, G. *et al.* Statistical control of peptide and protein error rates in large-scale
1107   targeted data-independent acquisition analyses. *Nature methods* **14**, 921-927,
1108   doi:10.1038/nmeth.4398 (2017).
1109

a CD: r=0.59, p<.0001
HFD: r=0.76, p<.0001

b CD: r=0.58, p<.0001
HFD: r=0.76, p<.0001

c CD: r=0
HFD: r=0.31, p=0.0004

d UT319 (0.6 y)
UT573 (2.3 y)

Diet
○ CD
+ HFD

e p = 0.001

f CD: r=0.31, p<.0001
HFD: r=0.31, p=0.0004

g Age-gain entropy   Age-loss entropy

Variance source: Age, Diet, Chol, BWF, Gluc, LS

h p<0.0001
R^2=0.06

i p=0.0002
R^2=0.05

j p=0.0004
R^2=0.05

a int.EAA, pan

Genotypes @ Chr11 92.75 Mb
BB
DD
int.EAA, pan
CD HFD

Stxbp4, Erbb2, Grb7, Ikzf3

b int.EAA, liver

Genotypes @ Chr19 38.65 Mb
BB
DD
int.EAA, liver
CD HFD

Cyp2c29, Cyp2c37, Kif11, Cep55, Nkx2-3, Cutc, Chuk

c Methylome-wide entropy

Genotypes @ Chr19 47.51 Mb
BB
DD
Methylome-wide Entropy
CD HFD

Pcgf6, Elovl3

Chromosome
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 X

a Liver transcriptome

CLUSTER 3
Oxidation-reduction

CLUSTER 2
Cell cycle

Cyp2c29
Sod1
Hsd11b1
Ndufb5

Y 0

Kif11
Kif2c
Aspm

Tnfrsf1b

−300

Nfkb2   *Cyp2c39*
       Tnfrsf21

CLUSTER 1

−350   0   350
              X

d Liver proteome

CLUSTER 3
Oxidation-reduction,
steroid metabolism

APOC1   CYP1A1
                          CYP1A2
                          FMO1
APOE   FMO3   GAPDH
                          ENO1
                          ENO2
Y 0
CLUSTER 2
                    CLUSTER 4
                    Glycolysis/
                    gluconeogenesis

CYP2C39
                    CYP2C29
APOA1
    APOA4   CLUSTER 1
                  Oxidation-reduction,
−700   steroid metabolism

−250   0   250
              X

b
CD: r=0.34, p=0.0006
HFD: r=0.60, p<.0001

int.EAA, liver

−1
              CD+HDF
              r=0.56, p<.0001
       17   18
  APOE protein (Liver)

c
CD: r=0.27, p=0.006
HFD: r=0.50, p<.0001

EAA, pan

−1
              CD+HDF
              r=0.48, p<.0001
       17   18
  APOE protein (Liver)

e
CD: r=0.36, p=0.03
HFD: r=0.36, p=0.1

0.5

int.EAA, liver

−0.5
              CD+HDF
              r=0.44, p=.0005
  16   17   18
  APOE protein (Adipose)

f
CD: r=0.23, p=0.16
HFD: r=0.54, p=0.01

0.5

EAA, pan

−0.5
              CD+HDF
              r=0.36, p=0.005
  16   17   18
  APOE protein (Adipose)