1 Multiple mechanisms impact fluconazole resistance of mutant Erg11 proteins in

2 *Candida glabrata*

5 Bao Gia Vu and W. Scott Moye-Rowley[*]

7 Running title: Cellular effects of Erg11 mutants

From: Department of Molecular Physiology and Biophysics, Carver College of Medicine,

University of Iowa, Iowa City, IA 52242 USA.

[*]Corresponding author. E-mail: scott-moye-rowley@uiowa.edu

Abstract (249 words)

Azoles remain the most common used antifungal drugs for invasive candidiasis worldwide. They

specifically inhibit the fungal lanosterol α-14 demethylase enzyme, which is commonly referred

to as Erg11 in fungi. Inhibition of Erg11 ultimately leads to a reduction in ergosterol production,

an essential fungal membrane sterol. Many *Candida* species, such as *Candida albicans*,

develop mutations in this enzyme which reduces the azole binding affinity and results in

increased azole resistance. *Candida glabrata* is also a pathogenic yeast that has a low intrinsic

susceptibility to azole drugs and easily develops elevated resistance. These azole resistant

mutations are almost exclusively found to cause hyperactivity of the Pdr1 transcription factor

and rarely lie within the *ERG11* gene. Here, we generated *C. glabrata ERG11* mutations that

were analogous to azole resistance associated mutations in *C. albicans ERG11*. Three different

Erg11 forms (Y141H, S410F, and the corresponding double mutant (DM)) conferred azole

resistance in *C. glabrata* with the DM Erg11 form causing the strongest phenotype. The DM

Erg11 also induced cross-resistance to amphotericin B and caspofungin. The azole resistance

caused by the DM allele of *ERG11* imposed a fitness cost that was not observed with

hyperactive *PDR1* alleles. These data support the view that *C. glabrata* does not typically

acquire *ERG11* mutations owing to growth defects associated with these lesions while

hyperactive *PDR1* alleles have no obvious growth issues. Understanding the physiology linking

ergosterol biosynthesis with Pdr1-mediated regulation of azole resistance is crucial for ensuring

the continued efficacy of azole drugs against *C. glabrata*.

Importance (141 words)

Azole drugs target the Erg11 enzyme and lead to a reduction in fungal ergosterol, a vital sterol

in yeast. Mutations in Erg11 are common among azole resistant *Candida albicans* clinical

isolates, but not in *C. glabrata*, a major human pathogen. In this study, we showed that *ERG11*

mutations were tolerated in *C. glabrata*, and these mutations could confer azole resistance. We

also showed that *ERG11* mutations were less efficient in inducing azole resistance in

comparison to *PDR1* gain-of-function mutations, the most common mutation found among *C.*

glabrata azole resistant clinical isolates. Our data suggest that growth issues associated with

azole resistant alleles of *ERG11* in *C. glabrata* may limit the appearance of this class of

resistant strains. Furthermore, these data illustrate the deep connections between ergosterol

biosynthesis and regulation of membrane transporter proteins via Pdr1 and the ergosterol-

responsive transcription factor Upc2A.

Introduction

Ergosterol is essential for fungal plasma membrane homeostasis (1, 2). It maintains

membrane integrity and fluidity, as well as facilitates membrane-bound enzymatic reactions (1).

The azole class of antifungal therapy targets the fungal ability to synthesize ergosterol through

inhibition of the function of lanosterol 14-alpha-demethylase, which is encoded by the *ERG11*

gene (3, 4). The Erg11 enzyme catalyzes a three-step reaction that results in the demethylation

of lanosterol. In each step, one oxygen molecule is oxidized, leading to the production of one

NADPH molecule (4, 5). The nucleophilic N-4 atom of the azole ring binds to heme, while the N-

1 group interacts with the enzyme active site, directly competing with the normal substrate

(lanosterol) binding (6, 7). All azoles bind selectively to the Erg11 enzyme (8). However, the K_d

(dissociation constant) values show a 2-fold to 4-fold lower affinity to fluconazole than that to

itraconazole and voriconazole (4).

In *Candida albicans* (Ca), mutations in the Erg11 enzyme have been shown to alter

azole binding capacity and significantly reduce the drug inhibitory effect resulting in an

enhanced azole resistant phenotype (9-14). The majority of these mutations cluster into three

hot spots, located within residues 105 to 165, 266 to 287, and 405 to 488 (15). The ability to

confer azole resistance by some of these mutations has also been confirmed by introducing

these mutations on plasmids into a model *Saccharomyces cerevisiae* system and then

assessing azole sensitivity. With this approach, single amino acid mutations, such as: Y132H,

S405F, G464S, R467K, and double mutations, such as: Y132H with S405F, Y132H with

G464S, R467K with G464S have been shown to confer azole resistance in *S. cerevisiae* (9). In

addition, several *CaERG11* constructs of the enzyme catalytic domain containing amino acid

substitutions, including Y132H, F145L, I471T, S279F, and G464S have been expressed in

Escherichia coli, purified, and showed to exhibit a lower binding capacity to azoles (15). Finally,

many of these mutant alleles have also been introduced back into an azole-susceptible *C.*

albicans strain, at the *ERG11* gene native locus, to confirm the relative contributions of

individual and combined mutation to azole resistance (10).

Unlike in *C. albicans*, mutations in the *ERG11* gene have been rarely reported among

Candida glabrata (Cg) azole resistant clinical isolates. The most common amino acid

substitution mutations in *C. glabrata* are in the gene encoding a Zn_2Cys_6 zinc cluster-containing

transcription factor called Pdr1 (16). These mutations yield a gain-of-function (GOF) phenotype

and lead to the elevated transcription of downstream target genes, such as the ATP-binding

cassette (ABC) transporter-encoding *CDR1* gene. The Cdr1 transporter is required for the azole

resistant mechanism in both wild-type and GOF *PDR1 C. glabrata* isolates (17).

It has been recently reported that there is biological crosstalk between the ergosterol

biosynthesis pathway and the drug efflux pump system in *C. glabrata* (18)*.* Alteration in

ergosterol synthesis directly induced expression of both Pdr1 and Cdr1. The Zn_2Cys_6 zinc

cluster-containing factor Upc2A is essential for such cross-interaction, through its functions as a

master transcription factor regulating the expression levels of both ergosterol synthesis genes

(*ERG*) and *PDR1-CDR1* genes (19) (20) . Considering the conserved inhibitory effects of azoles

on Erg11 protein in both *C. albicans* and *C. glabrata*, we wanted to examine whether the

mutations equivalent to those in the *CaERG11* gene would confer azole resistance in *C.*

glabrata. To do that, we first generated isogenic *C. glabrata* isolates carrying mutant *ERG11*

alleles, analogs of the *CaERG11* Y132H, S405F or Y132H-S405F mutations. Then we

examined the effects of these mutant alleles on ergosterol synthesis, antifungal drug resistance,

and abilities to cross-regulate the drug efflux pump Pdr1-Cdr1 system. Finally, we compared the

azole induction capabilities between the mutations in *ERG11* and *PDR1* genes. Our data

suggest that growth defects caused by changes in *ERG11* that lead to azole resistance

approaching levels supported by GOF alleles of *PDR1* restrict the appearance of lesions in this

key ergosterol biosynthetic enzyme. The most azole resistant allele of ERG11 also triggers

constitutive induction of the Pdr1/CDR1 regulatory circuit, providing further evidence directly

tying expression of this drug resistance pathway to ergosterol biosynthesis.

Results

Generation of azole-resistant forms of *C. glabrata* Erg11. Detailed structural information is

available for the Erg11 enzymes from these two pathogenic *Candida* species as their three

dimensional structures have recently been solved by X-ray crystallography, along with the

previously determined structure of *S. cerevisiae* Erg11 (6). Figure 1A shows an alignment of the

amino acid sequences of these three closely related proteins.

Previous analyses of azole resistant clinical isolates of *C. albicans* identified changes in

residue tyrosine 132 to histidine (Y132H), serine 405 to phenylalanine (S405F) and a double

mutant containing both of these lesions (Y132H S405F, referred to as double mutant or DM

here) to trigger a decrease in fluconazole susceptibility when present in the *C. albicans* Erg11

enzyme (9). Inspection of the aligned sequences shown in Figure 1A indicated that both Y141

and S410 (in *C. glabrata*) were conserved residues across these three different proteins. To

determine if these same substitution mutations would cause similar fluconazole resistance

phenotypes when introduced into the *C. glabrata* Erg11, we used site directed mutagenesis to

generate the appropriate mutant enzymes. These were returned to *C. glabrata* as the sole

source of Erg11 using a plasmid shuffling approach (see Materials and Methods). Appropriate

transformants were tested for their resistance to fluconazole, caspofungin and amphotericin B

(Figure 1B).

The presence of either single mutant form of Erg11 (Y141H or S410F) led to an increase

in fluconazole resistance compared to the wild-type enzyme while the double mutant (Y141H

and S410F: DM) caused the strongest elevation in fluconazole resistance. Only the DM Erg11

was able to enhance resistance to caspofungin or amphotericin B. These data indicated that

these mutant forms of *C. glabrata* Erg11 were able to increase fluconazole resistance like their

C. albicans counterparts and, when combined in the DM Erg11, increased resistance to two

other antifungal drugs. We also tested these strains for their susceptibility to voriconazole and

itraconazole (Supplemental Figure 1) and found that the mutants caused voriconazole

phenotypes similar to those seen for fluconazole but had no significant effect on itraconazole

susceptibility.

The multidrug resistant nature of the DM *ERG11* mutant strain was not expected. This

finding prompted us to evaluate the levels of ergosterol supported by each of these mutant

strains as well as expression of genes that could explain the increased resistance to

caspofungin. The three mutant strains along with their wild-type *ERG11* counterpart were grown

to mid-log phase and allowed to grow for 3 or 24 hours with or without fluconazole challenge.

Total sterols were extracted from these cultures and assayed for the level of ergosterol

produced in each strain (Figure 2A, 2B and 2C). Total RNA was also prepared from isogenic

wild-type and DM *ERG11* strains without fluconazole challenge and assayed by RT-qPCR for

the mRNA levels of the genes indicated (Figure 2D).

Ergosterol levels were the lowest in the DM Erg11-expressing strains in the absence of

fluconazole stress (Figure 2A). The two single mutants produced ergosterol levels that were

very similar to those seen in the wild-type strain. A 3 hour treatment with fluconazole caused a

dramatic change in relative ergosterol levels as at this point, the DM Erg11 strain had the

highest level of ergosterol while the other 3 strains dropped below this level. No differences

were observed between the two mutant forms of Erg11 compared to the wild-type protein until

the fluconazole treatment was extended to 24 hours (Figure 2C). At this time, the DM Erg11 still

produced the highest level of ergosterol but now both the Y141H and S410F single mutant

strains had levels of ergosterol that were higher than the wild-type but still lower than the DM

Erg11.

Analysis of mRNA levels from genes directly involved in production of β-glucan synthase

(*FKS1*, *FKS2*, *FKS3*, *KNH1*) as well as regulators of these genes (*CRZ1*, *CNA1*) indicated that

only the mRNA for the *FKS2* gene was elevated in the DM *ERG11* mutant compared to its

cognate wild-type strain. This transcriptional induction of *FKS2* could serve to explain the

increase in caspofungin resistance seen in the DM Erg11-expressing strain.

Communication between *ERG11* and Pdr1/*CDR1* pathway. Our previous work

demonstrated that reduction in Erg11 expression led to an induction of the Pdr1 transcription

factor and its target genes such as *CDR1* (19) . To determine if these mutant forms of Erg11

triggered a similar response, we measured expression of Pdr1 and Cdr1 at both the protein and

mRNA levels. Transformants containing the various *ERG11* alleles were grown to mid-log

phase and either treated with fluconazole for 3 hours or left untreated. Total RNA was prepared

from these cultures and mRNA for *CDR1*, *PDR1* and *ERG11* measured by RT-qPCR (Figure

3A).

In the absence of fluconazole stress, only the DM *ERG11* allele led to induction of all

three of these mRNAs. Upon challenge with fluconazole, all of these genes were induced to

similar relative levels. The enhanced fluconazole resistance seen in the DM *ERG11* strain may

be explained by elevated *PDR1* and *CDR1* transcription, at least in part.

The levels of these proteins were also evaluated by western blots using appropriate

rabbit polyclonal antibodies for each *C. glabrata* protein (Figure 3B) and quantitated (Figure 3C).

Both Pdr1 and Cdr1 were elevated in untreated cells along with Erg11. Treatment with

fluconazole led to similar levels of induction of all these proteins. Upc2A protein levels were not

changed under any of these conditions or genetic backgrounds. This is consistent with Upc2A

regulation occurring at the post-translational step as has been suggested (21-23) .

The levels of mRNA of a range of genes in the ergosterol biosynthetic pathway were

also determined in both the DM and wild-type Erg11 strains (Figure 3D). These data are

presented as a ratio of mRNA level in the DM Erg11 strain over the wild-type level for each

transcript. Most of these *ERG* genes were induced at least 2-fold in the DM Erg11-expressing

cells, consistent with the ergosterol limitation in this strain although several *ERG* genes

including *ERG9*, *ERG7*, *ERG10* and *HMG1* showed little to no increase. Clearly, the presence

of the DM Erg11 enzyme induced a wide range of transcriptional activation across the

ergosterol pathway in addition to its effects on *PDR1* and *CDR1* expression.

Upc2A function is increased in the *ERG11* mutants. Our previous data showed that genetic

inhibition of *ERG11* gene expression could induce the expression of the drug efflux pump

system Pdr1-Cdr1 (19). Upc2A was essential for this interaction in which it directly induced the

expression levels of genes in both the *ERG* and Pdr1/*CDR1* pathways (19, 20). Since these two

pathways were also induced in the DM *ERG11* mutant, we wanted to determine if Upc2A

function was activated in response to these changes in the *ERG11* gene. To test this, we

generated an Upc2A reporter construct that contains five copies of the Upc2A binding site

(Sterol Response Element: SRE) from the *ERG1* promoter. These concatemerized SREs were

placed upstream from the *S. cerevisiae CYC1* promoter that was fused to *E. coli lacZ*. To

confirm that this reporter system was sensitive to the *UPC2A* allele, we introduced this plasmid

into isogenic *C. glabrata* strains containing the 3X HA-*UPC2A*, 3X HA-G898D *UPC2A* gain-of-

function mutation (20) or a null allele of *UPC2A*. A plasmid containing only the *CYC1-lacZ*

fusion gene with no upstream activation sequence was also used as negative control (Figure

4A).

The introduction of the gain-of-function form of Upc2A (G898D *UPC2A*) led to production

of the highest level of β-galactosidase seen from the 5X SRE-*CYC1-lacZ* plasmid with nearly

1200 Miller units of activity compared to approximately 800 Miller units when the wild-type

UPC2A gene was present. Loss of *UPC2A* lowered expression to approximately 80 Miller units.

All *C. glabrata* isolates carrying the *CYC1-lacZ* control plasmid lacking an upstream activation

sequence showed minimal levels of β-galactosidase activity. These data are consistent with

this reporter system faithfully detecting the status of the *UPC2A* gene.

Next, we transformed these two plasmids into strains containing the various *ERG11*

alleles. Appropriate transformants were grown to mid-log phase and enzyme levels determined

(Figure 4B). The highest level of β-galactosidase activity was produced in the presence of the

DM *ERG11* allele, consistent with this strain triggering the highest level of Upc2A activity. Both

single mutant *ERG11* alleles induced smaller but significant increases in expression of the 5X

SRE-*CYC1-lacZ* plasmid compared to the wild-type *ERG11*-containing strain. Only minimal

levels of β-galactosidase activity were produced in strains containing the *CYC1-lacZ* reporter

plasmid lacking any SREs as upstream control elements. These data support the view that the

DM *ERG11* allele caused the largest increase in Upc2A function, likely through production of the

most compromised form of the Erg11 enzyme, but each single mutant had a less dramatic but

still detectable impact on Upc2A. To directly assess Upc2A function, we carried out single gene

chromatin immunoprecipitation analysis of Upc2A binding to target promoters

Previous work from several labs including ours (20, 23) has provided evidence that

ergosterol limitation increased Upc2A DNA-binding to nuclear target genes. We also found that

these Upc2A target genes included the *PDR1* and *CDR1* genes (19), linking control of

ergosterol biosynthesis with expression of genes involved in drug resistance. To determine if

the presence of the fluconazole-resistant, double mutant allele of *ERG11* caused increased

levels of Upc2A DNA binding, chromatin immunoprecipitation was used to detect binding of this

factor to SREs present in the *ERG11, CDR1* and *PDR1* promoters. Non-specific binding was

also evaluated by examining Upc2A enrichment on the HO promoter and *ERG11* coding

sequence. Isogenic wild-type and DM *ERG11* strains were grown to mid-log phase and Upc2A

DNA-binding analyzed by ChIP using anti-Upc2A or a non-specific control polyclonal antiserum

(Figure 4C).

Upc2A binding to the *ERG11* SRE was strongly induced when cells contained the DM

ERG11 allele compared to the wild-type gene. Upc2A association with the *PDR1* SRE was also

induced under these conditions although binding to *CDR1* did not change. Control ChIP

reactions confirmed that the Upc2A binding was antibody-specific and showed no enrichment

with either the *HO* gene or the *ERG11* coding sequence. The increased Upc2A-dependent

transcriptional activation and DNA-binding indicate that Upc2A function was increased in strains

containing the DM *ERG11* allele compared to the wild-type gene.

Phenotypic comparison of *PDR1* GOF and DM *ERG11* mutations indicate mutant Pdr1

factors drive higher level azole resistance. Since the DM *ERG11* mutant was capable of

inducing strong fluconazole resistance, we wanted to compare its behavior to that seen in the

presence of *PDR1* GOF alleles. These GOF mutants of *PDR1* represent the vast majority of

azole resistant mutants found in *C. glabrata* isolates (24-26). An isogenic series of strains was

produced that produced wild-type or DM mutant forms of Erg11 containing the following alleles

of *PDR1*: *pdr1Δ*, wild-type, D1082G or R376W. These strains were grown to mid-log phase and

tested for their relative susceptibility to 30 μg/ml fluconazole in a serial dilution plate assay

(Figure 5A).

Both GOF forms of *PDR1* lowered fluconazole susceptibility more than the DM *ERG11*

strain. To probe the relative effects of mutations in either *PDR1* or *ERG11*, we carried out a

competitive growth assay in which equal numbers of cells containing mutations of interest were

mixed together. These mixed cultures were then allowed to grow in the presence or absence of

fluconazole challenge and the relative contribution of each mutant strain to the final population

determined. This competitive growth assay allows subtle differences to be more readily

detected than on a plate-based assay.

We prepared mixed cultures of isogenic wild-type and DM *ERG11* cells (Figure 5B) and

allowed these cultures to grow under untreated conditions (YPD) or in YPD medium containing

fluconazole to late log phase. Growth in rich medium led to an increase in the population of the

wild-type (60% final) compared to the DM *ERG11* strain (40% final). This analysis indicated that

the DM *ERG11* had a growth defect compared to wild-type cells in the absence of fluconazole

but this mutant allele conferred a significant growth advantage in the presence of this drug.

The relative growth of the DM *ERG11* strain was then compared to each of the *PDR1*

GOF mutant strains (Figures 5C and 5D). In both cases, the GOF *PDR1* mutant outcompeted

the DM *ERG11* strain in the presence of fluconazole (D1082G-74%: DM-26%; R376W-65%:

DM-35%) while the only difference in growth in the absence of drug was seen in comparison of

the D1082G *PDR1* strain compared to the DM *ERG11*: 58% to 42%, respectively. Both GOF

PDR1 mutants were more effective in conferring fluconazole resistance while growing at least

as well (R376W) if not better (D1082G) than the DM *ERG11* strain. These growth differences

may help explain the predominance of GOF mutants of *PDR1* in clinical azole-resistant isolates

of *C. glabrata*.

Increased activation of *PDR1* and *CDR1* transcription seen in presence of GOF *PDR1*

mutants compared to DM *ERG11*. The phenotypic data above indicated that both GOF *PDR1*

strains produced a stronger effect on fluconazole resistance than the presence of the DM

ERG11 allele. To probe the basis of this phenotypic difference, we examined transcription of

CDR1, *PDR1* and *ERG11* by RT-qPCR. Strains containing wild-type, D1082G or R376W forms

of *PDR1* or the DM *ERG11* were grown to mid-log phase in the absence or presence of

fluconazole. Levels of mRNA from these three genes were assayed as before.

CDR1 mRNA levels were strongly elevated in the presence of both GOF *PDR1*

mutations, well above those induced by the presence of the DM *ERG11* mutation (Figure 6A).

This increase was seen both in the absence as well as the presence of fluconazole. *PDR1*

mRNA levels were significantly higher in the presence of the two GOF alleles than in the DM

ERG11 but this difference was restricted to the absence of fluconazole. Drug treatment

elevated both wild-type and DM *ERG11* strains to levels similar to those of the GOF mutants.

Finally, *ERG11* mRNA levels were higher in the DM *ERG11* strain than in the other three strains

in the absence of fluconazole. Strikingly, fluconazole exposure induced *ERG11* mRNA to

higher levels in the strains containing wild-type *PDR1* but was less effective in the two GOF

PDR1-containing strains. These data suggest that the increased *PDR1/CDR1* expression seen

in the GOF *PDR1* strains may reduce the level of fluconazole stress (and associated *ERG11*

induction) seen. The changes seen in mRNA levels were well-correlated with western blot

analyses for each protein (Figures 6B and 6C). Upc2A protein levels were constant under all

conditions tested, consistent with changes due to this factor being caused by alterations in

Upc2A function rather than its expression.

Elevated levels of Upc2A exacerbate the growth defect of the DM *ERG11* strain. Our

analysis of the cellular response to the presence of the DM *ERG11* mutation demonstrated the

involvement of Upc2A as activity of this factor was induced in this genetic background

compared to wild-type cells. We introduced the highly repressible *MET3* promoter (27) in place

of the chromosomal *UPC2A* cognate region in order to test the effect of changing Upc2A

production via repression in the presence of exogenous methionine. This *MET3-UPC2A* fusion

gene was generated in both wild-type and DM *ERG11* strains. Appropriate transformants were

placed on media lacking (SC) or containing excess methionine (SC+methionine) and allowed to

grow. Two independent isolates of each *MET3-UPC2A* transformant were tested to evaluate

strain variability in these assays.

Addition of methionine to strains containing the *MET3-UPC2A* fusion gene in addition to

the DM *ERG11* allele caused the growth of this strain to be further reduced when compared to

the presence of the wild-type *UPC2A* gene (Figure 7A, right hand panel). This result suggests

that reduction of Upc2A levels (by repression of the *MET3* promoter with methionine) caused a

growth defect in the presence of the DM *ERG11* mutation. When these *UPC2A* and *MET3-*

UPC2A strains containing the DM *ERG11* mutation were grown in the absence of exogenous

methionine, their growth was similar (Figure 7A left hand panel).

To determine the levels of Upc2A protein in these different strains, we prepared whole

cell protein extracts from these strains. These extracts were analyzed by western blotting using

the anti-Upc2A antiserum. The *MET3* promoter activity is highest in SC medium (27) and we

will consider the data from these media conditions first (Figure 7B, left hand panel). The *MET3*

promoter drove much higher levels of Upc2A protein than were produced from the native

UPC2A promoter in the presence of wild-type *ERG11* (compare lanes 1 and 3). The elevated

expression of Upc2A seen in wild-type cells containing *MET3-UPC2A* did not lead to a

corresponding increase in Erg11 expression (lanes 1 and 3). Interestingly, Upc2A levels were

dramatically reduced when the *MET3-UPC2A* fusion gene was present in the DM *ERG11* strain,

albeit to levels similar to those seen when Upc2A was produced by the native *UPC2A* gene

(compare lanes 2 and 4). The presence of the *MET3-UPC2A* allele did not affect the observed

induction of Erg11 in the presence of the DM *ERG11* allele (compare lanes 2 and 4), potentially

due to the surprisingly low levels of Upc2A driven by the *MET3-UPC2A* allele when the DM

ERG11 mutant was present. The lack of any detectable Upc2A expression differences in the

two DM *ERG11* strains (varying due to their different *UPC2A* promoters) also correlated well

with the equivalent ability of these strains to grow on SC medium (Figure 7A above).

When methionine was added to repress *MET3-UPC2A* transcription, the levels of the

corresponding Upc2A protein were strongly reduced as expected (Figure 7B, right hand panel).

However, the reduction of Upc2A upon methionine repression led to different levels of this

transcription factor in a manner dependent on the *ERG11* allele present in the cell. Methionine

repression of the *MET3-UPC2A* in the DM *ERG11* strain blocked the production of a detectable

level of Upc2A while this same treatment of cells carrying wild-type *ERG11* still produced readily

detectable Upc2A (compare lanes 7 and 8). The extremely low level Upc2A production seen in

the methionine-repressed *MET3-UPC2A* with DM *ERG11* may explain the methionine-

dependent growth defect seen in this strain (see Figure 7A). These low levels of Upc2A also

blocked the normal increase in Erg11 caused by the DM *ERG11* allele (compare lanes 6 and 8).

We carried out this same analysis but using a *MET3-PDR1* gene (Supplemental Figure

2). There was no significant effect of the *MET3-PDR1* fusion gene on growth of either the wild-

type or DM *ERG11* strain irrespective of the presence of methionine (Supplemental Figure 2A).

However, steady-state levels of Pdr1 were much higher in the presence of the wild-type *ERG11*

gene than in the DM *ERG11*-containing strain (compare lanes 1 and 3). The expression of Pdr1

in these two strains was inversely correlated with expression of the Pdr1 target gene *CDR1* as

evidenced by the high Cdr1 levels in the DM *ERG11* strain compared to the wild-type strain

(compare lanes 1 and 3). This finding suggests that Pdr1 is activated in the presence of the DM

ERG11 allele. Additionally, the reduced accumulation of Pdr1 in the presence of DM Erg11

supported the view that this mutant enzyme led to Pdr1 activation as evidenced by the lower

steady-state levels of this transcription factor. Previously, we used the *MET3* promoter to

produce either wild-type or GOF Pdr1 proteins (28). The GOF proteins were found to be less

stable than the corresponding wild-type factor. Production of this hypomorphic form of Erg11, in

the absence of any azole drug, was sufficient to strongly induce Pdr1 activity and Cdr1

synthesis.

Discussion

Fluconazole resistant clinical isolates of *C. albicans* are readily recovered with lesions in

the Ca*ERG11* gene but azole resistant isolates with mutations in this locus have rarely been

reported in *C. glabrata* (29). We constructed *C. glabrata ERG11* mutations based on cognate

changes in the *C. albicans* gene. We selected positions Y141 and S410 in the *C. glabrata*

enzyme as the analogous positions in the *C. albicans* Erg11 could be found mutated either

alone or together in azole-resistant clinical isolates (9). The *C. glabrata* equivalent mutations

caused azole resistance as expected but the DM allele also exhibited a growth defect, even on

rich YPD medium, which was not reported for the *C. albicans* mutant. This growth defect, at

least for the DM *ERG11* mutant, may be part of the reason that mutations in *ERG11* are not

found in *C. glabrata* isolates.

This DM *ERG11* allele was also found to cause several other drug phenotypes. The

increased resistance to caspofungin was not expected and may involve induction of the *FKS2*

gene (Figure 2D). We have preliminary evidence that Upc2A may act to control expression of

β-glucan synthase (20) but have not yet linked this to *FKS2*. Crz1 is a well-established

transcriptional activator of *FKS* gene expression (30, 31) and may be involved in the crosstalk

between changes in ergosterol biosynthesis and β-glucan synthase levels.

The DM *ERG11* seems to be acting as a reduced function or hypomorphic allele of this

gene. This was evidenced by the reduced level of ergosterol accumulation in the DM mutant

background and the increased amphotericin B resistance. We suggest that this reduction in

function is critical for the observed induction of the *ERG* pathway genes as well as the

Pdr1/*CDR1* regulatory circuit. Previous work from our lab showed that genetically depleting

Erg11 by repression of two different artificial promoters driving *ERG11* caused the same

changes in gene regulation (19). The behavior of the DM *ERG11* mutant strain provides further

support for our hypothesis that a physiological link exists between the ergosterol biosynthetic

pathway and Pdr1 (18).

Comparison of the DM *ERG11* allele with each of the two single mutant alleles suggests

that there is a synthetic interaction between the Y141H and S410F mutations. In neither of the

two single mutants was evidence obtained that endogenous downstream gene expression was

being induced in the presence of these altered enzymes. However, when both of these lesions

were present in the same protein, a range of downstream responses were triggered. These

include *FKS2* induction and Pdr1 activation along with Upc2A. This was not observed with the

single mutants. We suggest that the single mutants seem to cause increased fluconazole

resistance primarily through their action on activity of the altered enzyme. The very sensitive 5X

SRE-*CYC1-lacZ* reporter plasmid was modestly induced in the presence of each single mutant

Erg11 (Figure 4B) but their effect was small compared to the induction caused by the DM Erg11

protein. The combination of the two single mutations led to production of a more defective

Erg11 protein that in turn caused intracellular responses such as activation of Upc2A and Pdr1.

The activation of Upc2A-dependent transcription is a central feature of the cellular

response to the presence of the DM *ERG11* allele. We produced a conditional *MET3-UPC2A*

strain that contained the DM *ERG11* mutation to avoid any problematic interactions caused by

introduction of a *upc2AΔ* null allele into this strain. The essential contribution of Upc2A to the

maintenance of the DM *ERG11*-containing strain could be shown by the severe growth defect

caused by methionine repression of this *UPC2A* allele (Figure 7A). We suggest that the

presence of the DM *ERG11* allele led to induction of Upc2A function which in turn is linked to an

increased level of degradation of this active transcription factor. This suggestion is supported

by the inhibition of *MET3-UPC2A* transcription by the addition of methionine leading to a drop in

Upc2A levels below the limit of detection in the DM *ERG11* background (Figure 7B). This is

roughly the equivalent of a direct protein stability assay as we block synthesis at the level of

transcription while allowing degradation to proceed without intervention. The apparent instability

of the wild-type Upc2A after activation by the presence of the DM Erg11 protein is very similar to

the instability found when GOF forms of Pdr1 are compared to wild-type Pdr1 (17) . These

factors are evidently deleterious for the cell to maintain in their activated states unless their

presence is crucial for a given stress response. We have previously observed this as a

hyperactive mutant form of Pdr1 can be lethal (17). These data suggest that an important

control point for Upc2A may lie at the level of protein stability.

One reason for constructing these different fluconazole resistant forms of Erg11 in *C.*

glabrata was to see if their presence would influence fluconazole induction of Pdr1. It is

possible that fluconazole resistant forms of Erg11 would no longer respond to challenge by this

azole drug. However, that was not the case as all strains expressing fluconazole resistant

ERG11 allele still showed induction of the Pdr1/*CDR1* regulatory axis upon fluconazole

exposure (Figure 3). The striking induction of *CDR1*, *PDR1* and *ERG11* expression when the

DM *ERG11* allele is present with no fluconazole addition argues that these genes and the

pathways they define are normally co-regulated, irrespective of the presence of drug.

The simplest interpretation for azole resistant *ERG11* mutations in multiple organisms

has been that the altered enzymes may no longer be able to interact with azole drugs and this

can explain their resistance phenotype. While this is certainly true for some mutants (such as

the two single mutant alleles of *ERG11* analyzed here), there are likely to be more complex

physiological responses to these mutant enzymes that can alter the resistance phenotype in

ERG11 mutant strains. Ergosterol is an essential constituent of the fungal cell membrane and

limiting its biosynthesis triggers transcriptional correction in most fungi examined (5, 32, 33). In

C. glabrata, there appears to be a threshold of ergosterol limitation that is not crossed by the

single mutant forms of Erg11 but is exceeded by the DM Erg11 enzyme. Exceeding this

threshold leads to activation of downstream pathways to deal with the detected ergosterol

limitation.

Our finding of these deeper regulatory connections between ergosterol biosynthesis and

transcription circuitry including but not restricted to factors directly involved in *ERG* gene control

suggests that additional linkages of this type may be participating in azole resistance in other

fungi. Azole-resistant forms of *Aspergillus fumigatus* are commonly associated with substitution

mutations within the *cyp51A* gene (*ERG11* equivalent in *Aspergilli*) (33). Some of the most

common azole-resistant alleles found in *A. fumigatus* are the result of linked changes in which a

residue in the coding sequence is changed and a region of the promoter is duplicated

(Reviewed in (34)). These linked changes act together to enhance azole resistance more than

either alone (35), suggesting that alleles of this sort engage more than one mechanism to affect

azole resistance. We have previously shown that the L98H form of Cyp51A in *A. fumigatus*

produces lower levels of this enzyme than the wild-type yet is still resistant to voriconazole (36).

Additionally, mutations in the HMGCoA reductase gene (*hmg1*) from *A. fumigatus* also causes a

strong decrease in voriconazole susceptibility although *cyp51A* gene expression appeared to be

unaffected (37). These observations suggest that other fungi may also employ additional

transcriptional circuitry to impact azole resistance in response to alterations in ergosterol

biosynthesis. Identifying and characterizing these other transcriptional contributors to azole

resistance are important goals in the dissection of the molecular basis of azole resistance.

Materials and Methods

Strains and growth conditions.

C. glabrata was grown in rich YPD medium (1% yeast extract, 2% peptone, 2% glucose) or

under amino acid-selective conditions in complete supplemental medium (CSM) (Difco yeast

nitrogen extract without amino acids, amino acid powder from Sunrise Science Products, 2%

glucose). YPD media supplemented with 50 μg/ml nourseothricin (Jena Bioscience, Jena,

Germany) and 2 mM methionine was used to select strains containing pBV65 vector derivatives

(38). CSM media (2% glucose, 1mM estradiol) without methionine was used to recycle the

selection cassette on pBV65. YPD solid agar supplemented with 1 mg/ml 5-fluoro-orotic acid

was used to cure the pBV43 plasmid. All strains used in this study are listed in Table 1.

Plasmid construction and promoter mutagenesis.

All constructs used for homologous recombination into the chromosome were constructed in the

pUC19 vector (New England Biolabs, Ipswich, MA). PCR was used to amplify DNA fragments.

PCR products were run on 1.5 % agarose gel (RPI, Troy, NY), excised and purified with

Purelink quick gel extraction and PCR cleaning combo kit (Invitrogen, Carlsbad, CA). Gibson

assembly cloning (New England Biolabs) was routinely used to assemble fragments together

into appropriate plasmid backbones. All isogenic deletion constructs were made by assembling

the recyclable cassette from pBV65 and fragments from the immediate upstream and

downstream regions of the target genes. Fragments of the target genes were amplified from

CBS138 background. Eviction of the recyclable cassette left a single copy of *loxP* in place of the

excised target gene coding region.

PDR1 constructs were made by using Gibson assembly (39) to produce fragments

corresponding to the wild-type, R376W or D1082G alleles of *PDR1* with a recyclable cassette

located downstream from the natural stop codon. Eviction of the recyclable cassette in the

integrated constructs left a single copy of *loxP* 250-300 base-pairs downstream of the target

gene stop codons. R376W and D1082G forms of *PDR1* were amplified from pSK70 and pSK71

backgrounds (17).

ERG11 wild-type and mutant integration constructs were made as above with the

resulting fragments corresponding to the wild-type, Y141H, S410F or the Y141H S410F alleles

of *ERG11* with the *his3MX6* (19) marker located 270 bp from the natural stop codon.

Conversion of strains to *LEU2* was done by PCR amplifying the *LEU2* coding region

along with 500 base pairs immediately upstream and downstream from the CBS138

background. Linear DNA was then transformed into KKY2001 and the colonies were selected

on CSM agar without leucine.

To generate the *S. cerevisiae ERG11* complementing plasmid (pBV43), the Sc*ERG11*

promoter (800 bps upstream of AUG), Sc*ERG11* coding sequence, and Sc*ERG11* terminator

(350 bps downstream of the stop codon) were PCR amplified from the an S288c wild-type strain

(SEY6210). This fragment was then cloned into the pCU-MET3 vector (27), which was pre-

digested with SacI and XhoI.

The Sterol Response Element (SRE) reporter construct (pBV382) was produced by

cloning the E. coli *lacZ* gene as a PCR fragment from pSK80 (17). This fragment was then

cloned into the pCL vector (27), which was pre-digested with SalI and XhoI. A Upc2A-

responsive promoter cassette, containing the 5 concatemerized copies of the *ERG1* promoter

SRE (-725 to -775) located downstream from the *ADH1* terminator and upstream of the *CYC1*

minimal promoter, was generated by Genscript (Piscataway, NJ). The Upc2A responsive

promoter was PCR amplified and cloned into the pCL-*lacZ* vector, digested with SacI and SalI.

The control vector (pBV378) was constructed by cloning the *ADH1* terminator and the *CYC1*

minimal promoter, PCR amplified from BVGC7 (19), into the pCL-*lacZ* vector, digested with SacI

and SalI.

To generate the *MET3*-driven *UPC2A* and *PDR1* strains, the *MET3* promoter and the

URA3 gene were PCR amplified from the pUC-*MET3* plasmid. This *URA3-MET3* promoter

cassette was placed immediately upstream of the target gene AUG codon by recombination into

the gene of interest.

C. glabrata transformation.

Cell transformations were performed using a lithium acetate method. After being heat shocked,

cells were either directly plated onto selective CSM agar plates (for auxotrophic

complementation) or grown at 30°C at 200 rpm overnight (for nourseothricin selection).

Overnight cultures were then plated on YPD agar plates supplemented with 50 µg/ml of

nourseothricin. Plates were incubated at 30°C for 24 to 48 hours. In case of chromosomal

insertion, individual colonies were isolated and screened by PCR for correct insertion of the

targeted construct.

Construction of mutant *ERG11* strains

Since Erg11 is essential for *C. glabrata* aerobic growth (40) ,A *URA3* (*ScURA3*)-containing

plasmid shuffling technique was used to construct *ERG11* mutant alleles in the SPG96 (*ura3Δ*)

C. glabrata strain (17). An autonomous plasmid pBV43 marked with *ScURA3* and containing the

wild-type Sc*ERG11* was first transformed into the SPG96 strain. Then *ERG11* was deleted in

the chromosome by homologous recombination, leaving Sc*ERG11* on pBV43 functioning as the

sole copy of the *ERG11* gene. Next, the wildtype *C. glabrata ERG11* and the 3 mutant alleles

were individually reintroduced to the normal locus by homologous recombination with selection

for His+ transformants. Finally, the pBV43 plasmid was cured with 1 mg/ml 5-FOA on a YPD

agar plate to construct the *C. glabrata* strains, BVGC344 (*ERG11*), BVGC346 (Y141H *ERG11*),

BVGC336 (S410F *ERG11*), and a strain containing an ERG11 gene with both of these

mutations present (BVGC340: DM *ERG11*).

Total sterol estimation.

Cell total sterol was extracted and measured as previously described (19). In short, cell pellets

were lysed in 25% alcoholic potassium hydroxide at 90°C for 2 hours. Total sterol was detected

by spectrophotometric scanning between the wavelengths of 240 nm and 300nm. The presence

of ergosterol in the extracted sample resulted in a four-peak curve with peaks located at

approximately 262, 270, 281, and 290 nm.

Quantification of transcript levels by RT-qPCR.

Total RNA was extracted from cells by extraction using TRIzol (Invitrogen) and chloroform

(Fisher Scientific, Hampton, NH) followed by purification with RNeasy minicolumns (Qiagen,

Redwood City, CA). 500 ng -1 µg total RNA was reverse-transcribed using an iScript cDNA

synthesis kit (Bio-Rad, Des Plaines, IL). Assay of RNA via quantitative PCR (qPCR) was

performed with iTaq universal SYBR green supermix (Bio-Rad). Target gene transcript levels

were normalized to transcript levels of 18S rRNA.

Spot test assay.

Cells were grown in YPD medium to mid-log-phase. Cultures were then 10-fold serially diluted

and spotted onto YPD agar plates containing different concentrations of fluconazole (LKT

laboratories, St Paul, MN), caspofungin (Apexbio, Houston, TX) or Amphotericin B (Sigma). All

agar plates were incubated at 30°C for 24 to 48 h before imaging was performed.

β-galactosidase assay.

Harvested cells were lysed with glass beads (Scientific Industries Inc) in breaking buffer (100

mM Tris pH8, 1 mM Dithiothreitol, and 20% Glycerol) at 4°C for 10 min. Lysate was collected

and β-galactosidase enzymatic reaction was carried out in Z-buffer (60 mM Na_2HPO_4, 40 mM

NaH_2PO_4, 10 mM KCl, 1 mM $MgSO_4$, 50 mM 2-Mercaptoethanol) with 650 µg/ml O-nitrophenyl-

β-D-galactopyranoside (ONPG). Miller units were calculated based on the equation: (OD_{420} x

1.7) / (0.0045 x total protein concentration x used extract volume x time). Bradford assay (Bio-

Rad) was used to measure the total protein concentration in the lysate.

Chromatin immunoprecipitation (ChIP).

The detailed ChIP protocol was previously described (19). The sheared chromatin was

incubated with rabbit polyclonal anti-Upc2A antibody (1:50 dilution) or rabbit IgG control

antibody, NBP2-24891 (Novus Biologicals, Centennial, CO) for 2 h before being incubated

together with 30 µl of washed protein G Dynabeads (Life Technologies) overnight on a nutator

at 4°C. Washing, reversal of cross-links, and purification of DNA processed by the use of ChIP

were performed as described in (19).

Real-time PCR was performed on ChIP purified DNA, under the following conditions: 1 cycle of

95°C for 30 s followed by 40 cycles of 95°C for 15 s and 57°C for 30 s on a MyiQ2 Bio-Rad

machine. A 1 µl volume of the DNA processed by the use of ChIP (diluted 5-fold) or of input

(diluted 20-fold) DNA was used in a reaction mixture with a 20 µl total volume using SYBR

green master mix (Bio-Rad) and a 0.4 µM concentration of each primer. The percent input

method was used to calculate the signal of enrichment of the promoter region for each gene.

ERG11, *CDR1*, and *PDR1* promoters were analyzed with primers specifically targeting the

ERG11 promoter (−561 to −694 relative to the ATG as +1), *CDR1* promoter (−476 to −665), and

PDR1 promoter (−551 to −651) regions. A region of *ERG11*, located within the coding sequence

(+939 to +1042) and the *HO* promoter (-585 to -751) region were used as negative controls.

Competitive growth assay.

The *LEU2* coding region along with its immediate 500 bps up- and downstream sequences were

amplified from CBS138 genomic DNA. The product was then used to transform the *leu2Δ*::FRT

allele to *LEU2* in DM *ERG11* background at the *LEU2* native locus to generate BVGC365 (DM

ERG11/*LEU2)*. A mid-log growth culture [between O.D. of 1 and 2] was diluted to 0.2 O.D. in

fresh YPD. Each culture to be tested: wild-type, R376W *PDR1* or D1082G *PDR1*, was mixed at

a 1:1 ratio with DM *ERG11*/*LEU2*, and the mixed cultures were treated either with fluconazole

(20 µg/ml) or ethanol for 24 hours at 30°C. Cultures were collected, serially diluted, and plated

on YPD and CSM media without leucine for colony forming unit analysis.

Antibodies and western immunoblotting.

Cells were lysed with lysis buffer (1.85 M NaOH, 7.5% 2-Mercaptoethanol). Proteins were

precipitated with 50% Trichloroacetic acid and resuspended in Urea buffer (40 mM Tris pH8, 8.0

M Urea, 5% SDS, 1% 2-Mercaptoethanol). Cdr1, Pdr1, and Upc2A rabbit polyclonal antibodies

were previously described (19). Anti-HA monoclonal antibody was purchased from Invitrogen.

Secondary antibodies were purchased from LI-COR Biosciences. Imaging was performed with

Odyssey CLx Imaging System (LI-COR Biosciences) and analyzed by Image Studio Lite

Software (LI-COR Biosciences). Detected target band fluorescence intensity was normalized

against tubulin fluorescence intensity and compiled from two biological replicate experiments

and two technical replicates in each experiment, giving four replicates in total. Erg11 anti-

peptide rabbit polyclonal antibody was produced by GenScript (Piscataway, NJ). The peptide

sequence was AKIYWEKRHPEQKY, and it covered the 520th to 533rd amino-acids in the

Erg11 protein primary sequence. The peptide antibody was confirmed by western blotting

comparing wild-type and a strain carrying an *ERG11*-3X HA allele. Anti-peptide antibody

detected native Erg11 from wild-type cells and the increased molecular mass of the Erg11-3X

HA protein. Dual color LI-COR secondary antibodies were used to overlap the band intensity of

Erg11 protein detected with anti-HA mouse monoclonal antibody and Erg11 peptide rabbit

polyclonal antibody

Statistics.

The Student T-test was used to assess the statistical significance of results of comparisons of

samples. Paired conditions were used for comparisons of results from the same isolate obtained

under different treatment conditions, while unpaired conditions were used for comparisons of

results from isolates obtained under the same treatment conditions (*, $P < 0.5$; **, $P < 0.01$; ***,

$P < 0.001$).

Acknowledgements

This work was supported by NIH AI152494 to WSM. We thank Dr. Damian Krysan for useful

suggestions and Drs. Tom Conway and Lucia Simonicova for critically reading this manuscript.

References

1.   Rodrigues ML. 2018. The Multifunctional Fungal Ergosterol. mBio 9.

2.   Sokolov SS, Trushina NI, Severin FF, Knorre DA. 2019. Ergosterol Turnover in Yeast: An
Interplay between Biosynthesis and Transport. Biochemistry (Mosc) 84:346-357.
3.   Zhang J, Li L, Lv Q, Yan L, Wang Y, Jiang Y. 2019. The Fungal CYP51s: Their Functions,
Structures, Related Drug Resistance, and Inhibitors. Front Microbiol 10:691.
4.   Warrilow AG, Martel CM, Parker JE, Melo N, Lamb DC, Nes WD, Kelly DE, Kelly SL. 2010.
Azole binding properties of Candida albicans sterol 14-alpha demethylase (CaCYP51).
Antimicrob Agents Chemother 54:4235-45.
5.   Jorda T, Puig S. 2020. Regulation of Ergosterol Biosynthesis in Saccharomyces cerevisiae.
Genes (Basel) 11.
6.   Keniya MV, Sabherwal M, Wilson RK, Woods MA, Sagatova AA, Tyndall JDA, Monk BC.
2018. Crystal Structures of Full-Length Lanosterol 14alpha-Demethylases of Prominent
Fungal Pathogens Candida albicans and Candida glabrata Provide Tools for Antifungal
Discovery. Antimicrob Agents Chemother 62.
7.   Sagatova AA, Keniya MV, Wilson RK, Monk BC, Tyndall JD. 2015. Structural Insights into
Binding of the Antifungal Drug Fluconazole to Saccharomyces cerevisiae Lanosterol
14alpha-Demethylase. Antimicrob Agents Chemother 59:4982-9.
8.   Warrilow AG, Parker JE, Kelly DE, Kelly SL. 2013. Azole affinity of sterol 14alpha-
demethylase (CYP51) enzymes from Candida albicans and Homo sapiens. Antimicrob
Agents Chemother 57:1352-60.
9.   Sanglard D, Ischer F, Koymans L, Bille J. 1998. Amino acid substitutions in the
cytochrome P-450 lanosterol 14alpha-demethylase (CYP51A1) from azole-resistant
Candida albicans clinical isolates contribute to resistance to azole antifungal agents.
Antimicrob Agents Chemother 42:241-53.
10. Flowers SA, Colon B, Whaley SG, Schuler MA, Rogers PD. 2015. Contribution of clinically
derived mutations in ERG11 to azole resistance in Candida albicans. Antimicrob Agents
Chemother 59:450-60.
11. Kelly SL, Lamb DC, Kelly DE. 1999. Y132H substitution in Candida albicans sterol 14alpha-
demethylase confers fluconazole resistance by preventing binding to haem. FEMS
Microbiol Lett 180:171-5.
12. Kelly SL, Lamb DC, Loeffler J, Einsele H, Kelly DE. 1999. The G464S amino acid
substitution in Candida albicans sterol 14alpha-demethylase causes fluconazole
resistance in the clinic through reduced affinity. Biochem Biophys Res Commun 262:174-
9.
13. Warrilow AG, Mullins JG, Hull CM, Parker JE, Lamb DC, Kelly DE, Kelly SL. 2012. S279
point mutations in Candida albicans Sterol 14-alpha demethylase (CYP51) reduce in vitro
inhibition by fluconazole. Antimicrob Agents Chemother 56:2099-107.
14. Marichal P, Koymans L, Willemsens S, Bellens D, Verhasselt P, Luyten W, Borgers M,
Ramaekers FCS, Odds FC, Vanden Bossche H. 1999. Contribution of mutations in the
cytochrome P450 14alpha-demethylase (Erg11p, Cyp51p) to azole resistance in Candida
albicans. Microbiology (Reading) 145 (Pt 10):2701-2713.
15. Warrilow AG, Nishimoto AT, Parker JE, Price CL, Flowers SA, Kelly DE, Rogers PD, Kelly
SL. 2019. The Evolution of Azole Resistance in Candida albicans Sterol 14alpha-
Demethylase (CYP51) through Incremental Amino Acid Substitutions. Antimicrob Agents
Chemother 63.

16. Moye-Rowley WS. 2019. Multiple interfaces control activity of the Candida glabrata
Pdr1 transcription factor mediating azole drug resistance. Curr Genet 65:103-108.
17. Khakhina S, Simonicova L, Moye-Rowley WS. 2018. Positive autoregulation and
repression of transactivation are key regulatory features of the Candida glabrata Pdr1
transcription factor. Mol Microbiol 107:747-764.
18. Moye-Rowley WS. 2020. Linkage between genes involved in azole resistance and
ergosterol biosynthesis. PLoS Pathog 16:e1008819.
19. Vu BG, Thomas GH, Moye-Rowley WS. 2019. Evidence that Ergosterol Biosynthesis
Modulates Activity of the Pdr1 Transcription Factor in Candida glabrata. MBio 10.
20. Vu BG, Stamnes MA, Li Y, Rogers PD, Moye-Rowley WS. 2021. Azole resistance is
mediated by integration of sterol gene regulation and membrane transporter
production by the zinc cluster-containing transcription factor Upc2A in Candida
glabrata. bioRxiv doi:10.1101/2021.05.07.443069:2021.05.07.443069.
21. Davies BS, Wang HS, Rine J. 2005. Dual activators of the sterol biosynthetic pathway of
Saccharomyces cerevisiae: similar activation/regulatory domains but different response
mechanisms. Mol Cell Biol 25:7375-85.
22. Marie C, Leyde S, White TC. 2008. Cytoplasmic localization of sterol transcription factors
Upc2p and Ecm22p in S. cerevisiae. Fungal Genet Biol 45:1430-8.
23. Yang H, Tong J, Lee CW, Ha S, Eom SH, Im YJ. 2015. Structural mechanism of ergosterol
regulation by fungal sterol transcription factor Upc2. Nat Commun 6:6129.
24. Vermitsky JP, Edlind TD. 2004. Azole resistance in Candida glabrata: coordinate
upregulation of multidrug transporters and evidence for a Pdr1-like transcription factor.
Antimicrob Agents Chemother 48:3773-81.
25. Tsai HF, Krol AA, Sarti KE, Bennett JE. 2006. Candida glabrata PDR1, a transcriptional
regulator of a pleiotropic drug resistance network, mediates azole resistance in clinical
isolates and petite mutants. Antimicrob Agents Chemother 50:1384-92.
26. Ferrari S, Ischer F, Calabrese D, Posteraro B, Sanguinetti M, Fadda G, Rohde B, Bauser C,
Bader O, Sanglard D. 2009. Gain of function mutations in CgPDR1 of Candida glabrata
not only mediate antifungal resistance but also enhance virulence. PLoS Pathog
5:e1000268.
27. Zordan RE, Ren Y, Pan SJ, Rotondo G, De Las Penas A, Iluore J, Cormack BP. 2013.
Expression plasmids for use in Candida glabrata. G3 (Bethesda) 3:1675-86.
28. Simonicova L, Moye-Rowley WS. 2020. Functional information from clinically-derived
drug resistant forms of the Candida glabrata Pdr1 transcription factor. PLoS Genet
16:e1009005.
29. Hull CM, Parker JE, Bader O, Weig M, Gross U, Warrilow AG, Kelly DE, Kelly SL. 2012.
Facultative sterol uptake in an ergosterol-deficient clinical isolate of Candida glabrata
harboring a missense mutation in ERG11 and exhibiting cross-resistance to azoles and
amphotericin B. Antimicrob Agents Chemother 56:4223-32.
30. Matheos DP, Kingsbury TJ, Ahsan US, Cunningham KW. 1997. Tcn1p/Crz1p, a
calcineurin-dependent transcription factor that differentially regulates gene expression
in Saccharomyces cerevisiae. Genes Dev 11:3445-58.
31. Stathopoulos AM, Cyert MS. 1997. Calcineurin acts through the CRZ1/TCN1-encoded
transcription factor to regulate gene expression in yeast. Genes Dev 11:3432-44.

32.   Lv QZ, Yan L, Jiang YY. 2016. The synthesis, regulation, and functions of sterols in
707   Candida albicans: Well-known but still lots to learn. Virulence 7:649-59.
708   33.   Dhingra S, Cramer RA. 2017. Regulation of Sterol Biosynthesis in the Human Fungal
709   Pathogen Aspergillus fumigatus: Opportunities for Therapeutic Development. Front
710   Microbiol 8:92.
711   34.   Chowdhary A, Kathuria S, Xu J, Meis JF. 2013. Emergence of azole-resistant aspergillus
712   fumigatus strains due to agricultural azole use creates an increasing threat to human
713   health. PLoS Pathog 9:e1003633.
714   35.   Mellado E, Garcia-Effron G, Alcazar-Fuoli L, Melchers WJ, Verweij PE, Cuenca-Estrella M,
715   Rodriguez-Tudela JL. 2007. A new Aspergillus fumigatus resistance mechanism
716   conferring in vitro cross-resistance to azole antifungals involves a combination of
717   cyp51A alterations. Antimicrob Agents Chemother 51:1897-904.
718   36.   Paul S, Diekema D, Moye-Rowley WS. 2017. Contributions of both ATP-Binding Cassette
719   Transporter and Cyp51A Proteins Are Essential for Azole Resistance in Aspergillus
720   fumigatus. Antimicrob Agents Chemother 61.
721   37.   Rybak JM, Ge W, Wiederhold NP, Parker JE, Kelly SL, Rogers PD, Fortwendel JR. 2019.
722   Mutations in hmg1, Challenging the Paradigm of Clinical Triazole Resistance in
723   Aspergillus fumigatus. mBio 10.
724   38.   Vu BG, Moye-Rowley WS. 2018. Construction and Use of a Recyclable Marker To
725   Examine the Role of Major Facilitator Superfamily Protein Members in Candida glabrata
726   Drug Resistance Phenotypes. mSphere 3.
727   39.   Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, 3rd, Smith HO. 2009.
728   Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat Methods
729   6:343-5.
730   40.   Geber A, Hitchcock CA, Swartz JE, Pullen FS, Marsden KE, Kwon-Chung KJ, Bennett JE.
731   1995. Deletion of the Candida glabrata ERG3 and ERG11 genes: effect on cell viability,
732   cell growth, sterol composition, and antifungal susceptibility. Antimicrob Agents
733   Chemother 39:2708-17.

735   Figure legends

Figure 1. **Construction of mutant *ERG11* alleles in *Candida glabrata*.** A. Amino acid

sequence comparison between Erg11 proteins from *Saccharomyces cerevisiae*, *Candida*

glabrata and *Candida albicans*. Graphic indication of amino acid identity between these three

yeast enzymes is shown with a red bar indicating 100%, while light and dark green denotes

66% or no conservation. Expanded regions with conserved residues are shown to illustrate the

positions that were mutated in *C. glabrata* to match changes associated with fluconazole

resistance originally found in *C. albicans*. B. Isogenic *C. glabrata* strains containing the

indicated *ERG11* alleles were grown to mid-log phase and plated as serial dilutions on rich

medium (YPD) or the same medium containing the indicated concentrations of antifungal drugs.

The strain containing both the Y141H- and S410F-encoding alleles was designated double

mutant (DM). Plates were allowed to develop at 30°C and then photographed. Antifungal drugs

are abbreviated as fluconazole (FLC), caspofungin (CSF) and amphotericin B (AmB).

Figure 2. **Ergosterol levels and caspofungin resistance-related mRNA production in**

***ERG11* mutant strains.** Mid-log phase *C. glabrata* strains containing the indicated *ERG11*

alleles were either not treated (A) with fluconazole (-FLC) or exposed to this drug for 3 (B) or 24

(C) hours. Total lipid extracts were prepared and analyzed for the levels of ergosterol in each

sample. D. Total RNA was extracted from isogenic wild-type and DM *ERG11* strains without

fluconazole treatment. Samples were then assayed for levels of the indicated mRNAs using

RT-qPCR. Relative expression level refers to the ratio of mRNA produced in DM *ERG11*

strains/wild-type strain. Baseline indicates equivalent expression in the wildtype strain.

Figure 3. **Expression changes in response to *ERG11* alleles**. A. Levels of mRNA for *PDR1,*

CDR1, and *ERG11* were assayed by RT-qPCR in the indicated *ERG11* mutant strains during

growth in the absence (Control) or presence of fluconazole stress (Fluconazole). B. Whole cell

protein extracts were prepared from the same strains in A. These extracts were analyzed by

western blotting using the indicated antisera against Cdr1, Pdr1, Erg11, Upc2A or Tubulin.

Equal loading and transfer of these extracts was ensured by staining the nitrocellulose

membrane with Ponceau S prior to antisera incubation. C. Quantitation of 4 replicates of this

western blotting is shown with the data presented as the ratio of each strain relative to the wild-

type cells. D. RT-qPCR analysis of *ERG* gene expression in the DM *ERG11* mutant

background relative to the wild-type level. HMGCoA (HMG-CoA reductase) refers to the *HMG1*

or *CAGL0L11506g* locus.

Figure 4. **Regulation of Upc2A in response to *ERG11* alleles**. A. Two different reporter

constructs carried on low-copy-number *C. glabrata* vectors that contained either the *S.*

cerevisiae CYC1 promoter region with a translational fusion to *E. coli lacZ* (*CYC1-lacZ*) or the

same reporter construct with five copies of the *C. glabrata ERG1* sterol response element (SRE)

cloned upstream of the *CYC1* promoter (5X SRE-*CYC1-lacZ*) were introduced into the 3

different *C. glabrata* strains indicated at the top. These strains varied at their *UPC2A* allele

corresponding to a wild-type strain (*UPC2A*), gain-of-function form of *UPC2A* (G898D *UPC2A*)

or a null mutation (*upc2AΔ*). Transformants were grown to mid-log phase and then assayed for

the level of β-galactosidase produced in each strain. B. The two different reporter plasmids

from A were introduced into the 4 isogenic *ERG11* mutant strains indicated. Transformants

were grown to mid-log phase and assayed for β-galactosidase levels. C. Chromatin

immunoprecipitation (ChIP) of Upc2A DNA-binding to genomic target sites. Total sheared and

crosslinked chromatin was prepared from either wild-type or DM *ERG11* cells. ChIP reactions

were carried out with either anti-Upc2A or a non-specific control antisera. Immunopurified DNA

was quantitated with qPCR using primer pairs that detected the promoters of *ERG11* (*ERG11*),

CDR1 (*CDR1*), *PDR1* (*PDR1*) or *HO* (*HO*) along with a primer pair that detected a segment of

the *ERG11* coding sequence (*ERG11* cds). Data are plotted as the percentage of DNA

recovered in the immunopurified sample/total input DNA.

Figure 5. **Growth phenotypic comparison of DM *ERG11* with gain-of-function (GOF)**

alleles of *PDR1*. A. Isogenic mid-log phase *C. glabrata* strains containing *ERG11* wild-type or

DM alleles, *PDR1* null (*pdr1Δ*), wild-type *PDR1* (*PDR1*::loxP), or two GOF alleles of *PDR1*

D1082G and R376W were serially diluted and spotted on YPD agar plates supplemented with

or without fluconazole. Plates were incubated at 30°C and photographed. B. Fitness

comparisons between the DM *ERG11* strain compared with wild-type (B), the GOF D1082G (C)

and R376W (D) *PDR1* mutant strains. Equal number of mid-log phase cells from each strain

were mixed together (Pretreatment). Cultures were then allowed to grow in the absence (YPD)

or presence of azole drug (Fluconazole). At the end of incubation, samples of each culture

were plated and final distribution of each strain determined by plating. Deviations from the

starting 50:50 mix of strains would indicate a relative fitness advantage for one strain versus the

other.

Figure 6. **Relative gene expression responses to the DM *ERG11* or two different GOF**

***PDR1* strains**. A. Strains varying at the *ERG11* or *PDR1* loci as indicated were grown in the

presence or absence of fluconazole as in Figure 3 and processed for RT-qPCR analyses of the

indicated mRNAs. B. The strains above were analyzed by western blotting as in Figure 3 using

antisera for Cdr1, Pdr1, Erg11, Upc2A or Tubulin (Tub). The membrane was stained after

transfer with Ponceau S as above. Quantitation of 4 western experiments is provided on the

right hand side of the figure.

Figure 7. **Depletion of Upc2A enhances growth defect of a DM *ERG11* strain**. A. The

ERG11 promoter was replaced with the methionine-repressible *MET3* promoter in wild-type and

DM *ERG11* cells. Two different isolates of each *MET3* promoter replacement were grown to

mid-log phase, along with the isogenic wild-type and DM *ERG11* strains. These cells were

serially diluted onto plates containing either synthetic complete (SC) medium or the same

medium containing an excess of methionine (SC + methionine). Transformants were grown at

30°C and the plates photographed. B. The indicated strains were grown to mid-log phase in

the presence or absence of methionine and whole cell protein extracts prepared. These

extracts were analyzed as before using the indicated antisera.

Supplemental Figure Legends

Supplemental Figure 1. **Voriconazole and itraconazole resistance among *ERG11***

mutations. Mid-log phase *C. glabrata* strains containing the indicated *ERG11* alleles were

serially diluted and spotted on YPD agar plates supplemented with various concentrations of

voriconazole (A) or itraconazole (B).

Supplemental Figure 2. **Interaction of DM *ERG11* allele and Pdr1.** A. The *PDR1* promoter

was replaced with the *MET3* promoter in strains containing either the wild-type or DM *ERG11*

allele. Strains were grown to mid-log phase and then serial dilutions plated on SC medium

lacking or containing methionine. B. The strains above were grown to mid-log phase in SC

medium lacking methionine, methionine was either omitted (-) or added (+) and cultures allowed

to continue to grow for equal times. Whole cell protein extracts were then prepared and

analyzed by western blotting using the indicated antibodies.

Table

Strain name	Parent	Genotype
SPG96	KKY2001	*his3Δ::FRT leu2Δ::FRT trp1Δ::FRT ura3Δ(-85+932)::Tn903NeoR*
BVGC14	SPG96	*his3Δ::FRT trp1Δ::FRT CEN-ScERG11/URA3 erg11Δ::LEU2*
BVGC344	BVGC14	*leu2Δ::FRT trp1Δ::FRT ura3Δ(-85+932)::Tn903NeoR ERG11::HIS3*
BVGC346	BVGC14	*leu2Δ::FRT trp1Δ::FRT ura3Δ(-85+932)::Tn903NeoR* Y141H *ERG11::HIS3*
BVGC336	BVGC14	*leu2Δ::FRT trp1Δ::FRT ura3Δ(-85+932)::Tn903NeoR* S410F *ERG11::HIS3*
BVGC340	BVGC14	*leu2Δ::FRT trp1Δ::FRT ura3Δ(-85+932)::Tn903NeoR* Y141H S410F *ERG11::HIS3*
BVGC367	BVGC344	*leu2Δ::FRT trp1Δ::FRT ura3Δ(-85+932)::Tn903NeoR ERG11::HIS3 pdr1Δ::loxP*

BVGC383	BVGC367	*leu2Δ::FRT trp1Δ::FRT ura3Δ(-85+932)::Tn903NeoR ERG11::HIS3 PDR1::loxP*
BVGC387	BVGC367	*leu2Δ::FRT trp1Δ::FRT ura3Δ(-85+932)::Tn903NeoR ERG11::HIS3* D1082G *PDR1::loxP*
BVGC391	BVGC367	*leu2Δ::FRT trp1Δ::FRT ura3Δ(-85+932)::Tn903NeoR ERG11::HIS3* R376W *PDR1::loxP*
BVGC361	BVGC340	*leu2Δ::FRT trp1Δ::FRT ura3Δ(-85+932)::Tn903NeoR* Y141H S410F *ERG11::HIS3 PDR1::loxP*
BVGC365	BVGC361	*leu2Δ::LEU2 trp1Δ::FRT ura3Δ(-85+932)::Tn903NeoR* Y141H S410F *ERG11::HIS3 PDR1-LOXP*
BVGC374	BVGC344	*leu2Δ::FRT trp1Δ::FRT ura3Δ(-85+932)::Tn903NeoR ERG11::HIS3 URA3::MET3-PDR1*
BVGC378	BVGC344	*leu2Δ::FRT trp1Δ::FRT ura3Δ(-85+932)::Tn903NeoR*

		ERG11::HIS3 URA3::MET3-UPC2A
BVGC395	BVGC340	leu2Δ::FRT trp1Δ::FRT ura3Δ(-85+932)::Tn903NeoR Y141H S410F ERG11::HIS3 URA3::MET3-PDR1
BVGC404	BVGC340	leu2Δ::FRT trp1Δ::FRT ura3Δ(-85+932)::Tn903NeoR Y141H S410F ERG11::HIS3 URA3::MET3-UPC2A

845   Table 1. Strains used in this work.

A.

-FLC

B.

+FLC, 3 hr

C.

+FLC, 24 hr

D.

Figure 6. Vu and Moye-Rowley

A.

SC   SC + methionine

B.

A.

| | YPD | Voriconazole 1 μg/ml | Voriconazole 1.5 μg/ml |

ERG11 wt, Y141H, S410F, DM

B.

Itraconazole 0.2 μg/ml   Itraconazole 1 μg/ml

ERG11 wt, Y141H, S410F, DM

A.

MET3-PDR1

ERG11 Isolate #1
Isolate #2

DM *ERG11* Isolate #1
Isolate #2

SC   SC + methionine

B.

MET3-PDR1

ERG11   DM *ERG11*

Methionine   −   +   −   +

Pdr1

Cdr1

Erg11

Tubulin

Ponceau S

1   2   3   4