# A single-cell atlas of *de novo* β-cell regeneration reveals the contribution of hybrid β/δ cells to diabetes recovery in zebrafish

**Authors:** Sumeet Pal Singh [1*], Prateek Chawla [2*], Alisa Hnatiuk [2**], Margrit Kamel [2**], Luis Delgadillo Silva [2**], Bastiaan Spanjard [3], Sema Elif Eski[1], Sharan Janjuha [4], Pedro Olivares [3], Oezge Kayisoglu [5], Fabian Rost [6], Juliane Bläsche [6], Annekathrin Kränkel [6], Andreas Petzold [6], Thomas Kurth [7], Susanne Reinhardt [6], Jan Philipp Junker [3], and Nikolay Ninov [2,8] **

1. IRIBHM, Université Libre de Bruxelles (ULB), Brussels, Belgium.
2. Center for Regenerative Therapies Dresden, Technische Universität Dresden, Dresden, Germany
3. Berlin Institute for Medical Systems Biology, Max Delbrück Center for Molecular Medicine, Berlin, Germany.
4. Institute of Pharmacology and Toxicology, University of Zurich, Zurich, Switzerland
5. The Julius Maximilian University of Wurzburg, Wurzburg, Germany

6. DRESDEN-concept Genome Center, DFG NGS Competence Center, c/o Center for Molecular and Cellular Bioengineering (CMCB), Technische Universität Dresden, Dresden, Germany
7. TUD, Center for Molecular and Cellular Bioengineering (CMCB), Technology Platform, EM-Facility

8. Paul Langerhans Institute Dresden of the Helmholtz Zentrum München at the University Hospital and Faculty of Medicine Carl Gustav Carus of Technische Universität Dresden, Dresden, Germany

\* Equal contribution as first authors
\** Corresponding author

**Abstract:**

Regeneration-competent species possess the ability to reverse the progression of severe diseases by restoring the function of the damaged tissue. However, the cellular dynamics underlying this capability remain unexplored. Here, we use single-cell transcriptomics to map the cellular dynamics underlying *de novo* β-cell regeneration during induction and recovery from diabetes in zebrafish. We show that the zebrafish has evolved two distinct types of somatostatin-producing δ-cells, which we term δ1- and δ2-cells. Moreover, we characterize a small population of glucose-responsive islet cells, which share the hormones and fate-determinants of both β- and δ1-cells. The transcriptomic analysis of β-cell regeneration reveals that the development of β/δ hybrid cells constitutes a prominent source of insulin-expression during diabetes recovery. Using *in vivo* calcium imaging and cell tracking, we further show that the hybrid cells form *de novo* upon β-cell loss and acquire glucose-responsiveness in the course of regeneration. The overexpression of *dkk3*, a gene enriched in hybrid cells, increases their formation in the absence of β-cell injury. Our data provide an atlas of β-cell regeneration and indicate that the rapid formation of glucose-responsive hybrid cells contributes to the resolution of diabetes in zebrafish.

## Introduction

The pancreas, a glandular organ, is composed of exocrine and endocrine compartments. The exocrine part of the pancreas is involved in food digestion via the function of acinar and ductal cells. The endocrine part is responsible for metabolic homeostasis, specifically regulation of blood glucose levels. The endocrine compartment is organized into the islets of Langerhans, which consist of three major cell types: α-, β- and δ-cells. The three cell-types produce and secrete specific hormones: α-cells generate glucagon (*gcg*), β-cells generate insulin (*ins*), and δ-cells generate somatostatin (*sst*). Blood glucose homeostasis is achieved by regulated secretion of endocrine hormones in the blood stream; insulin activates glucose uptake in peripheral tissue, while glucagon opposes insulin action by stimulating the production of sugar in the liver (Noguchi and Huising, 2019). Somatostatin release has an inhibitory role for both insulin and glucagon secretion, as its loss triggers acute hypoglycemia (Li et al., 2018). As the selective autoimmune destruction of β-cells causes Type 1 Diabetes, enhancing the limited regenerative potential of the pancreas has become an important quest towards developing new regenerative therapies.

Regeneration and cell plasticity are intimately connected. Upon near-complete β-cell destruction, a small proportion of glucagon-expressing α-cells into β-cells in adult mice and embryonic zebrafish (Chera et al., 2014; Thorel et al., 2010). In addition, a subset of terminal duct cells, called centroacinar cells can give raise to β-cells in adult fish and mice (Delaspre et al., 2015; Mameishvili et al., 2019). Moreover, in juvenile mice, δ-cells initiate insulin production upon ablation of the β-cells, however this cellular makeover ceases to operate in older animals. To define the full spectrum of cellular dynamics underlying β-cell regeneration requires

technologies that can track gene expression in multiple cells over time during the process of β-cells regeneration.

In this study, we applied a unique blend of *in vivo* imaging, single-cell genomics, and genetics to study *de novo* β-cell regeneration in zebrafish. While regeneration is incomplete in mammals, zebrafish can naturally recover from extreme β-cell destruction and hyperglycemia (Moss et al., 2009). However, the reasons underlying this ability remain unexplored. By categorizing the cells in the zebrafish pancreas, we identify two discrete populations of δ-cells, which express different *sst* paralogues. Moreover, we find that a subset of islet cells exist under a hybrid identity, sharing the hormones and fate-determinants of both δ- and β-cells. During ablation and subsequent regeneration, the hybrid cells form *de novo*, acquire glucose-responsiveness and serve as a major source of insulin-expression. We also show that the secreted protein Dkk3b can increase the formation of hybrid cells in the absence of injury. We propose that the rapid formation of glucose-responsive hybrid cells provides a shortcut to restoring glucose homeostasis and contributes to the ability of zebrafish to reverse the course of diabetes.

## Results

### Single-cell RNA-sequencing of the pancreas identifies two populations of δ-cells in zebrafish

We first characterized the cell types in the zebrafish pancreas under homeostasis. We analyzed data from a droplet based single-cell transcriptome profiling of the islets and surrounding exocrine tissues from 2 months post-fertilization (mpf) zebrafish (Salem et al., 2019). For this experiment, the tissue surrounding the principal islet was dissected from the pancreas (n = 6 animals). The

dissected tissue was enzymatically dissociated, sorted by flow cytometry to remove damaged cells and profiled using the 10x Genomics pipeline (Fig. 1A). The transcriptome data was subjected to quality control that included number of unique molecular identifiers (UMI), genes detected per cell, and the percentage of mitochondrial reads detected per cell. This yielded a total of 1969 cells for downstream analysis. Clustering of the single-cell transcriptome data using Seurat identified eight major clusters in the data (Fig. 1B-D). Based on the expression of marker genes, the clusters corresponded to acinar (*ela2*), ductal (*cftr, hnf1ba, her15.1, anxa4;* Supp. Fig. 1A), endothelial (*vsg1*), β-cells (*insa*), α-cells (*gcgb*), ɛ-cells (*ghrl*), and two discrete types of δ-cells (*sst1.1* and *sst2*) (Fig. 1B-C) (Table 1). The two groups of δ-cells include a population marked by *sst2 (δ2)* (Li et al., 2009) and a distinct type that expresses *sst1.1 (δ1)*. Each of the two *sst* genes, which arose through gene-duplication in teleost fish is predicted to encode for a full-length and functional somatostatin protein (Devos et al., 2002; Liu et al., 2010). One notable difference between the δ1- and δ2-cell types is the preferential expression *of pancreatic and duodenal homeobox 1* (*pdx1)* in the former group (Fig. 2A). A number of other genes were differentially expressed between the two populations (Fig. 2B and Table 2). Thus, the zebrafish islet is equipped with two distinct δ-cell types, each expressing a different *sst* paralogue.

**The zebrafish islets contain a persistent bi-hormonal cell population**

Most cell populations we identified contained discrete hormonal cell-types. One exception occurs in the cells of the *ins-* and *sst1.1-*lineages. Notably, ~8 % of all *ins-*expressing cells (60 cells out of 746) were positive for both *ins* and *sst1.1* (Fig. 1E). The bi-hormonal cells had further distinguishing features. The cells expressed *dkk3b*, a Dickkopf-related secreted protein, and *pyyb*, a short peptide related to

Protein YY (Fig. 1E). Interestingly, *dkk3b* and *pyyb* were also enriched in δ1-cells (Fig. 2).

To confirm the co-expression of *sst1.1* and *ins*, we utilized the single-cell transcriptomic profiles of sorted β-cells from seven stages of zebrafish life including the juvenile and adult periods (Singh et al., 2018). We detected above-background expression of *sst1.1* in a fraction of *ins*-expressing cells at each of the seven stages (Supp. Fig. 2A). The fraction of *sst1.1*-expressing cells ranged from 5 – 9 % of the *ins*-positive population throughout the life of the fish, showing that the bi-hormonal population maintains steady proportions.

Using Assay for Transposase-Accessible Chromatin (ATAC) of sorted β-cells, we detected open chromatin peaks in regulatory elements of *ins*, *sst1.1* and *dkk3b* (Supp. Fig. 2B). This suggests that *ins*-expressing cells have transcriptionally favorable chromatin for expression of genes enriched in the bi-hormonal cells.

**Histological assessment of the bi-hormonal cells in the zebrafish pancreas.**

To directly visualize the bi-hormonal cells in the zebrafish pancreas, we took advantage of a transgenic reporter line, *Tg(sst1.1: EGFP-Ras)* (Löhr et al., 2018) in which the regulatory sequences of the *sst1.1* gene drive a membrane localized green fluorescent protein (Fig. 3A). Within islets, the GFP-positive δ1-cells formed an intricate mesh of cells interconnected by long cellular processes. We stained the pancreas from *Tg(sst1.1:EGFP-Ras)* larvae with antibodies against insulin to mark the β-cells. Confocal imaging of the islets demonstrated the presence of individual cells that co-expressed both EGFP and insulin (Fig. 3B). We further confirmed the co-expression of somatostatin and insulin proteins using immunofluorescence in pancreatic sections from adult fish aged 7 and 16 mpf (Fig. 3C-D). We clearly

detected immuno-reactivity for the two hormones along with the membrane GFP maker in single cells, corroborating the presence of bi-hormonal cells in the islet. In these cells, insulin showed higher concentration in the cytoplasm whereas somatostatin was localized closer to the cell membrane, which may reflect differential distribution of secretary granules for the two hormones. Based on these results, along with the single-cell transcriptomic data, we have identified cells that express at the RNA and protein levels both insulin and somatostatin and persist from the early stages of development to adulthood.

**The bi-hormonal cells exhibit glucose-stimulated calcium influx *in vivo***

In order to investigate if the bi-hormonal cells are glucose-responsive and present influx of calcium ions upon glucose stimulation, we performed 2-Photon calcium imaging *in vivo* using *Tg(sst1.1:EGFP-Ras); Tg(ins:K-GECO1)* double transgenic larvae. We used our previously developed method for calcium imaging (Salem et al., 2019). We found that cells co-expressing GFP and K-GECO1 show glucose-stimulated calcium influx, which was comparable to the response of β-cells in terms of speed and amplitude of response (n=3 islets) (Fig. 3E-F). Thus, the hybrid cells are functionally similar to β-cells in their capacity to respond to glucose.

**The bi-hormonal cells are molecular hybrids between δ1- and β-cells.**

To characterize the bi-hormonal cells at a better molecular resolution than the one provided by the 10x Genomics pipeline, we utilized the *Tg(sst1.1:EGFP-Ras); Tg(ins:BB1.0L)* double transgenic line to separate the hybrid cells from β- and δ1-cells using index flow cytometry (Fig. 4A). We performed single-cell transcriptomic using the Smart-Seq2 protocol, which has higher sensitivity than droplet-based single-cell sequencing (Ziegenhain et al., 2017). The single-cell sequencing

revealed enrichment of *ins* and *sst1.1* in δ1-cells and β-cells, respectively, validating the utility of the reporter lines to distinguish the two cell populations (Fig. 4B-C). Further, we observed a strong Pearson correlation in the co-expression of *sst1.1* and *dkk3b* as well as of *sst1.1* and *pyyb* in individual cells (Fig. 4D), validating our previous observations. Differential gene expression analysis showed that β-cells express significantly higher ($p_{adj} < 0.05$) levels of *nkx6.2* and *mnx1* (Fig. 4E), genes involved in their specification and maturation (Binot et al., 2010; Pan et al., 2015). Conversely, δ1-cells expressed HHEX (Fig. 4E), a transcription factor involved in the maintenance of δ-cell identity (Zhang et al., 2014). Interestingly, we observed intermediate expression levels for *nkx6.2*, *mnx1* and *HHEX* in the bi-hormonal cells, revealing an overlap of β- and δ-cell identity genes within the mixed population.

**The bi-hormonal cells express endocrine-progenitor markers**

Next, we applied pseudotemporal analysis using Monocle (Trapnell et al., 2014) to model cellular transitions. The pseudotemporal model showed that the bi-hormonal cells were positioned in between β- and δ1-cells (Fig. 4F), confirming the hybrid signature of these cells. We further characterized gene expression dynamics by calculating expression changes along the pseudotime (Fig. 4F'). As expected, transcription factors implicated in β-cell lineage and identity (*nkx6.2*, *mnx1*) were enriched towards the δ1- side of the pseudotime. Moreover, *proprotein convertase subtilisin/kexin type 1* (*pcsk1*) (Benjannet et al., 1991), a gene necessary for conversion of pre-proinsulin to functional insulin hormone displayed a similar trend. In contrast, genes implicated in δ-cell identity (*HHEX*), along with pathways controlling endocrine progenitor maintenance such a Wnt, Fgf, Notch, and Sphingolipid signaling (*s1pr1* and *s1pr2*) were associated with the trajectory towards

the δ1-cells (Ninov et al., 2012; Serafimidis et al., 2017; Seymour et al., 2012; Sharon et al., 2019; Zhang et al., 2014). The hybrid cells expressed intermediate levels of these factors and pathway components, as compared to either β or δ1-cells. The expression of the pathway components involved in progenitor maintenance was confirmed by bulk RNA-seq of sorted cells (Fig. 4G), which also revealed enrichment of *fev*, a transcription factor that was recently shown to be expressed in endocrine progenitors in mice (Byrnes et al., 2018). Altogether, we conclude that the hybrid cells display the transcriptional profiles of intermediate cell states between δ1- and β-cells, and exhibit an enrichment of pathways associated with endocrine progenitors.

**The hybrid cells increase during β-cell regeneration**

Intrigued by the association of precursor characteristics of hybrid cells, we further characterized their behaviors during β-cell regeneration in larval zebrafish. Near-complete β-cell death was induced using the Nitroreductase (NTR) system (Curado et al., 2007; Pisharath et al., 2007). In this system, *Tg(ins:FLAG-NTR)* transgenic zebrafish, expressing NTR (nitroreductase) specifically in the β-cells, are exposed to Mtz (metronidazole). NTR converts Mtz to a cytotoxin, which specifically kills the β-cells while sparing all other cells in the body. To label *sst1.1*-expressing cells during β-cell regeneration, we utilized *Tg(ins:FLAG-NTR)*; *TgBAC(sst1.1:EGFP-Ras)* double transgenic lines. We exposed *Tg(ins:FLAG-NTR)* larvae to Mtz for 24 hours, and collected samples for observation 4 and 8 days post ablation (dpa) (Supp. Fig. 3). At both stages, we observed a dramatic increase in the percentage of hybrid cells within the regenerating islet (Supp. Fig. 3B-C). Of note, when we examined the expression of *sst2*:RFP, which marks the population of

δ2-cells (δ2) in zebrafish (Li et al., 2009), we did not observe the formation of hybrid cells (unpublished data),

**A single-cell map of adult β-cell regeneration reveals a major contribution of hybrid cells to insulin expression**

To assess the dynamics of hybrid cells during the more physiologically-relevant setting of adult β-cell regeneration, we performed single-cell sequencing (using the 10X genomics pipeline) at different times following β-cell destruction and regeneration in adult zebrafish.  We selectively ablated the β-cells and dissected the primary islets and surrounding tissues, including the exocrine cells from adult zebrafish at 0, 2, 7, and 14 dpa.  These time-points were chosen as they include the time of β-cell destruction, the subsequent hyperglycemia, followed by normalization of glucose levels after two weeks of regeneration (Supp. Fig. 4) (Delaspre et al., 2015).  To induce efficient ablation of insulin-expressing cells, we used a new ablation line, which employs an improved and more effective version of the NTR enzyme (Supp. Fig. 4) (Singh et al., 2017).  We identified all major cell types, including discrete populations of *ins*-expressing β-cells, *sst1.1*-expressing δ-cells, *sst2*-expressing δ2-cells, ductal and acinar cells (Fig. 5).  Shortly upon β-cell ablation (at 0 and 2 dpa), the *ins*-only population observed in controls was nearly-completely lost based on single-cell gene expression analysis.  Instead, *ins* expression was detected mainly within the δ1-population, in which cells co-expressed both *ins* and *sst1.1* (Fig. 5 and Supp. Fig. 5).  At 7 and 14 dpa, the presence of hybrid cells was even more prominent with higher proportions of *ins*$^{hi}$ cells in the cluster corresponding to δ1-cells (Supp. Fig. 6).  Importantly, clustering of the integrated data from all time-points revealed that the hybrid cells within the *sst1.1* cluster are the main source of insulin expression during regeneration (Fig. 5D).  Other endocrine

clusters, such as the *sst2*-expressing δ2-cells, as well as α-cells, show a relatively minor contribution to insulin expression (Fig. 5D,E). Analyzing expression profiles across time-points and cell types showed that insulin expression increases in the cluster corresponding to δ1-cells, whereas the rest of the pancreatic cell types retained low *ins* levels over time (Supp. Fig. 6). Moreover, upon β-cell destruction, we detected increased expression in *pcsk1* and *pcsk2* in the *sst1.1*-expressing cells, which are necessary for insulin maturation (Supp. Fig. 7). These changes in gene-expression were accompanied by an increase in the number of *sst1.1*-expressing cells relative to ablation-invariant α-cells (Supp. Fig. 8). At 7 dpa, the relative number of *sst1.1*-expressing cells has stabilized at roughly twice as many as in controls.

To investigate if the hybrid cells may resolve into *sst1.1+* and *ins+* populations during regeneration, we analyzed the single-cell transcriptome data from 7 and 14 dpa (Fig. 6). 7 dpa represents an intermediate stage while 14 dpa corresponds to the resolution of diabetes. By specifically focusing on cells that express the two hormones, we observed overlap between the *sst1.1+* and *ins+* expressing cells at 7 dpa (Fig. 6A,B). By contrast, at 14 dpa the cells segregated into two clusters (Fig. 6D,E). Notably, the two clusters observed at 14 dpa were enriched for the expression of either *sst1.1* or *ins* (Fig. 6E), indicating that the two cell populations had diverged. We further performed gene correlation analysis to examine the distribution of *sst1.1* and *ins* gene expression level at cellular resolution. At 7 dpa, a positive correlation was observed between *ins* and *sst1.1*-expression (Fig. 6C), suggesting the presence of hybrid cells at this stage. At 14 dpa, however, individual cells displayed a negative correlation between *sst1.1* and *ins*-expression (Fig. 6F), suggesting a trend towards the resolution of gene expression into β- or δ1-cells.

Altogether, the single-cell atlas of β-cell regeneration reveals that the hybrid cells form during initial stages of pancreas regeneration, followed by β-cell formation.

**Live-cell tracking reveals hybrid-cell transition during regeneration**

To assess if the bi-hormonal cells arise *de novo* during β-cell regeneration or whether they might reflect pre-existing cells that have survived the ablation procedure, we performed live imaging of islets undergoing regeneration during the optically translucent larval stage. To label specifically the insulin-expressing cells, we used an *ins*:tdTomato reporter, whereas to label the *sst.1*-expressing cells, we used the *sst1.1*:EGFP-Ras reporter. Ablation of β-cells was induced using *Tg(ins:FLAG-NTR)* by Mtz-treatment from 3 to 4 dpf. First, we measured the efficiency of cell ablation and found that both β-cells and hybrid cells were efficiently ablated (Fig. 7A). We then monitored longitudinally the regeneration process in individual larvae. Strikingly, δ1-cells showed dramatic changes in morphology, as they accumulated prominent membrane-bound vacuolar structures within their cytosol evident both under confocal and electron microscopy (Supp. Fig. 9). Moreover, within islets that previously had lost any tdTomato expression, we now observed its re-expression in EGFP-positive cells (Fig. 7A). These data show that NTR-mediated ablation efficiently kills the pre-existing insulin-expressing cells and that in a matter of two days following β-cell destruction, new hybrid cells emerged *de novo.*

**The newly emerging bi-hormonal cells exhibit glucose-stimulated calcium influx *in vivo***

In order to investigate if the emerging bi-hormonal cells, which arise post β-cell ablation, are glucose-responsive and present a calcium influx upon glucose

stimulation, we performed *in vivo* calcium imaging using *Tg(sst1.1:EGFP-Ras);*

*Tg(ins: Flag-NTR);* and *Tg(ins:K-GECO1)* triple-transgenic larvae. We specifically

ablated β-cells by exposing the larvae to Mtz from 3 to 4 days post-fertilization (24

hours) and performed *in vivo* calcium imaging at 2, 4- and 6-days post β-cell ablation

(Movies 1 and 2). Interestingly, we found that the emerging bi-hormonal cells

became glucose-responsive as early as 4 days post-β cell ablation (Fig. 7B-C)

(Movie 2). The bi-hormonal cells showed a comparable glucose-responsiveness to

GFP-negative but insulin-positive cells (Fig. 7D), which likely arise via constitutive

developmental neogenesis in larvae. The glucose responsiveness of bi-hormonal

cells, however, was lower than that of β-cells in unablated animals, likely because

the cells were undergoing maturation. Thus, our results indicated that the emerging

bi-hormonal cells acquire responsiveness to glucose stimulus.

**Overexpression of *dkk3b* increases hybrid-cell formation**

Our initial analysis of gene-expression of the hybrid cells indicated that the cells

are enriched for the expression of *dkk3b*, which is also shared with δ1-cells.

Moreover, upon β-cell ablation, *dkk3b* was among the genes that showed a trend of

increasing expression within the δ1-cell cluster over time (Supp. Fig. 7). To test if

*dkk3b* plays a role in hybrid-cell formation, we generated transgenic lines with

constitutive expression of *dkk3b* under the *insulin* promoter, or with inducible

expression based on the *heat-shock promoter* (hsp). In both cases, we observed

that the upregulated expression of *dkk3b* in the islets let to increased proportions of

insulin and somatostatin double-positive cells during development (Fig. 8A,B). Thus,

the forced overexpression of *dkk3b*, which is normally enriched in hybrid and δ1-

cells, was sufficient to promote the formation of bi-hormonal cells without injury.

## Discussion

In this study, we utilized single-cell sequencing to characterize cellular dynamics in a regeneration-competent animal model – the zebrafish (Fig. 8C). Using this approach, we defined the gene-expression of different pancreatic cells upon β-cell destruction and established an atlas of β-cell regeneration, spanning the period from cell destruction to emergence of new insulin-expressing cells. Among the different cell types, we characterized a novel population of δ-cells, expressing the *sst* paralogue *sst1.1*, which is consistent with previous studies showing its expression in the zebrafish pancreas (Dalgin et al., 2011; Devos et al., 2002; Spanjaard et al., 2018). In addition, the single-cell resolution of our analysis allowed us to pin-point a smaller population of bi-hormonal cells, which exist under a hybrid β/δ-cell identity.

We show that the hybrid cells share the hormones, fate-determinants and glucose responsiveness of both δ1- and β-cells. While the cells are transcriptionally distinct from β-cells, they show glucose-stimulated $Ca^{2+}$ influx. We validated the presence of hybrid cells at all the stages of zebrafish life we have investigated: from 4 dpf to 16 mpf. Moreover, using single-cell RNA-Sequencing, we compared the transcriptional profile of the hybrid cells to mono-hormonal β-cells and δ1-cells. The differential gene expression analysis revealed an overlapping expression of genes involved in both β and δ-cell fate-specification and function. For example, the cells exhibit intermediate expression of genes involved in insulin prohormone processing, as compared to β-cells. Moreover, these cells express at intermediate levels transcription factors that differentiate between the β- and δ-cell lineages, such as the evolutionary conserved homeobox *Mnx1* and *HHEX*, respectively (Dalgin et al., 2011; Pan et al., 2015; Zhang et al., 2014). In parallel-unpublished work aimed to

define the lineage decisions in the developing endoderm, we find that the population of *sst1.1/ins* cells becomes evident as early as the embryonic islet forms, maintaining similar proportions as in the mature adult islet (~8% of the insulin-expressing cells). Together, the deep-sequencing experiments suggest that the hybrid cells are an intermediate-lineage population of pancreatic cells.

Polyhormonal endocrine cells have been observed during embryonic pancreas development in mammals and during regeneration (De Krijger et al., 1992; Teitelman et al., 1993). However, it remains unknown if such cells play a significant role in the pancreas. To test their role, we performed two parallel experiments. First, we asked if the hybrid cells are glucose-responsive. We found that under steady state, the cells behave similar to β-cells, as they exhibit a robust glucose-stimulated calcium influx. Moreover, during regeneration, the emerging hybrid cells acquire the capacity to undergo calcium influx in response to glucose-stimulation. The functional role of the cells in the context of regeneration is supported by the second experiment we carried out: to follow the time course of regeneration using single-cell transcriptomics and to correlate insulin-expression with reversal of diabetes. While there was no evidence for the presence of a major group of regenerated pure β-cells even 14 days after destruction, by this time, the fasting glucose was back to normal, with insulin being expressed predominately in the hybrid cells. Moreover, there was increased expression of *pdx1* and *pcsk1*, which is in line with them acting to restore the islet's insulin-producing function. Therefore, we show that the hybrid cells fulfill a role in glucose-regulation in the absence of β-cells.

An important open question remains as to the cellular sources of the hybrid cells during regeneration. Our observations point towards the subpopulation of *sst1.1*-expressing cells, which we suspect may increase insulin production upon β-

cell destruction, thereby forming hybrid cells. Intriguingly, in mice, δ-cells have been shown to be plastic in juvenile animal but lose their plasticity in adulthood (Chera et al., 2014). It is possible that adult zebrafish maintain an active population of δ1-cells, which covert into hybrid cells and make it possible to restore normoglycemia. However, we cannot exclude the possibility that the centroacinar subpopulation of terminal duct cells, also undergo a process of endocrine differentiation, passing through a bi-hormonal state, which may further resolve into β-cells over time. Differentiating between these two sources will required the development of tools for lineage tracing of δ1-cells in the zebrafish pancreas.

A second important question will be to define the signaling pathways the promote the formation and resolution of such hybrid cells. Here we show that hybrid cells express *dkk3b,* and its overexpression is sufficient to generate hybrid cells in the absence of injury. Since *dkk3b* is predicted to be a secreted factor, the precise cell population that responds to its overexpression to generate hybrid cells remains to be defined. *dkk3b* was previously shown to be expressed in an unknown group of islet cells in zebrafish (Untergasser et al., 2011), and we now reveal its association with the *sst1.1*-expressing δ-cells. Moreover, the human DKK3 is one of the first examples of a heterogeneously expressed gene in islets (Hermann et al., 2007). Notably, *DKK3* is also upregulated in islets from donors with type 2 diabetes (Solimena et al., 2018). The role of this interesting gene in the context of diabetes remains to be studied.

One of the take-home messages of our study is that regenerating imperfect yet functional hybrid cells capable of insulin-production may offer the opportunity to elicit glucose control. Perhaps, such surrogate cells would be stealthy in terms of escaping immune cell recognition and destruction in type 1 diabetes, as they exist

under a hidden identity, resembling two different hormonal cell types at the same time.

## MATERIALS AND METHODS

### Zebrafish strains and husbandry.

Wild-type or transgenic zebrafish of the outbred AB, WIK, or a hybrid WIK/AB strain were used in all experiments. Zebrafish were raised under standard conditions at 28 °C. Animals were chosen at random for all experiments. Published transgenic strains used in this study were *Tg(ins:BB1.0L; cryaa:RFP)* (Singh et al., 2017), *Tg(sst1.1:EGFP-Ras)[fr40Tg]* (Löhr et al., 2018), *Tg(fabp10:dsRed, ela3l:GFP)[gz12]* (Wan et al., 2006), *Tg(ins:FLAG-NTR)[s950]* (Andersson et al., 2012), *Tg(ins:YFP-2A-NTR3,cryaa:mCherry)[tud201Tg]* (Singh et al., 2017), and *Tg(sst2:RFP)* (Li et al., 2009). Experiments were conducted in accordance with the Animal Welfare Act and with permission of the Landesdirektion Sachsen, Germany (permits: TVV38/2015, TVV 45/2018, TVV33/2019, TVV 32/2020 and all corresponding amendments)

### Single-cell suspension of zebrafish pancreatic islets and liver cells.

Single cell suspension of zebrafish pancreas was performed according to the protocol outlined previously (Singh et al., 2018). In brief, primary islets and substantial surround tissue were dissected and dissociated into single cells by incubation in TrypLE (ThermoFisher, 12563029) with 0.1% Pluronic F-68 (ThermoFisher, 24040032) at 37 °C in a benchtop shaker set at 450 rpm for 30 min. During the dissection, we used the fluorescent signal of beta-cells in the *Tg(ins:BB1.0L; cryaa:RFP)* or *Tg(ins:YFP-2A-NTR3,cryaa:mCherry)* reporters in order to orient ourselves in respect to the anatomical position of the main pancreatic islets. Following dissociation, TrypLE was inactivated with 10% FBS, and the cells pelleted by centrifugation at 500g for 10min at 4 °C. The supernatant was carefully

discarded and the pellet re-suspended in 500 uL of HBSS (without Ca, Mg) + 0.1%

Pluronic F-68. To remove debris, the solution was passed over a 30 μm cell filter

(Miltenyi Biotec, 130-041-407). To remove dead cells, calcein violet (ThermoFisher,

C34858) was added at a final concentration of 1 μM and the cell suspension

incubated at room temperature for 20 minutes.  The single cell preparation was

sorted with the appropriate gates for identification of β-cells and alive cells

(calcein+). FACS was performed through 100 μm nozzle.  Dissociation of liver tissue

was performed in the same manner, except liver was used instead of the primary

islet. For this, liver was collected from *Tg(fabp10:dsRed, ela3l:GFP)*, in which the

hepatocytes are labeled with red fluorescence.

For samples from β-cell regeneration time-points, the adult fish were treated

with Mtz (Sigma-Aldrich, M3761) which was dissolved at 10mM concentration in fish

water. The animals were treated for 24 hours, protected from light. After 24h, Mtz

solution was removed and the animals were placed 3 times in fresh fish water before

returning them to standard maintenance conditions. Both untreated and Mtz-treated

animals were euthanized by an overdose of tricaine. Samples for each time-point

comprised of 6 animals (3 males and 3 females). Single-cell suspension was

prepared in the same manner (as highlighted above) and 15000 live-cells were

FACS sorted for 10X Genomics.

**Single-cell RNA-Seq. of the zebrafish pancreas.**

The sorted cells (38 μl) were carefully mixed with reverse transcription mix

before loading the cells on the 10X Genomics Chromium system (Zheng et al., 2017)

in a Chromium Single Cell B Chip and processed further following the guidelines of

the 10x Genomics user manual. In short, the droplets were directly subjected to

reverse transcription, the emulsion was broken and cDNA was purified using Silane

beads. After the amplification of cDNA with 11 cycles, it underwent purification and quantification.  The 10X Genomics single cell RNA-seq library preparation - involving fragmentation, dA-Tailing, adapter ligation and a 13 cycles indexing PCR – was performed based on the manufacturer's protocol. After quantification, the libraries were sequenced on an Illumina Novaseq6000 in paired-end mode (R1: 29 cycles; I1: 8 cycles; R2: 93 cycles), thus generating ~50-150 mio. fragments. The raw sequencing data was then processed with the 'count' command of the Cell Ranger software (v4.0.0) provided by 10X Genomics with the option '--expect-cells' set to 7500 (all other options were used as per default). To build the reference for Cell Ranger, zebrafish genome (GRCz11) as well as gene annotation (Ensembl 95) were downloaded from Ensembl and the annotation was filtered with the 'mkgtf' command of Cell Ranger (options: '--attribute=gene_biotype:protein_coding --attribute=gene_biotype:lincRNA –attribute=gene_biotype:antisense').  Genome sequence and filtered annotation were then used as input to the 'mkref' command of Cell Ranger to build the appropriate Cellranger Reference.

**Analysis of single-cell RNA-Seq of the zebrafish pancreas.**

We re-analyzed the single-cell RNA-Seq of the zebrafish pancreatic cells, which was generated using the 10x Chromium pipeline and previously published (Salem et al., 2019).  The data is deposited at the GEO repository with an accession number GSE123662.  For clustering and gene expression analysis, we utilized Seurat and followed the recommended analysis pipeline.  Briefly, the raw data as UMI-counts was log-normalized, regressed to remove the effect of library size, and scaled. Highly variable genes were identified for PCA analysis and graph-based clustering. Marker genes identified for each cluster were used to classify the exocrine,

endocrine and ductal population.  The endocrine cluster was isolated and sub-

clustered to identify and label individual endocrine populations.

To define the changes the abundance of δ1-cells during regeneration, at each time

point, we calculated the ratio between the number of *δ1-cells* and α-cells, filtering out

any clusters that expressed multiple hormones apart from the combination of *sst1.1*

and *ins*. We calculated 99% confidence intervals from a 1000-fold random sampling

of the total number of cells present at every time from a binomial distribution with the

observed cell type frequencies as probabilities, using the function rbinom in base R.

**ATAC-Seq. of zebrafish β-cells and hepatocytes.**

ATAC-Seq. was carried out using 50,000 cells collected using FACS.  For FACS,

single-cell suspension was generated as mentioned above.  For ATAC-Seq. of β-

cells, tissue for dissociation was collected from *Tg(ins:BB1.0L; cryaa:RFP)* line and

β-cells sorted using the red fluorescence.  For hepatocytes, liver was collected from

the *Tg(fabp10:dsRed, ela3l:GFP)* line and cells sorted using the red fluorescence.

Cells were processed according to the protocol outlined in (Buenrostro et al., 2013).

In brief, cells were homogenized in a 2 ml Dounce homogenizer with 5 strokes in 1X

homogenization buffer on ice.  The nuclei suspension was centrifuged at 500g for 10

min at 4 °C.  After removing the supernatant, transposition was carried out by

incubating the nuclei in the transposition reaction mix for 30 min at 37 °C.

Transposed DNA was purified using Qiagen MinElute Reaction Cleanup Kit.

Barcoded libraries were generated from the isolated DNA by PCR amplification for

11 cycles. Libraries were sequenced on Ilumina HiSeq2500 in 2 X 75 bp paired-end

mode.  We conducted two biological replicates for β-cells and one biological

replicate for hepatocytes.  Processing of ATAC-seq. raw data was done according to

the pipeline from the Kundaje lab (Koh et al., 2016). MACS2 (Feng et al., 2012) was

used to identify the peak regions. Only peaks present in both the biological replicates of β-cells and absent in the hepatocytes were retained. The β-cell specific peaks were exported as a bed file and visualized in IGV viewer (Robinson et al., 2011).

**Transcriptional profiling and analysis of β-, δ1-, and hybrid cells.**

Collection of β-, δ1-, and hybrid cells for transcriptional profiling was performed using FACS of single-cell suspension of dissected islets from *Tg(ins:BB1.0L); Tg(sst1.1:EGFP-Ras)*. For single-cell profiling using Smart-Seq., cells were collected in 96-well plate using index sorting and processed according to the previously published pipeline (Singh et al., 2018). Analysis was performed using Seurat (Satija et al., 2015), with cells classified using their position on the FACS plot (δ1-cells: Green+, Red-; hybrid cells: Green+, Red+; and β-cells: Green-, Red+). For bulk-sequencing and analysis, 2,000 cells of each type were collected in a 1.5 ml tube and processed according to the pipeline published in (Janjuha et al., 2018). Differential gene expression analysis was analyzed using DESeq2 (Love et al., 2014).

**Pseudotemporal analysis of single-cell RNA-seq. data.**

Pseudotemporal analysis was performed using Monocle 2 (Trapnell et al., 2014) following the recommended analysis protocol (http://cole-trapnell-lab.github.io/monocle-release/docs/). Briefly, raw count data was imported into a Monocle object, normalized using the estimated size factors, and filtered to remove low quality genes and cells. Gene dispersion was estimated using negative binomial distribution with fixed variance,and used to identify the highly variable genes in an unbiased manner for pseudotemporal ordering. Pseudotemporal trajectory was constructed using the 'DDRTree' algorithm, and plotted on a 1-D graph with jitter to enhance visualization. Cells in the graph were colored according to their FACS plot

identity.  Gene Expression dynamics in relation to pseudotime was calculated using the command 'differentialGeneTest (Monocle_object, fullModelFormulaStr = "~sm.ns(Pseudotime)")', and plotted using the heatmap function for visualization.

**Diffusion pseudotime analysis of single-cell RNA-seq. data**

For single cell trajectory inference, diffusion pseudotime was computed for 7 dpa and 14 dpa and solely on selected cells (*ins+* and *sst1.1+* cells) following the standard analysis pipeline (Haghverdi et al., 2016) provided by Scanpy package.  Since a single lineage of a timepoint was selected, the number of branchings was set to 1. For cell trajectory analysis, diffusion maps were developed.  For 14 dpa, the *sst1.1-*high cells were selected as root cells to capture the transition of δ1-cells. To visualize the marker genes trajectory on trendline plot, plot.wishbone_marker_trajectory function was used provided by the Wishbone package (Setty et al., 2016).

**Tissue collection for imaging.**

To facilitate confocal imaging of the islets, the pancreas was dissected from fish (larval) or the gut (juvenile and adults) after fixation.  Fish were killed in Tricaine prior to either direct fixation or dissection of gut, and the samples immersed in 4% paraformaldehyde + 1% Triton-X for 2 days at 4 °C.  The pancreas was then manually dissected and washed multiple times in PBS.

**Immunofluorescence and image acquisition.**

Whole-mount immunofluorescence was performed on pancreas collected as described above*.*  The collected samples were permeabilized in 1% PBT (Triton-X-100) and blocked in 4% PBTB (BSA).  Primary and secondary antibody staining were performed overnight at 4 °C.  Primary antibodies used in this study were anti-insulin (guinea pig, Dako A0564)/insulin (rabbit, Genetex 128490) at 1:200, anti-

somatostatin (rat, Genetex GTX3906) at 1:200, and anti-EGFP (chicken, Abcam ab13970) at 1:250. Secondary antibodies at 1:500 dilutions used in this study were Alexa Fluor 568 anti-guinea pig, Alexa Fluor 568 anti-rabbit, Alexa Fluor 647 anti-rabbit and Alexa Fluor 647 anti-rat. Samples were mounted in Vectashield and imaged using a Zeiss LSM 780. ImageJ was used to add scale bars and PowerPoint was used for adding arrows and labels.

**Live-imaging during β-cell regeneration.**

From 1dpf onwards, *Tg (ins: FLAG-NTR)*; *TgBAC (sst1.1: EGFP Ras), Tg (ins: BB1.0L)*-triple transgenic zebrafish larvae were treated with 0.003% 1-phenyl thio-urea (PTU) (200μM) to inhibit pigmentation. For β-cell ablation, 3 dpf larvae were incubated with 10 mM Mtz (Sigma-Aldrich, M3761) dissolved in E3 medium containing 1% DMSO, and maintained for 24 h in the dark before they were rinsed and returned to fresh E3 medium. E3 medium with 1% DMSO was used as control. Following the ablation, individual larvae was mounted in a glass-bottom micro-well dish (MakTek Corporation) in 1% low-melting agarose (LMA) containing 0.4g/litre tricaine. After the agarose was solidified, the dish were filled with E3 medium containing 0.4g/litre tricaine. Live-imaging was performed on an upright laser-scanning confocal microscope, Zeiss LSM 780. EGFP and dsRed signals were acquired simultaneously using the 488nm and 561 nm laser lines. Laser power was maintained as low as possible to minimize photo-toxicity.

***In vivo* calcium-imaging in zebrafish larvae.**

Live-imaging and glucose-injections were performed as described previously (Salem et al., 2019)

**Generation of *Tg(ins:nls-mCerulean-T2A-dkk3b; cryaa:mCherry)* zebrafish line.**

The dkk3b coding sequence (CDS) was purchased from Zebrafish Gene Collection

(ZGC) cDNA clones (MGC: 162332). The CDS was PCR amplified to flank the

cDNA with SpeI/PacI restriction sites. The PCR product was digested with SpeI/PacI

restriction enzymes. The ins:nls-mCerulean*; cryaa:mCherry* construct (used for

generating *Tg(ins:nls-mCerulean; cryaa:mCherry)* line) was digested with SpeI/PacI.

The two DNA fragments were ligated and transformed to generate ins:nls-

mCerulean-T2A-dkk3b; cryaa:mCherry plasmid. The entire construct is flanked with

I-SceI sites to facilitate transgenesis. I-SceI based transgenesis was carried out, and

founders selected based on mCherry (red) fluorescence in the eye, and mCerulean

(blue) expression in the pancreatic islet.

**Generation of *Tg(Tol2-hsp70l:mCherry-T2A-dkk3b)* zebrafish line.**

The dkk3b coding sequence (CDS) was purchased from Zebrafish Gene Collection

(ZGC) cDNA clones (MGC: 162332). The CDS was PCR amplified to flank the cDNA

with NheI/BglII restriction sites. The PCR product was digested with NheI/BglII

restriction enzymes and was fused to mCherry using the T2A sequence. Thereafter,

mCherry-t2a-dkk3b sequence was cloned downstream of the hsp70l promoter

region. The entire construct is flanked with Tol2 sites to facilitate transgenesis. Tol2

based transgenesis was carried out, and founders selected based on ubiquitous

expression of mCherry (red) fluorescence in the zebrafish larvae upon heat-shock at

37 degrees.

**Generation of *Tg(ins:K-GECO1; cryaa:mCherry)* zebrafish line.**

For the construction of the *Tg(ins:K-GECO1; cryaa:mCherry)* line, we used PCR

amplification with primers designed to introduce 5' EcoRI and 3' PacI restriction

enzymes sites in the cDNA for K-GECO1. The established plasmid backbone

containing ins:mAG-zGeminin; cryaa:mCherry (Ninov et al., 2013) which was

digested with EcoRI/PacI and K-GECO1 was ligated using the EcoRI/PacI sites. The construct was flanked with I-SceI sites to facilitate transgenesis. Several founders were screening and a founder with Mendelian segregation were selected for expanding the transgenic line.

**Electron Microscopy**

Correlative Light and Electron Microscopy (CLEM) of immunolabeled resin sections was performed as previously described (Fabig et al., 2012).  In brief, WT and ablated zebrafish larvae were embedded in Lowicryl K4M using the progressive lowering of temperature method (Carlemalm et al., 1982).  Ultrathin resin sections were cut on Leica UC6 ultramicrotrome (Leica Microsystems Wetzlar, Germany), labeled with primary anti-GFP (rabbit-anti-GFP, TP401 from Torrey Pines) followed by consecutive incubations with protein A 10 nm gold, goat-anti-rabbit Alexa 488, and DAPI. Sections were analysed on a Keyence BZ 8000 fluorescence microscope, contrasted with uranyl acetate, dried, and imaged on a Jeol JEM 1400-Plus transmission electron microscope (Jeol, Freising, Germany) running at 80 kV.

**Statistical analysis.**

Statistical analysis was performed using R and GraphPad Prism. No animals were excluded from analysis.  Blinding was not performed during analysis.  Analysis of normal distribution was not performed.  To compare the proportion of *sst1.1* expressing β-cells during regeneration, an unpaired two-tailed t-test with unequal variance (t-test(x = dataframe, alternative = "two.sided", paired = FALSE, var.equal = FALSE)) was used.  A p-value of <0.05 was considered statistically significant. For plotting, Tukey style boxplots showing the median along with the 25th percentile (lower quartile - Q1) to 75th percentile (upper quartile - Q3) range were used.  For all boxplots, whiskers extend to 1.5 times the interquartile range (IQR = Q3 − Q1). In

addition, individual data points are plotted on the graphs. To compare the proportions of bi-hormonal cells during time-course imaging, to evaluate the AUC during in-vivo calcium imaging and the bi-hormonal cell number upon dkk3b over-expression, Mann-Whitney unpaired t-test (non-parametric test) was used for statistical analysis. The graphs were prepared using Graph Pad prism software and p-value of $<0.05$ was considered statistically significant.

**Data availability**

The combined data set of the time course of regeneration will be provided with the manuscript as a publically available and searchable database in which gene expression and cell types can be visualized.

Datasets generated during this study are publicly available.

1. Single-cell RNA sequencing of zebrafish pancreatic cells
   GSE123662 Single-cell RNA sequencing of zebrafish pancreatic cells (GSE123662) (Salem et al., 2019).

2. SuperSeries (Collection of two datasets)
   Single-cell and Bulk analysis of pancreatic somatostatin and β-cells from zebrafish (GSE152697, Reviewer Token: chshyqkofvmhnyf)

   - Transcriptional profiling of pancreatic somatostatin and β-cells from zebrafish [scRNA-seq]
     RNA-Sequencing on FACS sorted cells expressing sst1.1, ins, or both sst1.1 and ins. (GSE152695)

   - Transcriptional profiling of pancreatic somatostatin and β-cells from zebrafish [bulk RNA-seq]
     RNA-Sequencing on FACS sorted cells expressing sst1.1, ins, or both sst1.1 and ins (GSE152696)

3. ATAC sequencing of zebrafish pancreatic β-cells and hepatocytes (GSE152199, reviewer token wzspysiqbnylvax)

**Contributions:**

Conceptualization: S.P.S, N.N; methodology: S.P.S, P.C., A.H, M.K., L.D.S, S.J., O.K., S.B., J.B., A.K., A.P., T.K., S.,R.; formal analysis: S.P.S., P.C. L.D.S, B.S., P.O.; S.E.E; resources: N.N, J.P.J.; writing (original draft): S.P.S, N.N.; supervision: J.P.J. and N.N.; funding acquisition: J.P.J. and N.N.

**Acknowledgements:**

Work in J.P.J.'s laboratory was funded by a European Research Council Starting Grant (ERC-StG 715361 SPACEVAR). We are grateful to M. Hammersmith for sharing *Tg(sst1:GFP-Ras)*. Work by S.P.S. is supported by the Fonds de la Recherche Scientifique-FNRS under Grant(s) n° 34772792 – SCHISM. N.N. receives funding from the Center for Regenerative Therapies Dresden at TU Dresden, the German Center for Diabetes Research (DZD), as well as research grants from the German Research Foundation (DFG), and the International Research Training Group (IRTG 2251): 'Immunological and Cellular Strategies in Metabolic Disease'. We are grateful to the following facilities at the CMCB and the CRTD: Deep Sequencing, Flow Cytometry Facility, Electron Microscopy, Light Microscopy, Zebrafish.

# References

Andersson, O., Adams, B.A., Yoo, D., Ellis, G.C., Gut, P., Anderson, R.M., German, M.S., Stainier, D.Y.R., 2012. Adenosine signaling promotes regeneration of pancreatic β cells in vivo. Cell Metab. 15, 885–894. https://doi.org/10.1016/j.cmet.2012.04.018

Benjannet, S., Rondeau, N., Day, R., Chrétien, M., Seidah, N.G., 1991. PC1 and PC2 are proprotein convertases capable of cleaving proopiomelanocortin at distinct pairs of basic residues. Proc. Natl. Acad. Sci. U. S. A. 88, 3564–3568. https://doi.org/10.1073/pnas.88.9.3564

Binot, A.-C., Manfroid, I., Flasse, L., Winandy, M., Motte, P., Martial, J.A., Peers, B., Voz, M.L., 2010. Nkx6.1 and nkx6.2 regulate alpha- and beta-cell formation in zebrafish by acting on pancreatic endocrine progenitor cells. Dev. Biol. 340, 397–407. https://doi.org/10.1016/j.ydbio.2010.01.025

Buenrostro, J.D., Giresi, P.G., Zaba, L.C., Chang, H.Y., Greenleaf, W.J., 2013. Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. Nat. Methods 10, 1213–8. https://doi.org/10.1038/nmeth.2688

Byrnes, L.E., Wong, D.M., Subramaniam, M., Meyer, N.P., Gilchrist, C.L., Knox, S.M., Tward, A.D., Ye, C.J., Sneddon, J.B., 2018. Lineage dynamics of murine pancreatic development at single-cell resolution. Nat. Commun. 9, 3922. https://doi.org/10.1038/s41467-018-06176-3

Carlemalm, E., Garavito, R.M., Villiger, W., 1982. Resin development for electron microscopy and an analysis of embedding at low temperature*. J. Microsc. 126, 123–143. https://doi.org/10.1111/j.1365-2818.1982.tb00362.x

Chera, S., Baronnier, D., Ghila, L., Cigliola, V., Jensen, J.N., Gu, G., Furuyama, K., Thorel, F., Gribble, F.M., Reimann, F., Herrera, P.L., 2014. Diabetes Recovery By Age-Dependent Conversion of Pancreatic δ-Cells Into Insulin Producers. Nature 514, 503–507. https://doi.org/10.1038/nature13633

Curado, S., Anderson, R.M., Jungblut, B., Mumm, J., Schroeter, E., Stainier, D.Y.R., 2007. Conditional targeted cell ablation in zebrafish: A new tool for regeneration studies. Dev. Dyn. 236, 1025–1035. https://doi.org/10.1002/dvdy.21100

Dalgin, G., Ward, A.B., Hao, L.T., Beattie, C.E., Nechiporuk, A., Prince, V.E., 2011. Zebrafish mnx1 controls cell fate choice in the developing endocrine pancreas. Development 138, 4597–4608. https://doi.org/10.1242/dev.067736

De Krijger, R.R., Aanstoot, H.J., Kranenburg, G., Reinhard, M., Visser, W.J., Bruining, G.J., 1992. The midgestational human fetal pancreas contains cells coexpressing islet hormones. Dev. Biol. 153, 368–75.

Delaspre, F., Beer, R.L., Rovira, M., Huang, W., Wang, G., Gee, S., Vitery, M. del C., Wheelan, S.J., Parsons, M.J., 2015. Centroacinar Cells Are Progenitors That Contribute to Endocrine Pancreas Regeneration. Diabetes 64, 3499–3509. https://doi.org/10.2337/db15-0153

Devos, N., Deflorian, G., Biemar, F., Bortolussi, M., Martial, J.A., Peers, B., Argenton, F., 2002. Differential expression of two somatostatin genes during zebrafish embryonic development. Mech. Dev. 115, 133–137. https://doi.org/10.1016/s0925-4773(02)00082-5

Fabig, G., Kretschmar, S., Weiche, S., Eberle, D., Ader, M., Kurth, T., 2012. Labeling of ultrathin resin sections for correlative light and electron microscopy. Methods Cell Biol. 111, 75–93. https://doi.org/10.1016/B978-0-12-416026-2.00005-4

Feng, J., Liu, T., Qin, B., Zhang, Y., Liu, X.S., 2012. Identifying ChIP-seq enrichment using MACS. Nat. Protoc. 7, 1728–1740. https://doi.org/10.1038/nprot.2012.101

Gannon, M., Ables, E.T., Crawford, L., Lowe, D., Offield, M.F., Magnuson, M.A., Wright, C.V.E., 2008. pdx-1 function is specifically required in embryonic β cells to generate appropriate numbers of endocrine cell types and maintain glucose homeostasis. Dev. Biol. 314, 406–417. https://doi.org/10.1016/j.ydbio.2007.10.038

Haghverdi, L., Büttner, M., Wolf, F.A., Buettner, F., Theis, F.J., 2016. Diffusion pseudotime robustly reconstructs lineage branching. Nat. Methods 13, 845–848. https://doi.org/10.1038/nmeth.3971

Hermann, M., Pirkebner, D., Draxl, A., Berger, P., Untergasser, G., Margreiter, R., Hengster, P., 2007. Dickkopf-3 is expressed in a subset of adult human pancreatic beta cells. Histochem. Cell Biol. 127, 513–521. https://doi.org/10.1007/s00418-007-0278-6

Janjuha, S., Singh, S.P., Tsakmaki, A., Mousavy Gharavy, S.N., Murawala, P., Konantz, J., Birke, S., Hodson, D.J., Rutter, G.A., Bewick, G.A., Ninov, N., 2018. Age-related islet inflammation marks the proliferative decline of pancreatic beta-cells in zebrafish. eLife 7, e32965. https://doi.org/10.7554/eLife.32965

Jiang, Z., Song, J., Qi, F., Xiao, A., An, X., Liu, N., Zhu, Z., Zhang, B., Lin, S., 2008. Exdpf is a key regulator of exocrine pancreas development controlled by retinoic acid and ptf1a in zebrafish. PLoS Biol. 6, e293. https://doi.org/10.1371/journal.pbio.0060293

Koh, P.W., Sinha, R., Barkal, A.A., Morganti, R.M., Chen, A., Weissman, I.L., Ang, L.T., Kundaje, A., Loh, K.M., 2016. An atlas of transcriptional, chromatin accessibility, and surface marker changes in human mesoderm development. Sci. Data 3, 160109. https://doi.org/10.1038/sdata.2016.109

Kragl, M., Knapp, D., Nacu, E., Khattak, S., Maden, M., Epperlein, H.H., Tanaka, E.M., 2009. Cells keep a memory of their tissue origin during axolotl limb regeneration. Nature 460, 60–65. https://doi.org/10.1038/nature08152

Li, N., Yang, Z., Li, Q., Yu, Z., Chen, X., Li, J.-C., Li, B., Ning, S.-L., Cui, M., Sun, J.-P., Yu, X., 2018. Ablation of somatostatin cells leads to impaired pancreatic islet function and neonatal death in rodents. Cell Death Dis. 9, 1–12. https://doi.org/10.1038/s41419-018-0741-4

Li, Z., Wen, C., Peng, J., Korzh, V., Gong, Z., 2009. Generation of living color transgenic zebrafish to trace somatostatin-expressing cells and endocrine pancreas organization. Differ. Res. Biol. Divers. 77, 128–134. https://doi.org/10.1016/j.diff.2008.09.014

Liu, Y., Lu, D., Zhang, Y., Li, S., Liu, X., Lin, H., 2010. The evolution of somatostatin in vertebrates. Gene 463, 21–28. https://doi.org/10.1016/j.gene.2010.04.016

Löhr, H., Hess, S., Pereira, M.M.A., Reinoß, P., Leibold, S., Schenkel, C., Wunderlich, C.M., Kloppenburg, P., Brüning, J.C., Hammerschmidt, M., 2018. Diet-Induced Growth Is Regulated via Acquired Leptin Resistance and Engages a Pomc-Somatostatin-Growth Hormone Circuit. Cell Rep. 23, 1728–1741. https://doi.org/10.1016/j.celrep.2018.04.018

Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. 15, 550. https://doi.org/10.1186/s13059-014-0550-8

Mameishvili, E., Serafimidis, I., Iwaszkiewicz, S., Lesche, M., Reinhardt, S., Bölicke, N., Büttner, M., Stellas, D., Papadimitropoulou, A., Szabolcs, M., Anastassiadis, K., Dahl, A., Theis, F., Efstratiadis, A., Gavalas, A., 2019. Aldh1b1 expression defines progenitor cells in the adult pancreas and is required for Kras-induced pancreatic cancer. Proc. Natl. Acad. Sci. 116, 20679–20688. https://doi.org/10.1073/pnas.1901075116

Merrell, A.J., Stanger, B.Z., 2016. Adult cell plasticity in vivo : de-differentiation and transdifferentiation are back in style. Nat. Rev. Mol. Cell Biol. 17, 413–425. https://doi.org/10.1038/nrm.2016.24

Moss, J.B., Koustubhan, P., Greenman, M., Parsons, M.J., Walter, I., Moss, L.G., 2009. Regeneration of the pancreas in adult zebrafish. Diabetes 58, 1844–1851. https://doi.org/10.2337/db08-0628

Ninov, N., Borius, M., Stainier, D.Y.R., 2012. Different levels of Notch signaling regulate quiescence, renewal and differentiation in pancreatic endocrine progenitors. Dev. Camb. Engl. 139, 1557–1567. https://doi.org/10.1242/dev.076000

Noguchi, G.M., Huising, M.O., 2019. Integrating the inputs that shape pancreatic islet hormone release. Nat. Metab. 1, 1189–1201. https://doi.org/10.1038/s42255-019-0148-2

Pan, F.C., Brissova, M., Powers, A.C., Pfaff, S., Wright, C.V.E., 2015. Inactivating the permanent neonatal diabetes gene Mnx1 switches insulin-producing β-cells to a δ-like fate and reveals a facultative proliferative capacity in aged β-cells. Dev. Camb. Engl. 142, 3637–48. https://doi.org/10.1242/dev.126011

Pisharath, H., Rhee, J.M., Swanson, M.A., Leach, S.D., Parsons, M.J., 2007. Targeted ablation of beta cells in the embryonic zebrafish pancreas using E. coli nitroreductase. Mech. Dev. 124, 218–229. https://doi.org/10.1016/j.mod.2006.11.005

Robinson, J.T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G., Mesirov, J.P., 2011. Integrative genomics viewer. Nat. Biotechnol. 29, 24–26. https://doi.org/10.1038/nbt.1754

Salem, V., Silva, L.D., Suba, K., Georgiadou, E., Neda Mousavy Gharavy, S., Akhtar, N., Martin-Alonso, A., Gaboriau, D.C.A., Rothery, S.M., Stylianides, T., Carrat, G., Pullen, T.J., Singh, S.P., Hodson, D.J., Leclerc, I., Shapiro, A.M.J., Marchetti, P., Briant, L.J.B., Distaso, W., Ninov, N., Rutter, G.A., 2019. Leader β-cells coordinate Ca 2+ dynamics across pancreatic islets in vivo. Nat. Metab. 1, 615–629. https://doi.org/10.1038/s42255-019-0075-2

Satija, R., Farrell, J.A., Gennert, D., Schier, A.F., Regev, A., 2015. Spatial reconstruction of single-cell gene expression data. Nat. Biotechnol. 33, 495–502. https://doi.org/10.1038/nbt.3192

Serafimidis, I., Rodriguez-Aznar, E., Lesche, M., Yoshioka, K., Takuwa, Y., Dahl, A., Pan, D., Gavalas, A., 2017. Pancreas lineage allocation and specification are regulated by sphingosine-1-phosphate signalling. PLOS Biol. 15, e2000949. https://doi.org/10.1371/journal.pbio.2000949

Setty, M., Tadmor, M.D., Reich-Zeliger, S., Angel, O., Salame, T.M., Kathail, P., Choi, K., Bendall, S., Friedman, N., Pe'er, D., 2016. Wishbone identifies bifurcating developmental trajectories from single-cell data. Nat. Biotechnol. 34, 637–645. https://doi.org/10.1038/nbt.3569

Seymour, P.A., Shih, H.P., Patel, N.A., Freude, K.K., Xie, R., Lim, C.J., Sander, M., 2012. A Sox9/Fgf feed-forward loop maintains pancreatic organ identity. Dev. Camb. Engl. 139, 3363–3372. https://doi.org/10.1242/dev.078733

Sharon, N., Vanderhooft, J., Straubhaar, J., Mueller, J., Chawla, R., Zhou, Q., Engquist, E.N., Trapnell, C., Gifford, D.K., Melton, D.A., 2019. Wnt Signaling Separates the Progenitor and Endocrine Compartments during Pancreas Development. Cell Rep. 27, 2281-2291.e5. https://doi.org/10.1016/j.celrep.2019.04.083

Singh, S.P., Janjuha, S., Chaudhuri, S., Reinhardt, S., Kränkel, A., Dietz, S., Eugster, A., Bilgin, H., Korkmaz, S., Zararsız, G., Ninov, N., Reid, J.E., 2018. Machine learning based classification of cells into chronological stages using single-cell transcriptomics. Sci. Rep. 8, 17156. https://doi.org/10.1038/s41598-018-35218-5

Singh, S.P., Janjuha, S., Hartmann, T., Kayisoglu, Ö., Konantz, J., Birke, S., Murawala, P., Alfar, E.A., Murata, K., Eugster, A., Tsuji, N., Morrissey, E.R., Brand, M., Ninov, N., 2017. Different developmental histories of beta-cells generate functional and proliferative heterogeneity during islet growth. Nat. Commun. 8, 664. https://doi.org/10.1038/s41467-017-00461-3

Solimena, M., Schulte, A.M., Marselli, L., Ehehalt, F., Richter, D., Kleeberg, M., Mziaut, H., Knoch, K.-P., Parnis, J., Bugliani, M., Siddiq, A., Jörns, A., Burdet, F., Liechti, R., Suleiman, M., Margerie, D., Syed, F., Distler, M., Grützmann, R., Petretto, E., Moreno-Moral, A., Wegbrod, C., Sönmez, A., Pfriem, K., Friedrich, A., Meinel, J.,

Wollheim, C.B., Baretton, G.B., Scharfmann, R., Nogoceke, E., Bonifacio, E., Sturm, D., Meyer-Puttlitz, B., Boggi, U., Saeger, H.-D., Filipponi, F., Lesche, M., Meda, P., Dahl, A., Wigger, L., Xenarios, I., Falchi, M., Thorens, B., Weitz, J., Bokvist, K., Lenzen, S., Rutter, G.A., Froguel, P., von Bülow, M., Ibberson, M., Marchetti, P., 2018. Systems biology of the IMIDIA biobank from organ donors and pancreatectomised patients defines a novel transcriptomic signature of islets from individuals with type 2 diabetes. Diabetologia 61, 641–657. https://doi.org/10.1007/s00125-017-4500-3

Spanjaard, B., Hu, B., Mitic, N., Olivares-Chauvet, P., Janjuha, S., Ninov, N., Junker, J.P., 2018. Simultaneous lineage tracing and cell-type identification using CRISPR-Cas9-induced genetic scars. Nat. Biotechnol. 36, 469–473. https://doi.org/10.1038/nbt.4124

Teitelman, G., Alpert, S., Polak, J.M., Martinez, A., Hanahan, D., 1993. Precursor cells of mouse endocrine pancreas coexpress insulin, glucagon and the neuronal proteins tyrosine hydroxylase and neuropeptide Y, but not pancreatic polypeptide. Dev. Camb. Engl. 118, 1031–9.

Thorel, F., Népote, V., Avril, I., Kohno, K., Desgraz, R., Chera, S., Herrera, P.L., 2010. Conversion of Adult Pancreatic α-cells to β-cells After Extreme β-cell Loss. Nature 464, 1149–1154. https://doi.org/10.1038/nature08894

Trapnell, C., Cacchiarelli, D., Grimsby, J., Pokharel, P., Li, S., Morse, M., Lennon, N.J., Livak, K.J., Mikkelsen, T.S., Rinn, J.L., 2014. The dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. Nat. Biotechnol. 32, 381–386. https://doi.org/10.1038/nbt.2859

Untergasser, G., Martowicz, A., Hermann, M., Töchterle, S., Meyer, D., 2011. Distinct expression patterns of dickkopf genes during late embryonic development of Danio rerio. Gene Expr. Patterns GEP 11, 491–500. https://doi.org/10.1016/j.gep.2011.08.005

Wan, H., Korzh, S., Li, Z., Mudumana, S.P., Korzh, V., Jiang, Y.-J., Lin, S., Gong, Z., 2006. Analyses of pancreas development by generation of gfp transgenic zebrafish using an exocrine pancreas-specific elastaseA gene promoter. Exp. Cell Res. 312, 1526–1539. https://doi.org/10.1016/j.yexcr.2006.01.016

Zhang, J., McKenna, L.B., Bogue, C.W., Kaestner, K.H., 2014. The diabetes gene Hhex maintains δ-cell differentiation and islet function. Genes Dev. 28, 829–34. https://doi.org/10.1101/gad.235499.113

Ziegenhain, C., Vieth, B., Parekh, S., Reinius, B., Guillaumet-Adkins, A., Smets, M., Leonhardt, H., Heyn, H., Hellmann, I., Enard, W., 2017. Comparative Analysis of Single-Cell RNA Sequencing Methods. Mol. Cell 65, 631-643.e4. https://doi.org/10.1016/j.molcel.2017.01.023

**Figure Legends:**

**Figure 1. Single-cell RNA-sequencing of the zebrafish pancreas reveals two types of islet δ-cells. (A)** Schematic of the high-throughput 10x Droplet-Seq. approach to classify the cell populations in the zebrafish pancreas. **(B)** t-SNE plot

representation of 1,969 pancreatic cells with clusters representing distinct cell-types.
**(C)** t-SNE plot representation of 1,530 endocrine cells with clusters based on the expression of hormonal genes.  **(D)** Violin Plot for expression of hormonal genes in the endocrine clusters.  **(E)** t-SNE plot representation of 746 insulin-expressing pancreatic cells with specific gene expression overlaid as a heat-map.  Cells with co-expression of *insulin* and *sst1.1* are outline

**Figure 2. Markers of δ1- and δ2- cells.  (A)** t-sne plot representation of the β-, δ1-  and δ2-cell populations, expressing *ins, sst1.1* and *sst2*, respectively. glucokinase (*gck*) is expressed in both δ1- and δ2- cells, whereas *dkk3b* and *pdx1* are enriched in the δ1-cells.  **(B)** Violin plot highlighting the expression levels of *ins*, *sst1.1*, *sst2*, *dkk3b*, *pdx1* and *gck* in the δ1- and δ2- cell clusters.

**Figure 3.  Characterization of a small population of glucose-responsive β/δ hybrid cells.  (A)** Schematic of the *TgBAC(sst1.1:EGFP-Ras)* reporter line used to mark specifically the *sst1.1* population, where a BAC containing the *sst1.1* promoter drives membrane GFP expression.  **(B)** Confocal projection of whole-mounted islets from the *TgBAC(sst1.1: EGFP-Ras)* reporter line stained with an anti-insulin antibody.  The dashed line indicates a GFP and insulin co-expressing cell. The high-magnification images in B' and B'' shows a single confocal plane of the same cell co-expressing GFP and insulin.  **(C-D)** Thin pancreatic sections from the *TgBAC(sst1.1: EGFP-Ras)* reporter line at 7 months post-fertilization (mpf) and 16 mpf.  The sections were stained with insulin (red) and somatostatin (magenta) antibodies.  Arrows mark EGFP+ cells co-stained with insulin and somatostatin antibody.  The white box indicates the region displayed at higher magnification in C, revealing cells expressing each protein along with GFP.  **(E-E')** Two-Photon in-vivo Ca$^{2+}$ imaging in 5 dpf double-transgenic zebrafish larvae of the genotype *Tg (sst1.1:*

*EGFP-Ras); Tg(ins: K-GECO1)* showing the glucose-stimulated response of mono-hormonal and bi-hormonal cells.  **(E'')** Normalized *K-GECO* fluorescent traces, showing glucose-stimulated calcium influx of bi-hormonal and mono-hormonal β-cells.

**Figure 4.  Molecular analysis of β/δ cells reveals a progenitor signature.**

**(A)** FACS plot of cells from *Tg(ins:BB1.0L)*; *Tg(sst1.1: EGFP-Ras)* adult fish.  Cells were marked based on the presence of red and green fluorescence, and index-sorted into 96-well plates.  β-cells are RFP+ (R), δ1-cells are GFP+ (G) and bi-hormonal cells are green+ and red+ (GR).  **(B)** A t-sne plot of the profiled cells with colors corresponding to their respective location on the FACS plot.  The cells are segregated into three groups: R (β-cells), G (δ1-cells), and GR (bi-hormonal cells).  **(C)** Violin Plot showing *insulin* and *sst1.1* expression in the three cell populations.  The Y-axis represents normalized gene expression.  **(D)** A dot-plot displaying the Pearson correlation coefficient between *sst1.1* and *dkk3b*, as well as, between *sst1.1* and *pyyb* expression levels.  **(E)** Violin Plot showing the expression levels of control genes (*dkk3b*, *pyyb*, *EGFP*) and selected genes related to β-cell differentiation and function in the three populations.  *Mnx1* and *nkx6.2* are enriched in β-cells, whereas *HHEX* is expressed in δ1-cells.  The Y-axis represents normalized gene expression.  **(F)** Pseudo-temporal ordering of the cells, with (F') showing the expression trend of selected genes that show significant correlation with the pseudo-time.  δ1-cells show enrichment of genes involved in the Wnt, FGF, Notch, and sphingolipid signaling pathways compared to β-cells.  The bi-hormonal cells exhibit intermediate expression of genes involved in β- and δ-cell fate-specification and function compared to the pure cell populations.  **(G)** To validate the single-cell RNA-Seq. results, the three endocrine cell populations were profiled using bulk-RNA

sequencing.  The results from the experiment are depicted as a heatmap demonstrating the expression levels of selected genes including those involved in Wnt, FGF and Notch and sphingolipid signaling.  All depicted genes show a statistically significant difference in expression between the δ1 and β-cells.

**Figure 5. β/δ cells constitute a prominent source of insulin expression during diabetes recovery in adult zebrafish.  (A-B)** UMAP-plot representation of integrated 10X scRNASeq data from all β-cell regeneration time-points: 0, 2, 7 and 14 days post-ablation (dpa) showing distinct clusters for δ1, δ2, alpha, acinar and duct cells.  **(C)** Feature-plot highlighting the expression of *sst1.1 (δ1), sst2 (δ2), gcgb (alpha), cftr (duct)* and *ela2 (acinar)* genes.  **(D)** Feature-plot highlighting *ins* expression and percentage of *ins*-positive cells in different pancreatic cells during regeneration. **(D)** Feature-plot highlighting *ins* expression in different pancreatic cells during regeneration  **(E)** VInplot highlighting *ins* expression in different pancreatic cells during regeneration. The hybrid cells from the δ1-cell-cluster serve as the main source of insulin-expression during the course of regeneration.

**Figure 6. Gene-correlation analysis indicates resolution of the β/δ cells towards *ins+* and *sst1.1+* populations during regeneration**
(**A**, **D**) UMAP displaying selected cells from 7 dpa (A) and 14 dpa (F).  (**B-E**) UMAP overlaid with gene expression plots for *sst1.1* and *ins* at 7 dpa (B) and 14 dpa (E). (**C**,**F**) Expression trends for *sst1.1* and *ins* at 7 dpa (C) and 14 dpa (F).  At 14 dpa, individual cells along display a negative correlation between *sst1.1* and *ins* expression, whereas at 7 dpa the cells co-express both hormones.

**Figure 7.  Live imaging of regeneration reveals that the β/δ cells arise *de novo* and acquire glucose-responsiveness.  (A)** To visualize the dynamics of bi-hormonal cells during β-cell regeneration, *Tg (ins: FLAG-NTR)*; *TgBAC (sst1.1:*

*EGFP- Ras), Tg(ins: BB1.0L)*-triple transgenic animals were treated with vehicle or Mtz for 24 hours. Following Mtz-mediated ablation of β-cells, a time-course live imaging was performed from 0 to 4 dpa. **(A)** 3D-projections of the zebrafish islet in control and at different time-points post β-cell ablation in the same larva. Following MTZ addition, all tdTomato-expressing cells were ablated and no longer present in the islet. New tdTomato- and GFP-positive cell appear in the islet over time, indicating a *de novo* origin (arrows). **(A')** Orientation of the zebrafish islet during the live-imaging process. **(A'')** Quantification of the proportion of bi-hormonal cells during the regeneration process (Brown-Forsythe and Welch ANOVA). **(B, C)** Images from the time-lapse recording of glucose-stimulated calcium influx (6 frames/sec, single plane) of the zebrafish islet in control (Unablated) and at 4-days post β-cell ablation (Ablated). **(B', C')** Normalized *K-GECO* fluorescent traces, showing glucose-stimulated calcium influx of bi-hormonal and mono-hormonal cells. **(D, D')** Quantification of changes in normalized *K-GECO* fluorescence intensity (recorded over 400 frames after glucose-injection) based on area under the curve (AUC) for single-cells. **(D)** Scatterplot for *K-GECO*+/GFP- and *K-GECO*+/GFP+ from unablated samples (Mann-Whitney unpaired t-test, p-value = 0.0907, non-significant). **(D')** Scatterplot for *K-GECO*+/GFP- and *K-GECO*+/GFP+ from ablated samples (Mann-Whitney unpaired t-test, p-value = 0.222, non-significant)

**Figure 8. Over-expression of *dkk3b* leads to increased formation of β/δ cells. (A)** Single-plane confocal images of islets from *Tg(ins: mCerulean_dkk3b)*; *TgBAC (sst1.1: EGFP- Ras)*-double transgenic animals (5 dpf) stained with insulin antibody (red). The yellow dashed-box outlines the *sst1.1-insulin* double-positive cells. **(A')** Scatter plot showing the number of bi-hormonal cells in wild-type and *ins:mCerulean-t2a-dkk3b* animals (Mann-Whitney unpaired t-test, p-value = 0.0005,

significant).  **(B)** Single-plane confocal images of islets from *Tg(hsp70l:mCherry-t2a-dkk3b)*; *TgBAC (sst1.1: EGFP- Ras)*-double transgenic animals stained with an insulin antibody (magenta).  The larvae were given a heat-shock for 2 hours at 37°C at 3 and 4dpf, thereby inducing the expression of *dkk3b*.  The yellow dashed-box outlines the *sst1.1-insulin* double positive cells.  **(B')** Scatter plot showing the number of bi-hormonal cells in wild-type and *Tg(hsp70l:mCherry-t2a-dkk3b)* larvae (Mann-Whitney unpaired t-test, p-value = 0.0352, significant).  **(C)** Cartoon model based on the experimental observations in this study.

# Figure 1: Single-cell RNA-Sequencing of the zebrafish pancreas reveals two types of islet δ-cells.

**Figure 2: Markers of δ1- and δ2- cells**

# Figure 3: Characterization of a small population of glucose-responsive hybrid β/δ cells.

# Figure 4: Molecular analysis of β/δ cells reveals a progenitor signature.

**Figure 5. β/δ cells constitute a prominent source of insulin expression during diabetes recovery in adult zebrafish.**

# Figure 6. Gene-correlation analysis indicates resolution of the β/δ-cells towards *ins*+ and *sst1.1*+ populations during regeneration

## Figure 7. Live imaging of regeneration reveals that β/δ-cells arise *de novo* and acquire glucose-responsiveness.

# Figure 8. Over-expression of *dkk3b* leads to increased formation of β/δ cells

**Supplementary figures:**

**A**

**Supplementary figure S1: Markers for duct cells (A)**. t-SNE plot representation of duct cell population expressing *her15.1, cftr, epcam and anxa4*

**A**

sst1.1

**B**

**Supplementary figure S2. Persistence of bi-hormonal cells during various developmental stages of zebrafish and ATAC seq analysis of chromatin in β-cells. (A)** Boxplot depicting the percentage of β-cells expressing *sst1.1* in a previously published single-cell smart2 RNA-Seq. dataset. The dataset contains single-cell RNA-Seq. profiles of β-cells isolated from the *Tg(ins:BB1.0L)* transgenic line using FACS at different stages of the zebrafish juvenile and adult life. **(B)** Plots for selected genes depicting the open genomic regions (in orange) identified using ATAC-Seq. specifically for β-cells. Regions proximal to the promoters of *ins, sst1.1* and *dkk3b* genes display open chromatin.

**Supplementary figure S3. The proportion of hybrid cells increase during β-cell regeneration. (A)** To assess the dynamics of hybrid cells during β-cell regeneration, *Tg(ins: FLAG-NTR)*; *TgBAC (sst1.1: EGFP-Ras)* double transgenic animals were treated with Mtz at 4 dpf for 24 hours. Following Mtz-based ablation of NTR-expressing β-cells, the samples were collected at 4- and 8-days post-ablation (dpa) for quantification. **(B)** Single confocal stack of islets at 4 dpa. Arrows mark bi-hormonal cells, which express EGFP and are also marked with the Insulin antibody. **(C)** Barplots displaying the percentage of bi-hormonal cells among all insulin+ cells in the islet at 4 and 8 dpa. n = 6 animals for each condition. Comparison based on two-tailed Heteroscedastic t-test. p-value: * < 0.05; *** < 0.001.

**Supplementary figure S4. An efficient NTR system highlighting β-cell ablation and regeneration in the adult zebrafish pancreas. (A)** Confocal projection of a pancreatic islet from a 3-month-old *Tg (ins:YFP-NTR3)* transgenic zebrafish showing β-cell destruction and regeneration following Mtz-treatment and wash-out. **(B)** Fasting blood glucose level measurements (mmol/L) show a return to normoglycemic levels by 14-days post β-cell ablation.

**Supplementary figure S5. δ1- but not δ2-cells express insulin upon β-cell loss in adults. (A-E)** UMAP plot representation of adult zebrafish pancreatic cells showing expression of *ins, sst1.1*, and *sst2* at various β-cell regeneration time-points: Control, 0 dpa, 2 dpa, 7 dpa, and 14 dpa.

**Supplementary figure S6: Time-course of insulin expression in different cell types during β-cell regeneration. (A-E)** The 10X regeneration datasets were mined to evaluate the proportions of ins-positive cells in the cell clusters expressing δ1, δ2, α, acinar and ductal markers. Based on the insulin expression level, cells were categorized into low, medium and high ins-expressing cells. The bar-plots highlighting *ins* expression profile across different β-cell regeneration time-points and cell types (control, 0, 2, 7 and 14 dpa respectively). Insulin expression increases (from low to high) in δ1-cells during β-cell regeneration, whereas other cell types retain low ins expression across time.

**Supplementary figure S7. Average expression profile of genes in the δ1-cluster during β-cell regeneration. (A)** The 10X regeneration dataset was mined to determine the average expression profile of various β-cell specific genes and progenitor markers in the δ1-cluster. The graph shows the change in the average expression of specific genes (*ins, sst1.1, pdx1, ppdpfa, ppdpfb, pcsk1, pcsk2, dkk3b, pyyb,* and *wif1*) during β-cell regeneration. The fold-change values on the Y-Axis have been normalized to the average expression profile of the control sample.

**Supplementary figure S8. The δ1-cell population expands following β-cell ablation.** Graph showing the increase of *sst1.1*-expressing cells relative to α-cells after β cell ablation with 99% confidence intervals.

**Supplementary figure S9.  Accumulation of vacuolar structures in the δ1-cells upon β-cell loss.  (A)** 3D-confocal projection of control and β-cell-ablated samples in *Tg (ins: FLAG-NTR)*; *TgBAC (sst1.1: EGFP Ras), Tg(ins: BB1.0L)*-triple transgenic animals.  Upon β-cell ablation, there is emergence of prominent vacuolar structures in the GFP-positive cells.  **(A')** Scatter-plot showing the total number of vacuolar structures in the δ1-cells post β-cell loss in the zebrafish larvae. **(B)**  CLEM (Correlative Light and Electron Microscopy) section obtained from the pancreatic islet larvae following β-cell ablation.  The animals are *Tg (ins: FLAG-NTR)*; *TgBAC (sst1.1:EGFP-Ras), Tg(ins:BB1.0L)*-triple transgenic.  The animals were treated with vehicle or Mtz from 2.5 to 3dpf and were fixed at 5dpf.  During the CLEM protocol, the sections were labelled with immuno-gold particles labelling the GFP protein (indicated by the yellow arrow).  The section highlights the appearance of vacuolar structures (highlighted with the red-box) in δ1-cells at 2-days post β-cell ablation.