Superior colliculus saccade motor bursts do not dictate movement kinematics

Tong Zhang[1,2], Tatiana Malevich[1,2], Matthias Baumann[1,2], and Ziad M. Hafed[1, 2]

[1] Werner Reichardt Centre for Integrative Neuroscience, Tübingen University,
72076, Tübingen, Germany

[2] Hertie Institute for Clinical Brain Research, Tübingen University,
72076, Tübingen, Germany

Correspondence

Ziad M. Hafed
Werner Reichardt Centre for Integrative Neuroscience
Tübingen University
Otfried-Müller Str. 25
Tübingen, 72076, Germany
Tel: +49 7071 29 88819
ziad.m.hafed@cin.uni-tuebingen.de

Abstract

The primate superior colliculus (SC) contains a topographic map of space, such that the anatomical location of active neurons defines a desired eye movement vector. Complementing such a spatial code, SC neurons also exhibit saccade-related bursts that are tightly synchronized with movement onset. Current models suggest that such bursts constitute a rate code dictating movement kinematics. Here, using two complementary approaches, we demonstrate a dissociation between the SC rate code and saccade kinematics. First, we show that SC burst strength systematically varies depending on whether saccades of the same amplitude are directed towards the upper or lower visual fields, but the movements themselves have similar kinematics. Second, we show that for the same saccade vector, when saccades are significantly slowed down by the absence of a visible saccade target, SC saccade-related burst strengths can be elevated rather than diminished. Thus, SC saccade-related motor bursts do not necessarily dictate movement kinematics.

Introduction

The superior colliculus (SC) plays an important role in saccade generation, as evidenced by the ease with which low-current electrical microstimulation of SC neurons evokes saccades[1,2]. Anatomically, SC neurons are organized to form a spatial code of eye movement displacement vectors[1,3,4], such that the location of an active neuron in the SC defines the amplitude and direction of a desired saccade. Robustness and accuracy of saccade vector representation are ensured through population coding[4-6], with the aggregate activity of a large number of simultaneously active neurons defining a given movement's metrics.

The SC spatial code necessarily entails a temporal synchrony of SC activity at the time of saccades. Indeed, saccade-related neurons show a highly characteristic temporal evolution of spiking[7-11], dominated by a burst tightly locked to movement onset. Interestingly, the strength of such a burst can vary, suggesting that SC neurons may encode additional properties beyond the saccadic displacement vector represented by the spatial code. For example, blink-perturbed saccades can have weaker, but prolonged, bursts[12,13]. Moreover, burst evolution during a saccade may be related to the remaining motor error of an ongoing eye movement[14,15] (i.e. how much more the eye needs to keep moving), or it may be related to the speed profile of the ensuing saccade[16]. Additionally, saccade-related burst strength can be modulated by audio-visual sensory combinations[17]. Thus, there is an SC rate code for saccades, the role of which is less well understood than that of the spatial code.

The most recent SC models posit an important role for the rate code in dictating saccade kinematics[13,16,18]. In these models, the locus of an active neuron (i.e. the spatial code) defines how each individual spike in a motor burst moves the eye along the amplitude dimension; parameters like eye speed or time to movement end would reflect the strength of (i.e. number of spikes in) the motor burst (i.e. the rate code). While appealing in their combination of both spatial and rate codes for movement specification, these models suggest a very tight relationship between saccade-related burst strength and movement kinematics. However, this may not necessarily always be the case. For example, we recently explored a situation in which saccade kinematics were altered by a simultaneity condition between a motor burst somewhere on the SC map and an irrelevant visual burst somewhere else[19,20]. When we recorded at both the motor and visual burst locations[21], we found a lawful relationship between the saccade changes and the number of additional spikes injected by the visual burst (consistent with the spatial code); however, critically, the simultaneous motor burst was minimally affected[21]. Thus, the rate code of the original movement commands was essentially unaltered even though the movements themselves were. This, along with other evidence[22], motivates investigating whether saccade kinematics are indeed dictated by the SC rate code or not.

We approached this question using two complementary approaches. In the first, we exploited a large asymmetry in how the SC represents the upper versus lower visual fields in its visual sensitivity[23]. If such an asymmetry still holds, but now for saccade-related motor burst strength, then there should be (at least according to current models of the rate code) systematic differences in the (amplitude-matched) saccades' kinematics. We confirmed a neural asymmetry in SC motor burst strengths, but found no concomitant kinematic differences between amplitude-matched saccades towards the upper and lower visual fields. In the second approach, we instead used vector-matched saccades, but of clearly different kinematics. Specifically, we exploited the fact that saccades to a blank can have significantly slower speeds than saccades to a clear, punctate visual target[24-30]. We, therefore, compared SC neuron motor bursts in these two conditions, sometimes recording multiple neurons simultaneously in the two behavioral contexts. Surprisingly, we found no correlation between SC motor burst modifications and the kinematic alterations of the saccades. More importantly, approximately one quarter of the neurons actually increased, rather than decreased, their motor burst strengths for the slower saccades. Our observations highlight the need to explore other potential functional roles for the saccade-related SC rate code.

Results

We first identified a dissociation between SC motor burst strengths and their associated eye movements' kinematics. Specifically, we explored how SC motor burst strength might differ as a function of visual field location. When we recently described an asymmetry in how the SC represents the upper and lower visual fields[23], we found that SC visual response properties were different across the fields. We also briefly mentioned that the strength of saccade-related motor bursts may also be asymmetric[23]. Here, we investigated the robustness of this saccade-related neural asymmetry in more detail, and we then asked whether it predicted an asymmetry in amplitude-matched saccade kinematics between movements towards the upper and lower visual fields. That is, we were motivated by a common assumption in existing models[13,16,18] that the relationship between SC motor burst strengths and movement kinematics is directionally symmetric and only depends on movement amplitudes. If so, then amplitude-matched movements of different directions, which have different SC motor burst strengths, should also have different kinematics. In a second set of experiments, we then used the complementary approach: we compared vector-matched saccades (where both the amplitude and direction were the same and towards the response field hotspot), and we asked whether alterations in these saccades' kinematics under different behavioral contexts were systematically related to alterations in the SC motor burst strengths.

In what follows, we first describe the amplitude-matched upper and lower visual field saccade results, and we then turn to the experiments with the vector-matched saccades having different kinematics.

Difference in superior colliculus motor burst strengths for saccades towards the upper versus lower visual fields

Visual sensitivity is significantly stronger in SC neurons representing the upper visual field[23]. That is, if a neuron's visual response field has a preferred (hotspot) location above the retinotopic horizontal meridian and we present a stimulus at this location, then the neuron's response is stronger than that of a neuron with a lower visual field hotspot location (and a stimulus presented at its preferred location). Curiously, in our earlier study[23], we noticed that saccade-related motor bursts showed the opposite asymmetry: saccade-related motor bursts (for preferred hotspot locations) were stronger for neurons representing saccades towards the lower visual field than for neurons representing saccades towards the upper visual field. However, in that study[23], we did not control for the depths of the recorded neurons from the SC surface when we analyzed the neurons' motor bursts. Since the strength of SC motor bursts can vary with depth from the SC surface[6], here, we wanted to first confirm whether the asymmetry alluded to above[23] was still present when carefully controlling for neuron depth (Methods). If this was the case, we could then ask whether saccade kinematics were systematically different or not.

We re-analyzed the neural database of ref. [23] (monkeys P and N) by first matching the depths of neurons from the SC surface between the upper and lower visual fields (Methods). For each extra-foveal neuron in this database (here referred to as dataset 1), we identified

whether the neuron was saccade-related or not[23]. We then classified the saccade-related
neurons as having a movement-related preferred response field location (or hotspot
location) in the upper or lower visual field. Preference was defined as the location for which
saccades were associated with the highest firing rates, similarly to how visual preference
was defined as the location for which visual stimuli evoked the strongest visual bursts[23].
Finally, we picked a range of neural depths from the SC surface that was overlapping
between the upper and lower visual field neurons. This final step was the critical step for the
present analysis, and it resulted in us having a total of 136 SC neurons with depths from the
SC surface between 600 and 1850 μm (Methods). The distribution of these depths is
represented in Fig. 1a, where each neuron's depth is plotted against a measure of whether
the neurons' preferred movement-related response field location was above or below the
horizontal meridian (the x-axis shows the angular direction of the preferred location from
the horizontal meridian; positive means above the meridian, and negative means below). As
can be seen, we found saccade-related SC activity at a range of depths from the SC surface
that was consistent with prior observations[6-10,31,32]. Critically, the neural depths were
overlapping between the upper and lower visual field neurons (Fig. 1a; p = 0.1257, t-test, t-
statistic: -1.5408, df: 134); note (as an aside) that upper visual field directions were
compressed relative to lower visual field directions, which is consistent with the idea of
upper visual field neural tissue magnification in the SC[23]. Therefore, we were now in a
position to check whether an asymmetry of saccade-related burst strengths alluded to
earlier[23] still held after controlling for neuron depth.
165
Having established that we now had a neural database with matched depths from the SC
surface, we proceeded to comparing motor burst strengths between the upper and lower
visual field neurons. We plotted the peri-saccadic firing rates of the neurons of Fig. 1a, from
a delayed, visually-guided saccade task (Methods). We employed such a delayed saccade
paradigm to allow analyzing motor bursts in isolation, without the recently occurring visual
bursts associated with target onset, which would have come too close to saccade onset in an
immediate, visually-guided saccade version of the task (Methods). We picked, for each
neuron, the preferred saccades of the neuron and plotted its firing rate for these
movements, as we did previously[23]. We then averaged across all neurons (Fig. 1b). There
was indeed an asymmetry in SC motor burst strengths, such that neurons representing the
lower visual field had significantly stronger motor bursts than neurons representing the
upper visual field (Fig. 1b). To statistically assess the difference in burst strengths after
matching for neural depths from the SC surface, we measured the average firing rate in the
final 50 ms before saccade onset for each neuron's preferred saccades[23] (Methods). We then
compared the population of measurements for the upper and lower visual field neurons of
Fig. 1a using a t-test. Across neurons, average firing rate for the upper visual field neurons
was 99 spikes/s, and it was 121 spikes/s for the lower visual field neurons. This difference
was statistically significant (p=0.039, t-test, t-statistic: -2.0844, df: 134). Therefore, even
after controlling for the depths of neurons from the SC surface, we confirmed a potential
asymmetry in saccade-related burst strength between upper and lower visual field
saccades[23].
187
188
189

Figure 1 Superior colliculus (SC) saccade-related motor bursts are stronger for downward saccades than for upward saccades, even when controlling for depth from the SC surface. (a) We picked neurons matched for depth from the SC surface (between 600 and 1850 μm) but having movement field hotspot locations in either the upper (light blue) or lower (light red) visual fields (i.e. positive or negative directions from the horizontal meridian, respectively). Note that the upper visual field neurons appear compressed along the direction dimension (i.e. in visual coordinates), likely due to upper visual field neural tissue magnification[23]. Such magnification is similar, in principle, to foveal magnification of SC neural tissue[33]. **(b)** For the neurons in **a**, we plotted peri-saccadic firing rates for saccades towards each neuron's preferred movement field location[23]. Saccade-related bursts were stronger for lower visual field than upper visual field neurons. **(c)** To better constrain errors in depth estimates from the SC surface (due to surface curvature), we further restricted the choice of neurons to those primarily near the horizontal meridian and at an eccentricity range associated with quasi-constant tissue curvature between upper and lower visual field locations[33]; the ranges of amplitudes, directions, and depths are shown in the inset. The motor bursts of the resulting 31 neurons were still stronger for lower visual field than upper visual field neurons (left panel). Error bars in all panels denote s.e.m.

A potential concern related to the above interpretation might be the curvature associated with the SC's 3-dimensional shape. Since all electrode paths were constant and defined by the recording chamber's orientation (Methods), it could still be possible that lateral recording sites (representing the lower visual field) could have had systematically different depths from the SC surface than medial recording sites (representing the upper visual field), by virtue of the different SC surface curvature at the two groups of sites. We therefore decided to analyze a stricter grouping of SC neurons. We picked a smaller range of eccentricities (5-15 deg), directions from the horizontal meridian (<30 deg), and depths from the SC surface (1100-1900 μm) for comparing upper and lower visual field neurons' motor bursts. Our prior work on SC surface topography and 3-dimensional anatomical shape[33] suggested that this range of selection should reduce potential systematic differences in estimates of depths from the SC surface between the upper and lower visual field groups of neurons. We found 31 neurons (20 upper visual field and 11 lower visual field) satisfying the above strict criteria. When we analyzed their saccade-related firing rates, we still found a similar asymmetry between upper and lower visual field locations (Fig. 1c). The average firing rate (in our same measurement interval) for the upper visual field neurons was 92 spikes/s, and it was 147 spikes/s for the lower visual field neurons; this difference was, again, statistically significant (p=0.0162, t-test, t-statistic: -2.5524, df: 29). Therefore, it is likely, given both analyses in Fig. 1, that there is indeed a systematic asymmetry in SC motor burst strength between saccades towards the upper and lower visual fields. We were now in a position to ask whether such an asymmetry was reflected in saccade kinematics, as might be predicted from some recent as well as classic models of the SC rate code[13,14,16].

232   *Similarity of movement kinematics for saccades towards the upper and lower visual fields*
233
According to existing models, which assume directional symmetry in the SC movement
commands, amplitude-matched saccades to the upper and lower visual fields should have
different movement kinematics given the different SC motor burst strengths that we saw in
Fig. 1. To test this, we analyzed saccades from both monkeys (P and N) from the same
delayed, visually-guided saccade task that was used to analyze the peri-saccadic SC firing
rates above (Methods). To compare size- and direction-matched movements, we picked, in
each monkey, 5 saccade sizes (3, 5, 7, 10, and 13 deg radial amplitude), and two example
directions from the horizontal meridian (+45 and -45 deg; i.e. oblique saccades; note that we
also made similar observations for example directions that were nearer to or farther away
from the horizontal meridian than +/- 45 deg). For each of the saccade sizes, we picked
movements landing within a radius of 0.5, 0.8, 1, 2, and 3 deg, respectively for the increasing
saccade amplitude categories listed above. Therefore, we ensured that the movement
endpoints were matched for landing accuracy. Example such movements are shown in Fig.
2a. In this figure, we only plotted rightward movements in monkey N and leftward
movements in monkey P, for simplicity, but Fig. 2b shows both rightward and leftward
saccades in each of the two monkeys. As can be seen from Fig. 2a, there was no clear
difference in the trajectories of upward (light blue) versus downward (light red) oblique
saccades, despite the significant SC neural asymmetry in Fig. 1. In fact, the pink upward
traces in Fig. 2a are the same as the light red downward traces in the figure, but now
reflected across the horizontal meridian for easier comparison to the upward saccades
shown in light blue. These pink traces clearly overlapped strongly with the upward saccades.
255
Across the population of measurements from the above saccades, we plotted radial eye
speed as a function of saccade amplitude and direction (Fig. 2b). This kind of plot
summarizes the kinematics of the eye movements[34,35]. For each saccade size and right/left
direction in each monkey, we plotted the radial eye speed for either upward (light blue) or
downward (light red) oblique saccades (error bars denote 95% confidence intervals). There
were no systematic differences in the saccadic profiles of the two groups of movements
(across all sizes tested), despite the systematically stronger SC motor bursts for downward
saccades seen in Fig. 1 (compare light blue and light red profiles for each saccade size). For
example, stronger motor bursts in Fig. 1 could have predicted systematically higher peak
speeds for the saccades directed towards the lower visual field[16]. This was clearly not the
case (Fig. 2b). In fact, lower visual field neurons possess larger movement fields than upper
visual field neurons[23], which should further increase the number of "active" spikes during
saccade-related bursting for saccades towards the lower visual field; nonetheless, the
kinematics of the movements were largely the same as those of upper visual field saccades
(Fig. 2). Therefore, the results so far are consistent with a dissociation between SC saccade-
related motor burst strength (Fig. 1) and saccade kinematics (Fig. 2).
272
273

Figure 2 Upward and downward oblique saccades from the same sessions as the neural recordings in Fig. 1 exhibited similar kinematic properties, despite the neural asymmetry of Fig. 1. (a) Oblique saccades of different directions and amplitudes in both monkeys (rightward for N and leftward for P). Each line plots the horizontal and vertical displacement of eye position for a given saccade having a +45 deg (light blue) or -45 deg (light red) direction from the horizontal meridian. Saccades of similar sizes are grouped together (to start from the same origin) and displaced horizontally in the figure from saccades of different size ranges (the scale bars apply to all sizes). The pink traces overlaying the light blue traces are the same as the light red traces of the lower visual field saccades, but now reflected along the vertical dimension for better comparison to the upper visual field saccades. The trajectories of the saccades were largely similar regardless of direction from the horizontal meridian, despite the asymmetry in motor bursts in Fig. 1. **(b)** For each monkey, we plotted radial eye speed as a function of time from saccade onset for leftward (left pair of plots) or rightward (right pair of plots) saccades. In each case, we separated upward and downward movements by color (as in **a**). The different saccade sizes in **a** are reflected in the different peak speeds[34,35]. The kinematic time courses of saccade acceleration, peak speed, and deceleration were largely similar for upward and downward saccades, despite the asymmetry of SC motor bursts in Fig. 1. Error bars denote 95% confidence intervals. The numbers of trials per condition in monkey N ranged from 7 to 362 (mean 121.2), and the numbers of trials per condition in monkey P ranged from 5 to 252 (mean 72.25).

Similarity of upper and lower visual field saccade kinematics for a variety of behavioral contexts

To further assess the dissociation between upper/lower visual field SC motor burst asymmetries and upward/downward saccade kinematics, we next turned to another, larger database of saccades for analyzing kinematics in more detail (dataset 2; Methods). In this case, we used: 1) immediate, visually-guided, 2) delayed, visually-guided, and 3) memory-guided saccades of different sizes and directions, with the sizes ranging from those associated with fixational microsaccades (approximately 0.1-0.2 deg) to approximately 15-20 deg. Aspects of these movements were analyzed previously for other purposes than movement kinematics[36,37]. Here, we wanted to confirm that the results of Fig. 2 still held for a larger range of movement amplitudes and directions, and also under different behavioral contexts. In other words, we analyzed the movements' kinematic properties in dataset 2, properties which were not analyzed in the prior publications. Moreover, dataset 2 allowed

308 us to include data from a third monkey, M, when assessing potential differences (or lack
309 thereof) in saccade kinematics between the upper and lower visual fields.
310
In our recent work with this dataset[36], we reported that saccadic reaction times were
systematically shorter for upper visual field target locations when compared to lower visual
field target locations, consistent with the asymmetry of SC visual neural sensitivity[23], and
also consistent with other behavioral evidence[38-41]. For example, in Fig. 3a, we plotted
example oblique saccades from monkey N from this dataset (but in a format similar to that
used in plotting the data of Fig. 2a). In Fig. 3b (left), we plotted the absolute value of vertical
eye position from the saccades shown in Fig. 3a (to facilitate comparing the upward and
downward movements). Here, we temporally aligned the movements to the time of the go
signal for triggering the saccades (the peripheral targets were continuously visible). Even
though the saccade trajectories looked similar in Fig. 3a (save for the upward and downward
distinction), the reaction times of the movements were markedly different (Fig. 3b, left).
Saccades towards the upper visual field were triggered significantly earlier than saccades
towards the lower visual field[36]. We then replotted the same saccades, but this time by
aligning them to the time of peak radial eye speed during the movements (Fig. 3b, right). The
movement kinematics were largely overlapping, with similar acceleration and deceleration
profiles.
327
328
329
330

331
Figure 3 Upward and downward saccades exhibit strong differences in reaction times, but not kinematics. (a)
From the second dataset, we plotted, by way of example, similarly-sized oblique saccades from one monkey, and
we separated them as being either upward or downward (as in Fig. 2a). **(b)** Relative to stimulus onset (left panel),
the saccades were very different from each other in terms of their reaction times, as we characterized in detail
earlier[36]; saccades towards the upper visual field (light blue) had significantly shorter reaction times than
saccades towards the lower visual field (light red). However, kinematically, the saccades were very similar when
aligned to the time of peak intra-saccadic eye speed (right panel). In both panels, we plotted the absolute value
of vertical eye position displacement for each saccade (for better comparison of the upward and downward
movements). Figures 4, 5 summarize the saccade kinematic results that we obtained for a much larger number
of movements, and for two monkeys. N=21 upward saccades and N=18 downward saccades in this figure.
342
343
344
345
To summarize this kinematic similarity across all saccades in this dataset, we generated plots
of peak eye speed as a function of saccade amplitude for each monkey[34,35]. In the first row of
Fig. 4, these plots were made for the immediate, visually-guided saccade task, in which the
fixation spot was extinguished at the same time as the appearance of the eccentric stimulus

(Methods; this task was not used during SC recordings because the visual and motor bursts
would occur too close to each other for proper neural analysis). For both monkeys N and M,
there was very minimal difference in the main sequence relationship between saccades
towards the upper (light blue) or lower (light red) visual fields, and any difference was
certainly much smaller than the neural effects in Fig. 1. In fact, the insets in the first row of
Fig. 4 show the reaction time results for the very same saccades, which are replicated from
our recent work[36] for clarity. Despite a large effect of the visual field location on the
movements' reaction times (also seen in Fig. 3b, left), there was minimal difference in
saccade kinematics. This is again supportive of a dissociation between saccade-related SC
motor burst strengths (Fig. 1) and movement kinematics (Figs. 2, 3); also see Figs. 6-8 below.
360
With an even larger database of visually-guided movements in this dataset, now from the
delayed, visually-guided saccade task, the same conclusion could be reached: the middle
row of Fig. 4 shows virtually no difference in the saccade kinematics between upward and
downward visually-guided saccades, despite a clear effect size for SC motor-related neural
responses in Fig. 1 and ref. [23]. The peak speeds in this row were also consistent with the
peak speeds in the first row of Fig. 4 obtained with the immediate, visually-guided saccade
task, as might be expected given the presence of a visual target for the saccades in both
tasks.
369
We also tested memory-guided saccades. Even though such saccades were generally slower
than visually-guided saccades (compare the bottom row of Fig. 4 to the two rows above it;
also see Figs. 6-8 below)[24-30], the above-mentioned kinematic similarity between movements
towards the upper and lower visual fields still persisted in the memory-guided saccade task.
Only in monkey N (left column of the bottom row of Fig. 4) was there a reduction in
downward saccade peak eye speed when compared to upward saccade peak eye speed.
However, even in this case, such a reduction was inconsistent with the stronger saccade-
related motor bursts for lower visual field saccades seen in the SC neural analyses (Fig. 1). If
anything, stronger lower visual field SC motor bursts (along with larger response fields[23])
might predict higher, rather than lower, peak speeds for downward saccades.
380
381
382

Figure 4 The main sequence relationship between peak eye speed and saccade amplitude does not depend on whether saccades are upward or downward for a variety of behavioral task contexts. In each monkey (left and right columns), we plotted the main sequence from dataset 2 but after first separating saccades as being directed towards either the upper visual field (light blue) or lower visual field (light red). The first two rows show visually-guided saccades (immediate for the first row, and delayed based on a task instruction in the second row; Methods). The third row shows memory-guided saccades towards a blank region of the display. The insets in the first row show saccadic reaction time (RT) data (as in Fig. 3b, left) for the same saccades as in the main sequence plots, to highlight the strong presence of a visual field effect on reaction times and a concomitant absence of a visual field effect on saccade kinematics. In all saccade contexts (across rows), the visual field location of the saccade endpoint had minimal effect on saccade kinematics (despite a large effect on saccadic reaction times and despite an asymmetry in SC motor bursts; Fig. 1). The insets in the first row directly replicate the plots in Fig. 4a, c of ref. [36] for easier comparison of effect sizes for reaction times and kinematics. Error bars denote s.e.m. Note that monkey N showed a small reduction of peak eye speed for downward saccades when compared to upward saccades only in the memory-guided condition (bottom row), but this effect is opposite from what would be expected if SC motor bursts (Fig. 1) dictated kinematics. The insets were replotted with permission from ref. [36].

Therefore, across a large range of movement sizes and directions, we found minimal kinematic differences between amplitude-matched upper and lower visual field saccades, even though other aspects of saccade generation (such as reaction times; insets in Fig. 4) were strongly different, and even though SC saccade-related motor bursts were also different (Fig. 1).

Finally, we also checked saccade durations as a function of saccade amplitudes (Fig. 5), and we reached similar conclusions. Saccade duration versus amplitude curves strongly overlapped for saccades towards the upper (light blue) and lower (light red) visual fields (Fig. 5), and this was true across task contexts. Note how monkey N compensated for the slightly

slower downward memory-guided saccade peak speeds (when compared to upward
memory-guided saccade peak speeds) with mildly longer durations for these movements
(left column of the bottom row of Fig. 5). This might suggest that there was lower drive for
generating this monkey's downward memory-guided saccades in general, which was then
compensated for by increased movement durations. Nonetheless, as stated above, this is an
opposite effect from what might be expected from the neural burst strengths in the SC (Fig.
1).
420
421
422

423
Figure 5 Saccade durations as a function of saccade amplitude are also largely insensitive to whether saccades
are directed towards the upper or lower visual field. Same as Fig. 4 but now plotting saccade duration as a
function of saccade amplitude. Similar conclusions were reached concerning the minimal influence of upper
versus lower visual field saccade target locations on saccade kinematics, despite asymmetries in SC motor bursts
(Fig. 1). For monkey N in the memory-guided condition (left panel of the bottom row), the slower lower visual
field saccade peak speeds (Fig. 4) meant slightly longer saccade durations when compared to upper visual field
saccades (a speed-duration tradeoff). Error bars denote s.e.m.
431
432
433
434 *Dissociation between SC motor burst strengths and movement kinematics also for vector-*
435 *matched saccades*
436
In the above experiments, and as stated above, we were motivated by an assumption of
directional isotropy in models of saccade control by the SC rate code[13,16,18]. In such models,
saccades are implemented (in terms of efferent connection strengths towards the
brainstem) according to their amplitude not vector; as a result, analyses of experimental

data often collapse measurements across different directions. We reasoned that if this was
indeed the case, then different SC burst strengths for upward and downward saccades (Fig.
1) should lead to different saccade kinematics, which we did not observe (Figs. 2-5). Having
said that, it may be argued that our observations so far merely suggest a different efferent
mapping to the downstream oculomotor control circuitry from the upper and lower visual
field SC representations, rather than a dissociation between SC motor burst strengths and
movement kinematics. While such a different efferent mapping between the upper and
lower visual fields would indeed be interesting, we elected to further test our original
hypothesis using a complementary approach, this time by employing vector-matched
saccades of different kinematics.
451
We exploited the fact that saccades to a blank (as in memory-guided saccades) can be
slower than visually-guided saccades[24-30]. We thus instructed 3 monkeys (M, N, and A) to
perform delayed, visually-guided saccades and memory-guided saccades towards the
response field hotspot locations of SC neurons. This meant that we now had even more SC
recording data from monkey N (beyond those shown in Fig. 1), as well as additional SC
motor burst measurements from two more monkeys (M and A), always comparing vector-
matched visually-guided and memory-guided saccades towards hotspot locations.
Moreover, in all 3 monkeys in this additional dataset (referred to here as dataset 3), we also
recorded neurons using linear electrode arrays (monkey M also contributed some single-
electrode sessions as well; Methods). This meant that we sometimes had simultaneously
recorded neurons for the same behavioral trials. For each isolated neuron, we first selected
saccades matched by direction and amplitude across the two tasks (data filtering
procedures, and minimum trial count requirements, are detailed in Methods). We then
checked the movement kinematics across the two tasks, and we evaluated how SC motor
bursts were potentially modified.
467
Figure 6a shows an example saccade vector from one of our sessions. In purple, we show the
delayed, visually-guided saccades from the session, and in green, we show the memory-
guided saccades. Despite being matched in direction and amplitude (as per our experimental
design), almost all memory-guided saccades from this session were slower than all visually-
guided saccades, as can be seen from Fig. 6b. Thus, we had vector-matched saccades with
clearly differing kinematics. We then checked how the SC motor bursts were altered. In Fig.
6c, we show the spike waveforms of 3 different SC neurons that we recorded simultaneously
from the same session in the two tasks. Each sub-plot in Fig. 6c shows the mean and
standard deviation of a random sampling of spike waveforms from a given isolated neuron in
both tasks (Methods). As can be seen, the waveforms were almost completely overlapping
for each neuron, suggesting that isolation quality was sufficiently stable for each of them as
we sequentially ran the two behavioral tasks (Methods). Therefore, we were now in a
position to compare the motor bursts of the neurons in the two behavioral contexts.
481
Surprisingly, there was a large diversity of motor burst modulations between the visually-
guided and memory-guided saccades, despite the highly consistent kinematic effects seen in
Fig. 6b. For example, Neuron 1 in Fig. 6d had a weaker burst in the memory-guided saccade
condition than in the visually-guided saccade condition, consistent with the kinematic effect
across the two conditions. However, Neuron 2 was much less affected by the behavioral
manipulation, and, most surprisingly, Neuron 3 had a much stronger motor burst in the
memory-guided condition instead of the visually-guided condition (Fig. 6d). Thus, there was

no systematic reduction in SC motor burst strengths (Fig. 6d) for the systematically slower
(but vector-matched) memory-guided saccades (Fig. 6b), as would be predicted by current
models of kinematic control by SC motor bursts.

497
Figure 6 Superior colliculus motor burst strength is dissociated from saccade kinematics even for vector-
matched movements. (a) Example saccade trajectories from one session of the vector-matched experiments.
Thin lines show individual movements, and thick lines show averages across trials. Purple denotes delayed,
visually-guided saccades, and green denotes memory-guided saccades. The vectors of the two types of saccades
were matched (Methods). **(b)** Memory-guided saccades were systematically slower than their visually-guided
counterparts, as might be expected[24-30]. **(c)** Spike waveforms from 3 example neurons recorded simultaneously
during the same session. Each thick line shows the average of a random sampling of waveforms from each of the
two tasks. The error bars denote the s.d. across observations. In all cases, the waveforms were stable across the
two task types, suggesting that each neuron was successfully recorded in the two tasks. The numbers of
waveforms included in the averages for the visually-guided and memory-guided conditions, respectively, are:
70/97 (Neuron 1), 79/74 (Neuron 2), and 42/54 (Neuron 3). **(d)** The neurons of **c** had a large diversity of effects,
in terms of peri-saccadic motor burst strengths, as a function of saccade type. Neuron 1 had a weaker motor
burst strength for memory-guided saccades; Neuron 2 was less affected by the condition; and Neuron 3 had a
surprisingly stronger motor burst in the memory-guided condition, despite the significantly slower saccades seen
in **b**. Error bars denote s.e.m. and numbers of trials are evident from the shown individual-trial spike rasters.

The above observations were repeatedly seen across our experiments. For example, in Fig.
7, we plotted the results from another example SC site, this time in the SC's lower visual field
representation. There was still a diversity of SC motor burst strength modulations as we
went from the visually-guided to the memory-guided saccade paradigms (Fig. 7d), despite
the matched saccade vectors (Fig. 7a), and also despite the clear kinematic differences
between the two conditions (Fig. 7b). There was also, again, a neuron in this session (Neuron

6) that violated the clearly slower saccades observed in Fig. 7b for the memory-guided
saccade condition. Thus, this additional example site revealed the very same patterns as
those shown in Fig. 6, and it suggests a dissociation between SC motor bursts and saccade
kinematics. A similar conclusion was also reached in earlier comparisons of the two tasks[24].

Figure 7 Similar observations to Fig. 6 from another example SC site. The figure is formatted identically to Fig.
6. The site of this example session was now in the SC's lower visual field representation, as evidenced by the
downward oblique saccades in **a**. Note how the saccade speed was clearly different between visually-guided and
memory-guided saccades (**b**), but the neurons still had a diversity of effects in terms of motor burst strengths
(**d**). The numbers of spike waveforms included in the averages of **c** for the visually-guided and memory-guided
conditions, respectively, are: 117/82 (Neuron 4), 76/65 (Neuron 5), and 85/81 (Neuron 6).

In total, we analyzed 114 SC neurons from 71 sites in these vector-matched experiments
(from monkeys M, N, and A; Methods). To summarize their results, we first confirmed that
all saccades were vector-matched across the visually-guided and memory-guided saccade
conditions, as per our experimental design. For each of the 71 sites, each having a unique
saccade vector, we collected the average saccade vector from each of the two conditions.
We then plotted the amplitude (Fig. 8a) and direction (Fig. 8b) of the memory-guided
saccade vector against the amplitude and direction of the visually-guided saccade vector.
There was no difference between the two conditions in either amplitude (Fig. 8a) or
direction (Fig. 8b) (amplitude comparison: p = 0.5313, paired t-test, t-statistic: -0.6292, df:
70; direction comparison: p = 0.9735, paired t-test, t-statistic: -0.0334, df: 70), as expected
(we explicitly matched the vectors of the saccades in these experiments; Methods). We then
confirmed that the saccades were significantly slower in the memory-guided condition than
in the visually-guided condition[24-30], and we did so by plotting in Fig. 8c the peak speeds from
all experiments against each other, in a fashion similar to the amplitude and direction plots

of Fig. 8a, b. Memory-guided saccades were significantly slower than vector-matched visually-guided saccades (p = 6.21x10^{-13}, paired t-test, t-statistic: -8.7979, df: 70). Most importantly, we then related the peak speed effect (for the vector-matched movements) to the neural motor burst effect. To do so, we measured each neuron's peak firing rate in the interval -25 to +25 ms from saccade onset (Methods). We then created a neural modulation index as the burst strength in the memory-guided condition minus the burst strength in the visually-guided condition, divided by the sum of the two burst strengths (Methods). Values of the index >0 would indicate that motor bursts were actually stronger in the memory-guided saccade condition than in the visually-guided saccade condition. Similarly, we created a behavioral modulation index as the peak saccade speed in the memory-guided condition minus the peak saccade speed in the visually-guided condition, divided by the sum of the two peak speeds (Methods). Across all neurons, there was no correlation between the neural and behavioral modulation indices (Fig. 8d) (Pearson correlation coefficient: -0.1374, p = 0.1449). Rather, there was a more-or-less constant behavioral effect (slower saccades in the memory-guided condition) irrespective of SC neural modulation effect, as evidenced by the vertical scatter of points across all neurons in Fig. 8d.

Figure 8 Population summary demonstrating how SC motor burst strength is dissociated from saccade kinematics even for vector-matched movements. (a) Average saccade amplitude in the memory-guided saccade task versus the visually-guided saccade task across all unique sessions in this experiment. Each symbol indicates a single session. As per the experimental design, the saccade amplitudes were matched across the two conditions. **(b)** Same as **a** but for the directions of the saccades. Negative means downward saccades, and positive means upward saccades. Again, there was no difference in saccade angles across the visually-guided and memory-guided conditions. Thus, **a** and **b** indicate that the saccades in the two conditions were vector-matched. **(c)** Despite the vector matching, peak speeds were consistently lower for memory-guided saccades. **(d)** For all recorded neurons from the same sessions, we calculated a behavioral modulation index, in which negative values indicated slower saccades in the memory-guided condition. We also created a neural modulation index, in which negative values indicated weaker motor bursts in the memory-guided condition. Note that some neurons were recorded simultaneously during the same trials, as in Figs. 6, 7 (Methods). Thus, there were more neurons than sessions, explaining the different numbers of symbols from **a-c**; this also meant that there could be multiple symbols with different y-axis values but having the same x-axis value (because multiple neurons were collected for the same behavioral trials). There was no correlation between neural and behavioral modulation indices.

Interestingly, there was a large dynamic range of neural modulation indices. Some neurons were almost completely suppressed in the memory-guided condition. These can qualify as visually-dependent saccade-related neurons[24,31,37]. Alternatively, 46.49% of the neurons (53/114) were above the diagonal line in Fig. 8d, thus violating the predictions of current models of kinematic control by SC motor bursts. Most intriguingly, almost one quarter of the neurons (23.7%, 27/114) had a neural modulation index >0, suggesting that these neurons

601 actually exhibited stronger motor bursts for memory-guided saccades than for vector-
602 matched visually-guided saccades (Fig. 8d), despite the significantly slower speeds of the
603 former (Fig. 8c). Thus, the motor bursts were independent of the actual triggered saccades.
604
605 These results, combined with those of Figs. 1-5 above, suggest that there is indeed a
606 dissociation between SC motor burst strengths and saccade kinematics.
607
608

609 **Discussion**

610
We described a dissociation between SC saccade-related motor burst strengths and
movement kinematics. In particular, we confirmed an asymmetry in motor burst strengths
between upper and lower visual field saccade target locations (Fig. 1). We then found that
the kinematics of amplitude-matched saccades towards upper and lower visual field
locations were not different from each other across a range of movement sizes, directions,
and behavioral contexts (Figs. 2-5). Finally, we demonstrated how there was no correlation
between SC motor burst effects and kinematic alterations even in vector-matched saccades
towards response field hotspot locations (Figs. 6-8).

619
Other examples of dissociations between SC motor burst strengths and movement
properties are consistent with our interpretation that the SC rate code does not necessarily
dictate movement kinematics, as might be suggested by some recent models[13,16]. For
example, and as we have confirmed (Fig. 8d), for memory-guided saccades towards
response field hotspot locations, a significant fraction of SC neurons becomes silent at the
time of movement onset[24,31,37]. This, in addition to neurons that exhibit potentially altered
response field profiles when making saccades towards a blank[29], suggests a significantly
modified SC representation during these movements. Indeed, we even found neurons that
exhibited stronger, rather than weaker, motor bursts for slower memory-guided saccades
(Fig. 8). Therefore, the relationship between movement kinematics and SC motor bursts is
relatively loose when making saccades towards a blank, and it was also relatively loose in
our analyses of upper and lower visual field target locations.

632
Another example of a dissociation between saccade motor burst strength and movement
kinematics was observed when saccades were driven by combinations of visual and auditory
sensory signals, as opposed to only visual signals[17]. Interestingly, it was again the case in this
example that a sensory scenario was relevant and critical for revealing a potential separation
between the SC rate code and movement kinematics. That is, in both the example above of
saccades towards a blank as well as the current example of multi-sensory target
specification, it was a modification of a sensory property of saccade targets that has allowed
observing a dissociation between motor burst strengths and eye movement properties. This
clear context-dependence of the bursts indicates that SC "motor" bursts are likely not pure
motor controllers, in the strictest sense of the word. In the current manuscript's context as
well, we were originally motivated by the fact that it was visual sensitivity that was strongly
variable between upper and lower visual field locations[23]. Indeed, given that stronger visual
responses occur in the upper visual field whereas stronger motor responses occur in the
lower visual field, it is intriguing to consider the possibility that there might be a general anti-
correlation property between visual sensitivity and saccade-related motor burst strength in
the SC, for example, in the ubiquitous visual-motor neurons of this structure.

More recently, Peel and colleagues also identified a dissociation between SC motor burst
strength and executed saccade properties[22]. Specifically, these authors causally perturbed
top-down inputs towards the SC through reversible cooling of the cortex, and they found
reduced SC burst strengths for metrically similar executed saccades. This study, along with
refs. [17,24,31,37], all showed that for the very same saccade vector (i.e. only within the upper
visual field SC representation or only within the lower visual field SC representation),
sensory[17,24,31,37] or physiological[22] manipulations can indeed significantly alter SC burst
strengths without fully accounting for the altered saccade properties. Therefore, whether
one considers a single saccade vector like in these studies or a comparison of upper versus
lower visual field saccades like in the first half of our study, a dissociation between SC
saccade-related motor bursts and saccade execution still exists.

One potential criticism of our comparison of upper and lower visual field saccades could be
that the existing models of the rate code dictating saccade kinematics[13,16] have primarily
focused on a single saccade vector, whereas we compared upward and downward saccades.
According to this potential counter argument to our results, by comparing upper and lower
visual field saccades, we might have been comparing saccades in which structures
downstream from the SC would compensate for the SC asymmetry shown in Fig. 1 and in ref.
[23]. Because the above-mentioned studies all showed dissociations between the SC rate code
and saccade properties for the very same saccade vector, just like the models of the SC rate
code[13,16], this counter argument does not seem to be too strong. In addition, we explicitly
performed single vector experiments (to response field hotspot locations) in Figs. 6-8, and
we found results that were strongly consistent with our original hypothesis. Perhaps most
critically, this potential counter argument might imply that the large SC asymmetry in both
saccade-related movement burst strengths (Fig. 1) and movement response field sizes[23] just
becomes nulled downstream of the SC, which creates the problem of why the asymmetry
exists in the SC at all in the first place. It also ignores the fact that the asymmetry has real
consequences for saccade latencies, saccade accuracy, and the likelihood of express
saccades[23]. Rather, we believe that the SC asymmetry motivates investigating what the
functional role of SC motor bursts is, and in a more general framework than just one of
controlling saccade kinematics.

Our results also provide complementary evidence to a phenomenon that we recently
studied[21]. In that recent study, we altered saccade metrics and kinematics by strategically
injecting "visual" bursts into the SC at spatial sites beyond the vector endpoints of the
currently executed movements[21]. We found alterations in movement metrics and
kinematics[20], which were lawfully related to the amounts of visual spikes that we injected
onto the SC map around the time of movement triggering[21]. These results were consistent
with the spatial code of the SC because the movement metric changes reflected the
topographic locations of the injected spikes. However, critically, the movements' kinematic
alterations occurred in the absence of strong alterations in the motor bursts themselves (for
the neurons generating the originally planned saccades). This was surprising for a variety of
reasons, including ideas related to lateral connectivity patterns in the SC, as we recently
discussed[21]. However, it also represented an opportunity for us to explicitly ask, in the
current study, whether or not the rate code was indeed as tightly linked to individual
movement kinematics. Therefore, here, in the first half of the study (Figs. 1-5), we took the
opposite approach from our recent study: we identified a situation in which the motor

bursts were different from each other for two different sets of amplitude-matched saccades, and we showed that the saccade kinematics in the two groups of movements were the same (Figs. 1-5). Of course, we also considered the case in which the same saccade vector was made towards the response field hotspot location but with different kinematics (Figs. 6-8). The net result is that either with altered movements and minimally-altered movement commands[21], or with minimally-altered movements and significantly altered movement commands (Figs. 1-5), or with vector-matched movements of different kinematics and a diversity of SC motor burst effects (Figs. 6-8), there does indeed seem to be a clear dissociation between saccade kinematics and SC motor burst strengths.

Recent work has suggested that the SC can support high level perceptual and cognitive phenomena[42,43]. For example, the SC causally influences selective behaviors[44,45], and it even shapes object-related visual representations in the ventral visual processing stream[46]. This is in addition to established roles for the SC in target selection[47-49]. All of this evidence suggests the SC might occupy a functional level that is slightly more abstract than that of specifying individual movement kinematics, consistent with our results. Thus, it might suffice for the SC to specify desired movement metrics, via the spatial code, and also potentially contribute to the decision of when to trigger an eye movement, as recently suggested[11]. The rest can be handled by downstream oculomotor control structures. If this is indeed the case, then a critical and urgent question for research in the immediate future is: what is, ultimately, the functional role of the SC rate code in visual-motor behavior and perception? One possibility could be that it allows providing a differential gain signal for cortical visual processing. For example, it is known that visual perception[50-53] and attention[54,55] are better in the lower visual field under conditions of gaze fixation. However, peri-saccadic perceptual mislocalization performance is different for upward saccades[56]. Moreover, when we recently measured perceptual sensitivity in peri-saccadic intervals, at the time of saccadic suppression[57], we found such sensitivity to be better in the upper visual field instead - consistent with a stronger peri-saccadic suppression of visual sensitivity in the lower visual field[58]. If SC motor bursts contribute to saccadic suppression, perhaps via inhibitory projections to the frontal cortex[32,59,60], then a possible functional role for stronger motor bursts in the SC's lower visual field representation could be to differentially modulate cortical visual processing at the time of saccades. It would be interesting to investigate this hypothesis in future studies.

Methods

In this study, we described results from three different sets of experiments, referred to as dataset 1, dataset 2, and dataset 3, respectively.

In dataset 1, we analyzed data from our previously published study[23]. Specifically, neural activity from the SC and saccadic behavior were recorded from two adult, male rhesus macaque monkeys (P and N)[23]. We analyzed both neural activity and behavior from that study, using a delayed, visually-guided saccade task.

In dataset 2, we analyzed saccadic behavior that was recorded from monkey N and a third adult, male monkey (M), again from a previously published experiment[36]; here, we analyzed additional behavioral parameters from that study that were not previously described. We also analyzed multiple behavioral tasks.

In dataset 3, we analyzed both saccadic behavior and SC neural activity from adult, male monkeys N, M, and A. The experiments consisted of either single-electrode recordings in monkey M or linear electrode array recordings in all 3 monkeys. The linear electrode array recordings in monkeys N and M were re-analyzed from a previous study[37], whereas the linear electrode array recordings from monkey A (aged 10 years and weighing 10 kg), as well as the single-electrode recordings in monkey M, were all newly-performed experiments not previously described in any other publication.

Thus, we had SC neural recordings from a total of 4 monkeys (M, N, A, and P) and behavior from a total of 3 monkeys (M, N, and P) in this study.

All experiments were approved by the Regierungspräsidium Tübingen, under licenses CIN3/13 and CIN4/19G, and they were in accordance with the German and European directives on the use of animals in research.

In what follows, we describe detailed methods relevant for the current work.

Animal preparation

For SC recording, a recording chamber was implanted centered on the midline in all 4 monkeys. The midline positioning of the chamber allowed recording from both the right and left SC in each animal. Magnetic resonance images (MRI's) obtained prior to the experiments aided in chamber implant alignment. We aimed for quasi-orthogonal electrode penetrations (relative to the SC curvature) at eccentricities we typically use in experiments (e.g. 5-15 deg).

Before receiving the chamber implants, the animals were also implanted with head-holding apparatuses and scleral search coils for eye tracking, as described earlier[61-63]. The scleral search coils allowed using the magnetic induction technique for measuring eye positions[64,65]. Specifically, a coil of wire was implanted around the sclera of the eye and below the conjunctiva. The animals were then seated near the middle of a cube in which alternating magnetic fields induced electrical current (which depended on ocular position) in the implanted scleral coil; we measured and calibrated this electrical current.

Behavioral tasks

For both neural and behavioral analyses, the monkeys performed classic saccade generation tasks.

In the immediate, visually-guided saccade task of dataset 2 (which was only used in behavioral experiments and not neurophysiological experiments), the monkey first fixated a central spot. After a variable delay, the spot was jumped to another location, and a saccade to follow the spot was triggered.

In the delayed version of the same task, during initial fixation, the fixation spot remained visible while an eccentric spot was presented. The monkey was required to maintain gaze fixation and withhold any reflexive orienting towards the eccentric spot for as long as the central fixation spot was visible. After the fixation spot was removed, the monkey generated a saccade towards the (still visible) eccentric spot.

Finally, in the memory-guided saccade task, during initial fixation, the eccentric spot was only flashed briefly (for approximately 50 ms). A delay period then ensued in which only the fixation spot was visible, and the monkey was required to maintain gaze fixation on it. At the end of this so-called memory period, the fixation spot was extinguished, instructing the monkey to generate a saccade towards the remembered location of the previous flash (i.e. towards a blank location on the display).

The delayed, visually-guided saccade task was used for all neural analyses reported in this study (dataset 1 and dataset 3). This was important because this task allows dissociating visual burst intervals from the saccade-related motor burst intervals that we were interested in analyzing. The memory-guided saccade task was also used for neural analyses in dataset 3. For behavioral analyses, we used the delayed saccade task in dataset 1 (e.g. Fig. 2), as well as all 3 saccade tasks in dataset 2.

For mapping response fields, we generally employed the delayed, visually-guided and memory-guided saccade tasks. However, in the newly-acquired portions of dataset 3 (monkey M single-electrode recordings and monkey A linear electrode array recordings), we first mapped response fields with a fixation variant of the delayed, visually-guided saccade task. That is, at the end of the trial, instead of fixation spot removal to release a visually-guided saccade towards the eccentric target, the monkey was simply rewarded for fixating until trial end. This allowed us to obtain visual response fields, and we then later tested for saccade-related bursts using the delayed, visually-guided and memory-guided saccade tasks.

In all cases, stimuli were presented on cathode ray tube (CRT) displays, with stimulus luminances and dimensions having been described earlier[23,36,66]. The timing of trial events in the tasks was also described earlier. For the present study, the primary focus was on the individual saccade kinematics at the ends of all trials, irrespective of timing parameters, such as the length of the delay or memory period, and irrespective of the exact stimulus visual properties. The effects of these factors (such as trial timing or visual stimulus properties) were described earlier[23,36,67,68].

828
829
Behavioral data analyses

All saccades from datasets 1 and 2 were detected for the previous two studies[23,36]. Here, we
analyzed the kinematic properties of the movements. For dataset 3, the saccades from the
electrode array recordings of monkeys M and N were also detected previously[37]. The
saccades from the newly-acquired monkey M and monkey A recordings were detected using
our standard approaches[61,69].

For behavioral analyses in dataset 1, we picked saccades having +45 or -45 deg direction
from the horizontal meridian (i.e. oblique saccades). We then picked 5 radial amplitude
categories to characterize 5 different ranges of saccade sizes (Fig. 2). The categories were: 3,
5, 7, 10, and 13 deg. For each of these categories, we picked all saccades landing within a
radius of 0.5, 0.8, 1, 2, and 3 deg from the designated amplitude/direction category,
respectively. For example, for saccades of 7 deg amplitude and +45 deg direction, we picked
all saccades that were upward and oblique, and that were directed towards an eccentricity
of 7 deg, and that landed within a radius of 1 deg from this eccentricity. Similarly, for 3 deg
saccades of +45 deg direction, we picked all upward oblique movements towards an
eccentricity of 3 deg and landing within a radius of 0.5 deg from it. This meant that we had
amplitude- and direction-matched saccades for either the oblique upward or the oblique
downward movements. We then plotted the trajectories (Fig. 2a) and radial speed profiles
(Fig. 2b) of all of these saccades. Since the speeds of temporally-directed saccades could be
different from the speeds of nasally-directed saccades for a given tracked eye, we analyzed
rightward and leftward saccades separately in this dataset (Fig. 2b). However, in dataset 2,
all saccade directions were combined, and with similar conclusions.

For dataset 2, we had a large range of saccade amplitudes and directions to analyze[36]. We
plotted the main sequence relationship[34,35] for these saccades after separating them into
two groups: saccades towards the upper visual field and saccades towards the lower visual
field. We plotted both the main sequence relationship of peak speed versus movement
amplitude (Fig. 4) and saccade duration versus movement amplitude (Fig. 5). For
comparison, we included a plot of saccadic reaction times for the same saccades in Fig. 4.
This was a replotting of the reaction time data already reported earlier[36], and we included it
here for easier comparison of the difference in effects of visual field location on saccade
kinematics and saccade reaction times. In total, we analyzed 1246, 928 visually-guided
saccade trials, 6147, 5871 delayed, visually-guided saccade trials, and 6428, 9631 memory-
guided saccade trials from monkeys N and M, respectively. The numbers of trials for the
behavioral analyses from dataset 1 are reported in the figure legend of Fig. 2.

For dataset 3, our behavioral analyses consisted of first ensuring vector matching and then
checking the movement kinematics to set the stage for neural data analyses. For the
previously collected data[37] (monkey M and monkey N linear electrode array recordings),
response fields were mapped with both the delayed, visually-guided and memory-guided
saccade tasks. Therefore, in offline analyses, we obtained the average firing rate in the
interval of -25 ms to +25 ms from saccade onset for each movement. We then plotted heat
maps of firing rate as a function of saccade horizontal and vertical amplitudes to confirm the
response fields. We identified the hotspot location of each neuron from the visually-guided

saccade response field, and we then picked all saccades in both tasks landing within 2 deg, 1
deg, or 0.5 deg of this location depending on the neuron's preferred eccentricity (within 2
deg for neurons with preferred eccentricity > 3 deg, within 1 deg for neurons in the range of
2-3 deg preferred eccentricity, and within 0.5 deg for foveal neurons). We only included
neurons for which we had at least 5 vector-matched saccades in each of the visually-guided
and memory-guided saccade conditions.
882
For the newly-collected measurements of dataset 3 (monkey M single-electrode recordings
and monkey A linear electrode array recordings), after mapping visual response fields with
the fixation task (described above), we ran delayed, visually-guided saccades and memory-
guided saccades towards the hotspot location (as assessed online during the experiment),
collecting at least 20 trials per task. We then checked for endpoint matching. We found the
median landing position from the delayed, visually-guided saccade task. Then, we only
included saccades in both tasks that landed within 1 deg from this position. Once again, we
only included neurons for which we had at least 5 vector-matched saccades in each of the
visually-guided and memory-guided saccade conditions (typically much more).
892
After finding vector-matched saccades in dataset 3, we then proceeded to plot radial eye
speed for the delayed, visually-guided and memory-guided saccade tasks. We also collected
measurements per session as follows: average saccade amplitude, average saccade
direction, and average saccade peak speed. This allowed us to plot these parameters across
the two tasks (e.g. Fig. 8a-c), to confirm vector matching as well as to confirm different
saccade kinematics across the two conditions.
899
To obtain a single behavioral modulation index across the two tasks in dataset 3, we
measured, in each session, the average peak saccade speed in the memory-guided condition
and the average peak saccade speed in the delayed, visually-guided condition. We
subtracted the latter from the former, and then divided by the sum of the two. This gave us
an index that ranged in values from -1 to +1, with indices >0 indicating that peak saccade
speed was higher in the memory-guided condition and indices <0 indicating that peak
saccade speed was higher in the delayed, visually-guided saccade condition. This gave us a
single number that we could relate to a similar single number for SC motor burst strength
modulation by the behavioral task (as described later below).
909
910
911 *Dataset 1 neural data analyses (Fig. 1)*
912
We analyzed peri-saccadic firing rates, as we did previously[23]. We obtained firing rates by
convolving individual spike times with a gaussian kernel of 40 ms σ. For each neuron in the
database of the previous study (containing >400 neurons), we had identified (for saccade-
related neurons) the saccades towards the neuron's preferred movement-related response
field location (i.e. the locations for which the neuron's saccade-related bursts were the
strongest). In the present study, we analyzed the firing rates for these preferred saccades.
However, we constrained the choice of neurons according to the needs of the current study.
Specifically, besides only considering extra-foveal neurons with saccade-related bursts, we
matched neural depths between neurons from the upper and lower visual field
representations of the SC (e.g. Fig. 1a). Specifically, since saccade-related motor bursts in the
SC can vary in strength as a function of depth of the neurons from the SC surface[6], we only

compared motor bursts after selecting neurons from the upper and lower visual field
representations that had matched depths.

To do so, we first considered all neurons in the upper and lower visual field representations
having a depth of 600-1850 μm from the SC surface. This range of depths is consistent with
known depths of saccade-related activity in the SC[6]. Importantly, for the present purposes,
this range of depths contained clear overlap between neurons in the upper and lower visual
field SC representations (Fig. 1a). This allowed comparing the strengths of motor bursts
between the selected depth-matched neurons. The resulting neural database had 136
neurons (Fig. 1).

To further confirm that there was no confound of neural depth from the SC surface in
interpreting a visual field asymmetry in motor burst strength, we were concerned that the
curvature of the SC surface could introduce systematic biases in depths of upper versus
lower visual field neurons from the SC surface. For example, it could potentially be the case
that the 3-dimensional SC surface curvature combined with a constant electrode approach
angle dictated by the recording chamber might systematically skew depth estimates: medial
(upper visual field) electrode locations might potentially have depth estimates that could be
systematically different from lateral (lower visual field) electrode locations in the chamber.
This could simply be a function of whether or not a given electrode track was more or less
perpendicular to the local SC surface topography at a given site. In our second analysis of
neural activity, we therefore picked a range of electrode locations in which we expected
minimal changes in SC curvature between upper and lower visual field representations. For
example, mapping the SC surface topography on the anatomical SC[33] might suggest a similar
relationship between electrode angle and SC surface for upper and lower visual field
representations near the horizontal meridian and within a specific range of movement
amplitudes. We therefore specifically picked neurons with movement-related response field
hotspots near the horizontal meridian (within 30 deg direction in either the upper or lower
visual fields) and with radial eccentricities of only 5-15 deg. We also picked a narrower depth
of neurons for the comparison (1100-1900 μm from SC surface). With this stricter neural
database (31 neurons), we again plotted peri-saccadic firing rates for neurons in the upper
and lower visual field representations (Fig. 1c).

In all cases, a neuron was considered to be part of the upper or lower visual field
representation if its preferred saccade (i.e. the movement-related response field hotspot
location) was in the upper or lower visual field, respectively. This was also consistent with
the known SC topographic representation[1,3,23,33], and it was already done in our previous
study[23].

To statistically compare saccade-related activity strength between the upper and lower
visual field representations in the SC, we measured the average firing rate in the final 50 ms
before saccade onset for each neuron. We then statistically compared the firing rates of all
neurons having movement-related response field hotspot locations in the upper visual field
to the firing rates of all neurons having response field hotspot locations in the lower visual
field (using t-tests). Note that measuring average firing rates is equivalent to counting spikes,
which has been the standard method to analyze the rate code[13,18]. Also note that in our
analyses of dataset 3 recordings (described below), we also picked a peri-movement burst

measurement (that is, including epochs also after saccade onset) rather than only a pre-
movement measurement as in dataset 1, with similar conclusions to this dataset's results.
973
974
975   *Dataset 3 neural data analyses (Figs. 6-8)*
976
977   We collected 25 sessions of linear electrode array recordings in monkey N, 16 sessions of
978   linear electrode array recordings and 32 sessions of single-electrode recordings in monkey
979   M, and 12 sessions of linear electrode array recordings in monkey A (all array recordings
980   were performed with V-probes from Plexon, Inc.). The linear electrode array recordings from
981   monkeys N and M were a subset of those described for a previous study[37].
982
We used offline sorting to identify single neurons. For the single-electrode recordings, we
used Plexon's Offline Sorter utility. The visually-guided and memory-guided saccade tasks
were collected (in sequence) together in the same file, and we sorted both tasks together.
For the linear electrode array recordings, we performed offline sorting using Kilosort[70],
followed by manual curation using the phy software. For the data from ref. [37], we used the
same sorting results that were obtained for the original study. Isolated neurons that
exceeded an estimated false positive rate (ISI violation) of 10% or had an isolation distance
below 30 were excluded from further analysis. We sorted linear electrode array recording
data from an entire session simultaneously, thus tracking neurons across the different tasks
that we ran (typically much more than the visually-guided and memory-guided saccade
tasks). To check isolation stability (e.g. for Fig. 6c), we collected 2000 spike waveforms
selected randomly from the same session. For our two tasks of interest, we took the
waveforms from this sampling of waveforms that happened to come from either of the two
tasks, and we plotted their distributions. Because the two tasks were typically run in
succession, it was usually very likely that isolation was stable throughout both of them. This
was also the case in the single-electrode recordings.
999
For each of the vector-matched saccades, we defined a motor interval as the 50-ms interval
centered on saccade onset (that is, the interval spanning -25 to +25 ms from saccade onset).
We also defined a baseline interval as the final 50-ms interval before stimulus onset (at trial
beginning). We then statistically compared the baseline interval firing rate to the motor
interval firing rate using either a t-test (for all the newly-collected data) or a ranksum test
(for the data from ref. [37]). The choice of test was dictated by the fact that the old data had
fewer numbers of trials because we selected saccades from response field mapping data,
whereas in the newly-collected sessions, we explicitly collected repeated saccades from the
same response field hotspot location. We only included a neuron if it had a significantly
elevated average firing rate in the motor interval relative to the average firing rate in the
baseline interval ($p<0.05$) in either the delayed, visually-guided saccade task or the memory-
guided saccade task or both.
1012
To check for changes in SC motor bursts across visually-guided and memory-guided
saccades, we calculated the peak firing rate in the motor interval (-25 to +25 ms from
saccade onset) in each condition. We then calculated a neural modulation index similar to
how we computed the behavioral modulation index above. That is, we subtracted peak firing
rate in the visually-guided saccades from peak firing rate in the memory-guided saccades,

and we divided by the sum of peak firing rates. Thus, a neural modulation index >0 indicated
stronger SC motor bursts for memory-guided than visually-guided saccades.
1020
1021
1022 *Statistics and reproducibility*
1023
1024 We analyzed SC recording data from 4 different monkeys, with consistent results. Similarly,
1025 we analyzed behavioral measurements from 3 monkeys. These numbers of animals increase
1026 confidence in the generalizability of the results, especially given how some observations
1027 were highly consistent with a large literature on SC saccade-related bursts (e.g. the depth
1028 profile shown in Fig. 1a).
1029
1030 For neural analyses, we statistically compared upper and lower visual field motor bursts on a
1031 neuron per neuron basis (Fig. 1). Similarly, in Figs. 6-8, we compared burst strengths on a
1032 neuron per neuron basis. In all cases, we collected a large enough sample of neurons to
1033 increase statistical confidence in the observations.
1034
1035 For eye movement analyses, we employed minimum count requirements (e.g. for the vector
1036 matching in the experiments of Figs. 6-8) to ensure enough replicates. Similarly, we used
1037 large numbers of replicates in the behavioral measurements of Figs. 2-5.
1038
1039 All relevant statistical tests are indicated in the figure legends or the associated Results text.
1040 Also, numbers of observations are indicated in the figures, in the figure legends, in Results,
1041 or in the Methods text.
1042
1043
1044 **Data availability**
1045
1046 The datasets generated during and/or analysed during the current study are available from
1047 the corresponding author on reasonable request.
1048
1049

<div style="text-align:center">References</div>

Robinson, D. A. Eye movements evoked by collicular stimulation in the alert monkey. *Vision Res* **12**, 1795-1808 (1972). https://doi.org:0042-6989(72)90070-3 [pii]

Katnani, H. A. & Gandhi, N. J. The relative impact of microstimulation parameters on movement generation. *J Neurophysiol* **108**, 528-538 (2012). https://doi.org:10.1152/jn.00257.2012

Cynader, M. & Berman, N. Receptive-field organization of monkey superior colliculus. *J Neurophysiol* **35**, 187-201 (1972).

Lee, C., Rohrer, W. H. & Sparks, D. L. Population coding of saccadic eye movements by neurons in the superior colliculus. *Nature* **332**, 357-360 (1988). https://doi.org:10.1038/332357a0

Munoz, D. P. & Wurtz, R. H. Saccade-related activity in monkey superior colliculus. II. Spread of activity during saccades. *J Neurophysiol* **73**, 2334-2348 (1995).

Massot, C., Jagadisan, U. K. & Gandhi, N. J. Sensorimotor transformation elicits systematic patterns of activity along the dorsoventral extent of the superior colliculus in the macaque monkey. *Commun Biol* **2**, 287 (2019). https://doi.org:10.1038/s42003-019-0527-y

Wurtz, R. H. & Goldberg, M. E. Superior colliculus cell responses related to eye movements in awake monkeys. *Science* **171**, 82-84 (1971).

Schiller, P. H. & Koerner, F. Discharge characteristics of single units in superior colliculus of the alert rhesus monkey. *J Neurophysiol* **34**, 920-936 (1971).

Munoz, D. P. & Wurtz, R. H. Saccade-related activity in monkey superior colliculus. I. Characteristics of burst and buildup cells. *J Neurophysiol* **73**, 2313-2333 (1995).

Sparks, D. L., Holland, R. & Guthrie, B. L. Size and distribution of movement fields in the monkey superior colliculus. *Brain Res* **113**, 21-34 (1976). https://doi.org:10.1016/0006-8993(76)90003-2

Jagadisan, U. K. & Gandhi, N. J. Population temporal structure supplements the rate code during sensorimotor transformations. *Current Biology* **32**, 1010-1025 (2022). https://doi.org:10.1016/j.cub.2022.01.015

Goossens, H. H. & Van Opstal, A. J. Blink-perturbed saccades in monkey. II. Superior colliculus activity. *J Neurophysiol* **83**, 3430-3452 (2000).

13   Goossens, H. H. & van Opstal, A. J. Optimal control of saccades by spatial-temporal
activity patterns in the monkey superior colliculus. *PLoS computational biology* **8**,
e1002508 (2012). https://doi.org:10.1371/journal.pcbi.1002508

14   Waitzman, D. M., Ma, T. P., Optican, L. M. & Wurtz, R. H. Superior colliculus neurons
mediate the dynamic characteristics of saccades. *J Neurophysiol* **66**, 1716-1737
(1991).

15   Waitzman, D. M., Ma, T. P., Optican, L. M. & Wurtz, R. H. Superior colliculus neurons
provide the saccadic motor error signal. *Exp Brain Res* **72**, 649-652 (1988).

16   Smalianchuk, I., Jagadisan, U. K. & Gandhi, N. J. Instantaneous Midbrain Control of
Saccade Velocity. *J Neurosci* **38**, 10156-10167 (2018).
https://doi.org:10.1523/JNEUROSCI.0962-18.2018

17   Frens, M. A. & Van Opstal, A. J. Visual-auditory interactions modulate saccade-
related activity in monkey superior colliculus. *Brain Res Bull* **46**, 211-224 (1998).
https://doi.org:S0361-9230(98)00007-0 [pii]

18   van Opstal, A. J. & Goossens, H. H. Linear ensemble-coding in midbrain superior
colliculus specifies the saccade kinematics. *Biol Cybern* **98**, 561-577 (2008).
https://doi.org:10.1007/s00422-008-0219-z

19   Buonocore, A., McIntosh, R. D. & Melcher, D. Beyond the point of no return: effects
of visual distractors on saccade amplitude and velocity. *J Neurophysiol* **115**, 752-762
(2016). https://doi.org:10.1152/jn.00939.2015

20   Buonocore, A. *et al.* Alteration of the microsaccadic velocity-amplitude main
sequence relationship after visual transients: implications for models of saccade
control. *J Neurophysiol* **117**, 1894-1910 (2017).
https://doi.org:10.1152/jn.00811.2016

21   Buonocore, A., Tian, X., Khademi, F. & Hafed, Z. M. Instantaneous movement-
unrelated midbrain activity modifies ongoing eye movements. *eLife* **10** (2021).
https://doi.org:10.7554/eLife.64150

22   Peel, T. R., Dash, S., Lomber, S. G. & Corneil, B. D. Frontal eye field inactivation alters
the readout of superior colliculus activity for saccade generation in a task-dependent
manner. *J Comput Neurosci* (2020). https://doi.org:10.1007/s10827-020-00760-7

23   Hafed, Z. M. & Chen, C. Y. Sharper, Stronger, Faster Upper Visual Field
Representation in Primate Superior Colliculus. *Curr Biol* **26**, 1647-1658 (2016).
https://doi.org:10.1016/j.cub.2016.04.059

24   Edelman, J. A. & Goldberg, M. E. Dependence of saccade-related activity in the
primate superior colliculus on visual target presence. *J Neurophysiol* **86**, 676-691
(2001). https://doi.org:10.1152/jn.2001.86.2.676

25   Gnadt, J. W. & Andersen, R. A. Memory related motor planning activity in posterior
parietal cortex of macaque. *Exp Brain Res* **70**, 216-220 (1988).
https://doi.org:10.1007/BF00271862

26   Hikosaka, O. & Wurtz, R. H. Visual and oculomotor functions of monkey substantia
nigra pars reticulata. III. Memory-contingent visual and saccade responses. *J
Neurophysiol* **49**, 1268-1284 (1983).

27   Powers, A. S., Basso, M. A. & Evinger, C. Blinks slow memory-guided saccades. *J
Neurophysiol* **109**, 734-741 (2013). https://doi.org:10.1152/jn.00746.2012

28   Smit, A. C., Van Gisbergen, J. A. & Cools, A. R. A parametric analysis of human
saccades in different experimental paradigms. *Vision Res* **27**, 1745-1762 (1987).
https://doi.org:10.1016/0042-6989(87)90104-0

29   Stanford, T. R. & Sparks, D. L. Systematic errors for saccades to remembered targets:
evidence for a dissociation between saccade metrics and activity in the superior
colliculus. *Vision Res* **34**, 93-106 (1994).

30   White, J. M., Sparks, D. L. & Stanford, T. R. Saccades to remembered target locations:
an analysis of systematic and variable errors. *Vision Res* **34**, 79-92 (1994).
https://doi.org:10.1016/0042-6989(94)90259-3

31   Mohler, C. W. & Wurtz, R. H. Organization of monkey superior colliculus:
intermediate layer cells discharging before eye movements. *J Neurophysiol* **39**, 722-
744 (1976).

32   Sommer, M. A. & Wurtz, R. H. What the brain stem tells the frontal cortex. I.
Oculomotor signals sent from superior colliculus to frontal eye field via mediodorsal
thalamus. *J Neurophysiol* **91**, 1381-1402 (2004).
https://doi.org:10.1152/jn.00738.2003
00738.2003 [pii]

33   Chen, C. Y., Hoffmann, K. P., Distler, C. & Hafed, Z. M. The Foveal Visual
Representation of the Primate Superior Colliculus. *Curr Biol* **29**, 2109-2119 e2107
(2019). https://doi.org:10.1016/j.cub.2019.05.040

34   Zuber, B. L., Stark, L. & Cook, G. Microsaccades and the velocity-amplitude
relationship for saccadic eye movements. *Science* **150**, 1459-1460 (1965).

35   Bahill, A. T., Clark, M. R. & Stark, L. The main sequence, a tool for studying human eye
movements. *Mathematical Biosciences* **24**, 191-204 (1975).

36   Hafed, Z. M. & Goffart, L. Gaze direction as equilibrium: more evidence from spatial
and temporal aspects of small-saccade triggering in the rhesus macaque monkey. *J*
Neurophysiol **123**, 308-322 (2020). https://doi.org:10.1152/jn.00588.2019

37   Willeke, K. F. *et al.* Memory-guided microsaccades. *Nat Commun* **10**, 3710 (2019).
https://doi.org:10.1038/s41467-019-11711-x

38   Greenwood, J. A., Szinte, M., Sayim, B. & Cavanagh, P. Variations in crowding,
saccadic precision, and spatial localization reveal the shared topology of spatial
vision. *Proc Natl Acad Sci U S A*, E3573-E3582 (2017).
https://doi.org:www.pnas.org/cgi/doi/10.1073/pnas.1615504114

39   Schlykowa, L., Hoffmann, K. P., Bremmer, F., Thiele, A. & Ehrenstein, W. H. Monkey
saccadic latency and pursuit velocity show a preference for upward directions of
target motion. *Neuroreport* **7**, 409-412 (1996).

40   Zhou, W. & King, W. M. Attentional sensitivity and asymmetries of vertical saccade
generation in monkey. *Vision Res* **42**, 771-779 (2002).

41   Greene, H. H., Brown, J. M. & Dauphin, B. When do you look where you look? A visual
field asymmetry. *Vision Res* **102**, 33-40 (2014).
https://doi.org:10.1016/j.visres.2014.07.012

42   Krauzlis, R. J., Lovejoy, L. P. & Zenon, A. Superior colliculus and visual spatial
attention. *Annu Rev Neurosci* **36**, 165-182 (2013). https://doi.org:10.1146/annurev-
neuro-062012-170249

43   Basso, M. A. & May, P. J. Circuits for Action and Cognition: A View from the Superior
Colliculus. *Annu Rev Vis Sci* (2017). https://doi.org:10.1146/annurev-vision-102016-
061234

44   Lovejoy, L. P. & Krauzlis, R. J. Inactivation of primate superior colliculus impairs covert
selection of signals for perceptual judgments. *Nat Neurosci* **13**, 261-266 (2010).
https://doi.org:nn.2470 [pii]
10.1038/nn.2470

45   Zenon, Z. & Krauzlis, R. J. Attention deficits without cortical neuronal deficits. *Nature*
489, 434-437 (2012).

46   Bogadhi, A. R., Katz, L. N., Bollimunta, A., Leopold, D. A. & Krauzlis, R. J. Midbrain
activity shapes high-level visual properties in the primate temporal cortex. *Neuron*
109, 690-699 e695 (2021). https://doi.org:10.1016/j.neuron.2020.11.023

47   Horwitz, G. D. & Newsome, W. T. Separate signals for target selection and movement
specification in the superior colliculus. *Science* **284**, 1158-1161 (1999).
https://doi.org:10.1126/science.284.5417.1158

48 McPeek, R. M. & Keller, E. L. Deficits in saccade target selection after inactivation of
superior colliculus. *Nat Neurosci* **7**, 757-763 (2004). https://doi.org:10.1038/nn1269
nn1269 [pii]

49 Krauzlis, R. J., Liston, D. & Carello, C. D. Target selection and the superior colliculus:
goals, choices and hypotheses. *Vision Res* **44**, 1445-1451 (2004).
https://doi.org:10.1016/j.visres.2004.01.005
S0042698904000264 [pii]

50 Barbot, A., Xue, S. & Carrasco, M. Asymmetries in visual acuity around the visual
field. *J Vis* **21**, 2 (2021). https://doi.org:10.1167/jov.21.1.2

51 Himmelberg, M. M., Winawer, J. & Carrasco, M. Stimulus-dependent contrast
sensitivity asymmetries around the visual field. *J Vis* **20**, 18 (2020).
https://doi.org:10.1167/jov.20.9.18

52 Talgar, C. P. & Carrasco, M. Vertical meridian asymmetry in spatial resolution: visual
and attentional factors. *Psychonomic bulletin & review* **9**, 714-722 (2002).

53 Montaser-Kouhsari, L. & Carrasco, M. Perceptual asymmetries are preserved in short-
term memory tasks. *Attention, perception & psychophysics* **71**, 1782-1792 (2009).
https://doi.org:10.3758/APP.71.8.1782

54 He, S., Cavanagh, P. & Intriligator, J. Attentional resolution and the locus of visual
awareness. *Nature* **383**, 334-337 (1996). https://doi.org:10.1038/383334a0

55 Rubin, N., Nakayama, K. & Shapley, R. Enhanced perception of illusory contours in
the lower versus upper visual hemifields. *Science* **271**, 651-653 (1996).

56 Grujic, N., Brehm, N., Gloge, C., Zhuo, W. & Hafed, Z. M. Peri-saccadic perceptual
mislocalization is different for upward saccades. *J Neurophysiol* (2018).
https://doi.org:10.1152/jn.00350.2018

57 Hafed, Z. M. & Krauzlis, R. J. Microsaccadic suppression of visual bursts in the primate
superior colliculus. *J Neurosci* **30**, 9542-9547 (2010). https://doi.org:30/28/9542 [pii]
10.1523/JNEUROSCI.1137-10.2010

58 Fracasso, A., Buonocore, A. & Hafed, Z. M. Peri-saccadic visual sensitivity is higher in
the upper visual field. *bioRxiv* (2022). https://doi.org:10.1101/2022.07.05.498850

59 Berman, R. A., Joiner, W. M., Cavanaugh, J. & Wurtz, R. H. Modulation of presaccadic
activity in the frontal eye field by the superior colliculus. *J Neurophysiol* **101**, 2934-
2942 (2009). https://doi.org:00053.2009 [pii]
10.1152/jn.00053.2009

60 Shin, S. & Sommer, M. A. Division of labor in frontal eye field neurons during
presaccadic remapping of visual receptive fields. *J Neurophysiol* **108**, 2144-2159
(2012). https://doi.org:10.1152/jn.00204.2012

61 Chen, C. Y. & Hafed, Z. M. Postmicrosaccadic enhancement of slow eye movements.
The Journal of neuroscience : the official journal of the Society for Neuroscience **33**,
5375-5386 (2013). https://doi.org:10.1523/JNEUROSCI.3703-12.2013

62 Hafed, Z. M. & Ignashchenkova, A. On the dissociation between microsaccade rate
and direction after peripheral cues: microsaccadic inhibition revisited. *J Neurosci* **33**,
16220-16235 (2013). https://doi.org:10.1523/JNEUROSCI.2240-13.2013

63 Chen, C. Y., Ignashchenkova, A., Thier, P. & Hafed, Z. M. Neuronal Response Gain
Enhancement prior to Microsaccades. *Curr Biol* **25**, 2065-2074 (2015).
https://doi.org:10.1016/j.cub.2015.06.022

64 Fuchs, A. F. & Robinson, D. A. A method for measuring horizontal and vertical eye
movement chronically in the monkey. *J Appl Physiol* **21**, 1068-1070 (1966).

65 Judge, S. J., Richmond, B. J. & Chu, F. C. Implantation of magnetic search coils for
measurement of eye position: an improved method. *Vision Res* **20**, 535-538 (1980).

66 Malevich, T., Zhang, T., Baumann, M. P., Bogadhi, A. R. & Hafed, Z. M. Faster
detection of "darks" than "brights" by monkey superior colliculus neurons. *BioRxiv*
(2022). https://doi.org:10.1101/2022.08.03.502615

67 Chen, C. Y., Sonnenberg, L., Weller, S., Witschel, T. & Hafed, Z. M. Spatial frequency
sensitivity in macaque midbrain. *Nat Commun* **9**, 2852 (2018).
https://doi.org:10.1038/s41467-018-05302-5

68 Chen, C. Y. & Hafed, Z. M. Orientation and Contrast Tuning Properties and Temporal
Flicker Fusion Characteristics of Primate Superior Colliculus Neurons. *Front Neural*
Circuits **12**, 58 (2018). https://doi.org:10.3389/fncir.2018.00058

69 Bellet, M. E., Bellet, J., Nienborg, H., Hafed, Z. M. & Berens, P. Human-level saccade
detection performance using deep neural networks. *J Neurophysiol* **121**, 646-661
(2019). https://doi.org:10.1152/jn.00601.2018

70 Pachitariu, M., Steinmetz, N. A., Kadir, S. N., Carandini, M. & Harris, K. D. Fast and
accurate spike sorting of high-channel count probes with KiloSort. *Advances in Neural*
Information Processing Systems (NIPS 2016) **29** (2016).
1249
1250

Acknowledgements

We were funded by the Deutsche Forschungsgemeinschaft (DFG) through Research Unit:
FOR1847 (project: A6: HA6749/2-1). The following DFG-funded projects also contributed to
the current study: HA6749/3-1, HA6749/4-1, and SFB1233, Robust Vision: Inference
Principles and Neural Mechanisms (TP 11, project number: 276693517).

Author contributions

ZMH collected and analyzed datasets 1 and 2. TZ, TM, MPB, and ZMH collected and analyzed
dataset 3. All authors wrote the manuscript.

Declaration of interests

The authors declare no competing interests.