Machine Learning based Genome-Wide Association Studies for Uncovering QTL Underlying Soybean Yield and its Components

Mohsen Yoosefzadeh-Najafabadi,[1] Sepideh Torabi,[1] Davoud Torkamaneh,[2] Dan Tulpan,[3] Istvan Rajcan,[1] and Milad Eskandari[1*]

[1.] Department of Plant Agriculture, University of Guelph, Guelph, ON N1G 2W1, Canada

[2.] Département de Phytologie, Université Laval, Québec City, QC G1V 0A6, Canada

[3.] Department of Animal Biosciences, University of Guelph, Guelph, ON N1G 2W1, Canada

[*]Corresponding author: Email: meskanda@uoguelph.ca

Highlight

Implementing sophisticated mathematical approaches such as machine learning algorithms in GWAS can simultaneously consider a wide range of interconnected biological processes and mechanisms that shape the phenotype of complex traits such as yield and its components in soybean.

Abstract

Genome-wide association study (GWAS) is currently one of the important approaches for discovering quantitative trait loci (QTL) associated with traits of interest. However, insufficient statistical power is the limiting factor in current conventional GWAS methods for characterizing quantitative traits, especially in narrow genetic bases plants such as soybean. In this study, we evaluated the potential use of machine learning (ML) algorithms such as support vector machine (SVR) and random forest (RF) in GWAS, compared with two conventional methods of mixed linear models (MLM) and fixed and random model circulating probability unification (FarmCPU), for identifying QTL associated with soybean yield components. In this study, important soybean yield component traits, including the number of reproductive nodes (RNP), non-reproductive nodes (NRNP), total nodes (NP), and total pods (PP) per plant along with yield and maturity were assessed using 227 soybean genotypes evaluated across four environments. Our results indicated SVR-mediated GWAS outperformed RF, MLM and FarmCPU in discovering the most relevant QTL associated with the traits, supported by the functional annotation of candidate gene analyses. This study for the first time demonstrated the potential

31 benefit of using sophisticated mathematical approaches such as ML algorithms in GWAS for
32 identifying QTL suitable for genomic-based breeding programs.

33 **Keywords:** Data-driven Models; FarmCPU; Genome-wide association study; MLM; Soybean
34 Breeding; Support vector machine.

Introduction

Soybean (*Glycine max* [L.] Merr.) is known as one of the most important legume crops with
substantial economic value (Rębilas *et al.*, 2020). Soybean is widely used for food, feed, fiber,
biodiesel, and green manure (Temesgen and Assefa, 2020). Despite the importance of genetic
improvement in soybean yield, the soybean germplasm has in general a narrow genetic basis,
especially within North American germplasm, which has resulted in limited enhancement of the
genetic gain, historically (Xavier and Rainey, 2020). Therefore, there is a great need for
analytical breeding to explore the optimum genetic potential of soybean (Mangena, 2020; Suhre
et al., 2014).

Analytical breeding strategy as an alternate breeding approach requires a better understanding of
the factors, or individual traits, responsible for the development, growth, and yield (Richards,
1982). This strategy considers highly correlated secondary traits with the trait of interest as the
selection criteria that can make empirical selection more efficient for improving the genetic gain
(Reynolds, 2001; Richards, 1982; Xavier and Rainey, 2020). The application of the analytical
approaches in plant breeding programs has been limited due mainly to lack of sufficient
resources, as they are time and labor-consuming (Richards, 1982; Xavier *et al.*, 2018). Therefore,
breeders are restricted to evaluating secondary traits in a small number of genotypes, which
results in the limitation of the knowledge in the genome-to-phenome analysis process (Kahlon *et*
al., 2011; Nico *et al.*, 2019; Robinson *et al.*, 2009).

Yield potential in soybean is mainly determined by the following yield component traits: the
total number of pods, nodes, reproductive nodes, non-reproductive nodes, and pods per plant
(Pedersen and Lauer, 2004; Reynolds, 2001; Xavier *et al.*, 2018; Xavier and Rainey, 2020;
Yoosefzadeh-Najafabadi *et al.*, 2021b). Of these, the total number of nodes and pods play more
important roles in seed yield production than other yield components (Robinson *et al.*, 2009;
Yoosefzadeh-Najafabadi *et al.*, 2021b). Several studies reported a steady increase in the total
number of nodes and the total number of pods in soybean cultivars from 1920 to 2010 (Kahlon *et*
al., 2011; Suhre *et al.*, 2014; Xavier and Rainey, 2020). These findings may highlight the
importance and potential use of the phenotypic and genotypic information on these traits, along
with yield per se, as selection criteria in cultivar development programs (Ma *et al.*, 2001).

Genetic studies of soybean yield component traits can accelerate the breeding process more
accurately (Xavier and Rainey, 2020). Genome-Wide Association Studies (GWAS), as one of
the common genetic approaches, can be implemented on diverse populations to detect the
quantitative trait loci (QTL) associated with the soybean yield component traits (Kaler *et al.*,
2020). Associated QTL can be used for screening large soybean populations in a short time with
less elaborate efforts (Xavier *et al.*, 2018). Several GWAS algorithms have been developed for
genetic studies, such as mixed linear models (MLM), multiple loci linear mixed model
(MLMM), and fixed and random model circulating probability unification (FarmCPU) (Kaler *et*

al., 2020). However, due to the narrow genetic base of some plant species, including soybean, the conventional approaches may not have enough statistical power to detect reliable QTL (Kaler *et al.*, 2020; Mohammadi *et al.*, 2020; Xavier and Rainey, 2020). Therefore, the development of more sophisticated statistical methods is required in order to establish effective GWAS methods for plant species with a narrow genetic base.

Current GWAS methods are based on the conventional statistical methods that are useful for studying less complex traits in plant species with broader genetic bases (Lipka *et al.*, 2015; Pasaniuc and Price, 2017). Machine learning (ML) algorithms as powerful and reliable mathematical methods can be considered as an alternative to conventional statistical methods for performing GWAS, which are efficient for studying more complex traits in plants with narrow genetic base (Xavier and Rainey, 2020). Recently, the use of ML algorithms has been reported in different areas such as plant science (Hesami *et al.*, 2020; Yoosefzadeh-Najafabadi *et al.*, 2021a), animal science (Tulpan, 2020), human science (Chen and Verghese, 2020), engineering (Kim *et al.*, 2020), and computer science (Jordan and Mitchell, 2015). The application of ML algorithms in GWAS was previously investigated in humans by Szymczak *et al.* (2009). They explained a possible use of different ML algorithms such as artificial neural networks (ANN), Bayesian network analysis (BNA), and random forests (RF) in GWAS for human disease studies (Szymczak *et al.*, 2009). One of the most common used ML algorithms is RF developed by Breiman (2001), which generates a series of trees from the independent samples for better prediction performance (Meinshausen, 2006). The latter algorithm has been widely used in plant genomics (Ogutu *et al.*, 2011), phenomics (Yoosefzadeh-Najafabadi *et al.*, 2021a), proteomics (Jamil *et al.*, 2020), and metabolomics (Sun *et al.*, 2020).

The first and only use of the RF-mediated GWAS in soybean, for detecting the genomics association in soybean yield component traits, was reported by Xavier and Rainey (2020). Support vector machine (SVM) is another common algorithm that can detect behavior and patterns of nonlinear relationships (Auria and Moro, 2008; Hesami and Jones, 2020; Su *et al.*, 2017). Theoretically, SVM should have high performance due to the use of structural risk minimization instead of the empirical risk minimization inductive principles (Belayneh *et al.*, 2014; Yoosefzadeh-Najafabadi *et al.*, 2021a). There is a significant number of reports on the successful using of SVM in prediction problems (Denton and Salleb-Aouissi, 2020; Duan *et al.*, 2005; Hesami *et al.*, 2020; Tulpan, 2020; Yoosefzadeh-Najafabadi *et al.*, 2021a). Support vector regression (SVR) is known as the regression version of SVM that commonly used for continuous dataset. There are also reports on the successful use of SVR for addressing plant prediction problems (Awad and Khanna, 2015). However, the possible use of SVR in GWAS is still unexplored in plant science area.

In this study we aimed to: (1) gain a better understanding of the genetic relationships between soybean yield and its component traits, and (2) investigate the potential use of RF and SVM algorithms in GWAS for discovering QTL underlying soybean yield components as compared to conventional GWAS methods of MLM and FarmCPU. The results of this study will help soybean breeders to have a better perspective of exploiting ML algorithms in GWAS studies, and may offer them new genomic tools for screening high yielding genotypes with improved genetic gain based on genomic regions associated with yield components.

Materials and Methods

Population and experimental design

An GWAS panel of 250 soybean genotypes was grown at the University of Guelph, Ridgetown Campus in two locations, Palmyra (42°25'50.1"N 81°45'06.9"W, 195 m above sea level) and Ridgetown (42°27'14.8"N 81°52'48.0"W, 200m above sea level) in Ontario, Canada, in two consecutive years, 2018 and 2019. The panel used in this study consisted of the main germplasm of the soybean breeding program at the University of Guelph, Ridgetown Campus, that has been established over 35 years for cultivar development and genetic studies. The randomized complete block design (RCBD) with two replications was used for all four environments. In general, there were 500 and 1000 research plots per environment and year, respectively. Each plot consisted of five 4.2 m long rows with 57 seeds per m^2 seeding rate.

Phenotyping

In this experiment, soybean seed yield (t ha^{-1} at 13% moisture) for each plot was estimated by harvesting three middle rows. Soybean seed yield components, including the total number of reproductive nodes per plant (RNP), the total number of non-reproductive nodes per plant (NRNP), the total nodes per plant (NP), and the total number of pods per plant (PP), were measured using 10 randomly selected plants from each plot. The maturity was recorded as the number of days from planting to physiological maturity (R7, (Fehr and Caviness, 1971) for each genotype.

Genotyping

Young trifoliate leaf tissue for each soybean genotype from the first replication of the trail at the Ridgetown in 2018, were collected and in a 2 mL screw-cap tube. The leaf samples were freeze-dried for 72 hours, using the Savant ModulyoD Thermoquest (Savant Instruments, Holbrook, NY). By using the DNA Extraction Kit (SIGMA®, Saint Louis, MO), DNA was extracted for soybean genotypes, and the quantity of DNAs was checked via Qubit® 2.0 fluorometer (Invitrogen, Carlsbad, CA). For genotyping-by-sequencing (GBS), DNA samples were sent to Plate-forme D'analyses Génomiques at Université Laval (Laval, Quebec, Canada). The GWAS panel was genotyped via a GBS protocol based on the enzymatic digestion with *ApeKI* (Sonah *et al.*, 2013). Single-nucleotide polymorphisms (SNPs) were called by the Fast GBS pipeline (Torkamaneh *et al.*, 2020), using Gmax_275_v2 reference genome. Markov model was used to impute the missing loci, and SNPS with a minor allele frequency (MAF) less than 0.05 were removed below the threshold. In total, after checking the quality of reading sequence and removing SNPs with more than 50% heterozygosity, 23 genotypes were eliminated from the experiment and 17,958 high-quality SNPs from 227 soybean genotypes used for genetic analysis.

Statistical analyses

The best linear unbiased prediction (BLUP) as one of the common linear mixed models (Goldberger, 1962) was used to estimate the genetic values of each soybean genotype. Also, R package *lme4* (Bates *et al.*, 2014) was used to analyze yield and yield components with 'environment' as a fixed effect and 'genotype' as a random effect. To control for the possible

soil heterogeneity among the plots within a given block and reduce the associated experimental errors, nearest-neighbor analyses (NNA) was used as one of the common error control methods (Bowley, 1999; Katsileros *et al.*, 2015; Stroup and Mulitze, 1991). Outliers were determined in the raw dataset based on the protocols proposed by Bowley (1999) and treated the same as missing data points in the analysis. Overall, the following statistical model was used in this study:

$$Y_{ij} = \mu + f(s) + G_i + E_j + GE_{ij} + \varepsilon_{ij} , i = 1, ..., k; j = 1, ..., n \quad (Eq\ 5.1)$$

Where Y_{ij} stands for the trait of interest (soybean seed yield and yield component traits) as a function of an intercept μ, f(s) stands for the spatial covariate, G_i is the random genotype effect, E_j stands for the fixed environment effect, GE_{ij} is the genotype x environment interaction effect, and ε_{ij} stands for the residual effect.

The heritability was calculated for soybean seed yield and yield components using *lme4* open-source R package (Bates *et al.*, 2007) based on the following equation:

$$H^2 = \frac{\sigma^2_G}{\sigma^2_G + \sigma^2_E} \quad (Eq\ 5.2)$$

where σ^2_G stands for the genotypic variance, and σ^2_E is the environmental variance.

Analysis of population structure

A total of 17,958 high-quality SNPs from 227 soybean genotypes were used to conduct population structure analysis using fastSTRUCTURE (Raj *et al.*, 2014). Five runs were conducted for K set from 1 and 15 to estimate the most appropriate number of subpopulations by using the K tool from the fastSTRUCTURE software.

Association studies

Since different GWAS methods may capture different genomic regions (Yang *et al.*, 2018). Therefore, MLM and FarmCPU (two most common GWAS methods) and RF and SVM (two most common machine learning algorithms) were used in this study. MLM and FarmCPU were implemented by using *GAPIT* package (Lipka *et al.*, 2012), and RF, as well as SVM, were conducted through the *Caret* package (Kuhn *et al.*, 2020) in R software version 3.6.1. A brief description of each of the GWAS methods is provided below:

Mixed Linear Model (MLM): This GWAS is based on the likelihood ratio between the full model, consisting of the marker of interest, and the reduced model, which is known as the model without the marker of interest (Wen *et al.*, 2018).

Fixed and random model circulating probability unification (FarmCPU): This GWAS takes the advantages of using MLM as the random model, and stepwise regression as the fixed model iteratively (Liu *et al.*, 2016). False discovery rate (FDR) is used for setting the threshold both in the FarmCPU and MLM models (Benjamini and Hochberg, 1995).

Random Forest (RF): This machine-learning algorithm was first implemented by Xavier and
Rainey (2020) in a soybean GWAS study. This method is known as the powerful non-parametric
regression approach that is derived from aggregating the bootstrapping in various decision trees
(Breiman, 2001). In this experiment, a 1000-set of decision trees constructed the forest, and the
GWAS analysis was done by measuring the importance of each feature (Botta *et al.*, 2014),
which was an SNP in this study.

Support vector regression (SVR): This machine learning algorithm is known as one of the
common supervised learning methods in prediction problems (Cortes and Vapnik, 1995). This
algorithm is based on constructing a set of hyperplanes that can be useful in regression problems
(Fletcher, 2009). The association statistics in this algorithm can be achieved by estimating the
feature importance that was previously proposed by Weston *et al.* (2001). In this experiment,
SNP markers were selected as inputs, and the traits were selected as target variables for
estimating the feature importance.

Variable Importance measurement

As one of the common indices for tree-based algorithms, the impurity index was chosen as the
metric of the feature importance for the RF algorithm. Regarding the SVR algorithm, the
variable importance method for SVR Weston *et al.* (2001) was implemented in this dataset. For
both algorithms, the importance of each SNP was scaled based on 0 to 100 percent scale. Since
there is no confirmed way of defining the significant threshold in the tested algorithms, the
global empirical threshold that provides the empirical distribution of the null hypothesis
(Churchill and Doerge, 1994; Doerge and Churchill, 1996) was used for establishing threshold in
this study. The global empirical threshold was estimated based on fitting the ML algorithm,
storing the highest variable importance, repeating 1000 times, and select the SNPs based on
α=0.05.

Data-driven model processes

In order to estimate the feature importance in RF and SVR algorithms, a five-fold cross-
validation strategy (Siegmann and Jarmer, 2015) with ten repetitions was applied on the dataset.
All of the tested machine learning algorithms were optimized for their parameters for this dataset
accordingly.

Functional annotation of candidate SNPs

For each tested GWAS model, the flanking regions of each QTL was determined using LD decay
distance (Fig.1), and then potential candidate genes were retrieved using the *G. max* cv. William
82 reference genome, gene models 2.0 in SoyBase (https://www.soybase.org). After listing
potential candidate genes in defined windows around each significant SNP, at the peak of each
QTL, Gene Ontology annotation, GO term enrichment (https://www.soybase.org), and the report
from previous studies were used as the criteria to select and report the most relevant candidate
genes associated with the identified QTL. The Electronic Fluorescent Pictograph (eFP) browser
for soybean (www.bar.utoronto.ca) was also used to generate additional information such as
tissue- and developmental-stage dependent expression (based on transcriptomic data from
Severin *et al.* (2010)) for the identified candidate genes.

Visualization

All of the visualizations in this study were conducted using the *ggplot2* package (Wickham, 2011) in R version 3.6.1 software and Microsoft Excel software (2016).

Results

Phenotyping evaluations

The tested GWAS panel of 227 soybean genotypes showed significant variations among the genotypes for seed yield, maturity, and yield component traits. The distribution of the phenotypic measures for the traits across the four environments is presented in Fig. 2. The highest heritability was observed for maturity with an estimate of 0.78 followed by 0.34, 0.33, 0.31, and 0.30 for NP, RNP, NRNP, and PP, respectively (Fig. 2). The lowest heritability was estimated for yield with a value of 0.24 (Fig. 2). Soybean seed yield and PP showed the highest variability across the environments (Fig. 2).

The linear correlations among all the measured traits were estimated using the coefficients of correlation (r). Based on the results (Fig. 3), all traits were positively correlated with each other, except the NRNP that was negatively associated with yield, maturity, RNP, NP, and PP. NP showed the highest correlation with the RNP ($r= 0.97$) and NRNP ($r= -0.63$). RNP had the highest correlation ($r =0.86$) with yield among all the tested yield components (Fig. 3).

Genotyping evaluations

For the tested GWAS panel, high-quality SNPs were obtained from 210M single-end Ion Torrent reads that were proceeded with Fast-GBS.v2. From a total of 40,712 SNPs, 17,958 SNPs were polymorphic and mapped to 20 soybean chromosomes. The minimum and maximum number of SNPs were 403 and 1780 on chromosomes 11 and 18, respectively. Overall, the average number of SNPs across all the 20 chromosomes was 898, with the mean density of one SNP for every 0.12 cM across the genome.

Population structure and kinship

The structure profile for the tested population is presented in Fig. 4. The result of genotypic evaluations suggested that the tested GWAS panel was composed of four to seven subpopulations. Therefore, we chose to conduct the structure analysis using K=7 as the appropriate K for the structure profile of the tested GWAS panel (Fig. 4). In order to reduce the confounding, the kinship was also estimated between genotypes of the GWAS panel.

GWAS analysis

The average value for soybean maturity in the tested GWAS panel was 106 days with a standard deviation of 5 days (Fig. 3). Association analysis by the MLM method identified nine associated SNP markers located on chromosomes 2 and 19 (Fig. 5A). Using FarmCPU, a total of nine associated SNP markers were located on chromosomes 2, 19, and 20 (Fig. 5A). By using the RF method, the total of three SNP markers on chromosomes 3, 7, and 16 were associated with the

soybean maturity, whereas SVR-mediated GWAS detected 10 associated SNP markers located
on chromosomes 2, 6, 10, 16, 19, and 20 (Fig. 5A).

SVR-mediated GWAS detected five QTL directly related to the reproductive period and R8 full
maturity (Table 1). The average soybean seed yield in the GWAS panel was 3.5 t ha^{-1} with a
standard deviation of 0.45 (Fig. 3). Using MLM, FarmCPU, RF, and SVR approach, we
identified two, three, five, and 18 SNP markers associated with the yield, respectively (Fig. 5B).
The SNP markers identified by MLM and FarmCPU were located on chromosomes 6 and 8.
Using the RF-mediated GWAS method, associated SNP markers were located on chromosomes
4, 7, 12, and 17. By using the SVR-mediated GWAS method, the SNP markers were located on
chromosomes 3, 4, 6, 7, 15, 19, and 20 (Fig. 5B). In SVR-mediated GWAS, the identified QTL
were co-localized with eight previously reported related QTL such as seed yield, seed weight,
and seed set (Table 2). However, other tested GWAS methods could not co-localized with any
QTL associated with seed yield (Table 2).

The average NP in the tested GWAS panel was 15.21 nodes with a standard deviation of 0.77
nodes (Fig. 3). By using the MLM and FarmCPU methods, one and two associated SNP markers
were detected, respectively (Fig. 6A). Four and ten associated SNP markers were detected by NP
using RF and SVR methods, respectively. SVR-mediated GWAS was the only method that were
co-localized with three previously reported NP-related QTL (Table 3). The average NRNP was
3.33 nodes with a standard deviation of 0.28 nodes (Fig. 3). A total of two, three, five, and ten
associated SNP markers were detected using the MLM, FarmCPU, RF, and SVR methods,
respectively (Fig. 6B). The detected SNP markers using the SVR method were located on
chromosomes 4, 7, 18, 19, and 20, whereas SNP markers identified through RF were located on
chromosomes 1, 4, 7, 18, and 19 (Fig. 6B). Chromosomes number 4, 8, and 15 were identified as
carrying SNP markers with NRNP using FarmCPU. The MLM method identified SNP markers
located on chromosomes 8 and 15, which most of the detected QTL co-localized with previously
reported QTL related to seed weight, seed protein, water use efficiency, first flower, and soybean
cyst nematode (Table 4).

The average RNP was 11.89 nodes with a standard deviation of 0.98 nodes (Fig. 3). Based on the
results of MLM and FarmCPU methods, four associated SNP markers with RNP were located on
chromosomes 8 and 19. Using the RF method, four associated SNP markers were identified on
chromosomes 8, 9, 15, and 20. Using the SVR method, 11 SNP markers were associated with
RNP located on chromosomes 4, 7, 8, 15, 18, 19, and 20 (Fig. 7A). Regardless of the type of
GWAS methods used in this study, we found SNP markers associated with the trait on
chromosome 8. The position of the associated SNP marker on chromosome 8 was identical both
in SVR and RF (462.3 Kbp) and MLM and FarmCPU (481.6 Kbp). The list of detected QTL for
RNP is presented in Table 5. The average value for PP in the tested GWAS panel was 45.02 pods
with a standard deviation of 8.54 pods. We did not detect any SNP marker associated with PP
using the MLM and FarmCPU methods. However, by using the RF method, four SNP markers
were found to be associated with PP and located on chromosomes 7, 10, 18, and 20 (Fig. 7B).
Twelve associated SNP markers were found by SVR that were located on chromosomes 6, 9, 10,
11, 15, 18, and 19 (Fig. 7B). The GWAS of chromosome 10 with PP were found both in RF and
SVR with 4.6 cM distance far from each other. In PP, MLM and FarmCPU did not detect any

related QTL for this trait, while SVR-mediated GWAS was identified seven QTL directly related
to the pod number (Table 6).

Identification of candidate genes within QTL

According to the flanking regions of each QTL which was determined using LD decay distance,
150-kbp upstream and downstream of each SNP's peak were considered to identify potential
candidate genes (Fig. 1). Candidate genes were extracted for each significant peak SNP with
high allelic effect and based on the gene annotation, enrichment tools and previous studies
(Table S1). For maturity, three peak SNPs (Chr2_695362, Chr2_720134, and Chr19_47513536)
had the highest allelic effect than other detected peak SNPs (Fig. 8A). On the basis of the gene
annotation and expression within QTL, *Glyma.02g006500* (GO:0015996) and *Glyma.19g224200*
(GO:0010201) were identified as the strong candidate genes for maturity, which encode
chlorophyll catabolic process and phytochrome A (PHYA) related genes, respectively.
Glyma.02g006500 (GO:0015996) was exactly detected in the peak SNP position of
Chr2_695362, whereas *Glyma.19g224200* (GO:0010201) was 119 Kbp far from the detected
peak SNP at Chr19_47513536. In yield, the peak SNP with the position of Chr7_1032587 had
the highest allelic effect in comparison with other detected peak SNPs (Fig. 8B). Within a 77
Kbp above from the detected peak SNP (Chr7_1032587), *Glyma.07G014100* (GO:0010817) was
identified, which encodes the regulation of hormone levels, as the strongest candidate genes in
yield. For NP, two peak SNPs (Chr7_1032587 and Chr7_1092403) had the highest allelic effect
among all detected peak SNPs (Fig. 8C). SNP peak position of Chr7_1032587 was detected in
common for yield, NP, and NRNP. *Glyma.07G205500* (GO:0009693) and *Glyma.08G065300*
(GO:0042546) were detected as the strongest candidate genes both in NP and NRNP, which
encode UBP1-associated protein 2C and cell wall biogenesis, respectively. Both detected gene
candidates were exactly at the associated peak SNPs at Chr7_1032587 and Chr8_5005929 (Fig.
8D). Regarding peak SNPs associated with RNP, the highest allelic effects were found in peak
SNPs of Chr9_40285014 and Chr15_34958361 (Fig. 8E). *Glyma.15G214600* (GO:0009920) and
Glyma.15G214700 (GO:0009910), which encode cell plate formation involved in plant-type cell
wall biogenesis and acetyl-CoA biosynthetic process, as strong candidate genes in NRNP.
Glyma.15G214600 (GO:0009920) and *Glyma.15G214700* (GO:0009910) were 127 and 90 Kbp
far from the detected peak SNP at Chr15_3495836, respectively. In PP, the highest allelic effects
were found in peak SNPs at Chr7_15331676, Chr11_5245870, and Chr18_55469601 (Fig. 8F).
Glyma.07G128100 (GO:0009909) was the strongest candidate genes for PP, which encodes
regulation of flower development. *Glyma.07G128100* (GO:0009909) was detected exactly in the
peak SNP position of Chr7_15331676.

Discussion

One of the objectives of this study was to gain a better understanding of the roles of soybean
yield component traits in the production of total seed yield and how these traits can be used for
facilitating the development of high-yielding soybeans. The genetic dissection of soybean yield
component traits in order to develop genetic and genomics toolkits can be useful for designing
breeding population and selection criteria aiming at improving yield genetic gains in new
cultivars (Cooper *et al.*, 2009; Hu *et al.*, 2020; Xavier and Rainey, 2020). For this aim, a wide
range of analyses, including Pearson correlation, normality and distribution plots, GWAS both in

combined and separate environments, and functional annotation of candidate genes and QTL, were performed in this study. The collective evaluation of the mentioned analysis contributes to building the wide perspectives of the genetic architecture of the soybean yield component traits. One of the important factors for genetic studies is to evaluate the phenotypic variation within genotypes and environments. High phenotypic variation was observed for yield and PP, while maturity and NP had the lowest phenotypic variation across the tested environments. These findings are in line with the results of previous research on yield component traits (Kahlon and Board, 2012; Xavier and Rainey, 2020). The heritability and correlation analyses showed that NP had the highest heritability and significant linear correlations with RNP and PP. Also, PP had the highest correlation with yield among all the tested soybean yield components. The number of nodes and pods in soybean are known as the two of the key soybean yield components that play an important role in determining the final soybean seed yield (Herbert and Litchfield, 1982; Kahlon and Board, 2012; Xavier and Rainey, 2020). However, studies showed the low heritability rates for soybean yield components, especially NP and PP (KUSWANTORO, 2017; Sulistyo and Sari, 2018; Xavier *et al.*, 2016a; Xavier and Rainey, 2020). The nature of these traits can explain low heritability rates as they are mostly affected by environmental factors (Price and Schluter, 1991). Although heritability indicates the strength of the relationship between phenotype and genetic variation of the particular trait, it does not indicate the value of the trait for genetic study (Cassell, 2009). Different low heritable traits are highly correlated with significant economic traits (Cassell, 2009). In soybean, yield can be considered as the most important economic trait that is highly determined by yield components. Therefore, any genetic and environmental studies around yield components can open the possibility of overall yield improvement in major crops such as soybean.

GWAS is known as one of the most important genetic toolkits for detecting QTL associated with quantitate traits (Kaler *et al.*, 2020). There are several statistical methods implemented in GWAS for improving the detection of associated SNP markers with the trait of interest. While conventional GWAS are appropriate approaches for detecting SNP markers with large effects on complex traits, they are, however, underpowered for the simultaneous consideration of a wide range of interconnected biological processes and mechanisms that shape the phenotype of complex traits (Lee *et al.*, 2020). Therefore, using variable importance values in ML algorithms for identifying SNP-trait associations may improve the power of ML-mediated GWAS for discovering variant-trait association with higher resolution (Szymczak *et al.*, 2009). The variable importance methods based on linear and logistic regressions, support vector machines, and random forests are well established in the literature (Grömping, 2009; Williamson *et al.*, 2020; Wu and Liu, 2009; Yoosefzadeh-Najafabadi *et al.*, 2021a). Among all the tested GWAS methods in this study, SVR-mediated GWAS was the best method to detect SNP markers with high allelic effects associated with the tested traits. The advantage of SVR-mediated GWAS over conventional GWAS models can be explained by the presence of a nonlinear relationship between input and output variables, which is used to build an algorithm with accurate prediction ability (Kaneko, 2020). Therefore, genomic regions could be better detected by SVR-mediated GWAS because of its ability to consider the interaction effects between SNPs rather than *p*-values for individual SNP-trait GWAS tests.

None of the detected QTL by MLM, FarmCPU, and RF were reported to be associated directly with soybean maturity. However, using SVR-mediated GWAS, five QTL were detected on

chromosomes 16 and 19 specifically related to the soybean maturity. Those QTL were
previously reported by Sonah *et al.* (2015) and Copley *et al.* (2018) in separate studies. Also, the
peak SNP position of Chr19_47513536 detected by SVR-mediated GWAS had the highest allelic
effect among all the detected SNPs in soybean maturity, which is in line with Sonah *et al.*
(2015). For soybean seed yield, five QTL detected by SVR-mediated GWAS were reported
previously (Copley *et al.*, 2018; Hu *et al.*, 2014), while none of the detected QTL from other
tested GWAS methods was previously reported for this trait. There was no previous study on the
genetic structure of NRNP and RNP, therefore, all the detected QTL in this study are presented
for the first time. For PP, conventional GWAS methods were not able to detect any associated
QTL. However, using SVR-mediated GWAS, a total of seven QTL were detected to be related to
pod numbers based on previous studies (Zhang *et al.*, 2015a). It would be necessary to
emphasize that the average allelic effects of all detected QTL presented in Fig. 8 was not directly
estimated by the tested GWAS methods. The RF and SVR-mediated GWAS methods do not
specifically provide an allele effect therefore, the aim of this study was mostly focused on
detecting the associated genes and QTL underlying the soybean yield, maturity, and yield
components.

The results of candidate gene identifications within identified QTL by SVR-mediated GWAS
analyses reveled important information. For example, from all the detected genes using SVR-
mediated GWAS for maturity, candidate gene *Glyma.02g006500* (GO:0015996) is a protein
ABC transporter 1, that is annotated as a chlorophyll catabolic process and located exactly in the
peak SNP position at Chr02_695362. ATP-binding cassette (ABC) transporter genes play
conspicuous roles in different plant growth and developmental stages by transporting different
phytochemicals across endoplasmic reticulum (ER) membranes (Hwang *et al.*, 2016). Because of
the central roles of ABC transporters in transporting biomolecules such as phytohormones,
metabolites, and lipids, they play important roles in plant growth and development as well as
maturity (Block and Jouhet, 2015; Hwang *et al.*, 2016). Moreover, recent studies revealed that
ER uses fatty acid building blocks made in the chloroplast to synthesize Triacylglycerol (TAG).
Therefore, ABC transporter genes are important for the normal accumulation of Triacylglycerol
(TAG) during the seed-filling stage and maturity (Block and Jouhet, 2015; Kim *et al.*, 2013).
Additionally, *Glyma.19g224200* (GO:0010201) in E3 locus, which was previously discovered by
Buzzell (1971) and molecularly characterized as a phytochrome A (PHYA) gene (Watanabe *et*
al., 2009), was detected through the SVR-mediated GWAS. Phytochromes, through
PHYTOCHROME INTERACTING FACTOR (PIF), regulate the expression of some specific
genes encoding rate-limiting catalytic enzymes of different plant growth regulators (e.g., abscisic
acid, gibberellins, auxin) and, therefore, play crucial roles in plant maturity (Legris *et al.*, 2019).
In addition, PHYB is inactivated after imbibition shade signals, which repress PHYA-dependent
signaling in the embryo that results in the maturity of seeds by preventing germination (Casal,
2013; De Wit *et al.*, 2016). This is obtained by regulating the balance between abscisic acid and
gibberellin. Subsequently, abscisic acid transports from the endosperm to the embryo by ABC
transporter (De Wit *et al.*, 2016).

Regarding NRNP, candidate gene *Glyma.07G205500* (GO:0009693- UBP1-associated protein
2C) that annotated as ethylene biosynthetic process was located exactly in the peak SNP position
of *Chr7_37469678*, was detected by SVR-mediated GWAS. An interaction screen with the
heterogeneous   nuclear   ribonucleoprotein   (hnRNP)   results   in   the   production   of

oligouridylatebinding protein 1 (UBP1)-associated protein (Lambermon *et al.*, 2002). It has been
well documented that this protein plays important roles in several physiological processes such
as responses to abiotic stresses (Li *et al.*, 2002), leaf senescence(Kim *et al.*, 2008), floral
development (Streitner *et al.*, 2008), and chromatin modification (Liu *et al.*, 2007). In addition,
previous studies showed that the production of productive or non-reproductive nodes is
completely accompanied by the upregulation or downregulation of this protein (Bäurle and
Dean, 2008; Na *et al.*, 2015). In addition, *Glyma.08G065300* (GO:0042546- MADS-box
transcription factor) that is associated with cell wall biogenesis, was located in the SNP position
of Chr8_5005929. The genes of the MADS-box family can be considered as the main regulators
for cell differentiation and organ determination (Lee *et al.*, 2013). The floral organ recognition
MADS-box family has been categorized into A, B, C, D, and E classes. Among these classes,
class E was shown to be associated with reproductive organ development (Hussin *et al.*, 2021).
Indeed, activation or repression of this transcription factor leads to the development of nodes to
productive or non-productive nodes (Ditta *et al.*, 2004; Gao *et al.*, 2010; Liu *et al.*, 2013).

Gene expression data provided by Severin *et al.* (2010) noted that 20 candidate genes for PP that
were detected using the SVR-mediated GWAS were expressed in flowers, 1 cm pod (7 DAF),
pod shell (10-13 DAF), pod shell (14-17 DAF) and seeds. In PP, most of the genes detected by
SVR-mediated GWAS are associated with auxin influx carrier or auxin response factors (ARFs),
gibberellin synthesis, and response to brassinosteroid (Lin *et al.*, 2020; Yin *et al.*, 2018). Song *et*
al. (2020) and Li *et al.* (2018a) also reported that some genes related to PP were associated with
embryo development, stamen development, ovule development, cytokinin biosynthesis, and
response gibberellin that we also identified in our study. Soybean seed yield significantly
depends on seed number and seed size (Liu *et al.*, 2010; Rotundo *et al.*, 2009). These two factors
are determined from fertilization to seed maturity. Therefore, soybean seed development can be
divided into three stages or phases: pre-embryo or seed set, embryo growth or seed growth, and
desiccation stages or seed maturation phases (Ruan *et al.*, 2012; Weber *et al.*, 2005). In
Arabidopsis, a complex signaling pathway and regulatory networks, including sugar and
hormonal signaling, transcription factors, and metabolic pathway, have been reported to be
involved in seed development (Le *et al.*, 2010; Orozco-Arroyo *et al.*, 2015). Several key genes
and transcription factors (e.g., LEAFY COTYLEDON 1 (LEC1), LEC2, FUSCA3 (FUS3),
AGAMOUS-LIKE15 (AGL15), ABSCISIC ACID INSENSITIVE 3 (ABI3), YUCCA10
(YUC10), ARFs) have been determined to control several downstream plant growth regulators
pathways to the seed development (Lepiniec *et al.*, 2018; Pelletier *et al.*, 2017; Sun *et al.*, 2010).
Indeed, a high ratio of abscisic acid to gibberellic acid can regulate seed development
(Figueiredo and Köhler, 2018; Wang *et al.*, 2016). The downregulation of FUS3 obtains this
through repressing GA3ox1 and GA3ox2 and activating ABA biosynthesis (Weber *et al.*, 2005).
In soybean, RNA seq analysis for the seed set, embryo growth, and early maturation stages of
developing seeds in two soybeans with contrasting seed size showed cell division and growth
genes, hormone regulation, transcription factors, and metabolic pathway are involved in seed
size and numbers (Du *et al.*, 2017).

Conclusion

A better understanding of the genetic architecture of the yield component traits in soybean may
enable breeders to establish more efficient selection strategies for developing high-yielding

cultivars with improved genetic gains. Major yield components such as maturity, NP, NRNP,
RNP, and PP play important roles in determining the overall yield production in soybean. This
study verified the importance of those traits, using correlation and distribution analyses, in
determining of the total soybean seed yield. Furthermore, by testing different conventional and
ML-mediated GWAS methods, this study demonstrated the potential benefit of using ML-
mediated methods in GWAS. SVR-mediated GWAS outperformed all the other methods tested
in this study, and therefore, it is recommended as an alternative to conventional GWAS methods
with a greater power for detecting genomic regions associated with complex traits such as yield
and its components in soybean, and possibly other crop species. To the best of our knowledge,
this study is the first attempt in which SVR was used for GWAS analyses in plants. In order to
verify the causal relationship between identified QTL and the target phenotypic traits, we
identified candidate genes within each QTL using gene annotation procedures and information.
The results demonstrated the efficiency of SVR-mediated GWAS in detecting reliable QTL that
can be used in marker-assisted selection. Nevertheless, further investigation is recommended to
confirm the efficiency of SVR-mediated GWAS in detecting associated genomic regions in other
plant species.

Supplementary Data

Table S1 The full list of detected genes using different GWAS methods for soybean seed yield,
maturity, and yield component traits.

Acknowledgments

The authors are grateful to the past and current members of Eskandari laboratory at the
University of Guelph, Ridgetown, Bryan Stirling, John Kobler, and Robert Brandt for their
technical support. We would like to thank Maryam Vazin and Mohsen Hesami for their
assistance with the field data collection and reviewing the manuscript, respectively.

Author Contribution

ME conceptualized, designed and directed the experiments. MY-N performed the experiments,
modeled, summed up, and wrote the manuscript. ST participated in candidate gene analyses; DT,
DTOR, ST, IR, and ME revised the manuscript and validated the results. All authors have read
and approved the final manuscript.

Data Availability Statement

The raw data supporting the conclusions of this article will be made available by the authors,
without undue reservation.

Reference

Auria L, Moro RA. 2008. Support vector machines (SVM) as a technique for solvency analysis.
Awad M, Khanna R. 2015. Support vector regression. *Efficient learning machines*: Springer,
67-80.

Bao Y, Kurle JE, Anderson G, Young ND. 2015. Association mapping and genomic prediction
for resistance to sudden death syndrome in early maturing soybean germplasm. Molecular
Breeding **35**, 1-14.
Bates D, Mächler M, Bolker B, Walker S. 2014. Fitting linear mixed-effects models using
lme4. arXiv preprint arXiv:1406.5823.
Bates D, Sarkar D, Bates MD, Matrix L. 2007. The lme4 package. R package version **2**, 74.
Bäurle I, Dean C. 2008. Differential interactions of the autonomous pathway RRM proteins and
chromatin regulators in the silencing of Arabidopsis targets. PLoS ONE **3**, e2733.
Belayneh A, Adamowski J, Khalil B, Ozga-Zielinski B. 2014. Long-term SPI drought
forecasting in the Awash River Basin in Ethiopia using wavelet neural network and wavelet
support vector regression models. Journal of Hydrology **508**, 418-429.
Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful
approach to multiple testing. Journal of the Royal Statistical Society: Series B (Methodological)
57, 289-300.
Block MA, Jouhet J. 2015. Lipid trafficking at endoplasmic reticulum–chloroplast membrane
contact sites. Current opinion in cell biology **35**, 21-29.
Botta V, Louppe G, Geurts P, Wehenkel L. 2014. Exploiting SNP correlations within random
forest for genome-wide association studies. PLoS ONE **9**, e93379.
Bowley S. 1999. *A hitchhiker's guide to statistics in plant biology*: Guelph, Ont.: Any Old
Subject Books.
Breiman L. 2001. Random forests. Machine learning **45**, 5-32.
Buzzell R. 1971. Inheritance of a soybean flowering response to fluorescent-daylength
conditions. Canadian Journal of Genetics and Cytology **13**, 703-707.
Casal JJ. 2013. Photoreceptor signaling networks in plant responses to shade. Annual review of
plant biology **64**, 403-427.
Cassell BG. 2009. Using heritability for genetic improvement.
Chang H-X, Hartman GL. 2017. Characterization of insect resistance loci in the USDA
soybean germplasm collection using genome-wide association studies. Frontiers in plant science
8, 670.
Che Z, Liu H, Yi F, Cheng H, Yang Y, Wang L, Du J, Zhang P, Wang J, Yu D. 2017.
Genome-Wide Association Study Reveals Novel Loci for SC7 Resistance in a Soybean Mutant
Panel. Frontiers in plant science **8**.
Chen JH, Verghese A. 2020. Planning for the Known Unknown: Machine Learning for Human
Healthcare Systems. The American Journal of Bioethics **20**, 1-3.
Churchill GA, Doerge RW. 1994. Empirical threshold values for quantitative trait mapping.
Genetics **138**, 963-971.
Contreras-Soto RI, Mora F, de Oliveira MAR, Higashi W, Scapim CA, Schuster I. 2017. A
genome-wide association study for agronomic traits in soybean using SNP markers and SNP-
based haplotype analysis. PLoS ONE **12**, e0171105.
Cook DE, Bayless AM, Wang K, Guo X, Song Q, Jiang J, Bent AF. 2014. Distinct copy
number, coding sequence, and locus methylation patterns underlie Rhg1-mediated soybean
resistance to soybean cyst nematode. Plant Physiology **165**, 630-647.
Cooper M, van Eeuwijk FA, Hammer GL, Podlich DW, Messina C. 2009. Modeling QTL for
complex traits: detection and context for plant breeding. Current Opinion in Plant Biology **12**,
231-240.

Copley TR, Duceppe M-O, O'Donoughue LS. 2018. Identification of novel loci associated with maturity and yield traits in early maturity soybean plant introduction lines. BMC Genomics **19**, 1-12.

Cortes C, Vapnik V. 1995. Support vector machine. Machine learning **20**, 273-297.

De Wit M, Galvão VC, Fankhauser C. 2016. Light-mediated hormonal regulation of plant growth and development. Annual review of plant biology **67**, 513-537.

Denton SM, Salleb-Aouissi A. 2020. A Weighted Solution to SVM Actionability and Interpretability. arXiv preprint arXiv:2012.03372.

Dhanapal AP, Ray JD, Singh SK, Hoyos-Villegas V, Smith JR, Purcell LC, King CA, Cregan PB, Song Q, Fritschi FB. 2015a. Genome-wide association study (GWAS) of carbon isotope ratio (δ 13 C) in diverse soybean [Glycine max (L.) Merr.] genotypes. Theoretical and Applied Genetics **128**, 73-91.

Dhanapal AP, Ray JD, Singh SK, Hoyos-Villegas V, Smith JR, Purcell LC, King CA, Fritschi FB. 2015b. Association mapping of total carotenoids in diverse soybean genotypes based on leaf extracts and high-throughput canopy spectral reflectance measurements. PLoS ONE **10**, e0137213.

Dhanapal AP, Ray JD, Smith JR, Purcell LC, Fritschi FB. 2018. Identification of Novel Genomic Loci Associated with Soybean Shoot Tissue Macro and Micronutrient Concentrations. The Plant Genome **11**, 170066.

Ditta G, Pinyopich A, Robles P, Pelaz S, Yanofsky MF. 2004. The SEP4 gene of Arabidopsis thaliana functions in floral organ and meristem identity. Current Biology **14**, 1935-1940.

Doerge RW, Churchill GA. 1996. Permutation tests for multiple loci affecting a quantitative character. Genetics **142**, 285-294.

Du J, Wang S, He C, Zhou B, Ruan Y-L, Shou H. 2017. Identification of regulatory networks and hub genes controlling soybean seed set and size using RNA sequencing analysis. Journal of experimental botany **68**, 1955-1972.

Duan K-B, Rajapakse JC, Wang H, Azuaje F. 2005. Multiple SVM-RFE for gene selection in cancer classification with expression data. IEEE transactions on nanobioscience **4**, 228-234.

Fang C, Ma Y, Wu S, Liu Z, Wang Z, Yang R, Hu G, Zhou Z, Yu H, Zhang M. 2017. Genome-wide association studies dissect the genetic networks underlying agronomical traits in soybean. Genome Biology **18**, 1-14.

Fehr W, Caviness C. 1971. Burmood DT, Penington J S. Development description of soybean, Glycine max (L) Mer. Crop science **11**, 929-931.

Figueiredo DD, Köhler C. 2018. Auxin: a molecular trigger of seed development. Genes & development **32**, 479-490.

Fletcher T. 2009. Support vector machines explained. Tutorial paper., Mar, 28.

Gao X, Liang W, Yin C, Ji S, Wang H, Su X, Guo C, Kong H, Xue H, Zhang D. 2010. The SEPALLATA-like gene OsMADS34 is required for rice inflorescence and spikelet development. Plant Physiology **153**, 728-740.

Goldberger AS. 1962. Best linear unbiased prediction in the generalized linear regression model. Journal of the American Statistical Association **57**, 369-375.

Grömping U. 2009. Variable importance assessment in regression: linear regression versus random forest. The American Statistician **63**, 308-319.

Herbert S, Litchfield G. 1982. Partitioning Soybean Seed Yield Components 1. Crop science **22**, 1074-1079.

Hesami M, Jones AMP. 2020. Application of artificial intelligence models and optimization algorithms in plant cell and tissue culture. Applied Microbiology and Biotechnology, 1-37.

Hesami M, Naderi R, Tohidfar M, Yoosefzadeh-Najafabadi M. 2020. Development of support vector machine-based model and comparative analysis with artificial neural network for modeling the plant tissue culture procedures: effect of plant growth regulators on somatic embryogenesis of chrysanthemum, as a case study. Plant methods **16**, 1-15.

Hu D, Zhang H, Du Q, Hu Z, Yang Z, Li X, Wang J, Huang F, Yu D, Wang H. 2020. Genetic dissection of yield-related traits via genome-wide association analysis across multiple environments in wild soybean (Glycine soja Sieb. and Zucc.). Planta **251**, 39.

Hu Z, Zhang D, Zhang G, Kan G, Hong D, Yu D. 2014. Association mapping of yield-related traits and SSR markers in wild soybean (Glycine soja Sieb. and Zucc.). Breeding science **63**, 441-449.

Hussin SH, Wang H, Tang S, Zhi H, Tang C, Zhang W, Jia G, Diao X. 2021. SiMADS34, an E-class MADS-box transcription factor, regulates inflorescence architecture and grain yield in Setaria italica. Plant Molecular Biology **105**, 419-434.

Hwang J-U, Song W-Y, Hong D, Ko D, Yamaoka Y, Jang S, Yim S, Lee E, Khare D, Kim K. 2016. Plant ABC transporters enable many unique aspects of a terrestrial plant's lifestyle. Molecular Plant **9**, 338-355.

Jamil IN, Remali J, Azizan KA, Muhammad NAN, Arita M, Goh H-H, Aizat WM. 2020. Systematic Multi-Omics Integration (MOI) Approach in Plant Systems Biology. Frontiers in plant science **11**.

Jordan MI, Mitchell TM. 2015. Machine learning: Trends, perspectives, and prospects. Science **349**, 255-260.

Kahlon CS, Board JE. 2012. Growth dynamic factors explaining yield improvement in new versus old soybean cultivars. Journal of crop improvement **26**, 282-299.

Kahlon CS, Board JE, Kang MS. 2011. An analysis of yield component changes for new vs. old soybean cultivars. Agronomy Journal **103**, 13-22.

Kaler AS, Dhanapal AP, Ray JD, King CA, Fritschi FB, Purcell LC. 2017. Genome-wide association mapping of carbon isotope and oxygen isotope ratios in diverse soybean genotypes. Crop science **57**, 3085-3100.

Kaler AS, Gillman JD, Beissinger T, Purcell LC. 2020. Comparing different statistical models and multiple testing corrections for association mapping in soybean and maize. Frontiers in plant science **10**, 1794.

Kan G, Zhang W, Yang W, Ma D, Zhang D, Hao D, Hu Z, Yu D. 2015. Association mapping of soybean seed germination under salt stress. Molecular Genetics and Genomics **290**, 2147-2162.

Kaneko H. 2020. Support vector regression that takes into consideration the importance of explanatory variables. Journal of Chemometrics, e3327.

Katsileros A, Drosou K, Koukouvinos C. 2015. Evaluation of nearest neighbor methods in wheat genotype experiments. Communications in Biometry and Crop Science **10**, 115-123.

Kim CY, Bove J, Assmann SM. 2008. Overexpression of wound-responsive RNA-binding proteins induces leaf senescence and hypersensitive-like cell death. New Phytologist **180**, 57-70.

Kim GB, Kim WJ, Kim HU, Lee SY. 2020. Machine learning applications in systems metabolic engineering. Current opinion in biotechnology **64**, 1-9.

Kim S, Yamaoka Y, Ono H, Kim H, Shim D, Maeshima M, Martinoia E, Cahoon EB, Nishida I, Lee Y. 2013. AtABCA9 transporter supplies fatty acids for lipid synthesis to the endoplasmic reticulum. Proceedings of the National Academy of Sciences **110**, 773-778.

Kuhn M, Wing J, Weston S, Williams A, Keefer C, Engelhardt A, Cooper T, Mayer Z, Kenkel B, Team RC. 2020. Package 'caret'. The R Journal.

KUSWANTORO H. 2017. Genetic variability and heritability of acid-adaptive soybean promising lines. Biodiversitas Journal of Biological Diversity **18**.

Lambermon MH, Fu Y, Kirk DAW, Dupasquier M, Filipowicz W, Lorković ZJ. 2002. UBA1 and UBA2, two proteins that interact with UBP1, a multifunctional effector of pre-mRNA maturation in plants. Molecular and Cellular Biology **22**, 4346-4357.

Le BH, Cheng C, Bui AQ, Wagmaister JA, Henry KF, Pelletier J, Kwong L, Belmonte M, Kirkbride R, Horvath S. 2010. Global analysis of gene activity during Arabidopsis seed development and identification of seed-specific transcription factors. Proceedings of the National Academy of Sciences **107**, 8063-8070.

Leamy LJ, Zhang H, Li C, Chen CY, Song B-H. 2017. A genome-wide association study of seed composition traits in wild soybean (Glycine soja). BMC Genomics **18**, 1-15.

Lee JH, Ryu H-S, Chung KS, Posé D, Kim S, Schmid M, Ahn JH. 2013. Regulation of temperature-responsive flowering by MADS-box transcription factor repressors. Science **342**, 628-632.

Lee S, Liang X, Woods M, Reiner AS, Concannon P, Bernstein L, Lynch CF, Boice JD, Deasy JO, Bernstein JL. 2020. Machine learning on genome-wide association studies to predict the risk of radiation-associated contralateral breast cancer in the WECARE Study. PLoS ONE **15**, e0226157.

Legris M, Ince YÇ, Fankhauser C. 2019. Molecular mechanisms underlying phytochrome-controlled morphogenesis in plants. Nature communications **10**, 1-15.

Lepiniec L, Devic M, Roscoe T, Bouyer D, Zhou D-X, Boulard C, Baud S, Dubreucq B. 2018. Molecular and epigenetic regulations and functions of the LAFL transcriptional regulators that control seed development. Plant reproduction **31**, 291-307.

Li C, Zou J, Jiang H, Yu J, Huang S, Wang X, Liu C, Guo T, Zhu R, Wu X. 2018a. Identification and validation of number of pod□and seed□related traits QTL s in soybean. Plant Breeding **137**, 730-745.

Li J, Kinoshita T, Pandey S, Ng CK-Y, Gygi SP, Shimazaki K-i, Assmann SM. 2002. Modulation of an RNA-binding protein by abscisic-acid-activated protein kinase. Nature **418**, 793-797.

Li X, Tian R, Kamala S, Du H, Li W, Kong Y, Zhang C. 2018b. Identification and verification of pleiotropic QTL controlling multiple amino acid contents in soybean seed. Euphytica **214**, 1-14.

Li Y-h, Reif JC, Ma Y-s, Hong H-l, Liu Z-x, Chang R-z, Qiu L-j. 2015. Targeted association mapping demonstrating the complex molecular genetics of fatty acid formation in soybean. BMC Genomics **16**, 1-13.

Li Yh, Shi Xh, Li Hh, Reif JC, Wang Jj, Liu Zx, He S, Yu Bs, Qiu Lj. 2016. Dissecting the genetic basis of resistance to soybean cyst nematode combining linkage and association mapping. The Plant Genome **9**, plantgenome2015.2004.0020.

Lin F, Wani SH, Collins PJ, Wen Z, Li W, Zhang N, McCoy AG, Bi Y, Tan R, Zhang S. 2020. QTL mapping and GWAS for identification of loci conferring partial resistance to Pythium sylvaticum in soybean (Glycine max (L.) Merr). Molecular Breeding **40**, 1-11.

Lipka AE, Kandianis CB, Hudson ME, Yu J, Drnevich J, Bradbury PJ, Gore MA. 2015. From association to prediction: statistical methods for the dissection and selection of complex traits in plants. Current Opinion in Plant Biology **24**, 110-118.

Lipka AE, Tian F, Wang Q, Peiffer J, Li M, Bradbury PJ, Gore MA, Buckler ES, Zhang Z. 2012. GAPIT: genome association and prediction integrated tool. Bioinformatics **28**, 2397-2399.

Liu B, Liu X, Wang C, Li Y, Jin J, Herbert S. 2010. Soybean yield and yield component distribution across the main axis in response to light enrichment and shading under different densities. Plant, Soil and Environment **56**, 384-392.

Liu C, Teo ZWN, Bi Y, Song S, Xi W, Yang X, Yin Z, Yu H. 2013. A conserved genetic pathway determines inflorescence architecture in Arabidopsis and rice. Developmental cell **24**, 612-622.

Liu F, Quesada V, Crevillén P, Bäurle I, Swiezewski S, Dean C. 2007. The Arabidopsis RNA-binding protein FCA requires a lysine-specific demethylase 1 homolog to downregulate FLC. Molecular cell **28**, 398-407.

Liu X, Huang M, Fan B, Buckler ES, Zhang Z. 2016. Iterative usage of fixed and random effect models for powerful and efficient genome-wide association studies. PLoS Genetics **12**, e1005767.

Ma B, Dwyer LM, Costa C, Cober ER, Morrison MJ. 2001. Early prediction of soybean yield from canopy reflectance measurements. Agronomy Journal **93**, 1227-1234.

Mangena P. 2020. Phytocystatins and their Potential Application in the Development of Drought Tolerance Plants in Soybeans (Glycine max L.). Protein and Peptide Letters **27**, 135-144.

Mao T, Li J, Wen Z, Wu T, Wu C, Sun S, Jiang B, Hou W, Li W, Song Q. 2017. Association mapping of loci controlling genetic and environmental interaction of soybean flowering time under various photo-thermal conditions. BMC Genomics **18**, 1-17.

Meinshausen N. 2006. Quantile regression forests. Journal of Machine Learning Research **7**, 983-999.

Moellers TC, Singh A, Zhang J, Brungardt J, Kabbage M, Mueller DS, Grau CR, Ranjan A, Smith DL, Chowda-Reddy R. 2017. Main and epistatic loci studies in soybean for Sclerotinia sclerotiorum resistance reveal multiple modes of resistance in multi-environments. Scientific reports **7**, 1-13.

Mohammadi M, Xavier A, Beckett T, Beyer S, Chen L, Chikssa H, Cross V, Moreira FF, French E, Gaire R. 2020. Identification, Deployment, and Transferability of Quantitative Trait Loci from Genome-Wide Association Studies in Plants. Current Plant Biology, 100145.

Na J-K, Kim J-K, Kim D-Y, Assmann SM. 2015. Expression of potato RNA-binding proteins StUBA2a/b and StUBA2c induces hypersensitive-like cell death and early leaf senescence in Arabidopsis. Journal of experimental botany **66**, 4023-4033.

Nico M, Miralles DJ, Kantolic AG. 2019. Natural post-flowering photoperiod and photoperiod sensitivity: Roles in yield-determining processes in soybean. Field crops research **231**, 141-152.

Ogutu JO, Piepho H-P, Schulz-Streeck T. 2011. A comparison of random forests, boosting and support vector machines for genomic selection. *BMC proceedings*, Vol. 5: Springer, S11.

Orozco-Arroyo G, Paolo D, Ezquer I, Colombo L. 2015. Networks controlling seed size in Arabidopsis. Plant reproduction **28**, 17-32.

Pasaniuc B, Price AL. 2017. Dissecting the genetics of complex traits using summary association statistics. Nature reviews genetics **18**, 117-127.

Pedersen P, Lauer JG. 2004. Response of soybean yield components to management system
and planting date. Agronomy Journal **96**, 1372-1381.
Pelletier JM, Kwong RW, Park S, Le BH, Baden R, Cagliari A, Hashimoto M, Munoz MD,
Fischer RL, Goldberg RB. 2017. LEC1 sequentially regulates the transcription of genes
involved in diverse developmental processes during seed development. Proceedings of the
National Academy of Sciences **114**, E6710-E6719.
Price T, Schluter D. 1991. On the low heritability of life history traits. Evolution **45**, 853-861.
Priolli RHG, Campos J, Stabellini N, Pinheiro J, Vello N. 2015. Association mapping of oil
content and fatty acid components in soybean. Euphytica **203**, 83-96.
Qin J, Song Q, Shi A, Li S, Zhang M, Zhang B. 2017. Genome-wide association mapping of
resistance to Phytophthora sojae in a soybean [Glycine max (L.) Merr.] germplasm panel from
maturity groups IV and V. PLoS ONE **12**, e0184613.
Raj A, Stephens M, Pritchard JK. 2014. fastSTRUCTURE: variational inference of population
structure in large SNP data sets. Genetics **197**, 573-589.
Ray JD, Dhanapal AP, Singh SK, Hoyos-Villegas V, Smith JR, Purcell LC, King CA,
Boykin D, Cregan PB, Song Q. 2015. Genome-wide association study of ureide concentration
in diverse maturity group IV soybean [Glycine max (L.) Merr.] accessions. G3: Genes,
Genomes, Genetics **5**, 2391-2403.
Rębilas K, Klimek-Kopyra A, Bacior M, Zając T. 2020. A model for the yield losses
estimation in an early soybean (Glycine max (L.) Merr.) cultivar depending on the cutting height
at harvest. Field crops research **254**, 107846.
Reynolds M. 2001. *Application of physiology in wheat breeding*: Cimmyt.
Richards R. 1982. Breeding and selecting for drought resistant wheat. p. 303–316. Drought
resistance in crops with emphasis on rice. IRRI, Manila, Philippines. Breeding and selecting for
drought resistant wheat. p. 303–316. In Drought resistance in crops with emphasis on rice. IRRI,
Manila, Philippines., -.
Robinson AP, Conley SP, Volenec JJ, Santini JB. 2009. Analysis of high yielding,
early planted soybean in Indiana. Agronomy Journal **101**, 131-139.
Rotundo JL, Borrás L, Westgate ME, Orf JH. 2009. Relationship between assimilate supply
per seed during seed filling and soybean seed composition. Field crops research **112**, 90-96.
Ruan Y-L, Patrick JW, Bouzayen M, Osorio S, Fernie AR. 2012. Molecular regulation of
seed and fruit set. Trends in plant science **17**, 656-665.
Severin AJ, Woody JL, Bolon Y-T, Joseph B, Diers BW, Farmer AD, Muehlbauer GJ,
Nelson RT, Grant D, Specht JE. 2010. RNA-Seq Atlas of Glycine max: a guide to the soybean
transcriptome. BMC plant biology **10**, 1-16.
Siegmann B, Jarmer T. 2015. Comparison of different regression models and validation
techniques for the assessment of wheat leaf area index from hyperspectral data. International
Journal of Remote Sensing **36**, 4519-4534.
Sonah H, Bastien M, Iquira E, Tardivel A, Légaré G, Boyle B, Normandeau É, Laroche J,
Larose S, Jean M, Belzile F. 2013. An Improved Genotyping by Sequencing (GBS) Approach
Offering Increased Versatility and Efficiency of SNP Discovery and Genotyping. PLoS ONE **8**,
e54603.
Sonah H, O'Donoughue L, Cober E, Rajcan I, Belzile F. 2015. Identification of loci
governing eight agronomic traits using a GBS GWAS approach and validation by QTL
mapping in soya bean. Plant biotechnology journal **13**, 211-221.

Song J, Sun X, Zhang K, Liu S, Wang J, Yang C, Jiang S, Siyal M, Li X, Qi Z. 2020. Identification of QTL and genes for pod number in soybean by linkage analysis and genome-wide association studies. Molecular Breeding **40**, 1-14.

Streitner C, Danisman S, Wehrle F, Schöning JC, Alfano JR, Staiger D. 2008. The small glycine☐rich RNA binding protein AtGRP7 promotes floral transition in Arabidopsis thaliana. The Plant Journal **56**, 239-250.

Stroup W, Mulitze D. 1991. Nearest neighbor adjusted best linear unbiased prediction. The American Statistician **45**, 194-200.

Su Q, Lu W, Du D, Chen F, Niu B, Chou K-C. 2017. Prediction of the aquatic toxicity of aromatic compounds to tetrahymena pyriformis through support vector regression. Oncotarget **8**, 49359.

Suhre JJ, Weidenbenner NH, Rowntree SC, Wilson EW, Naeve SL, Conley SP, Casteel SN, Diers BW, Esker PD, Specht JE. 2014. Soybean yield partitioning changes revealed by genetic gain and seeding rate interactions. Agronomy Journal **106**, 1631-1642.

Sulistyo A, Sari K. 2018. Correlation, path analysis and heritability estimation for agronomic traits contribute to yield on soybean. *IOP Conference Series: Earth and Environmental Science*, Vol. 102, 012034.

Sun S, Wang C, Ding H, Zou Q. 2020. Machine learning and its applications in plant molecular studies. Briefings in Functional Genomics **19**, 40-48.

Sun X, Shantharaj D, Kang X, Ni M. 2010. Transcriptional and hormonal signaling control of Arabidopsis seed development. Current Opinion in Plant Biology **13**, 611-620.

Szymczak S, Biernacka JM, Cordell HJ, González☐Recio O, König IR, Zhang H, Sun YV. 2009. Machine learning in genome☐wide association studies. Genetic epidemiology **33**, S51-S57.

Temesgen D, Assefa F. 2020. Inoculation of native symbiotic effective Sinorhizobium spp. enhanced soybean [Glycine max (L.) Merr.] grain yield in Ethiopia. Environmental Systems Research **9**, 1-19.

Torkamaneh D, Laroche J, Belzile F. 2020. Fast-GBS v2.0: an analysis toolkit for genotyping-by-sequencing data. Genome **63**, 577-581.

Tulpan D. 2020. 311 A brief overview, comparison and practical applications of machine learning models. Journal of animal science **98**, 44-45.

Vuong T, Sonah H, Meinhardt C, Deshmukh R, Kadam S, Nelson R, Shannon J, Nguyen H. 2015. Genetic architecture of cyst nematode resistance revealed by genome-wide association study in soybean. BMC Genomics **16**, 1-13.

Wang L, Hu X, Jiao C, Li Z, Fei Z, Yan X, Liu C, Wang Y, Wang X. 2016. Transcriptome analyses of seed development in grape hybrids reveals a possible mechanism influencing seed size. BMC Genomics **17**, 1-15.

Watanabe S, Hideshima R, Xia Z, Tsubokura Y, Sato S, Nakamoto Y, Yamanaka N, Takahashi R, Ishimoto M, Anai T. 2009. Map-based cloning of the gene associated with the soybean maturity locus E3. Genetics **182**, 1251-1262.

Weber H, Borisjuk L, Wobus U. 2005. Molecular physiology of legume seed development. Annu. Rev. Plant Biol. **56**, 253-279.

Wen Y-J, Zhang H, Ni Y-L, Huang B, Zhang J, Feng J-Y, Wang S-B, Dunwell JM, Zhang Y-M, Wu R. 2018. Methodological implementation of mixed linear models in multi-locus genome-wide association studies. Briefings in bioinformatics **19**, 700-712.

Wen Z, Tan R, Yuan J, Bales C, Du W, Zhang S, Chilvers MI, Schmidt C, Song Q, Cregan PB. 2014. Genome-wide association mapping of quantitative resistance to sudden death syndrome in soybean. BMC Genomics **15**, 1-11.

Weston J, Mukherjee S, Chapelle O, Pontil M, Poggio T, Vapnik V. 2001. Feature selection for SVMs. *Advances in neural information processing systems*, 668-674.

Wickham H. 2011. ggplot2. Wiley Interdisciplinary Reviews: Computational Statistics **3**, 180-185.

Williamson BD, Gilbert PB, Simon NR, Carone M. 2020. A unified approach for inference on algorithm-agnostic variable importance. arXiv preprint arXiv:2004.03683.

Wu Y, Liu Y. 2009. Variable selection in quantile regression. Statistica Sinica, 801-817.

Xavier A, Jarquin D, Howard R, Ramasubramanian V, Specht JE, Graef GL, Beavis WD, Diers BW, Song Q, Cregan PB. 2018. Genome-wide analysis of grain yield stability and environmental interactions in a multiparental soybean population. G3: Genes, Genomes, Genetics **8**, 519-529.

Xavier A, Muir WM, Rainey KM. 2016a. Assessing predictive properties of genome-wide selection in soybeans. G3: Genes, Genomes, Genetics **6**, 2611-2616.

Xavier A, Muir WM, Rainey KM. 2016b. Impact of imputation methods on the amount of genetic variation captured by a single-nucleotide polymorphism panel in soybeans. BMC Bioinformatics **17**, 1-9.

Xavier A, Rainey KM. 2020. Quantitative Genomic Dissection of Soybean Yield Components. G3: Genes, Genomes, Genetics **10**, 665-675.

Yang J, Yeh C-TE, Ramamurthy RK, Qi X, Fernando RL, Dekkers JC, Garrick DJ, Nettleton D, Schnable PS. 2018. Empirical comparisons of different statistical models to identify and validate kernel row number-associated variants from structured multi-parent mapping populations of maize. G3: Genes, Genomes, Genetics **8**, 3567-3575.

Yin Z, Qi H, Mao X, Wang J, Hu Z, Wu X, Liu C, Xin D, Zuo X, Chen Q. 2018. QTL mapping of soybean node numbers on the main stem and meta-analysis for mining candidate genes. Biotechnology & Biotechnological Equipment **32**, 915-922.

Yoosefzadeh-Najafabadi M, Earl HJ, Tulpan D, Sulik J, Eskandari M. 2021a. Application of Machine Learning Algorithms in Plant Breeding: Predicting Yield From Hyperspectral Reflectance in Soybean. Frontiers in plant science **11**.

Yoosefzadeh-Najafabadi M, Tulpan D, Eskandari M. 2021b. Application of machine learning and genetic optimization algorithms for modeling and optimizing soybean yield using its component traits. PLoS ONE **16**, e0250665.

Zhang H, Hao D, Sitoe HM, Yin Z, Hu Z, Zhang G, Yu D. 2015a. Genetic dissection of the relationship between plant architecture and yield component traits in soybean (Glycine max) by association analysis across multiple environments. Plant Breeding **134**, 564-572.

Zhang J, Song Q, Cregan PB, Jiang G-L. 2016. Genome-wide association study, genomic prediction and marker-assisted selection for seed weight in soybean (Glycinemax). Theoretical and Applied Genetics **129**, 117-130.

Zhang J, Song Q, Cregan PB, Nelson RL, Wang X, Wu J, Jiang G-L. 2015b. Genome-wide association study for flowering time, maturity dates and plant height in early maturing soybean (Glycine max) germplasm. BMC Genomics **16**, 1-11.

Zhang J, Wang X, Lu Y, Bhusal SJ, Song Q, Cregan PB, Yen Y, Brown M, Jiang G-L. 2018. Genome-wide Scan for Seed Composition Provides Insights into Soybean Quality Improvement and the Impacts of Domestication and Breeding. Molecular Plant **11**, 460-472.

Tables

Table 1. The list of detected QTL for soybean maturity using different GWAS methods in the tested soybean population.

GWAS Method	Chromosome	Peak SNP position	Detected QTL	Environment [a]	Reference
MLM	2	2212910	Sclero 3-g31	NA	(Moellers et al., 2017)
		8233782	Seed Weight 6-g1	NA	(Sonah et al., 2015)
FarmCPU	2	2212910	Sclero 3-g31	NA	(Moellers et al., 2017)
		8233766	Seed Weight 6-g1	NA	(Sonah et al., 2015)
	20	37765851	WUE 2-g53	NA	(Kaler et al., 2017)
RF	3	2978272	Leaflet area 1-g2.1	NA	(Fang et al., 2017)
			Leaflet width 1-g4.1	NA	(Fang et al., 2017)
			Leaflet area 1-g2.2	NA	(Fang et al., 2017)
			Leaflet width 1-g4.2	NA	(Fang et al., 2017)
			Salt tolerance 1-g12	NA	(Kan et al., 2015)
	16	5730281	Plant height 6-g17	NA	(Zhang et al., 2015b)
			Plant height 1-g17	NA	(Zhang et al., 2015b)
			First flower 4-g63	NA	(Mao et al., 2017)
	17	34757372	SDS root retention 1-g6	NA	(Bao et al., 2015)
SVR	2	695362	Seed linolenic 2-g1	NA	(Leamy et al., 2017)
			Seed linolenic 2-g2	NA	(Leamy et al., 2017)
		720134	SDS 1-g12.1	2	(Wen et al., 2014)
			SDS 1-g12.2	2	(Wen et al., 2014)
			Ureide content 1-g2	2	(Ray et al., 2015)
		827374	SDS 1-g12.3	NA	(Wen et al., 2014)
	10	1595239	Shoot Cu 1-g8	NA	(Dhanapal et al., 2018)
		1689395	Seed oil 5-g3	NA	(Sonah et al., 2015)
	16	2438652	Reproductive period 4-g16	NA	(Zhang et al., 2015b)
			R8 full maturity 9-g2	NA	(Zhang et al., 2015b)
		2460921	Reproductive period 2-g16	NA	(Zhang et al., 2015b)
			R8 full maturity 2-g2	NA	(Zhang et al., 2015b)
	19	47513536	R8 full maturity 4-g1	NA	(Sonah et al., 2015)
		47513572	First flower 4-g81	NA	(Mao et al., 2017)

[a] Detected in separate environments (1: 2018Ridgetown, 2:2019Ridgetwon, 3:2018Palmyra, 4:2019Palmyra, NA: Not found in any environment). MLM: Mixed Linear Model, FarmCPU: Fixed and random model circulating probability unification, RF: Random Forest, SVR: Support Vector Regression

Table 1. The list of detected QTL for soybean yield using different GWAS methods in the tested soybean population.

GWAS Method	Chromosome	Peak SNP position	Detected QTL	Environment [a]	Reference
MLM	5	34391386	Ureide content 1-g16.1	NA	(Ray *et al.*, 2015)
			Ureide content 1-g16.2	NA	(Ray *et al.*, 2015)
FarmCPU	5	34391386	Ureide content 1-g16.1	NA	(Ray *et al.*, 2015)
			Ureide content 1-g16.2	NA	(Ray *et al.*, 2015)
RF	7	1032587	WUE 2-g18	NA	(Kaler *et al.*, 2017)
SVR	3	36309302	First flower 4-g10	NA	(Mao *et al.*, 2017)
			First flower 3-g2	NA	(Hu *et al.*, 2014)
			Seed weight 4-g3	NA	(Hu *et al.*, 2014)
			Seed yield 4-g2	NA	(Hu *et al.*, 2014)
			R8 full maturity 3-g3	NA	(Hu *et al.*, 2014)
			Plant height 3-g17	NA	(Contreras-Soto *et al.*, 2017)
		37617293	Leaflet shape 1-g1.1	NA	(Fang *et al.*, 2017)
			Leaflet shape 1-g1.2	NA	(Fang *et al.*, 2017)
			Leaflet shape 1-g1.3	NA	(Fang *et al.*, 2017)
			Seed set 1-g32.1	NA	(Fang *et al.*, 2017)
			Seed set 1-g32.2	NA	(Fang *et al.*, 2017)
	7	44488152	Seed yield 4-g4	NA	(Hu *et al.*, 2014)
		1032587	WUE 2-g18	NA	(Kaler *et al.*, 2017)
	15	34958361	SCN 5-g35	NA	(Li *et al.*, 2016)
	19	41385139	Seed weight 5-g20	NA	(Zhang *et al.*, 2016)
			Seed weight 4-g18	NA	(Hu *et al.*, 2014)
			Seed yield 4-g5	NA	(Hu *et al.*, 2014)
			Shoot Zn 1-g28.1	NA	(Dhanapal *et al.*, 2018)
			Shoot Zn 1-g28.2	NA	(Dhanapal *et al.*, 2018)

	Shoot Zn 1-g29.1	NA	(Dhanapal *et al.*, 2018)
	Shoot Zn 1-g29.2	NA	(Dhanapal *et al.*, 2018)
	Shoot Zn 1-g29.3	NA	(Dhanapal *et al.*, 2018)

[a] Detected in separate environments (1: 2018Ridgetown, 2:2019Ridgetwon, 3:2018Palmyra, 4:2019Palmyra, NA: Not found in any environment).

MLM: Mixed Linear Model, FarmCPU: Fixed and random model circulating probability unification, RF: Random Forest, SVR: Support Vector Regression

Table 2. The list of detected QTL for soybean total number of nodes per plant (NP) using different GWAS methods in the tested soybean population.

GWAS Method	Chromosome	Peak SNP position	Detected QTL	Environment [a]	Reference
FarmCPU	19	40131952	Pubescence density 1-g17	NA	(Chang and Hartman, 2017)
			Seed weight 9-g5.1	NA	(Copley *et al.*, 2018)
RF	4	1205787	Shoot Ca 1-g10	NA	(Dhanapal *et al.*, 2018)
	6	50570624	Seed set 1-g51.1	NA	(Fang *et al.*, 2017)
			Seed set 1-g43.1	NA	(Fang *et al.*, 2017)
			Seed set 1-g25.1	NA	(Fang *et al.*, 2017)
			Seed set 1-g43.2	NA	(Fang *et al.*, 2017)
			Seed set 1-g25.2	NA	(Fang *et al.*, 2017)
			Seed set 1-g51.2	NA	(Fang *et al.*, 2017)
			Seed set 1-g43.3	NA	(Fang *et al.*, 2017)
		50570473	Seed set 1-g51.3	NA	(Fang *et al.*, 2017)
			Seed set 1-g25.3	NA	(Fang *et al.*, 2017)
			Pod number 1-g3	NA	(Fang *et al.*, 2017)
			Seed palmitic 2-g2	NA	(Fang *et al.*, 2017)
			Seed long-chain faty acid 1-g22	NA	(Fang *et al.*, 2017)
SVR	6	50570624	Seed set 1-g51.1	NA	(Fang *et al.*, 2017)
			Seed set 1-g43.1	NA	(Fang *et al.*, 2017)
			Seed set 1-g25.1	NA	(Fang *et al.*, 2017)
			Seed set 1-g43.2	NA	(Fang *et al.*, 2017)
			Seed set 1-g25.2	NA	(Fang *et al.*, 2017)
			Seed set 1-g51.2	NA	(Fang *et al.*, 2017)
			Seed set 1-g43.3	NA	(Fang *et al.*, 2017)
		50570473	Seed set 1-g51.3	NA	(Fang *et al.*, 2017)
			Seed set 1-g25.3	NA	(Fang *et al.*, 2017)

		Pod number 1-g3	NA	(Fang *et al.*, 2017)
		Seed palmitic 2-g2	NA	(Fang *et al.*, 2017)
		Seed long-chain faty acid 1-g22	NA	(Fang *et al.*, 2017)
7	1032587	WUE 2-g18	NA	(Kaler *et al.*, 2017)
	1092403	WUE 2-g18	NA	(Kaler *et al.*, 2017)
		First flower 3-g4	NA	(Fang *et al.*, 2017)
		Leaflet shape 1-g4.1	NA	(Fang *et al.*, 2017)
		Leaflet shape 1-g4.2	NA	(Fang *et al.*, 2017)
		Leaflet shape 1-g4.3	NA	(Fang *et al.*, 2017)
		Seed stearic 4-g5	NA	(Li *et al.*, 2015)
		Node number 1-g6.1	NA	(Fang *et al.*, 2017)
		Node number 1-g6.2	NA	(Fang *et al.*, 2017)
		Pod number 1-g1.1	NA	(Fang *et al.*, 2017)
		Pod number 1-g1.2	NA	(Fang *et al.*, 2017)
18	55645699	Pode number 1-g1.3	NA	(Fang *et al.*, 2017)
		WUE 3-g31	NA	(Kaler *et al.*, 2017)
		Seed weight, SoyNAM 14-g28	NA	(Xavier *et al.*, 2016b)
		Lodging, SoyNAM 4-g15	NA	(Cook *et al.*, 2014)
		Branching 1-g1.1	NA	(Fang *et al.*, 2017)
		Plant height 5-g4.2	NA	(Fang *et al.*, 2017)
		Plant height 5-g4.3	NA	(Fang *et al.*, 2017)
		Shoot p 1-g30	NA	(Dhanapal *et al.*, 2018)
19	47350110	Node number 1-g2.3	NA	(Fang *et al.*, 2017)

[a] Detected in separate environments (1: 2018Ridgetown, 2:2019Ridgetwon, 3:2018Palmyra, 4:2019Palmyra, NA: Not found in any environment). MLM: Mixed Linear Model, FarmCPU: Fixed and random model circulating probability unification, RF: Random Forest, SVR: Support Vector Regression

Table 3. The list of detected QTL for soybean total number of non-reproductive nodes per plant (NRNP) using different GWAS methods in the tested soybean population.

GWAS Method	Chromosome	Peak SNP position	Detected QTL	Environment [a]	Reference
MLM	15	10193796	Seed protein 6-g2	NA	(Zhang *et al.*, 2018)
			Seed Arg 1-g4	NA	(Zhang *et al.*, 2018)
			Seed coat luster 1-g1.3	NA	(Fang *et al.*, 2017)
FarmCPU	15	10193796	Seed protein 6-g2	NA	(Zhang *et al.*, 2018)
			Seed Arg 1-g4	NA	(Zhang *et al.*, 2018)
			Seed coat luster 1-g1.3	NA	(Fang *et al.*, 2017)
RF	1	54647498	First flower 4-g2	NA	(Mao *et al.*, 2017)
	7	329800	Phytoph 2-g32	NA	(Qin *et al.*, 2017)
			Phytoph 2-g7	NA	(Qin *et al.*, 2017)
	18	12945778	SCN 4-g14	NA	(Vuong *et al.*, 2015)
	19	40218800	Seed weight 9-g5.1	NA	(Copley *et al.*, 2018)
SVR	7	1032587 [2]	WUE 2-g18	2	(Kaler *et al.*, 2017)
	19	40218800	Seed weight 9-g5.1	NA	(Copley *et al.*, 2018)

[a] Detected in separate environments (1: 2018Ridgetown, 2:2019Ridgetwon, 3:2018Palmyra, 4:2019Palmyra, NA: Not found in any environment).

MLM: Mixed Linear Model, FarmCPU: Fixed and random model circulating probability unification, RF: Random Forest, SVR: Support Vector Regression

Table 4. The list of detected QTL for soybean total number of reproductive nodes per plant (RNP) using different GWAS methods in the tested soybean population.

GWAS Method	Chromosome	Peak SNP position	Detected QTL	Environment [a]	Reference
RF	9	40285014	Shoot Fe 1-g8.1	NA	(Dhanapal *et al.*, 2018)
			Shoot Fe 1-g8.2	NA	(Dhanapal *et al.*, 2018)
			Shoot Fe 1-g8.3	NA	(Dhanapal *et al.*, 2018)
			Shoot Fe 1-g9	NA	(Dhanapal *et al.*, 2018)
			Shoot Fe 1-g10	NA	(Dhanapal *et al.*, 2018)
			Shoot Fe 1-g11	NA	(Dhanapal *et al.*, 2018)
			Soybean mosaic virus 2-g5	NA	(Che *et al.*, 2017)
	15	34958361	SCN 5-g35	NA	(Li *et al.*, 2016)
SVR	7	1032587	WUE 2-g18	NA	(Kaler *et al.*, 2017)
	15	34958361 [1]	SCN 5-g35	1	(Li *et al.*, 2016)

[a] Detected in separate environments (1: 2018Ridgetown, 2:2019Ridgetwon, 3:2018Palmyra, 4:2019Palmyra, NA: Not found in any environment).

MLM: Mixed Linear Model, FarmCPU: Fixed and random model circulating probability unification, RF: Random Forest, SVR: Support Vector Regression

Table 5. The list of detected QTL for soybean total number of pods per plant (PP) using different GWAS methods in the tested soybean population.

GWAS Method	Chromosome	Peak position	SNP	Detected QTL	Environment [a]	Reference
RF	7	15331676		Seed weight, SoyNAM 14-g11	NA	(Xavier *et al.*, 2016b)
	19	42300695		First flower 4-g77	NA	(Mao *et al.*, 2017)
				Lodging, SoyNAM 4-g17	NA	(Cook *et al.*, 2014)
SVR	9	39366957		Pod number 1-g4.1	NA	(Fang *et al.*, 2017)
				Pod number 1-g4.2	NA	(Fang *et al.*, 2017)
				Pod number 1-g4.3	NA	(Fang *et al.*, 2017)
				Seed thickness 2-g4	NA	(Fang *et al.*, 2017)
	9	39372117		Seed Thr 2-g1	NA	(Li *et al.*, 2018b)
				Seed Ser 2-g1	NA	(Li *et al.*, 2018b)
				Seed Tyr 2-g2	NA	(Li *et al.*, 2018b)
				Seed Lys 2-g2	NA	(Li *et al.*, 2018b)
				Seed leu 2-g2	NA	(Li *et al.*, 2018b)
				Seed ile 2-g2	NA	(Li *et al.*, 2018b)
				Seed Ala 2-g2	NA	(Li *et al.*, 2018b)
				Seed Gly 2-g2	NA	(Li *et al.*, 2018b)
	11	5245870		Ureide content 1-g29	NA	(Ray *et al.*, 2015)
				Pod number 1-g6	NA	(Fang *et al.*, 2017)
		55645699		Leaflet shape 1-g4.1	NA	(Fang *et al.*, 2017)
				Leaflet shape 1-g4.2	NA	(Fang *et al.*, 2017)
				Leaflet shape 1-g4.3	NA	(Fang *et al.*, 2017)
				Seed stearic 4-g5	NA	(Li *et al.*, 2015)
				Node number 1-g6.1	NA	(Fang *et al.*, 2017)
				Node number 1-g6.2	NA	(Fang *et al.*, 2017)
				Pode number 1-g1.1	NA	(Fang *et al.*, 2017)
				Pode number 1-g1.2	NA	(Fang *et al.*, 2017)
				Pode number 1-g1.3	NA	(Fang *et al.*, 2017)
	18	55469601		WUE 3-g31	NA	(Dhanapal *et al.*, 2015a)
				Seed weight, SoyNAM 14-g28	NA	(Xavier *et al.*, 2016b)
				Lodging, SoyNAM 4-g15	NA	(Cook *et al.*, 2014)
				Branching 1-g1.1	NA	(Fang *et al.*, 2017)
				Plant height 5-g4.2	NA	(Fang *et al.*, 2017)
				Plant height 5-g4.3	NA	(Fang *et al.*, 2017)
				Shoot p 1-g30	NA	(Dhanapal *et al.*, 2018)
				Seed yield, SoyNAM 7-g19	NA	(Cook *et al.*, 2014)

		R8 full maturity, SoyNAM 13-g19	NA	(Cook *et al.*, 2014)
		Plant height 5-g4.3	NA	(Fang *et al.*, 2017)
	43077182	Seed weight 9-g5.2	NA	(Copley *et al.*, 2018)
		Seed weight 5-g21	NA	(Copley *et al.*, 2018)
		First flower 5-g3	NA	(Fang *et al.*, 2017)
		First flower 5-g17	NA	(Fang *et al.*, 2017)
19	47235604	First flower 4-g77	NA	(Mao *et al.*, 2017)
		Seed palmitic 1-g19	NA	(Priolli *et al.*, 2015)
	47350110	Leaf carotenoid content 1-g14	NA	(Dhanapal *et al.*, 2015b)
		Ureide content 1-g50.3	NA	(Ray *et al.*, 2015)
		Ureide content 1-g50.4	NA	(Ray *et al.*, 2015)
	47224293	Node number 1-g2.3	NA	(Fang *et al.*, 2017)

[a] detected in separate environments (1: 2018Ridgetown, 2:2019Ridgetwon, 3:2018Palmyra, 4:2019Palmyra, NA: Not found in any environment). MLM: Mixed Linear Model, FarmCPU: Fixed and random model circulating probability unification, RF: Random Forest, SVR: Support Vector Regression

Figure legends

Fig. 1 LD decay distance in the tested 227 soybean genotypes

Fig. 2 The distribution of seed yield (A), maturity (B), NP (C), NRNP (D), RNP (E), and PP (F)
in 227 soybean genotypes across four environments. The estimated heritability is provided for
each of the six traits. RNP: Total number of reproductive nodes per plant, NRNP: The total
number of non-reproductive nodes per plant, NP: The total nodes per plant, PP: The total number
of pods per plant.

Fig. 3 The distributions and Pearson correlations among the soybean seed yield, maturity, and
yield component traits. RNP: Total number of reproductive nodes per plant, NRNP: The total
number of non-reproductive nodes per plant, NP: The total nodes per plant, PP: The total number
of pods per plant. The heat map scale for values is provided by colour for the panel.

Fig. 4 Structure and kinship plots for the 227 soybean genotypes. The x-axis is the number of
genotypes used in this GWAS panel, and the y axis is the membership of each subgroup. G1-G7
stands for the subpopulation.

Fig. 5 Genome-wide Manhattan plots for GWAS studies of A) maturity and B) seed yield in
soybean using MLM, FarmCPU, RF, and SVR methods, from top to bottom, respectively.

Fig. 6 Genome-wide Manhattan plots for GWAS studies of A) the total number of nodes (NP)
and B) the total number of non-reproductive nodes (NRNP) in soybean using MLM, FarmCPU,
RF, and SVR methods, from top to bottom, respectively.

Fig. 7 Genome-wide Manhattan plots for GWAS studies of A) The total number of reproductive
nodes (RNP) and B) the total number of pods (PP) in soybean using MLM, FarmCPU, RF, and
SVR methods, from top to bottom, respectively.

Fig. 8 The average effects of reference allele and alternative allele from the detected SNP's peak
for seed yield (A), maturity (B), NP (C), NRNP (D), RNP (E), and PP (F) in 227 soybean
genotypes across four environments. RNP: Total number of reproductive nodes per plant, NRNP:
The total number of non-reproductive nodes per plant, NP: The total nodes per plant, PP: The
total number of pods per plant

Fig. 1 LD decay distance in the tested 227 soybean genotypes

Fig. 2 The distribution of seed yield (A), maturity (B), NP (C), NRNP (D), RNP (E), and PP (F) in 227 soybean genotypes across four environments. The estimated heritability is provided for each of the six traits. RNP: Total number of reproductive nodes per plant, NRNP: The total number of non-reproductive nodes per plant, NP: The total nodes per plant, PP: The total number of pods per plant.

Fig. 3 The distributions and Pearson correlations among the soybean seed yield, maturity, and yield component traits. RNP: Total number of reproductive nodes per plant, NRNP: The total number of non-reproductive nodes per plant, NP: The total nodes per plant, PP: The total number of pods per plant. The heat map scale for values is provided by colour for the panel.

Fig. 4 Structure and kinship plots for the 227 soybean genotypes. The x-axis is the number of genotypes used in this GWAS panel, and the y axis is the membership of each subgroup. G1-G7 stands for the subpopulation.

Fig. 5 Genome-wide Manhattan plots for GWAS studies of A) maturity and B) seed yield in soybean using MLM, FarmCPU, RF, and SVR methods, from top to bottom, respectively.

Fig. 6 Genome-wide Manhattan plots for GWAS studies of A) the total number of nodes (NP) and B) the total number of non-reproductive nodes (NRNP) in soybean using MLM, FarmCPU, RF, and SVR methods, from top to bottom, respectively.

Fig. 7 Genome-wide Manhattan plots for GWAS studies of A) The total number of reproductive nodes (RNP) and B) the total number of pods (PP) in soybean using MLM, FarmCPU, RF, and SVR methods, from top to bottom, respectively.

Fig. 8 The average effects of reference allele and alternative allele from the detected SNP's peak for seed yield (A), maturity (B), NP (C), NRNP (D), RNP (E), and PP (F) in 227 soybean genotypes across four environments. RNP: Total number of reproductive nodes per plant, NRNP: The total number of non-reproductive nodes per plant, NP: The total nodes per plant, PP: The total number of pods per plant