Neural activity in human brainstem adapts to sound-level statistics

Björn Herrmann[1,2,3]*, Sonia Yasmin[3], Kurdo Araz[3], David W. Purcell[4], & Ingrid S. Johnsrude[3,4]

[1] Rotman Research Institute,

Baycrest, M6A 2E1, North York, ON, Canada

[2] Department of Psychology,

University of Toronto, M5S 1A1, Toronto, ON, Canada

[3] Department of Psychology and Brain & Mind Institute,

The University of Western Ontario, N6A 3K7, London, ON, Canada

[4] School of Communication Sciences & Disorders,

The University of Western Ontario, N6A 5B7, London, ON, Canada

* Correspondence concerning this article should be addressed to Björn Herrmann, Rotman Research Institute, Baycrest, 3560 Bathurst St, North York, ON, M6A 2E1, Canada. E-mail: bherrmann@research.baycrest.org

Abstract

Optimal perception requires adaptation to sounds in the environment. Adaptation involves representing the acoustic stimulation history in neural response patterns, for example, by altering response magnitude or latency as sound-level statistics change. Neurons in the auditory brainstem of rodents are sensitive to acoustic stimulation history and sound-level statistics, but the degree to which the human brainstem exhibits such neural adaptation is unclear. In six electroencephalography experiments with over 125 participants, we demonstrate that acoustic stimuli within a time window of at least 40 ms are represented in response latency of the human brainstem. We further show that human brainstem responses adapt to sound-level statistical information, but that neural sensitivity to sound-level statistics is less reliable when acoustic stimuli need to be integrated over periods of ~40 ms. Our results provide evidence of adaptation to sound-level statistics in the human brainstem and of the timescale over which sound-level statistics affect neural responses to sound. The research delivers an important link to studies on neural adaptation in non-human animals.

Keywords: Human brainstem, Wave V, sound-level statistics, electroencephalography, neural adaptation

47   Introduction

Optimal sound perception requires an individual to adapt flexibly to their changing acoustic

environments[1-5]. One crucial mechanism underlying such flexibility is neural adaptation. Neural

adaptation describes a decrease in neural response magnitude and/or an increase in response latency

with sustained or repeated acoustic stimulation[6-8]. Adaptation provides a means by which the history of

acoustic information is represented in neural circuits, which, in turn, influences neural responses to

future sounds. Through adaptation, neural circuits represent when when a sound has occurred in time[9-

11] as well as statistical properties of entire acoustic feature dimensions, such as spectral variance[12-14] or

the modal/mean sound level of acoustic environments[15-17].

56   Work in rodents demonstrates that auditory nerve fibers and neurons in the brainstem and

auditory cortex are capable of adapting to environmental sound-level statistics[15,18-23]. Work in humans

has documented adaptation to sound-level statistics and sensitivity to violations of expected sound

levels in primary and secondary auditory cortices[16,24-27]. However, it is currently unclear whether

neurons at lower stages of the human auditory system, such as the brainstem, adapt to ambient sound-

level information and, if so, over what timescale statistical information is represented in neural

responses. In a series of electroencephalography (EEG) experiments, we evaluate neural adaptation in

the human brainstem to statistical structures of acoustic environments.

64   Investigations of neural adaptation often rely on manipulations of the interval duration between

65   two sounds[10,28]. The longer the interval between two sounds, the more time neurons will have to recover

66   from adaptation after responding to the first sound. Accordingly, responses to the second sound are

67   faster and larger as the interval between the two sounds increases[9-11,29-34]. Adaptation of human

68   brainstem responses appears to manifest most prominently in the response latency, while the response

69   amplitude appears to adapt less[35-38]; but see[33].

70   When neural response properties change in response to recent stimulation, this may reflect

simply the interval since the last sound, or it may reflect something more complex about the history of

stimulation, spanning a period incorporating several sounds. The latter case is more relevant for

adaptation to sound statistics. Some previous studies have demonstrated that the latency of human

brainstem responses adapts to the stimulation history beyond the influence of the directly preceding

sound[28,39], whereas others have found weak longer-term adaptation of neurons in the human

brainstem[40]. In a few instances, human brainstem activity has been shown to integrate previous sound

stimulation over several hundreds of milliseconds[28,39], suggesting that the human brainstem may be

capable of representing sound-level statistical information.

The current study investigates whether brainstem responses in humans, similar to rodents[15,21,22],

adapt to sound-level statistics. In order to investigate neural adaptation to sound-level statistics one

must first establish the extent to which acoustic stimulation history is represented in brainstem

responses, given a specific experimental setup. For example, human brainstem neurons demonstrate

adaptation over a longer window for high- compared to low-level sounds[35,41]. Accordingly, we first

conduct three EEG experiments (two detailed in the Supplemental Materials) to establish with our

experimental setup to what extent brainstem responses in human listeners adapt to stimulation history.

We use regular, predictable sound sequences to investigate adaptation of brainstem responses (two

experiments), and then use irregular, unpredictable sound sequences (one experiment) to exclude the

possibility that prediction-related processes (and not neural adaptation) underpin sensitivity of

brainstem responses to acoustic stimulation history. These three experiments enable us, in three

subsequent experiments, to investigate whether neurons in the human brainstem adapt to sound-level

statistics. This will reveal the time scale over which sound-level statistical information is represented in

human brainstem responses.

Results

Human brainstem activity is sensitive to stimulation history incorporating several sounds

A prerequisite for adaptation to sound-level statistical context is the capacity of neurons to represent

the sound history over several stimuli in their activity patterns. To investigate whether the human

brainstem is sensitive to sound history, we adapted a paradigm that has been successfully utilized to

investigate adaptation of the human auditory cortex[42,43]. In Experiment I, normal-hearing younger adults

(N=17) listened to click sequences presented at 60 dB sensation level (SL) that continuously

"accelerated" and "decelerated": the click onset-to-onset interval decreased logarithmically from 60 ms

to 10 ms and then increased back to 60 ms (Figure 1A). Analyses focused on the Wave V potential at a

latency of around 6.6 ms (Figure 1B), which is known to originate from the brainstem[41,44-47]. Wave V

amplitude and latency were binned by the preceding onset-to-onset interval, and responses were

averaged within bins. Wave V amplitude increased (t_{16} = 2.421, p = 0.028, r_e = 0.518) and latency

decreased (t_{16} = -9.621, p = 4.7×10^{-8}, r_e = 0.923) with increasing time since the previous click, as expected

(Figure 1B/C).

Figure 1: Stimulus design and results for Experiment I. A: Onset-to-onset intervals (logarithmic y-axis) of accelerating and decelerating click sequences. B: Response time courses to clicks, separately for different onset-to-onset intervals to a preceding click. Topographical distribution reflects the mean Wave V (~6.6 ms). C: Wave V amplitude and latency as a function of onset-to-onset interval. Box plots and individual data points (dots) reflect the slope from a linear function fit relating Wave V amplitude/latency to onset-to-onset intervals. Wave V latency and amplitude were sensitive to the inter-click-interval (slope significantly different from zero, p ≤ 0.05). D: Response time courses for a short (13.5 ms) and a long interval (44.5 ms), separately for decelerating and accelerating sequence contexts. Topographical distribution reflects the mean Wave V. E/F: Wave V amplitude and latency as a function of onset-to-onset interval, separately for decelerating and accelerating contexts. Box plots and individual data points (dots) reflect the mean difference between amplitudes/latencies for decelerating and accelerating contexts. Amplitudes were larger and latencies smaller in accelerating compared to decelerating contexts (p ≤ 0.05), demonstrating longer-term influences of the acoustic stimulation on the Wave V response. Error bars reflect the standard error of the mean (removal of between-subject variance[48]). *p ≤ 0.05, n.s. – not significant

In order to investigate the influence of stimulation history – beyond the interval directly

preceding a click – on brainstem responses, trials were separated into accelerating and decelerating

sequence contexts (Figure 1D). Both accelerating and decelerating contexts comprise the same onset-

to-onset intervals and, as a result, any differences in brainstem responses between contexts could only

arise from the longer-term stimulation history: clicks in the accelerating context are preceded by fewer

clicks in the same time window compared to the decelerating context, despite the same interval directly

preceding any given click. Across stimuli with different preceding intervals, Wave V latency, but not

amplitude, was smaller in the accelerating compared to the decelerating context (latency: $F_{1,16}$ = 17.831,

p = 6.5×10^{-4}, n_p^2 = 0.527; amplitude: $F_{1,16}$ = 1.23, p = 0.284, n_p^2 = 0.071; Figure 1E/F), demonstrating the

influence of longer-term stimulation history on brainstem activity. No interaction between onset-to-

onset interval and context (accelerating, decelerating) was observed for either latency or amplitude (ps

> 0.05), suggesting that the stimulation history of at least 40 ms is represented in human brainstem

responses (Figure 1F).

We replicated the effects of stimulation history on Wave V latency in Supplementary Experiment

A (N=28; using data from a previous study with a different focus[49]) using accelerating-decelerating click

sequences with onset-to-onset intervals ranging between 4 ms and 40 ms (Figure S1; Supplementary

Materials).

The effect of context on response latency may be due to the density of clicks in an extended time

period preceding the click for which the response was analyzed, or it may be related to some kind of

prediction, given the orderly presentation of stimuli we used. In Supplementary Experiment B (N=19;

Supplementary Materials), we pseudo-randomly vary onset-to-onset intervals between 4 ms and 40 ms

to examine whether the Wave V latency is indeed affected by click density in a given time period (Figure

S2). Click density had the same effect in this experiment as in the accelerating/decelerating experiment,

suggesting that Wave V latency changes are not due to any prediction-related process.

In sum, the results of Experiment I and Supplementary Experiments A and B show that the latency

of brainstem activity (Wave V) in response to a click is strongly influenced not only by the interval directly

preceding a click, but also by click density before that interval (see also[28,39]). In our data, preceding sound

information appears to be integrated over at least the preceding 40 ms (Figure 1F). Our data are

consistent with neural adaptation underlying changes in brainstem response latency. Such influence of

sound history on neural responses should enable the brainstem to be sensitive to statistical sound

features.

Human brainstem activity is affected by sound-level statistics

In Experiment II, we investigated whether neural activity in the human brainstem is sensitive to sound-

level statistics. To this end, we adapted a paradigm that we have recently used to investigate adaptation

to sound-level statistics in the human auditory cortex[16,24]. Young, normal hearing adults (N=18) listened

to clicks presented every 9 ms in two statistical contexts that differed in the modal sound level: In one

context, the modal level was 25 dB SL, whereas in the other context, the modal level was 55 dB SL (Figure

2A). Critically, the sound level of a small set of target clicks (ranging from 15 to 65 dB SL) were identical

in both contexts (black dots in Figure 2B/C) as was the level of the click directly preceding each target

click (i.e., N-1). Because target and N-1 preceding clicks were identical in the 25-dB and the 55-dB

contexts, any differences in neural responses must be due to the longer-term sound-level statistics of

the stimulation.

Figure 2: **Stimulus design and results for Experiment II. A:** Example of the two sound-level contexts (distributions with either a 25-dB or a 55-dB modal sound level) utilized to investigate adaptation to sound-level statistics. **B/C:** Examples of sound-level stimulation in the two statistical contexts. Each dot represents the sound level of one click. The black dots represent the sound level of clicks that were identical in both statistical contexts and for which the sound level of the directly preceding click was also identical in both contexts. **D:** Response time courses to clicks were binned according to their sound level. Topographical distribution reflects the mean Wave V. **E/F:** Wave V amplitude and latency as a function sound level (binned into five categories). Box plots and individual data points (dots) reflect the slope from a linear function fit relating Wave V amplitude/latency to sound level. **G:** Response time courses for low (25.7 dB SL) and high sound levels (54.3 dB SL), separately for the sound-level context with a mode of 25 dB SL and a mode of 55 dB SL. Topographical distribution reflects the mean Wave V. **H/I:** Wave V amplitude and latency as a function sound level, separately for the 25-dB and the 55-dB context. Box plots and individual data points (dots) reflect the mean difference between amplitudes/latencies for the 25-dB and the 55-dB context. Latencies were smaller in the 25-dB compared to the 55-dB context (p ≤ 0.05), demonstrating adaptation to sound-level

statistics in the human brainstem. Error bars reflect the standard error of the mean (removal of between-subject variance[48]). *p ≤ 0.05, n.s. – not significant

We first confirmed that the sound level of clicks modulates the Wave V. All trials, independent

of statistical context, were binned into five sound-level categories (centered on 20, 30, 40, 50, and 60

dB SL) and averaged. Figure 2D depicts the time courses for each sound-level category. A linear function

was fit to Wave V amplitude and latency as function of sound level, separately for each participant, and

the resulting linear coefficient was tested against zero. Consistent with previous work, Wave V amplitude

increased (t_{17} = 13.289, p = 2.1×10^{-10}, r_e = 0.955; Figure 2E) and Wave V latency decreased (t_{17} = -11.061,

p = 3.5×10^{-9}, r_e = 0.937; Figure 2F) with increasing sound level[32,41,50,51].

Next, we limited the analysis to target clicks – that is, the clicks for which level, and the level of

the immediately preceding click were identical in both statistical contexts (25-dB and 55-dB contexts;

black dots in Figure 2B). A repeated-measures ANOVA (rmANOVA) revealed that Wave V amplitude

increased with click level, consistent with the previous analysis ($F_{4,68}$ = 59.103, p < 0.001, n_p^2 = 0.777),

but there was no effect of statistical context nor an interaction between target click level and context

(both ps > 0.5; Figure 2H). Wave V latencies were earlier for clicks presented at a high compared to low

level, consistent with the previous analysis ($F_{4,68}$ = 76.544, p < 0.001, n_p^2 = 0.818) and were also earlier

for clicks presented in the 25-dB compared to the 55-dB context ($F_{1,17}$ = 18.077, p = 5.4×10^{-4}, n_p^2 = 0.515).

Again, no interaction was observed between target click level and context (p > 0.5; Figure 2I). Experiment

II shows that the latency, but not amplitude, of the human brainstem Wave V is sensitive to sound-level

statistics.

Sensitivity of the brainstem to sound-level statistics decreases with longer intervals between stimuli

169   In Experiment II, we used an onset-to-onset interval of 9 ms and the levels of both target and preceding

170   clicks were constant across contexts. Given this timing, contexts could differ 18 ms prior to target clicks.

171   In order to examine the time window over which sound-level statistics affect human brainstem

172   responses, we conducted Experiments III (N=16) and IV (N=19) using the same experimental design as

173   in Experiment II (Figure 2A-C), but with onset-to-onset intervals of 14 ms and 22 ms, respectively.

Figure 3: Neural adaptation to sound-level statistics for different click presentation rates. A: Wave V latencies for target clicks with different sound levels in two statistical contexts (25-dB SL and 55-dB SL contexts) and for different click presentation rates (9 ms [Experiment II], 14 ms [Experiment III], 22 ms [Experiment IV]). **B:** Overall Wave V latency for target clicks presented at different presentation rates, independent of sound level. **C:** Latency difference between statistical contexts (55-dB context minus 25-dB context) for each target click sound level (left) and averaged across target click sound levels (right). Asterisks below the bar graphs indicate a significant difference from zero (i.e., between the two statistical contexts). Error bars reflect the standard error of the mean. *p ≤ 0.05, n.s. − not significant.

Wave V latencies for the 25-dB and the 55-dB contexts for each of the three onset-to-onset

intervals (9, 14, 22 ms) are shown in Figure 3A. Mean latency, independent of context, decreased with

increasing onset-to-onset interval (9 ms vs. 14 ms: t_{32} = 4.392, p = 1.1×10^{-4}, r_e = 0.613; 9 ms vs. 22 ms:

t_{35} = 6.266, p = 3.4×10^{-7}, r_e = 0.727; Figure 3B), indicating more adaptation of brainstem responses for

short intervals between clicks (see also Experiment I and Supplementary Experiments A and B[28,31,41]).

The difference between statistical contexts (55-dB minus 25-dB context), indicating adaptation

to sound-level statistics, was significant for 9 ms (t_{17} = 4.252, p = 5.4×10^{-4}, r_e = 0.718) and 14 ms (t_{15} =

4.381, p = 5.4×10^{-4}, r_e = 0.749) onset-to-onset intervals, but not for 22 ms onset-to-onset interval (t_{18} =

1.163, p = 0.260, r_e = 0.264; Figure 3C); although the absence of a context effect for the 22-ms interval

appears to be driven by one outlier data point (Figure 3C, right), as its removal (t_{17} = 3.428, p = 0.003, r_e

= 0.639) and a non-parametric test (p = 0.049, df = 18) revealed a significant difference. The effect of

statistical context (55-dB minus 25-dB context) was larger for the 9 ms compared to the 22 ms onset-to-

onset interval (t_{35} = 2.721, p = 0.01, r_e = 0.418; Figure 3C), but did not differ between onset-to-onset

intervals of 9 ms and 14 ms (t_{32} = 1.554, p = 0.13, r_e = 0.265; Figure 3C) or 14 ms and 22 ms (t_{33} = 1.601,

p = 0.119, r_e = 0.269; Figure 3C).

The results from Experiments II-IV demonstrate that human brainstem responses are sensitive
to sound-level statistics in near continuous stimulation (i.e., for onset-to-onset intervals of 9 ms and 14
ms), but less so when sounds are more spread out in time (for the 22-ms onset-to-onset interval). That
brainstem responses were sensitive to sound-level statistics for the 14 ms interval – that is, to contextual
differences in sound levels 28 ms or further in the past – but less for the 22-ms interval (i.e., contextual
differences of 44 ms or more) suggests that sound-level information over at least a period of 28 ms (but
less than 44 ms) are represented in human brainstem responses. This is also consistent with the
identified temporal evolution of neural adaptation observed in Experiments I and Supplementary
Experiments A and B (i.e., integration of ~40 ms of stimulation history).

Discussion

In the current study, we conducted a series of four electroencephalography experiments and two
supplementary experiments to investigate the extent to which neural units in the human brainstem
represent the history of acoustic stimuli, whether the human brainstem is sensitive to sound-level
statistics of acoustic environments, and the timescale over which statistical information is integrated by
the brainstem. We show that acoustic stimuli within a time window of at least 40 ms are represented in
human brainstem responses through neural adaptation (Experiment I and Supplementary Experiment
A), while we exclude prediction-related processes as an underlying mechanism (Supplementary
Experiment B). We further demonstrate that brainstem circuits represent sound-level statistics
(Experiment II and III) and that the sensitivity to sound-level statistics decreases as stimuli are more
spread out in time (Experiment IV). Our results provide information about the timecourses of neural
adaptation in the human brainstem and are an important link to studies on neural adaptation in non-
human animals.

Human brainstem responses are sensitive to stimulation history spanning several sounds

Adaptation to sound-level statistics or other properties of the acoustic environment requires that
neurons represent stimulation history across several stimuli. For Experiment I and Supplementary
Experiments A and B, we adapted experimental paradigms from our previous work on longer-term
adaptation in the human auditory cortex[42,43] to investigate the extent to which stimulation history is

reflected in adaptation of human brainstem responses. We observed that the Wave V of the auditory

evoked potential – originating from the brainstem[41,44-47] – peaked later when more clicks occurred within

a preceding time window ("decelerating" sequence) compared to fewer clicks ("accelerating"

sequence), despite an identical interval directly preceding each target click and identical sound levels

(Experiments I and Supplementary Experiment A; Figures 1 and S1). We estimate that, given the current

stimulation protocols, acoustic stimuli over a period of at least 40 ms are represented in human

brainstem responses (e.g., Figure 1F). Finally, showing that the longer-term stimulation history affects

the Wave V latency in sequences with clicks occurring at random, unpredictable times (Supplementary

Experiment B), rules out that prediction-related processes underlie the changes in Wave V latency. Our

data are more consistent with neural adaptation by, for example, affecting the spike latency of brainstem

neurons[6,52].

Our results demonstrating that Wave V latency is sensitive to stimulation history are consistent

with previous work, using isochronous click sequences and click trains, showing that Wave V latency

increases with increasing click rate[38,41] and that Wave V latency is affected by the stimulation history

spanning several sounds[28,35,39]. The degree to which the longer-term stimulation history modulates

Wave V latency varies across studies[28,35,36,40] and appears to depend to some extent on the sound level

at which acoustic stimuli are presented[35,41]. Louder sounds appear to have a longer-term influence on

the latency of human brainstem responses than softer sounds[35]. Given this prior work, we expected to

observe adaptation that reflected stimulation history in the current study. However, the variability in

previous work and the potential dependence on experimental parameters, such as sound level, required

a detailed investigation in Experiment I and Supplementary Experiments A and B to establish effects of

stimulation history before investigating adaptation to sound-level statistics. We show here that

brainstem responses represent the acoustic stimulation history in accelerating-decelerating sequences

and random-interval sequences for at least 40 ms.

Human brainstem responses adapt to sound-level statistics

In three EEG experiments, we investigated the extent to which neurons in the human brainstem adapt

to sound-level statistics. We utilized a paradigm in which participants listened to clicks in two contexts,

where the sound level was drawn from distributions with different modal sound levels (25 dB SL and 55

dB SL) and where target clicks and clicks preceding target clicks were identical in the two contexts with

different modal sound levels[16,24]. Controlling the immediate history of clicks across contexts allowed us

to investigate the effect of longer-term sound-level statistics on brainstem responses. We observed that

Wave V latency, but not amplitude was affected by sound-level context, such that Wave V peaked later

in the 55-dB context compared to the 25-dB context (Figures 2 and 3). This context effect was present

when at least 28 ms of sound stimulation prior to target clicks was identical in both statistical contexts

(Figure 3C; Experiment III; 14 ms onset-to-onset interval), but was less reliable when at least 44 ms of

stimulation history prior to target clicks was identical between sound-level contexts (Figure 3C;

Experiment IV; 22 ms onset-to-onset interval). Our data suggest that human brainstem responses

represent sound-level statistics at time scales of at least 28 ms, and to a higher degree when stimuli are

closer together in time (9 ms onset-to-onset interval).

255   Our observation of sensitivity to sound-level statistics in the human brainstem is consistent with

256   previous findings of adaptation to sound-level statistics in rodent inferior colliculus[15,18,21,22] and auditory

257   nerve fibers[19,20], and with work on the human auditory cortex[16,24-27]. Our findings also complement a

258   broader literature showing adaptation to other features and distributions of acoustic environments, in

259   addition to sound level, such as acoustic contrast[53,54], mean and variance of interaural level[55], spectral

260   variance[6,12,13], and other complex distributional properties[14].

261   Previous work suggests that timescales of adaptation increase along the auditory pathway, with

262   shorter timescales in the brainstem compared to the auditory cortex[54]. This is consistent with the

263   relatively short timescale in the range of a few tens of milliseconds over which the human brainstem

264   integrates sound-level statistics in the current study, and previous work that demonstrates that the

265   human auditory cortex shows adaptation lasting for 10 s or more[9,10,30].

266   One difference between the current work and previous work is that we show adaptation of

response latency, but not amplitude, to sound-level statistics (Figure 2H). Wave V amplitude increased

with increasing sound level but was not sensitive to statistical context. In contrast, previous

investigations of adaptation to sound-level statistics report sensitivity of firing rate in non-human

mammals or magneto-/electroencephalographic response magnitude in humans[15,16,18,21,22,24], whereas

no effects of spike latency or response latency were reported; but see[6]. Although it is well known that

Wave V amplitude adapts less than Wave V latency[32,36], the reasons for this are unknown. One previous

study revealed frequency-specific adaptation of spike latency in the inferior colliculus of rats, but no

difference in latency adaptation between two statistical contexts differing in spectral variance[6].

However, in this previous study, stimulus presentation was much slower (at an onset-to-onset interval

of 300 ms) and so no context effect on latency would be expected, based on the current results. Future

work in rodents may investigate more closely whether spike latency also adapts to sound-level statistics

to enable more direct comparisons with the current study.

Conclusions

In the current study, we investigated the extent to which neural responses in the human brainstem

adapt to the acoustic stimulation history beyond the preceding stimulus and whether the sound-level

statistical context is represented in the response. We also explored over which timescale statistical

information is reflected in human brainstem activity. We demonstrate that acoustic stimuli within a time

window of at least 40 ms are represented in the latency of human brainstem responses, likely via neural

adaptation mechanisms but not through prediction-related processes. We further show that brainstem

responses represent sound-level statistical information, but that this neural sensitivity to sound-level

statistics decreases when acoustic information over periods beyond ~40 ms need to be integrated. Our

results demonstrate the timescale of neural adaptation in the human brainstem and provide a link to

studies on neural adaptation in non-human animals.

Methods and Materials

Consent and ethics approval

A hundred and twenty-eight normal hearing adults (17–32 years) participated in the six

electroencephalography experiments of the current study. Demographic information about participants

is provided separately for each experiment. Four of the experiments are detailed here, whereas two

others – Supplementary Experiments A and B – are presented in the supplementary materials.

Participants gave prior written informed consent and were paid $5 CAD per half-hour for their

participation or received course credits. The study was conducted in accordance with the Declaration of

Helsinki, the Canadian Tri-Council Policy Statement on Ethical Conduct for Research Involving Humans

(TCPS2-2014), and was approved by the Nonmedical Research Ethics Board of the University of Western

Ontario (protocol ID: 106570). Participants reported no neurological disease or hearing problems and

were naïve to the purposes of the experiment.

General procedure and acoustic stimulation setup

Sounds used in the current series of experiments consisted of clicks of 0.1 ms duration (rectangle

function) presented at different onset-to-onset intervals and sound levels depending on the specific

experiment described below. Click polarity was inverted on half of the click presentations and polarity

inversion was tightly controlled across conditions within each experiment. Clicks were presented to the

right-ear via an Etymotic ER1 insert earphone, using a Fireface 400 external sound card controlled by

Psychtoolbox (version 3.14) in MATLAB (MathWorks Inc.). The sensation level for click trains was

determined for each participant using a method of limits procedure[56], and all experimental stimuli were

presented relative to this sensation level (see details below).

Electroencephalographic recordings and preprocessing

Electroencephalographic activity was recorded using a 16-channel BioSemi (Active 2) system. Additional

electrodes were placed at the left and right mastoids. EEG was recorded at 16,384 Hz with an online

low-pass filter of 3,334 Hz. Electrodes were referenced online to a monopolar reference feedback loop

connecting a driven passive sensor and a common-mode-sense (CMS) active sensor, both located

posteriorly on the scalp.

MATLAB software (v7.14; MathWorks, Inc.) was used for offline data analysis. Raw data were

filtered with an elliptic filter to suppress line noise at 60 Hz. Data were re-referenced by averaging the

two mastoid channels and subtracting the average separately from each of the 16 channels. Data were

high-pass filtered at 130 Hz (2743 points, Hann window) and low-pass filtered at 2500 Hz (101 points,

Hann window). Data were divided into epochs ranging from 2 ms pre-click-onset to 10 ms post-click

onset. Epochs were excluded if the signal change exceeded 60 µV in the 0 to 10 ms time window in

either the Cz or C3 electrode. These electrodes were of particular interest because they are known to

capture the Wave V most strongly, given right ear acoustic stimulation[57]. The peak latency of the Wave

V was identified, and Wave V amplitude calculated by averaging the amplitude within a 0.2 ms time

window centered on Wave V latency.

Experiment I

Participants. Seventeen individuals participated in Experiment I (age range: 18–28 years; median age:

22 years; 11 females). Data from three additional participants were recorded but excluded at the analysis

stage, because Wave V was absent (N=2) or an atypically late Wave V latency was exhibited, peaking

later than 8.5 ms (N=1). Results were unaffected by inclusion vs. exclusion of the latter participant.

Acoustic stimulation. Participants listened to six blocks of stimulation (each about 8 min long) that

comprised click-train stimuli, in which the onset-to-onset interval of clicks changed from 60 ms to 10 ms

(in 7 logarithmically-spaced steps) and back from 10 ms to 60 ms (in 7 logarithmically-spaced steps),

creating an accelerating-decelerating stimulation cycle (Figure 1A; see also[42,43]). Across the six blocks,

participants listened to 7,200 accelerating-decelerating stimulation cycles. All clicks were presented at

60 dB SL.

Data analysis. Single trials were separated into an 'accelerating' context (60 ms to 10 ms) and a

'decelerating' context (10 ms to 60 ms). As such, each context (accelerating, decelerating) comprised

the same onset-to-onset intervals and, consequently, any response differences between contexts

cannot be attributed to the duration of interval directly preceding a click, but must be attributed to the

longer-term stimulation history (in which the decelerating context compared to the accelerating context

comprised more clicks in the same preceding time). Single-trial time courses for a unique onset-to-onset

interval and one of its direct neighbors (i.e., shorter and longer intervals) were binned and averaged[43,58].

The Wave V latency and amplitudes were calculated for each onset-to-onset interval and each context

(accelerating, decelerating). For statistical analyses, repeated-measured analyses of variance

(rmANOVA) with the factors Interval and Context (accelerating, decelerating) were calculated separately

for the Wave V latency and the Wave V amplitude.

Experiment II

Participants. Eighteen individuals participated in Experiment II (age range: 17–32 years; median age: 18

years; 15 females). None of them had participated in Experiment I. Data from two additional participants

were excluded because of large stimulus artifacts in the recordings (N=1) and the absence of an

identifiable Wave V (N=1).

Acoustic stimulation. The experimental paradigm was derived from our previous work on neural

adaptation to sound-level statistics in the auditory cortex[16,24]. Clicks were presented in two types of

contexts that differed with respect to the sound-level distributions from which a click's sound level was

drawn: for one context, the modal sound level was 25 dB SL, for the other context it was 55 dB SL (Figure

2A). To investigate the effects of sound-level statistical context on human brainstem responses, we

eliminated the confounding effects of different acoustics by ensuring that the clicks for which we

analyzed neural responses as well as their local context (i.e., directly preceding clicks) were identical

across contexts (i.e., 25 dB SL vs. 55 dB SL), whereas the longer-term sound-level statistics were different

between the two contexts.

Participants listened to six blocks, each approximately 7 min long. In each block, 44,800 clicks

were presented at a constant onset-to-onset interval of 9 ms. In each block, eight target clicks with

sound levels ranging from 15 dB SL to 65 dB SL (step size: 7.143 dB SL) were each presented 960 times,

yielding 7,680 target clicks per block. These stimuli were identical across all six blocks (black dots in

Figure 2B/C). The 7,680 clicks immediately preceding each of these $8 \cdot 960 = 7,680$ target clicks were

also fixed, such that for each of the 960 presentations of one of the eight sound levels, the sound level

of the preceding tone took on one of 30 sound levels (range: 15 dB SL to 65 dB SL; step size: 1.724 dB

SL) without replacement. Thus, the same 7,680 pairs of experimental clicks were presented in each

block.

The sound levels for the remaining 29,440 clicks (out of the 44,800 clicks per block) were chosen

randomly (range: 15 dB SL to 65 dB SL; step size: 0.1 dB SL) depending on the sound-level distribution

(Figure 2A). For three out of the six blocks, sound levels were randomly chosen from a sound-level

distribution with a 25-dB high-probability region (Figure 2B). For the other three blocks, sound levels

were randomly chosen from a sound-level distribution with a 55-dB high-probability region (Figure 2C).

High probability regions had a width of 10 dB centered on 25 dB SL or 55 dB SL (Figure 2A-C). The 15,360

experimental clicks (i.e., 7,680 pairs) and the 29,440 filler clicks were randomly intermixed in each block

and presented such that at least one filler tone occurred between each of the 7,680 pairs of

experimental clicks. Blocks with different sound-level contexts alternated, and the starting sound-level

context was counter-balanced across participants. Analysis of the influence of sound-level statistical

context on neural responses focused on the second click in each pair of experimental clicks: these were

identical across the two sound-level distributions (black dots in Figure 2B/C).

Data analysis. The number of trials for each of the 8 sound levels in the different contexts was relatively

low for the analysis of the small Wave V brainstem potential (2880 trials = 960 trials × 3 blocks). In order

to increase the number of trials in the response average, thereby increasing the signal-to-noise ratio[16,58],

we averaged trials for four adjacent sound levels out of the 8 sound levels (separately for the 25-dB SL

context and the 55-dB SL context). For example, single trials for sounds with the 15, 22.1, 29.3, and 36.4

dB SL sound level were averaged. Averaging trials for four adjacent sound levels reduced the number of

unique sound levels from 8 to 5, while drastically increasing the number of trials in a unique response

average. Our choice of smoothing across different sound-level conditions also ensured that a sufficiently

high number of trials were averaged in Experiments III and IV, in which longer onset-to-onset intervals

reduced the number trials per sound level that were presented (see below). Wave V latency was

extracted for each sound level and statistical context, and a rmANOVA with factor Sound Level and

Context (25-dB SL, 55-dB SL) was calculated.

Experiment III

Participants. Sixteen individuals participated in Experiment III (age range: 18–32 years; median age: 18.5

years; 12 females). None of them had participated in Experiment I or II.

Acoustic stimulation. Stimulation procedures were identical to Experiment II with the following

exceptions. Participants listened to six blocks, each approx. 8.5 min long. In each block, 36,400 clicks

were presented at a constant onset-to-onset interval of 14 ms. 6,240 experimental click pairs (8 sound

levels × 780 trials) and 23,920 filler clicks were presented per block.

Data analysis. The same analyses as for Experiment II were carried out.

Experiment IV

Participants. Nineteen individuals participated in Experiment IV (age range: 17–23 years; median age:

18 years; 13 females and 5 males; one person did not provide information about their sex). None of

them had participated in any of the other experiments.

Acoustic stimulation. Stimulation procedures were identical to Experiment II with the following

exceptions. Participants listened to six blocks, each approx. 8.3 min long. In each block, 22,400 clicks

were presented at a constant onset-to-onset interval of 22 ms. 3,840 experimental click pairs (8 sound

levels × 480 trials) and 14,720 filler clicks were presented per block.

Data analysis. The same analyses as for Experiment II were carried out.

Effect sizes

Throughout the manuscript, effect sizes are provided as partial eta squared (η_p^2) for repeated-measures

ANOVAs and as r_e ($r_{equivalent}$) for t-tests[59].

416   **Acknowledgements**

417   BH was supported by a BrainsCAN postdoctoral fellowship (Canada First Research Excellence Fund;

418   CFREF) and by the Canada Research Chair program.

419   **Author contributions**

420   The experiments were performed in the Brain and Mind Institute at the University of Western Ontario,

421   London, Canada. B.H., S.Y.., I.S.J. designed the study. B.H., S.J., K.A. recorded the data. B.H. programmed

422   the experiments and analyzed the data. S.Y. and D.P. supported data analysis. B.H., S.Y., D.P., and I.S.J.

423   interpreted the results. B.H. wrote the manuscript. All authors reviewed and revised the manuscript.

424   **Competing interests**

425   The authors declare no competing interests.

426   **Data availability**

427   The datasets generated during the current study are not publicly available because participant consent

428   for public-data sharing was not obtained. Data are available from the corresponding author on

429   reasonable request.

430   **References**

1   Barlow, H. B. in *Sensory Communication* (ed W. Rosenblith) 217-234 (MIT Press., 1961).

2   Kluender, K. R., Stilp, C. E. & Kiefte, M. in *Vowel Inherent Spectral Change*   (eds Geoffrey Stewart
Morrison & Peter F. Assmann) 117-151 (Springer, 2013).

3   Lewicki, M. S. Efficient coding of natural sounds. *Nature* **5**, 356-363 (2002).

4   Wark, B., Lundstrom, B. N. & Fairhall, A. Sensory adaptation. *Current Opinion in Neurobiology* **17**,
423-429 (2007).

5   Whitmire, C. J. & Stanley, G. B. Rapid Sensory Adaptation Redux: A Circuit Perspective. *Neuron* **92**,
298-315 (2016).

6   Herrmann, B., Parthasarathy, A., Han, E. X., Obleser, J. & Bartlett, E. L. Sensitivity of rat inferior
colliculus neurons to frequency distributions. *Journal of Neurophysiology* **114**, 2941-2954 (2015).

7   Nelken, I. Stimulus-specific adaptation and deviance detection in the auditory system: experiments
and models. *Biological Cybernetics* **108**, 655-663 (2014).

8   Nelken, I. & Ulanovsky, N. Mismatch Negativity and Stimulus-Specific Adaptation in Animal Models.
Journal of Psychophysiology **21**, 214-223 (2007).

9   Sams, M., Hari, R., Rif, J. & Knuutila, J. The Human Auditory Sensory Memory Trace Persists about
10 sec: Neuromagnetic Evidence. *Journal of Cognitive Neuroscience* **5**, 363-370 (1993).

10   Hari, R., Kaila, K., Katila, T., Tuomisto, T. & Varpula, T. Interstimulus interval dependence of the
auditory vertex response and its magnetic counterpart: Implications for their neural generation.
Electroencephalography and clinical Neurophysiology **54**, 561-569 (1982).

11   Butler, R. A. The cumulative effects of different stimulus repetition rates on the auditory evoked
response in man. *Electroencephalography and clinical Neurophysiology* **35**, 337-345 (1973).

12   Herrmann, B., Henry, M. J. & Obleser, J. Frequency-specific adaptation in human auditory cortex
depends on the spectral variance in the acoustic stimulation. *Journal of Neurophysiology* **109**, 2086-
2096 (2013).

13   Garrido, M. I., Sahani, M. & Dolan, R. J. Outlier Responses Reflect Sensitivity to Statistical Structure
in the Human Brain. *PLoS Computational Biology* **9**, e1002999 (2013).

14   Kvale, M. N. & Schreiner, C. E. Short-Term Adaptation of Auditory Receptive Fields to Dynamic
Stimuli. *Journal of Neurophysiology* **91**, 604-612 (2004).

15   Dean, I., Harper, N. S. & McAlpine, D. Neural population coding of sound level adapts to stimulus
statistics. *Nature Neuroscience* **8**, 1684-1689 (2005).

16   Herrmann, B., Maess, B. & Johnsrude, I. S. Aging Affects Adaptation to Sound-Level Statistics in
Human Auditory Cortex. *The Journal of Neuroscience* **38**, 1989-1999 (2018).

17   Hildebrandt, K. J., Benda, J. & Hennig, R. M. Multiple Arithmetic Operations in a Single Neuron: The
Recruitment of Adaptation Processes in the Cricket Auditory Pathway Depends on Sensory Context.
The Journal of Neuroscience **31**, 14142-14150 (2011).

18   Dean, I., Robinson, B. L., Harper, N. S. & McAlpine, D. Rapid Neural Adaptation to Sound Level
Statistics. *The Journal of Neuroscience* **28**, 6430-6438 (2008).

19   Wen, B., Wang, G. I., Dean, I. & Delgutte, B. Dynamic Range Adaptation to Sound Level Statistics in
the Auditory Nerve. *The Journal of Neuroscience* **29**, 13797-13808 (2009).

20   Wen, B., Wang, G. I., Dean, I. & Delgutte, B. Time course of dynamic range adaptation in the auditory
nerve. *Journal of Neurophysiology* **108**, 69-82 (2012).

21   Robinson, B. L., Harper, N. S. & McAlpine, D. Meta-adaptation in the auditory midbrain under
cortical influence. *Nature Communications* **7**, 13442 (2016).

22   Bakay, W. M. H., Anderson, L. A., Garcia-Lazaro, J. A., McAlpine, D. & Schaette, R. Hidden hearing
loss selectively impairs neural adaptation to loud sound environments. *Nature Communications* **9**,
4298 (2018).

23   Scholl, B., Gao, X. & Wehr, M. Level Dependence of Contextual Modulation in Auditory Cortex.
Journal of Neurophysiology **99**, 1616-1627 (2008).

24   Herrmann, B., Augereau, T. & Johnsrude, I. S. Neural Responses and Perceptual Sensitivity to Sound
Depend on Sound-Level Statistics. *Scientific Reports* **10**, 9571 (2020).

25   Näätänen, R., Paavilainen, P., Alho, K., Reinikainen, K. & Sams, M. Do event-related potentials reveal
the mechanism of the auditory sensory memory in the human brain? *Neuroscience Letters* **98**, 217-
221 (1989).

26   Schröger, E. & Winkler, I. Presentation rate and magnitude of stimulus deviance effects on human
pre-attentive change detection. *Neuroscience Letters* **193**, 185-188 (1995).

27   Althen, H., Grimm, S. & Escera, C. Fast Detection of Unexpected Sound Intensity Decrements as
Revealed by Human Evoked Potentials. *PLoS ONE* **6**, e28522 (2011).

28   Lasky, R. E., Maier, M. M. & Hecox, K. Auditory Evoked Brain Stem Responses to Trains of Stimuli in
Human Adults. *Ear & Hearing* **17**, 544-551 (1996).

29   Thornton, A. R. & Coleman, M. J. The adaptation of cochlear and brainstem auditory evoked
potentials in humans. *Electroencephalography and Clinical Neurophysiology* **39**, 399-406 (1975).

30   Zacharias, N., König, R. & Heil, P. Stimulation-history effects on the M100 revealed by its differential
dependence on the stimulus onset interval. *Psychophysiology* **49**, 909-919 (2012).

31   Burkard, R. F. & Sims, D. The Human Auditory Brainstem Response to High Click Rates: Aging Effects.
American Journal of Audiology **10**, 1-9 (2001).

32   Pratt, H. & Sohmer, H. Intensity and Rate Functions of Cochlear and Brainstem Evoked Responses
to Click Stimuli in Man. *Archives of Oto-Rhino-Laryngology* **212**, 85-92 (1976).

33   Skoe, E. & Tufts, J. Evidence of noise-induced subclinical hearing loss using auditory brainstem
responses and objective measures of noise exposure in humans. *Hearing Research* **361**, 80-91
(2018).

34   Yagi, T. & Kaga, K. The Effect of the Click Repetition Rate on the Latency of the Auditory Evoked
Brain Stem Response and its Clinical Use for a Neurological Diagnosis. *Archives of Oto-Rhino-
Laryngology* **222**, 91-97 (1979).

35   Ballachanda, B. B., Moushegian, G. & Stillman, R. D. Adaptation of the Auditory Brainstem Response:
Effects of Click Intensity, Polarity, and Position. *Journal of the American Academy of Audiology* **3**,
275-282 (1992).

36   Polyakov, A. & Pratt, H. The cumulative effect of high click rate on monaural and binaural processing
in the human auditory brainstem. *Clinical Neurophysiology* **114**, 366-375 (2003).

37   Scott, M. L. & Harkins, S. W. Amplitude of the Brainstem Auditory Evoked Response: The Effect of
Interstimulus Interval. *International Journal of Neuroscience* **8**, 147-152 (1978).

38   Harkins, S. W., McEvoy, T. M. & Scott, M. L. Effects of interstimulus interval on latency of the
brainstem auditory evoked potential. *International Journal of Neuroscience* **10**, 7-14 (1979).

39   Lasky, R. E. Rate and adaptation effects on the auditory evoked brainstem response in human
newborns and adults. *Hearing Research* **111**, 165-176 (1997).

40   Wynne, D. P. *et al.* Loudness adaptation accompanying ribbon synapse and auditory nerve
disorders. *Brain* **136**, 1626-1638 (2013).

41   Don, M., Allen, A. R. & Starr, A. Effect of Click Rate on the Latency of Auditory Brain Stem Responses
in Humans. *Annals of Otology, Rhinology & Laryngology* **86**, 186-195 (1977).

42   Herrmann, B., Buckland, C. & Johnsrude, I. S. Neural signatures of temporal regularity processing in

520   sounds differ between younger and older adults. *Neurobiology of Aging* **83**, 73-85,

521   doi:https://doi.org/10.1101/522375 (2019).

522   43   Herrmann, B., Henry, M. J., Johnsrude, I. S. & Obleser, J. Altered temporal dynamics of neural

523   adaptation in the aging human auditory cortex. *Neurobiology of Aging* **45**, 10-22 (2016).

524   44   Parkkonen, L., Fujiki, N. & Mäkelä, J. P. Sources of auditory brainstem responses revisited:

525   Contribution by magnetoencephalography. *Human Brain Mapping* **30**, 1772-1782 (2009).

526   45   Møller, A. R. Neural Generators of the Brainstem Auditory Evoked Potentials. *Seminars in Hearing*

527   **19**, 11-27 (1998).

528   46   Møller, A. R. & Jannetta, P. J. Interpretation of brainstem auditory evoked potentials: results from

529   intracranial recordings in humans. *Scandinavian Audiology* **12**, 125-133 (1983).

530   47   Møller, A. R. & Jannetta, P. J. Evoked potentials from the inferior colliculus in man.

531   *Electroencephalography and Clinical Neurophysiology* **53**, 612-620 (1982).

532   48   Masson, M. E. J. & Loftus, G. R. Using confidence intervals for graphically based data interpretation.

533   *Canadian Journal of Experimental Psychology and Aging* **57**, 203-220 (2003).

534   49   Yasmin, S., Purcell, D. W., Veeranna, S. A., Johnsrude, I. S. & Herrmann, B. A novel approach to

535   investigate subcortical and cortical sensitivity to temporal structure simultaneously. *Hearing*

536   *Research* **398**, 108080 (2020).

537   50   Verhulst, S., Jagadeesh, A., Mauermann, M. & Ernst, F. Individual Differences in AuditoryBrainstem

538   Response Wave Characteristics:Relations to Different Aspectsof Peripheral Hearing Loss. *Trends in*

539   *Hearing* **20**, 1-20 (2016).

540   51   Stockard, J. E., Stockard, J. J., Westmoreland, B. F. & Corfits, J. L. Brainstem auditory-evoked

541   responses: Normal variation as a function of stimulus and subject characteristics. *Archives of*

542   *Neurology* **36**, 823-831 (1979).

543   52   Heil, P. First-spike latency of auditory neurons revisited. *Current Opinion in Neurobiology* **14**, 461-

544   647 (2004).

545   53   Rabinowitz, N. C., Willmore, B. D. B., Schnupp, J. W. H. & King, A. J. Contrast Gain Control in Auditory

546   Cortex. *Neuron* **70**, 1178-1191 (2011).

54   Lohse, M., Bajo, V. M., King, A. J. & Willmore, B. D. B. Neural circuits underlying auditory contrast
gain control and their perceptual implications. *Nature Communications* **11**, 324 (2020).

55   Dahmen, J. C., Keating, P., Nodal, F. R., Schulz, A. L. & King, A. J. Adaptation to Stimulus Statistics in
the Perception and Neural Representation of Auditory Space. *Neuron* **66**, 937-948 (2010).

56   Herrmann, B. & Johnsrude, I. S. Attentional State Modulates the Effect of an Irrelevant Stimulus
Dimension on Perception. *Journal of Experimental Psychology: Human Perception and Performance*
44, 89-105 (2018).

57   Picton, T. W. in *Human auditory evoked potentials*   (ed Terence W. Picton) 213-246 (Plural
Publishing, 2010).

58   Ingham, N. J. & McAlpine, D. GABAergic Inhibition Controls Neural Gain in Inferior Colliculus
Neurons Sensitive to Interaural Time Differences. *The Journal of Neuroscience* **25**, 6187-6198
(2005).

59   Rosenthal, R. & Rubin, D. B. requivalent: A Simple Effect Size Indicator. *Psychological Methods* **8**,
492-496 (2003).

561