Experience-dependent weakening of callosal synaptic connections in the absence of postsynaptic FMRP

Zhe Zhang, *Jay R. Gibson and *Kimberly M. Huber

Department of Neuroscience, O'Donnell Brain Institute, University of Texas Southwestern Medical Center, Dallas, TX, United States.

*Correspondence: Jay.Gibson@UTSouthwestern.edu and Kimberly.Huber@UTSouthwestern.edu

Abstract

Reduced structural and functional interhemispheric connectivity correlates with the severity of Autism Spectrum Disorder (ASD) behaviors in humans. Little is known of how ASD-risk genes regulate callosal connectivity. Here we show that *Fmr1*, whose loss-of-function leads to Fragile X Syndrome (FXS), cell autonomously promotes maturation of callosal excitatory synapses between somatosensory barrel cortices in mice. Postnatal, cell-autonomous deletion of *Fmr1* in postsynaptic Layer (L) 2/3 or L5 neurons results in a selective weakening of AMPA receptor- (R), but not NMDA receptor-, mediated callosal synaptic function, indicative of immature synapses. Sensory deprivation by contralateral whisker trimming normalizes callosal input strength, suggesting that experience-driven activity of postsynaptic *Fmr1* KO L2/3 neurons weakens callosal synapses. In contrast to callosal inputs, synapses originating from local L4 and L2/3 circuits are normal, revealing an input-specific role for postsynaptic *Fmr1* in regulation of synaptic connectivity within local and callosal neocortical circuits. These results suggest direct cell autonomous and postnatal roles for FMRP in development of specific cortical circuits and suggest a synaptic basis for long-range functional underconnectivity observed in FXS patients.

Introduction

Disrupted structural and functional brain connectivity has been widely observed in patients with autism spectrum disorder (ASD) (Dimond et al., 2019; Holiga et al., 2019; Rane et al., 2015). A common finding in ASD is reduced corpus callosum integrity and interhemispheric functional connectivity, the latter of which correlates with autistic symptoms (Li et al., 2019; O'Reilly et al., 2017; Yao et al., 2021). The corpus callosum connects bilateral hemispheres and functions to synchronize cortical circuits necessary for sensory-motor processing, attention, perception and higher cognitive functions (Schulte and Müller-Oehring, 2010). Little is known of how autism-risk genes regulate development of callosal connectivity and the cellular or synaptic basis of reduced functional connectivity in ASD.

To provide insight into these questions, we have studied the role of the Fragile X Mental Retardation gene (*Fmr1*) in development of callosal synaptic connections in mice. Loss of function mutations in *FMR1* in humans cause Fragile X Syndrome (FXS), the most common inherited form of intellectual disability and leading monogenic cause of ASD (Garber et al., 2008; Niu et al., 2017). Children, age 6-24 months, with FXS have reduced structural integrity of white matter tracts, including the corpus callosum (Swanson et al., 2018). *Fmr1* knockout (KO) mice, an animal model for FXS, have a similar reduction in white matter tract integrity as well as decreased functional coherence among different cortical regions as measured with functional MRI (Haberl et al., 2015; Zerbi et al., 2018). Specifically, neural networks involved in sensory processing show severe functional underconnectivity, including both intra- and inter-hemispheric cortical circuits (Zerbi et al., 2018). The cellular or synaptic basis for decreased inter-region functional coherence in FXS is unknown, and whether this is due to direct or indirect roles for *Fmr1* in cortical neurons is unclear.

In addition to reduced long-range connectivity, there are reports of increased connectivity and hyperexcitability of local cortical circuits in humans with ASD and FXS (Ciarrusta et al., 2020; Courchesne and Pierce, 2005). In the *Fmr1* KO mouse, there is strong evidence for hyperexcitable local cortical circuits in *Fmr1* KO mice, including in visual, auditory and somatosensory cortices (Contractor et al., 2015; Gibson et al., 2008; Goncalves et al., 2013; Hays et al., 2011; Osterweil et al., 2013). Hyperactive cortical circuits are also observed in humans with FXS, and *Fmr1* KO mice as an increase in resting state gamma power in the resting state EEG (Jonak et al., 2020; Lovelace et al., 2018; Wang et al., 2017). Multiple cellular and synaptic alterations likely contribute to hyperexcitability of local circuits including synaptically hyperconnected pyramidal neurons, reduced inhibitory neuron activity and changes in intrinsic excitability (Gibson et al., 2008; Goel et al., 2018; He et al., 2014; Zhang et al., 2014). Layer (L) 2/3 and L5 cortical pyramidal neurons receive and integrate excitatory synaptic inputs from homotopic contralateral hemisphere (callosal) as well as other long-range inputs

from ipsilateral cortical regions and local cortical circuits. Little is known if or how the development of local and long-range synaptic inputs is balanced or if this balance is regulated by ASD-risk genes.

Fmr1 encodes Fragile X Mental Retardation Protein (FMRP), an RNA binding protein that interacts with many mRNAs including those encoding pre- and post-synaptic proteins (Darnell et al., 2011). It is perhaps through this diversity of mRNA targets that FMRP regulates multiple properties of synapses, including maturation, pruning and acute forms of synaptic plasticity (Huang et al., 2013; Pfeiffer and Huber, 2009). Regarding excitatory cortical synapses in somatosensory cortex, *Fmr1* regulates maturation of thalamocortical inputs to L4 neurons as well as between local cortical circuits. A common finding is a delayed maturation of excitatory synapses on *Fmr1* KO neurons, as observed by the delayed presence of NMDA receptor-only, or "silent", synapses and acquisition of AMPAR-mediated synaptic transmission at thalamocortical synapses onto L4 neurons and between locally connected L5 neurons (Contractor et al., 2015; Patel et al., 2014). Furthermore, L4 to L2/3 synaptic inputs are weak and delayed in their developmental strengthening, and dendritic spines on cortical pyramidal neurons are thin and filopodial-like, resembling immature spines (Cruz-Martin et al., 2010; Huang et al., 2013; Nimchinsky et al., 2001). L5 local synaptic connections ultimately mature and appear normal during the 4[th] postnatal week but then fail to prune, which results in hyperconnectivity of L5 neurons at 4-5 weeks of age as compared to wild-type cortex (Patel et al., 2014). Importantly, the delayed development and hyperconnectivity of L5 circuits are due to a cell autonomous, postsynaptic role for FMRP in L5 neurons, suggesting a direct role of FMRP in coordinating multiple synapse development processes. The cellular locus of FMRP function in development of L4 to L2/3 inputs or whether postsynaptic FMRP similarly regulates development of long-range synaptic connections is unknown.

Using viral mediated expression of Channelrhodopsin 2 (ChR2) in callosal projecting cortical neurons (Petreanu et al., 2009), we observe weak callosal synaptic inputs onto L2/3 and L5 *Fmr1* KO neurons that are mediated by a cell autonomous, postsynaptic, and postnatal role of FMRP. Callosal inputs have a selective deficit in AMPA receptor (R), but not NMDAR-, mediated synaptic transmission, indicative of synapse maturation deficit in *Fmr1* KO neurons. Sensory deprivation by whisker trimming normalized callosal input strength suggesting that sensory experience-driven activity of postsynaptic *Fmr1* KO neurons weakens callosal synapses. Surprisingly, local excitatory inputs were normal on L2/3 neurons with postsynaptic *Fmr1* deletion, revealing differential regulation of local and callosal synaptic connections by FMRP. These results reveal a cellular and synaptic substrate for reduced interhemispheric connectivity in FXS as well as imbalanced activity with local circuits.

Results

Optogenetic activation of callosal axons shows weak excitatory synaptic inputs onto L2/3 pyramidal neurons in barrel cortex of *Fmr1* KO mice

To measure callosal synaptic function and connectivity between hemispheres, mCherry-tagged Channelrhodpsin-2 (ChR2) was expressed in callosal projecting cortical neurons by injecting AAV9 into in one hemisphere of primary somatosensory cortex of postnatal day (P) 1 pups. At P18-25, acute coronal slices containing barrel cortex contralateral to the AAV injected side were prepared and whole cell patch clamp recordings of L2/3 pyramidal neurons were performed in the region innervated by fluorescently labelled callosal axons (Fig. 1A-B). Monosynaptic excitatory postsynaptic currents (EPSCs) were elicited by stimulating ChR2 expressing callosal axons with brief (2ms) blue LED light pulses in the presence of tetrodotoxin (TTX) and 4-aminopyridine (4-AP) as described previously (Petreanu et al., 2007; Rajkovich et al., 2017) (Fig. 1C). The blue LED was delivered through a 40X lens centered around the soma of approximately 350 µm diameter in size and thus the amplitude of LED-evoked EPSCs likely reflects the overall strength of callosal synaptic inputs onto the recorded neuron. The amplitude of LED-evoked EPSCs in *Fmr1* KO neurons was reduced by ~40% compared to WT. Similarly, reduced EPSC amplitudes were observed by comparing raw values or when normalized to LED power (Fig. 1D). To determine if weak callosal synaptic inputs were localized to a subcellular region on recorded L2/3 neurons (e.g. apical vs. distal dendrites), we utilized a method termed subcellular Channelrhodopsin2-assisted circuit mapping (sCRACM (Petreanu et al., 2009)). A blue laser was flashed across an array of locations, relative to the soma of the recorded neuron, in a pseudorandom order to elicit glutamate release from ChR2-expressing callosal axon terminals and EPSCs at localized dendritic sites (Fig. 1E-F). The mean amplitude of EPSCs at each site was converted into a color map for each recorded neuron and these maps were then aligned by the position of soma and averaged within genotype. The resulting average map depicts the subcellular distribution of callosal synaptic input strengths onto WT or *Fmr1* KO neurons (Fig. 1G). Results reveal that callosal synaptic inputs onto WT L2/3 neurons are strongest at the proximal apical dendrites (Fig. 1H), consistent with previous reports (Petreanu et al., 2009) and there is an interaction of *Fmr1* and vertical position (*p<0.05, $F_{(15, 849)}=1.723$; mixed-effects ANOVA). Comparison of mean EPSC amplitudes from proximal apical dendrites reveals a 50% reduction in callosal synaptic input strength in *Fmr1* KO mice (Fig. 1H).

Postsynaptic *Fmr1* cell-autonomously promotes callosal synapses

FMRP is known to have both pre- and postsynaptic, cell autonomous roles in excitatory synapse

development, depending on the cell type and synaptic input (Patel et al., 2013; Patel et al., 2014;

Pfeiffer et al., 2010). To determine the cell-autonomous and synaptic locus of FMRP function in

development of callosal synapses, we deleted *Fmr1* in a sparse population (~3-5%) (Fig. 2 – figure

supplement 1) of postsynaptic L2/3 pyramidal neurons by injecting AAV expressing GFP-tagged Cre

(AAV9.GFP-Cre) into the left lateral ventricle of P1 pups (Kim et al., 2013) with a floxed *Fmr1* gene

(*Fmr1^{fl/fl}* or *Fmr1^{fl/y}*) (Fig. 2A) (Mientjes et al., 2006). In the same animals, AAV.ChR2-mCherry was

injected into the right barrel cortex to label callosal axons as in Fig. 1. This experimental design allowed

us to measure the cell autonomous effect of postsynaptic FMRP deletion in L2/3 neurons in a primarily

WT cortex on synapses from WT callosal axons. At P18-30, acute cortical slices were prepared and

simultaneous whole cell recordings were performed on pairs of neighboring L2/3 pyramidal neurons in

left barrel cortex with one being a GFP (-), or WT neuron, and the other a GFP (+), or *Fmr1* KO neuron

(Fig. 2A-B). Monosynaptic EPSCs from callosal axons were elicited with either bulk LED stimulation or

blue laser, to perform sCRACM, as in Fig. 1 (Fig. 2C, Fig. 2 - figure supplement 2 A-B). Interestingly, at

the earliest age for recording (P18-20), overall callosal synaptic input strengths were not different

between WT and postsynaptic *Fmr1* KO neurons (Fig. 2D). However, at P23-30, postsynaptic *Fmr1* KO

neurons had a 40% reduction in callosal synaptic input strength compared to neighboring WT neurons

(Fig. 2D). A 2 way ANOVA indicates a significant interaction between genotype and age (*Fmr1* x Age

***$p<0.001$, $F_{(1, 32)}=13.87$)(Fig. 2E). In addition, we observed a strong trend of callosal input strength

to increase with developmental age in WT, but not *Fmr1* KO, neurons (Fig. 2E). Similar results are

obtained with sCRACM which reveal a ~45% reduction in callosal input strength onto proximal

dendrites of *Fmr1* KO neurons (Fig. 2 – figure supplement 2 C-E). These data demonstrate that *Fmr1* in

postsynaptic L2/3 neurons cell-autonomously promotes the development and/or strengthening of

callosal synaptic inputs. Weak callosal synaptic strength in *Fmr1* KO neurons at P23-30 could be a

consequence of deficient or delayed callosal synapse maturation and may normalize in the adult. To

test this possibility, we repeated experiments deleting postsynaptic *Fmr1* with a P1 AAV Cre-GFP

injection and recorded LED-evoked EPSCs onto pairs of neighboring WT and *Fmr1* KO L2/3 neurons in

adult mice (P57-65). Similar to young mice, callosal mediated EPSCs were weak in *Fmr1* KO neurons;

reduced by 25%, in comparison to WT neurons (Fig. 2F-G). These results indicate that weak callosal

synaptic transmission persists into adulthood with postsynaptic *Fmr1* deletion.

Selective weakening of AMPAR-, but not NMDAR-, mediated synaptic transmission from callosal

inputs onto *Fmr1* KO L2/3 pyramidal neurons

Weak callosal mediated EPSCs onto postsynaptic *Fmr1* KO L2/3 pyramidal neurons could be due to
reductions in the number of synaptic connections, presynaptic release probability, the strength of
individual synapses, or any combination of these. To further investigate the synaptic basis of weak
callosal inputs in P23-30 slices, we recorded in strontium (Sr^{2+}), substituted for Ca^{2+}, in the external
ASCF which results in asynchronous vesicle release and measurement of quantal synaptic events
evoked from callosal axons by the LED (Oliet et al., 1996). Changes in the amplitude of quantal events
reflect synapse strength, whereas changes in the frequency of events reflect changes in synapse
number and/or presynaptic release probability. Postsynaptic *Fmr1* KO neurons received >20% fewer
evoked events in Sr^{2+} in comparison to neighboring WT neurons with no change in event amplitude (Fig.
3A-C). The baseline frequency and amplitude of spontaneous EPSCs, prior to LED stimulation, were
not different between genotypes. These results suggest the reduced EPSCs onto postsynaptic *Fmr1*
KO neurons are a consequence of reduced presynaptic release probability, functional synapse number
or both. To further test this conclusion, we measured the coefficient of variance (C.V.) of LED-evoked
callosal EPSCs onto WT and *Fmr1* KO neurons (from experiments in Figs, 2D (P23-30) and Fig. 5B$_1$).
C.V. is inversely proportional to release probability and synapse number (Manabe et al., 1993) and was
increased by 18% in *Fmr1* KO neurons (Fig. 3 – figure supplement 1). This result together with
reduced frequency of events evoked in Sr^{2+} indicate that the weakening of callosal mediated EPSCs in
Fmr1 KO is due in part to decreased synapse number and/or presynaptic release probability.

To further differentiate between release probability and functional synapse number, we measured
NMDA receptor (R) mediated EPSCs from callosal axons onto WT or postsynaptic *Fmr1* KO L2/3
neurons. Because NMDARs are colocalized with AMPARs at excitatory synapses, reduced glutamate
release probability or callosal synapse number onto *Fmr1* KO neurons would be expected to result in
weak NMDAR-EPSCs. To evoke isolated NMDAR EPSCs from callosal axons, we included the
AMPAR antagonist DNQX, glycine, and low Mg^{2+} (0.1 mM) in the ACSF and voltage clamped neurons
at -70 mV. In contrast to AMPAR-mediated EPSCs, amplitudes of NMDAR EPSCs evoked from callosal
axons were not different between WT and neighboring postsynaptic *Fmr1* KO neurons (Fig. 3D). This
observation suggests that the NMDAR content within callosal synapses are similar for WT and KO
neurons and that there is no change in presynaptic release probability or synapse number for the
callosal axons. Taken together with the reduced frequency of AMPAR quantal events and increased
C.V. (Fig. 3C, Fig. 3 – figure supplement 1), our results suggest that L2/3 *Fmr1* KO neurons have an
increased proportion of synapses with NMDARs, but not functional AMPARs, indicative of immature
synapses.

Postsynaptic *Fmr1* KO pyramidal neurons in both L2/3 and L5 receive weak callosal synaptic inputs with action-potential-driven synaptic transmission

For experiments described in Figs. 1 and 2, we included TTX and 4-AP in the bath to isolate evoked EPSCs from ChR2-expressing callosal axons and block potential contamination from polysynaptic local L2/3 inputs. A caveat of this approach is that callosal synaptic transmission is not triggered by action potentials, but by direct depolarization and activation of voltage-gated Ca^{2+} channels at the callosal axon terminal. To determine if similar results are observed with action potential-evoked synaptic transmission, we repeated experiments without TTX/4AP, but increased Ca^{2+}/Mg^{2+} and added the NMDAR blocker, CPP, to reduce polysynaptic responses from local circuits. We also used a blue laser to depolarize a local (30µm) area of callosal axons while limiting activation of local circuits (Fig. 4A). Quantification of mean EPSC amplitude within the region covering soma and major dendrites shows that L2/3 pyramidal neurons with postsynaptic *Fmr1* deletion have a 45% reduction in callosal synaptic input strength as compared to neighboring WT neurons (Fig. 4A-B).

L5 pyramidal neurons are another major target of callosal axons (Petreanu et al., 2007; Wang et al., 2007). To determine if weak callosal synaptic inputs to postsynaptic *Fmr1* KO neurons are specific to L2/3, we performed the same experiment on L5 pyramidal neurons. Similar to L2/3, L5 pyramidal neurons with postsynaptic *Fmr1* deletion have a 40% reduction in the strength of callosal inputs stimulated around the soma and apical dendrites (Fig. 4C-D). Together, these results show that postsynaptic deletion of *Fmr1* in pyramidal cortical neurons generally weakens callosal synaptic inputs and this is observed with action potential driven synaptic transmission and across different layers.

Sensory deprivation by whisker trimming normalizes callosal synaptic strengths in *Fmr1* KO L2/3 neurons

Targeting, branching and elaboration of callosal axons into the contralateral neocortex occurs postnatally (~P5-15) and depends on activity of presynaptic and postsynaptic cortical neurons as well as whisker sensory experience (Huang et al., 2013; Mizuno et al., 2010; Wang et al., 2007). This suggests that sensory experience may interact with FMRP to regulate development of callosal synapses. To test this idea, we sparsely deleted *Fmr1* in postsynaptic L2/3 neurons in the left barrel cortex and express ChR2-mCherry in callosal projecting neurons in the right hemisphere (as in Fig. 5A). Beginning at P15, we unilaterally trimmed whiskers daily on the right whisker pad which would reduce the most direct ascending sensory-driven patterned activity to the left hemisphere containing L2/3 neurons with postsynaptic *Fmr1* deletion (postsynaptic deprivation condition). In littermates, we

trimmed whiskers on the left whisker pad which would primarily deprive the L2/3 callosal projection
neurons of sensory driven patterned activity (presynaptic deprivation condition) (Fig 5A). Whisker
trimming began at P15 to reduce effects of sensory deprivation on the early growth and branching of
callosal projection axons and continued until the day before slice recordings. Recordings of EPSCs
evoked from callosal axons by either LED stimulation (Fig 5B) or sCRACM (Fig 5 - figure supplement 1)
were obtained from pairs of neighboring WT and *Fmr1* KO L2/3 neurons. Because we recorded from
pairs of WT and postsynaptic *Fmr1* KO neuron neighbors, any effects of sensory deprivation on callosal
axon innervation, growth, or branching in a given cortical region would be expected to similarly affect
each genotype. Thus, our results reflect the cell autonomous effects of *Fmr1* on synaptic function or
connectivity. In the "presynaptic deprivation" condition, callosal synaptic input strength was weak (27%
reduction) onto postsynaptic *Fmr1* KO L2/3 neurons as compared to WT, similar to that observed in
whisker intact mice (Fig 2D). In contrast, in the "postsynaptic deprivation" condition, callosal synaptic
inputs strengths were similar between WT and *Fmr1* KO neurons. To compare the callosal synaptic
inputs strength within genotypes and across sensory deprivation paradigms, we normalized each LED
induced EPSC to its stimulation power (Fig 5C). A 2-way ANOVA revealed an interaction of deprivation
condition and *Fmr1* (*$p<0.05$, $F(1, 31)=4.977$, ANOVA). Surprisingly, callosal synaptic input strengths in
WT neurons were not different between deprivation conditions. In contrast, in *Fmr1* KO neurons
callosal input strengths were weaker, decreased by ~45%, in the presynaptic deprivation condition, as
compared to postsynaptic deprivation. These data suggest that whisker experience-driven activity of
postsynaptic *Fmr1* KO L2/3 neurons weakens callosal synaptic inputs.

Input specific strengthening of callosal synaptic connections by postsynaptic FMRP

A previous study reported weak L4 to L2/3 synaptic inputs in the barrel cortex of global *Fmr1* KO mice
(Bureau et al., 2008), suggesting that *Fmr1* generally promotes excitatory synapse strength onto L2/3
neurons regardless of whether they are from local or long-range sources. To determine if postsynaptic
Fmr1 promotes excitatory synapse development from local cortical circuits in a cell autonomous
manner as it does for callosal inputs, we assessed local input strengths using Laser Scanning Photo-
Stimulation (LSPS) with glutamate uncaging. Slices were bathed in MNI-caged glutamate and
pseudorandom flashes of a UV laser (355 nm) beam at individual locations (20 µm diameter) within a
16-by-16 grid surrounding the recorded L2/3 neurons focally released glutamate to evoke action
potentials and synaptic transmission from neurons at that location. The grid for UV laser flashing was
positioned to stimulate neurons in Layers 2/3-5 of home and adjacent barrels. At P23-30, LSPS was
performed on pairs of simultaneously recorded WT and neighboring L2/3 neurons with postsynaptic

Fmr1 deletion (Fig. 6A-B). The amplitude of monosynaptic EPSCs evoked from each position in the
slice for an individual neuron were converted into a color map and then individual maps were aligned to
the home barrel to create an average color map per genotype (Fig. 6C-D). Responses from direct
glutamate activation onto recorded neurons, as described in methods, were excluded from analysis,
and represented by black pixels. EPSCs evoked from L4 home or adjacent barrels, L5A or L2/3 were
unaffected on *Fmr1* KO L2/3 neurons (Fig. 6E), in contrast to the weak EPSCs from callosal inputs.

Bureau and colleagues demonstrated weak L4 to L2/3 synaptic strength in the global *Fmr1* KO when
measured at 2 weeks of age, but the difference was diminished at 3 weeks, suggestive of a
developmental delay. To test if loss of postsynaptic *Fmr1* weakens local L4 to L2/3 synaptic inputs at
early developmental stages, we performed LSPS on pairs of WT and postsynaptic *Fmr1* KO L2/3
pyramidal neurons at P14-17. Similar to results obtained at P23-30, we observed normal L4 and
adjacent L2/3 synaptic input strengths onto postsynaptic *Fmr1* L2/3 neurons recorded at P14-17 (Fig. 6
– figure supplement 1A). We also tested if embryonic deletion of postsynaptic *Fmr1* was necessary to
affect L4 to L2/3 synapse development, using slices from females with heterozygous (het) and mosaic
expression of *Fmr1*, as we have described (Patel et al., 2013; Patel et al., 2014). Briefly, X-linked GFP
mice were bred with *Fmr1* KO males. Due to random X-chromosome inactivation in the embryo, female
Fmr1 het mice offspring have a mosaic expression of GFP(+) WT and GFP(-) *Fmr1* KO neurons. LSPS
maps were performed on pairs of WT and *Fmr1* KO neurons at P14-17 where we observed normal L4
input strengths onto L2/3 *Fmr1* KO neurons (Fig. 6 – figure supplement 1B). In contrast to results with
postsynaptic *Fmr1* deletion, we observed weak L4-to-L2/3 synaptic strengths with LSPS in the global
Fmr1 KO (Fig. 6 – figure supplement 1C) similar to that reported by Bureau et al. (Bureau et al., 2008).
However, unlike Bureau et al, we observed weak L4 inputs at later ages (P18-25 ; Fig. 6 - figure
supplement 1C). Taken together our results implicate FMRP in postsynaptic L2/3 neurons in the input
specific development of callosal synapses. Although synaptic inputs from L4 to L2/3 are weak in the
global *Fmr1* KO, this is not a postsynaptic, cell autonomous function of FMRP, but instead is either a
non-cell autonomous function of FMRP or a role in presynaptic L4 neurons.

Discussion
Reduced interhemispheric connectivity, observed both structurally and functionally, is a hallmark of
ASD in humans and correlated with symptoms (Dimond et al., 2019; Holiga et al., 2019; Li et al., 2019;
O'Reilly et al., 2017; Rane et al., 2015; Yao et al., 2021). However, little is known of the cellular,
synaptic, and molecular mechanisms by which this occurs in ASD and any direct role of ASD-risk
genes. Here, we demonstrate a direct, postsynaptic and postnatal role for *Fmr1* in maturation and/or
stability of callosal synaptic inputs in L2/3 and L5 cortical neurons and this change is also observed in

the FXS- mouse model, the global *Fmr1* KO. Surprisingly, postsynaptic deletion of *Fmr1* did not
weaken synaptic inputs from local columnar circuits onto L2/3, revealing that postsynaptic FMRP
differentially regulates development of select synaptic inputs (Fig. 6 – figure supplement 2). Sensory
deprivation of postsynaptic *Fmr1* KO neurons prevented weakening of callosal synaptic inputs
suggesting that experience-driven patterned activity of postsynaptic L2/3 neurons without FMRP is
necessary for synaptic weakening and/or prevents maturation. In conclusion, our results reveal a
postsynaptic mechanism by which *Fmr1* regulates callosal connectivity that likely contributes to the
reduced interhemispheric structural and functional connectivity in *Fmr1* KO mice and humans with FXS
(Haberl et al., 2015; Swanson et al., 2018; Zerbi et al., 2018).

A synaptic basis for reduced functional long-range connectivity in FXS.

Functional MRI studies show that *Fmr1* KO mice have reduced corticocortical long-range connectivity,
especially among the sensory and motor cortices (Haberl et al., 2015; Zerbi et al., 2018) as well as
reduced corpus callosum structural integrity. By investigating callosal synaptic connections between
bilateral barrel cortices, we find weak functional synaptic inputs at these long-range connections in L2/3
in *Fmr1* KO mice (Fig. 1). This result suggests that reduced functional coherence between bilateral
barrel cortices in *Fmr1* KO mice could be due to a disrupted communication between the cortices
through the weakened monosynaptic transmission. Interhemispheric functional connectivity is thought
to shape perceptual integration including those involved in speech comprehension and global form
processing, domains that are impaired in ASD (Booth and Happé, 2018; Friederici et al., 2007; Happe
and Frith, 2006; Peiker et al., 2015; Preisig et al., 2021; Simon and Wallace, 2016). Based on fMRI
results in the *Fmr1* KO (Haberl et al., 2015; Zerbi et al., 2018), it is likely that FMRP promotes other
"long-range" synaptic connections between cortical areas, such as ipsilateral connectivity between S1
and M1/M2, as well as cortical-subcortical structures.

Postsynaptic FMRP in L2/3 neurons promotes maturation of callosal synapses

Using *in vivo* sparse deletion and simultaneous recording paradigm, we can stimulate the same set of
callosal axons for a pair of WT and *Fmr1* KO neurons and directly compare their synaptic inputs. Our
observation of weaker callosal synaptic inputs onto cell autonomous *Fmr1* KO L2/3 neurons (Fig. 2)
confirms results in *Fmr1* global KO mice and further reveals an essential role of postsynaptic FMRP in
promoting callosal synapses development. With postsynaptic *Fmr1* KO we observe weak callosal
synapses at P23-30, but not at P18-20. To determine if a similar developmental profile is observed in
the global *Fmr1* KO, we analyzed a subgroup of data collected at P18-20 and observe weak callosal
synaptic strength in the *Fmr1* KO at this early time point (WT =181 ±22 pA, n=15; KO=106.1±17.77 pA,

n=7, *p<0.05, unpaired t-test). The later developmental onset of weak callosal synaptic transmission

with postsynaptic *Fmr1* deletion may be due to later postnatal deletion of *Fmr1* using P1 AAV-Cre-GFP

injection which we estimate to occur about P7-P9 (Rajkovich et al., 2017). However, we cannot rule out

a role for FMRP in presynaptic, callosal projecting neurons, or other cell types, such as

oligodendrocytes, within the first postnatal weeks to establish callosal synaptic connections (Doll et al.,

2020; Hanson and Madison, 2007; Patel et al., 2013).

Our results indicate that callosal synapses do not mature or are not maintained without FMRP in

postsynaptic L2/3 neurons. In support of a role in maturation, callosal synaptic strength tends to

increase from P18-20 to P23-30 in WT neurons but not in postsynaptic *Fmr1* KO neurons (Fig. 2E).

Furthermore, we observe a selective weakening of AMPAR, but not NMDAR-, mediated synaptic

transmission at callosal inputs onto *Fmr1* KO neurons (Fig. 3). This result, together with a decreased

frequency of evoked quantal events in Sr^{2+} and increased coefficient of variation, suggests a reduced

number of mature synapses with functional AMPARs. As synapses mature, they acquire NMDARs prior

to AMPARs and NMDAR-only, immature synapses are often termed "silent" synapses (Ashby and

Isaac, 2011; Hanse et al., 2013). Although we did not measure "silent" callosal synapses here, our

findings would predict more "silent" and immature callosal synapses in *Fmr1* KO L2/3 neurons. In

support of this idea, thalamocortical inputs to L4, another "long-range" synaptic pathway, are delayed in

their development in the *Fmr1* KO, as measured by acquisition of AMPARs (Harlow et al., 2010).

Another possible explanation for our results is the optogenetic stimulation paradigm we used to evoke

glutamate release from callosal axons saturates synaptic NMDARs, but not AMPARs, due to the higher

affinity of glutamate for NMDARs (Patneau and Mayer, 1990). Additional experiments and/or methods

are needed to confirm a selective decrease in AMPARs at *Fmr1* KO callosal synapses. The

observations of reduced corpus callosum structural integrity and interhemispheric coherence with fMRI

in adult *Fmr1* KO suggest weak callosal connectivity in adults (Haberl et al., 2015; Zerbi et al., 2018).

Weak callosal synapses persist in adults with postsynaptic *Fmr1* deletion, suggesting a deficit in

maturation or AMPAR insertion/stability, as opposed to developmental delay (Fig. 2F-G). Reduced

callosal axon diameter is observed in adult *Fmr1* KO which may be a consequence of weak or

immature callosal synapses and contribute to reduced interhemispheric coherence in FXS (Haberl et al.,

2015).

Postsynaptic FMRP differentially regulates synaptic inputs from local and long-range cortical

circuits

Neocortical pyramidal neurons integrate excitatory synaptic inputs from local and long-range circuits,
including ipsilateral and contralateral cortical areas (Feldmeyer, 2012; Gerfen et al., 2018). An
imbalance in local and long-range connectivity has been hypothesized to contribute to ASD in humans;
specifically, hyperconnectivity of local circuits and underconnectivity of long-range circuits or between
brain regions (Belmonte et al., 2004; Courchesne and Pierce, 2005; O'Reilly et al., 2017; Rane et al.,
2015). If or how ASD genes regulate the balance of local and long-range synaptic connectivity is
unknown. Here we demonstrate that postsynaptic FMRP differentially regulates development and/or
maintenance of synaptic inputs from local and long-range cortical sources. L2/3 neurons with
postsynaptic deletion of *Fmr1* have weak and immature callosal synaptic inputs but normal synaptic
inputs from L4, adjacent L2/3 and L5 (Fig. 6). Bureau et al. (Bureau et al., 2008) and we (Fig. 6 – figure
supplement 1C) find that L4 to L2/3 synapses are weak in the global *Fmr1* KO, but we do not observe
this with embryonic or postnatal cell autonomous deletion of *Fmr1* in L2/3 neurons. Together these
results suggest a role for FMRP in presynaptic L4 neurons in synapse development onto L2/3 neurons,
which is consistent with the reported deficits in L4 axon morphology in the global *Fmr1* KO (Bureau et
al., 2008).

Results in L5 also indicate differential regulation of local and long-range cortical connectivity by
postsynaptic FMRP. Our previous work using multiple simultaneous recordings of locally connected
L5A neurons, revealed hyperconnectivity of *Fmr1* KO L5 neurons with their immediate neighbors (<40
μm apart) in S1 at 4 weeks of age (Patel et al., 2014). Hyperconnectivity of L5 local subnetworks
resulted from deficient developmental pruning between *Fmr1* KO L5 neurons and was observed in both
the global *Fmr1* KO and with postsynaptic *Fmr1* deletion. Prefrontal L5 cortical neurons in *Fmr1* KO
mice are similarly hyperconnected (Testa-Silva et al., 2012). In contrast to local hyperconnectivity, here
we find that postsynaptic deletion of *Fmr1* in L5 neurons results in weak callosal synaptic inputs. Thus,
Fmr1 KO L5 pyramidal neurons are hyperconnected locally and under-connected to contralateral cortex;
an effect that is mediated by cell-autonomous and postsynaptic deletion of FMRP. Such an effect may
promote the reported imbalances in local and long-range functional connectivity observed in ASD
individuals. Postsynaptic deletion of FMRP in L2/3 neurons did not affect synaptic inputs from other
layers or between columns within L2/3 (Fig. 6) suggesting that FMRP may selectively promote pruning
of local connections in L5 neurons. Alternatively, FMRP may prune synaptic connections within very
local cortical subnetworks (within 40 μm) in both L2/3 and L5, but not connections between layers and
columns.

The molecular mechanisms by which postsynaptic FMRP differentially regulates L4 and callosal
inputs to L2/3 neurons is unclear. L4 neurons synapse primarily on basal dendrites of L2/3 neurons,

whereas callosal inputs are primarily on apical dendrites (Bosman et al., 2011)(Fig. 1G). Therefore,
localized expression and translational regulation of specific dendritic mRNAs by FMRP at either basal
or apical dendritic compartments could differentially affect L4 and callosal inputs. Alternatively, FMRP
translational regulation of postsynaptic cell adhesion molecules, such as neuroligins, could differentially
impact specific presynaptic inputs based on their expression of binding partners such as neurexin
splice variants (Südhof, 2017).

Bidirectional regulation of callosal synaptic function by MEF2C and FMRP

Differential regulation of local and long-range cortical synaptic connectivity is observed with another
ASD-risk gene, Myocyte Enhancer Factor 2C (*Mef2c*). Loss of function mutations in *MEF2C*, which
encodes an activity-dependent transcription factor, are implicated in intellectual disability, ASD and
schizophrenia (Assali et al., 2019; Mitchell et al., 2018; Rocha et al., 2016). In contrast to FMRP,
postsynaptic deletion of *Mef2c* in L2/3 neurons results in fewer and weak inputs from local circuits (L4,
L2/3 and L5), but strengthened callosal inputs (Rajkovich et al., 2017). The differential regulation of
local and callosal synaptic connections by postsynaptic MEF2C and FMRP, albeit in different directions,
suggest that imbalances in local and long-range synaptic connectivity may contribute to different
genetic causes of neurodevelopmental disorders. Our results also implicate roles for transcription and
translational control in the input-specific development of cortical circuits. Bidirectional regulation of local
and long-range connectivity by *Mef2c* in L2/3 or by *Fmr1* in L5 could be an effect of homeostasis or
competition between local and long-range synaptic connections to maintain optimal cortical circuit
function. For example, weakening of callosal synaptic input in *Fmr1* KO neurons may be compensatory
and an attempt to normalize hyperconnected or hyperexcitable local circuits.

Interaction of sensory experience and ASD-risk genes in regulation of long-range cortical
circuits

The regulation of local or callosal connectivity by FMRP and MEF2C requires normal sensory
experience suggesting that both of these genes function in experience and activity-regulated pathways
necessary for cortical circuit development. Sensory deprivation by whisker trimming normalizes callosal
synaptic inputs in L2/3 neurons with postsynaptic deletion of *Fmr1* (Fig. 5) or *Mef2c* (Rajkovich et al.,
2017). Specifically, trimming whiskers contralateral to L2/3 neurons with postsynaptic *Fmr1* deletion
(postsynaptic deprivation) prevented callosal synaptic weakening. In contrast, "presynaptic deprivation"
or trimming whiskers contralateral to ChR2-expressing, callosal projecting neurons, had no effect (Fig.
5). This result suggests that sensory-driven patterned activity of postsynaptic *Fmr1* KO L2/3 neurons
weakens callosal synapses. An intriguing possibility is that an activity-dependent long-term synaptic
depression (LTD) process is enhanced at *Fmr1* KO L2/3 neurons as observed in hippocampal CA1

(Huber et al., 2002). Sensory deprivation also induces homeostatic synaptic scaling in primary sensory
cortices, as measured with spontaneous or miniature (m) EPSCs (Feldman, 2009; Hooks and Chen,
2020). Neither the frequency nor amplitude of mEPSCs was different between neighboring WT and
Fmr1 KO L2/3 neurons in mice with normal sensory experience or "presynaptic deprivation". However,
with "postsynaptic deprivation", mEPSC amplitude and frequency was increased in *Fmr1* KO neurons
relative to WT (Fig. 5 – figure supplement 2) suggestive of enhanced homeostatic up scaling in *Fmr1*
KO neurons that may contribute to callosal input strengthening in this condition.

Bilateral underconnectivity is common in ASD as well as disconnection of other long-range cortical
connections with other brain regions such as hippocampus and cerebellum. Bilateral connectivity and
synchrony between cortical regions are necessary for speech comprehension, sensory processing and
cognition (Bland et al., 2020; Castro et al., 2014; Friederici et al., 2007; Fries, 2009; 2015; Panzica et
al., 2019), domains impaired in FXS and ASD. Our present findings contribute to the understanding of
the cellular and synaptic mechanisms by which ASD-risk genes, such as *FMR1,* regulate long-range
connectivity, how this is coregulated and balanced with local circuit connectivity and interacts with
experience-dependent brain development.   Such information may contribute to therapies to aid
abnormal brain connectivity in ASD.

 Materials and Methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
genetic reagent (*M. musculus*)	*Fmr1$^{-/y}$* (male)	Jackson Laboratory	003025	
genetic reagent (*M. musculus*)	*Fmr1$^{fl/fl}$* (female) *Fmr1$^{fl/y}$* (male)	PMID:16257225		Dr. David Nelson (Baylor College of Medicine)
genetic reagent (*M. musculus*)	X-linked GFP	Jackson Laboratory	003116	
strain, strain background (*AAV*)	AAV9.CMV.HI. eGFP-Cre. WPRE.SV40	Addgene	105545	
strain, strain background (*AAV*)	AAV9.CAG.hChR2(H134R)-mCherry.WPRE.SV40	Addgene	100054	
chemical compound, drug	MNI-caged-L-glutamate	Tocris / HelloBio	1490 / HB0423	
chemical compound, drug	(RS)-CPP	Tocris / HelloBio	0173 / HB0036	
chemical compound, drug	4-Aminopyridine (4-AP)	Sigma-Aldrich	A78403	
chemical compound, drug	DNQX disodium salt	Tocris	2312	
software, algorithm	LabView	National Instruments	RRID:SCR_014325	
software, algorithm	Multiclamp 700A	Molecular Devices	RRID:SCR_021040	
software, algorithm	Prism 8	Graphpad Software	RRID:SCR_002798	

Animals

Fmr1 KO (*Fmr1$^{-/y}$*) and X-linked GFP mice were obtained from Jackson laboratories (Stock No: 003025

and 003116, respectively). *Fmr1$^{fl/fl}$* were obtained from Dr. David Nelson (Baylor College of Medicine)

(Mientjes et al., 2006). Mice were maintained on a C57BL/6J background and reared on a 12 hr light-

dark cycle with access to food and water ad libitum. Male *Fmr1* WT (*Fmr1$^{+/y}$*) with *Fmr1* KO (*Fmr1$^{-/y}$*)

littermates were used for experiments and both male and female pups were used for *Fmr1* flox (*Fmr1$^{fl/y}$*

and *Fmr1$^{fl/fl}$*). All animal experiments were conducted in accordance with the Institutional Animal Care

and Use Committee (IACUC) at University of Texas Southwestern Medical Center.

Viral transfections in neonatal mice

Commercially   made   AAV9.CMV.HI.eGFP-Cre.WPRE.SV40   (Addgene   #105545)   and

AAV9.CAG.hChR2(H134R)-mCherry.WPRE.SV40 (Addgene #100054) were diluted to a titer ~10^{12}

vg/mL using sterile saline. Traces of Fast Green FCF dye (Sigma) were added to facilitate visualization

of virus spreading. Neonatal mouse pups (P1) were first anesthetized by hypothermia, then fixed on a

customized mold and placed on a stereotaxic frame. AAV9.eGFP-Cre (420-560 nL) was delivered to

left ventricle at a depth of approximately 1.1 mm underneath skull through a beveled glass pipette using

Nanoject II injector (Drummond Scientific, Inc.). AAV9.ChR2-mCherry (400 nL) was delivered to

superficial layers (0.5 mm underneath skull) of somatosensory cortex in right hemisphere at a speed of

1.2 µL/min using syringe pump (Harvard Apparatus, Inc.).

Acute slice preparation

Mice were anesthetized by intraperitoneal (i.p.) injection of Ketamine/Xylazine mixture and decapitated

upon irresponsiveness to toe-pinch. Acute coronal slices (300 µm thickness) containing somatosensory

barrel cortex were prepared using vibrating microtome (Leica VT1200S). During sectioning, tissue

blocks were submerged in ice-cold dissection buffer containing (in mM): 110 choline chloride, 25

NaHCO$_3$, 25 dextrose, 11.6 ascorbic acid, 2.5 KCl, 1.25 NaH$_2$PO$_4$, 3.1 Na-pyruvate, 7 MgCl$_2$ and 0.5

CaCl$_2$, continuously aerated with 95%CO$_2$/5%O$_2$. For mice older than P23, transcardial perfusion of ice-

cold dissection buffer was performed prior to decapitation to increase slice quality. Slices were then

transferred to artificial cerebrospinal fluid (ACSF) solution containing (in mM): 125 NaCl, 25 NaHCO$_3$,

10 dextrose, 2.5 KCl, 1.25 NaH$_2$PO$_4$, 2 MgCl$_2$ and 2 CaCl$_2$ (aerated with 95%CO$_2$/5%O2) and

recovered at 34°C for 30 min followed by 30 min at room temperature.

Electrophysiology

After recovery, slices were transferred to a recording chamber at room temperature and perfused with ACSF aerated with 95%CO_2/5%O_2. Slices were visualized by infrared differential interference contrast (IR-DIC) optics (Olympus BX51W1). Whole cell recordings of L2/3 and L5 pyramidal neurons were obtained using borosilicate pipettes (4-7 MΩ) and a Multiclamp 700A amplifier (Molecular Devices). Internal solution contained (in mM): 130 K-gluconate, 10 HEPES, 6 KCl, 3 NaCl, 0.2 EGTA, 14 phosphocreatine-tris, 4 Mg-ATP and 0.4 Na-GTP. All recordings were conducted in voltage clamp, holding at -70 mV unless otherwise specified, and data were collected and analyzed using custom Labview programs (Labview 8.6, National Instruments Inc.). Spiking patterns upon current injection were used as criteria to identify excitatory neurons. For experiments where spiking was blocked by TTX, rise time to hyperpolarizing current (> 50 ms) and kinetics of mPSC (width at half height > 2 ms) were used as criteria (Povysheva et al., 2006). Excitatory neurons with resting membrane potential < -50 mV and a series resistance < 35 MΩ were included in analysis. Voltages were not corrected for junction potential. For simultaneous patch clamp recordings, the distance between the pair of cells (center-to-center) is 10-40 μm.

Optogenetic bulk stimulation of callosal axons

Mice with either cortical injection of AAV9.ChR2-mCherry or unilateral ventricular injection of AAV9.eGFP-Cre and contralateral cortical injection of AAV9.ChR2-mCherry were used for this experiment. Fluorescence of GFP-positive soma and mCherry-labeled axons were visualized using a fluorescent mercury lamp (Excelitas Technologies Corp.). Slices containing clearly labeled mCherry+ axons from contralateral barrel cortex were used for recording. Slices with somatic infection of ChR2-mCherry due to virus leaked from contralateral hemisphere were discarded to avoid contamination from local inputs. The infection rate of recorded Cre-GFP+ neurons in barrel cortex was 3-5% (Figure 2 – figure supplement 1). Only neurons residing in an area with densely labeled callosal axons were subject to recording to achieve a reliable magnitude and reduced variability of light-induced responses. For all LED experiments, responses were evoked by a 2 ms flash from a digitally controlled blue LED (final beam diameter: 350 μm; power: 0.1-4.6 mW; wavelength: 470 nm; M470L4-C1, Thorlabs Inc.) through a 40X water-immersed objective. The LED flash was centered on soma and proximal apical dendrites of recorded neurons.

To measure LED-evoked EPSCs, each cell (or cell pair) was stimulated 3-10 times with 20-30 s intervals. LED power was adjusted to obtain an EPSC amplitude of 100-1000pA in WT neurons (for cell pairs). The external solution for both LED and sCRACM experiments contained ACSF with 1 μM TTX, 100 μM 4-aminopyridine (4-AP), 10 μM (±)-3-(2-carboxypiperazin-4-yl)propyl-1-phosphonic acid (CPP) and 100 μM picrotoxin to isolate monosynaptic AMPAR-mediated excitatory inputs. Callosal input

strength was measured as the peak amplitude of an average EPSC of 3 - 10 evoked EPSCs for each
neuron.

To measure Sr^{2+} evoked quantal events, each cell pair was stimulated with an LED flash every 30s, 12-
30 times in ACSF containing 4 mM $MgCl_2$, 4 mM $SrCl_2$, 10 µM CPP and 100 µM picrotoxin. Events
occurring within a 1 s window prior to the LED were defined as spontaneous events and those
occurring 50-350 ms post-LED were defined LED-evoked events. The frequency and amplitude of
events were analyzed using MiniAnalysis (Synaptosoft).

NMDAR-mediated EPSCs from callosal inputs were pharmacologically isolated in ACSF containing: 3
mM $CaCl_2$, 0.1 mM $MgCl_2$, 20 µM DNQX, 20 µM glycine, 100 µM picrotoxin, 1 µM TTX, 100 µM 4-AP.
An LED flash was delivered every 30s, 9 times. Evoked EPSCs were averaged and peak amplitude
was measured.

Subcellular Channelrhodopsin-assisted Circuit Mapping (sCRACM) and action potential dependent activation of ChR2

After collection of LED-evoked responses, the objective was switched to 4X to visualize a broader area.
ChR2 expressing axons were then stimulated by a blue laser (1 ms; wavelength: 473 nm; power range:
0.7 - 16 mW; final beam diameter: 25 µm; CrystaLaser) scanning through a 12 x 12 grid (50 µm
spacing) in a pseudorandom order to avoid repeated activation of neighboring locations. The grid was
aligned along the pia and centered in the medial-lateral position on the recorded somas. The grid was
repeatedly scanned 2 - 4 times at 40 s intervals. Laser-evoked EPSCs were collected from each grid
spot. Most neuron pairs (>90%) were homogeneously distributed between 150 and 300 µm from pia
surface.

For action potential-dependent callosal synaptic strength measurements, an independent cohort of
mice was used. The recording ACSF was similar to sCRACM experiments, except TTX and 4-AP were
omitted, and divalent cations were increased (4 mM $MgCl_2$; 3 mM $CaCl_2$) and CPP was added to
reduce polysynaptic activation of local circuits. For L2/3 neurons, an 8 x 8 grid with 75 × 100µm x-y
spacing was used. For L5 neurons, an 8 x 8 grid with 75 × 125µm x-y spacing was used.

Analysis of blue laser evoked EPSCs

For sCRACM, EPSCs evoked from 2-4 laser-stimulation at each grid spot from an individual neuron
were averaged. The peak amplitude of the average EPSC (between 5-80 ms after laser onset) was
determined as the input strength for that spot. A spatial map of input strengths was then generated for
each individual neuron. The spatial maps for all neurons of each genotype were then aligned to soma

location, oriented with respect to the pial surface, and averaged to generate an average spatial map of
input strengths for each genotype aligned to the soma. A color representation of the average spatial
map for each genotype were generated for the figures. Pixel size for the genotype-averaged color maps
was halved through pixel interpolation (25x25 µm) to provide better spatial resolution for soma
alignment. Vertical profiling of input strength was achieved by averaging inputs from each horizontal
row and plotting against the vertical distance from soma. Input strength from a specific area was
calculated by averaging input strengths within an area for each neuron and averaging according to
genotype.

For action-potential-dependent ChR2 activation and circuit mapping, analysis was performed as for
sCRACM, except EPSC amplitudes were averaged during 2-30 ms after laser onset to exclude
contamination from polysynaptic responses.

Laser Scanning Photostimulation (LSPS) with glutamate uncaging

LSPS experiments were performed similar to that described previously (Rajkovich et al., 2017;
Shepherd et al., 2003). For all experiments, only brain slices with L2/3 apical dendrites parallel to the
slice surface were used to ensure preservation of the planar barrel cortical geometry of cross-layer
synaptic pathways spanning at least 3 barrel columns. Usually 2-4 brain slices per animal met such
criteria. ACSF included 4 mM $MgCl_2$, 4 mM $CaCl_2$, and CPP (10 µM) to reduced polysynaptic local
circuit activity (Rajkovich et al., 2017; Shepherd et al., 2003), and 4-Methoxy-7-nitroindolinyl-caged-L-
glutamate, MNI glutamate (MNI, 0.3 mM, either Tocris-1490 or HelloBio-HB0423). A 1 ms UV laser
flash (wavelength: 355 nm; power range: 30 - 40 mW; final beam diameter: 20 µm; DPSS Lasers Inc.)
was delivered at individual points within a 16 x 16 grid (50 × 60µm x-y spacing) in a pseudorandom
order. The grid was aligned along the pia surface and centered medial-laterally on the soma location.
The entire grid was repeatedly scanned 2 - 4 times at 40 s intervals. 2-4 maps were acquired for each
neuron included in all datasets.

For each neuron, a single average map was calculated from acquired LSPS maps, where at each
stimulation point the averaged light evoked EPSC area was calculated within a time window of 5-80 ms
following the laser pulse. If a response was observed within 5 ms of LSPS and displayed kinetics visibly
faster than the longer-latency EPSC then it was considered to be a non-synaptic, 'direct' response to
uncaging of glutamate on the recorded neuron. Direct responses were removed from the map and not
included in analysis. For responses with a major component of monosynaptic transmission and minor
contamination from 'direct' activation, 'direct' response component was subtracted from the whole
response by fitting a double-exponential decay equation. An IR-DIC image of the slice with patch

pipettes in place and stimulation grid was acquired prior to LSPS for marking soma location and
anatomical features of the slice (i.e. barrels, etc.). Finally, a color map for each neuron was created. All
individual color maps within genotype were then overlaid upon spatial alignment with respect to the
center of the "home" barrel directly beneath recorded L2/3 neurons. Superimposition of the average
maps was achieved by 1) transposing each map such that the home barrel center was located at the
origin of alignment grid, 2) preserving the medial-lateral orientation of the brain slice, and 3) stretching
the home barrel in x and y dimensions to normalize barrel size. Pixel size for the genotype-averaged
color maps was halved through pixel interpolation (25 x 30 µm) to provide better spatial resolution for
soma alignment. Black pixels within an averaged color map in the figures indicate deleted direct
responses or pixels that did not meet the minimum sampling threshold (minimum of n = 8 neurons per
stimulation point).

Whisker trimming

Unilateral whisker trimming was performed daily on mice starting at P15 until the day before experiment
(P23-30) using a miniature electric shaver. All whiskers from the trimmed pad were maintained at a
length < 1mm. For a litter of mice, half of the mice would be trimmed on the right whisker pad,
ipsilateral to the AAV9.ChR2-mCherry injection (postsynaptic deprivation), while the other half would be
trimmed on the left whisker pad, contralateral to the AAV9.ChR2-mCherry injection (presynaptic
deprivation).

Statistics

All the statistical tests and graphs were performed using GraphPad Prism 8 (GraphPad Software Inc.).
Prior to statistical tests of significance, each dataset underwent normality tests (D'Agostino-Pearson
and Anderson-Darling) to determine if parametric or non-parametric tests should be used. All statistical
tests are two-sided. Only datasets tested to be normally distributed by both tests were subject to
parametric statistics – unpaired or paired t-test. Otherwise, Mann-Whitney test or Wilcoxon matched
pairs test were used as indicated in the figure legends. For data collected from double-patch
experiments, paired statistics were used as each pair of neurons would be considered correlated. For
comparing more than two groups of data, 2-way ANOVA or mixed effect analysis with multiple
comparisons was used. Repeated measures ANOVA was used for vertical profile maps in Figures 1
and 2.

Acknowledgements

We thank Patricia Hahn, Jacob Eli Bowles, and Christopher Williams for technical assistance with the

mice. JRG, KMH, ZZ, conceived various aspects of the project. ZZ and JRG performed experiments

and analyzed data. ZZ and KMH wrote the manuscript with input from all authors. This work was

supported by grants from the NIH grants to KMH and JRG (R01 HD052731, U54HD082008,

U54HD104461) and JRG (R03MH104366).

The authors declare no competing interests.

References

Ashby, M.C., and Isaac, J.T. (2011). Maturation of a recurrent excitatory neocortical circuit by experience-
dependent unsilencing of newly formed dendritic spines. Neuron *70*, 510-521.
10.1016/j.neuron.2011.02.057.
Assali, A., Harrington, A.J., and Cowan, C.W. (2019). Emerging roles for MEF2 in brain development and mental
disorders. Current Opinion in Neurobiology *59*, 49-58. 10.1016/j.conb.2019.04.008.
Belmonte, M.K., Allen, G., Beckel-Mitchener, A., Boulanger, L.M., Carper, R.A., and Webb, S.J. (2004). Autism and
abnormal development of brain connectivity. J Neurosci *24*, 9228-9231. 10.1523/jneurosci.3340-
04.2004.
Bland, N.S., Mattingley, J.B., and Sale, M.V. (2020). Gamma coherence mediates interhemispheric integration
during multiple object tracking. J Neurophysiol *123*, 1630-1644. 10.1152/jn.00755.2019.
Booth, R.D.L., and Happé, F.G.E. (2018). Evidence of Reduced Global Processing in Autism Spectrum Disorder. J
Autism Dev Disord *48*, 1397-1408. 10.1007/s10803-016-2724-6.
Bosman, L.W., Houweling, A.R., Owens, C.B., Tanke, N., Shevchouk, O.T., Rahmati, N., Teunissen, W.H., Ju, C.,
Gong, W., Koekkoek, S.K., and De Zeeuw, C.I. (2011). Anatomical pathways involved in generating and
sensing rhythmic whisker movements. Front Integr Neurosci *5*, 53. 10.3389/fnint.2011.00053.
Bureau, I., Shepherd, G.M., and Svoboda, K. (2008). Circuit and plasticity defects in the developing
somatosensory cortex of FMR1 knock-out mice. J Neurosci *28*, 5178-5188. 10.1523/JNEUROSCI.1076-
08.2008.
Castro, S., Cavelli, M., Vollono, P., Chase, M.H., Falconi, A., and Torterolo, P. (2014). Inter-hemispheric coherence
of neocortical gamma oscillations during sleep and wakefulness. Neurosci Lett *578*, 197-202.
10.1016/j.neulet.2014.06.044.
Ciarrusta, J., Dimitrova, R., Batalle, D., O'Muircheartaigh, J., Cordero-Grande, L., Price, A., Hughes, E., Kangas, J.,
Perry, E., Javed, A., et al. (2020). Emerging functional connectivity differences in newborn infants
vulnerable to autism spectrum disorders. Transl Psychiatry *10*, 131. 10.1038/s41398-020-0805-y.
Contractor, A., Klyachko, V.A., and Portera-Cailliau, C. (2015). Altered Neuronal and Circuit Excitability in Fragile
X Syndrome. Neuron *87*, 699-715. 10.1016/j.neuron.2015.06.017.
Courchesne, E., and Pierce, K. (2005). Why the frontal cortex in autism might be talking only to itself: local over-
connectivity but long-distance disconnection. Curr Opin Neurobiol *15*, 225-230.
10.1016/j.conb.2005.03.001.

Cruz-Martin, A., Crespo, M., and Portera-Cailliau, C. (2010). Delayed stabilization of dendritic spines in fragile X
mice. J Neurosci *30*, 7793-7803. 10.1523/JNEUROSCI.0577-10.2010.
Darnell, J.C., Van Driesche, S.J., Zhang, C., Hung, K.Y., Mele, A., Fraser, C.E., Stone, E.F., Chen, C., Fak, J.J., Chi,
S.W., et al. (2011). FMRP stalls ribosomal translocation on mRNAs linked to synaptic function and autism.
Cell *146*, 247-261. 10.1016/j.cell.2011.06.013.
Dimond, D., Schuetze, M., Smith, R.E., Dhollander, T., Cho, I., Vinette, S., Ten Eycke, K., Lebel, C., McCrimmon, A.,
Dewey, D., et al. (2019). Reduced White Matter Fiber Density in Autism Spectrum Disorder. Cereb Cortex
29, 1778-1788. 10.1093/cercor/bhy348.
Doll, C.A., Yergert, K.M., and Appel, B.H. (2020). The RNA binding protein fragile X mental retardation protein
promotes myelin sheath growth. Glia *68*, 495-508. 10.1002/glia.23731.
Feldman, D.E. (2009). Synaptic mechanisms for plasticity in neocortex. Annu Rev Neurosci *32*, 33-55.
10.1146/annurev.neuro.051508.135516.
Feldmeyer, D. (2012). Excitatory neuronal connectivity in the barrel cortex. Front Neuroanat *6*, 24.
10.3389/fnana.2012.00024.
Friederici, A.D., von Cramon, D.Y., and Kotz, S.A. (2007). Role of the corpus callosum in speech comprehension:
interfacing syntax and prosody. Neuron *53*, 135-145. 10.1016/j.neuron.2006.11.020.
Fries, P. (2009). Neuronal gamma-band synchronization as a fundamental process in cortical computation. Annu
Rev Neurosci *32*, 209-224. 10.1146/annurev.neuro.051508.135603.
Fries, P. (2015). Rhythms for Cognition: Communication through Coherence. Neuron *88*, 220-235.
10.1016/j.neuron.2015.09.034.
Garber, K.B., Visootsak, J., and Warren, S.T. (2008). Fragile X syndrome. Eur J Hum Genet *16*, 666-672.
10.1038/ejhg.2008.61.
Gerfen, C.R., Economo, M.N., and Chandrashekar, J. (2018). Long distance projections of cortical pyramidal
neurons. J Neurosci Res *96*, 1467-1475. 10.1002/jnr.23978.
Gibson, J.R., Bartley, A.F., Hays, S.A., and Huber, K.M. (2008). Imbalance of neocortical excitation and inhibition
and altered UP states reflect network hyperexcitability in the mouse model of fragile X syndrome. J
Neurophysiol *100*, 2615-2626. 10.1152/jn.90752.2008.
Goel, A., Cantu, D.A., Guilfoyle, J., Chaudhari, G.R., Newadkar, A., Todisco, B., de Alba, D., Kourdougli, N., Schmitt,
L.M., Pedapati, E., et al. (2018). Impaired perceptual learning in a mouse model of Fragile X syndrome is
mediated by parvalbumin neuron dysfunction and is reversible. Nat Neurosci *21*, 1404-1411.
10.1038/s41593-018-0231-0.
Goncalves, J.T., Anstey, J.E., Golshani, P., and Portera-Cailliau, C. (2013). Circuit level defects in the developing
neocortex of Fragile X mice. Nat Neurosci *16*, 903-909. 10.1038/nn.3415.
Haberl, M.G., Zerbi, V., Veltien, A., Ginger, M., Heerschap, A., and Frick, A. (2015). Structural-functional
connectivity deficits of neocortical circuits in the Fmr1 (-/y) mouse model of autism. Sci Adv *1*, e1500775.
10.1126/sciadv.1500775.
Hanse, E., Seth, H., and Riebe, I. (2013). AMPA-silent synapses in brain development and pathology. Nat Rev
Neurosci *14*, 839-850. 10.1038/nrn3642.
Hanson, J.E., and Madison, D.V. (2007). Presynaptic FMR1 genotype influences the degree of synaptic
connectivity in a mosaic mouse model of fragile X syndrome. J Neurosci *27*, 4014-4018.
10.1523/JNEUROSCI.4717-06.2007.
Happe, F., and Frith, U. (2006). The weak coherence account: detail-focused cognitive style in autism spectrum
disorders. J Autism Dev Disord *36*, 5-25. 10.1007/s10803-005-0039-0.
Harlow, E.G., Till, S.M., Russell, T.A., Wijetunge, L.S., Kind, P., and Contractor, A. (2010). Critical period plasticity
is disrupted in the barrel cortex of FMR1 knockout mice. Neuron *65*, 385-398.
10.1016/j.neuron.2010.01.024.

Hays, S.A., Huber, K.M., and Gibson, J.R. (2011). Altered neocortical rhythmic activity states in Fmr1 KO mice are
due to enhanced mGluR5 signaling and involve changes in excitatory circuitry. J Neurosci *31*, 14223-
14234. 10.1523/JNEUROSCI.3157-11.2011.

He, Q., Nomura, T., Xu, J., and Contractor, A. (2014). The developmental switch in GABA polarity is delayed in
fragile X mice. J Neurosci *34*, 446-450. 10.1523/JNEUROSCI.4447-13.2014.

Holiga, S., Hipp, J.F., Chatham, C.H., Garces, P., Spooren, W., D'Ardhuy, X.L., Bertolino, A., Bouquet, C., Buitelaar,
J.K., Bours, C., et al. (2019). Patients with autism spectrum disorders display reproducible functional
connectivity alterations. Sci Transl Med *11*. 10.1126/scitranslmed.aat9223.

Hooks, B.M., and Chen, C. (2020). Circuitry Underlying Experience-Dependent Plasticity in the Mouse Visual
System. Neuron *106*, 21-36. 10.1016/j.neuron.2020.01.031.

Huang, Y., Song, N.N., Lan, W., Zhang, Q., Zhang, L., Zhang, L., Hu, L., Chen, J.Y., Zhao, C.J., Li, L., et al. (2013).
Sensory input is required for callosal axon targeting in the somatosensory cortex. Mol Brain *6*, 53.
10.1186/1756-6606-6-53.

Huber, K.M., Gallagher, S.M., Warren, S.T., and Bear, M.F. (2002). Altered synaptic plasticity in a mouse model of
fragile X mental retardation. Proc Natl Acad Sci U S A *99*, 7746-7750. 10.1073/pnas.122205699.

Jonak, C.R., Lovelace, J.W., Ethell, I.M., Razak, K.A., and Binder, D.K. (2020). Multielectrode array analysis of EEG
biomarkers in a mouse model of Fragile X Syndrome. Neurobiol Dis *138*, 104794.
10.1016/j.nbd.2020.104794.

Kim, J.Y., Ash, R.T., Ceballos-Diaz, C., Levites, Y., Golde, T.E., Smirnakis, S.M., and Jankowsky, J.L. (2013). Viral
transduction of the neonatal brain delivers controllable genetic mosaicism for visualising and
manipulating neuronal circuits in vivo. Eur J Neurosci *37*, 1203-1220. 10.1111/ejn.12126.

Li, Q., Becker, B., Jiang, X., Zhao, Z., Zhang, Q., Yao, S., and Kendrick, K.M. (2019). Decreased interhemispheric
functional connectivity rather than corpus callosum volume as a potential biomarker for autism
spectrum disorder. Cortex *119*, 258-266. 10.1016/j.cortex.2019.05.003.

Lovelace, J.W., Ethell, I.M., Binder, D.K., and Razak, K.A. (2018). Translation-relevant EEG phenotypes in a mouse
model of Fragile X Syndrome. Neurobiol Dis *115*, 39-48. 10.1016/j.nbd.2018.03.012.

Manabe, T., Wyllie, D.J., Perkel, D.J., and Nicoll, R.A. (1993). Modulation of synaptic transmission and long-term
potentiation: effects on paired pulse facilitation and EPSC variance in the CA1 region of the
hippocampus. J Neurophysiol *70*, 1451-1459. 10.1152/jn.1993.70.4.1451.

Mientjes, E.J., Nieuwenhuizen, I., Kirkpatrick, L., Zu, T., Hoogeveen-Westerveld, M., Severijnen, L., Rife, M.,
Willemsen, R., Nelson, D.L., and Oostra, B.A. (2006). The generation of a conditional Fmr1 knock out
mouse model to study Fmrp function in vivo. Neurobiol Dis *21*, 549-555.

Mitchell, A.C., Javidfar, B., Pothula, V., Ibi, D., Shen, E.Y., Peter, C.J., Bicks, L.K., Fehr, T., Jiang, Y., Brennand, K.J.,
et al. (2018). MEF2C transcription factor is associated with the genetic and epigenetic risk architecture
of schizophrenia and improves cognition in mice. Mol. Psychiatry *23*, 123-132. 10.1038/mp.2016.254.

Mizuno, H., Hirano, T., and Tagawa, Y. (2010). Pre-synaptic and post-synaptic neuronal activity supports the axon
development of callosal projection neurons during different post-natal periods in the mouse cerebral
cortex. Eur J Neurosci *31*, 410-424. 10.1111/j.1460-9568.2009.07070.x.

Nimchinsky, E.A., Oberlander, A.M., and Svoboda, K. (2001). Abnormal development of dendritic spines in FMR1
knock-out mice. J Neurosci *21*, 5139-5146.

Niu, M., Han, Y., Dy, A.B.C., Du, J., Jin, H., Qin, J., Zhang, J., Li, Q., and Hagerman, R.J. (2017). Autism Symptoms in
Fragile X Syndrome. J Child Neurol *32*, 903-909. 10.1177/0883073817712875.

O'Reilly, C., Lewis, J.D., and Elsabbagh, M. (2017). Is functional brain connectivity atypical in autism? A
systematic review of EEG and MEG studies. PLoS One *12*, e0175870. 10.1371/journal.pone.0175870.

Oliet, S.H., Malenka, R.C., and Nicoll, R.A. (1996). Bidirectional control of quantal size by synaptic activity in the
hippocampus. Science *271*, 1294-1297.

Osterweil, E.K., Chuang, S.C., Chubykin, A.A., Sidorov, M., Bianchi, R., Wong, R.K., and Bear, M.F. (2013).
Lovastatin corrects excess protein synthesis and prevents epileptogenesis in a mouse model of fragile X
syndrome. Neuron *77*, 243-250. 10.1016/j.neuron.2012.01.034.

Panzica, F., Schiaffi, E., Visani, E., Franceschetti, S., and Giovagnoli, A.R. (2019). Gamma electroencephalographic
coherence and theory of mind in healthy subjects. Epilepsy Behav *100*, 106435.
10.1016/j.yebeh.2019.07.036.

Patel, A.B., Hays, S.A., Bureau, I., Huber, K.M., and Gibson, J.R. (2013). A target cell-specific role for presynaptic
Fmr1 in regulating glutamate release onto neocortical fast-spiking inhibitory neurons. J Neurosci *33*,
2593-2604. 10.1523/JNEUROSCI.2447-12.2013.

Patel, A.B., Loerwald, K.W., Huber, K.M., and Gibson, J.R. (2014). Postsynaptic FMRP promotes the pruning of
cell-to-cell connections among pyramidal neurons in the L5A neocortical network. J Neurosci *34*, 3413-
3418. 10.1523/JNEUROSCI.2921-13.2014.

Patneau, D.K., and Mayer, M.L. (1990). Structure-activity relationships for amino acid transmitter candidates
acting at N-methyl-D-aspartate and quisqualate receptors. J Neurosci *10*, 2385-2399.
10.1523/jneurosci.10-07-02385.1990.

Peiker, I., David, N., Schneider, T.R., Nolte, G., Schöttle, D., and Engel, A.K. (2015). Perceptual Integration Deficits
in Autism Spectrum Disorders Are Associated with Reduced Interhemispheric Gamma-Band Coherence. J
Neurosci *35*, 16352-16361. 10.1523/jneurosci.1442-15.2015.

Petreanu, L., Huber, D., Sobczyk, A., and Svoboda, K. (2007). Channelrhodopsin-2-assisted circuit mapping of
long-range callosal projections. Nat Neurosci *10*, 663-668. 10.1038/nn1891.

Petreanu, L., Mao, T., Sternson, S.M., and Svoboda, K. (2009). The subcellular organization of neocortical
excitatory connections. Nature *457*, 1142-1145. 10.1038/nature07709.

Pfeiffer, B.E., and Huber, K.M. (2009). The state of synapses in fragile X syndrome. Neuroscientist *15*, 549-567.
10.1177/1073858409333075.

Pfeiffer, B.E., Zang, T., Wilkerson, J.R., Taniguchi, M., Maksimova, M.A., Smith, L.N., Cowan, C.W., and Huber,
K.M. (2010). Fragile X mental retardation protein is required for synapse elimination by the activity-
dependent transcription factor MEF2. Neuron *66*, 191-197. 10.1016/j.neuron.2010.03.017.

Povysheva, N.V., Gonzalez-Burgos, G., Zaitsev, A.V., Kroner, S., Barrionuevo, G., Lewis, D.A., and Krimer, L.S.
(2006). Properties of excitatory synaptic responses in fast-spiking interneurons and pyramidal cells from
monkey and rat prefrontal cortex. Cereb Cortex *16*, 541-552. 10.1093/cercor/bhj002.

Preisig, B.C., Riecke, L., Sjerps, M.J., Kösem, A., Kop, B.R., Bramson, B., Hagoort, P., and Hervais-Adelman, A.
(2021). Selective modulation of interhemispheric connectivity by transcranial alternating current
stimulation influences binaural integration. Proc Natl Acad Sci U S A *118*. 10.1073/pnas.2015488118.

Rajkovich, K.E., Loerwald, K.W., Hale, C.F., Hess, C.T., Gibson, J.R., and Huber, K.M. (2017). Experience-
Dependent and Differential Regulation of Local and Long-Range Excitatory Neocortical Circuits by
Postsynaptic Mef2c. Neuron *93*, 48-56. 10.1016/j.neuron.2016.11.022.

Rane, P., Cochran, D., Hodge, S.M., Haselgrove, C., Kennedy, D.N., and Frazier, J.A. (2015). Connectivity in Autism:
A Review of MRI Connectivity Studies. Harv Rev Psychiatry *23*, 223-244.
10.1097/HRP.0000000000000072.

Rocha, H., Sampaio, M., Rocha, R., Fernandes, S., and Leao, M. (2016). MEF2C haploinsufficiency syndrome:
Report of a new MEF2C mutation and review. Eur J Med Genet *59*, 478-482. 10.1016/j.ejmg.2016.05.017.

Schulte, T., and Müller-Oehring, E.M. (2010). Contribution of callosal connections to the interhemispheric
integration of visuomotor and cognitive processes. Neuropsychol Rev *20*, 174-190. 10.1007/s11065-010-
9130-1.

Shepherd, G.M., Pologruto, T.A., and Svoboda, K. (2003). Circuit analysis of experience-dependent plasticity in
the developing rat barrel cortex. Neuron *38*, 277-289. 10.1016/s0896-6273(03)00152-1.

Simon, D.M., and Wallace, M.T. (2016). Dysfunction of sensory oscillations in Autism Spectrum Disorder.
Neurosci Biobehav Rev *68*, 848-861. 10.1016/j.neubiorev.2016.07.016.

Südhof, T.C. (2017). Synaptic Neurexin Complexes: A Molecular Code for the Logic of Neural Circuits. Cell *171*,
745-769. 10.1016/j.cell.2017.10.024.
Swanson, M.R., Wolff, J.J., Shen, M.D., Styner, M., Estes, A., Gerig, G., McKinstry, R.C., Botteron, K.N., Piven, J.,
Hazlett, H.C., and Infant Brain Imaging Study, N. (2018). Development of White Matter Circuitry in
Infants With Fragile X Syndrome. JAMA Psychiatry *75*, 505-513. 10.1001/jamapsychiatry.2018.0180.
Testa-Silva, G., Loebel, A., Giugliano, M., de Kock, C.P., Mansvelder, H.D., and Meredith, R.M. (2012).
Hyperconnectivity and slow synapses during early development of medial prefrontal cortex in a mouse
model for mental retardation and autism. Cereb Cortex *22*, 1333-1342. 10.1093/cercor/bhr224.
Wang, C.L., Zhang, L., Zhou, Y., Zhou, J., Yang, X.J., Duan, S.M., Xiong, Z.Q., and Ding, Y.Q. (2007). Activity-
dependent development of callosal projections in the somatosensory cortex. J Neurosci *27*, 11334-
11342. 10.1523/JNEUROSCI.3380-07.2007.
Wang, J., Ethridge, L.E., Mosconi, M.W., White, S.P., Binder, D.K., Pedapati, E.V., Erickson, C.A., Byerly, M.J., and
Sweeney, J.A. (2017). A resting EEG study of neocortical hyperexcitability and altered functional
connectivity in fragile X syndrome. J Neurodev Disord *9*, 11. 10.1186/s11689-017-9191-z.
Yao, S., Becker, B., and Kendrick, K.M. (2021). Reduced Inter-hemispheric Resting State Functional Connectivity
and Its Association With Social Deficits in Autism. Front Psychiatry *12*, 629870.
10.3389/fpsyt.2021.629870.
Zerbi, V., Ielacqua, G.D., Markicevic, M., Haberl, M.G., Ellisman, M.H., A, A.B., Frick, A., Rudin, M., and
Wenderoth, N. (2018). Dysfunctional Autism Risk Genes Cause Circuit-Specific Connectivity Deficits With
Distinct Developmental Trajectories. Cereb Cortex *28*, 2495-2506. 10.1093/cercor/bhy046.
Zhang, Y., Bonnan, A., Bony, G., Ferezou, I., Pietropaolo, S., Ginger, M., Sans, N., Rossier, J., Oostra, B., LeMasson,
G., and Frick, A. (2014). Dendritic channelopathies contribute to neocortical and sensory
hyperexcitability in Fmr1(-/y) mice. Nat Neurosci *17*, 1701-1709. 10.1038/nn.3864.

Figure 1. L2/3 pyramidal neurons in barrel cortex of *Fmr1* KO mice have weak callosal synaptic inputs. (A) Timeline and schematic of experimental paradigm. (B) Example image of S1 contralateral to AAV ChR2-mCherry injection. Left: DIC. Right: Red fluorescence of ChR2-mCherry labelled axons in corpus callosum (CC) and cortex. Recordings were performed on L2/3 neurons in an area of mCherry fluorescence. (C) LED stimulation paradigm (left) and example EPSCs from WT and *Fmr1* KO mice (right). Blue rectangle = 2ms blue LED flash. (D) Left: Raw LED-evoked EPSC amplitudes in WT and *Fmr1* KO animals (WT =172 ±18 pA, n=20; KO=105 ±11 pA, n=19; unpaired t-test); Right: LED-evoked EPSC amplitudes normalized to LED power (WT =2.2 ±0.1, n=20; KO=1.7±0.1, n=19; unpaired t-test). (E) Schematic (left) and example experiment (right) of grid of blue laser stimulation during sCRACM

relative to recorded neuron. (F) Example laser-evoked EPSCs at the locations highlighted in yellow in E from WT and *Fmr1* KO mice. (G) Group average of EPSC amplitudes evoked at different locations relative to the cell soma in WT and *Fmr1* KO mice. Individual maps are aligned by the location of soma (cyan dot). Pixel color represents the average amplitude of EPSCs evoked from that location. (H) Left: Vertical profile of mean synaptic input strength (mean input − average of EPSC amplitudes from individual locations within a specific area) (soma, x=0) (*Fmr1* n.s. p=0.16, F(1, 60)=2.060; *Fmr1* x vertical position, F(15, 849)=1.723; mixed-effects model). Right: Mean of EPSC amplitudes in the outlined area in G (white) and left graph (blue), normalized to laser power. (WT=0.674±0.098, n=40; KO=0.312±0.167, n=22; Mann Whitney). For this and all figures, error bars represent standard error mean (SEM). *p<0.05, **p<0.01.

Figure 2. Postnatal and postsynaptic deletion of *Fmr1* in L2/3 pyramidal neurons cell autonomously weakens callosal synaptic inputs. (A) Timeline and schematic of experimental paradigm for juvenile recordings. (B) Simultaneous patch clamping of a neighboring AAV Cre-GFP+, *Fmr1* KO (arrowhead) and GFP-, *Fmr1* WT (arrow) pyramidal neurons in L2/3 of barrel cortex. Left: DIC. Right: Green fluorescence. (C) LED bulk stimulation paradigm (left) and example EPSCs from a pair of WT and *Fmr1* KO neurons (right). Scale = 100pA, 20 ms. (D) Left: Group average of LED-induced EPSC amplitudes in WT and *Fmr1* KO neurons at P18-20 (top) (WT=238 ±30 pA; KO=281 ±27 pA,

n=19 pairs, n.s.) and P23-30 (bottom) (WT=424 ±42 pA; KO=250±46 pA, n=15 pairs; paired t-test);

Right: EPSC amplitudes from individual cell pairs (open circles). Mean ±SEM (filled circle). Diagonal

line represents equality. (E) EPSC amplitudes, normalized to LED power, across different ages (replot

from (D)) (*Fmr1* x Age ***$p<0.001$, $F_{(1, 32)}=13.87$; *Fmr1*, Age, ns, ANOVA; #$p<0.1$, multiple

comparison). (F) Timeline and schematic of 2-month old recordings. (G) Left: Group average of LED-

induced EPSC amplitudes in WT and *Fmr1* KO neurons at 2-month old (WT=346 ±47 pA; KO=260 ±43

pA, n=11 pairs, paired t-test); Right: Distribution of values from individual cell pairs. *$p<0.05$;

$p<0.01$;*$p<0.001$.

B. Transfection rate in L2/3   ■ GFP+   ■ GFP-

 Figure 2 – figure supplement 1. Sparse transfection of Cre-GFP in the recorded hemisphere. (A)

Top: Representative epifluorescence image of GFP for an acute coronal slice (300 µm thick) showing

barrel cortex from the Cre-GFP injected hemisphere under low magnification (4x). Cre-GFP is sparsely

expressed across all cortical layers in a similar level (see green arrowheads for example GFP+ cells).

The medial side (M) has higher transfection than the lateral side (L) because it's closer to the needle

tract (not shown in the image) going into the ventricle.   Bottom: representative composite images of

GFP (green) and DIC under high magnification (40x) for L2/3, L4 and L5. (B) Estimation of percentage

Cre-GFP transfection in L2/3 of the ipsilateral barrel cortex from 4 animals individually. These were

calculated from 5-15 field of views (0.05 mm^2 each) from at least 2 slices per animal. Transfection rates

in layers 4 and 5 are similar to L2/3.

Figure 2 – figure supplement 2. Weakening of callosal synaptic inputs onto postsynaptic *Fmr1* KO L2/3 neurons is confirmed by sCRACM with spatial distribution. (A) Schematic (left) and image (right) of blue laser stimulation grid configuration and dual recordings for sCRACM experiments. (B) Example of laser-induced EPSCs at location highlighted in A (yellow) from WT and *Fmr1* KO pair. Scale = 100pA, 50 ms (C) Averaged color map of subcellular distribution of callosal synaptic input strengths onto WT and *Fmr1* KO L2/3 pyramidal neurons at P23-30. (D) Vertical profile of mean synaptic inputs strength along the neuronal dendritic tree (soma, x=0) (*Fmr1* F(1, 13)=5.956; *Fmr1* x vertical position F(14, 144)=1.793; mixed-effects model). (E) Left: Quantification of mean synaptic inputs strength from the highlighted positions in C (white) and D (blue), (WT=46 ±7; KO= 26 ±6 pA, n=14 pairs; Wilcoxon test); Right: Distribution of values from individual cell pairs. *p<0.05;***p<0.001.

Figure 3. Callosal synapses onto *Fmr1* KO neurons have reduced quantal event frequency and a selective weakening of AMPA receptor-mediated synaptic transmission. (A) Example traces of Sr^{2+}-evoked quantal events. Scale = 25pA, 100 ms. Baseline, spontaneous events are defined as those which occur within 1s prior to LED flash (red dotted line) and evoked events occur 50-350 ms after LED

flash (red line). (B) Left: Baseline frequency (top) (WT=1.3±0.1, KO=1.2±0.2 Hz, n.s., paired t-test) and amplitude (bottom) (WT=12±1, KO=11±1 pA, n.s., Wilcoxon test) of quantal EPSCs for WT and *Fmr1* KO neuron pairs; Right: distribution of values from individual cell pairs. (C) Left: Evoked frequency (top) (WT=8.3±0.5, KO=6.7±0.5 Hz, *p<0.05, Wilcoxon test) and amplitude (bottom) (WT=15±1, KO= 15±1 pA, n.s., Wilcoxon test, n=13 pairs) of quantal EPSCs for WT and *Fmr1* KO neuron pairs; Right: distribution of values from individual cell pairs. (D) Top: Example NMDAR EPSCs from a WT and *Fmr1* KO pair. Scale = 25pA, 20 ms. Bottom: LED-induced NMDAR EPSC amplitudes of WT and *Fmr1* KO neuron pairs (WT=124±20, KO=120±18 pA, n.s., paired t-test, n=15 pairs); Right: distribution of values from individual cell pairs.

Figure 3 – figure supplement 1. Coefficient of variance (C.V.) of LED-evoked EPSCs from P23-30 cell pairs in figure 2D and 5B$_1$. Left: summary statistics (WT=0.153$\pm$0.013, KO=0.181$\pm$0.015 Hz, *p<0.05, paired t-test, n=30 pairs); Right: distribution of values from individual cell pairs.

Figure 4. Cell autonomous deletion of *Fmr1* in postsynaptic L2/3 or L5 neurons results in weak action-potential driven synaptic transmission from callosal inputs. (A) Top: Average color map of action-potential mediated callosal synaptic input strengths onto pairs of WT and *Fmr1* KO L2/3 pyramidal neurons at P23-30; Bottom: Example responses from the highlighted positions (white). Scale = 100pA, 50 ms. (B) Top: Mean callosal inputs strength from area highlighted in white in A (WT=61±13, KO=35±6 pA, n=11 pairs, *p<0.05, paired t-test); Bottom: Distribution of values from individual cell pairs. (C) Top: Average color map of callosal input strengths onto pairs of WT and *Fmr1* KO L5 pyramidal neurons at P23-30; Bottom: Example responses from the highlighted area (white). Scale = 200pA, 60 ms. (D) Top: Mean callosal input strengths onto L5 neurons from the area highlighted in white in C (WT=49±9, KO=31±4 pA, n=12 pairs, *p<0.05, paired t-test). Bottom: distribution of values from individual cell pairs.

Figure 5. Sensory deprivation by whisker trimming normalizes callosal input strength onto postsynaptic *Fmr1* KO neurons. (A) Experimental paradigm. Trimming the whisker pad either ipsilateral or contralateral to the AAV-Cre-GFP injected hemisphere deprived either the presynaptic callosal projection neurons or postsynaptic *Fmr1* KO neurons, respectively, of patterned sensory experience- driven activity. (B_1, B_2) Left: Raw LED-induced EPSC amplitudes in WT and *Fmr1* KO neurons with presynaptic deprivation (WT=510±25; KO=376±41 pA, n=16 pairs, **p<0.01, paired t-test) or with postsynaptic deprivation (WT=428±36; KO=433±31 pA, n=17 pairs, n.s., paired t-test): Right: Values from individual cell pairs. (C) LED-induced EPSC amplitudes normalized to LED power (replot

969   from (B)) (*Fmr1* x deprivation interaction *$p<0.05$, $F_{(1, 31)}=4.977$, ANOVA; presynaptic deprivation WT

970   vs. *Fmr1* KO, *$p<0.05$, *Fmr1* KO presynaptic deprivation vs. postsynaptic deprivation, *$p<0.05$, Sidak's

971   multiple comparisons).

Figure 5 – figure supplement 1. Sensory dependent weakening of callosal inputs is confirmed

by sCRACM. (A) Averaged color map of subcellular distribution of callosal synaptic inputs onto WT and

Fmr1 KO neurons with presynaptic (A_1) or postsynaptic (A_2) sensory deprivation. (B_1, B_2) Left: Mean

synaptic inputs strength onto soma and proximal apical dendrites (highlighted in white in A) for WT and

Fmr1 KO neurons with presynaptic (WT=43±7; KO=31±5 pA, n=16 pairs, *p<0.05, Wilcoxon test) or

postsynaptic (WT=42±4; KO=42±6 pA, n=17 pairs, n.s., Wilcoxon test) sensory deprivation; Right:

values from individual cell pairs.

1000

1001

1002

1003

1004

1005

1006 **Figure 5 – figure supplement 2.** Miniature (m) EPSC frequency and amplitude, as well as input
1007 resistance from pairs of WT and postsynaptic *Fmr1* KO L2/3 pyramidal neurons in P23-30 sensory
1008 intact, presynaptic sensory deprived and postsynaptic sensory deprived animals.

1009

1010

Figure 6. Postsynaptic deletion of *Fmr1* in L2/3 neurons does not affect excitatory synaptic inputs from local columnar inputs. (A) Experimental design schematic. (B) Left: Position of UV laser scanning photostimulation (LSPS) grid (grey dots) relative to cortical layers and recorded neurons in L2/3 and L4 barrels (grey squares). Right: IR-DIC image of dual recordings in L2/3 in a slice with laser stimulation grid (red). Yellow dots indicate L4 home barrel. (C) Example of EPSCs in a pair of WT and *Fmr1* KO L2/3 neurons in response to LSPS and glutamate uncaging in L4 home barrel (yellow dots in B; right). (D) Color map of spatial distribution of average synaptic input strengths in response to LSPS. L4 home barrel (white rectangle). Cyan dots represent locations of soma and black pixels are direct responses. (E) Mean synaptic input strength from L4 home barrel, L4 adjacent barrels, adjacent L2/3 and L5A (WT vs. *Fmr1* KO, n.s., n=17 pairs, Wilcoxon or paired t-test) and values from individual cell pairs.

1022

1023

Figure 6 – figure supplement 1. Postsynaptic deletion of *Fmr1* does not affect local columnar
circuit inputs onto L2/3 pyramidal neurons at 2 weeks of age. (A) Right: Sparse deletion of *Fmr1* in
L2/3 neurons using AAV.GFP-Cre injection at P1 does not affect local synaptic input strength from L4
home barrel (L4H) measured at P14-17 (n = 8 pairs). (B) Similar as (A), but experiment was done in
mosaic *Fmr1* females (*Fmr1$^{-/gfp}$*) (n = 11 pairs). Half of the cells are *Fmr1* KO GFP- while the other half
are *Fmr1* WT GFP+ due to X chromosome inactivation. (C) Averaged LSPS color maps (left) and
quantification of local synaptic input strength from L4 home and adjacent (L4A) barrels (right)
comparing across WT and *Fmr1* global KO animals at 3 weeks of age (P18-25) (n = 15 for WT, n = 20

for KO). Quantified inputs were normalized to the respective laser power and log transformed to enable

cross-animal comparison.

1034

Figure 6 – figure supplement 2. Summary model of the role of postsynaptic *Fmr1* on development of callosal and local neocortical synapses. During development of cortical circuits for mouse primary somatosensory barrel cortex (postnatal 4 weeks), *Fmr1* functions postnatally and cell autonomously in postsynaptic L2/3 pyramidal neurons to promote maturation or stability of callosal synaptic connections by promoting/maintaining AMPAR transmission. Without postsynaptic *Fmr1*, our results suggest that there are more "NMDAR-only" or "silent" immature callosal synapses onto *Fmr1* KO neurons and callosal synapses undergo synaptic silencing that depends on sensory experience driven activity of postsynaptic L2/3 neurons. Postsynaptic *Fmr1* selectively promotes callosal synaptic connections for L2/3 neurons as local circuit inputs from other layers or columns are not changed.

1044

1045

1046

1047

1048

1049

1050