FMRP regulates experience-dependent maturation of callosal synaptic connections and bilateral cortical synchrony

Zhe Zhang[1], Jeffrey Rumschlag[2], Carrie R. Jonak[3], Devin K. Binder[3], Khaleel A. Razak[2], *Jay R. Gibson[1], *Kimberly M. Huber[1]

[1]Department of Neuroscience, O'Donnell Brain Institute, University of Texas Southwestern Medical Center, Dallas, TX, United States.

[2]Department of Psychology, University of California, Riverside, Riverside, CA, United States.

[3]Division of Biomedical Sciences, School of Medicine, University of California, Riverside, Riverside, CA, United States.

*Correspondence: Jay.Gibson@UTSouthwestern.edu and Kimberly.Huber@UTSouthwestern.edu

Abstract

Reduced structural and functional interhemispheric connectivity correlates with the severity Autism Spectrum Disorder (ASD) behaviors in humans. Little is known of how ASD-risk genes regulate callosal connectivity. Here we show that *Fmr1*, whose loss-of-function leads to Fragile X Syndrome (FXS), cell autonomously promotes maturation of callosal excitatory synapses between somatosensory barrel cortices. Postnatal, cell-autonomous deletion of *Fmr1* in postsynaptic Layer (L) 2/3 or L5 neurons results in a selective weakening of AMPA receptor- (R), but not NMDA receptor-, mediated synaptic function, indicative of increased "silent", or immature, callosal synapses. Sensory deprivation by contralateral whisker trimming normalizes callosal input strength, suggesting that experience-driven activity of postsynaptic *Fmr1* KO L2/3 neurons weakens callosal synapses. In contrast to callosal inputs, synapses originating from local L4 and L2/3 circuits are normal, revealing an input-specific role for postsynaptic *Fmr1* in promoting callosal synaptic connectivity. Multielectrode EEG recordings in awake *Fmr1* KO mice find reduced coherence of bilateral neural oscillations in beta and gamma frequencies during rest and sensory stimulation, reflecting reduced interhemispheric integration. Our findings reveal the cellular and synaptic mechanisms by which loss of *Fmr1* leads to reduced interhemispheric connectivity and suggests a novel biomarker of brain dysfunction in FXS.

Introduction

Disrupted structural and functional brain connectivity has been widely observed in patients with autism spectrum disorder (ASD) (Dimond et al., 2019; Holiga et al., 2019; Rane et al., 2015). A common finding in ASD is reduced corpus callosum integrity and interhemispheric functional connectivity, the latter of which correlates with autistic symptoms (Li et al., 2019; O'Reilly et al., 2017; Yao et al., 2021). The corpus callosum connects bilateral hemispheres and functions to synchronize cortical circuits necessary for sensory-motor processing, attention, perception and higher cognitive functions (Schulte and Müller-Oehring, 2010). Little is known of how autism-risk genes regulate development of callosal connectivity and the cellular or synaptic basis of reduced functional connectivity in ASD.

To provide insight into these questions, we have studied the role of the Fragile X Mental Retardation gene (*Fmr1*) in development of callosal synaptic connections in mice. Loss of function mutations in *FMR1* in humans cause Fragile X Syndrome (FXS), the most common inherited form of intellectual disability and leading monogenic cause of ASD (Garber et al., 2008) (Niu et al., 2017). Children, age 6-24 months, with FXS have reduced structural integrity of white matter tracts, including the corpus callosum (Swanson et al., 2018). *Fmr1* knockout (KO) mice, an animal model for FXS, have a similar reduction in white matter tract integrity as well as decreased functional coherence among different cortical regions as measured with functional MRI (Haberl et al., 2015) (Zerbi et al., 2018). Specifically, neural networks involved in sensory processing show severe functional underconnectivity, including both intra- and inter-hemispheric cortical circuits (Zerbi *et al.*, 2018). The cellular or synaptic basis for decreased inter-region functional coherence in FXS is unknown, and whether this is due to direct or indirect roles for *Fmr1* in cortical neurons is unclear.

In addition to reduced long-range connectivity, there are reports of increased connectivity and hyperexcitability of local cortical circuits in humans with ASD and FXS (Courchesne and Pierce, 2005) (Ciarrusta et al., 2020). In the *Fmr1* KO mouse, there is strong evidence for hyperexcitable local cortical circuits in *Fmr1* KO mice, including in visual, auditory and somatosensory cortices (Contractor et al., 2015; Gibson et al., 2008; Goncalves et al., 2013; Hays et al., 2011; Osterweil et al., 2013). Hyperactive cortical circuits are also observed in humans with FXS, and *Fmr1* KO mice as an increase in resting state gamma power in the resting state EEG (Jonak et al., 2020; Lovelace et al., 2018; Wang et al., 2017). Observed changes that may likely contribute to hyperexcitability of local circuits include synaptically hyperconnected pyramidal neurons, reduced inhibitory neuron activity and changes in intrinsic excitability (Gibson *et al.*, 2008; Goel et al., 2018; He et al., 2014; Zhang et al., 2014). Layer (L) 2/3 and L5 cortical pyramidal neurons receive and integrate excitatory synaptic inputs from homotopic contralateral hemisphere (callosal) as well as other long-range inputs from ipsilateral cortical regions and local cortical

circuits. Little is known if or how the development of local and long-range synaptic inputs is balanced or if this balance is regulated by ASD-risk genes.

Fmr1 encodes Fragile X Mental Retardation Protein (FMRP), an RNA binding protein that interacts with many mRNAs including those encoding pre- and post-synaptic proteins (Darnell et al., 2011). It is perhaps through this diversity of mRNA targets that FMRP regulates multiple properties of synapses, including maturation, pruning and acute forms of synaptic plasticity (Pfeiffer and Huber, 2009) (He and Portera-Cailliau, 2013). Regarding excitatory cortical synapses in somatosensory cortex, *Fmr1* regulates maturation of thalamocortical inputs to L4 neurons as well as between local cortical circuits. A common finding is the slow maturation of excitatory synapses on *Fmr1* KO neurons, as observed by the delayed presence of NMDA receptor-only, or "silent", synapses and acquisition of AMPAR-mediated synaptic transmission at thalamocortical synapses onto L4 neurons and between locally connected L5 neurons. Furthermore, L4 to L2/3 synaptic inputs are weak and delayed in their developmental strengthening, and dendritic spines on cortical pyramidal neurons are thin and filopodial-like, resembling immature spines (Cruz-Martin et al., 2010; He and Portera-Cailliau, 2013; Nimchinsky et al., 2001). L5 local synaptic connections ultimately mature and appear normal during the 4th postnatal week but then fail to prune, which results in hyperconnectivity of L5 neurons at 4-5 weeks of age as compared to wild-type cortex. Importantly, the delayed development and hyperconnectivity of L5 circuits are due to a cell autonomous, postsynaptic role for FMRP in L5 neurons, suggesting a direct role of FMRP in coordinating multiple synapse development processes. The cellular locus of FMRP function in development of L4 to L2/3 inputs or whether postsynaptic FMRP similarly regulates development of long-range synaptic connections is unknown.

Using viral mediated expression of Channelrhodopsin 2 (ChR2) in callosal projecting cortical neurons (Petreanu et al., 2009), we observe weak callosal synaptic inputs onto L2/3 and L5 *Fmr1* KO neurons that are mediated by a cell autonomous, postsynaptic, and postnatal role of FMRP. Callosal inputs have a selective deficit in AMPA receptor (R) but not NMDAR- mediated synaptic transmission, suggesting the presence of silent, immature synapses onto mature *Fmr1* KO neurons. Sensory deprivation by whisker trimming normalized callosal input strength suggesting that sensory experience-driven activity of postsynaptic *Fmr1* KO neurons weakens callosal synapses. Surprisingly, local excitatory inputs were normal on L2/3 neurons with postsynaptic *Fmr1* deletion, revealing differential regulation of local and callosal synaptic connections by FMRP. Multielectrode array (MEA) EEG recordings in awake, adult *Fmr1* KO mice suggest that weak callosal synaptic connectivity leads to reduced synchrony between bilateral homotopic cortical circuits as measured by reduced inter-site phase clustering (ISPC) at fast time scales (in the beta-gamma frequency (20-60 Hz), both at rest and during sensory stimulation. These results

reveal a cellular and synaptic substrate for reduced interhemispheric connectivity in FXS as well as
imbalanced activity with local circuits.

Results

Optogenetic activation of callosal axons shows weak excitatory synaptic inputs onto L2/3
pyramidal neurons in barrel cortex of *Fmr1* KO mice

To measure callosal synaptic function and connectivity between hemispheres, mCherry-tagged
Channelrhodpsin-2 (ChR2) was expressed in callosal projecting cortical neurons by injecting AAV9 into
in one hemisphere of primary somatosensory cortex of postnatal day (P) 1 pups. At P18-25, acute coronal
slices containing barrel cortex contralateral to the AAV injected side were prepared and whole cell patch
clamp recordings of L2/3 pyramidal neurons were performed in the region innervated by fluorescently
labelled callosal axons (Fig. 1A-B). Monosynaptic excitatory postsynaptic currents (EPSCs) were elicited
by stimulating ChR2 expressing callosal axons with brief (2ms) blue LED light pulses in the presence of
tetrodotoxin (TTX) and 4-aminopyridine (4-AP) as described previously (Petreanu et al., 2007; Rajkovich
et al., 2017) (Fig. 1C). The blue LED was delivered through a 40X lens centered around the soma of
approximately 200 µm diameter in size and thus the amplitude of LED-evoked EPSCs likely reflects the
overall strength of callosal synaptic inputs onto the recorded neuron. The amplitude of LED-evoked
EPSCs in *Fmr1* KO neurons was reduced by ~40% compared to WT. Similarly reduced EPSC amplitudes
were observed by comparing raw values or when normalized to LED power (Fig. 1D). To determine if
weak callosal synaptic inputs were localized to a subcellular region on recorded L2/3 neurons (e.g. apical
vs. distal dendrites), we utilized a method termed subcellular Channelrhodopsin2-assisted circuit
mapping (sCRACM (Petreanu *et al.,* 2009)). A blue laser (473 nm) was applied across an array of
locations, relative to the soma of the recorded neuron, in a pseudorandom order (Fig. 1E) to elicit
glutamate release from ChR2-expressing callosal axon terminals and EPSCs at localized dendritic sites
(Fig. 1F). The mean amplitude of EPSCs at each site was converted into a color map for each recorded
neuron and these maps were then aligned by the position of soma and averaged within genotype. The
resulting average map depicts the subcellular distribution of callosal synaptic input strengths onto WT or
Fmr1 KO neurons (Fig. 1G). Results reveal that callosal synaptic inputs onto WT L2/3 neurons are
strongest at the proximal apical dendrites (Fig. 1H), consistent with previous reports (Petreanu *et al.*,
2009) and there is an interaction of *Fmr1* and vertical position (*$p<0.05$, $F_{(15, 849)}=1.723$; mixed-effects
ANOVA). Comparison of mean EPSC amplitudes from proximal apical dendrites reveals a 50% reduction
in callosal synaptic input strength in *Fmr1* KO mice (Fig. 1H).

Postsynaptic *Fmr1* cell-autonomously promotes callosal synapses

FMRP is known to have both pre- and postsynaptic, cell autonomous roles in excitatory synapse development, depending on the cell type and synaptic input (Patel et al., 2013; Patel et al., 2014; Pfeiffer et al., 2010). To determine the cell-autonomous and synaptic locus of FMRP function in development of callosal synapses, we deleted *Fmr1* in a sparse population of postsynaptic L2/3 pyramidal neurons by injecting AAV expressing GFP-tagged Cre (AAV9.GFP-Cre) into the left lateral ventricle of P1 pups(Kim et al., 2013) with a floxed *Fmr1* gene (*Fmr1^{fl/fl}* or *Fmr1^{fl/y}*) (Fig. 2A) (Mientjes et al., 2006). In the same animals, AAV.ChR2-mCherry was injected into the right barrel cortex to label callosal axons as in Fig. 1. This experimental design allowed us to measure the cell autonomous effect of postsynaptic FMRP deletion in L2/3 neurons in a primarily WT cortex on synapses from WT callosal axons. At P18-30, acute cortical slices were prepared and simultaneous whole cell recordings were performed on pairs of neighboring L2/3 pyramidal neurons in left barrel cortex with one being a GFP (-), or WT neuron, and the other a GFP (+), or *Fmr1* KO neuron (Fig. 2A, 2B). Monosynaptic EPSCs from callosal axons were elicited with either bulk LED stimulation or blue laser, to perform sCRACM, as in Fig. 1 (Fig. 2C, F-G). Interestingly, at the earliest age for recording (P18-20), overall callosal synaptic input strengths were not different between WT and postsynaptic *Fmr1* KO neurons (Fig. 2D). However, at P23-30, postsynaptic *Fmr1* KO neurons had a 40% reduction in callosal synaptic input strength compared to neighboring WT neurons (Fig. 2D). Comparison across age groups using ANOVA shows significant interaction between the genotype and age (*Fmr1* x Age ***p<0.001, $F_{(1, 32)}=13.87$)(Fig. 2E). In addition, we observed a strong trend of callosal input strength to increase with developmental age in WT, but not *Fmr1* KO, neurons (Fig. 2E). Similar results are obtained with sCRACM which reveal a ~45% reduction in callosal input strength onto proximal dendrites of *Fmr1* KO neurons (Fig. 2H-J). These data demonstrate that *Fmr1* in postsynaptic L2/3 neurons cell-autonomously promotes the development and/or strengthening of callosal synaptic inputs.

Selective weakening of AMPAR-, but not NMDAR-, mediated synaptic transmission from callosal inputs onto *Fmr1* KO L2/3 pyramidal neurons

Weak callosal synaptic EPSCs onto postsynaptic *Fmr1* KO L2/3 pyramidal neurons could be due to reductions in the number of synaptic connections, presynaptic release probability (Pr), the strength of individual synapses, or any combination of these. To further investigate the synaptic basis of weak callosal inputs in P23-30 slices, we recorded in strontium (Sr^{2+}), substituted for Ca^{2+}, in the external ASCF which results in asynchronous vesicle release and measurement of quantal synaptic events evoked from

callosal axons by the LED (Oliet et al., 1996). Changes in the amplitude of quantal events relate to synapse strength, whereas changes in the frequency of events relate to synapse number and/or presynaptic release probability. Postsynaptic *Fmr1* KO neurons received >20% fewer evoked events in Sr^{2+} in comparison to neighboring WT neurons with no change in event amplitude (Fig. 3A-C). The baseline frequency and amplitude of spontaneous EPSCs, prior to LED stimulation, were not different between genotypes. These results suggest the reduced EPSCs onto postsynaptic *Fmr1* KO neurons are a consequence of reduced presynaptic release probability, functional synapse number or both.

To further differentiate between release probability and functional synapse number, we measured NMDA receptor (R) mediated EPSCs from callosal axons onto WT or postsynaptic *Fmr1* KO L2/3 neurons. Because NMDARs are colocalized with AMPARs at excitatory synapses, reduced glutamate release probability from callosal axons onto *Fmr1* KO neurons would be expected to result in weak NMDAR-EPSCs. To evoke isolated NMDAR EPSCs from callosal axons, we included the AMPAR antagonist DNQX, glycine, and low Mg^{2+} (0.1 mM) in the ACSF and voltage clamped neurons at -70 mV. In contrast to AMPAR-mediated EPSCs, amplitudes of NMDAR EPSCs evoked from callosal axons were not different between WT and neighboring postsynaptic *Fmr1* KO neurons (Fig. 3D). This observation suggests that the NMDAR content within callosal synapses are similar for WT and KO neurons and that there is no change in presynaptic release probability for the callosal axons. Taken together with the reduced frequency of AMPAR quantal events (Fig. 3C), we conclude that there is a higher percentage of NMDAR-only, or 'silent' callosal synapses onto postsynaptic *Fmr1* KO L2/3 pyramidal neurons.

Postsynaptic *Fmr1* KO pyramidal neurons in both L2/3 and L5 receive weak callosal synaptic inputs with action-potential-driven synaptic transmission

For experiments described in Figs. 1 and 2, we included TTX and 4-AP in the bath to isolate evoked EPSCs from ChR2-expressing callosal axons and block potential contamination from polysynaptic local L2/3 inputs. A caveat of this approach is that callosal synaptic transmission is not triggered by action potentials, but by direct depolarization and activation of voltage-gated Ca^{2+} channels at the callosal axon terminal. To determine if similar results are observed with action potential-evoked synaptic transmission, we repeated experiments without TTX/4AP, but increased Ca^{2+}/Mg^{2+} and added the NMDAR blocker, CPP, to reduce polysynaptic responses from local circuits. We also used a blue laser to depolarize a local (30µm) area of callosal axons while limiting activation of local circuits (Fig. 4A). Quantification of mean EPSC amplitude within the region covering soma and major dendrites shows that L2/3 pyramidal

neurons with postsynaptic *Fmr1* deletion have a 45% reduction in callosal synaptic input strength as
compared to neighboring WT neurons (Fig. 4A-B).

L5 pyramidal neurons are another major target of callosal axons (Petreanu *et al.*, 2007; Wang et al.,
2007). To determine if weak callosal synaptic inputs to postsynaptic *Fmr1* KO neurons are specific to
L2/3, we performed the same experiment on L5 pyramidal neurons. Similar to L2/3, L5 pyramidal neurons
with postsynaptic *Fmr1* deletion have a 40% reduction in the strength of callosal inputs stimulated around
the soma and apical dendrites (Fig. 4C-D). Together, these results show that postsynaptic deletion of
Fmr1 in pyramidal cortical neurons generally weakens callosal synaptic inputs and this is observed with
action potential driven synaptic transmission and across different layers.

Sensory deprivation by whisker trimming normalizes callosal synaptic input strengths in *Fmr1*
KO L2/3 neurons

Targeting, branching and elaboration of callosal axons into the contralateral neocortex occurs postnatally
(~P5-15) and depends on activity of presynaptic and postsynaptic cortical neurons as well as whisker
sensory experience (Huang et al., 2013; Mizuno et al., 2010; Wang *et al.*, 2007). This suggests that
sensory experience may interact with FMRP to regulate development of callosal synapses. To test the
interaction of sensory experience and FMRP, we used the paradigm above to sparsely delete *Fmr1* in
postsynaptic L2/3 neurons in the left barrel cortex and express ChR2-mCherry in callosal axon projecting
neurons in right hemisphere (Fig. 5A). Beginning at P15, we unilaterally trimmed whiskers daily on the
right whisker pad which would reduce the most direct ascending sensory-driven patterned activity to the
left hemisphere containing *Fmr1* KO 2/3 neurons. In littermates, we trimmed whiskers on the left whisker
pad which would primarily deprive the L2/3 callosal projection neurons of sensory driven patterned activity.
Therefore, in an individual mouse, either the postsynaptic recorded L2/3 neurons receiving inputs from
ChR2-expressing callosal axons were sensory deprived (postsynaptic deprivation condition) or the
presynaptic, ChR2-expressing callosal projection neurons were deprived of sensory experience
(presynaptic deprivation condition) (Fig 5A). To reduce effects of sensory deprivation on the early growth
and branching of callosal projection axons, whisker trimming began at P15 and continued until the day
before slice recordings. Recordings of EPSCs evoked from callosal axons by either LED stimulation (Fig
5B) or sCRACM (Fig 5C-D) were obtained from pairs of neighboring WT and *Fmr1* KO L2/3 neurons.
Because we recorded from pairs of WT and postsynaptic *Fmr1* KO neuron neighbors, any effects of
sensory deprivation on callosal axon innervation, growth, or branching in a given cortical region would
be expected to similarly affect each genotype. Thus, our results reflect the cell autonomous effects of

Fmr1 on synaptic function or connectivity. In the "presynaptic deprivation" condition, callosal synaptic

input strength was weak (27% reduction) onto postsynaptic *Fmr1* KO L2/3 neurons as compared to WT,

similar to that observed in whisker intact mice (Fig 2D). In contrast, in the "postsynaptic deprivation"

condition, callosal synaptic inputs strengths are similar onto WT and *Fmr1* KO neurons. To compare the

callosal synaptic inputs strength within genotypes and across sensory deprivation paradigms, we

normalized each LED induced EPSC to its stimulation power, log transformed the value and plot all the

data in a single graph (Fig 5E). A 2 way ANOVA and multiple comparison tests of these data revealed

an interaction of deprivation condition and *Fmr1* (*$p<0.05$, $F_{(1, 31)}=4.977$, ANOVA;). Surprisingly, in WT

neurons callosal synaptic input strengths were not different between pre- and postsynaptic deprivation

conditions. Instead, callosal input strengths onto postsynaptic *Fmr1* KO neurons were weak, decreased

by ~45%, in the presynaptic deprivation condition as compared to postsynaptic deprivation. These data

suggest that whisker experience-driven activity of *Fmr1* KO L2/3 neurons weakens callosal synaptic

inputs.

Input specific strengthening of callosal synaptic connections by postsynaptic FMRP

A previous study reported weak L4 to L2/3 synaptic inputs in the barrel cortex of global *Fmr1* KO mice

(Bureau et al., 2008), suggesting that *Fmr1* generally promotes excitatory synapse strength onto L2/3

neurons regardless of whether they are from local or long-range sources. To determine if postsynaptic

Fmr1 promotes excitatory synapse development from local cortical circuits in a cell autonomous manner

as it does for callosal inputs, we assessed local input strengths using Laser Scanning Photo-Stimulation

(LSPS) with glutamate uncaging. Slices were bathed in MNI-caged glutamate and pseudorandom flashes

of a UV laser (355 nm) beam at individual locations (20 μm diameter) within a 16-by-16 grid surrounding

the recorded L2/3 neurons focally released glutamate to evoke action potentials and synaptic

transmission from neurons at that location. The grid for UV laser flashing was positioned to stimulate

neurons in Layers 2/3-5 of home and adjacent barrels. At P23-30, LSPS was performed on pairs of

simultaneously recorded WT and neighboring L2/3 neurons with postsynaptic *Fmr1* deletion (Fig. 6A-B).

The amplitude of monosynaptic EPSCs evoked from each position in the slice for an individual neuron

were converted into a color map and then individual maps were aligned to the home barrel to create an

average color map per genotype (Fig. 6C-D). Responses from direct glutamate activation onto recorded

neurons, as described in methods, were excluded from analysis, and represented by black pixels. EPSCs

evoked from L4 home or adjacent barrels, L5A or L2/3 were unaffected on *Fmr1* KO L2/3 neurons (Fig.

6E), in contrast to the weak EPSCs from callosal inputs.

Bureau and colleagues demonstrated weak L4 to L2/3 synaptic strength in the global *Fmr1* KO when
measured at 2 weeks of age, but the difference was diminished at 3 weeks, suggestive of a
developmental delay. To test if loss of postsynaptic *Fmr1* weakens local L4 to L2/3 synaptic inputs at
early developmental stages, we performed LSPS on pairs of WT and postsynaptic *Fmr1* KO L2/3
pyramidal neurons at P14-17. Similar to results obtained at P23-30, we observed normal L4 and adjacent
L2/3 synaptic input strengths onto postsynaptic *Fmr1* L2/3 neurons recorded at P14-17 (Fig. S1A). We
also tested if embryonic deletion postsynaptic of *Fmr1* was necessary to affect L4 to L2/3 synapse
development, using slices from females with heterozygous (het) and mosaic expression of *Fmr1*, as we
have described (Patel *et al.*, 2013; Patel *et al.*, 2014). Briefly, X-linked GFP mice were bred with *Fmr1*
KO males. Due to random X-chromosome inactivation in the embryo, female *Fmr1* het mice offspring
have a mosaic expression of GFP(+) WT and GFP(-) *Fmr1* KO neurons. LSPS maps were performed on
pairs of WT and *Fmr1* KO neurons at P14-17 where we observed normal L4 input strengths onto L2/3
Fmr1 KO neurons (Fig. S1B). In contrast to results with postsynaptic *Fmr1* deletion, we observed weak
L4-to-L2/3 synaptic strengths with LSPS in the global *Fmr1* KO (Fig. S1C) similar to that reported by
Bureau et al. (Bureau *et al.*, 2008). However, unlike Bureau et al, we observed weak L4 inputs at P18-
25 (Fig. S1C). Taken together our results suggests that FMRP functions in presynaptic L4 neurons or in
a non-cell autonomous fashion to promote development of local L4 to L2/3 synaptic inputs. *Fmr1* in
postsynaptic L2/3 neurons selectively promotes development of callosal synaptic inputs.

Reduced synchrony of bilateral homotopic cortices in Fmr1 KO mice
To assess the consequences of reduced callosal synaptic connectivity on integration of sensory cortex
function *in vivo*, we measured the synchrony of EEG signals between bilateral homotopic cortices using
multi-electrode array (MEA) recordings in awake WT and *Fmr1* KO mice, as described (Jonak et al., 2018;
2020). Functional connectivity between hemispheres is observed by the coherence or phase-locking of
cortical oscillations. We measured the phase synchrony across hemispheres by calculating the mean
inter-site phase clustering (ISPC) of EEG signals at each frequency (3-80 Hz) from two bilateral
electrodes placed over homotopic temporal cortex (Cohen, 2014) (Fig. 7A). ISPC measures the
consistency of the phase-relationship between two signals. EEG recordings were previously acquired for
another study and analyzed here for ISPC motivated by the in vitro slice recordings (Jonak *et al.*, 2020).
Consistent with weak callosal synaptic connections, interhemispheric phase synchrony in the beta (15-
30 Hz) and gamma frequency (31-50 Hz) ranges during rest was significantly lower in *Fmr1* KO mice
compared to WT (Figure 7B, Student's t-test p_β=0.008, p_γ=0.013, respectively). To determine alterations
in bilateral functional connectivity during sensory stimulation, we calculated ISPC of the EEG during

presentation of acoustic stimuli, a brief, broad-band noise stimulus (BBN). During sound presentation, interhemispheric phase synchrony was significantly lower in *Fmr1* KO mice compared to WT (Figure 7C). As in the resting phase synchrony results, the significant differences in phase synchrony appear in the beta and gamma frequency ranges. These results are consistent with reduced callosal synaptic connectivity and reveal reduced synchrony and integration of bilateral cortical circuits in *Fmr1* KO mice.

Discussion

Reduced interhemispheric connectivity, observed both structurally and functionally, is a hallmark of ASD in humans and correlated with symptoms (Dimond *et al.*, 2019; Holiga *et al.*, 2019; Li *et al.*, 2019; O'Reilly *et al.*, 2017; Rane *et al.*, 2015; Yao *et al.*, 2021). However, little is known of the cellular, synaptic, and molecular mechanisms by which this occurs in ASD and any direct role of ASD-risk genes. Here, we observed that loss of *Fmr1,* the most common monogenetic cause of ASD, leads to weak and immature callosal synaptic inputs in global *Fmr1* KO and with postnatal, postsynaptic cell-autonomous *Fmr1* deletion in L2/3 or L5 neurons. Surprisingly, postsynaptic deletion of *Fmr1* did not weaken synaptic inputs from local columnar circuits onto L2/3 revealing that postsynaptic FMRP differentially regulates development of select synaptic inputs. Sensory deprivation of postsynaptic *Fmr1* KO neurons prevented weakening of callosal synaptic inputs suggesting that experience-driven patterned activity of postsynaptic L2/3 neurons without FMRP drives synaptic weakening and/or prevents maturation. Using MEA recordings of EEG signals, we show reduced phase synchrony between bilateral cortical circuits within the beta and gamma bands (15-50 Hz), indicative of poor integration between cortical regions. In conclusion, our results reveal a postsynaptic mechanism by which *Fmr1* regulates callosal connectivity that likely contributes to the reduced interhemispheric structural and functional connectivity in *Fmr1* KO mice and perhaps humans with FXS.

A synaptic basis for reduced functional long-range connectivity in FXS.

Functional MRI studies show that *Fmr1* KO mice have reduced corticocortical long-range connectivity, especially among the sensory and motor cortices (Haberl *et al.*, 2015; Zerbi *et al.*, 2018) as well as reduced corpus callosum structural integrity. By investigating callosal synaptic connections between bilateral barrel cortices, we show that functional synaptic inputs at these long-range connections in L2/3 are weak in *Fmr1* KO mice compared to WTs (Fig. 1). This result suggests that reduced functional coherence between bilateral barrel cortices in *Fmr1* KO mice could be due to a disrupted communication between the cortices through the weakened monosynaptic transmission. In support of this idea, we find that phase synchrony of EEG signals in bilateral temporal cortices is reduced when measured in the beta

and gamma frequencies (20-60 Hz). Interhemispheric coherence in beta-gamma band is abolished by

callosal cutting (Kiper et al., 1999; Munk et al., 1995), suggesting that the changes in interhemispheric

synchrony we observe are due to decreased callosal connectivity. Interhemispheric functional

connectivity is thought to shape perceptual integration including those involved in speech comprehension

and global form processing, domains that are impaired in ASD (Booth and Happé, 2018; Friederici et al.,

2007; Happe and Frith, 2006; Peiker et al., 2015; Preisig et al., 2021; Simon and Wallace, 2016). Based

on fMRI results in the *Fmr1* KO (Haberl *et al.*, 2015; Zerbi *et al.*, 2018), it is likely that FMRP promotes

development of long-range connectivity between ipsilateral cortical areas, such as S1 and M1/M2, as

well as cortical-subcortical structures.

Postsynaptic FMRP in L2/3 neurons promotes maturation of callosal synapses

Using *in vivo* sparse deletion and simultaneous recording paradigm, we are able to stimulate the same

set of callosal axons for a pair of WT and *Fmr1* KO neurons and directly compare their synaptic inputs.

Our observation of weaker callosal synaptic inputs onto cell-autonomous *Fmr1* KO L2/3 neurons (Fig. 2)

confirms results in *Fmr1* global KO mice and further reveals an essential role of postsynaptic FMRP in

promoting callosal synapses development. We observe a later onset of weak callosal synapses with

postsynaptic *Fmr1* KO (P23-30) in comparison to the global KO (P18-25) which may be due to the later,

postnatal, AAV-Cre mediated, deletion of *Fmr1* in postsynaptic KO condition, likely about P7-

P9(Rajkovich *et al.*, 2017). However, we cannot rule out a role for FMRP in presynaptic, callosal

projecting neurons, or other cell types within the first postnatal weeks to establish callosal synaptic

connections. Nevertheless, callosal synapses do not mature or are not maintained without FMRP in

postsynaptic L2/3 neurons. In support of a role in maturation, callosal synaptic strength tends to increase

from P18-20 to P23-30 in WT neurons but not in postsynaptic *Fmr1* KO neurons (Fig. 2E). Furthermore,

we observe a selective weakening of AMPAR, but not NMDAR-, mediated synaptic transmission at

callosal inputs onto *Fmr1* KO neurons (Fig. 3). This result, together with a decreased frequency of evoked

quantal events in Sr^{2+}, suggests a higher proportion of NMDAR-only, or silent, callosal synapses on *Fmr1*

KO neurons. Silent synapses are characteristic of immature synapses reflecting the acquisition of

NMDARs prior to AMPARs during synapse maturation (Ashby and Isaac, 2011; Hanse et al., 2013).

Thalamocortical inputs to L4, another "long-range" synaptic pathway, are delayed in their development

in the *Fmr1* KO, as measured by acquisition of AMPARs (Harlow et al., 2010). Whether the silent callosal

synapses on *Fmr1* KO neurons reflects a developmental delay or persist in the adult remains an open

question. The observations of reduced corpus callosum structural integrity and interhemispheric

coherence with fMRI and EEG (Fig. 7) in adult *Fmr1* KO suggest weak callosal connectivity in adults

(Haberl *et al.*, 2015; Zerbi *et al.*, 2018). Reduced corpus callosum integrity and axon diameter observed

in adult *Fmr1* KO may be a later consequence of the failure of maturation and maintenance of callosal
synapses. Alternatively, or in addition, FMRP may function in axons and oligodendrocytes to maintain
callosal axon size and integrity (Antar et al., 2006; Doll et al., 2020; Hanson and Madison, 2007).

Postsynaptic FMRP differentially regulates synaptic inputs from local and long-range cortical
circuits

Neocortical pyramidal neurons integrate excitatory synaptic inputs from local and long-range circuits,
including ipsilateral and contralateral cortical areas (Feldmeyer, 2012; Gerfen et al., 2018). An imbalance
in local and long-range connectivity has been hypothesized to contribute to ASD in humans; specifically,
hyperconnectivity of local circuits and underconnectivity of long-range circuits or between brain regions
(Belmonte et al., 2004; Courchesne and Pierce, 2005; O'Reilly *et al.*, 2017; Rane *et al.*, 2015). If or how
ASD genes regulate the balance of local and long-range synaptic connectivity is unknown. Here we
demonstrate that postsynaptic FMRP differentially regulates development and/or maintenance of
synaptic inputs from local and long-range cortical sources. L2/3 neurons with postsynaptic deletion of
Fmr1 have weak and immature callosal synaptic inputs but normal synaptic inputs from L4, adjacent L2/3
and L5 (Fig. 5). Bureau et al. (Bureau *et al.*, 2008) and we (Fig. S1C) find that L4 to L2/3 synapses are
weak in the global *Fmr1* KO, but we do not observe this with embryonic or postnatal cell autonomous
deletion of *Fmr1* in L2/3 neurons. Together these results suggest a role for FMRP in presynaptic L4
neurons in synapse development onto L2/3 neurons, which is consistent with the reported deficits in L4
axon morphology in the global *Fmr1* KO (Bureau *et al.*, 2008).

Results in L5 also indicate differential regulation of local and long-range cortical connectivity by
postsynaptic FMRP. Our previous work using multiple simultaneous recordings of locally connected L5A
neurons, revealed hyperconnectivity of *Fmr1* KO L5 neurons with their immediate neighbors (<40 μm
apart) in S1 at 4 weeks of age (Patel *et al.*, 2014). Hyperconnectivity of L5 local subnetworks resulted
from deficient developmental pruning between *Fmr1* KO L5 neurons and was observed in both the global
Fmr1 KO and with postsynaptic *Fmr1* deletion. Prefrontal L5 cortical neurons in *Fmr1* KO mice are
similarly hyperconnected (Testa-Silva et al., 2012). In contrast to local hyperconnectivity, here we find
that postsynaptic deletion of *Fmr1* in L5 neurons results in weak callosal synaptic inputs. Thus, *Fmr1* KO
L5 pyramidal neurons are hyperconnected locally and under connected to contralateral cortex; an effect
that is mediated by cell-autonomous and postsynaptic deletion of FMRP. Such an effect may promote an
imbalance in local and long-range connectivity observed in some ASD individuals. Postsynaptic deletion
of FMRP in L2/3 neurons did not affect synaptic inputs from other layers or between columns within L2/3
(Fig. 6) suggesting that FMRP may selectively promote pruning of local connections in L5 neurons.

Alternatively, FMRP may prune synaptic connections within very local cortical subnetworks (within 40 μm)

in both L2/3 and L5, but not connections between layers and columns.

Bidirectional regulation of callosal synaptic function by MEF2C and FMRP

Differential regulation of local and long-range cortical synaptic connectivity is observed with another ASD-

risk gene, Myocyte Enhancer Factor 2C (*Mef2c*). Loss of function mutations in *MEF2C*, which encodes

an activity-dependent transcription factor, are implicated in intellectual disability, ASD and schizophrenia

(Assali et al., 2019; Mitchell et al., 2018; Rocha et al., 2016). In contrast to FMRP, postsynaptic deletion

of *Mef2c* in L2/3 neurons results in fewer and weak inputs from local circuits (L4, L2/3 and L5), but

strengthened callosal inputs (Rajkovich *et al.*, 2017). The differential regulation of local and callosal

synaptic connections by postsynaptic MEF2C and FMRP, albeit in different directions, suggest that

imbalances in local and long-range synaptic connectivity may contribute to different genetic causes of

neurodevelopmental disorders. Our results also implicate roles for transcription and translational control

in the input-specific development of cortical circuits. Bidirectional regulation of local and long-range

connectivity by MEF2c in L2/3 or by *Fmr1* in L5 could be an effect of homeostasis of local and long-range

synaptic connectivity to maintain optimal cortical circuit function. For example, weakening of callosal

synaptic input in *Fmr1* KO neurons may be compensatory and an attempt to normalize hyperconnected

or hyperexcitable local circuits.

Interaction of sensory experience and ASD-risk genes in regulation of long-range cortical

circuits

The regulation of local callosal connectivity by FMRP and MEF2C requires normal sensory experience

suggesting that both of these genes function in experience and activity-regulated pathways necessary

for cortical circuit development. Whisker trimming (at P9-P29) normalizes callosal synaptic inputs in L2/3

neurons with postsynaptic deletion of *Fmr1* (Fig. 5) or *Mef2c* (Rajkovich *et al.*, 2017). Specifically,

trimming whiskers contralateral to L2/3 neurons with postsynaptic *Fmr1* deletion normalized callosal

synaptic input strengths as compared to WT L2/3 neighbors. Because L2/3 neurons receive the most

direct ascending inputs from contralateral whiskers via the thalamus and L4, contralateral trimming

deprives postsynaptic *Fmr1* KO neurons of most patterned, whisker- driven activity, so we termed this

"postsynaptic deprivation". In contrast, in mice with trimmed whiskers contralateral to ChR2-expressing,

callosal axon projection neurons, or "presynaptic deprivation", we observed weak callosal synapses onto

postsynaptic *Fmr1* KO neurons (Fig. 6). Callosal synaptic strengths onto WT neurons were not different

between pre- and postsynaptic deprivation conditions; whereas in *Fmr1* KO neurons callosal synapses

were weak in the postsynaptic deprivation condition as compared to presynaptic deprivation. This result

suggests that sensory-driven patterned activity of postsynaptic *Fmr1* KO L2/3 neurons weakens callosal

synapses. An intriguing possibility is that an activity-dependent long-term synaptic weakening process is

enhanced at *Fmr1* KO L2/3 neurons as observed in hippocampal CA1(Huber et al., 2002). Sensory

deprivation also induces homeostatic synaptic scaling in primary sensory cortices, as measured with

spontaneous or miniature (m) EPSCs(Feldman, 2009) (Hooks and Chen, 2020). Neither the frequency

nor amplitude of mEPSCs was different between neighboring WT and *Fmr1* KO L2/3 neurons in mice

with normal sensory experience or "presynaptic deprivation". However, with "postsynaptic deprivation",

mEPSC amplitude and frequency was increased in *Fmr1* KO neurons relative to WT (Fig. S2) suggestive

of enhanced homeostatic up scaling in *Fmr1* KO neurons that may contribute to callosal input

strengthening in this condition.

Bilateral underconnectivity is common in ASD as well as disconnection of other long-range cortical

connections with other brain regions such as hippocampus and cerebellum. Bilateral connectivity and

gamma synchrony between cortical regions is necessary for speech comprehension, sensory processing

and cognition (Bland et al., 2020; Castro et al., 2014; Friederici *et al.*, 2007; Fries, 2009; 2015; Panzica

et al., 2019) and it is likely that reduced connectivity and synchrony between regions contributes to

dysfunction in these domains in FXS and ASD. Our present findings contribute to the understanding of

the cellular and synaptic mechanisms by which ASD-risk genes, such as *FMR1,* regulate long-range

connectivity, how this is coregulated and balanced with local circuit connectivity and interacts with

experience-dependent brain development. Such information may contribute to therapies to aid abnormal

brain connectivity in ASD.

Materials and Methods

Animals

Fmr1 KO (*Fmr1$^{-/y}$*) and X-linked GFP mice were obtained from Jackson laboratories (Stock No: 003025 and 003116, respectively). *Fmr1$^{fl/fl}$* were obtained from Dr. David Nelson (Baylor College of Medicine) (Mientjes *et al.*, 2006). Mice were maintained on a C57BL/6J background and reared on a 12 hr light-dark cycle with access to food and water ad libitum. Male *Fmr1* WT (*Fmr1$^{+/y}$*) with *Fmr1* KO (*Fmr1$^{-/y}$*) littermates were used for experiments and both male and female pups were used for *Fmr1* flox (*Fmr1$^{fl/y}$* and *Fmr1$^{fl/fl}$*). All animal experiments were conducted in accordance with the Institutional Animal Care and Use Committee (IACUC) at University of Texas Southwestern Medical Center.

Viral transfections in neonatal mice

Commercially made AAV9.CMV.HI.eGFP-Cre.WPRE.SV40 (Addgene #105545) and AAV9.CAG.hChR2(H134R)-mCherry.WPRE.SV40 (Addgene #100054) were diluted to a titer ~10^{12} vg/mL using sterile saline. Traces of Fast Green FCF dye (Sigma) were added to facilitate visualization of virus spreading. Neonatal mouse pups (P1) were first anesthetized by hypothermia, then fixed on a customized mold and placed on a stereotaxic frame. AAV9.eGFP-Cre (420-560 nL) was delivered to left ventricle at a depth of approximately 1.1 mm underneath skull through a beveled glass pipette using Nanoject II injector (Drummond Scientific, Inc.). AAV9.ChR2-mCherry (400 nL) was delivered to superficial layers (0.5 mm underneath skull) of somatosensory cortex in right hemisphere at a speed of 1.2 µL/min using syringe pump (Harvard Apparatus, Inc.).

Acute slice preparation

Mice were anesthetized by intraperitoneal (i.p.) injection of Ketamine/Xylazine mixture and decapitated upon irresponsiveness to toe-pinch. Acute coronal slices (300 µm thickness) containing somatosensory barrel cortex were prepared using vibrating microtome (Leica VT1200S). During sectioning, tissue blocks were submerged in ice-cold dissection buffer containing (in mM): 110 choline chloride, 25 NaHCO$_3$, 25 dextrose, 11.6 ascorbic acid, 2.5 KCl, 1.25 NaH$_2$PO$_4$, 3.1 Na-pyruvate, 7 MgCl$_2$ and 0.5 CaCl$_2$, continuously aerated with 95%CO$_2$/5%O$_2$. For mice older than P23, transcardial perfusion of ice-cold dissection buffer was performed prior to decapitation to increase slice quality. Slices were then transferred to artificial cerebrospinal fluid (ACSF) solution containing (in mM): 125 NaCl, 25 NaHCO$_3$, 10 dextrose, 2.5 KCl, 1.25 NaH$_2$PO$_4$, 2 MgCl$_2$ and 2 CaCl$_2$ (aerated with 95%CO$_2$/5%O2) and recovered at 34°C for 30 min followed by 30 min at room temperature.

Electrophysiology

After recovery, slices were transferred to a recording chamber at room temperature and perfused with ACSF aerated with 95%CO_2/5%O_2. Slices were visualized by infrared differential interference contrast (IR-DIC) optics (Olympus BX51W1). Whole cell recordings of L2/3 and L5 pyramidal neurons were obtained using borosilicate pipettes (4-7 MΩ) and a Multiclamp 700A amplifier (Molecular Devices). Internal solution contained (in mM): 130 K-gluconate, 10 HEPES, 6 KCl, 3 NaCl, 0.2 EGTA, 14 phosphocreatine-tris, 4 Mg-ATP and 0.4 Na-GTP. All recordings were conducted in voltage clamp, holding at -70 mV unless otherwise specified, and data were collected and analyzed using custom Labview programs (Labview 8.6, National Instruments Inc.). Spiking patterns upon current injection were used as criteria to identify excitatory neurons. For experiments where spiking was blocked by TTX, rise time to hyperpolarizing current (> 50 ms) and kinetics of mPSC (width at half height > 2 ms) were used as criteria (Povysheva et al., 2006). Excitatory neurons with resting membrane potential < -50 mV and a series resistance < 35 MΩ were included in analysis.

Optogenetic bulk stimulation of callosal axons

Mice with either cortical injection of AAV9.ChR2-mCherry or unilateral ventricular injection of AAV9.eGFP-Cre and contralateral cortical injection of AAV9.ChR2-mCherry were used for this experiment. Fluorescence of GFP-positive soma and mCherry-labeled axons were visualized using a fluorescent mercury lamp (Excelitas Technologies Corp.). Slices containing clearly labeled mCherry+ axons from contralateral barrel cortex were used for recording. Slices with somatic infection of ChR2-mCherry due to virus leaked from contralateral hemisphere were discarded to avoid contamination from local inputs. The infection rate of recorded Cre-GFP+ neurons in barrel cortex was < 10%. Only neurons residing in an area with densely labeled callosal axons were subject to recording to achieve a reliable magnitude and reduced variability of light-induced responses. For all LED experiments, responses were evoked by a 2 ms flash from a digitally controlled blue LED (final beam diameter: 350 μm; power: 0.1-4.6 mW; wavelength: 470 nm; M470L4-C1, Thorlabs Inc.) through a 40X water-immersed objective. The LED flash was centered on soma and proximal apical dendrites of recorded neurons.

To measure LED-evoked EPSCs, each cell (or cell pair) was stimulated 3-10 times with 20-30 s intervals. LED power was adjusted to obtain an EPSC amplitude of 100-1000pA in WT neurons (for cell pairs). The external solution for both LED and sCRACM experiments contained ACSF with 1 μM TTX, 100 μM 4-aminopyridine (4-AP), 10 μM (±)-3-(2-carboxypiperazin-4-yl)propyl-1-phosphonic acid (CPP) and 100 μM picrotoxin to isolate monosynaptic AMPAR-mediated excitatory inputs. Callosal input strength was measured as the peak amplitude of an average EPSC of 3 - 10 evoked EPSCs for each neuron.

To measure Sr^{2+} evoked quantal events, each cell pair was stimulated an LED flash every 30s, 12-30

times in ACSF containing 4 mM $MgCl_2$, 4 mM $SrCl_2$, 10 µM CPP and 100 µM picrotoxin. Events occurring

within a 1 s window prior to the LED were defined as spontaneous events and those occurring 50-350

ms post-LED were defined LED-evoked events. The frequency and amplitude of events were analyzed

using MiniAnalysis (Synaptosoft).

NMDAR-mediated EPSCs from callosal inputs were pharmacologically isolated in ACSF containing: 3

mM $CaCl_2$, 0.1 mM $MgCl_2$, 20 µM DNQX, 20 µM glycine, 100 µM picrotoxin, 1 µM TTX, 100 µM 4-AP.

An LED flash was delivered every 30s, 9 times. Evoked EPSCs were averaged and peak amplitude

measured.

Subcellular Channelrhodopsin-assisted Circuit Mapping (sCRACM) and action potential dependent activation of ChR2

After collection of LED-evoked responses, the objective was switched to 4X to visualize a broader area.

ChR2 expressing axons were then stimulated by a blue laser (1 ms; wavelength: 473 nm; power range:

0.7 - 16 mW; final beam diameter: 25 µm; CrystaLaser) scanning through a 12 x 12 grid (50 µm spacing)

in a pseudorandom order to avoid repeated activation of neighboring locations. The grid was aligned

along the pia and centered in the medial-lateral position on the recorded somas. The grid was repeatedly

scanned 2 - 4 times at 40 s intervals. Laser-evoked EPSCs were collected from each grid spot.

For action potential-dependent callosal synaptic strength measurements, an independent cohort of mice

was used. The recording ACSF was similar to sCRACM experiments, except TTX and 4-AP were omitted,

and divalent cations were increased (4 mM $MgCl_2$; 3 mM $CaCl_2$) and CPP was added to reduce

polysynaptic activation of local circuits. For L2/3 neurons, an 8 x 8 grid with 75 × 100µm x-y spacing. For

L5 neurons, an 8 x 8 grid with 75 × 125µm x-y spacing.

Analysis of blue laser evoked EPSCs

For sCRACM, EPSCs evoked from 2-4 laser-stimulation at each grid spot from an individual neuron were

averaged. The peak amplitude of the average EPSC (between 5-80 ms after laser onset) was determined

as the input strength for that spot. A spatial map of input strengths was then generated for each individual

neuron. The spatial maps for all neurons of each genotype were then aligned to soma location, oriented

with respect to the pial surface, and averaged to generate an average spatial map of input strengths for

each genotype aligned to the soma. A color representation of the average spatial map for each genotype

were generated for the figures. Pixel size for the genotype-averaged color maps was halved through pixel

interpolation (25x25 µm) to provide better spatial resolution for soma alignment. Vertical profiling of input

strength was achieved by averaging inputs from each horizontal row and plotting against the vertical distance from soma. Input strength from a specific area was calculated by averaging input strengths within an area for each neuron and averaging according to genotype.

For action-potential-dependent ChR2 activation and circuit mapping, analysis was performed as for sCRACM, except EPSC amplitudes were averaged during 2-30 ms after laser onset to exclude contamination from polysynaptic responses.

Laser Scanning Photostimulation (LSPS) with glutamate uncaging

LSPS experiments were performed similar to that described previously (Rajkovich *et al.*, 2017; Shepherd et al., 2003). For all experiments, only brain slices with L2/3 apical dendrites parallel to the slice surface were used to ensure preservation of the planar barrel cortical geometry of long-range synaptic pathways spanning at least 3 barrel columns. Usually 2-4 brain slices per animal met such criteria. ACSF included 4 mM $MgCl_2$, 4 mM $CaCl_2$, and CPP (10 µM) to reduced polysynaptic local circuit activity (Rajkovich *et al.*, 2017; Shepherd *et al.*, 2003), and 4-Methoxy-7-nitroindolinyl-caged-L-glutamate, MNI glutamate (MNI, 0.3 mM, either Tocris-1490 or HelloBio-HB0423). A 1 ms UV laser flash (wavelength: 355 nm; power range: 30 - 40 mW; final beam diameter: 20 µm; DPSS Lasers Inc.) was delivered at individual points within a 16 x 16 grid (50 × 60µm x-y spacing) in a pseudorandom order. The grid was aligned along the pia surface and centered medial-laterally on the soma location. The entire grid was repeatedly scanned 2 - 4 times at 40 s intervals. 2-4 maps were acquired for each neuron included in all datasets.

For each neuron, a single average map was calculated from acquired LSPS maps, where at each stimulation point the averaged light evoked EPSC area was calculated within a time window of 5-80 ms following the laser pulse. If a response was observed within 5 ms of LSPS and displayed kinetics visibly faster than the longer-latency EPSC then it was considered to be a non-synaptic, 'direct' response to uncaging of glutamate on the recorded neuron. Direct responses were removed from the map and not included in analysis. For responses with a major component of monosynaptic transmission and minor contamination from 'direct' activation, 'direct' response component was subtracted from the whole response by fitting a double-exponential decay equation and subtracted from the trace obtained at all stimulation points at which they were observed. An IR-DIC image of the slice with patch pipettes in place and stimulation grid was acquired prior to LSPS for marking soma location and anatomical features of the slice (i.e. barrels, etc.). Finally, a color map for each neuron was created. All individual color maps within genotype were then overlaid upon spatial alignment with respect to the center of the "home" barrel directly beneath recorded L2/3 neurons. Superimposition of the average maps was achieved by 1) transposing each map such that the home barrel center was located at the origin of alignment grid, 2)

569   preserving the medial-lateral orientation of the brain slice, and 3) stretching the home barrel in x and y

570   dimensions to normalize barrel size. Pixel size for the genotype-averaged color maps was halved through

571   pixel interpolation (25 x 30 μm) to provide better spatial resolution for soma alignment. Black pixels within

572   an averaged color map in the figures indicate deleted direct responses or pixels that did not meet the

573   minimum sampling threshold (minimum of n = 8 neurons per stimulation point).

Whisker trimming

576   Unilateral whisker trimming was performed daily on mice starting at P15 until the day before experiment

577   (P23-30) using a miniature electric shaver. All whiskers from the trimmed pad were maintained at a length

578   < 1mm. For a litter of mice, half of the mice would be trimmed on the right whisker pad, ipsilateral to the

579   AAV9.ChR2-mCherry injection (postsynaptic deprivation), while the other half would be trimmed on the

580   left whisker pad, contralateral to the AAV9.ChR2-mCherry injection (presynaptic deprivation).

Statistics

All the statistical tests and graphs were performed using GraphPad Prism 8 (GraphPad Software Inc.).

Prior to statistical tests of significance, each dataset underwent normality tests (D'Agostino-Pearson and

Anderson-Darling) to determine if parametric or non-parametric tests should be used. All statistical tests

are two-sided. Only datasets tested to be normally distributed by both tests were subject to parametric

statistics – unpaired or paired t-test. Otherwise, Mann-Whitney test or Wilcoxon matched pairs test were

used as indicated in the figure legends. For data collected from double-patch experiments, paired

statistics were used as each pair of neurons would be considered correlated. For comparing more than

two groups of data, 2-way ANOVA or mixed effect analysis with multiple comparisons was used.

Repeated measures ANOVA was used for vertical profile maps in Figures 1 and 2.

Multielectrode Array Recordings

MEA recordings were performed as previously described (Jonak *et al.*, 2018) and the recordings collected

in this former study were analyzed here (Jonak *et al.*, 2020). Briefly, for MEA studies, *Fmr1* KO

(B6.129P2-Fmr1tm1Cgr/J, stock #003025) and C57BL/6J WT (stock #000664) mice were obtained from

Jackson Laboratories at 8 weeks of age and housed prior to surgery and recording. MEAs were implanted

onto the skull of 10 WT and 11 *Fmr1* KO male mice between 2 and 3 months of age. Recordings were

performed from awake mice 2- 4 days after surgery.

Phase Synchrony Analysis

Data analyses were done using a combination of Analyzer 2.1 (BrainVision Inc.) and MATLAB. Data were extracted from the Smartbox files and saved in a file format compatible with Analyzer 2.1 software. Data were first down sampled to 625 Hz and EEG artifacts were removed using a semi-automatic procedure in Analyzer 2.1 for all recordings. Less than 20% of data were rejected due to artifacts from any single mouse. We applied a complex dynamic Morlet wavelet transformation to the resting (no stimulus) and broadband noise (BBN) stimulus EEG recordings.

Resting:

We measured the phase synchrony across the temporal electrodes at rest by calculating the mean inter-site phase clustering (ISPC) between electrodes 7 and 24 (Figure 7A) at each frequency from 3-80 Hz, according to the formula below. ISPC measures the consistency of the phase relationship between two electrodes across time. We used Student's t-test to compare ISPC values for WT and *Fmr1* KO mice in each frequency range.

$$\text{ISPC}_f = \left| n^{-1} \sum_{t=1}^{n} e^{i(\phi_{xt} - \phi_{yt})} \right|$$ (Cohen, 2014).

Broad-band noise (BBN) Stimulus:

Phase synchrony was measured across the 2 temporal electrodes (Figure 7A) in relation to auditory stimulus presentation. Acoustic stimulation is described in (Jonak *et al.*, 2020). Briefly, trains of broadband noise with each train consisting of 10 repetitions (1 Hz repetition rate). Each noise stimulus was 100 ms in duration, with a 5 ms rise/fall time and was presented at 70 dB SPL. The inter-train interval was 8 s. The total duration of broadband stimuli was 30 min. Phase synchrony was measured during the first noise stimulus in the train and averaged across 10 repetitions for each mouse. In contrast to resting ISPC, which was calculated across continuous time, stimulus-related ISPC was calculated across trials. This measures how consistent the phase-relationship was between the two electrodes at each time-frequency point in the trial window. We used a nonparametric permutation clustering method to test for significant differences between the groups (Cohen, 2014; Maris and Oostenveld, 2007). A t-test was run on each time-frequency point. The resulting p-values <0.025 were considered to be significant. We then measured the area of all clusters of adjacent significant p-values. The group assignments were randomly shuffled 2000 times, and significant clusters were measured for each shuffle. This yielded a distribution of cluster sizes expected by chance. Clusters in the original comparison that were larger than 95% of the surrogate clusters were considered to be significant. This method allows for the discovery of significant

differences between groups without performing an unwieldy number of multiple comparisons and without

making assumptions about normality or homogeneity of variance between groups.

Acknowledgements

We thank Patricia Hahn, Jacob Eli Bowles, and Christopher Williams for technical assistance with the

mice. JRG, KMH, ZZ, DKB and KAR conceived various aspects of the project. ZZ, JRG and CRJ

performed experiments and analyzed data. JR performed the ISPC analysis. ZZ and KMH wrote the

manuscript with input from all authors. This work was supported by grants from the NIH (R01 HD052731

(KMH, JRG), U54HD082008 and U54HD104461 (KMH, JRG, DKB and KAR)).

The authors declare no competing interests.

References

Antar, L.N., Li, C., Zhang, H., Carroll, R.C., and Bassell, G.J. (2006). Local functions for FMRP in axon
growth cone motility and activity-dependent regulation of filopodia and spine synapses. Mol Cell Neurosci
32, 37-48. 10.1016/j.mcn.2006.02.001.
Ashby, M.C., and Isaac, J.T. (2011). Maturation of a recurrent excitatory neocortical circuit by experience-
dependent unsilencing of newly formed dendritic spines. Neuron *70*, 510-521.
10.1016/j.neuron.2011.02.057.
Assali, A., Harrington, A.J., and Cowan, C.W. (2019). Emerging roles for MEF2 in brain development and
mental disorders. Current Opinion in Neurobiology *59*, 49-58. 10.1016/j.conb.2019.04.008.
Belmonte, M.K., Allen, G., Beckel-Mitchener, A., Boulanger, L.M., Carper, R.A., and Webb, S.J. (2004).
Autism and abnormal development of brain connectivity. J Neurosci *24*, 9228-9231.
10.1523/jneurosci.3340-04.2004.
Bland, N.S., Mattingley, J.B., and Sale, M.V. (2020). Gamma coherence mediates interhemispheric
integration during multiple object tracking. J Neurophysiol *123*, 1630-1644. 10.1152/jn.00755.2019.
Booth, R.D.L., and Happé, F.G.E. (2018). Evidence of Reduced Global Processing in Autism Spectrum
Disorder. J Autism Dev Disord *48*, 1397-1408. 10.1007/s10803-016-2724-6.
Bureau, I., Shepherd, G.M., and Svoboda, K. (2008). Circuit and plasticity defects in the developing
somatosensory cortex of FMR1 knock-out mice. J Neurosci *28*, 5178-5188. 10.1523/JNEUROSCI.1076-
08.2008.
Castro, S., Cavelli, M., Vollono, P., Chase, M.H., Falconi, A., and Torterolo, P. (2014). Inter-hemispheric
coherence of neocortical gamma oscillations during sleep and wakefulness. Neurosci Lett *578*, 197-202.
10.1016/j.neulet.2014.06.044.
Ciarrusta, J., Dimitrova, R., Batalle, D., O'Muircheartaigh, J., Cordero-Grande, L., Price, A., Hughes, E.,
Kangas, J., Perry, E., Javed, A., et al. (2020). Emerging functional connectivity differences in newborn
infants vulnerable to autism spectrum disorders. Transl Psychiatry *10*, 131. 10.1038/s41398-020-0805-
y.
Cohen, M.X. (2014). Analyzing Neural Time Series Data: Theory and Practice (The MIT Press).
https://doi.org/10.7551/mitpress/9609.001.0001.
Contractor, A., Klyachko, V.A., and Portera-Cailliau, C. (2015). Altered Neuronal and Circuit Excitability
in Fragile X Syndrome. Neuron *87*, 699-715. 10.1016/j.neuron.2015.06.017.

Courchesne, E., and Pierce, K. (2005). Why the frontal cortex in autism might be talking only to itself: local over-connectivity but long-distance disconnection. Curr Opin Neurobiol *15*, 225-230. 10.1016/j.conb.2005.03.001.

Cruz-Martin, A., Crespo, M., and Portera-Cailliau, C. (2010). Delayed stabilization of dendritic spines in fragile X mice. J Neurosci *30*, 7793-7803. 10.1523/JNEUROSCI.0577-10.2010.

Darnell, J.C., Van Driesche, S.J., Zhang, C., Hung, K.Y., Mele, A., Fraser, C.E., Stone, E.F., Chen, C., Fak, J.J., Chi, S.W., et al. (2011). FMRP stalls ribosomal translocation on mRNAs linked to synaptic function and autism. Cell *146*, 247-261. 10.1016/j.cell.2011.06.013.

Dimond, D., Schuetze, M., Smith, R.E., Dhollander, T., Cho, I., Vinette, S., Ten Eycke, K., Lebel, C., McCrimmon, A., Dewey, D., et al. (2019). Reduced White Matter Fiber Density in Autism Spectrum Disorder. Cereb Cortex *29*, 1778-1788. 10.1093/cercor/bhy348.

Doll, C.A., Yergert, K.M., and Appel, B.H. (2020). The RNA binding protein fragile X mental retardation protein promotes myelin sheath growth. Glia *68*, 495-508. 10.1002/glia.23731.

Feldman, D.E. (2009). Synaptic mechanisms for plasticity in neocortex. Annu Rev Neurosci *32*, 33-55. 10.1146/annurev.neuro.051508.135516.

Feldmeyer, D. (2012). Excitatory neuronal connectivity in the barrel cortex. Front Neuroanat *6*, 24. 10.3389/fnana.2012.00024.

Friederici, A.D., von Cramon, D.Y., and Kotz, S.A. (2007). Role of the corpus callosum in speech comprehension: interfacing syntax and prosody. Neuron *53*, 135-145. 10.1016/j.neuron.2006.11.020.

Fries, P. (2009). Neuronal gamma-band synchronization as a fundamental process in cortical computation. Annu Rev Neurosci *32*, 209-224. 10.1146/annurev.neuro.051508.135603.

Fries, P. (2015). Rhythms for Cognition: Communication through Coherence. Neuron *88*, 220-235. 10.1016/j.neuron.2015.09.034.

Garber, K.B., Visootsak, J., and Warren, S.T. (2008). Fragile X syndrome. Eur J Hum Genet *16*, 666-672. 10.1038/ejhg.2008.61.

Gerfen, C.R., Economo, M.N., and Chandrashekar, J. (2018). Long distance projections of cortical pyramidal neurons. J Neurosci Res *96*, 1467-1475. 10.1002/jnr.23978.

Gibson, J.R., Bartley, A.F., Hays, S.A., and Huber, K.M. (2008). Imbalance of neocortical excitation and inhibition and altered UP states reflect network hyperexcitability in the mouse model of fragile X syndrome. J Neurophysiol *100*, 2615-2626. 10.1152/jn.90752.2008.

Goel, A., Cantu, D.A., Guilfoyle, J., Chaudhari, G.R., Newadkar, A., Todisco, B., de Alba, D., Kourdougli, N., Schmitt, L.M., Pedapati, E., et al. (2018). Impaired perceptual learning in a mouse model of Fragile X syndrome is mediated by parvalbumin neuron dysfunction and is reversible. Nat Neurosci *21*, 1404-1411. 10.1038/s41593-018-0231-0.

Goncalves, J.T., Anstey, J.E., Golshani, P., and Portera-Cailliau, C. (2013). Circuit level defects in the developing neocortex of Fragile X mice. Nat Neurosci *16*, 903-909. 10.1038/nn.3415.

Haberl, M.G., Zerbi, V., Veltien, A., Ginger, M., Heerschap, A., and Frick, A. (2015). Structural-functional connectivity deficits of neocortical circuits in the Fmr1 (-/y) mouse model of autism. Sci Adv *1*, e1500775. 10.1126/sciadv.1500775.

Hanse, E., Seth, H., and Riebe, I. (2013). AMPA-silent synapses in brain development and pathology. Nat Rev Neurosci *14*, 839-850. 10.1038/nrn3642.

Hanson, J.E., and Madison, D.V. (2007). Presynaptic FMR1 genotype influences the degree of synaptic connectivity in a mosaic mouse model of fragile X syndrome. J Neurosci *27*, 4014-4018. 10.1523/JNEUROSCI.4717-06.2007.

Happe, F., and Frith, U. (2006). The weak coherence account: detail-focused cognitive style in autism spectrum disorders. J Autism Dev Disord *36*, 5-25. 10.1007/s10803-005-0039-0.

Harlow, E.G., Till, S.M., Russell, T.A., Wijetunge, L.S., Kind, P., and Contractor, A. (2010). Critical period plasticity is disrupted in the barrel cortex of FMR1 knockout mice. Neuron *65*, 385-398. 10.1016/j.neuron.2010.01.024.

Hays, S.A., Huber, K.M., and Gibson, J.R. (2011). Altered neocortical rhythmic activity states in Fmr1 KO
mice are due to enhanced mGluR5 signaling and involve changes in excitatory circuitry. J Neurosci *31*,
14223-14234. 10.1523/JNEUROSCI.3157-11.2011.
He, C.X., and Portera-Cailliau, C. (2013). The trouble with spines in fragile X syndrome: density, maturity
and plasticity. Neuroscience *251*, 120-128. 10.1016/j.neuroscience.2012.03.049.
He, Q., Nomura, T., Xu, J., and Contractor, A. (2014). The developmental switch in GABA polarity is
delayed in fragile X mice. J Neurosci *34*, 446-450. 10.1523/JNEUROSCI.4447-13.2014.
Holiga, S., Hipp, J.F., Chatham, C.H., Garces, P., Spooren, W., D'Ardhuy, X.L., Bertolino, A., Bouquet,
C., Buitelaar, J.K., Bours, C., et al. (2019). Patients with autism spectrum disorders display reproducible
functional connectivity alterations. Sci Transl Med *11*. 10.1126/scitranslmed.aat9223.
Hooks, B.M., and Chen, C. (2020). Circuitry Underlying Experience-Dependent Plasticity in the Mouse
Visual System. Neuron *106*, 21-36. 10.1016/j.neuron.2020.01.031.
Huang, Y., Song, N.N., Lan, W., Zhang, Q., Zhang, L., Zhang, L., Hu, L., Chen, J.Y., Zhao, C.J., Li, L., et
al. (2013). Sensory input is required for callosal axon targeting in the somatosensory cortex. Mol Brain *6*,
53. 10.1186/1756-6606-6-53.
Huber, K.M., Gallagher, S.M., Warren, S.T., and Bear, M.F. (2002). Altered synaptic plasticity in a mouse
model of fragile X mental retardation. Proc Natl Acad Sci U S A *99*, 7746-7750. 10.1073/pnas.122205699.
Jonak, C.R., Lovelace, J.W., Ethell, I.M., Razak, K.A., and Binder, D.K. (2018). Reusable Multielectrode
Array Technique for Electroencephalography in Awake Freely Moving Mice. Front Integr Neurosci *12*, 53.
10.3389/fnint.2018.00053.
Jonak, C.R., Lovelace, J.W., Ethell, I.M., Razak, K.A., and Binder, D.K. (2020). Multielectrode array
analysis of EEG biomarkers in a mouse model of Fragile X Syndrome. Neurobiol Dis *138*, 104794.
10.1016/j.nbd.2020.104794.
Kim, J.Y., Ash, R.T., Ceballos-Diaz, C., Levites, Y., Golde, T.E., Smirnakis, S.M., and Jankowsky, J.L.
(2013). Viral transduction of the neonatal brain delivers controllable genetic mosaicism for visualising and
manipulating neuronal circuits in vivo. Eur J Neurosci *37*, 1203-1220. 10.1111/ejn.12126.
Kiper, D.C., Knyazeva, M.G., Tettoni, L., and Innocenti, G.M. (1999). Visual stimulus-dependent changes
in interhemispheric EEG coherence in ferrets. J Neurophysiol *82*, 3082-3094. 10.1152/jn.1999.82.6.3082.
Li, Q., Becker, B., Jiang, X., Zhao, Z., Zhang, Q., Yao, S., and Kendrick, K.M. (2019). Decreased
interhemispheric functional connectivity rather than corpus callosum volume as a potential biomarker for
autism spectrum disorder. Cortex *119*, 258-266. 10.1016/j.cortex.2019.05.003.
Lovelace, J.W., Ethell, I.M., Binder, D.K., and Razak, K.A. (2018). Translation-relevant EEG phenotypes
in a mouse model of Fragile X Syndrome. Neurobiol Dis *115*, 39-48. 10.1016/j.nbd.2018.03.012.
Maris, E., and Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data. J
Neurosci Methods *164*, 177-190. 10.1016/j.jneumeth.2007.03.024.
Mientjes, E.J., Nieuwenhuizen, I., Kirkpatrick, L., Zu, T., Hoogeveen-Westerveld, M., Severijnen, L., Rife,
M., Willemsen, R., Nelson, D.L., and Oostra, B.A. (2006). The generation of a conditional Fmr1 knock
out mouse model to study Fmrp function in vivo. Neurobiol Dis *21*, 549-555.
Mitchell, A.C., Javidfar, B., Pothula, V., Ibi, D., Shen, E.Y., Peter, C.J., Bicks, L.K., Fehr, T., Jiang, Y.,
Brennand, K.J., et al. (2018). MEF2C transcription factor is associated with the genetic and epigenetic
risk architecture of schizophrenia and improves cognition in mice. Mol. Psychiatry *23*, 123-132.
10.1038/mp.2016.254.
Mizuno, H., Hirano, T., and Tagawa, Y. (2010). Pre-synaptic and post-synaptic neuronal activity supports
the axon development of callosal projection neurons during different post-natal periods in the mouse
cerebral cortex. Eur J Neurosci *31*, 410-424. 10.1111/j.1460-9568.2009.07070.x.
Munk, M.H., Nowak, L.G., Nelson, J.I., and Bullier, J. (1995). Structural basis of cortical synchronization.
II. Effects of cortical lesions. J Neurophysiol *74*, 2401-2414. 10.1152/jn.1995.74.6.2401.
Nimchinsky, E.A., Oberlander, A.M., and Svoboda, K. (2001). Abnormal development of dendritic spines
in FMR1 knock-out mice. J Neurosci *21*, 5139-5146.
Niu, M., Han, Y., Dy, A.B.C., Du, J., Jin, H., Qin, J., Zhang, J., Li, Q., and Hagerman, R.J. (2017). Autism
Symptoms in Fragile X Syndrome. J Child Neurol *32*, 903-909. 10.1177/0883073817712875.

O'Reilly, C., Lewis, J.D., and Elsabbagh, M. (2017). Is functional brain connectivity atypical in autism? A
systematic review of EEG and MEG studies. PLoS One *12*, e0175870. 10.1371/journal.pone.0175870.
Oliet, S.H., Malenka, R.C., and Nicoll, R.A. (1996). Bidirectional control of quantal size by synaptic activity
in the hippocampus. Science *271*, 1294-1297.
Osterweil, E.K., Chuang, S.C., Chubykin, A.A., Sidorov, M., Bianchi, R., Wong, R.K., and Bear, M.F.
(2013). Lovastatin corrects excess protein synthesis and prevents epileptogenesis in a mouse model of
fragile X syndrome. Neuron *77*, 243-250. 10.1016/j.neuron.2012.01.034.
Panzica, F., Schiaffi, E., Visani, E., Franceschetti, S., and Giovagnoli, A.R. (2019). Gamma
electroencephalographic coherence and theory of mind in healthy subjects. Epilepsy Behav *100*, 106435.
10.1016/j.yebeh.2019.07.036.
Patel, A.B., Hays, S.A., Bureau, I., Huber, K.M., and Gibson, J.R. (2013). A target cell-specific role for
presynaptic Fmr1 in regulating glutamate release onto neocortical fast-spiking inhibitory neurons. J
Neurosci *33*, 2593-2604. 10.1523/JNEUROSCI.2447-12.2013.
Patel, A.B., Loerwald, K.W., Huber, K.M., and Gibson, J.R. (2014). Postsynaptic FMRP promotes the
pruning of cell-to-cell connections among pyramidal neurons in the L5A neocortical network. J Neurosci
34, 3413-3418. 10.1523/JNEUROSCI.2921-13.2014.
Peiker, I., David, N., Schneider, T.R., Nolte, G., Schöttle, D., and Engel, A.K. (2015). Perceptual
Integration Deficits in Autism Spectrum Disorders Are Associated with Reduced Interhemispheric
Gamma-Band Coherence. J Neurosci *35*, 16352-16361. 10.1523/jneurosci.1442-15.2015.
Petreanu, L., Huber, D., Sobczyk, A., and Svoboda, K. (2007). Channelrhodopsin-2-assisted circuit
mapping of long-range callosal projections. Nat Neurosci *10*, 663-668. 10.1038/nn1891.
Petreanu, L., Mao, T., Sternson, S.M., and Svoboda, K. (2009). The subcellular organization of
neocortical excitatory connections. Nature *457*, 1142-1145. 10.1038/nature07709.
Pfeiffer, B.E., and Huber, K.M. (2009). The state of synapses in fragile X syndrome. Neuroscientist *15*,
549-567. 10.1177/1073858409333075.
Pfeiffer, B.E., Zang, T., Wilkerson, J.R., Taniguchi, M., Maksimova, M.A., Smith, L.N., Cowan, C.W., and
Huber, K.M. (2010). Fragile X mental retardation protein is required for synapse elimination by the activity-
dependent transcription factor MEF2. Neuron *66*, 191-197. 10.1016/j.neuron.2010.03.017.
Povysheva, N.V., Gonzalez-Burgos, G., Zaitsev, A.V., Kroner, S., Barrionuevo, G., Lewis, D.A., and
Krimer, L.S. (2006). Properties of excitatory synaptic responses in fast-spiking interneurons and
pyramidal cells from monkey and rat prefrontal cortex. Cereb Cortex *16*, 541-552. 10.1093/cercor/bhj002.
Preisig, B.C., Riecke, L., Sjerps, M.J., Kösem, A., Kop, B.R., Bramson, B., Hagoort, P., and Hervais-
Adelman, A. (2021). Selective modulation of interhemispheric connectivity by transcranial alternating
current stimulation influences binaural integration. Proc Natl Acad Sci U S A *118*.
10.1073/pnas.2015488118.
Rajkovich, K.E., Loerwald, K.W., Hale, C.F., Hess, C.T., Gibson, J.R., and Huber, K.M. (2017).
Experience-Dependent and Differential Regulation of Local and Long-Range Excitatory Neocortical
Circuits by Postsynaptic Mef2c. Neuron *93*, 48-56. 10.1016/j.neuron.2016.11.022.
Rane, P., Cochran, D., Hodge, S.M., Haselgrove, C., Kennedy, D.N., and Frazier, J.A. (2015).
Connectivity in Autism: A Review of MRI Connectivity Studies. Harv Rev Psychiatry *23*, 223-244.
10.1097/HRP.0000000000000072.
Rocha, H., Sampaio, M., Rocha, R., Fernandes, S., and Leao, M. (2016). MEF2C haploinsufficiency
syndrome: Report of a new MEF2C mutation and review. Eur J Med Genet *59*, 478-482.
10.1016/j.ejmg.2016.05.017.
Schulte, T., and Müller-Oehring, E.M. (2010). Contribution of callosal connections to the interhemispheric
integration of visuomotor and cognitive processes. Neuropsychol Rev *20*, 174-190. 10.1007/s11065-010-
9130-1.
Shepherd, G.M., Pologruto, T.A., and Svoboda, K. (2003). Circuit analysis of experience-dependent
plasticity in the developing rat barrel cortex. Neuron *38*, 277-289. 10.1016/s0896-6273(03)00152-1.
Simon, D.M., and Wallace, M.T. (2016). Dysfunction of sensory oscillations in Autism Spectrum Disorder.
Neurosci Biobehav Rev *68*, 848-861. 10.1016/j.neubiorev.2016.07.016.

Swanson, M.R., Wolff, J.J., Shen, M.D., Styner, M., Estes, A., Gerig, G., McKinstry, R.C., Botteron, K.N.,
Piven, J., Hazlett, H.C., and Infant Brain Imaging Study, N. (2018). Development of White Matter Circuitry
in Infants With Fragile X Syndrome. JAMA Psychiatry *75*, 505-513. 10.1001/jamapsychiatry.2018.0180.
Testa-Silva, G., Loebel, A., Giugliano, M., de Kock, C.P., Mansvelder, H.D., and Meredith, R.M. (2012).
Hyperconnectivity and slow synapses during early development of medial prefrontal cortex in a mouse
model for mental retardation and autism. Cereb Cortex *22*, 1333-1342. 10.1093/cercor/bhr224.
Wang, C.L., Zhang, L., Zhou, Y., Zhou, J., Yang, X.J., Duan, S.M., Xiong, Z.Q., and Ding, Y.Q. (2007).
Activity-dependent development of callosal projections in the somatosensory cortex. J Neurosci *27*,
11334-11342. 10.1523/JNEUROSCI.3380-07.2007.
Wang, J., Ethridge, L.E., Mosconi, M.W., White, S.P., Binder, D.K., Pedapati, E.V., Erickson, C.A., Byerly,
M.J., and Sweeney, J.A. (2017). A resting EEG study of neocortical hyperexcitability and altered
functional connectivity in fragile X syndrome. J Neurodev Disord *9*, 11. 10.1186/s11689-017-9191-z.
Yao, S., Becker, B., and Kendrick, K.M. (2021). Reduced Inter-hemispheric Resting State Functional
Connectivity and Its Association With Social Deficits in Autism. Front Psychiatry *12*, 629870.
10.3389/fpsyt.2021.629870.
Zerbi, V., Ielacqua, G.D., Markicevic, M., Haberl, M.G., Ellisman, M.H., A, A.B., Frick, A., Rudin, M., and
Wenderoth, N. (2018). Dysfunctional Autism Risk Genes Cause Circuit-Specific Connectivity Deficits
With Distinct Developmental Trajectories. Cereb Cortex *28*, 2495-2506. 10.1093/cercor/bhy046.
Zhang, Y., Bonnan, A., Bony, G., Ferezou, I., Pietropaolo, S., Ginger, M., Sans, N., Rossier, J., Oostra,
B., LeMasson, G., and Frick, A. (2014). Dendritic channelopathies contribute to neocortical and sensory
hyperexcitability in Fmr1(-/y) mice. Nat Neurosci *17*, 1701-1709. 10.1038/nn.3864.

Figure 1. L2/3 pyramidal neurons in barrel cortex of *Fmr1* KO mice have weak callosal synaptic inputs. (A) Timeline and schematic of experimental paradigm (B) Example image of S1 contralateral to AAV ChR2-mCherry injection. Left: DIC. Right: Red fluorescence of ChR2-mCherry labelled axons in corpus callosum (CC) and cortex. Recordings were performed on L2/3 neurons in an area of mCherry fluorescence. (C) LED stimulation paradigm (left) and example EPSCs from WT and *Fmr1* KO mice (right). Blue rectangle = 2ms blue LED flash. (D) Left: Raw LED-evoked EPSC amplitudes in WT and *Fmr1* KO animals (WT =172 ±18 pA, n=20; KO=105 ±11 pA, n=19; unpaired t-test); Right: LED-evoked EPSC amplitudes normalized to LED power (WT =2.2 ±0.1, n=20; KO=1.7±0.1, n=19; unpaired t-test). (E)

Schematic and example experiment of grid of blue laser stimulation during sCRACM (left) relative to

recorded neuron (right). (F) Example laser-evoked EPSCs at the locations highlighted in yellow in E from

WT and *Fmr1* KO mice. (G) Group average of EPSC amplitudes evoked at different locations relative to

the cell soma in WT and *Fmr1* KO mice. Individual maps are aligned by the location of soma (cyan dot).

Pixel color represents the average amplitude of EPSCs evoked from that location. (H) Left: Vertical profile

of mean synaptic input strength (mean input – average of EPSC amplitudes from individual locations

within a specific area) Left: Mean EPSC amplitudes along vertical axis of the cortex (soma, x=0) (*Fmr1*

n.s. $p=0.16$, $F_{(1, 60)}=2.060$; *Fmr1* x vertical position, $F_{(15, 849)}=1.723$; mixed-effects model). Right:

Mean of EPSC amplitudes in the outlined area in G (white) and left graph (blue), normalized to laser

power. (WT=0.674±0.098, n=40; KO=0.312±0.167, n=22; Mann Whitney). For this and all figures, error

bars represent standard error mean (SEM) and *$p<0.05$, **$p<0.01$.

Figure 2. Postnatal and postsynaptic deletion of *Fmr1* in L2/3 pyramidal neurons cell
autonomously weakens callosal synaptic inputs. (A) Timeline and schematic of experimental

paradigm (B) Simultaneous patch clamping of a neighboring AAV Cre-GFP+, *Fmr1* KO (arrowhead) and

GFP-, *Fmr1* WT (arrow) pyramidal neurons in L2/3 of barrel cortex. Left: DIC. Right: Green fluorescence.

(C) LED bulk stimulation paradigm (left) and example EPSCs from a pair of WT and *Fmr1* KO neurons

(right). Scale = 100pA, 20 ms. (D) Left: Group average of LED-induced EPSC amplitudes in WT and

Fmr1 KO neurons at P18-20 (top) (WT=238 ±30 pA; KO=281 ±27 pA, n=19 pairs, n.s.) and P23-30

(bottom) (WT=424 ±42 pA; KO=250±46 pA, n=15 pairs; paired t-test); Right: EPSC amplitudes from

individual cell pairs (open circles). Mean ±SEM (filled circle). Diagonal line represents equality. (E) EPSC

amplitudes, normalized to LED power, across different ages (replot from (D)) (*Fmr1* x Age ***$p<0.001$,

$F_{(1, 32)}=13.87$; *Fmr1*, Age, ns, ANOVA; #$p<0.1$, multiple comparison). (F) Schematic (left) and image

(right) of blue laser stimulation grid configuration and dual recordings for sCRACM experiments. (G)

Example of laser-induced EPSCs at location highlighted in E (yellow) from WT and *Fmr1* KO pair. Scale

= 100pA, 50 ms (H) Averaged color map of subcellular distribution of callosal synaptic input strengths

onto WT and *Fmr1* KO L2/3 pyramidal neurons at P23-30. (I) Vertical profile of mean synaptic inputs

strength along the neuronal dendritic tree (soma, x=0) (*Fmr1* $F_{(1, 13)}=5.956$; *Fmr1* x vertical position

$F_{(14, 144)}=1.793$; mixed-effects model). (J) Left: Quantification of mean synaptic inputs strength from

the highlighted positions in H (white) and I (blue), (WT=46 ±7; KO= 26 ±6 pA, n=14 pairs; Wilcoxon test);

Right: Distribution of values from individual cell pairs. *$p<0.05$; **$p<0.01$;***$p<0.001$

Figure 3. Callosal synapses onto *Fmr1* KO neurons have reduced quantal event frequency and a

selective weakening of AMPA receptor-mediated synaptic transmission (A) Example traces of Sr²⁺-

evoked quantal events. Scale = 25pA, 100 ms. Baseline, spontaneous events occur within 1s prior to

LED flash (red dotted line) and evoked events occur 50-350 ms after LED flash (red line). (B) Left:

Baseline frequency (top) (WT=1.3±0.1, KO=1.2±0.2 Hz, n.s., paired t-test) and amplitude (bottom)

(WT=12±1, KO=11±1 pA, n.s., Wilcoxon test) of quantal EPSCs for WT and *Fmr1* KO neuron pairs;

Right: distribution of values from individual cell pairs. (C) Left: Evoked frequency (top) (WT=8.3±0.5,

KO=6.7±0.5 Hz, *p<0.05, Wilcoxon test) and amplitude (bottom) (WT=15±1, KO= 15±1 pA, n.s.,

Wilcoxon test, n=13 pairs) of quantal EPSCs for WT and *Fmr1* KO neuron pairs; Right: distribution of

values from individual cell pairs. (D) Top: Example NMDAR EPSCs from a WT and *Fmr1* KO pair. Scale

= 25pA, 20 ms. Bottom: LED-induced NMDAR EPSC amplitudes of WT and *Fmr1* KO neuron pairs

(WT=124±20, KO=120±18 pA, n.s., paired t-test, n=15 pairs); Right: distribution of values from individual

cell pairs.

Figure 4. Cell autonomous deletion of *Fmr1* in postsynaptic L2/3 or L5 neurons results in weak action-potential driven synaptic transmission from callosal inputs. (A) Top: Average color map of action-potential mediated callosal synaptic input strengths onto pairs of WT and *Fmr1* KO L2/3 pyramidal neurons at P23-30; Bottom: Example responses from the highlighted positions (white). Scale = 100pA, 50 ms. (B) Top: Mean callosal inputs strength from area highlighted in white in A) (WT=61±13, KO=35±6 pA, n=11 pairs, *p<0.05, paired t-test); Bottom: Distribution of values from individual cell pairs. (C) Top: Average color map of callosal input strengths onto pairs of WT and *Fmr1* KO L5 pyramidal neurons at P23-30; Bottom: Example responses from the highlighted area (white). Scale = 200pA, 60 ms. (D) Top: Mean callosal input strengths onto L5 neurons from the area highlighted in white in C) (WT=49±9, KO=31±4 pA, n=12 pairs, *p<0.05, paired t-test). Bottom: distribution of values from individual cell pairs.

Figure 5. Sensory deprivation by whisker trimming normalizes callosal input strength onto postsynaptic *Fmr1* KO neurons. (A) Experimental paradigm. Trimming the whisker pad either

ipsilateral or contralateral to the AAV-Cre-GFP injected hemisphere deprived either the presynaptic
callosal projection neurons or postsynaptic *Fmr1* KO neurons, respectively, of patterned sensory
experience- driven activity. (B) Left: Raw LED-induced EPSC amplitudes in WT and *Fmr1* KO neurons
with presynaptic deprivation (WT=510±25; KO=376±41 pA, n=16 pairs, **p<0.01, paired t-test) or with
postsynaptic deprivation (WT=428±36; KO=433±31 pA, n=17 pairs, n.s., paired t-test): Right: Values from
individual cell pairs. (C and D) Left: Averaged color map of subcellular distribution of callosal synaptic
inputs onto WT and *Fmr1* KO neurons with presynaptic (C) or postsynaptic (D) sensory deprivation. Right:
Mean synaptic inputs strength onto soma and proximal apical dendrites (highlighted in white in C) for WT
and *Fmr1* KO neurons with presynaptic (WT=43±7; KO=31±5 pA, n=16 pairs, *p<0.05, Wilcoxon test) or
postsynaptic (WT=42±4; KO=42±6 pA, n=17 pairs, n.s., Wilcoxon test) sensory deprivation and values
from individual cell pairs. (E) LED-induced EPSC amplitudes normalized to LED power (replot from (B))
(*Fmr1* X deprivation interaction *p<0.05, F(1, 31)=4.977, ANOVA; presynaptic deprivation WT vs. *Fmr1*
KO, *p<0.05, *Fmr1* KO presynaptic deprivation vs. postsynaptic deprivation, *p<0.05, Sidak's multiple
comparisons).

Figure 6. Postsynaptic deletion of *Fmr1* in L2/3 neurons does not affect excitatory synaptic inputs from local columnar inputs. (A) Experimental design schematic. (B) Left: Position of UV laser scanning photostimulation (LSPS) grid (grey dots) relative to cortical layers and recorded neurons in L2/3 and L4 barrels (grey squares). Right: IR-DIC image of dual recordings in L2/3 in a slice with laser stimulation grid (red). Yellow dots indicate L4 home barrel. (C) Example of EPSCs in a pair of WT and *Fmr1* KO L2/3 neurons in response to LSPS and glutamate uncaging in L4 home barrel (yellow dots in B; right). (D) Color map of spatial distribution of average synaptic input strengths in response to LSPS. L4 home barrel (white rectangle). Cyan dots represent locations of soma and black pixels are direct responses. (E) Mean synaptic input strength from L4 home barrel, L4 adjacent barrels, adjacent L2/3 and L5A (WT vs. Fmr1 KO, n.s., n=17 pairs, Wilcoxon or paired t-test) (top) and values from individual cell pairs (bottom).

Figure 7. Reduced bilateral phase synchrony in EEG recordings of awake *Fmr1* KO mice. A.
Schematic of multielectrode array (MEA) electrode. Red circles indicate homotopic temporal sites
(electrodes 7 and 24) analyzed. B. Lower interhemispheric ISPC in beta and gamma frequency bands
during rest in *Fmr1* KOs compared to WT. (p_β=0.008, p_γ=0.013; unpaired t-test). C. *Fmr1* KO mice exhibit
lower interhemispheric ISPC during sound presentations compared to WTs. Color representation of
average ISPC values during a 100ms acoustic stimulus presentation at time = 0 (see methods). Using a
nonparametric permutation clustering method, we discovered a significant cluster of lower ISPC values
in the beta and gamma frequency ranges (black outline) in *Fmr1* KO mice during acoustic stimulus
presentation compared to WT mice.

Supplemental Figures and Legends:

Figure S1. Postsynaptic deletion of *Fmr1* does not affect local columnar circuit inputs onto L2/3

pyramidal neurons at 2 weeks of age. (A) Right: Sparse deletion of *Fmr1* in L2/3 neurons using

AAV.GFP-Cre injection at P1 does not affect local synaptic input strength from L4 home barrel (L4H)

measured at P14-17 (n = 8 pairs). (B) Similar as (A), but experiment was done in mosaic *Fmr1* females

(*Fmr1^-/gfp*) (n = 11 pairs). Half of the cells are *Fmr1* KO GFP- while the other half are *Fmr1* WT GFP+ due

to X chromosome inactivation. (C) Averaged LSPS color maps (left) and quantification of local synaptic

input strength from L4 home and adjacent (L4A) barrels (right) comparing across WT and *Fmr1* global

KO animals at 3 weeks of age (P18-25) (n = 15 for WT, n = 20 for KO). Quantified inputs were normalized

to the respective laser power and log transformed to enable cross-animal comparison.

Figure S2. Miniature (m) EPSC frequency and amplitude, as well as input resistance from pairs of WT and postsynaptic *Fmr1* KO L2/3 pyramidal neurons in P23-30 sensory intact, presynaptic sensory deprived and postsynaptic sensory deprived animals.

Figure S3. Summary model. During postnatal development of cortical circuits for mouse primary somatosensory barrel cortex (postnatal 4 weeks), *Fmr1* in the postsynaptic L2/3 pyramidal neurons promotes callosal synaptic connections from the contralateral side by maintaining the AMPAR transmission in the synapse. Without postsynaptic *Fmr1*, some callosal synapses undergo synaptic silencing which is dependent on sensory experience driven activity in the postsynaptic neuron.