Coupling of pupil- and neuronal population dynamics reveals diverse influences of arousal on cortical processing

Thomas Pfeffer[1,2*@], Christian Keitel[3,4*], Daniel S. Kluger[5,6], Anne Keitel[7], Alena Russmann[2], Gregor Thut[4], Tobias H. Donner[2**], Joachim Gross[4,5,6**]

1 Universitat Pompeu Fabra, Center for Brain and Cognition, Computational Neuroscience Group, Barcelona, Spain | **2** University Medical Center Hamburg-Eppendorf, Department of Neurophysiology and Pathophysiology, Martinistr. 52, 20246 Hamburg, Germany | **3** University of Stirling, Psychology, FK9 4LA Stirling, UK | **4** Centre for Cognitive Neuroimaging, Institute of Neuroscience and Psychology, University of Glasgow, 62 Hillhead Street, G12 8QB Glasgow, UK | **5** Institute for Biomagnetism and Biosignal Analysis, University of Münster, Malmedyweg 15, 48149 Muenster, Germany | **6** Otto Creutzfeldt Center for Cognitive and Behavioral Neuroscience, University of Münster, Fliednerstrasse 21, 48149 Muenster, Germany | **7** University of Dundee, Psychology, DD1 4HN Dundee, UK

* both authors contributed equally, joint first authorship; ** both authors contributed equally, joint senior authorship; @ corresponding author - email: thms.pfffr@gmail.com

Abstract

Fluctuations in arousal, controlled by subcortical neuromodulatory systems, continuously shape cortical state, with profound consequences for information processing. Yet, how arousal signals influence cortical population activity in detail has only been characterized for a few selected brain regions so far. Traditional accounts conceptualize arousal as a homogeneous modulator of neural population activity across the cerebral cortex. Recent insights, however, point to a higher specificity of arousal effects on different components of neural activity and across cortical regions. Here, we provide a comprehensive account of the relationships between fluctuations in arousal and neuronal population activity across the human brain. Exploiting the established link between pupil size and central arousal systems, we performed concurrent magnetoencephalographic (MEG) and pupillographic recordings in a large number of participants, pooled across three laboratories. We found a cascade of effects relative to the peak timing of spontaneous pupil dilations: Decreases in low-frequency (2-8 Hz) activity in temporal and lateral frontal cortex, followed by increased high-frequency (>64 Hz) activity in mid-frontal regions, followed by linear and non-linear relationships with intermediate frequency-range activity (8-32 Hz) in occipito-parietal regions. The non-linearity resembled an inverted U-shape whereby intermediate pupil sizes coincided with maximum 8-32 Hz activity. Pupil-linked arousal also coincided with widespread changes in the structure of the aperiodic component of cortical population activity, indicative of changes in the excitation-inhibition balance in underlying microcircuits. Our results provide a novel basis for studying the arousal modulation of cognitive computations in cortical circuits.

Keywords: Pupillometry, pupil-linked arousal, pupil, brain rhythms, MEG, locus coeruleus (LC), basal forebrain, noradrenaline, acetylcholine, cortex, brainstem

Introduction

Variations in cortical state profoundly shape information processing and, thus, cognition (Busse et al., 2017; Fu et al., 2014; Zagha et al., 2013). These variations occur continuously due to subtle fluctuations in the level of arousal, and even in the absence of changes in overt behavior (Harris & Thiele, 2011; McGinley, Vinck, et al., 2015). Two key mediators of arousal-dependent variations are the brainstem nucleus locus coeruleus (LC), which supplies noradrenaline (NE), and the basal forebrain (BF), which supplies acetylcholine (ACh)(Harris & Thiele, 2011; Hasselmo, 1995; Lee & Dan, 2012; Steriade, 1996). It has long been thought that these nuclei are organized homogeneously and innervate cortical target regions diffusely, and indiscriminately across cortical regions. This has led to the idea that the arousal system acts as a global "broadcast signal" and uniform controller of cortical state (Aston-Jones & Cohen, 2005; Harris & Thiele, 2011; Leopold et al., 2003; Turchi et al., 2018).

More recently, however, this view has been challenged: Several neuronal subpopulations with distinct projection targets have been found in both the LC and BF (Chandler et al., 2013, 2019; Sarter et al., 2009; Schwarz & Luo, 2015; Totah et al., 2018; Zaborszky et al., 2015; Záborszky et al., 2018). In addition, neuromodulator receptors exhibit a rich and diverse distribution across the cortex (Burt et al., 2018; van den Brink et al., 2019). In this report, we comprehensively show how this structural and molecular heterogeneity translates into a multitude of arousal effects on neuronal population activity across the human cortex.

We used pupil diameter as a proxy for arousal (Beatty, 1982; Bradshaw, 1967; Hess & Polt, 1964; Kahneman et al., 1967; Kahneman & Beatty, 1966). Pupil diameter has recently attracted interest in neuroscience as a peripheral index of cortical state (McGinley, David, et al., 2015; Reimer et al., 2014; Vinck et al., 2015). Spontaneous changes in pupil diameter mimic the effects seen during the behavioral state transition from quiet wakefulness to locomotion in rodents (Crochet & Petersen, 2006; Niell & Stryker, 2010; Polack et al., 2013; Poulet & Petersen, 2008): Suppression of low (< 10 Hz) and elevation of higher frequency (> 30 Hz) activity (McGinley, Vinck, et al., 2015; Reimer et al., 2014; Vinck et al., 2015). Critically, fluctuations in pupil diameter under constant retinal illumination track LC-NE activity (Breton-Provencher & Sur, 2019; de Gee et al., 2017; Joshi et al., 2016; P. R. Murphy et al., 2014; Reimer et al., 2016) as well as BF-ACh activity (Reimer et al., 2016). Fluctuations in pupil diameter are reflected more strongly in cholinergic activity, whereas the rate of change in pupil diameter (i.e., its temporal derivative) is most strongly associated with noradrenergic activity (Reimer et al., 2016).

Rodent research has mostly been limited to circumscribed sensory cortical areas (Joshi & Gold, 2020; McGinley, David, et al., 2015; Reimer et al., 2014; but see Shimaoka et al., 2018). Insights from non-invasive magnetic resonance imaging in humans, on the other hand, have revealed spatially rich correlations of cortical activity with pupil-linked arousal (Groot et al., 2021; Schneider et al., 2016; Yellin et al., 2015). However, the low temporal resolution of these measurements complicates direct comparisons with results from electrophysiological recordings in rodents and non-human primates. Therefore, we still lack a comprehensive understanding of which features of cortical population activity are shaped by arousal and how such effects are distributed across cortex.

To close these gaps, we assessed the co-variation of spontaneous fluctuations in pupil-linked arousal and whole-head magnetoencephalography (MEG) recordings in resting human participants across brain regions (Figure 1), components of cortical activity (oscillatory and aperiodic), and temporal lags. Our findings provide a comprehensive picture of linear and nonlinear relationships between intrinsic fluctuations in pupil-linked arousal and neural population activity in different brain regions. First, we demonstrate opposite effects of pupil-linked arousal on low- and high-frequency components of cortical population activity across large parts of the brain, similar to previously observed effects in rodent sensory cortices. These distinct effects occurred in a systematic temporal sequence. Second, we also uncovered hitherto unknown, non-linear (inverted U-shape) arousal effects in occipito-parietal cortex. Third, we identified spatially widespread correlations between pupil-linked arousal and the structure of aperiodic activity, suggesting underlying changes in cortical excitation-inhibition balance (Gao et al., 2017; Waschke et al., 2021).

Figure 1. Rationale of our approach and summary results. **(A)** Even at rest and under constant illumination, pupil size (diameter) fluctuates spontaneously. **(B)** The Locus Coeruleus (LC) in the brainstem and the Nucleus Basalis of Meynert in the basal forebrain (BF) affect pupil size. LC releases noradrenergic (NE) neuromodulation to the cortex. BF releases Acetylcholine (ACh). Synchronous local neuronal population activity across the brain can be measured with MEG. **(C)** Rationale: Studying the coupling of pupil size and local population activity recorded with MEG provides insights into the functional influences of deep-brain structures in brainstem (LC) and basal forebrain (BF). Modified from Van den Brink et al. (2019).

Results

We analysed data from concurrent whole-head MEG and pupil recordings in 81 human participants, collected at three MEG laboratories (Glasgow, UK; Hamburg & Münster, Germany). All recordings were carried out under constant illumination and in sound-attenuated environments, thus removing fluctuations in pupil size due to changes in sensory input. Participants were instructed to keep their eyes open, while fixating and resting otherwise. The length of individual recordings varied between 5 - 10 min (see Methods for further details).

For each of the three MEG laboratories, we obtained time courses of MEG activity and of pupil diameter fluctuations with highly similar spectral characteristics (Supplementary Figures S1 and S2). In what follows, we refer to the time course of pupil diameter as 'pupil'. Previous work in rodents suggests that this time course and its temporal derivative indicate distinct (cholinergic and noradrenergic, respectively) neuromodulatory inputs to cortex (Reimer et al., 2016). Thus, we also tested for relationships of MEG activity with the first temporal derivative

of the pupil diameter time course (hereafter: 'pupil-derivative'). We found that, in our data, the results were largely similar for pupil and pupil-derivative. Therefore, the main Figures show only results for pupil, whereas the results for pupil-derivative are presented in the Supplement (Supplementary Figures S3-8).

Robust pupil-brain couplings across the spectrum of cortical population activity

We evaluated the strength of pupil-brain coupling across a wide range of frequency bands in MEG sensor space by means of mutual information (MI) which measures all (linear and nonlinear) associations between two signals (Ince et al., 2017). We calculated the MI between pupil traces and power time courses of band-limited cortical activity across 25 log-spaced center frequencies from 2 to 128 Hz at each MEG sensor (see Supplementary Figures S1 and S2 for power spectra of MEG and pupil signals). These MI values were standardised against permutation distributions of surrogate MI values obtained by time-shifting the pupil- against the power time courses, then averaged across all sensors for a first summary statistic by recording site (see Methods).

For all three MEG laboratories, standardised-MI spectra showed similar patterns of widespread pupil-brain associations across frequencies with prominent peaks in the 8-32 Hz range (Figure 2A). MI scalp distributions within this frequency range further underpinned the commonalities between recording sites despite different MEG systems and setups (Figure 2B). Significant relationships with the pupil were also evident for the low-frequency (4-8 Hz) and high-frequency (64-128 Hz) components of MEG-power (Figure 2A).

Figure 2. (A) Mutual information (MI) spectra demonstrate the covariation between pupil-size and power fluctuations across canonical frequency bands from 2 - 128 Hz (note the log-spaced frequency axis). MI values are standardised against individual permutation distributions based on a temporal re-shuffling of pupil-size and power time courses (for details see Methods). Filled/open markers in plots signify frequency-specific t-tests of individual standardised MI values against zero (two-sided) at an FDR-corrected/uncorrected threshold of *P*<0.05. Data pooled across all MEG sensors. Shaded areas depict the standard error of the mean across subjects. Color-coding refers to the recording site; left: Glasgow data in red; center: Hamburg data in blue; right: Münster data in yellow. N denotes the number of samples recorded at each site. **(B)** Scalp topographies illustrate the commonalities of the spatial distribution of pupil-size power covariations for two prominent frequency ranges. As in A, colors code the recording site - from left to right: Glasgow, Hamburg, Münster.

Fluctuations in pupil diameter lag behind changes in cortical population activity in a frequency-dependent manner

Having established robust pupil-brain couplings, we quantified the temporal relationship between both signals. Changes of neural activity in brainstem arousal centers such as the LC generate changes in neural activity that exhibit considerable temporal lag (~500 ms in

macaques) and temporal low-pass characteristics (Breton-Provencher & Sur, 2019; de Gee et al., 2017; Hoeks & Levelt, 1993; Joshi et al., 2016; Korn & Bach, 2016). This temporal smoothing is produced by the peripheral pathways (nerves and muscles) controlling pupil motility (Hoeks & Levelt, 1993; Korn & Bach, 2016). Thus, one might expect the effect of
160 neuromodulatory arousal signals to occur earlier in cortical population activity (a single synapse from LC to cortex, unless mediated by a third structure, such as the thalamus) than in pupil diameter.

We computed cross-correlations between fluctuations in frequency-resolved sensor-level power and pupil (Methods). We found the maximum correlation magnitude between pupil
165 diameter and power fluctuations at a frequency-dependent lag (Figure 3A, B): For activity in the low (around 2-4 Hz) and high frequency bands (around 64-128 Hz), correlation magnitudes peaked at lags of 832 ms (negative peak) and 562 ms (positive peak), respectively. This result was highly consistent across the three MEG sites (Figure 3A). These latencies indicated power fluctuations that preceded corresponding variations in pupil
170 diameter. Correlations with power fluctuations in the 8 - 16 Hz range, on the other hand, peaked at a lag closer to zero (210 ms; negative peak), indicating a temporally closer relationship (Figure 3A, B; see Discussion).

Fluctuations in pupil, and pupil-derivative, may primarily reflect cholinergic and noradrenergic activity, respectively (Reimer et al., 2016). Consistent for all three data sets,
175 the cross-correlation analysis for pupil-derivative revealed peaks closer to zero lag for low and high frequencies (2 - 4 Hz negative peak at 277 ms, 64 - 128 Hz positive peak at -90 ms; Supplementary Figure S3). In contrast, correlations with activity in the 8 - 16 Hz range peaked somewhat later (700 ms; negative peak; Supplementary Figure S3). The shorter lags for the pupil derivative are consistent with previous findings in rodents (Reimer et al., 2016).

180 The cortical innervation profiles of subcortical arousal centers and the cortical receptor composition for the neuromodulators released by these centers exhibit substantial heterogeneity across cortex (Ramos & Arnsten, 2007; van den Brink et al., 2018, 2019). We, therefore, reasoned that the nature (frequency-dependence and/or temporal lag) of pupil-brain couplings may differ considerably across cortical areas. We next tested the
185 topographical profile of the cross-correlations. To account for the different channel layouts of the three MEG systems, we averaged cross-correlations, computed for individual MEG channels, within 39 non-overlapping channel groups, consisting of channels that were grouped based on their location along the anterior-to-posterior axis (see Methods). In keeping with the previous analyses, we collapsed the data into three non-overlapping frequency
190 bands of interest: 2 - 4 Hz, 8 - 16 Hz, and 64 - 128 Hz.

Figure 3. Sensor level cross-correlations between fluctuations in pupil diameter and band-limited power fluctuations. **(A)** Correlation between fluctuations in pupil diameter and band-limited power fluctuations across lags, ranging from -5 to 5s and for all three recording sites (left: Glasgow; center: Hamburg; right: Muenster). Colors indicate the MEG frequency-band, with darker colors representing lower frequencies and lighter colors representing higher frequencies. Negative lags are indicative of "pupil preceding MEG", whereas positive lags are indicative of the opposite. The vertical dashed black line denotes a lag of 930 ms (Hoeks & Levelt, 1993) and the dashed light gray line a lag of 0 ms. **(B)** Same as in (A), but averaged across the three recording sites and for three frequency-bands of interest: 2-4 Hz (black), 8-16 Hz (dark gray) and 64-128 Hz (gray dashed). **(C)** Correlation values across all channel groups, sorted from anterior to posterior, and for lags ranging from -5 to +5s. The dashed gray line depicts a lag of 0 ms and the dashed black line shows a lag of 930 ms (Hoeks & Levelt, 1993). Correlation values were averaged within three frequency-bands of interest: 2-4 Hz (left), 8 - 16 Hz (center) and 64-128 Hz (right).

Consistent with the results obtained from sensor averages, a negative correlation between pupil diameter fluctuations and power in the low frequencies (2-4 Hz) peaked at a lag of around 900 ms, with a spatial profile that was relatively uniform across brain areas (Figure 3C, left). For the high frequencies (64-128 Hz), on the other hand, the positive peak was confined to posterior regions, with a lag similar to the low frequency band (Figure 3C, right). In the 8 to 16 Hz-range, the picture was more nuanced (Figure 3C, center): The close-to-zero-lag positive correlation dominated posterior sensors whereas anterior sensors registered a negative correlation with a peak around 1000 ms. This suggests that the correlation between pupil diameter and power in the 8 - 16 Hz range may be driven by independent mechanisms that operate on distinct time scales (see Discussion).

The spatio-temporal correlation patterns between the pupil-derivative and band-limited power is shown in Supplementary Figures S3 and S4.

In summary, we identified lags between spontaneous fluctuations in pupil diameter and low- and high-frequency activity that are consistent with previously reported lags between noradrenergic and cholinergic activity, respectively, and spontaneous pupil diameter fluctuations in rodents. In addition, we found that pupil-brain coupling in the intermediate 8-16 Hz range peaked with a time lag closer to maximum pupil dilation for posterior sensors. Our findings thus suggest that the correlations in different frequency bands, and for different

cortices within the intermediate frequency range, may be mediated by different neuromodulatory systems (see Discussion).

Spatial and spectral dissociations of pupil-power correlations

230   Having established the distinct temporal coupling profiles for different frequency bands, we now tested which cortical areas received the strongest modulation. Sensor space analyses suggested that effects of pupil-brain coupling were not uniformly distributed along the anterior-posterior axis (Figure 3). To map out cortical loci of frequency-specific pupil-brain coupling in detail we used spectrally resolved source-reconstructed MEG data. Given

235   the high consistency of the sensor-space results, we report data pooled across the three recording sites for the following analyses.

Based on the results obtained from the cross-correlation analysis and in line with previous reports (Hoeks & Levelt, 1993; Joshi et al., 2016; Reimer et al., 2016), we shifted the pupil signal 930 ms forward (with respect to the MEG signal). In other words, we shifted the pupil

240   signal to compensate for the lag that had been observed between external manipulations of arousal (Hoeks & Levelt, 1993) as well as spontaneous noradrenergic activity (Reimer et al., 2016) and changes in pupil diameter. Using the forward-shifted pupil signal, we computed correlations with fluctuations in spectral power (Figure 4), separately for 8799 source locations covering the 246 regions of the Brainnetome atlas (Fan et al., 2016; see Methods).

245   For each region, pupil-power associations were quantified by means of Spearman's rank correlation. All steps were repeated for correlations between pupil-derivative and power fluctuations, however without accounting for the lag as previous research showed near-instantaneous correlations between neuromodulatory activity and fluctuations in pupil derivative (Reimer et al., 2016).

250   Averaged across all source locations, correlations between intrinsic pupil and power fluctuations were frequency-dependent (Figure 4A): Consistent with findings from invasive recordings in primary visual cortex of rodents (Reimer et al., 2014; Vinck et al., 2015), we found a robust negative correlation between pupil diameter and power in the low frequencies (2-8 Hz) and a numerically small, albeit statistically robust, positive correlation in the high

255   frequencies (between 50 Hz and 100 Hz). In order to investigate the spatial distribution of the correlations, we sorted the regions of the Brainnetome atlas along the anterior-posterior axis. Whereas the correlations in the low frequencies covered large parts of the brain (except for a few posterior regions, Figure 4B), pupil-brain coupling in other frequency ranges exhibited a spatially richer structure, with positive (8 - 32 Hz) and negative (> 50 Hz) correlations in

260   more occipital regions and correlations of opposite sign in more anterior sources (Figure 4B).

Previous studies have reported largely similar effects of pupil-linked arousal on population activity in the different sensory cortices (McGinley, David, et al., 2015; Reimer et al., 2014; Vinck et al., 2015; but see Shimaoka et al., 2018). Capitalising on our source reconstructions, we tested if the correlations between pupil diameter and cortical activity varied across cortical

265   regions of interest (see inset maps in Figure 4C), including three sensory areas (visual cortex, auditory cortex and somatosensory cortex) as well as anterior cingulate cortex, a region receiving strong innervations from locus coeruleus (Chandler et al., 2013). Patterns of pupil-power correlations were highly region-specific: Correlations in the primary visual cortex were

positive within the 8-32 Hz range (Figure 4C, top). Correlations in the auditory cortex, on the

270 other hand, were dominated by negative correlations in the low frequency range (< 8 Hz; Figure 4C, second to top). Somatosensory cortex exhibited negative correlations in the low and intermediate frequencies (2-16 Hz) and, at the same time, positive correlations in the high frequency range (~ 64 Hz; Figure 4C, second to bottom). Correlations with activity in anterior cingulate were significant across most frequencies, with a spectral pattern similar to auditory

275 cortex (Figure 4C, bottom).

Figure 4. Source space pupil-power correlations. **(A)** Spectrum of pupil-power correlations, averaged across all 8799 source locations. Filled black circles denote significant correlations (P < 0.05; two-tailed paired t-test; FDR-
280 adjusted, q=0.1). The gray shaded area depicts the standard error of the mean across subjects. **(B)** Pupil-power correlations, averaged across the 246 regions of the Brainnetome atlas and sorted from anterior to posterior regions. Masked at *P* = 0.05 (two-tailed paired t-test; FDR-adjusted, q=0.1). **(C)** Pupil-power correlations across four selected regions-of-interest (see Methods for details): visual cortex auditory cortex, somatosensory cortex and anterior cingulate cortex (from top to bottom). Insets depict the spatial extent of the respective region of
285 interest. **(D)** Spatial distribution of pupil-power correlations, across four frequency bands of interest. Correlations between pupil and power (in clockwise direction, starting from the top left) at 4 Hz, 11.3 Hz, 19.0 Hz, and 64 Hz. All maps are masked at *P* = 0.05 (two-tailed paired t-test; FDR-adjusted, q=0.1).

For further insights into the spatial distribution of the observed correlations, we mapped
290 correlations of pupil and pupil-derivative time courses with band-limited power fluctuations across all 8799 source locations, again estimated at a lag of 930 ms for pupil and a lag of 0 ms for pupil-derivative. To this end, we focused on four frequency bands of interest, with the following center frequencies (Figure 4D): 4 Hz, 11.3 Hz, 19 hz, and 64 Hz, corresponding to the classical theta, alpha, beta, and gamma frequency ranges. In the low frequencies
295 (centered at 4 Hz), correlations were negative in the majority of source locations, with bilateral peaks in the anterior sections of the hippocampus (Figure 4D, top left). Significant positive correlations between pupil and activity in the alpha-/beta range (11.3 Hz and 19 Hz) were located in the visual cortices of both hemispheres (Figure 4D, top right and bottom left). Significant negative correlations were found in left and right motor and somatosensory
300 cortices (Figure 4D, top right), although limited to activity in the alpha-range (centered at 11.3

Hz). Similarly, correlations with high frequency activity (centered at 64 Hz) were non-uniform across space, with negative correlations located in visual cortices bilaterally, and positive correlations across large parts of the prefrontal cortex (Figure 4D, bottom right).

305   Repeating the analyses for the pupil derivative, we found largely similar spatial patterns for the correlations, albeit with reduced magnitudes (Supplementary Figure S5; see Supplementary Figure S6 for statistical comparison of pupil-power and pupil-derivative-power correlations). McGinley et al. (2015) reported a similar reduction when comparing pupil-derivative with pupil-based correlations in rodents.

310   Briefly summarised, source localisation of pupil-brain coupling revealed frequency-band specific patterns of arousal influences on cortical processing. While coupling with low-frequency activity was relatively uniform, coupling with intermediate- and high-frequency activity showed effects of opposite signs for different cortices, even within frequency bands, indicating that arousal has discernible effects on different sensory cortices, as well as higher-order associative regions of the human brain at rest.

315   **Nonlinear relations between pupil-linked arousal and band-limited cortical activity**

Above described analyses focused on quantifying linear relationships between intrinsic fluctuations in pupil diameter, as well as pupil-derivative, and band-limited cortical population activity. However, arousal may exert more complex influences on cortical activity that roughly follow a nonlinear, inverted-U shape, known as the Yerkes-Dodson law (Yerkes & Dodson, 1908; Aston-Jones & Cohen, 2005; McGinley, David, et al., 2015). Evidence suggests that gain of neuronal input-output functions as well as behavioural performance peak at intermediate arousal levels (He & Zempel, 2013; McGinley, Vinck, et al., 2015; Waschke et al., 2019), flanked by lower gain and performance at lower (e.g. disengagement and drowsiness) and higher arousal states (e.g. stress). We, therefore, reasoned that local pupil-power couplings may exhibit similar nonlinear relationships. To test this, we divided the data into non-overlapping temporal segments of 2s length and computed power spectra as well as mean pupil size (or mean pupil-derivative) for each segment. Next, we averaged the power spectra within each of 14 evenly sized pupil bins, ranging from smallest to largest pupil diameter on each block.

330   Focusing on three predefined frequency bands, the relation between pupil diameter and pupil-derivative with low-frequency (2-4 Hz) and high-frequency activity (64-128 Hz) was well-characterized by a linear relationship (Figure 5A, left and right) when averaged across all source locations. In contrast, activity in the 8-16 Hz range exhibited a high degree of nonlinearity that followed an inverted-U-relationship (Figure 5A, center).

335   To test for potential nonlinear relationships exhaustively across frequencies and space, we fitted a quadratic polynomial to the power-by-pupil bin functions, separately for each source region and frequency (see Methods). Next, we mapped the coefficient of the quadratic term of the polynomial (β_2) across space and frequencies. In order to account for differences in absolute power (offset), we normalized (i.e., z-scored) the power-by-pupil functions prior to model fitting. The analysis revealed that inverted-U relationships were exclusive to the range of 8-32 Hz (i.e., β_2 significantly smaller than zero; two-sided paired t-test; Figure 5B).

Figure 5. Nonlinear relations between pupil-linked arousal and band-limited power fluctuations. (A) Normalized spectral power, averaged across three frequency ranges (2-4 Hz, 8-16 Hz and 64-128 Hz; from left to right), as a function of mean pupil diameter. **(B)** Normalized coefficient of the quadratic term (β_2; See Methods for details), averaged across all subjects and atlas regions. Black dots denote $P<0.05$, red dots denote $P<0.01$ and yellow dots denote $P<0.05$ after adjusting for false discoveries (q = 0.1; two-sided paired t-test). The gray shaded area depicts the standard error of the mean across subjects. Positive and negative values are indicative of a U-shaped and inverted-U-shaped relationship, respectively **(C)** Normalized β_2 coefficient, averaged across all subjects, separately for four regions of interest: visual cortex, auditory cortex, somatosensory cortex and anterior cingulate cortex (from left to right). Insets show the approximate extent of the regions of interest. **(D)** Spatial distribution of the coefficient of the quadratic term (see Methods for details), at center frequencies 11 Hz (left) and 19 Hz (right). Masked at $P = 0.05$ (two-sided paired t-test, adjusted for a false discovery rate of 0.1).

Focusing on the quadratic nonlinearity in four regions of interest (as used in the previous analyses) pointed towards locally emphasised intermediate-frequency inverted-U shaped relationships (auditory and somatosensory cortices, Figure 5C). A spatially more fine-grained analysis of these effects revealed that the inverted-U relationship in the alpha band (~11 Hz) peaked in the left superior parietal lobule (Figure 5D, center left). The inverted-U relationship between activity in the low beta band (~19 Hz) and pupil diameter, on the other hand, additionally comprised bilateral parietal and temporal regions (Figure 5D, right). We found similar non-linear effects, with largely comparable spatial patterns when analysing the pupil derivative instead of the pupil diameter (Supplementary Figure S7).

Taken together, we found evidence for a non-linear inverted-U relationship between pupil-linked arousal and band-limited cortical activity that was confined to the classical alpha/beta frequency band (8-32 Hz) and parieto-occipital cortices. Put differently, cortical activity in this frequency range, and in the respective cortical regions, was maximal at medium pupil

diameters that indicate intermediate optimal levels of arousal. Activity in surrounding low- and high-frequency ranges showed exclusively linear effects.

370 **Pupil-linked arousal also predicts changes in aperiodic components of cortical population activity**

So far, we have shown how pupil-linked arousal co-varies with band-limited cortical activity. However, cortical electrophysiological processes can also be characterised by a broadband, or aperiodic component. This component can be understood as the decay rate
375 of power with increasing frequency, and is therefore also known as the 1/f component. The slope of the 1/f component reflects fluctuations in attentional state (Waschke et al., 2021) and has been suggested to track the ratio between excitation and inhibition (in short: E/I) in the underlying neuronal circuits (Gao et al., 2017). As neuromodulators linked with the regulation of arousal have been shown to change cortical E/I (Froemke, 2015; Pfeffer et al., 2018, 2020),
380 we next investigated the relation between intrinsic fluctuations in pupil diameter (as well as its temporal derivative) and the aperiodic component in the power spectrum. Our goal was to characterize correlations between intrinsic fluctuations in pupil-linked arousal and the slope of the aperiodic component, as a potential marker of cortical E/I.

To this end, we parameterized the power spectra of consecutive data epochs of the
385 source-projected MEG data in a frequency range from 3 to 40 Hz, separately for all source locations (Donoghue et al., 2020). The resulting parameter estimates described the periodic and the aperiodic components per epoch (data segmentation as in the previous section; see Methods).

We correlated fluctuations in the estimated spectral exponent of the aperiodic
390 components, a measure that expresses the steepness of the 1/f slope, with fluctuations in pupil diameter. The spectral exponent decreased linearly as a function of pupil diameter (Figure 6A) - note that an decrease in the spectral exponent means a shallower decay of power as a function of frequency (see Methods for details). This resulted in a significant negative correlation when averaged across all source locations (Figure 6B, grey dots), with
395 little variability when zooming in on four regions of interest (Visual cortex: mean r = -0.0266; P = 0.0013; Auditory cortex: mean r = -0.0484 , P = 8.12*10^{-8} ; Somatosensory cortex: mean r = -0.0412, P = 3.75*10^{-5}; Anterior cingulate cortex: r = -0.0325, P = 1.13*10^{-5}; two-tailed paired t-tests against zero). In fact, the negative correlation between the spectral exponent and pupil diameter was significant in the vast majority of brain regions (8197/8799 of source
400 locations, or 93.2%, after adjusting for a false discovery rate of 0.1; Figure 6C). A correlation was notably absent in several extrastriate occipital cortices.

We found a similarly widespread negative correlation between the spectral exponent and pupil-derivative. Albeit weaker in magnitude, this effect additionally encompassed the entire occipital cortex and showed stronger local maxima in parietal cortices, when compared with
405 the correlation with pupil (Supplementary Figure S8A-C).

In sum, these results speak to a global effect of pupil-linked arousal on the excitation-inhibition ratio across the brain, with few potential exceptions (occipital cortices). Minor

differences between pupil- and pupil-derivative-based spatial patterns of these effects may again point towards the involvement of different neuromodulators.

Figure 6. Pupil-linked arousal correlates with aperiodic component of the power spectrum. (A) Spectral exponent (i.e., the slope of the aperiodic component) for 20 pupil bins (sorted from small to large). **(B)** Correlation coefficients of the correlation between the spectral exponent and pupil diameter fluctuations, averaged across space (black dots, showing the individual subjects) and for the four regions of interest: visual cortex (red), auditory cortex (yellow) somatosensory cortex (purple) and anterior cingulate cortex (green). The horizontal lines highlight the mean correlation coefficient (*** $P<0.001$, ** $P<0.01$; two-sided paired t-tests). **(C)** Spatial map of the correlation between the slope of the aperiodic component and intrinsic fluctuations in pupil diameter. Spatial maps are masked at $P = 0.05$ after accounting for the false discovery rate (two-sided paired t-test). **(D)** Correlations between fluctuations in pupil diameter and band-limited power, after removing the aperiodic component. Note the truncated frequency axis, due to fitting the slope in the 3-40 Hz range. Black dots denote significant values ($P<0.05$; FDR-adjusted; two-sided paired t-test).

Couplings between pupil-linked arousal and periodic components of cortical population activity are not induced by couplings with aperiodic components

Given the pupil-linked changes in the slope of the (aperiodic component of) the cortical power spectrum, one concern is that these introduced spurious changes in band-limited power (Donoghue et al., 2020), thus potentially producing spurious correlations between fluctuations in pupil diameter and band-limited power. To test whether the correlations between pupil and spectral power reported above were merely a result of changes in the slope of the power spectra, we subtracted the fitted aperiodic component from the empirical power spectra (in the range from 3 Hz to 40 Hz), separately for each time segment and source location. Subsequently, we correlated the residual power estimates with pupil diameter, across all segments.

The pattern of correlations between pupil diameter and power remained largely unchanged (Figure 6D): Fluctuations in pupil diameter were negatively correlated with activity in the low frequencies, around 4 Hz, as well as with activity in a higher frequency range around 32 Hz. Band-limited activity in the range between 8 and 16 Hz exhibited robust positive correlations.

Correlations between residual power and pupil-derivative, on the other hand, were strongly reduced in magnitude and not significant for most frequencies (Supplementary Figure S8D).

These control analyses confirm that the couplings between pupil (and pupil-derivative) and band-limited cortical activity reported in the preceding sections were genuine, reflecting modulations of specific cortical oscillations by arousal.

Discussion

Pupil-linked arousal exerts pivotal influences on the states of sensory cortices (McGinley, David, et al., 2015; Reimer et al., 2014; Vinck et al., 2015) and associative cortical regions (Joshi & Gold, 2020). However, subcortical arousal systems also project to a wealth of other regions across the cerebral cortex. This warranted a detailed investigation of how these projections shape cortical population activity. To achieve this, we exhaustively mapped the relationship between intrinsic fluctuations in pupil-linked arousal and resting-state human MEG-recorded cortical activity. In a large sample (*N*=81), our mapping revealed diverse and consistent linear and non-linear arousal effects on band-limited, as well as aperiodic cortical activity.

Arousal modulates cortical activity across space, time and frequencies

Our findings are consistent with previously reported links between both locomotion-related and pupil-related changes in arousal and LFP recordings in sensory cortices of rodents (McGinley, David, et al., 2015; Reimer et al., 2014, 2016). Briefly put, periods of elevated arousal were accompanied by a decrease in low- and an increase in high-frequency activity, similar to cortical state changes typically observed during increased attention (Harris & Thiele, 2011; Schwalm & Jubal, 2017). The observed temporal relationships also align with the lags between manipulations of arousal state and pupil dilations of around 500-1000 msec in humans (Hoeks & Levelt, 1993; Wierda et al., 2012) and a ~500 msec delay between LC activation and peak pupil dilation in non-human primates (Joshi et al., 2016). In rodents, Reimer et al. (2016) and Breton-Provencher & Sur (2019) reported peak correlations between noradrenergic axonal activity and pupil diameter at around 900 ms, whereas the correlation between basal forebrain-regulated cholinergic activity and pupil diameter peaked around 500 ms (Reimer et al., 2016). Here, we report comparable peak lags for low (832 ms) and high frequency activity (562 ms), suggesting that different neuromodulatory systems may drive correlations in different frequency ranges in the human brain. Future research using brain-wide maps of the distribution of neuromodulatory receptors (van den Brink et al., 2019) may help to further disentangle the differential contributions of noradrenaline and acetylcholine as well as their various receptor subtypes.

In addition to the bipartite division into low- and high frequency effects known from rodent sensory cortices, we also report linear pupil-brain coupling in a third intermediate alpha/beta-range frequency band. Specifically, a negative correlation, localised to somatomotor cortices and specific to alpha-band (8-12 Hz) activity, peaked earlier than a positive correlation that localised to occipital visual cortices and encompassed alpha and beta bands (see Figures 3C & 4D). Of all effects found, the effect in visual cortices had the shortest lag with respect to pupil dilation (~200msec). This short-lag positive correlation may stem from interactions

between the locus coeruleus and regions of the thalamus (McCormick, 1989; McCormick et al., 1991; Stitt et al., 2018), in particular those implicated in the generation of occipital alpha-band activity via thalamo-cortical loops (e.g. the Pulvinar; Saalmann et al., 2012). Put differently, a more indirect multi-synaptic route to neuromodulation of occipital alpha activity may delay the lag relative to other effects and put it closer to the time of maximum pupil dilation. Another, not mutually exclusive, possibility is that the arousal-dependent low-lag alpha-/beta-band activity is linked to the activity of a distinct subpopulation of GABAergic neurons in the locus coeruleus, that have been shown to exhibit a temporally closer relationship with pupil diameter compared to the noradrenergic neurons (Breton-Provencher & Sur, 2019).

Note that reported changes in alpha activity in the visual cortex cannot be trivially explained by pupil dilation altering retinal illumination. In that case, one would expect a negative relationship, i.e. decreasing alpha power with increasing pupil size (e.g., Donner & Siegel, 2011). Moreover, changes in alpha activity precede changes in pupil dilation, albeit with a shorter lag compared to activity in other frequency ranges (see Figure 3).

Correlations between cortical activity and pupil derivative largely mirrored the results from pupil diameter, although most links were found to be weaker in magnitude. This is likely due to the fact that both signals (pupil diameter as well as its temporal derivative) correlate to some extent with noradrenergic and cholinergic activity (Reimer et al., 2016).

An arousal-triggered cascade of activity in the resting human brain

The temporal profile of the diverse relationships between pupil-linked arousal and local cortical activity described above points towards a cascade of interrelated effects that the resting brain may undergo routinely. Several scenarios may produce such a cascade: First, episodes of arousal could arise seemingly at random, from underlying complex non-linear neural dynamics (Robinson et al., 2003; Kringelbach et al., 2020). Second, they could be coupled to other physiological processes, such as respiration (Kluger & Gross, 2020; Zelano et al., 2016). Similarly to pupil fluctuations (Bouma & Baghuis, 1971; Pomè et al., 2020; Turnbull et al., 2017), respiration has an average base rate of ~0.2 Hz (Del Negro et al., 2018; Fleming et al., 2011). Finally, as a third scenario, the observed cascade may imply a specific cognitive sequence: Most participants will likely have 'mind-wandered', or engaged in 'task-unrelated thoughts' (Groot et al., 2021) while resting. The sequence of effects we observe may therefore be a transient re-orientation towards the external sensory environment - a 'monitoring sweep'. This is consistent with the increase in pupil-linked arousal, and its inverse relationship with an early dip in low-frequency activity, i.e. the cessation of a globally synchronised cortical state, and agrees with a concurrent widespread increase in high-frequency cortical activity that signifies increased cortical excitability to process external sensory input (Hanslmayr et al., 2011; Harris & Thiele, 2011).

One aspect of mind-wandering is the retrieval of information from memory that heavily relies on theta rhythms in the hippocampus (Siegle & Wilson, 2014). Here, we localised the strongest pupil (and pupil-derivative) correlations with theta-range power fluctuations in bilateral hippocampal regions (see Figure 4D), suggesting an arousal modulation of memory processes. In support, hippocampal locus-coeruleus-noradrenaline modulation has long

520   been established in the rodent brain (Sara, 2009; Segal & Bloom, 1976b, 1976a), and McGinley et al. (2015) reported negative correlations between pupil-linked arousal and the occurrence of theta-range hippocampal ripple events. Moreover, a recent human neuroimaging study found that pupil-linked arousal modulated blood oxygenation in (para-) hippocampal regions during memory processes (Clewett et al., 2018).

525   Further on, the observed sequence of effects concludes with a peak in alpha/beta activity that is triggered roughly at the time of maximum correlations of arousal with low- and high-frequency activity. The increase in synchronised alpha/beta activity marks the return to a strong regime where external sensory (visual) input is attenuated, potentially indicating a return to an introspective focus. A strong alpha rhythm thereby imposes widespread bouts

530 of "pulsed inhibition" on visual cortex that suppress local gamma band activity (Jensen & Mazaheri, 2010; Haegens et al., 2011; for a review see Van Diepen et al., 2019). This relationship between alpha and gamma activity has been well described in non-human primate- (Spaak et al., 2012) and human visual cortices, with a role for cholinergic neuromodulation (Bauer et al., 2012). Corroborating this scenario, we also observe a

535 decrease in local gamma-band activity in visual cortices.

     The three scenarios described here are not mutually exclusive and may explain one and the same phenomenon from different perspectives. A pivotal manipulation to test these assumptions will be to contrast this cascade with couplings of pupil-linked arousal and cortical activity between different behavioural states.

540   **Nonlinear relationship between periodic cortical activity and pupil-linked arousal**

     Nonlinear relations between arousal and other factors are well-characterized, most prominently in the form of the Yerkes-Dodson law, which posits an inverted-U-shaped relation between arousal and cognitive performance (Yerkes & Dodson, 1908). More recently, an inverted-U relation has also been identified for pupil-linked arousal fluctuations and

545 detection performance and low-frequency (2-10 Hz) membrane potential fluctuations in primary auditory cortex (McGinley, David, et al., 2015). Here, we found a pronounced and comparable inverted-U-shaped relationship between power in the alpha- and beta-bands, and pupil and pupil-derivative alike: The inverted-U alpha-band nonlinearity was most strongly expressed in inferior parietal cortex, extending towards higher visual as well as

550 temporal regions in the beta band. In contrast to McGinley et al. (2015), we found no evidence for an U-shaped relationship between pupil and high-frequency (50-100 Hz) activity.

     The inverted-U shaped relationship between arousal and behavior may result from the differential activation of neuromodulatory receptors in particular noradrenergic adrenoreceptors with varying affinity (Berridge & Spencer, 2016) or nicotinic and muscarinic

555 acetylcholine receptors (Bentley et al., 2011). Similar mechanisms may therefore underlie the quadratic relationship observed here. More specifically, in the case of noradrenaline, the activation of high-affinity α2 receptors at medium arousal level may (directly or indirectly) lead to increased alpha- and beta-band activity. At elevated arousal levels, on the other hand, correspondingly high levels of noradrenaline lead to the activation of low-affinity α1 receptors,

560 possibly resulting in a decrease of alpha- and beta-band activity.

From a functional perspective, an inverted-U relationship seems paradoxical. One would expect alpha activity to approach a minimum at intermediate states of pupil-linked arousal, i.e., show a U-shaped relationship instead, to optimally facilitate the processing of external sensory input (Jensen & Mazaheri, 2010). However, our participants were resting, while neither receiving nor expecting visual input. In this situation an inward focus of attention, suppressing external visual information may have reversed the expected relationship (also see Hong et al., 2014).

Of note, we find evidence for both, linear positive and non-linear inverted-U shape relationships between alpha/beta activity and pupil-linked arousal. Cortical sources of the two effects coincide spatially (cp. Figures 4D & 5C). Previous research has suggested that parieto-occipital cortices harbor at least two distinct alpha rhythms (Barzegaran et al., 2017; Keitel & Gross, 2016) with separable characteristics (Benwell et al., 2019) that may serve different functions (Capilla et al., 2014; Sokoliuk et al., 2019). Therefore, spatially overlapping linear and non-linear effects may reflect distinct influences of arousal on the generative processes underlying separate alpha rhythms.

Arousal modulation of cortical excitation-inhibition ratio

In addition to interactions between pupil-linked arousal and band-limited activity, we show that pupil diameter correlates systematically with aperiodic brain activity. Recent theoretical and empirical evidence suggest that the slope of the aperiodic component of neural power spectra, the spectral exponent, reflects the ratio between excitatory and inhibitory neuronal processes in the underlying neuronal circuits (Colombo et al., 2019; Gao et al., 2017).

Here, we report a negative linear relation between pupil diameter and the spectral exponent. In other words, periods of elevated arousal were accompanied by a flattening of the power spectrum. In simulated neural networks, a shallower slope of the aperiodic component is related to increased excitation relative to inhibition (Gao et al., 2017; Trakoshis et al., 2020). Thus, increased arousal (i.e., dilated pupils) may co-occur with increased E/I in distributed neuronal circuits, whereas periods of low arousal (i.e., constricted pupils) would indicate states of relatively decreased E/I. This interpretation is also consistent with previous findings arguing that noradrenaline increases cortical E/I (Pfeffer et al., 2018, 2020), possibly through a decrease in inhibition (Froemke, 2015; Martins & Froemke, 2015). We found this relationship of E/I and pupil-linked arousal in the majority of regions, thus suggesting a global and uniform effect. In sum, pupil-linked arousal may provide a window into yet another cortical circuit property, cortical E/I, relevant for cognitive computation (Cavanagh et al., 2020; Kosciessa et al., 2021; Lam et al., 2017; B. K. Murphy & Miller, 2003; Pettine et al., 2021; Pfeffer et al., 2020).

Conclusion

We exhaustively mapped how fluctuations in pupil-linked arousal and cortical activity co-vary in the human brain at rest. The diverse linear and non-linear relationships we describe in our data suggest profound influences of arousal on cortical states and challenge a global and unspecific role for arousal neuromodulation in cortical function. Instead, our results support

the view of specific neuromodulatory influences that differ depending on the targeted cortical region, can express in varying frequencies of band-limited activity - or as broadband effects, and vary in their timing. The present data provide the basis for studying how these influences change under cognitive task demands and when engaging in different behaviours. In fact, 605 neuromodulatory effects are often neglected when studying the cortical population activity underlying cognitive function - given a central role for arousal, our results argue for including it as a covariate in future studies.

Methods

610   **Participants and Data Acquisition (Hamburg)**

30 healthy human participants (16 women, mean age 26.7, range 20-36) participated in the study after informed consent. All included participants were healthy, with no current or previous diagnosis of psychiatric or neurological disorder (full list of exclusion criteria can be found in Pfeffer et al., 2020). The study was approved by the Ethics Committee of the Medical

615   Association Hamburg. Two participants were excluded from analyses, one due to excessive MEG artifacts, the other due to not completing all recording sessions (see below). Thus, we report results from N=28 participants of which 15 were women. The present dataset is a subset of a larger data set that entailed selective pharmacological manipulations (Pfeffer et al., 2018, 2020). For the present article, only data from the placebo / resting state condition

620   was analyzed.

Each participant completed three experimental sessions (scheduled at least 2 weeks apart), consisting of drug or placebo intake at two time points, a waiting period of 3 hours, and an MEG recording session. During the recordings, participants were seated on a chair inside a magnetically shielded chamber. Each recording session consisted of two resting

625   state measurements as well as four blocks consisting of two variants of a behavioral task. Each block was 10 minutes long and followed by a short break of variable duration.

Brain activity was recorded using a whole-head CTF 275 MEG system (CTF Systems, Inc., Canada) at a sampling rate of 1200 Hz. Eye movements and pupil diameter were recorded with an MEG-compatible EyeLink 1000 Long Range Mount system (SR Research, Osgoode,

630   ON, Canada) and electrocardiogram (ECG) as well as vertical, horizontal and radial EOG was acquired using Ag/AgCl electrodes. During all recordings, the participants were instructed to keep their eyes open and fixate a green fixation dot in the center of a gray background of constant luminance, which was projected onto a screen from outside the magnetically shielded recording chamber at a refresh rate of 60 Hz.

Participants and Data Acquisition (Münster)

Forty right-handed volunteers (21 women, age 25.1 ± 2.7 y (Mean ± SD), range 21-32 y) participated in the study. All participants reported having no respiratory or neurological disease and gave written informed consent prior to all experimental procedures. The study

640   was approved by the local ethics committee of the University of Muenster.

Participants were seated upright in a magnetically shielded room while we simultaneously recorded 5 minutes of MEG and eye tracking data. MEG data was acquired using a 275 channel whole-head system (OMEGA 275, VSM Medtech Ltd., Vancouver, Canada) at a sampling frequency of 600 Hz. Pupil area of the right eye was recorded at a sampling rate of

645   1000 Hz using an EyeLink 1000 plus eye tracker (SR Research). During recording, participants were to keep their eyes on a fixation cross centred on a projector screen placed in front of them. To minimise head movement, participants' heads were stabilised with cotton pads inside the MEG helmet.

Participants and Data Acquisition (Glasgow)

650   MEG resting-state data were acquired for 24 healthy, right-handed participants (9 women; age 23.5 ± 5.5 y (Mean ± SD), range 18-39 y). The study was approved by the local ethics committee (University of Glasgow, College of Science and Engineering; approval number 300140078). Participants gave written informed consent prior to testing.

655   Recordings were obtained with a 248-magnetometer whole-head MEG system (MAGNES 3600 WH, 4-D Neuroimaging), set to a sampling rate of 1017 Hz, and an EyeLink 1000 eye tracker (SR Research), sampling the pupil area of the left eye at 1000 Hz, both situated in a magnetically shielded room (Vacuumschmelze). Head position was measured at the start and end of the recording. To this end, a set of five coils were placed on the head of the participant.

660   Coil positions and head shape were digitised using a FASTRAK stylus (Polhemus Inc., VT, US). During recordings, participants sat upright and fixated on a green point (measure) projected centrally on a screen with a DLP projector (60 Hz refresh rate) for a minimum of 7 minutes (420 s).

     Resting state recordings were part of a study into audio-visual speech processing that

665   was recorded in two sessions. Audio-visual data have been published elsewhere (Keitel et al., 2018, 2020). In this study, the resting state block was recorded as the last block of the second session after 7 - 9 experimental blocks, or more than 65 min into the session. Note that the resting state data have not been reported in any of the previous publications.

     Data of two participants were excluded from analysis after screening recordings, one due

670   to strong intermittent environmental artifacts in their MEG recordings and another due to a technical fault in tracking the pupil.

MEG Preprocessing (Hamburg)

     The sensor-level MEG signal was cleaned of extra-cranial artifacts using semi-automatic

675   artifact detection routines implemented in the FieldTrip toolbox (Oostenveld et al., 2011). Broken and noisy recordings channels were identified through visual inspection and were marked as artifactual across all channels (+/- 500 ms). Subsequently, the data were downsampled to 400 Hz split into low ([0.5-2]-40 Hz; the lower cutoff was variable across (but identical within) subjects at 0.5, 1 or 2 Hz) and high (>40 Hz) frequency components,

680   using a 4th order Butterworth filter. The separate data segments were independently submitted to Independent Component Analysis (ICA; Hyvarinen, 1999). The split into low- and high-frequency data aimed to facilitate the detection of sustained muscle artifacts especially present in the high-frequency range. Artifactual independent components were identified through visual inspection of their topography and power spectra. On average, 23 ±

685   14 components (Mean ± SD) were subtracted from the raw data (placebo condition only). After artifactual components were removed, the low- and high-frequency segments were combined into one single data set.

MEG Preprocessing (Münster)

690   MEG preprocessing was done in Fieldtrip (Oostenveld et al., 2011) for MATLAB (The MathWorks Inc). First, noisy recording channels were identified by visual inspection and

695 rejected from further analyses. Next, we adapted the synthetic gradiometer order to the third order for better MEG noise balancing (ft_denoise_synthetic). Power line artefacts were removed using a discrete Fourier transform (DFT) filter on the line frequency of 50 Hz and all its harmonics (including spectrum interpolation; *dftfilter* argument in *ft_preprocessing*). Next, we applied independent component analysis (ICA) on the filtered data to capture eye blinks and cardiac artefacts (*ft_componentanalysis* with 32 extracted components). On average, artefacts were identified in 2.35 ± 0.83 components (M ± SD) per participant and removed from the data. Finally, cleaned MEG data were downsampled to 400 Hz.

MEG Preprocessing (Glasgow)

For MEG and eye tracking data preprocessing and analysis, we used Matlab (The MathWorks Inc), involving in-house MATLAB routines and the FieldTrip toolbox (Oostenveld et al., 2011). The MEG signal was resampled to the sampling rate of the eye tracker (1000
705 Hz) and denoised using in-built MEG reference sensors. We rejected between 2–27 noisy MEG channels (median = 7.5 ± 4 absolute deviation) by visual inspection using FieldTrip's *ft_rejectvisual*. Continuous MEG recordings were then screened for noisy segments. High-pass filtered (2 Hz cutoff, 4th-order Butterworth, forward-reverse two-pass) MEG time series were subjected to an independent component analysis (ICA; Hyvarinen, 1999). Prior to the
710 ICA, a principal component analysis (PCA) projected MEG data into a 32-dimensional subspace. For each participant, we visually identified 2–4 components (median = 3) capturing eye (blinks, movements) and heart-beat artefacts. Finally, resampled continuous MEG recordings were projected through the subspace spanned by the remaining components.

MEG and eye tracker time axes were realigned by cross-correlating traces of horizontal
715 and vertical eye movements that were simultaneously recorded by the eye tracker and the MEG. The lag corresponding to the maximum of the cross-correlation functions of horizontal and vertical eye movement traces was used as offset for the realignment.

Pupil preprocessing (all sites)

720 Pupil area traces were converted to pupil diameter to linearise our measure of pupil size. Using additional Matlab routines (available from https://github.com/anne-urai/pupil_preprocessing_tutorial, as used in (Urai et al., 2017), blinks were identified by an automatic (and visually validated) procedure and linearly interpolated. In a second pass, a similar procedure was used on smaller eye tracker artifacts (spikes, jumps). Next, canonical
725 responses to blinks and saccades were estimated and then removed from pupil time series (Hoeks & Levelt, 1993; Knapen et al., 2016; Wierda et al., 2012). To that end, pupil time series were band-pass filtered (pass band: 0.005 - 2 Hz, 2nd-order Butterworth, forward-reverse two-pass), then downsampled to a sampling rate of 400 Hz. Finally, we computed the first-order derivative of each band-pass filtered and down-sampled pupil trace.

Spectral decomposition of pupil time series (all sites)

The spectral content of pupil traces was evaluated using the multi-taper approach as implemented in FieldTrip. We used the default settings, except for a spectral smoothing of 0.035 Hz, and zero-padding time series to 2^{19} (minimum power of 2 that exceeded the length of any pupil trace) to unify frequency resolutions across centres and traces of different lengths. Power spectra were calculated on standardized (z-scored) pupil traces for the 0.005-2 Hz range, sampled in logarithmic steps.

Resulting power spectra were also subjected to the spectral parameterization toolbox ('fooof'; Donoghue et al., 2020) to extract Hippus peak frequency as well as the exponent describing the spectral slope as a means to characterise the aperiodic activity. We used the 'fooof' MATLAB wrapper with the default settings, except setting the maximum number of peaks ('max_n_peaks') to 5, and the peak detection threshold ('peak_threshold') to 0.5.

MEG spectral analysis (all sites)

In order to derive spectral estimates from the sensor-level data, we followed an approach outlined previously (Hipp et al., 2012), based on Morelet's wavelets (Tallon-Baudry & Bertrand, 1999):

$$w(t,f) \; = \; (\sigma_t \sqrt{\pi})^{-1/2} e^{-\frac{t^2}{2\sigma_t^2}} e^{-i2\pi ft}$$

(Eq. 1)

Following Hipp et al. (2012), we derived spectral estimates for 25 logarithmically spaced center frequencies, covering a broad frequency range from 2 Hz to 128 Hz. Spectral estimates were derived for successive temporal windows with an overlap of 80%. Windows that contained periods marked as artifactual (see Preprocessing) were discarded from all further analyses. From the spectral estimates, sensor-level power envelopes were obtained through:

$$Y(t,f) = \; |\, X_{sens}(t,f)\,|^2$$

(Eq. 2)

where X_{sens} denotes the complex spectral estimates for segment t and frequency f.

MEG source reconstruction (all sites)

Accurate source models were generated using the FieldTrip interface to SPM8 (Litvak et al., 2011), and the Freesurfer toolbox (Fischl, 2012). Source models were based on individual T1-weighted structural magnetic resonance images (MRIs), acquired on a 3T Siemens Tim Trio system with a 12-channel head coil (Glasgow), a Siemens 3T Prisma with a 20-channel head coil (Münster), and a Siemens Magnetom Trio (Hamburg). Anatomical scans were coregistered with the MEG coordinate system using a semiautomatic procedure (Gross et al., 2013), then segmented and linearly normalised to a template brain (MNI space). We used a single shell as the volume conduction model (Nolte, 2003).

770   We projected sensor-level data into source space using frequency-specific DICS (Dynamic Imaging of Coherent Sources) beamformers (Gross et al., 2001) with a regularisation parameter of 5% and optimal dipole orientation (singular value decomposition method). As the source model we used a volumetric grid. Grid points had a spacing of 5 mm, resulting in 8799 dipoles covering the whole brain. Source level power envelopes were computed from
775   the source-projected complex signal:

$$X_{src}(t,f) \ = \ | \, A(r,f) * X_{sens}(t,f) \, |^2$$

(Eq. 3)

        where A denotes the spatial filter, X_{src} the source-level power envelopes and X_{sens} the sensor-level analytic signal. The 5 mm spacing was chosen so as to allow for an integer
780   spatial downsampling of the Brainnetome atlas (Fan et al., 2016) which we used to map cortical (and subcortical) regions into 246 distinct areas.

Mutual information between pupil and spectral activity in sensor space

        To quantify global pupil-power coupling in sensor space we calculated mutual information (MI) between frequency-specific power envelopes and pupil time traces. To this end, pupil
785   time series were downsampled to the temporal resolution of the respective power envelope, and segments containing artifacts in power envelopes were also removed from the pupil traces. After normalising both time series by means of a Gaussian-copula based approach (Ince et al., 2017), mutual information was computed per frequency and MEG sensor.

        Observed MI was evaluated against permutation distributions of surrogate MIs based on
790   randomly time-shifting pupil traces 200 times, while avoiding a window of +/- 10 sec around the match of both original time series. Using the log-transformed observed MI, and the mean and standard deviation of equally log-transformed permutation distributions we computed z-values for each frequency and sensor. In a final step, these z-values were pooled across sensors and their mean across participants tested against zero, for each recording site
795   separately. Test results are reported for uncorrected and FDR-corrected thresholds in Figure 2.

Cross-correlation analysis (sensor space)

        In order to allow for the comparison across MEG recording sites (with different sensor layouts), we first ordered the recording channels into 39 equally sized, non-overlapping
800   spatial bins, sorted from anterior to posterior channels. We next averaged the cross correlograms per bin and additionally averaged the cross correlations across three distinct frequency ranges: 2-4 Hz, 8-16 Hz and 64- 128 Hz. Due to the varying width of the wavelets for each center frequency, the temporal resolution, and, therefore, the number of samples, varied across frequencies. Hence, prior to averaging, we linearly interpolated the frequency-
805   specific cross-correlograms such that the number of samples was equal to the number of samples of the highest frequency of each band (i.e., 4 Hz, 16 Hz and 128 Hz).

Source-level power spectra and spectral parametrization

810   In order to correlate fluctuations in pupil diameter (and its temporal derivative) with fluctuations in the slope of the aperiodic component of the power spectra (henceforth referred to as the 'spectral exponent') we first computed source-level power spectra. To this end, we source-projected the broad-band signal using linear beamforming (LCMV; Van Veen et al., 1997) instead of DICS (Gross et al., 2001). The procedure was similar to DICS, with the difference that the spatial filters were obtained using the cross-spectral density matrix

815   averaged across all 25 frequency-bands of interest. In keeping with the previous analysis, we used a regularization parameter of 5%.

     Next, we divided the resulting source-level broad-band signal into non-overlapping temporal segments of 2s length. For each segment, power spectra were computed using MATLAB's 'pwelch' function, ranging from 2 Hz to 128 Hz, with a frequency resolution of 0.5

820   Hz. In addition, for each temporal segment, the mean pupil diameter as well as the mean of the pupil-derivative were computed.

     To separate aperiodic and periodic components of the power spectra, we utilized the spectral parameterization toolbox 'fooof' for Python 3.7 (Donoghue et al., 2020). The algorithm iteratively fits neuronal power spectra as the sum of a Lorentzian function and a

825   variable number of Gaussians of varying width, height and mean (corresponding to the center frequency). To constrain the fitting procedure, we limited the number of periodic components to 6 and defined a minimum peak height of 0.05, with the minimum and maximum bandwidth of each peak set to 1 Hz and 8 Hz. The knee of the Lorentzian was set to zero.

     The algorithm was applied to the power spectra of each individual temporal segment (see

830   above), which provided time-resolved estimates of periodic and aperiodic components. While the slope of the aperiodic component is relatively stable for lower frequencies up until ~40-50 Hz, the rate at which power changes can differ substantially for higher frequencies. This means that fitting power spectra uniformly across the entire frequency range may result in poorer fits and, consequently, misleading parameter estimates. Thus, and largely consistent

835   with previous work (Pfeffer et al., 2020; Waschke et al., 2021), neuronal power spectra were fitted in the frequency range from 3 Hz to 40 Hz.

Polynomial model

     We quantified non-linear components of the relation between pupil diameter (and pupil derivative) and band-limited activity fluctuations in spectral power. To this end, we sorted the

840   aforementioned temporal segments (see section above: 'Source-level power spectra and spectral parametrization') based on pupil diameter (or pupil derivative) into 14 non-overlapping and equidistant bins. Next, we computed the average power for each of the bins and normalized (z-scored) the obtained values. Quadratic relationships were quantified by fitting the power vs. pupil (or pupil-derivative) function with a polynomial of degree 2,

845   separately for each of the 25 MEG center frequencies:

$$P_{src}(s,f) = \beta_0 + \beta_1 x + \beta_2 x^2$$

(Eq. 4)

where s denotes the source location (with s being 1 to 246 regions of the Brainnetome atlas),
850   f is the MEG center frequency (from 2 to 128 Hz, in 25 logarithmic steps) and x is the mean pupil diameter for each pupil bin. The coefficients for the quadratic and the linear component are denoted with β_2 and β_1, respectively, and β_0 describes the offset. The polynomial was fitted by minimizing the sum of the squares of the difference between the polynomial model and the power vs. pupil (pupil-derivative) functions (as implemented in MATLAB's 'polyfit'
855   function).

References

Aston-Jones, G., & Cohen, J. D. (2005). AN INTEGRATIVE THEORY OF LOCUS COERULEUS-NOREPINEPHRINE FUNCTION: Adaptive Gain and Optimal Performance. *Annual Review of Neuroscience*, *28*(1), 403–450. https://doi.org/10.1146/annurev.neuro.28.061604.135709

860   Barzegaran, E., Vildavski, V. Y., & Knyazeva, M. G. (2017). Fine Structure of Posterior Alpha Rhythm in Human EEG: Frequency Components, Their Cortical Sources, and Temporal Behavior. *Scientific Reports*, *7*(1), 8249. https://doi.org/10.1038/s41598-017-08421-z

Bauer, M., Kluge, C., Bach, D., Bradbury, D., Heinze, H. J., Dolan, R. J., & Driver, J. (2012). Cholinergic Enhancement of Visual Attention and Neural Oscillations in the Human Brain. *Current Biology*, *22*(5), 397–402.
865   https://doi.org/10.1016/j.cub.2012.01.022

Beatty, J. (1982). Task-evoked pupillary responses, processing load, and the structure of processing resources. *Psychological Bulletin*, *91*(2), 276–292. https://doi.org/10.1037/0033-2909.91.2.276

Bentley, P., Driver, J., & Dolan, R. J. (2011). Cholinergic modulation of cognition: Insights from human pharmacological functional neuroimaging. *Progress in Neurobiology*, *94*(4), 360–388.
870   https://doi.org/10.1016/j.pneurobio.2011.06.002

Benwell, C. S. Y., London, R. E., Tagliabue, C. F., Veniero, D., Gross, J., Keitel, C., & Thut, G. (2019). Frequency and power of human alpha oscillations drift systematically with time-on-task. *NeuroImage*, *192*, 101–114. https://doi.org/10.1016/j.neuroimage.2019.02.067

Berridge, C. W., & Spencer, R. C. (2016). Differential cognitive actions of norepinephrine a2 and a1 receptor
875   signaling in the prefrontal cortex. *Brain Research*, *1641*, 189–196. https://doi.org/10.1016/j.brainres.2015.11.024

Bouma, H., & Baghuis, L. C. J. (1971). Hippus of the pupil: Periods of slow oscillations of unknown origin. *Vision Research*, *11*(11), 1345–1351. https://doi.org/10.1016/0042-6989(71)90016-2

Bradshaw, J. (1967). Pupil Size as a Measure of Arousal during Information Processing. *Nature*, *216*(5114), 515–
880   516. https://doi.org/10.1038/216515a0

Brand, A., Allen, L., Altman, M., Hlava, M., & Scott, J. (2015). Beyond authorship: Attribution, contribution, collaboration, and credit. *Learned Publishing*, *28*(2), 151–155. https://doi.org/10.1087/20150211

Breton-Provencher, V., & Sur, M. (2019). Active control of arousal by a locus coeruleus GABAergic circuit. *Nature Neuroscience*, *22*(2), 218–228. https://doi.org/10.1038/s41593-018-0305-z

885   Burt, J. B., Demirtaş, M., Eckner, W. J., Navejar, N. M., Ji, J. L., Martin, W. J., Bernacchia, A., Anticevic, A., & Murray, J. D. (2018). Hierarchy of transcriptomic specialization across human cortex captured by structural neuroimaging topography. *Nature Neuroscience*, *21*(9), 1251–1259. https://doi.org/10.1038/s41593-018-0195-0

Busse, L., Cardin, J. A., Chiappe, M. E., Halassa, M. M., McGinley, M. J., Yamashita, T., & Saleem, A. B. (2017).
890   Sensation during Active Behaviors. *Journal of Neuroscience*, *37*(45), 10826–10834.

Capilla, A., Schoffelen, J.-M., Paterson, G., Thut, G., & Gross, J. (2014). Dissociated α-Band Modulations in the Dorsal and Ventral Visual Pathways in Visuospatial Attention and Perception. *Cerebral Cortex*, *24*(2), 550–561. https://doi.org/10.1093/cercor/bhs343

Cavanagh, S. E., Lam, N. H., Murray, J. D., Hunt, L. T., & Kennerley, S. W. (2020). A circuit mechanism for decision-
895   making biases and NMDA receptor hypofunction. *ELife*, *9*, e53664. https://doi.org/10.7554/eLife.53664

Chandler, D. J., Jensen, P., McCall, J. G., Pickering, A. E., Schwarz, L. A., & Totah, N. K. (2019). Redefining Noradrenergic Neuromodulation of Behavior: Impacts of a Modular Locus Coeruleus Architecture. *Journal of Neuroscience*, *39*(42), 8239–8249.

Chandler, D. J., Lamperski, C. S., & Waterhouse, B. D. (2013). Identification and distribution of projections from
900   monoaminergic and cholinergic nuclei to functionally differentiated subregions of prefrontal cortex. *Brain Research*, *1522*, 38–58. https://doi.org/10.1016/j.brainres.2013.04.057

Clewett, D. V., Huang, R., Velasco, R., Lee, T.-H., & Mather, M. (2018). Locus Coeruleus Activity Strengthens Prioritized Memories Under Arousal. *Journal of Neuroscience*, *38*(6), 1558–1574. https://doi.org/10.1523/JNEUROSCI.2097-17.2017

905   Colombo, M. A., Napolitani, M., Boly, M., Gosseries, O., Casarotto, S., Rosanova, M., Brichant, J.-F., Boveroux, P., Rex, S., Laureys, S., Massimini, M., Chieregato, A., & Sarasso, S. (2019). The spectral exponent of the resting EEG indexes the presence of consciousness during unresponsiveness induced by propofol, xenon, and

ketamine. *NeuroImage*, *189*, 631–644. https://doi.org/10.1016/j.neuroimage.2019.01.024

910 Crochet, S., & Petersen, C. C. H. (2006). Correlating whisker behavior with membrane potential in barrel cortex of awake mice. *Nature Neuroscience*, *9*(5), 608–610. https://doi.org/10.1038/nn1690

de Gee, J. W., Colizoli, O., Kloosterman, N. A., Knapen, T., Nieuwenhuis, S., & Donner, T. H. (2017). Dynamic modulation of decision biases by brainstem arousal systems. *ELife*, *6*, e23232. https://doi.org/10.7554/eLife.23232

Del Negro, C. A., Funk, G. D., & Feldman, J. L. (2018). Breathing matters. *Nature Reviews Neuroscience*, *19*(6), 915 351–367. https://doi.org/10.1038/s41583-018-0003-6

Donner, T. H., & Siegel, M. (2011). A framework for local cortical oscillation patterns. *Trends in Cognitive Sciences*, *15*(5), 191–199. https://doi.org/10.1016/j.tics.2011.03.007

Donoghue, T., Haller, M., Peterson, E. J., Varma, P., Sebastian, P., Gao, R., Noto, T., Lara, A. H., Wallis, J. D., Knight, R. T., Shestyuk, A., & Voytek, B. (2020). Parameterizing neural power spectra into periodic and aperiodic 920 components. *Nature Neuroscience*, *23*(12), 1655–1665. https://doi.org/10.1038/s41593-020-00744-x

Fan, L., Li, H., Zhuo, J., Zhang, Y., Wang, J., Chen, L., Yang, Z., Chu, C., Xie, S., Laird, A. R., Fox, P. T., Eickhoff, S. B., Yu, C., & Jiang, T. (2016). The Human Brainnetome Atlas: A New Brain Atlas Based on Connectional Architecture. *Cerebral Cortex*, *26*(8), 3508–3526. https://doi.org/10.1093/cercor/bhw157

Fischl, B. (2012). FreeSurfer. *NeuroImage*, *62*(2), 774–781. https://doi.org/10.1016/j.neuroimage.2012.01.021

925 Fleming, S., Thompson, M., Stevens, R., Heneghan, C., Plüddemann, A., Maconochie, I., Tarassenko, L., & Mant, D. (2011). Normal ranges of heart rate and respiratory rate in children from birth to 18 years of age: A systematic review of observational studies. *The Lancet*, *377*(9770), 1011–1018. https://doi.org/10.1016/S0140-6736(10)62226-X

Froemke, R. C. (2015). Plasticity of Cortical Excitatory-Inhibitory Balance. *Annual Review of Neuroscience*, *38*(1), 930 195–219. https://doi.org/10.1146/annurev-neuro-071714-034002

Fu, Y., Tucciarone, J. M., Espinosa, J. S., Sheng, N., Darcy, D. P., Nicoll, R. A., Huang, Z. J., & Stryker, M. P. (2014). A Cortical Circuit for Gain Control by Behavioral State. *Cell*, *156*(6), 1139–1152. https://doi.org/10.1016/j.cell.2014.01.050

Gao, R., Peterson, E. J., & Voytek, B. (2017). Inferring synaptic excitation/inhibition balance from field potentials. 935 *NeuroImage*, *158*, 70–78. https://doi.org/10.1016/j.neuroimage.2017.06.078

Groot, J. M., Boayue, N. M., Csifcsák, G., Boekel, W., Huster, R., Forstmann, B. U., & Mittner, M. (2021). Probing the neural signature of mind wandering with simultaneous fMRI-EEG and pupillometry. *NeuroImage*, *224*, 117412. https://doi.org/10.1016/j.neuroimage.2020.117412

Gross, J., Baillet, S., Barnes, G. R., Henson, R. N., Hillebrand, A., Jensen, O., Jerbi, K., Litvak, V., Maess, B., 940 Oostenveld, R., Parkkonen, L., Taylor, J. R., van Wassenhove, V., Wibral, M., & Schoffelen, J.-M. (2013). Good practice for conducting and reporting MEG research. *NeuroImage*, *65*, 349–363. https://doi.org/10.1016/j.neuroimage.2012.10.001

Gross, J., Kujala, J., Hämäläinen, M., Timmermann, L., Schnitzler, A., & Salmelin, R. (2001). Dynamic imaging of coherent sources: Studying neural interactions in the human brain. *Proceedings of the National Academy of* 945 *Sciences*, *98*(2), 694–699. https://doi.org/10.1073/pnas.98.2.694

Haegens, S., Nácher, V., Luna, R., Romo, R., & Jensen, O. (2011). α-Oscillations in the monkey sensorimotor network influence discrimination performance by rhythmical inhibition of neuronal spiking. *Proceedings of the National Academy of Sciences*, *108*(48), 19377–19382. https://doi.org/10.1073/pnas.1117190108

Hanslmayr, S., Gross, J., Klimesch, W., & Shapiro, K. L. (2011). The role of alpha oscillations in temporal attention. 950 *Brain Research Reviews*, *67*(1), 331–343. https://doi.org/10.1016/j.brainresrev.2011.04.002

Harris, K. D., & Thiele, A. (2011). Cortical state and attention. *Nature Reviews Neuroscience*, *12*(9), 509–523. https://doi.org/10.1038/nrn3084

Hasselmo, M. E. (1995). Neuromodulation and cortical function: Modeling the physiological basis of behavior. *Behavioural Brain Research*, *67*(1), 1–27. https://doi.org/10.1016/0166-4328(94)00113-T

955 He, B. J., & Zempel, J. M. (2013). Average Is Optimal: An Inverted-U Relationship between Trial-to-Trial Brain Activity and Behavioral Performance. *PLOS Computational Biology*, *9*(11), e1003348. https://doi.org/10.1371/journal.pcbi.1003348

Hess, E. H., & Polt, J. M. (1964). Pupil Size in Relation to Mental Activity during Simple Problem-Solving. *Science*, *143*(3611), 1190–1192. https://doi.org/10.1126/science.143.3611.1190

960 Hipp, J. F., Hawellek, D. J., Corbetta, M., Siegel, M., & Engel, A. K. (2012). Large-scale cortical correlation structure of spontaneous oscillatory activity. *Nature Neuroscience*, *15*(6), 884–890. https://doi.org/10.1038/nn.3101

Hoeks, B., & Levelt, W. J. M. (1993). Pupillary dilation as a measure of attention: A quantitative system analysis. *Behavior Research Methods, Instruments, & Computers*, *25*(1), 16–26. https://doi.org/10.3758/BF03204445

Hong, L., Walz, J. M., & Sajda, P. (2014). Your Eyes Give You Away: Prestimulus Changes in Pupil Diameter 965 Correlate with Poststimulus Task-Related EEG Dynamics. *PLOS ONE*, *9*(3), e91321. https://doi.org/10.1371/journal.pone.0091321

Hyvarinen, A. (1999). Fast and robust fixed-point algorithms for independent component analysis. *IEEE Transactions on Neural Networks*, *10*(3), 626–634. https://doi.org/10.1109/72.761722

Ince, R. A. A., Giordano, B. L., Kayser, C., Rousselet, G. A., Gross, J., & Schyns, P. G. (2017). A statistical framework 970 for neuroimaging data analysis based on mutual information estimated via a gaussian copula. *Human Brain Mapping*, *38*(3), 1541–1573. https://doi.org/10.1002/hbm.23471

Jensen, O., & Mazaheri, A. (2010). Shaping Functional Architecture by Oscillatory Alpha Activity: Gating by Inhibition. *Frontiers in Human Neuroscience*, *4*. https://doi.org/10.3389/fnhum.2010.00186

Joshi, S., & Gold, J. I. (2020). Context-Dependent Relationships between Locus Coeruleus Firing Patterns and Coordinated Neural Activity in the Anterior Cingulate Cortex. *BioRxiv*, 2020.09.26.314831. https://doi.org/10.1101/2020.09.26.314831

Joshi, S., Li, Y., Kalwani, R. M., & Gold, J. I. (2016). Relationships between Pupil Diameter and Neuronal Activity in the Locus Coeruleus, Colliculi, and Cingulate Cortex. *Neuron*, *89*(1), 221–234. https://doi.org/10.1016/j.neuron.2015.11.028

Kahneman, D., & Beatty, J. (1966). Pupil Diameter and Load on Memory. *Science*, *154*(3756), 1583–1585. https://doi.org/10.1126/science.154.3756.1583

Kahneman, D., Beatty, J., & Pollack, I. (1967). Perceptual Deficit during a Mental Task. *Science*, *157*(3785), 218–219. https://doi.org/10.1126/science.157.3785.218

Keitel, A., & Gross, J. (2016). Individual Human Brain Areas Can Be Identified from Their Characteristic Spectral Activation Fingerprints. *PLOS Biology*, *14*(6), e1002498. https://doi.org/10.1371/journal.pbio.1002498

Keitel, A., Gross, J., & Kayser, C. (2018). Perceptually relevant speech tracking in auditory and motor cortex reflects distinct linguistic features. *PLOS Biology*, *16*(3), e2004473. https://doi.org/10.1371/journal.pbio.2004473

Keitel, A., Gross, J., & Kayser, C. (2020). Shared and modality-specific brain regions that mediate auditory and visual word comprehension. *ELife*, *9*, e56972. https://doi.org/10.7554/eLife.56972

Kluger, D. S., & Gross, J. (2020). Respiration modulates oscillatory neural network activity at rest. *BioRxiv*, 2020.04.23.057216. https://doi.org/10.1101/2020.04.23.057216

Knapen, T., de Gee, J. W., Brascamp, J., Nuiten, S., Hoppenbrouwers, S., & Theeuwes, J. (2016). Cognitive and Ocular Factors Jointly Determine Pupil Responses under Equiluminance. *PLOS ONE*, *11*(5), e0155574. https://doi.org/10.1371/journal.pone.0155574

Korn, C. W., & Bach, D. R. (2016). A solid frame for the window on cognition: Modeling event-related pupil responses. *Journal of Vision*, *16*(3), 28–28. https://doi.org/10.1167/16.3.28

Kosciessa, J. Q., Lindenberger, U., & Garrett, D. D. (2021). Thalamocortical excitability modulation guides human perception under uncertainty. *Nature Communications*, *12*(1), 2430. https://doi.org/10.1038/s41467-021-22511-7

Lam, N. H., Borduqui, T., Hallak, J., Roque, A. C., Anticevic, A., Krystal, J. H., Wang, X.-J., & Murray, J. D. (2017). Effects of Altered Excitation-Inhibition Balance on Decision Making in a Cortical Circuit Model. *BioRxiv*, 100347. https://doi.org/10.1101/100347

Lee, S.-H., & Dan, Y. (2012). Neuromodulation of Brain States. *Neuron*, *76*(1), 209–222. https://doi.org/10.1016/j.neuron.2012.09.012

Leopold, D. A., Murayama, Y., & Logothetis, N. K. (2003). Very Slow Activity Fluctuations in Monkey Visual Cortex: Implications for Functional Brain Imaging. *Cerebral Cortex*, *13*(4), 422–433. https://doi.org/10.1093/cercor/13.4.422

Litvak, V., Mattout, J., Kiebel, S., Phillips, C., Henson, R., Kilner, J., Barnes, G., Oostenveld, R., Daunizeau, J., Flandin, G., Penny, W., & Friston, K. (2011, March 6). *EEG and MEG Data Analysis in SPM8* [Research Article]. Computational Intelligence and Neuroscience; Hindawi. https://doi.org/10.1155/2011/852961

Martins, A. R. O., & Froemke, R. C. (2015). Coordinated forms of noradrenergic plasticity in the locus coeruleus and primary auditory cortex. *Nature Neuroscience*, *18*(10), 1483–1492. https://doi.org/10.1038/nn.4090

McCormick, D. A. (1989). Cholinergic and noradrenergic modulation of thalamocortical processing. *Trends in Neurosciences*, *12*(6), 215–221. https://doi.org/10.1016/0166-2236(89)90125-2

McCormick, D. A., Pape, H.-C., & Williamson, A. (1991). Chapter 22 - Actions of norepinephrine in the cerebral cortex and thalamus: Implications for function of the central noradrenergic system. In C. D. Barnes & O. Pompeiano (Eds.), *Progress in Brain Research* (Vol. 88, pp. 293–305). Elsevier. https://doi.org/10.1016/S0079-6123(08)63817-0

McGinley, M. J., David, S. V., & McCormick, D. A. (2015). Cortical Membrane Potential Signature of Optimal States for Sensory Signal Detection. *Neuron*, *87*(1), 179–192. https://doi.org/10.1016/j.neuron.2015.05.038

McGinley, M. J., Vinck, M., Reimer, J., Batista-Brito, R., Zagha, E., Cadwell, C. R., Tolias, A. S., Cardin, J. A., & McCormick, D. A. (2015). Waking State: Rapid Variations Modulate Neural and Behavioral Responses. *Neuron*, *87*(6), 1143–1161. https://doi.org/10.1016/j.neuron.2015.09.012

Murphy, B. K., & Miller, K. D. (2003). Multiplicative Gain Changes Are Induced by Excitation or Inhibition Alone. *The Journal of Neuroscience*, *23*(31), 10040–10051. https://doi.org/10.1523/JNEUROSCI.23-31-10040.2003

Murphy, P. R., O'Connell, R. G., O'Sullivan, M., Robertson, I. H., & Balsters, J. H. (2014). Pupil diameter covaries with BOLD activity in human locus coeruleus. *Human Brain Mapping*, *35*(8), 4140–4154. https://doi.org/10.1002/hbm.22466

Niell, C. M., & Stryker, M. P. (2010). Modulation of Visual Responses by Behavioral State in Mouse Visual Cortex. *Neuron*, *65*(4), 472–479. https://doi.org/10.1016/j.neuron.2010.01.033

Nolte, G. (2003). The magnetic lead field theorem in the quasi-static approximation and its use for magnetoencephalography forward calculation in realistic volume conductors. *Physics in Medicine and Biology*, *48*(22), 3637–3652. https://doi.org/10.1088/0031-9155/48/22/002

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). FieldTrip: Open Source Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data. *Computational Intelligence and Neuroscience*, *2011*, 1–9. https://doi.org/10.1155/2011/156869

Pettine, W. W., Louie, K., Murray, J. D., & Wang, X.-J. (2021). Excitatory-inhibitory tone shapes decision strategies in a hierarchical neural network model of multi-attribute choice. *PLOS Computational Biology*, *17*(3),

1040   e1008791. https://doi.org/10.1371/journal.pcbi.1008791

Pfeffer, T., Avramiea, A.-E., Nolte, G., Engel, A. K., Linkenkaer-Hansen, K., & Donner, T. H. (2018). Catecholamines alter the intrinsic variability of cortical population activity and perception. *PLOS Biology*, *16*(2), e2003453. https://doi.org/10.1371/journal.pbio.2003453

Pfeffer, T., Ponce-Alvarez, A., Meindertsma, T., Gahnström, C., Brink, R. L. van den, Nolte, G., Tsetsos, K., Engel, A.

1045   K., Deco, G., & Donner, T. H. (2020). Circuit mechanisms for chemical modulation of cortex-wide network interactions and exploration behavior. *BioRxiv*, 2020.06.25.171199. https://doi.org/10.1101/2020.06.25.171199

Polack, P.-O., Friedman, J., & Golshani, P. (2013). Cellular mechanisms of brain state–dependent gain modulation in visual cortex. *Nature Neuroscience*, *16*(9), 1331–1339. https://doi.org/10.1038/nn.3464

1050   Pomè, A., Burr, D. C., Capuozzo, A., & Binda, P. (2020). Spontaneous pupillary oscillations increase during mindfulness meditation. *Current Biology*, *30*(18), R1030–R1031. https://doi.org/10.1016/j.cub.2020.07.064

Poulet, J. F. A., & Petersen, C. C. H. (2008). Internal brain state regulates membrane potential synchrony in barrel cortex of behaving mice. *Nature*, *454*(7206), 881–885. https://doi.org/10.1038/nature07150

Ramos, B. P., & Arnsten, A. F. T. (2007). Adrenergic pharmacology and cognition: Focus on the prefrontal cortex.

1055   *Pharmacology & Therapeutics*, *113*(3), 523–536. https://doi.org/10.1016/j.pharmthera.2006.11.006

Reimer, J., Froudarakis, E., Cadwell, C. R., Yatsenko, D., Denfield, G. H., & Tolias, A. S. (2014). Pupil Fluctuations Track Fast Switching of Cortical States during Quiet Wakefulness. *Neuron*, *84*(2), 355–362. https://doi.org/10.1016/j.neuron.2014.09.033

Reimer, J., McGinley, M. J., Liu, Y., Rodenkirch, C., Wang, Q., McCormick, D. A., & Tolias, A. S. (2016). Pupil

1060   fluctuations track rapid changes in adrenergic and cholinergic activity in cortex. *Nature Communications*, *7*(1), 13289. https://doi.org/10.1038/ncomms13289

Robinson, P. A., Rennie, C. J., Rowe, D. L., O'Connor, S. C., Wright, J. J., Gordon, E., & Whitehouse, R. W. (2003). Neurophysical Modeling of Brain Dynamics. *Neuropsychopharmacology*, *28*(1), S74–S79. https://doi.org/10.1038/sj.npp.1300143

1065   Saalmann, Y. B., Pinsk, M. A., Wang, L., Li, X., & Kastner, S. (2012). The Pulvinar Regulates Information Transmission Between Cortical Areas Based on Attention Demands. *Science*, *337*(6095), 753–756. https://doi.org/10.1126/science.1223082

Sara, S. J. (2009). The locus coeruleus and noradrenergic modulation of cognition. *Nature Reviews Neuroscience*, *10*(3), 211–223. https://doi.org/10.1038/nrn2573

1070   Sarter, M., Parikh, V., & Howe, W. M. (2009). Phasic acetylcholine release and the volume transmission hypothesis: Time to move on. *Nature Reviews Neuroscience*, *10*(5), 383–390. https://doi.org/10.1038/nrn2635

Schneider, M., Hathway, P., Leuchs, L., Sämann, P. G., Czisch, M., & Spoormaker, V. I. (2016). Spontaneous pupil dilations during the resting state are associated with activation of the salience network. *NeuroImage*, *139*,

1075   189–201. https://doi.org/10.1016/j.neuroimage.2016.06.011

Schwalm, M., & Jubal, E. R. (2017). Back to Pupillometry: How Cortical Network State Fluctuations Tracked by Pupil Dynamics Could Explain Neural Signal Variability in Human Cognitive Neuroscience. *ENeuro*, *4*(6). https://doi.org/10.1523/ENEURO.0293-16.2017

Schwarz, L. A., & Luo, L. (2015). Organization of the Locus Coeruleus-Norepinephrine System. *Current Biology*,

1080   *25*(21), R1051–R1056. https://doi.org/10.1016/j.cub.2015.09.039

Segal, M., & Bloom, F. E. (1976a). The action of norepinephrine in the rat hippocampus. III. Hippocampal cellular responses to locus coeruleus stimulation in the awake rat. *Brain Research*, *107*(3), 499–511. https://doi.org/10.1016/0006-8993(76)90140-2

Segal, M., & Bloom, F. E. (1976b). The action of norepinephrine in the rat hippocampus. IV. The effects of locus

1085   coeruleus stimulation on evoked hippocampal unit activity. *Brain Research*, *107*(3), 513–525. https://doi.org/10.1016/0006-8993(76)90141-4

Shimaoka, D., Harris, K. D., & Carandini, M. (2018). Effects of Arousal on Mouse Sensory Cortex Depend on Modality. *Cell Reports*, *22*(12), 3160–3167. https://doi.org/10.1016/j.celrep.2018.02.092

Siegle, J. H., & Wilson, M. A. (2014). Enhancement of encoding and retrieval functions through theta phase-

1090   specific manipulation of hippocampus. *ELife*, *3*, e03061. https://doi.org/10.7554/eLife.03061

Sokoliuk, R., Mayhew, S. D., Aquino, K. M., Wilson, R., Brookes, M. J., Francis, S. T., Hanslmayr, S., & Mullinger, K. J. (2019). Two Spatially Distinct Posterior Alpha Sources Fulfill Different Functional Roles in Attention. *Journal of Neuroscience*, *39*(36), 7183–7194.

Spaak, E., Bonnefond, M., Maier, A., Leopold, D. A., & Jensen, O. (2012). Layer-Specific Entrainment of Gamma-

1095   Band Neural Activity by the Alpha Rhythm in Monkey Visual Cortex. *Current Biology*, *22*(24), 2313–2318. https://doi.org/10.1016/j.cub.2012.10.020

Steriade, M. (1996). Arousal—Revisiting the Reticular Activating System. *Science*, *272*(5259), 225–225. https://doi.org/10.1126/science.272.5259.225

Stitt, I., Zhou, Z. C., Radtke-Schuller, S., & Fröhlich, F. (2018). Arousal dependent modulation of thalamo-cortical

1100   functional interaction. *Nature Communications*, *9*(1), 2455. https://doi.org/10.1038/s41467-018-04785-6

Tallon-Baudry, C., & Bertrand, O. (1999). Oscillatory gamma activity in humans and its role in object representation. *Trends in Cognitive Sciences*, *3*(4), 151–162. https://doi.org/10.1016/s1364-6613(99)01299-1

Totah, N. K., Neves, R. M., Panzeri, S., Logothetis, N. K., & Eschenko, O. (2018). The Locus Coeruleus Is a Complex

1105   and Differentiated Neuromodulatory System. *Neuron*, *99*(5), 1055-1068.e6.

https://doi.org/10.1016/j.neuron.2018.07.037

Trakoshis, S., Martínez-Cañada, P., Rocchi, F., Canella, C., You, W., Chakrabarti, B., Ruigrok, A. N., Bullmore, E. T., Suckling, J., Markicevic, M., Zerbi, V., MRC AIMS Consortium, Baron-Cohen, S., Gozzi, A., Lai, M.-C., Panzeri, S., & Lombardo, M. V. (2020). Intrinsic excitation-inhibition imbalance affects medial prefrontal cortex differently in autistic men versus women. *ELife*, *9*, e55684. https://doi.org/10.7554/eLife.55684

Turchi, J., Chang, C., Ye, F. Q., Russ, B. E., Yu, D. K., Cortes, C. R., Monosov, I. E., Duyn, J. H., & Leopold, D. A. (2018). The Basal Forebrain Regulates Global Resting-State fMRI Fluctuations. *Neuron*, *97*(4), 940-952.e4. https://doi.org/10.1016/j.neuron.2018.01.032

Turnbull, P. R. K., Irani, N., Lim, N., & Phillips, J. R. (2017). Origins of Pupillary Hippus in the Autonomic Nervous System. *Investigative Ophthalmology & Visual Science*, *58*(1), 197–203. https://doi.org/10.1167/iovs.16-20785

Urai, A. E., Braun, A., & Donner, T. H. (2017). Pupil-linked arousal is driven by decision uncertainty and alters serial choice bias. *Nature Communications*, *8*(1), 14637. https://doi.org/10.1038/ncomms14637

van den Brink, R. L., Nieuwenhuis, S., & Donner, T. H. (2018). Amplification and Suppression of Distinct Brainwide Activity Patterns by Catecholamines. *Journal of Neuroscience*, *38*(34), 7476–7491.

van den Brink, R. L., Pfeffer, T., & Donner, T. H. (2019). Brainstem Modulation of Large-Scale Intrinsic Cortical Activity Correlations. *Frontiers in Human Neuroscience*, *13*. https://doi.org/10.3389/fnhum.2019.00340

Van Diepen, R. M., Foxe, J. J., & Mazaheri, A. (2019). The functional role of alpha-band activity in attentional processing: The current zeitgeist and future outlook. *Current Opinion in Psychology*, *29*, 229–238. https://doi.org/10.1016/j.copsyc.2019.03.015

Van Veen, B. D., van Drongelen, W., Yuchtman, M., & Suzuki, A. (1997). Localization of brain electrical activity via linearly constrained minimum variance spatial filtering. *IEEE Transactions on Bio-Medical Engineering*, *44*(9), 867–880. https://doi.org/10.1109/10.623056

Vinck, M., Batista-Brito, R., Knoblich, U., & Cardin, J. A. (2015). Arousal and Locomotion Make Distinct Contributions to Cortical Activity Patterns and Visual Encoding. *Neuron*, *86*(3), 740–754. https://doi.org/10.1016/j.neuron.2015.03.028

Waschke, L., Donoghue, T., Fiedler, L., Smith, S., Garrett, D. D., Voytek, B., & Obleser, J. (2021). Modality-specific tracking of attention and sensory statistics in the human electrophysiological spectral exponent. *BioRxiv*, 2021.01.13.426522. https://doi.org/10.1101/2021.01.13.426522

Waschke, L., Tune, S., & Obleser, J. (2019). Local cortical desynchronization and pupil-linked arousal differentially shape brain states for optimal sensory performance. *ELife*, *8*, e51501. https://doi.org/10.7554/eLife.51501

Wierda, S. M., van Rijn, H., Taatgen, N. A., & Martens, S. (2012). *Pupil dilation deconvolution reveals the dynamics of attention at high temporal resolution. 109*(22), 8456–8460. https://doi.org/10.1073/pnas.1201858109

Yellin, D., Berkovich-Ohana, A., & Malach, R. (2015). Coupling between pupil fluctuations and resting-state fMRI uncovers a slow build-up of antagonistic responses in the human cortex. *NeuroImage*, *106*, 414–427. https://doi.org/10.1016/j.neuroimage.2014.11.034

Yerkes, R. M., & Dodson, J. D. (1908). The relation of strength of stimulus to rapidity of habit-formation. *Journal of Comparative Neurology and Psychology*, *18*(5), 459–482. https://doi.org/10.1002/cne.920180503

Zaborszky, L., Csordas, A., Mosca, K., Kim, J., Gielow, M. R., Vadasz, C., & Nadasdy, Z. (2015). Neurons in the Basal Forebrain Project to the Cortex in a Complex Topographic Organization that Reflects Corticocortical Connectivity Patterns: An Experimental Study Based on Retrograde Tracing and 3D Reconstruction. *Cerebral Cortex*, *25*(1), 118–137. https://doi.org/10.1093/cercor/bht210

Záborszky, L., Gombkoto, P., Varsanyi, P., Gielow, M. R., Poe, G., Role, L. W., Ananth, M., Rajebhosale, P., Talmage, D. A., Hasselmo, M. E., Dannenberg, H., Minces, V. H., & Chiba, A. A. (2018). Specific Basal Forebrain–Cortical Cholinergic Circuits Coordinate Cognitive Operations. *Journal of Neuroscience*, *38*(44), 9446–9458.

Zagha, E., Casale, A. E., Sachdev, R. N. S., McGinley, M. J., & McCormick, D. A. (2013). Motor Cortex Feedback Influences Sensory Processing by Modulating Network State. *Neuron*, *79*(3), 567–578. https://doi.org/10.1016/j.neuron.2013.06.008

Zelano, C., Jiang, H., Zhou, G., Arora, N., Schuele, S., Rosenow, J., & Gottfried, J. A. (2016). Nasal Respiration Entrains Human Limbic Oscillations and Modulates Cognitive Function. *Journal of Neuroscience*, *36*(49), 12448–12467.

Additional author information

#	Author	Affil	ORCID	Twitter@
1	Thomas Pfeffer	1,2	0000-0001-9561-3085	@thmspfffr
2	Christian Keitel	3,4	0000-0003-2597-5499	@KeiCetel
3	Daniel S. Kluger	5,6	0000-0002-0691-794X	@danlikesbrains
4	Anne Keitel	7	0000-0003-4498-0146	@anneke_sci
5	Alena Russmann	2		
6	Gregor Thut	4	0000-0003-1313-4262	@GregorThut
7	Tobias Donner	2	0000-0002-7559-6019	@donner_lab
8	Joachim Gross	4,5,6	0000-0002-3994-1006	@joachim__gross

1 Universitat Pompeu Fabra, Center for Brain and Cognition, Computational Neuroscience Group, Barcelona, Spain | **2** University Medical Center Hamburg-Eppendorf, Department of Neurophysiology and Pathophysiology, Martinistr. 52, 20246 Hamburg, Germany | **3** University of Stirling, Psychology, FK9 4LA Stirling, UK | **4** Centre for Cognitive Neuroimaging, Institute of Neuroscience and Psychology, University of Glasgow, 62 Hillhead Street, G12 8QB Glasgow, UK | **5** Institute for Biomagnetism and Biosignal Analysis, University of Münster, Malmedyweg 15, 48149 Muenster, Germany | **6** Otto Creutzfeldt Center for Cognitive and Behavioral Neuroscience, University of Münster, Fliednerstrasse 21, 48149 Muenster, Germany | **7** University of Dundee, Psychology, DD1 4HN Dundee, UK

Acknowledgments

TP and THD thank Christiane Reissmann and Karin Deazle for assistance with MEG recordings and the acquisition of structural brain scans. CK, AK, GT and JG thank Gavin Paterson and Frances Crabbe for expert assistance with MEG recordings and structural brain scans. CK thanks Linbi Hong, Josef Faller, and Paul Sajda for comments on the research. DSK and JG thank Hildegard Deitermann, Ute Trompeter, and Karin Wilken for assistance with MEG recordings and structural brain scans. All authors thank Ruud L. van den Brink for helpful comments on the manuscript.

Funding

TP has been supported by a Feodor-Lynen fellowship of the Alexander-von-Humboldt Foundation. CK and AK have been supported by a Wellcome Trust Senior Investigator Grant awarded to JG (#098433) and GT (#98434). CK received further support through BBSRC Flexible Talent Mobility funding (BB/R506576/1) awarded by the University of Glasgow. THD has been supported by the German Research Foundation (DFG, grants DO 1240/3-1, DO 1240/4-1, and SFB 936, project numbers A7 & Z3) and the German Federal Ministry of

Education and Research (BMBF, grants 01GQ1907 and 01EW2007B). JG has been supported by the Interdisciplinary Center for Clinical Research (IZKF) of the medical faculty

1185   of Münster (grant number Gro3/001/19) and the DFG (GR 2024/5-1)

Author contributions

Contributions coded according to the CRediT taxonomy (Brand et al., 2015). Conceptualization: TP, CK, AR, THD, JG. Data curation: TP, CK, DK. Formal analysis: TP, CK, DK. Funding acquisition: TP, THD, GT, JG, CK. Investigation: TP, AK, CK, DK.

1190   Methodology: TP, CK, TD, JG. Project administration: TP, CK. Resources: THD, JG, GT. Software: TP, CK. Supervision: THD, JG. Validation: TP, CK, DK, AK. Visualization: TP, CK. Writing – original draft: TP, CK. Writing – review & editing: All authors. Final version of the manuscript approved by all authors.

Declaration of conflicting interests

1195   The authors declare that there were no competing interests with respect to the authorship or publication of this research article.

Supplementary Figures

1200

Figure S1. **(A)** Power spectra across recording sites, separately shown for all three MEG laboratories (Glasgow, Hamburg and Muenster; from top to bottom). **(B)** The individual power spectra, across all 81 included participants, are highly correlated

1205

Figure S2. Power spectra and signal components of pupil time series, for all MEG laboratories. **(A)** Grand-average pupil power spectra for each centre. From top to bottom: Glasgow (red), Hamburg (blue), Münster (yellow). Shaded areas show standard error of the mean. **(B)** Scatter plot of two aperiodic parameters of individual pupil spectra, the spectral slope characterised by its 1/f exponent (y-axis) and the spectral offset (x-axis) as extracted with the FOOOF algorithm (Donoghue et al., 2020). Dots use the same color-coding for labs as in A. **(C)** Scatter plot of the center frequency (x-axis) and power (y-axis) of the prominent rhythmic component between 0.2-0.4 Hz in individual power spectra (see A), the so-called Hippus, as extracted with FOOOF. Dots use the same color-coding for labs as in A.

1210

1215

Figure S3. Sensor level cross-correlations between the temporal derivative of fluctuations in pupil diameter (pupil derivative) and band-limited power fluctuations. **(A)** Correlation between the pupil derivative and band-limited power fluctuations across lags, ranging from -5 to 5s and for all three recording sites (left: Glasgow; center: Hamburg; right: Muenster). Colors indicate the MEG frequency-band, with darker colors representing lower frequencies and lighter colors representing higher frequencies. Negative lags are indicative of "pupil derivative preceding MEG", whereas positive lags are indicative of the opposite. The vertical dashed black line denotes a lag of 930 ms (Hoeks & Levelt, 1993) and the dashed light gray line a lag of 0 ms. **(B)** Same as in (A), but averaged across the three recording sites and for three frequency-bands of interest: 2-4 Hz (black), 8-16 Hz (dark gray) and 64-128 Hz (gray dashed). **(C)** Correlation values across all channel groups, sorted from anterior to posterior, and for lags ranging from -5 to +5s. The dashed gray line depicts a lag of 0 ms and the dashed black line shows a lag of 930 ms (Hoeks & Levelt, 1993). Correlation values were averaged within three frequency-bands of interest: 2-4 Hz (left), 8 - 16 Hz (center) and 64-128 Hz (right). Masked at *P*<0.05 (two-sided paired t-test; FDR-adjusted, q=0.01).

Figure S4. Spatial distribution of cross correlations for all recording sites. **(A)** Cross correlation between spontaneous fluctuations in pupil diameter and band-limited power, across various temporal lags (from -5s to 5s), separately for the Glasgow (top), the Hamburg (middle) and the Münster (bottom) datasets and averaged across three frequency ranges of interest (2-4 Hz, 8-16 Hz and 64-128 Hz, from left to right). **(B)** Same as (A), but for pupil-derivative. All masked at $P = 0.05$ (two-sided paired t-test). Note that, compared to Figure 3, the P-values were not FDR-adjusted.

1235

1240

Figure S5. Source space pupil-derivative-power correlations. **(A)** Spectrum of pupil-derivative-power correlations, averaged across all 8799 source locations. Filled black circles denote significant correlations (P < 0.05; two-tailed paired t-test; FDR-adjusted, q=0.1). The gray shaded area depicts the standard error of the mean across subjects.

1245 **(B)** Pupil-power correlations, averaged across the 246 regions of the Brainnetome atlas and sorted from anterior to posterior regions. Masked at *P* = 0.05 (two-tailed paired t-test; FDR-adjusted, q=0.1). **(C)** Pupil-power correlations across four selected regions-of-interest: visual cortex, auditory cortex, somatosensory cortex and anterior cingulate cortex (from top to bottom). Insets depict the spatial extent of the respective region of interest. **(D)** Spatial distribution of pupil-power correlations, across four frequency bands of interest. Correlations between pupil and power (in clockwise direction, starting from the top left) at 4 Hz, 11.3 Hz, 19.0 Hz, and 64 Hz. All maps

1250 are masked at *P* = 0.05 (two-tailed paired t-test; FDR-adjusted, q=0.1).

Figure S6. Difference between pupil-power and pupil-derivative-power correlations. **(A)** Differences, pooled across all 8799 source locations (black dots indicate *P*<0.05; FDR-adjusted, q=0.01). **(B)** Differences, averaged across the 246 regions of the Brainnetome atlas, and sorted from anterior to posterior regions. Masked at *P* = 0.05 (two-sided paired t-test; FDR-adjusted, q=0.01). Red colors: correlation values are larger for pupil-power correlations; blue colors: correlation values larger for the pupil-derivative-power correlations. **(C)** Differences across four regions of interest (from top to bottom): visual cortex, auditory cortex, somatosensory cortex and anterior cingulate cortex. **(D)** Spatial distribution of difference in pupil-power and pupil-derivative-power correlations, across four frequency bands of interest. Red colors: correlation values are larger for pupil-power correlations; blue colors: correlation values larger for the pupil-derivative-power correlations. Filled black circles denote *P*<0.05 (Two-sided paired t-test; FDR-adjusted, q=0.01);

1255

1260

1265

Figure S7. Nonlinear relations between pupil-derivative and band-limited power fluctuations. (A) Normalized spectral power, averaged across three frequency ranges (2-4 Hz, 8-16 Hz and 64-128 Hz; from left to right), as a function of mean pupil diameter. (B) Normalized coefficient of the quadratic term (β_2; See Methods for details),
1270   averaged across all subjects and atlas regions. Black dots denote $P<0.05$, red dots denote $P<0.01$ and yellow dots denote $P<0.05$ after adjusting for false discoveries (q = 0.1; two-sided paired t-test). The gray shaded area depicts the standard error of the mean across subjects. Positive and negative values are indicative of a U-shaped and inverted-U-shaped relationship, respectively (C) Normalized β_2 coefficient, averaged across all subjects, separately for four regions of interest: visual cortex, auditory cortex, somatosensory cortex and anterior cingulate
1275   cortex (from left to right). Insets show the approximate extent of the regions of interest. (D) Spatial distribution of the coefficient of the quadratic term (see Methods for details), at center frequencies 11 Hz (left) and 19 Hz (right). Masked at $P = 0.05$ (two-sided paired t-test, adjusted for a false discovery rate of 0.1).

1280

Figure S8. Pupil-derivative correlates with aperiodic component of the power spectrum. (A) Spectral exponent (i.e., the slope of the aperiodic component) for 20 pupil bins (sorted from small to large). **(B)** Correlation coefficients of the correlation between the spectral exponent and pupil diameter fluctuations, averaged across space (black dots, showing the individual subjects) and for the four regions of interest: visual cortex (red), auditory

1285 cortex (yellow) somatosensory cortex (purple) and anterior cingulate cortex (green). The horizontal lines highlight the mean correlation coefficient (*** $P<0.001$; two-sided paired t-tests). **(C)** Spatial map of the correlation between the slope of the aperiodic component and intrinsic fluctuations in pupil derivative. Spatial maps are masked at $P = 0.05$ after adjusting for the false discovery rate (two-sided paired t-test). **(D)** Correlations between fluctuations in pupil derivative and band-limited power, after removing the aperiodic component. Black dots denote significant

1290 values ($P<0.05$; FDR-adjusted; two-sided paired t-test).