1

Solar simulated ultraviolet radiation inactivates HCoV-NL63

and SARS-CoV-2 coronaviruses at environmentally relevant doses

Georg T. Wondrak[a*], Jana Jandova[a], Spencer J. Williams[b], and Dominik Schenten[b*]

[a]Department of Pharmacology and Toxicology, College of Pharmacy and UA Cancer Center,

University of Arizona, Tucson, Arizona.

[b]Department of Immunobiology, College of Medicine, University of Arizona, Tucson, Arizona.

*Corresponding authors' email:

wondrak@pharmacy.arizona.edu (Georg T. Wondrak)

dschenten@arizona.edu (Dominik Schenten)

Keywords: Coronavirus; HCoV-NL63; SARS-CoV-2; Solar simulated ultraviolet radiation; Viral inactivation

ABSTRACT

The germicidal properties of short wavelength ultraviolet C (UVC) light are well established and used to inactivate many viruses and other microbes. However, much less is known about germicidal effects of terrestrial solar UV light, confined exclusively to wavelengths in the UVA and UVB regions. Here, we have explored the sensitivity of the human coronaviruses HCoV-NL63 and SARS-CoV-2 to solar-simulated full spectrum ultraviolet light (sUV) delivered at environmentally relevant doses. First, HCoV-NL63 coronavirus inactivation by sUV-exposure was confirmed employing (*i*) viral plaque assays, (*ii*) RT-qPCR detection of viral genome replication, and (*iii*) infection-induced stress response gene expression array analysis. Next, a detailed dose-response relationship of SARS-CoV-2 coronavirus inactivation by sUV was elucidated, suggesting a half maximal suppression of viral infectivity at low sUV doses. Likewise, extended sUV exposure of SARS-CoV-2 blocked cellular infection as revealed by plaque assay and stress response gene expression array analysis. Moreover, comparative (HCoV-NL63 versus SARS-CoV-2) single gene expression analysis by RT-qPCR confirmed that sUV exposure blocks coronavirus-induced redox, inflammatory, and proteotoxic stress responses. Based on our findings, we estimate that solar ground level full spectrum UV light impairs coronavirus infectivity at environmentally relevant doses. Given the urgency and global scale of the unfolding SARS-CoV-2 pandemic, these prototype data suggest feasibility of solar UV-induced viral inactivation, an observation deserving further molecular exploration in more relevant exposure models.

Abbreviations: MOI, multiplicity of infection; sUV, solar simulated ultraviolet light; UV, ultraviolet.

1. Introduction

The germicidal properties of short wavelength ultraviolet C (UVC) light are well established and widely used to inactivate many viruses and other microbes, and virucidal activity of solar UVC targeting pathogenic coronaviruses has been explored in much detail before [1-3]. Given the urgency and global scale of the unfolding SARS-CoV-2-caused COVID-19 pandemic, UV-induced inactivation of coronaviruses including SARS-CoV-2 has reemerged as a matter of much contemporary research interest [2-8]. Indeed, recently, rapid and complete inactivation of SARS-CoV-2 by UVC has been substantiated experimentally, and virucidal UVC light sources (254 nm emission) are used for surface disinfection and decontamination [5,8]. Moreover, far UVC (222 nm) has attracted considerable attention due to its potent virucidal activity [2]. However, much less is known about germicidal (and coronavirus-directed) effects of terrestrial (ground level) solar UV light, a matter of much interest given the airborne spread of coronaviruses including SARS-CoV-2 [2,6]. UVC (< 290 nm) is not present in the solar spectrum reaching the Earth's surface, and most of solar UV energy incident on the skin is from the UVA region (>95%; from 320–400 nm). Remarkably, the UVB (290–320 nm) proportion of total solar UV-flux received by skin can be well below 2% depending on the solar angle, which determines the atmospheric light path length and thereby the degree of ozone-filtering and preferential Rayleigh scattering of short wavelength UV light [9].

Recently, the role of ground level (environmentally relevant) solar UV has been explored in the context of SARS-CoV-2 disinfection, and a role of solar UVB in human coronavirus inactivation has been substantiated based on atmospheric and geophysical simulations [2,6,10,11]. Specifically, inactivation times of SARS coronaviruses exposed to environmental photons with wavelengths between 290-315 nm have been calculated using OMI (ozone monitoring instrument)

satellite data for the sunlit earth [10]. Moreover, recent research has demonstrated that simulated

sunlight rapidly inactivates SARS-CoV-2 on surfaces including human saliva when exposed to

simulated sunlight representative of the summer solstice at 40 °N latitude at sea level on a clear

day [10]. Also, indirect effects of solar UVB exposure in reducing COVID-19 deaths have been

substantiated, potentially mediated by UVB-driven cutaneous vitamin D synthesis, among other

factors [12-14]. In addition, a role of solar UVA photons in the inactivation of coronaviruses has

been proposed [7].

Given the complexity of virucidal activity as a function of spectral composition from ultraviolet

to infrared, a topic recently reviewed by various authors, a more detailed knowledge and direct

evidence of solar UV-induced coronavirus inactivation (achievable at ground level and

environmentally relevant doses) would offer improved options that inform decisions at the basic

research, clinical care, and public health levels [2,6,8]. Here, for the first time, we have explored

the sensitivity of the human coronaviruses HCoV-NL63 and SARS-CoV-2 to solar simulated

ultraviolet light (sUV). Our findings suggest that solar UV delivered at environmentally relevant

dose levels inactivates HCoV-NL63 and SARS-CoV-2 coronaviruses with pronounced blockade

of infectivity protecting mammalian host cells.

2. Materials and Methods

2.1. Chemicals

All chemicals were purchased from Sigma Aldrich (St. Louis, MO, USA).

2.2. Mammalian cell culture, viral propagation, and target cell infection

As established viral target cells infected by HCoV-NL63 and SARS-CoV-2, Calu-3 human

metastatic lung epithelial adenocarcinoma (HTB-55), Caco-2 human colorectal epithelial

adenocarcinoma (HTB-37) and Vero normal epithelial monkey kidney (CCL-81) cells (all from

ATCC, Manassas, VA, USA) were maintained according to published standard procedures [15-

18]. In brief, all cells (Calu-3, Caco-2 and Vero) were cultured in Eagle's Minimum Essential

Medium (MEM) medium (Corning, Manassas, VA) supplemented with 10% bovine calf serum

(BCS, HyClone™ Laboratories, Logan, UT).   Coronavirus HCoV-NL63 (NR-470) and its

genomic RNA (NR-44105) were obtained from BEI Resources (NIAID, NIH).   SARS-CoV-2

strain WA1 (NR-52281; BEI Resources) was propagated in Vero cells unless specified otherwise

[6]. For viral stocks, cells were infected at a multiplicity of infection (MOI) of 0.01 and cultured

for 48 h. At that point, cells were harvested, homogenized, subjected to a single freeze-thaw cycle,

and then combined with the culture supernatant followed by centrifugation (3000 rpm, 10 min).

The viral titers of the final supernatant (after serial dilution) was determined by plaque forming

assay. All work with SARS-CoV-2 was performed under BSL3 conditions in a facility with

negative pressure and PPE that included Tyvek suits and N95 masks for respiratory protection.

2.3. Viral irradiation with solar simulated UV light (sUV)

A KW large area light source solar simulator, model 91293, from Oriel Corp. (Stratford, CT)

was used, equipped with a 1000W xenon arc lamp power supply, model 68920, and a VIS-IR band

pass blocking filter plus either an atmospheric attenuation filter (output 290–400 nm plus residual

650–800 nm for solar simulated light) [19,20]. For viral irradiation, viral stocks were diluted

>1:100 in PBS and irradiated in a sealed UV-transparent cuvette [BrandTech™ BRAND™ UV-

Cuvets, providing transparency from 230 to 900 nm, widely used for DNA, RNA and protein

analysis (BrandTech™ 759170, Fisher Scientific)]. The cuvette was inserted into a fully UV-

transparent scintillation counter vial (Wheaton '180' low-potassium glass, SigmaAldrich

Z253081). The UV output was quantified using a dosimeter from International Light Inc.

(Newburyport, MA), model IL1700, with an SED240 detector for UVB (range 265–310 nm, peak

at 285 nm) or a SED033 detector for UVA (range 315–390 nm peak 365 nm) at a distance of 365

mm from the source, which was used for all experiments. In order to avoid artifactual thermal

effects of photon exposure on viral activity, cuvettes were placed on ice during irradiation. At 365

mm from the source, total solar UV intensity was 5.34 mJ/cm^2 s (UVA) and 0.28 mJ/cm^2 s (UVB).

2.4. HCoV-NL63 plaque forming assay and viral RNA quantification

A published standard procedure was followed [15,21]. For HCoV-NL63, target cells (CaCo-2

or Calu-3) were seeded in 6-well plates at approximately 4×10^5 cells per well and incubated until

the monolayer was 80–90% confluent. Prior to infection, cells were washed with phosphate

buffered saline (PBS). Virus inoculum (MOI=0.01) in 500 μL of growth media supplemented with

2% horse serum (with standard penicillin/streptomycin and L-glutamine supplementation) was

added to each well. Viral entry was performed by incubation at 4°C for 30-60 min with gentle

agitation followed by 1 h incubation in 33°C, 5% CO$_2$. Then, inoculum was removed and cells

were washed twice with PBS and replaced by 2 mL of normal growing media. After infection,

cells were washed twice with PBS and placed in the incubator and cultured in normal growth

media. Once plaques appeared (~5-7 d post infection), cells were fixed with 10% neutral buffered

formalin for 30 min at room temperature and stained with 1% crystal violet in 20% methanol for

20 min. Then, cells were washed several times with water, and plaques were counted and

representative pictures taken at 10x magnification using an inverted microscope (Nikon

Instruments, Melville, NY).   In addition, viral RNA was extracted from cells and the respective

culture supernatant with the QIAamp Viral RNA Mini Kit (QIAGEN, Hilden, Germany). One step

RT-qPCR for HCoV-NL63 with absolute virus RNA quantification was performed using the

following primer/probe set as published before [22]:

forward primer – 5′-ACGTACTTCTATTATGAAGCATGATATTAA-3′

reverse primer – 5′-AGCAGATCTAATGTTATACTTAAAACTACG-3′

probe – FAM-5′- ATTGCCAAGGCTCCTAAACGTACAGGTGTT -3′-NFQ-MGB

Briefly, RT-qPCR was carried out in a 20 µL reaction mixture with extracted RNA and One

step RT-qPCR 2x Master Mix containing ROX as a passive reference dye (Gold Biotechnology,

St. Louis, MO) and 300 nM forward and reverse primers and 200 nM MGB probe. Amplification

and detection were performed in ABI 7500 system (Applied Biosystems, Foster city, CA) under

the following conditions: first strand cDNA synthesis at 42°C for 30 min; initial denaturation/RT

inactivation at 95°C for 3 min; denaturation at 95°C for 10 sec and annealing/extension at 55°C

for 30 sec followed by 45 sec for data acquisition at 72°C. During amplification, the ABI PRISM

7500 sequence detector monitored real-time PCR amplification by quantitative analysis of the

fluorescence emissions. The reporter dye (FAM) signal was measured against the internal

reference dye (ROX) signal to normalize the signals for non-PCR-related fluorescence fluctuations

that occur from well to well. The cycle threshold (Ct) represented the refraction cycle number at

which a positive amplification was measured and was set at ten times the standard deviation of the

mean baseline emission calculated for PCR cycles 3 to 15. Genomic RNA from HCoV-NL63 was

used as a positive control.

2.5. SARS-CoV-2 plaque forming assay and viral RNA quantification

The quantification of infectious SARS-CoV-2 has been published before [18]. Target cells

(Vero or Calu-3) were infected in triplicates at an MOI of 0.005 (high titer) or 0.001 (low titer).

Briefly, cells were incubated with SARS-CoV-2 for 2 h and subsequently overlaid with 1%

methylcellulose in culture medium. After 3-4 days, the cells were fixed in 10% neutral buffered

formalin for 30 min, washed under tap water, and stained with 1% crystal violet. The number of

plaques was counted on a light table. Alternatively, infection of cells was determined by measuring

the amount of viral RNA. Cells were lysed in Trizol followed by RNA extraction with the

RNAeasy kit (Qiagen). After reverse transcription, cDNA corresponding to the gene encoding the

SARS-CoV-2 spike protein was quantified by qPCR with the Perfecta FastMix (QuantaBio) using:

forward primer (SARS-CoV-2) 5′-GCTGGTGCTGCAGCTTATTA-3′

reverse primer (SARS-CoV-2) 5′-AGGGTCAAGTGCACAGTCTA-3′

at an annealing temperature of 60 ºC. For normalization, *GAPDH* expression was measured using

the following primers:

forward primer (*GAPDH*) 5′-TGGTGAAGGTCGGTGTGAAC-3′

reverse primer (*GAPDH*) 5′-CCATGTAGTTGAGGTCAATGAAGG-3′ .

2.6. Human Stress & Toxicity PathwayFinder RT² Profiler^{TM} gene expression array analysis of

infected host cells

Seven days post infection of Calu-3 host cells with either HCoV-NL63 (MOI=0.01) or HCoV-

NL63 exposed to sUV (UVB portion: 706 mJ/cm^2), total mRNA from host cells was isolated using

the RNeasy Mini kit (Qiagen, Valencia, CA) following our published standard procedures. Reverse

transcription was then performed using the RT^2 First Strand kit (Qiagen) from 500 ng total RNA.

For gene expression array analysis, the human Stress & Toxicity PathwayFinder RT^2 Profiler^{TM}

technology (Qiagen), assessing expression of 84 stress response-related genes, was used as

published before [23,24]. Quantitative PCR was run using the following conditions: 95 °C (10

min), followed by 40 cycles at 95 °C (15 s) alternating with 60 °C (1 min) (Applied Biosystems,

Carlsbad, CA). Gene-specific products were normalized to a group of 5 housekeeping genes

(*ACTB*, *B2M*, *GAPDH*, *HPRT1*, *RPLP0)* and quantified using the comparative $\Delta\Delta Ct$ method (ABI

PRISM 7500 sequence detection system user guide). Expression values were averaged across at

least three independent array experiments, and standard deviation was calculated for graphing and

statistical analysis as published before.

2.7. Individual RT-qPCR analysis

Total cellular mRNA was isolated using the Qiagen RNeasy Mini Kit (Qiagen, Gaithersburg,

MD) according to the manufacturer's protocol as published by us before [24]. Human primer

probes [*CCL3* (Hs_00234142_m1), *CSF2* (Hs_00929873_m1), *HSPA6* (Hs_00275682_s1), *IL1B*

(Hs_00174097_m1), *IL6* (Hs_00985639_m1), *SOD2* (Hs_00167309_m1), *TNF*

(Hs_00174128_s1), and *RSP18 (*housekeeping gene; Hs_01375212_g1)], were obtained

from ThermoFisher Scientific (Waltham, MA). After cDNA synthesis, quantitative PCR reactions

were performed as follows: 10 min (95 °C) followed by 15 sec (95 °C), 1 min (60 °C), 40 cycles,

using the ABI7500 Real-Time PCR System (Applied Biosystems, Foster City, CA). Amplification

plots were generated, and Ct values were recorded as published before [24].

2.8. Statistical analysis

Unless stated differently, data sets were analyzed employing analysis of variance (ANOVA)

with Tukey's posthoc test using the GraphPad Prism 9.1.0 software (Prism Software Corp., Irvine,

CA); in respective bar graphs (analyzing more than two groups), means without a common letter

differ ($p < 0.05$) as published before [24]. For bar graphs comparing two groups only, statistical

significance was calculated employing the Student's two-tailed t-test, utilizing Excel (Microsoft,

Redmond, WA). Experiments were performed in sets of at least three independent repeats. The

level of statistical significance was marked as follows: *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

3. Results

3.1. Solar simulated UV exposure of HCoV-NL63 blocks subsequent viral infection and

replication in Calu-3 human epithelial lung cells

First, we examined the feasibility of UV-inactivation of a pathologically relevant coronavirus

by employing a single dose of solar simulated UV light using a commercial xenon light source

with quantified spectral power distribution (Fig. 1A). To this end, we exposed human coronoavirus

NL63 (HCoV-NL63) in PBS to a high dose of sUV [equivalent to approximately 6 minimal

erythemal doses (MEDs; UVA: 13.46 J/cm^2; UVB: 706 mJ/cm^2)] and subsequently used it to infect

Calu-3 target cells for 7 days [2, 6, 8]. We used unexposed virus as controls. Strikingly, sUV pre-

exposure strongly suppressed viral infectivity of target cells as demonstrated by quantitative

plaque assay analysis, indicating that sUV exposure caused a more than 8-fold decrease in viral

infectivity (Fig. 1B).

Next, we examined the dose-response relationship characterizing the inhibition of HCoV-NL63

viral replication (induced by sUV pre-exposure) by one step RT-qPCR analysis of the genomic

RNA copy number. We detected a significant inhibition at low sUV doses [UVA: 0.25 J/cm^2; 13

mJ/cm^2 UVB]. Viral inactivation of more than 98 % occurred at doses equal and above 480 mJ/cm^2

UVB (UVA: 9.04 J/cm^2; Fig. 1C).

230   >Figure 1<

3.2. Solar simulated UV exposure of HCoV-NL63 blocks subsequent infection of Caco-2 human epithelial colorectal cells

In order to explore sUV effects on HCoV-NL63 infectivity in another human target cell, we exposed the virus (in PBS) to a high dose of sUV [equivalent to approximately 6 MEDs (UVA: 13.46 J/cm^2; UVB: 706 mJ/cm^2)] and subsequently infected Caco-2 epithelial colon cells (Fig. 2). As observed before with Calu-3 cells (Fig. 1), our quantitative plaque assay analysis showed that the suppression of viral infectivity of Caco-2 target cells by sUV exposure caused a more than 4-fold decrease in plaque formation (Fig. 2A). Likewise, our dose response analysis by RT-qPCR of genomic RNA copy numbers indicated that sUV exposure caused a pronounced suppression of HCoV-NL63 viral replication at doses as low as 240 mJ/cm^2 UVB (UVA: 4.52 J/cm^2; Fig. 2B).

>Figure 2<

3.3. Stress response gene expression array analysis confirms solar UV-induced inhibition of HCoV-NL63 infectivity targeting Calu-3 human epithelial lung cells

Next, the cellular stress response of Calu-3 human epithelial lung cells, elicited by infection with either mock-irradiated or sUV pre-exposed HCoV-NL63, was examined at the gene expression level using the RT^2 Human Stress and Toxicity PathwayFinder^{TM} PCR Array technology. To this end, we infected Calu-3 target cells with sUV or mock-treated virus (doses as in Figs. 1, 2) and profiled the gene expression at the end of the experiment. We observed global HCoV-NL63-induced expression changes (antagonized by viral pre-exposure to sUV) as depicted by Volcano plot (Fig. 3). As expected, HCoV-NL63 viral infection caused a pronounced upregulation of stress response gene expression including genes encoding key regulators of

inflammatory signaling (such as *CSF2*, *TNF, IL1B*, *IL1A*, *CCL3*, *CXCL10*, *NFKBIA*, and *IL6*),

oxidative stress defense (such as *SOD2*), and heat shock response (such as *HSPA6*; Fig. 3). In

contrast, after viral sUV-exposure performed pre-infection, most of these infection-associated

expression changes were either attenuated or completely obliterated, an observation consistent

with pronounced suppression of HCoV-NL63 viral infectivity as a consequence of sUV-exposure.

Likewise, HCoV-NL63 viral infection-induced expression changes causing downregulation of

specific apoptotic modulators including *BCL2L1*, *EGR1*, *CASP8*, and *CASP1*, proliferation

markers such as *PCNA*, and heat shock response factors such as *HSPA4*, *HSPH1*, and *HSP90AA2P*

were completely absent in samples obtained from cells exposed to the pre-irradiated virus.

Strikingly, expression of seven specific genes (*CDKN1A*, *CYP1A1*, *MDM2*, *HMOX1*, *RAD50*,

HSPA1L, and *E2F1*) was modulated uniquely in response to exposure to sUV-preirradiated HCoV-

NL63, a finding consistent with gene expression changes responsive to sUV-induced chemical

damage to viral components (including ribonucleic acids, proteins, and lipids) [1-3].

268   >Figure 3<

3.4. Dose-response relationship of solar simulated UV-induced inhibition of SARS-CoV-2

infectivity targeting Vero and Calu-3 mammalian cells

After demonstrating HCoV-NL63 coronavirus inactivation by sUV at an environmentally

relevant dose level, we examined whether sUV-inactivation might also be applicable to SARS-

CoV-2. To this end, we exposed the virus with a dose range of sUV, subsequently infected Vero

monkey epithelial cells at two different multiplicities of infection (MOIs, high versus low titer),

and measured the number of infectious virions three days later by plaque forming assay.

Strikingly, as observed with HCoV-NL63, sUV exposure caused a pronounced suppression of viral

infectivity. This antiviral effect, observable over a broad range of sUV doses, followed an

exponential decay curve with an effective ED_{50} (sUV dose diminishing SARS-CoV-2 viral

infectivity by 50%) approximating 55 mJ/cm^2 (low titer) and 62 mJ/cm^2 (high titer) (Fig. 4A).

Next, we tested feasibility of achieving complete inhibition of SARS-CoV-2 replication by high

dose sUV [UVB portion: 1010 mJ/cm^2, a maximum dose level similar to the one used in the HCoV-

NL63-directed dose-response experiments (Fig. 1C)]. To this end, we pre-exposed SARS-CoV-2

to sUV and measured the amount of viral RNA (corresponding to the region of the viral genome

encoding the S protein) by RT-qPCR analysis. Indeed, complete inhibition was achieved at that

dose (Fig. 4B). We obtained similar results for sUV-exposed SARS-CoV-2 infections of Calu-3

human lung epithelial target cells with viral load in supernatants being monitored over three days

by RT-qPCR (Fig. 4C). Taken together, we conclude that SARS-CoV-2 is sensitive to sUV

suggesting viral inactivation at environmentally relevant exposure levels.

291   >Figure 4<

3.5. Solar simulated UV exposure of SARS-CoV-2 prevents stress response gene expression

elicited by viral infection of Calu-3 human epithelial lung cells as detected by array analysis

Next, to determine Calu-3 human epithelial lung cell stress response gene expression elicited

by SARS-CoV-2 as a function of viral pre-exposure to sUV, we employed expression analysis

using the Human Stress and Toxicity PathwayFinder™ PCR Array technology. To this end, we

infected Calu-3 target cells with sUV or mock-treated virus as outlined before, followed by

comparative gene expression profiling at the end of the experiment. We observed multiple SARS-

CoV-2-induced expression changes (antagonized by viral pre-exposure to sUV) as shown in the

Volcano plot depiction [displaying statistical significance (P value) versus magnitude of change

(fold change)] (Fig. 5). SARS-CoV-2 infection caused a pronounced upregulation of stress

response gene expression including genes encoding key regulators of inflammatory signaling

including *IL1A, IL1B*, *IL6* , *TNF*, *CCL3, CXCL10, CSF2, and NFKBIA*, oxidative stress defense

such as *SOD2*, and heat shock response such as *HSPA6* (Fig. 5). In contrast, after infection with

sUV-exposed virus, most of these infection-associated expression changes were either attenuated

or completely obliterated, an observation consistent with pronounced suppression of SARS-CoV-

2 infectivity as a consequence of sUV-exposure. Remarkably, these expression changes closely

mirrored those observed in response to HCoV-NL63 infection that occurred with or without viral

exposure to sUV (Fig. 3).

312   >Figure 5<

Likewise, we observed a striking similarity between the gene expression changes elicited by

HCoV-NL63 and SARS-CoV-2 (and blocked by viral sUV pre-exposure), modulating redox,

inflammatory, and proteotoxic stress responses in Calu-3 human epithelial lung cells (Fig. 6).

Specifically, sUV-induced (UVB portion: 706 mJ/cm^2) viral inactivation was apparent from

independent RT-qPCR assessment of mRNA levels ('no sUV' versus 'sUV') interrogating genes

encoding key regulators of redox (*SOD2*), inflammatory (*IL1B*, *TNF*, *CCL3*, *IL6*, *CSF2*), and

proteotoxic ('heat shock'; *HSPA6*) stress responses in Calu-3 target cells as detailed above. Thus,

our data suggest that similar to HCoV-NL63, sUV exposure of SARS-CoV-2 interrupts the viral

life cycle causing suppression of viral replication and virus-induced inflammatory and cellular

stress responses in mammalian target cells.

325   >Figure 6<

4. Discussion

Identification and mechanistic exploration of environmental factors that might determine

coronavirus infectivity are of significant interest with relevance to both basic molecular research

and public health-related preventive and interventional investigations [2]. Here, we have explored

for the first time the effects of full spectrum (UVA + UVB) solar ultraviolet radiation on

coronavirus infectivity and demonstrate that sUV inactivates HCoV-NL63 and SARS-CoV-2

coronaviruses at environmentally relevant doses. First, we observed that exposure of HCoV-NL63

and SARS-CoV-2 to sUV (performed at acute dose levels relevant to human populations

worldwide) blocks subsequent viral infection and replication in relevant primate target cells

[human: Calu-3 lung epithelial, Caco-2 colorectal epithelial; monkey: Vero kidney epithelial (Figs.

1, 2, 4)]. Blockade of viral infectivity in response to sUV pre-exposure was also confirmed using

stress response gene expression profiling in array (Figs. 3, 5) and independent RT-qPCR format

(Fig. 6) elicited in Calu-3 target cells by coronavirus infection (HCoV-NL63 and SARS-CoV-2).

Remarkably, dose levels used throughout this pilot study are representative of terrestrial ground

level exposure suggesting environmental relevance, and significant coronavirus inactivation was

detectable even at low exposure levels expected to be beneath the cutaneous sunburn-inducing

threshold (Figs. 1, 2, 4) [2,6,8]. In this context, it is remarkable that recent research has already

indicated that ground level solar UV displays significant virucidal effects targeting coronaviruses

including SARS-CoV-2 [2,6,11,13]. However, the complexity of human exposure levels to solar

UV as a function of solar zenith angle, seasonality, spectral distribution, and latitude remain to be

addressed before any firm conclusions relevant to human populations can be drawn. Specifically,

the anti-viral activity of specific spectral components of sUV remains to be determined since the

light source employed in our prototype studies emitted full spectrum simulated solar UV, and the

action spectrum of virus inactivation by solar UV remains largely undefined. For example, it is

possible that the UVA portion of ground level sUV significantly contributes to the coronavirus-

directed effects described by us [7]. It therefore remains to be seen if indirect impairment of viral

structure and infectivity occurs by alternative mechanisms, such as UVA-driven photosensitization

and oxidative stress (mediated by formation of reactive oxygen species including singlet oxygen),

that might be operative in addition to direct inactivation of viral genomic RNA through nucleic

acid base photodamage. It will also be interesting to explore potential mechanistic synergisms

underlying virucidal effects that occur upon combined UVA and UVB as compared to separate

spectral exposure. Likewise, experimental conditions used throughout our studies (including viral

irradiation in PBS and exposure performed in cell culture medium) might limit the applicability of

our conclusions in the context of relevant coronavirus transmission situations that involve more

complex determinants of infectivity including the role air-borne and aerosol transmission and

intermediate surface retention [6].

Addressing urgency and global scale of the unfolding SARS-CoV-2 pandemic requires an

improved understanding of environmental factors that modify viral infectivity [2,6,8]. Taken

together, our data suggest feasibility of sUV-induced viral inactivation targeting HCoV-NL63 and

SARS-CoV-2 coronaviruses, a finding to be substantiated by future mechanistic exploration

performed in more relevant *in vivo* exposure models.

Funding

Supported in part by grants from the National Institutes of Health (R21ES029579, ES007091, ES006694, CA023074). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Acknowledgement

The authors like to acknowledge technical support provided by Jennifer L. Uhrlaub.

Declaration of competing interest

The authors state no conflict of interest.

References

[1]   N.G. Reed, The history of ultraviolet germicidal irradiation for air disinfection, Public Health Rep. 125 (2010) 15-27, 10.1177/003335491012500105.

[2]   L. Horton, A.E. Torres, S. Narla, A.B. Lyons, I. Kohli, J.M. Gelfand, D.M. Ozog, I.H. Hamzavi, H.W. Lim, Spectrum of virucidal activity from ultraviolet to infrared radiation, Photochem. Photobiol. Sci. 19 (2020) 1262-1270, 10.1039/d0pp00221f.

[3]   M. Hessling, K. Hones, P. Vatter, C. Lingenfelder, Ultraviolet irradiation doses for coronavirus inactivation - review and analysis of coronavirus photoinactivation studies, GMS Hyg. Infect. Control. 15 (2020) Doc08, 10.3205/dgkh000343.

[4]   C. Huang, Y. Wang, X. Li, L. Ren, J. Zhao, Y. Hu, L. Zhang, G. Fan, J. Xu, X. Gu, Z. Cheng, T. Yu, J. Xia, Y. Wei, W. Wu, X. Xie, W. Yin, H. Li, M. Liu, Y. Xiao, H. Gao, L. Guo, J. Xie, G. Wang, R. Jiang, Z. Gao, Q. Jin, J. Wang, B. Cao, Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China, Lancet. 395 (2020) 497-506, 10.1016/S0140-6736(20)30183-5.

[5]   N. Storm, L.G.A. McKay, S.N. Downs, R.I. Johnson, D. Birru, M. de Samber, W. Willaert, G. Cennini, A. Griffiths, Rapid and complete inactivation of SARS-CoV-2 by ultraviolet-C irradiation, Sci. Rep. 10 (2020) 22421, 10.1038/s41598-020-79600-8.

[6]   S. Ratnesar-Shumate, G. Williams, B. Green, M. Krause, B. Holland, S. Wood, J. Bohannon, J. Boydston, D. Freeburger, I. Hooper, K. Beck, J. Yeager, L.A. Altamura, J. Biryukov, J. Yolitz, M. Schuit, V. Wahl, M. Hevey, P. Dabisch, Simulated Sunlight Rapidly Inactivates SARS-CoV-2 on Surfaces, J. Infect. Dis. 222 (2020) 214-222, 10.1093/infdis/jiaa274.

[7]   A. Rezaie, G.G.S. Leite, G.Y. Melmed, R. Mathur, M.J. Villanueva-Millan, G. Parodi, J. Sin, J.F. Germano, W. Morales, S. Weitsman, S.Y. Kim, J.H. Park, S. Sakhaie, M. Pimentel,

Ultraviolet A light effectively reduces bacteria and viruses including coronavirus, PLoS One. 15 (2020) e0236199, 10.1371/journal.pone.0236199.

[8] H.A. Aboubakr, T.A. Sharafeldin, S.M. Goyal, Stability of SARS-CoV-2 and other coronaviruses in the environment and on common touch surfaces and the influence of climatic conditions: A review, Transbound. Emerg. Dis. 68 (2021) 296-312, 10.1111/tbed.13707.

[9] G.T. Wondrak, M.K. Jacobson, E.L. Jacobson, Endogenous UVA-photosensitizers: mediators of skin photodamage and novel targets for skin photoprotection, Photochem. Photobiol. Sci. 5 (2006) 215-237, 10.1039/b504573h.

[10] J. Herman, B. Biegel, L. Huang, Inactivation times from 290 to 315 nm UVB in sunlight for SARS coronaviruses CoV and CoV-2 using OMI satellite data for the sunlit Earth, Air Qual. Atmos. Health. (2020) 1-17, 10.1007/s11869-020-00927-2.

[11] A. Seyer, T. Sanlidag, Solar ultraviolet radiation sensitivity of SARS-CoV-2, Lancet Microbe. 1 (2020) e8-e9, 10.1016/S2666-5247(20)30013-6.

[12] K.P. Marik P.E., and Varonc J., Does vitamin D status impact mortality from SARS-CoV-2 infection?, Med. Drug Discov. 6 (2020) 100041, 10.1016/j.medidd.2020.100041.

[13] S.G. Nicastro F., Antonello E., Bianco A., Biasin M., Brucato J.R., Ermolli I., Pareschi G., Salvati M., Tozzi P., Trabattoni D., and Clerici M., Modulation of COVID-19 Epidemiology by UV-B and –A Photons from the Sun, medRxiv (2020) https://doi.org/10.1101/2020.06.03.20121392.

[14] R.K. Moozhipurath, L. Kraft, B. Skiera, Evidence of protective role of Ultraviolet-B (UVB) radiation in reducing COVID-19 deaths, Sci. Rep. 10 (2020) 17705, 10.1038/s41598-020-74825-z.

[15] P. Herzog, C. Drosten, M.A. Muller, Plaque assay for human coronavirus NL63 using human colon carcinoma cells, Virol. J. 5 (2008) 138, 10.1186/1743-422X-5-138.

[16] H.R. Jonsdottir, R. Dijkman, Coronaviruses and the human airway: a universal system for virus-host interaction studies, Virol. J. 13 (2016) 24, 10.1186/s12985-016-0479-5.

[17] L. van der Hoek, K. Sure, G. Ihorst, A. Stang, K. Pyrc, M.F. Jebbink, G. Petersen, J. Forster, B. Berkhout, K. Uberla, Human coronavirus NL63 infection is associated with croup, Adv. Exp. Med. Biol. 581 (2006) 485-491, 10.1007/978-0-387-33012-9_86.

[18] T.J. Ripperger, J.L. Uhrlaub, M. Watanabe, R. Wong, Y. Castaneda, H.A. Pizzato, M.R. Thompson, C. Bradshaw, C.C. Weinkauf, C. Bime, H.L. Erickson, K. Knox, B. Bixby, S. Parthasarathy, S. Chaudhary, B. Natt, E. Cristan, T. El Aini, F. Rischard, J. Campion, M. Chopra, M. Insel, A. Sam, J.L. Knepler, A.P. Capaldi, C.M. Spier, M.D. Dake, T. Edwards, M.E. Kaplan, S.J. Scott, C. Hypes, J. Mosier, D.T. Harris, B.J. LaFleur, R. Sprissler, J. Nikolich-Zugich, D. Bhattacharya, Orthogonal SARS-CoV-2 Serological Assays Enable Surveillance of Low-Prevalence Communities and Reveal Durable Humoral Immunity, Immunity. 53 (2020) 925-933 e924, 10.1016/j.immuni.2020.10.004.

[19] M. Rojo de la Vega, D.D. Zhang, G.T. Wondrak, Topical Bixin Confers NRF2-Dependent Protection Against Photodamage and Hair Graying in Mouse Skin, Front. Pharmacol. 9 (2018) 287, 10.3389/fphar.2018.00287.

[20] R. Justiniano, L. de Faria Lopes, J. Perer, A. Hua, S.L. Park, J. Jandova, M.S. Baptista, G.T. Wondrak, The Endogenous Tryptophan-derived Photoproduct 6-formylindolo[3,2-b]carbazole (FICZ) is a Nanomolar Photosensitizer that Can be Harnessed for the Photodynamic Elimination of Skin Cancer Cells in Vitro and in Vivo, Photochem. Photobiol. 97 (2021) 180-191, 10.1111/php.13321.

[21] A. Baer, K. Kehn-Hall, Viral concentration determination through plaque assays: using
traditional and novel overlay systems, J. Vis. Exp. (2014) e52065, 10.3791/52065.
[22] B. Malhotra, M.A. Swamy, P.V. Reddy, N. Kumar, J.K. Tiwari, Evaluation of custom
multiplex real - time RT - PCR in comparison to fast - track diagnostics respiratory 21
pathogens kit for detection of multiple respiratory viruses, Virol. J. 13 (2016) 91,
10.1186/s12985-016-0549-8.
[23] S.D. Lamore, G.T. Wondrak, Zinc pyrithione impairs zinc homeostasis and upregulates
stress response gene expression in reconstructed human epidermis, Biometals. 24 (2011)
875-890, 10.1007/s10534-011-9441-6.
[24] J. Jandova, G.T. Wondrak, Genomic GLO1 deletion modulates TXNIP expression, glucose
metabolism, and redox homeostasis while accelerating human A375 malignant melanoma
tumor growth, Redox Biol. 39 (2021) 101838, 10.1016/j.redox.2020.101838.

 Figure legends

 Figure 1. Solar simulated UV pre-exposure antagonizes HCoV-NL63 viral infectivity

 targeting Calu-3 human epithelial lung cells. Virus in PBS was exposed to sUV or left

 unexposed followed by Calu-3 target cell infection (0.01 MOI) and post infection culture over 7

 days followed by analysis. (A) Spectral power distribution (irradiance) of the solar simulator light

 source equipped with appropriate cut-off filter (sUV: UVB + UVA, solid black line). (B) Plaque

 assay after viral exposure to sUV (UVB portion: 706 mJ/cm^2) as visualized by light microscopy

 (10 x magnification); bar graph summarizes numerical data. (C) RT-qPCR of viral genome

 replication in target cells [left panel: amplification curves as a function of sUV dose (UVB portion

 as indicated); right panel: bar graph summarizing numerical data].

 Figure 2. Solar simulated UV pre-exposure antagonizes HCoV-NL63 viral infectivity

 targeting Caco-2 human epithelial colorectal cells. Virus in PBS was exposed to sUV or left

 unexposed followed by Caco-2 target cell infection (0.01 MOI) and post infection culture (7 days)

 followed by analysis. (A) Plaque assay after viral exposure to sUV (UVB portion: 706 mJ/cm^2)

 as visualized by light microscopy (10 x magnification); bar graph summarizes numerical data. (B)

 RT-qPCR detection of viral genome replication in target cells; left panel: amplification curves (as

 a function of sUV-dose); right panel: bar graph summarized numerical data.

 Figure 3. Solar simulated UV pre-exposure of HCoV-NL63 prevents stress response gene

 expression elicited in Calu-3 human epithelial lung target cells. Treatments were performed as

 detailed in Fig. 1. (A) Target cell stress response [control (HCoV-NL63) versus sUV (UVB

 portion: 706 mJ/cm^2) pre-exposed virus] assessed by *RT^2 Profiler^{TM} Stress and Toxicity Pathway*

gene expression array analysis [volcano plot depiction: p value over log2 (fold expression change)]. (B) Scatter plot depiction comparing expression changes elicited by untreated control virus (top panel) or sUV pre-exposed virus (bottom panel). (C) Venn diagram depicting expression changes induced by mock-irradiated virus (control) versus sUV pre-irradiated virus. (D) Tabular summary of numerical values specifying gene expression changes at the mRNA level ($p<0.05$).

Figure 4. Solar simulated UV exposure of SARS-CoV-2 antagonizes subsequent viral infection and replication in African green monkey Vero and Calu-3 human epithelial lung cells. (A) SARS-CoV-2 was sUV-irradiated (UVB portion: up to 480 mJ/cm^2; or remained unirradiated) in PBS and subsequently used to infect Vero cells at two different MOIs (high versus low titer). Dose response of plaque formation as a function of sUV pre-exposure dose was assessed; a representative experiment (left panel, top and bottom rows) and quantification (right panels) are depicted. (B) Detection of viral genome replication in Vero cells with quantification of viral RNA after infection using mock or sUV pre-irradiated virus (UVB portion: 1010 mJ/cm^2) as assessed by RT-qPCR after 24 h. (C) Infection of Calu-3 cells with SARS-CoV-2 [sUV pre-exposed (UVB portion: 706 mJ/cm^2) versus unirradiated virus]. The presence of infectious virions in the supernatants was quantified over the course of three days post infection by RT-qPCR (nd: not detectable).

Figure 5. Solar simulated UV pre-exposure of SARS-CoV-2 prevents stress response gene expression elicited in Calu-3 human epithelial lung target cells. Treatment and analysis were performed as detailed in Fig. 3. (A) Target cell stress response [control (SARS-CoV-2) versus sUV (UVB portion: 706 mJ/cm^2)-preirradiated virus] assessed by *RT^2 Profiler^{TM} Stress and*

Toxicity Pathway gene expression array analysis [volcano plot depiction: p value over log2 (fold

expression change)]. (B) Scatter plot depiction comparing expression changes elicited by untreated

control virus (top panel) or sUV pre-exposed virus (bottom panel). (C) Venn diagram depicting

expression changes induced by mock-irradiated virus (control) versus sUV pre-irradiated virus.

(D) Tabular summary of numerical values of gene expression changes at the mRNA level ($p<0.05$).

Figure 6. Comparative analysis of redox, inflammatory, and proteotoxic stress response gene

expression in Calu-3 human epithelial lung cells elicited by HCoV-NL63 and SARS-CoV-2

(with and without viral sUV pre-exposure). Gene expression as assessed by single RT-qPCR

quantification in virus-exposed target cells as a function of viral pre-exposure ['no sUV' versus

'sUV' (UVB portion: 706 mJ/cm^2)]. Bar graphs depict fold change ('sUV' versus 'no sUV')

normalized to housekeeping gene expression (*RPS18*; gray bar: no sUV pretreatment; black bar:

sUV-pretreatment).

Fig. 1

Calu-3

A solar simulated UV [sUV]: spectral power distribution

B HCoV-NL63 viral plaques

solar simulated UV

+	+	HCoV-NL63
-	+	sUV

C HCoV-NL63 viral RNA

sUV [UVB (mJ/cm^2)]

0 — 13 — 240 — 480 — 960

HCoV-NL63 viral genome [vs. non-UV control]

sUV [UVB (mJ/cm^2)]

Fig. 2

Fig. 3

C

HCoV-NL63

no sUV			CXCL10	post sUV
BCL2L1	HSP90AA2P	NFKBIA		CDKN1A
CASP1	HSPA4	PCNA		CYP1A1
CASP8	HSPA6	PTGS1		E2F1
CCL3	HSPH1	SOD2		HMOX1
CSF2	IL1A	TNF		HSPA1L
EGR1	IL1B	UGT1A4		MDM2
GDF15	IL6	UNG		RAD50
21 genes			*1 gene*	*7 genes*

D

Calu-3

HCoV-NL63

Gene	Description	fold change (p≤0.05)	
		no sUV	post sUV
CSF2	Colony stimulating factor 2 (granulocyte-macrophage)	2452.4	-
TNF	Tumor necrosis factor	81.6	-
IL1B	Interleukin 1, beta	48.3	-
IL1A	Interleukin 1, alpha	45.9	-
SOD2	Superoxide dismutase 2, mitochondrial	27.3	-
CCL3	Chemokine (C-C motif) ligand 3	12.9	-
HSPA6	Heat shock 70kDa protein 6 (HSP70B')	10.8	-
CXCL10	Chemokine (C-X-C motif) ligand 10	7.4	3.6
NFKBIA	NFKB inhibitor alpha	6.5	-
CYP1A1	Cytochrome P450, family 1, subfamily A, polypeptide 1	-	5.5
IL6	Interleukin 6 (interferon, beta 2)	4.9	-
CDKN1A	Cyclin-dependent kinase inhibitor 1A (p21, Cip1)	-	3.0
MDM2	Mdm2 p53 binding protein homolog	-	2.4
HMOX1	Heme oxygenase (decycling) 1	-	-2.1
PTGS1	Prostaglandin-endoperoxide synthase 1	-2.4	-
RAD50	RAD50 homolog	-	-2.4
CASP1	Caspase 1, apoptosis-related cysteine peptidase	-3.0	-
HSPA1L	Heat shock 70kDa protein 1-like	-	-3.1
E2F1	E2F transcription factor 1	-	-4.0
UGT1A4	UDP glucuronosyltransferase 1 family, polypeptide A4	-4.9	-
HSP90AA2P	Heat shock protein 90kDa alpha, class A member 2	-5.8	-
PCNA	Proliferating cell nuclear antigen	-6.3	-
GDF15	Growth differentiation factor 15	-6.7	-
UNG	Uracil-DNA glycosylase	-7.3	-
HSPH1	Heat shock 105kDa/110kDa protein 1	-8.4	-
CASP8	Caspase 8, apoptosis-related cysteine peptidase	-9.8	-
HSPA4	Heat shock 70kDa protein 4	-10.3	-
EGR1	Early growth response 1	-40.5	-
BCL2L1	BCL2-like 1	-1078.6	-

Fig. 4

A

Vero

high titer

low titer

0   13   60   120   240   480

sUV [UVB (mJ/cm²)]

high titer

$ED_{50} = 62.4$

(95% CI 54.2 - 73.3)

$R^2 = 0.99$

pfu/well

sUV [UVB (mJ/cm²)]

low titer

$ED_{50} = 54.9$

(95% CI 45.2 - 69.8)

$R^2 = 0.97$

pfu/well

sUV [UVB (mJ/cm²)]

B

Vero

SARS-CoV-2 viral genome [relative abundance (fold)]

\-   +   **sUV**

***

C

Calu-3

infectious SARS-CoV-2 in supernatant [pfu/well]

nd   nd   nd

\-   +   \-   +   \-   +   **sUV**

day 1   day 2   day 3

***   ***

Fig. 5

B

2^{-ΔCt} [SARS-CoV-2] vs 2^{-ΔC} [untreated], sUV − and +

D

Gene	Description	SARS-CoV-2 fold change (p≤0.05)	
		no sUV	post sUV
IL1A	Interleukin 1, alpha	982.3	26.9
CXCL10	Chemokine (C-X-C motif) ligand 10	654.8	2.0
IL1B	Interleukin 1, beta	469.5	-
TNF	Tumor necrosis factor	284.0	-
CCL3	Chemokine (C-C motif) ligand 3	230.7	-
IL6	Interleukin 6 (interferon, beta 2)	89.6	-
CSF2	Colony stimulating factor 2 (granulocyte-macrophage)	51.1	-
NFKBIA	NFKB inhibitor alpha	50.6	-
CCL4	Chemokine (C-C motif) ligand 4	48.2	-
EPHX2	Epoxide hydrolase 2, cytoplasmic	37.4	-
TNFSF10	Tumor necrosis factor (ligand) superfamily, member 10	36.1	-
SOD2	Superoxide dismutase 2, mitochondrial	29.9	2.7
UGT1A4	UDP glucuronosyltransferase 1 family, polypeptide A4	23.4	-
CYP1A1	Cytochrome P450, family 1, subfamily A, polypeptide 1	23.2	-
MT2A	Metallothionein 2A	20.5	-
LTA	Lymphotoxin alpha (TNF superfamily, member 1)	19.6	-
HSP90AA2P	Heat shock protein 90kDa alpha, class A member 2	19.0	-
XRCC2	X-ray repair cross complementing 2	18.2	4.4
NOS2	Nitric oxide synthase 2, inducible	17.1	-
HSPA6	Heat shock 70kDa protein 6 (HSP70B')	11.5	-
HSPE1	Heat shock 10kDa protein 1 (chaperonin 10)	11.2	2.9
CRYAB	Crystallin, alpha B	9.3	-
PRDX1	Peroxiredoxin 1	9.1	-
HSPH1	Heat shock 105kDa/110kDa protein 1	8.5	2.2
CASP1	Caspase 1, apoptosis-related cysteine peptidase	7.5	-
SOD1	Superoxide dismutase 1, soluble	6.9	-
HSPA1L	Heat shock 70kDa protein 1-like	6.3	-
SERPINE1	Serpin peptidase inhibitor, clade E, member 1	6.0	-
PTGS1	Prostaglandin-endoperoxide synthase 1	5.5	-
FMO5	Flavin containing monooxygenase 5	5.5	-
HSPA4	Heat shock 70kDa protein 4	4.6	-
CASP10	Caspase 10, apoptosis-related cysteine peptidase	4.6	-
PCNA	Proliferating cell nuclear antigen	4.5	-
HSPA2	Heat shock 70kDa protein 2	4.4	-
NFKB1	Nuclear factor of kappa 1	3.9	-
DNAJA1	DnaJ (Hsp40) homolog, subfamily A, member 1	3.8	-
HSP90AB1	Heat shock protein 90kDa alpha, class B member 1	2.7	-
HSPA8	Heat shock 70kDa protein 8	2.3	-
IL18	Interleukin 18 (interferon-gamma-inducing factor)	2.2	-
GDF15	Growth differentiation factor 15	-3.1	-
EGR1	Early growth response 1	-12.4	-
RAD23A	RAD23 homolog A (S. cerevisiae)	-	-2.8
HSPA5	Heat shock 70kDa protein 5	-	-3.8
HSPB1	Heat shock 27kDa protein 1	-	-4.3

Fig. 6