Essential role of translation factor eIF5a in cytokine production and cell cycle regulation in primary CD8 T lymphocytes

Thomas CJ Tan[1], Van Kelly[2], Xiaoyan Zhou[1], Tony Ly[2,3] and *Rose Zamoyska[1]

Author list footnotes

1. Institute of Immunology and Infection Research, University of Edinburgh, Ashworth Laboratories, Charlotte Auerbach Road, Edinburgh, EH9 3FL, United Kingdom.
2. Wellcome Trust Centre for Cell Biology, University of Edinburgh, Michael Swann Building, Max Born Crescent, Edinburgh, EH9 3BF, United Kingdom.
3. Centre for Gene Regulation and Expression, Life Sciences Research Complex, University of Dundee, Dundee, DD1 5EH, United Kingdom.

Contact info

*. Correspondence: Rose.Zamoyska@ed.ac.uk

Abstract

Translational control adjusts protein production rapidly and facilitates local cellular responses to environmental conditions. Translation can be regulated through sequestration of mRNAs by regulatory proteins or RNAs, but also by the availability of ribosomes and translation factors which enable initiation and elongation of nascent polypeptides. Traditionally initiation of mRNA translation has been considered to be a major translational control point, however, control of peptide elongation can also play a role. Here we show that post-translational modification of the elongation factor, eIF5a, controls translation of subsets of proteins in naïve T-cells upon activation. Sequencing of nascent polypeptides indicated that functional eIF5a was required for the production of proteins which regulate T-cell proliferation and effector function. Control of

translation in multiple immune cell lineages is required to co-ordinate immune responses and these data illustrate that translational elongation can contribute to post-transcriptional regulons important for the control of inflammation.

Keywords

eIF5a; nascent proteomics; CRISPR-Cas9; translation control; T cell; immunosenescence; GC7

INTRODUCTION

Regulation of protein expression at the level of ribosome production and translation can rapidly rewire and co-ordinate responses to changing environmental cues. Poor correlations between transcriptomes and proteomes have been documented in numerous cell types, including stem cells (Munoz et al., 2011) and T lymphocytes (Hukelmann et al., 2015) indicating that a substantial proportion of their protein production is regulated translationally. Both these lineages can undergo rapid expansions and changes in cell identity in response to environmental triggers (Gabut et al., 2020). Regulation of ribosome biogenesis and global mRNA translation (Gabut et al., 2020), contribute to this plasticity together with more targeted post-transcriptional regulons, important for directing the production of proteins with related functions or involved in common biological pathways (Anderson, 2010). Regulation of such subsets of mRNAs frequently involves interactions with RNA-binding proteins or non-coding RNAs (eg miRNA and lncRNA) (Ciafrè and Galardi, 2013; Zhang et al., 2019b). In addition alterations in ribosome constituents or the proteins involved with the translation of mRNAs may play a role. For example, we showed that for naïve CD8 T cells the degree of ribosome biogenesis was dramatically affected by the extent of activation through the T cell receptor (TCR), which impacted global translation and also selectively affected production of some groups of proteins over others (Tan et al., 2017). How this plasticity of translational control is orchestrated around the ribosome is not well understood, but a number of proteins are directly involved with the ribosome during translation (Dever et al., 2018; Sonenberg and Dever, 2003) and these may themselves be regulated both translationally and post-translationally.

One such protein is the translation factor eIF5a, which was originally identified as a translation initiation factor. However, recent work in yeast has indicated a role in translation elongation and termination (Schuller et al., 2017). Depletion of eIF5a in *S. cerevisiae* reduced protein synthesis by approximately 30%, raising the possibility that eIF5a is required for translation of subsets of proteins, rather than being a ubiquitous translation factor (Kang and Hershey, 1994). *In vivo* reporter and *in vitro* translation assays revealed that eIF5a is required for efficient peptide bond formation between consecutive proline residues, a function described for its bacterial homolog, EF-P (Gutierrez et al., 2013). Two independent yeast ribosome profiling studies found the requirement for eIF5a is not limited to elongation between poly-proline sequences, but that it also influences translation of glycine and charged amino acids, suggesting a wider involvement of eIF5a in regulating translation of mRNAs with common motifs, (Pelechano and Alepuz, 2017; Schuller et al., 2017) and suggesting it might be an important player in influencing post-transcriptional regulons.

eIF5a isoforms in eukarya and archaea are the only proteins with post-translational attachment of a hypusine residue on Lys[50-51], a modification that is essential for eIF5a function (Park and Wolff, 2018). Hypusination involves enzymatic transfer of the 4-aminobutyl moiety of the polyamine spermidine to lysine, catalysed by deoxyhypusine synthase (DHPS) to produce the deoxyhypusinated intermediate form, and subsequent hydroxylation by deoxyhypusine hydroxylase (DOHH) to form the mature hypusinated protein. The site of hypusination is in contact with the acceptor stem of the P site tRNA on 80S ribosome (Schmidt et al., 2016), and it is postulated that hypusination is required for eIF5a activity as non-modified eIF5a is inefficient in facilitating peptide synthesis (Gutierrez et al., 2013) and inhibition of the hypusination pathway impairs proliferation in mammalian cell lines (Colvin et al., 2013; Hanauske-Abel et al., 1994). Further evidence for the importance of eIF5a hypusination comes from studies of spermidine, a natural polyamine which is the donor moiety for eIF5a hypusination and whose abundance is reduced in ageing cells. Specifically, spermidine dietary supplementation restored both the abundance of eIF5a hypusination and age-related T cell impairments (Puleston et al., 2014). Spermidine administration also extended the life span of yeast, flies, worms, and cultured human PBMCs (Eisenberg et al., 2009). Therefore strong evidence links the abundance of

eIF5a hypusination and its function in age-related immunosenescence, although it is currently unclear how these processes are associated.

The aim of this study was to determine firstly how the activity of eIF5a itself was regulated in T lymphocytes, and secondly, whether eIF5a has a selective role in regulating translation of specific subsets of mRNA during T cell activation and differentiation from naïve to effector cells. We used CRISPR knockout of eIF5a itself and of the key enzymes, DHPS and DOHH, involved in adding the hypusine modification to eIF5a and compared the effect of disrupting the latter to inhibition of hypusination by the deoxyhypusine synthase inhibitor, GC7. We show that although naïve T cells express abundant eIF5a it is largely unmodified by hypusination and therefore of limited functionality. Upon activation of T cells, DHPS and DOHH are expressed rapidly and hypusinated eIF5a becomes abundant. Total loss of eIF5a surprisingly increased production of some ribosomal proteins and eIF6, a possible translational inhibitor, while decreasing some translation initiation factors, together suggesting alterations in ribosome composition. Mass spectrometry analysis of nascent peptide production in these cells confirmed unequal impacts on synthesis of specific sets of proteins in the absence of eIF5a, particularly those involved in cell cycle regulation and cytokine production. We confirmed that loss of functional eIF5a resulted in a specific loss of proteins containing polyproline motifs as described previously. However treatment with GC7 did not faithfully reproduce the phenotypes of knockout cells indicating a significant off-target impact of this drug. Overall our data are consistent with a role for eIF5a in co-ordinating post-transcriptional regulons in T lymphocytes and reveal how these may influence primary T cell function.

RESULTS

T cell activation enhances production and functional maturation of eIF5a

A unique feature of T cells is that they exist for long periods as naïve cells which are functionally immature, non-dividing and metabolically inert, with barely detectable macromolecular synthesis. Upon stimulation through their antigen specific TCRs they rapidly turn on gene transcription and protein synthesis. In the initial 24h following stimulation CD8 T cells increase cytoplasmic mass by approximately 4-fold, indicating a rapid initiation of macromolecular synthesis, before commencing a burst of proliferation with a division time as short as 2-4h (Lawrence and Braciale, 2004; Yoon et al., 2007).

We examined the functional maturation of eIF5a in OT-1 CD8 T cells which express a single specificity transgenic TCR that recognises a peptide from chicken ovalbumin (SIINFEKL) presented by the MHC allele, H-2D[b]. Western blot of whole cell lysates from naïve unstimulated OT-1 T cells showed abundant expression of eIF5a, but very little that was modified by hypusination, as revealed by the paucity of staining with a hypusine specific Ab (Fig 1A and B). Upon stimulation of the T cells with peptide, the abundance of hypusinated eIF5a increased by ~50-fold, concurrent with ~10-fold increase in expression of DHPS, the enzyme that catalyses transfer of the 4-aminobutyl moiety of spermidine to eIF5a. The abundance of both hypusinated eIF5a and DHPS peaked approximately 48h after stimulation. There was a modest increase in the abundance of eIF5a, consistent with the increase of a control protein, ZAP-70, which reflected the increase in overall protein abundance upon activation.

Abrogation of eIF5a or its maturation pathway differentially disrupts key cellular functions

Hypusination of eIF5a allows efficient facilitation of translation elongation *in vitro* (Gutierrez et al., 2013) and may be responsible for its cytoplasmic localisation (Lee et al., 2009). However it is unclear whether or not the non-hypusinated precursor and deoxyhypusinated intermediate forms of eIF5a have any activity in primary cells. This question is important as it is uncertain whether targeting DHPS and DOHH as means to inhibit eIF5a are optimal strategies, given deletion of budding yeast DOHH had only a mild phenotype (Park et al., 2006) whereas embryonic targeting of DOHH in mouse

was lethal (Sievert et al., 2014). Moreover most studies of eIF5a function utilise the drug GC7 to inhibit hypusination and, therefore, function of eIF5a both *in vivo* and *in vitro*. GC7 is a spermidine derivative which competes with spermidine for binding to DHPS, thereby preventing the conversion of non-hypusinated eIF5a precursor to deoxyhypusinated intermediate, but it is undocumented whether GC7 has effects beyond inhibiting eIF5a.

In order to address these questions, naïve OT-1 cells were activated for 2 days, and then either treated with GC7 for 2 days, or transfected with CRISPR-Cas9 guides to disrupt *eIF5a* or *Dhps* or *Dohh* genes. KO cells were recovered after 3 days (*eIF5a* KO) or 4 days (*Dhps* and *Dohh* KO), with the duration of the culture adjusted for the different KOs to maximise viable cell recovery. CRIPSR-Cas9 transfection controls were targeted for the unrelated surface molecule, Thy 1. Given targeting efficiency was maximally 80%, both gene KO and untargeted cells from the same culture could be compared directly in the analysis by using intracellular staining for eIF5a or hypusine to identify positive cells. Fig 2A shows a representative experiment (n≥7) with a mean percentage KO of *eIF5a*, *Dhps*, and *Dohh* of 38.4%, 71.6% and 78.9%, respectively. eIF5a depletion is known to affect cellular proliferation (Kang and Hershey, 1994; Mathews and Hershey, 2015), and this likely accounts for the apparently reduced efficiency of targeting of eIF5a itself, as KO cells will be at a proliferative disadvantage relative to untargeted cells in the same culture. There was a decrease also in the number of cells recovered from cultures in which hypusination was impaired, including GC7-treated and *Dhps* and *Dohh* KOs relative to control populations, but this was not as profound as for the *eIF5a* KO.

To further characterise any proliferation defect, cell cycle analysis of was performed by staining with Hoechst 33342. Surprisingly GC7 inhibition of eIF5a hypusination resulted in cell cycle profiles that were distinct from those observed following knockout of *eIF5a*, *Dhps*, or *Dohh*. Fig 2A shows a pronounced cellular accumulation in S phase following GC7 treatment which was less apparent in the KOs and suggests possible GC7 off-target effects beyond its impact on eIF5a hypusination. Knockout of *Dohh* did not result in noticeable changes in the cell cycle profile despite negatively impacting hypusination and cell numbers and suggested the deoxyhypusinated intermediate

form of eIF5a is partially active, possibly slowing progression rather than arresting the cell cycle.

Cells lacking eIF5a, DHPS or DOHH were further analysed for their protein synthesis rates and cytokine production. After CRISPR-Cas9 transfection and maintenance in IL-2 cytokine, the cells were re-stimulated with peptide for 4h, in order to reactivate effector cytokine transcription and translation, and labelled with puromycin during the last 20 minutes. Incorporation of puromycin into nascent proteins can be analysed by flow cytometry and used to assess the rate of global protein synthesis. In addition, the production of specific CD8 T cell effector proteins, IFNγ and TNFα were measured by intracellular staining (Fig 2B). *eIF5a* KO cells reduced puromycin incorporation by ~84% while *Dhps* and *Dohh* KO cells both decreased synthesis by slightly less compared to untargeted cells (n=6). Cytokine production showed dramatic decrease in IFNγ production in *eIF5a* KO cells, with the majority of the population falling in the negative gate (Fig 2B). In contrast the reduction in IFNγ production was less pronounced in *Dhps* KO cells (~60% negative cells), while *Dohh* KO cells showed only a slight reduction in IFNγ (~25% negative cells). Interestingly the loss of IFNγ production was selective rather than reflecting a general shut down in cytokine translation, as all the targeted cells showed a much smaller, albeit significant, reduction in TNFα.

To determine at what stage of production these cytokines were regulated the abundance of IFNγ and TNFα mRNA was assessed. RT-qPCR was performed on cell lysates from unseparated *Thy1*, *Dhps* and *Dohh* targeted populations given the frequency of KO cells was routinely ≥80%. As the *eIF5a* targeted populations were closer to 50% of the total, the population was fixed for intracellular eIF5a staining and sorted into targeted (KO) and control (WT) subsets from the same culture (Fig 2C). In contrast to the non-restimulated control cells, all cells restimulated with peptide significantly upregulated IFNγ and TNFα mRNA as expected. Compared to their untargeted or *Thy-1* targeted control cells, IFNγ mRNA was decreased significantly in *eIF5a* KO and *Dhps* KO cells, but not in the *Dohh* targeted population. No significant reduction in TNFα mRNA abundance was observed in any population, however, curiously in the absence of DHPS TNFα mRNA was increased significantly. Taken together, these results indicate that lack of fully matured eIF5a does not impair TNFα

mRNA transcription and has a mild effect on its translation. By contrast, mature eIF5a is required both for transcription and translation of IFNγ. However, it is unclear whether this effect is direct, or else an indirect consequence of an eIF5a-dependent transcription factor.

Systematic analysis of eIF5a-regulated translation in CD8 T cells

To ask whether specific groups of proteins were regulated by functional eIF5a, global nascent peptide profiling by incorporation of 4-Azido-L-homoalanine (AHA) into nascent peptides in place of methionine was performed on *eIF5a* KO and GC7-treated T cells. Cultures were pulsed with AHA for 2h and *eIF5a* KO and WT populations were separated by cell sorting for eIF5a following fixation and staining, ensuring that untargeted WT cells were subjected to identical culture and labelling conditions as the KO cells. Following lysis of the sorted cells, AHA-labelled peptides were isolated using copper-mediated Click chemistry with Alkyne agarose beads and purified nascent proteomes were identified and quantified using LC-MS.

Single shot label-free proteomics identified approximately 5500 proteins that were at least 5 times more abundant in the AHA-labelled samples compared to non-labelled controls. Four biological replicate samples were used for each group (Fig 3A, left). iBAQ values were used to measure absolute quantities of protein comparisons between groups (Schwanhäusser et al., 2011). The iBAQ values were normalised further by the average of each sample within the same group to minimise variations caused by loading and acquisition, and then normalised again by the input cell number to allow comparison between groups (Fig 3A, right; Table S1). 3D MDS plot of all nascent proteins detected between *eIF5a* KO and WT cells and between GC7-treated and mock-treated control cells (Fig 3B) showed distinctive signatures separating the 4 groups with the largest source of variations (Dimension 1) separating KO/WT from GC7/Mock, most likely due to formaldehyde fixation and the subsequent reverse crosslinking in the KO/WT samples. The second largest source of variation (Dimension 2) separated KO from WT and GC7 from Mock so is most likely caused by the manipulation of eIF5a activity. Differential expression analyses were conducted between GC7- and mock-treated cells and between eIF5a KO and WT cells. GC7 had a systematic effect on protein production, with 3044 proteins down-regulated and only

31 proteins up-regulated (Fig S1A); while eIF5a KO had a more specific effect on protein production, with 369 down-regulated and 149 up-regulated proteins (Fig S1B).

Clustering of GO terms on molecular functions was performed (WebGestalt (Liao et al., 2019)) on the down- and up-regulated proteins in each comparison, compared to all MS-detected newly synthesised proteins as the background (Fig S1). Noticeably transcription factors, ribosomal proteins and translation factors were affected by both GC7 and *eIF5a* KO but in different ways (Fig 3C). Amongst all detected transcription factors from the mass spec data, notably production of E2F2 was profoundly impacted in *eIF5a* KO cells, which correlated with their impaired proliferation. Several key regulators of environmental response and cytokine production, including IRF4, RELA, NFκB1, STAT2, STAT4, STAT6, and TBET (Tbx21) were also downregulated in their production, with STAT2 being the most impacted. Some transcription factors were upregulated, such as the tumour suppressor p53, CNOT3 and TRPS1 which may control cell cycle (Shirai et al., 2019; Wu et al., 2014) and NFYA, whose function in T cells is undocumented.

Given the profound effect of *eIF5a* KO on puromycin incorporation, we investigated components of the cellular translation machinery in the nascent proteome. Amongst the 73 detected ribosomal proteins, 11 were up-regulated in *eIF5a* KO cells (FC>1.5, P<0.05, paired T test, n=4) while none was down-regulated. Up-regulation of specific ribosomal proteins might suggest that ribosomes with alternative composition were being produced in cells in the absence of eIF5a and these ribosomes may preferentially translate a different subset of mRNAs compared to those in WT cells (Segev and Gerst, 2018; Shi et al., 2017). Additionally, down-regulation of translation factors eIF1a, eIF3 components and eIF5 suggest the formation and function of the 43S pre-initiation complex may be reduced (Jackson et al., 2010), while an increase in eIF6 would prevent the 60S ribosome subunit from joining with the 40S subunit (Ceci et al., 2003). Together these data illustrate a general reduction in protein production, and an altered translation profile in *eIF5a* KO cells. All the aforementioned proteins were down-regulated upon GC7 treatment, illustrating the systematically inhibited nature of protein synthesis in these samples. Polyamines are known to modulate both the efficiency and fidelity of protein synthesis, by direct interaction with the ribosome (Dever and Ivanov, 2018). It is possible that the eIF5a-independent effects of GC7

may result from its competitive binding to ribosomes and thus displacing other polyamines present in the cell.

To further separate direct effects of eIF5a on translation from indirect effects following changes in transcription factor production, next generation transcriptomic sequencing of GC7- and Mock-treated samples was performed (Table S2). Although GC7-treatment has off target effects independent of eIF5a, the quality of RNA extracted from these fresh cells was suitable for transcriptomic analyses, whereas RNA extracted from eIF5a KO cells, which needed to be formaldehyde-fixed for sorting, was not.

Translationally down-regulated genes in the GC7 treated samples were defined as the 1635 genes with no significant reduction in RNA abundance (FDR>0.1, edgeR, n=4) but decreased in their nascent proteins (Fig 4A left Venn diagram and Table S3, FDR<0.05, FC<-1.5, unpaired T test with Benjamini-Hochberg correction, n=4). These genes were overlaid with those reduced in their nascent proteins in eIF5a KO compared to WT cells (Fig 4A, right Venn diagram and Table S3, P<0.05, FC<-1.5, paired T test, n=4) which gave 119 genes whose translation we could confidently assign as being dependent on eIF5a. Proteins translationally down-regulated only by GC7 treatment were enriched in GO terms "structural constituent of ribosome", "mRNA binding", "rRNA binding", and "translation factor activity" (FDR < 0.01); while the 119 proteins translationally reduced by both GC7 treatment and eIF5a KO did not significantly cluster into any GO terms, although some key genes regulating cellular functions such as mTORC1 modulators SLC3A2 and SLC7A5, cell cycle regulators Cyclin B1, CDK1 and RFC3/5, MHC Class I genes H2-D1, H2-K1 and H2-Q4, transcription factors FOXP1, RELA, IRF4 and TBET, and translation factors eIF3F/L/M were in this Venn group (Table S3).

Given the well studied role of eIF5a in translation through polyproline and charged amino acid regions of proteins, the frequencies of previously characterised eIF5a-controlled 3-mer motifs (PPP, PPG, DVG, DDG, GGT, and RDK (Schuller et al., 2017)) were counted in the coding sequence (CDS) of the eIF5a-regulated gene set (Fig 4C). Compared to the frequencies of 50 randomly selected 3-mer sequences, PPP is enriched (P<0.01, one-sided Fisher's exact test). PPG and DVG were possibly enriched but ejected by the statistic test (P>0.99). To assess the polyproline content

of these genes, they were overlaid with a published list of genes in the mouse genome with PPP or PPG motifs in their coding region (Mandal et al., 2014). We plotted the percentage of genes containing polyproline in 2900 randomly selected genes in the transcriptome (repeated 1000 times), all expressed proteins detected by LC-MS, and translationally down-regulated genes in GC7-treated or *eIF5a* KO cells in 4 biological samples shown in Fig 4D. In GC7-treated cells, the percentage of polyproline genes was similar between all expressed and translationally down-regulated proteins, suggesting no enrichment for polyproline while the translationally down-regulated proteins in *eIF5a* KO cells had a noticeable increase in polyproline content (Fig 4D), consistent with a more targeted inhibition of eIF5a in the KO versus GC7-treated samples.

We then focused our analyses and validation on proteins involved in regulating cytokine production and cell cycle progression whose translation we had identified as regulated by the availability of mature eIF5a (listed in Fig 4E-F), since defects of these cellular functions were observed in *eIF5a* KO cells. Moreover, these cellular functions are the major age-related impairments described in CD8 T cells (Po et al., 2002), and so might illustrate a possible role eIF5a plays in immunosenescence. Of these genes, we were able to obtain suitable antibodies against IRF4, TBET and CDK1 for flow cytometry which when stained together with antibody against eIF5a or hypusine allowed visualisation of these proteins in the KO and WT populations within the same culture (Fig 5A). Both IRF4 and TBET are transcription factors regulating production of IFNγ and TNFα in T cells (Raczkowski et al., 2013; Szabo et al., 2002), and CDK1 interacts with different cyclins to facilitate cell cycle progression (Santamaría et al., 2007). In agreement with the systematic datasets, the abundance of IRF4 dropped in *eIF5a* KO, *Dhps* KO and to a lesser extent in *Dohh* KO cells, with no reduction in its RNA levels in these cells (Fig 5B). Similarly, TBET and CDK1 proteins decreased in *eIF5a* KO and *Dhps* KO, but not in *Dohh* KO cells, in correlation with their cell cycle profiles in Fig 2A, and again their RNA abundances were unchanged.

Spermidine availability affects some but not all eIF5a-regulated proteins

The polyamine spermidine is produced naturally in cells and its abundance decreases with age. Exogenous supplementation of spermidine has been shown to increase

organismal longevity (Eisenberg et al., 2009) and restore age-related T cell functions in an autophagy-dependent manner (Puleston et al., 2014). In addition to being required for hypusination of eIF5a, spermidine is also known for inhibiting the acetyltransferase activity of Ep300 and thereby inducing autophagy (Pietrocola et al., 2015). This non-eIF5a-dependent activity of spermidine may have contributed to discrepancies between cellular phenotypes of GC7 (a spermidine derivative) treated cells and *eIF5a/Dhps* KO cells. To further understand how spermidine affects T cell function via eIF5a, activated T cells were treated with an ornithine decarboxylase inhibitor DFMO for 48h to halt spermidine biosynthesis. Additionally DMFO-treated cells were supplemented with exogenous spermidine or were treated with GC7.

Quantitation of cellular hypusine levels reflected the predicted outcome of these treatments. The DFMO-treated cells had decreased hypusine, which was restored to a level comparable to untreated cells by spermidine supplementation. Depletion of hypusination by DFMO was exacerbated by additional GC7 treatment (Fig 6A). However, when we examined the abundance of total and selected proteins whose production we identified as being sensitive to loss of eIF5a by flow cytometry, an altered pattern emerged (Fig. 6B). Puromycin incorporation and CDK1, IRF4, TBET, and IFNγ abundance decreased following DFMO treatment, and were restored by exogenous spermidine addition, indicating they are very sensitive to cellular spermidine levels. Addition of GC7 further decreased Puromycin incorporation, CDK1 and IFNγ expression but did not impact significantly on the abundance of TBET and IRF4 protein. Interestingly, although IFNγ production was affected by both DFMO and DFMO+GC7, the protein was still detectable in all cells which was in contrast to *eIF5a/Dhps* KO cells where a significant proportion of the population produced no detectable IFNγ (Fig 2B). In contrast, TNFα, CD25 and CDC45 were only decreased in DFMO+GC7-treated cells and not following treatment with DFMO alone, indicating that they are less sensitive to spermidine depletion, despite showing some regulation by eIF5a (Fig 2B and Table S1). It is likely that that GC7 treatment incompletely blocked eIF5a hypusination and, therefore, function and so had a milder effect on the down-regulation of IRF4 and TBET compared to *eIF5a* or *Dhps* KO (Fig S2). Together these data indicate that the maintenance of some cellular proteins such as CDK1, TBET and IRF4 are very sensitive to the availability of functional hypusinated eIF5a. On the other hand, other proteins such as the cytokine IFNγ, requires total ablation of

eIF5a or DHPS in order to show a profound reduction in abundance. We could further demonstrate eIF5a-independent effect of GC7 on CD25 (IL-2ra), a crucial receptor for T cell expansion and survival, which was only mildly reduced in *eIF5a* KO or following DFMO treatment, but significantly down-regulated by DMFO+GC7 treatment (Fig 6B and S2).

DISCUSSION

Our current understanding of eIF5a function largely comes from studies in yeast and transformed cell lines which proliferate continuously and therefore constitutively express active hypusinated eIF5a. In comparison, little has been reported on how eIF5a is regulated in primary mammalian cells. It was interesting therefore, that in primary T cells that transited between a naïve and an activated phenotype, the major regulation of eIF5a activity occurred post-translationally. We demonstrated that naïve murine T cells have abundant expression of eIF5a protein, but very little contains the hypusine modification. Upon T cell activation, eIF5a becomes hypusinated in a dynamic fashion by the transcriptional upregulation of DHPS and DOHH, the enzymes dedicated to the transfer the 4-aminobutyl moiety from spermidine onto Lys^{50} of eIF5a. Similarly, human peripheral blood lymphocyte were shown to increase eIF5a hypusination upon activation with mitogens (Cooper et al., 1982). Using CRISPR knockout of these enzymes or eIF5a itself, we confirmed that in T cells hypusination was important for eIF5a function, however the phenotype of the eIF5a knockout was more extreme than that of DHPS or DOHH knockouts. These studies suggest that either unmodified eIF5a has some residual activity or that a fraction of hypusinated eIF5a protein is very long-lived and that very small lingering quantities can still promote translation of some proteins. We also found that the deoxyhypusinated intermediate form of eIF5a had clearly detectable, albeit reduced functionality in T cells, as knockout of DOHH, which catalyses the last irreversible step in the conversion of lysine to hypusine, showed much less impact that the removal of DHPS. Two key pathways in T cells which we could verify were primarily controlled at the level of translation by functional eIF5A, were control of cell cycle and cytokine production, both essential for T cell differentiation and function.

Controlling protein production by regulating polypeptide elongation is emerging as an important means of translational control beyond that exerted by the more extensively studied regulators of translation initiation (Richter and Coller, 2015). Dysregulation of elongation has also been shown to contribute to diseases such as neurodegeneration and cancer (Knight et al., 2020; Richter and Coller, 2015). The availability of aminoacyl-transfer RNAs and the action of key proteins such as eEF1A, eEF2 and eEF2K have been well documented in controlling the rate of polypeptide formation

(Choi et al., 2018). In contrast the role of eIF5A is less well understood but it is thought to be particularly important for elongation through certain motifs in the mRNAs particularly those encoding proline-proline and other charged residues (Schuller et al., 2017). We confirmed that eIF5a regulates de novo translation of subsets of cellular proteins in T cells which are enriched in PPP motifs, as previously described (Gutierrez et al., 2013), although more than 50% of our identified eIF5a-regulated genes did not have these motifs in their coding sequences. Strong ribosome pausing has been observed in eIF5a-depleted yeast at non-proline, charged amino acid containing motifs such as DVG, DDG, GGT, and RDK (Schuller et al., 2017), although of these motifs, only DVG seemed slightly, but not significantly, enriched in primary T cells. Online motif discovery tools such as MEME Suite (Bailey et al., 2009) were not able to identify enriched motifs in the CDS of eIF5a-regulated genes compared to all expressed proteins. The mechanism for down-regulation of non-polyproline containing eIF5a-regulated proteins remains unresolved. One possibility is that those mRNAs are disadvantaged by changes in ribosome composition (Ferretti et al., 2017; Shi et al., 2017) which were suggested by alterations in nascent polypeptide abundance of ribosome associated proteins in *eIF5a* KO cells and which might further influence prioritisation of mRNA translation. In this case some of the more global down-regulation of translation may not be linked to the elongation activity of eIF5a directly but rather indirectly through its influence on ribosomal protein translation. Interestingly in this context is a recent description of a ribosomopathy that presents as immune deficiency and impaired T cell activation (Robertson et al., 2021), providing further evidence that successful T cell activation may be particularly sensitive to ribosome composition.

An important consequence of T cell activation is their differentiation to effector cells whose function requires cytokine secretion in a temporally and spatially regulated manner. It has been appreciated for some time that cytokine production in immune cells is controlled both transcriptionally and translationally, with the latter involving both miRNAs (Asirvatham et al., 2009) and RNA binding proteins (Anderson, 2010; Salerno et al., 2020). Production of some effector cytokines, including IFNγ, was found to be particularly sensitive to the availability of functional eIF5a. Our data indicate that eIF5a provides an additional layer of cytokine regulation able to influence both translation of cytokines directly and indirectly through translation of their transcription factors such

as TBET and IRF4. In this context there is an interesting correlation between the decline of cytokine production, T cell proliferation (Sievert et al., 2014) and memory T cell formation (Puleston et al., 2014) observed in ageing and an age-related decline in eIF5a protein abundance in lymphocytes (Zhang et al., 2019a). Not only is eIF5a protein reduced but spermidine availability also declines with age (Eisenberg et al., 2009) and dietary spermidine supplementation in old mice has been shown to promote and improve T memory cell formation (Puleston et al., 2014) and B cell antibody production (Zhang et al 2019). Although indirect, these publications support an active role for eIF5a in maintaining T cell functionality in ageing.

Inhibition of eIF5a hypusination with GC7 is in consideration as a treatment for several cancers due to its anti-proliferative activity (Coni et al., 2020) and ability to induce cell death (Schultz et al., 2018), so it was interesting to evaluate whether GC7 treatment mirrored the effects of genetic ablation of DHPS or eIF5a in T cells. GC7 produced a very different cellular phenotype from *Dhps* knockout, as has been shown for a fibrosarcoma cell line (Oliverio et al., 2014). As a structural derivative of polyamine spermidine, GC7 is known to compete with spermidine for binding to DHPS and has been widely used as an inhibitor for functional maturation of eIF5a. As spermidine has functions independent of eIF5a, such as inhibition of Ep300 activity, it is also likely that GC7 has similar effects. CD25 has been previously shown to decrease following GC7 treatment in CD4 T cells (Colvin et al., 2013) as was seen here in CD8 T cells. However knockout of *eIF5a* did not reduce CD25 to the same extent, suggesting GC7 at least partially reduces CD25 expression independently of eIF5a. In our LC-MS dataset, GC7 treatment resulted in 3044 down-regulated nascent proteins, while eIF5a KO only decreased 369 proteins. These observations indicate a substantial inhibitory effect of GC7 that does not act through eIF5a. Proteins down-regulated exclusively after GC7 treatment are not enriched in poly proline motifs, and they cluster significantly to GO terms associated with ribosome constituents, suggesting GC7 reduces cellular translation in a more global manner than eIF5a, of which the knockout resulted in increase, rather than decrease, in several ribosomal proteins.

Our data support the importance of eIF5a in regulating immune cell function. Targeting eIF5a hypusination has been shown to have anti-inflammatory effects in macrophages (Oedem Paulo de et al., 2014) and to regulate macrophage mitochondrial respiration

and activation (Puleston et al., 2019). Our current study supports a role for eIF5a in influencing cytokine production and proliferation in T cells also both through directly regulating translation of subsets of mRNAs and indirectly by influencing the production of key transcription factors. Together these studies lend weight to the existence of eIF5a directed RNA regulons important for modulating immune responses.

Acknowledgements:

This research was funded by the Wellcome Trust Investigator Award WT205014/Z/16/Z to RZ and byt the Wellcome Trust/Royal Society Sir Henry Dale Fellowship 20611/Z/17/Z to TL. For the purpose of open access, the author has applied a CC BY public copyright license to any author accepted manuscript version arising from this submission.

We would like to thank the following at the University of Edinburgh: Dr Martin Waterfall IIIR for cell sorting expertise, the BVS animal facility in Ashworth labs and CRM for support with animal husbandry, and the proteomics facility at the Wellcome Centre for Cell Biology for mass spec. We also thank Dr Alehandro Brenes Murillo at the University of Dundee for valuable discussion and help regarding dataset analysis.

Author contributions:

Conceptualization: TT & RZ; Methodology: TT, VK and TL; Investigation: TT, XZ and VK; Formal analysis: TT; Data curation: TT and VK; Writing – original draft: TT and RZ; Writing review and editing: TT, TL and RZ; Funding acquisition: TL and RZ; Resources: RZ; Supervision: RZ.

Declaration of Interests:

The authors declare no competing interests.

Online data deposition:

Original and processed RNASeq dataset have been deposited in the Gene Expression Omnibus (GEO) database, https://www.ncbi.nlm.nih.gov/geo (accession no. GSE168731).

Original files for the nascent proteomic dataset are available at EBI PRIDE database, https://www.ebi.ac.uk/pride/ (accession no. PXD021063).

STAR Methods

LEAD CONTACT AND MATERIALS AVAILABILITY

Requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Rose Zamoyska (Rose.Zamoyska@ed.ac.uk).

EXPERIMENTAL MODEL

Mice used in experiments were of C57BL/6 background strain carrying homozygous OT-1 transgenic T cell receptor on a Rag-1KO background. All mice were bred at the animal facility in the University of Edinburgh in accordance with the UK Home Office and local ethically approved guidelines. Mice of both sexes were used.

METHOD DETAILS

Cell culture and stimulation

Lymph node (LN) OT-I T cells were cultured in IMDM medium (Sigma) supplemented with 10% FCS, L-glutamine, 100U/ml penicillin, 100U/ml streptomycin and 50µM 2β-ME. 100nM SIINFEKL (N4) peptide (Cambridge Peptides) were added to culture media and the cells were incubated at 37°C for the duration specified in each experiment.

For GC7 treatment for transcriptomic and nascent proteomic profiling, 2-day cultured activated cells were washed and then cultured for 2 days in culture media supplemented with 20ng/mL recombinant human IL-2 (PeproTech) in the presence or absence of 10µM GC7 (Merck Chemicals). For DFMO experiment, activated cells expanded for 2 days in IL-2 were cultured for additional 2 days in fresh IL-2, or IL-2 supplemented with 0.5mM DFMO (Bio-Techne), DFMO+10µM spermidine (Sigma-Aldrich), or DFMO+10µM GC7.

CRISPR-Cas9 transfection

The following guide RNA sequences were purchased from Integrated DNA Technologies: *eIF5a* (TGCTCAGCATTACGTAAGAA and CTTCGAGACAGGAGATGCAG), *Dhps* (ATACCTCGTGCAGCACAACA), *Dohh* (GCAGTATTCTACGGACCCAG), *Thy1* (ACAGACAAGCTGGTCAAGTG), and TRAC guide RNA (CAGGGTTCTGGATATCTGT). Cas9 ribonucleoprotein complexes were

formed by mixing 2µL of 100µM tracrRNA and 2µL of 100µM crRNA in 25µL Nuclease-free duplex buffer (Integrated DNA Technologies, USA), incubating at 95°C for 5 min and then mixed with 2µL of 5mg/mL Truecut Cas9 protein (Thermo Fisher Scientific, USA) for 10 min at 37°. The RNP complexes were made immediately before the transfection and mixed with 10^6 2-day activated OT-1 cells. Transfections were performed with the Neon transfection system (Thermo Fisher Scientific, USA) following manufacturer's instructions, with the setting 1600V, 10ms, 3 pulses. Transfected cells were cultured in IL-2-containing media for 3 days for *eIF5a* KO, and 4 days for *Dhps* and *Dohh* KO.

AHA-labelling and cell sorting

Cells were transferred to IL-2-containing RPMI media in the absence of methionine, at a concentration of 5 x 10^6 cells/mL. Cells were starved of methionine for one hour then 4-Azido-L-homoalanine HCl (L-AHA) (Jena Bioscience) was added to the media (final concentration 40µM) and incubated for 2 hours at 37°C. For *eIF5a* CRISPR experiment, cells were fixed in 2% formaldehyde for 20 min and permeabilised with ice-cold 90% methanol for 1 hour. Staining was done using anti-eIF5a-C-terminus antibody (Abcam) and anti-rabbit secondary antibody at AlexaFluor 647 (Thermo Fisher Scientific). Sorting was undertaken with a FACS Aria (BD).

Click chemistry and preparation for LC-MS

Proteins from freshly frozen cell samples (12.5 x 10^6 of non-AHA-labelled, 32.5 x 10^6 of GC7 and 12.5 x 10^6 of Mock, 4 samples each) were extracted with CHAPS Lysis buffer from the Click Chemistry Capture Kit (Jena Bioscience). For sorted fixed cells, proteins from 10 x 10^6 cells were extracted using 310µL of 200mM Tris, 0.5% SDS, 200mM NaCl, pH8.0, with brief sonications to disturb the cell pellet, followed by reverse crosslinking at 95°C for 45 min and cooled on ice, and then mixed with equal volume of 16M Urea giving a final concentration of 8M Urea. Extracted proteins were incubated overnight with Alkyne agarose beads (Jena Bioscience), and downstream preparations were done using Click Chemistry Capture Kit following the manufacturer's guidelines. Tryptic peptides derived from pooled GC7 and Mock cells were used to generate a fractionated library using high pH reverse phase chromatography with concatenation into 12 fractions. All samples were analysed by LCMS with a 120min gradient over a 75um x 50cm easyspray C18 column. LC was coupled to an Orbitrap Elite™ Hybrid Ion Trap-Orbitrap Mass Spectrometer (Thermo

Scientific) where data was acquired with 120k resolution MS1 followed by top 20 CID MS2 in the ion trap.

Systematic Dataset handling

Pair-end RNA sequencing dataset reads were generated by Novogene (GC7- and Mock-treated samples, 4 biological replicates) and were trimmed of low-quality and adapter sequences. Reads were aligned to the mouse genome (GRCm38.96) using STAR (Dobin et al., 2013) and mapped reads were counted using featureCounts (Liao et al., 2013). Differential expression analysis was performed with edgeR (Robinson et al., 2010).

Nascent proteomic data was analysed using MaxQuant version 1.6.2.6. Match-between-runs was enabled to transfer identifications from the library and between samples. iBAQ values were used for analysis. The values were normalised using average sum within sample groups, and then by the input cell numbers. Statistical tests were performed using Log2-transformed normalised iBAQ values. Unpaired T test with Benjamini-Hochberg correction (Benjamini and Hochberg, 1995) was used to determine significantly down-regulated proteins between GC7- and Mock-treated samples. Paired T test was used for eIF5a KO versus WT samples.

Flow cytometry

Cells were labelled with LIVE/DEAD™ Aqua Dead Cell Stain Kit (Invitrogen) and then fixed in Intracellular Staining Fixation Buffer (BioLegend) and permeabilised in Intracellular Staining Permeabilization Wash Buffer (BioLegend). Antibodies against eIF5a (EP527Y, Abcam), hypusine (Hpu24, Creative Biolabs), IFNg (XMG1.2, BioLegend), TNFa (MP6-XT22, eBioscience), IRF4 (3E4, eBioscience), TBET (4B10, BioLegend), CDK1 (A17, Abcam), CD25 (PC61, BioLegend), CDC45 (EPR5759, Abcam) and puromycin (12D10, Millipore) were used; for cell cycle analysis, fixed-permeabilised cells were stained with Hoechst 33342 (Sigma) and Ki-67 antibody (BD Biosciences). Cells were incubated with primary antibodies for 16 hours at 4°C, washed, followed by staining with Goat anti-rabbit Alexa Fluor 647 secondary antibody (Thermo Fisher Scientific) for 1 hour at RT. All analyses were done on singlet live events.

All acquisitions were done on a MACSQuant flow cytometer (Miltenyi Biotec).

Protein extraction and Western blotting

Cells were lysed in RIPA Buffer (Thermo Fisher Scientific) supplemented with a protease inhibitors cocktail (Sigma). The lysates were centrifuged to remove the debris. Reducing Laemmli buffer was added to the lysates, and samples were heated to 95°C and separated by SDS-PAGE. Proteins were transferred to Immobilon-FL PVDF membrane (Millipore), and membranes were incubated in Odyssey blocking buffer (LI-COR Biosciences) prior to incubation with the primary antibodies. Quantitative signals were generated with secondary Abs: IRDye® 680RD Goat anti-Rabbit and IRDye® 800CW Goat anti-Mouse (LI-COR Biosciences), and visualised using the Odyssey Infrared Imaging System (LI-COR Biosciences).

The following primary Abs were used: anti-ZAP70 (BD Biosciences), DHPS (ab224134, Abcam), eIF5a (EP527Y, Abcam), and hypusine (Hpu24, Creative Biolabs).

Transcript analyses by RT-qPCR

Total cellular RNA was extracted using Direct-zol RNA Miniprep kit (Zymo Research) and reverse transcribed using Lunascript kit (New England Biolabs). qPCR was performed on a Light Cycler 480 (Roche) using Brilliant III SYBR® Master Mixes (Agilent). Transcription level of each gene was normalised against the quantity of beta actin (Actb) in the sample. For fixed sorted cells, total cellular RNA was isolated using RecoverAll™ Total Nucleic Acid Isolation Kit (Thermo Fisher Scientific) and analysed as above.

QUANTIFICATION AND STATISTICAL ANALYSIS

Prism software (GraphPad) was used for statistical analyses. All data points represent the measurement of biological replicates. Unpaired or paired two-tailed Student's t test was used for comparisons between two normally distributed datasets with equal variances unless specified. One-tailed Fisher's exact test was used for testing enrichment of amino acid trimers. In the GC7-Mock nascent proteome dataset, Benjamini–Hochberg method was used to adjust P values of a family of multiple t tests with a single null hypothesis. P value was used to quantify the statistical significance of the null hypothesis testing. $*p \leq 0.05$, $**p \leq 0.01$.

DATA AND CODE AVAILABILITY

Original data and code are available at https://doi.org/10.17632/vhyhnn96hf.1

Supplementary Tables as Excel files

Table S1:

Normalised iBAQ values of nascent proteomes, differential expression analysis, and selection of proteins for presentation

Legend: iBAQ values from LC-MS for GC7-treated, Mock-treated, eIF5a-KO and eIF5a-WT cells were normalised using average sum within sample groups, and then by the input cell numbers. Statistical tests were performed using Log2-transformed normalised iBAQ values. Unpaired T test with Benjamini-Hochberg correction was used to determine significantly down-regulated proteins between GC7- and Mock-treated samples. Paired T test was used for eIF5a KO versus WT samples.

Table S2:

Differential expression analysis of the RNASeq dataset between GC7- and Mock-treated cells, with top ten enriched KEGG pathways in significantly up- and down-regulated genes.

Legend: The dataset was trimmed of low-quality and adapter sequences. Reads were aligned to the mouse genome (GRCm38.96) using STAR and mapped reads were counted using featureCounts. Differential expression analysis was performed with edgeR. Pathway enrichment analysis was conducted with WebGestalt.

Table S3:

Determination of genes translationally regulated by eIF5a using Venn analyses.

Legend: Genes translationally down-regulated upon GC7-treatment was defined as those not down-regulated in RNA levels but decreased in nascent peptide levels following the treatment. This list of genes was overlaid with nascent proteins down-regulated in eIF5a-KO cells and the common genes between them were defined as genes translationally regulated by eIF5a.

References

Anderson, P. (2010). Post-transcriptional regulons coordinate the initiation and resolution of inflammation. Nature reviews. Immunology *10*, 24-35.

Asirvatham, A.J., Magner, W.J., and Tomasi, T.B. (2009). miRNA regulation of cytokine genes. Cytokine *45*, 58-69.

Babicki, S., Arndt, D., Marcu, A., Liang, Y., Grant, J.R., Maciejewski, A., and Wishart, D.S. (2016). Heatmapper: web-enabled heat mapping for all. Nucleic acids research *44*, W147-153.

Bailey, T.L., Boden, M., Buske, F.A., Frith, M., Grant, C.E., Clementi, L., Ren, J., Li, W.W., and Noble, W.S. (2009). MEME Suite: tools for motif discovery and searching. Nucleic Acids Res *37*, W202-W208.

Benjamini, Y., and Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society. Series B (Methodological) *57*, 289-300.

Ceci, M., Gaviraghi, C., Gorrini, C., Sala, L.A., Offenhäuser, N., Carlo Marchisio, P., and Biffo, S. (2003). Release of eIF6 (p27BBP) from the 60S subunit allows 80S ribosome assembly. Nature *426*, 579-584.

Choi, J., Grosely, R., Prabhakar, A., Lapointe, C.P., Wang, J., and Puglisi, J.D. (2018). How Messenger RNA and Nascent Chain Sequences Regulate Translation Elongation. Annual Review of Biochemistry *87*, 421-449.

Ciafrè, S.A., and Galardi, S. (2013). microRNAs and RNA-binding proteins: a complex network of interactions and reciprocal regulations in cancer. RNA Biol *10*, 935-942.

Colvin, S.C., Maier, B., Morris, D.L., Tersey, S.A., and Mirmira, R.G. (2013). Deoxyhypusine Synthase Promotes Differentiation and Proliferation of T Helper Type 1 (Th1) Cells in Autoimmune Diabetes. Journal of Biological Chemistry *288*, 36226-36235.

Coni, S., Serrao, S.M., Yurtsever, Z.N., Di Magno, L., Bordone, R., Bertani, C., Licursi, V., Ianniello, Z., Infante, P., Moretti, M.*, et al.* (2020). Blockade of EIF5A hypusination limits colorectal cancer growth by inhibiting MYC elongation. Cell Death & Disease *11*, 1045.

Cooper, H.L., Park, M.H., and Folk, J.E. (1982). Posttranslational formation of hypusine in a single major protein occurs generally in growing cells and is associated with activation of lymphocyte growth. Cell *29*, 791-797.

Dever, T.E., Dinman, J.D., and Green, R. (2018). Translation Elongation and Recoding in Eukaryotes. Cold Spring Harbor perspectives in biology *10*, a032649.

Dever, T.E., and Ivanov, I.P. (2018). Roles of polyamines in translation. The Journal of biological chemistry *293*, 18719-18729.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15-21.

Eisenberg, T., Knauer, H., Schauer, A., Büttner, S., Ruckenstuhl, C., Carmona-Gutierrez, D., Ring, J., Schroeder, S., Magnes, C., Antonacci, L.*, et al.* (2009). Induction of autophagy by spermidine promotes longevity. Nature Cell Biology *11*, 1305.

Ferretti, M.B., Ghalei, H., Ward, E.A., Potts, E.L., and Karbstein, K. (2017). Rps26 directs mRNA-specific translation by recognition of Kozak sequence elements. Nature Structural & Molecular Biology *24*, 700-707.

Gabut, M., Bourdelais, F., and Durand, S. (2020). Ribosome and Translational Control in Stem Cells. Cells *9*, 497.

Gutierrez, E., Shin, B.-S., Woolstenhulme, Christopher J., Kim, J.-R., Saini, P., Buskirk, Allen R., and Dever, Thomas E. (2013). eIF5A Promotes Translation of Polyproline Motifs. Molecular Cell *51*, 35-45.

Hanauske-Abel, H.M., Park, M.H., Hanauske, A.R., Popowicz, A.M., Lalande, M., and Folk, J.E. (1994). Inhibition of the G1-S transition of the cell cycle by inhibitors of deoxyhypusine hydroxylation. Biochimica et Biophysica Acta (BBA) - Molecular Cell Research *1221*, 115-124.

Hukelmann, J.L., Anderson, K.E., Sinclair, L.V., Grzes, K.M., Murillo, A.B., Hawkins, P.T., Stephens, L.R., Lamond, A.I., and Cantrell, D.A. (2015). The cytotoxic T cell proteome and its shaping by the kinase mTOR. Nat Immunol *17*, 104-112.

Jackson, R.J., Hellen, C.U.T., and Pestova, T.V. (2010). The mechanism of eukaryotic translation initiation and principles of its regulation. Nature Reviews Molecular Cell Biology *11*, 113-127.

Kang, H.A., and Hershey, J.W. (1994). Effect of initiation factor eIF-5A depletion on protein synthesis and proliferation of Saccharomyces cerevisiae. Journal of Biological Chemistry *269*, 3934-3940.

Knight, J.R.P., Garland, G., Pöyry, T., Mead, E., Vlahov, N., Sfakianos, A., Grosso, S., De-Lima-Hedayioglu, F., Mallucci, G.R., von der Haar, T.*, et al.* (2020). Control of translation elongation in health and disease. Dis Model Mech *13* (3):dmm043208

Lawrence, C.W., and Braciale, T.J. (2004). Activation, differentiation, and migration of naive virus-specific CD8+ T cells during pulmonary influenza virus infection. Journal of immunology *173*, 1209-1218.

Lee, S.B., Park, J.H., Kaevel, J., Sramkova, M., Weigert, R., and Park, M.H. (2009). The effect of hypusine modification on the intracellular localization of eIF5A. Biochemical and biophysical research communications *383*, 497-502.

Liao, Y., Smyth, G.K., and Shi, W. (2013). featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics *30*, 923-930.

Liao, Y., Wang, J., Jaehnig, E.J., Shi, Z., and Zhang, B. (2019). WebGestalt 2019: gene set analysis toolkit with revamped UIs and APIs. Nucleic acids research *47*, W199-W205.

Mandal, A., Mandal, S., and Park, M.H. (2014). Genome-Wide Analyses and Functional Classification of Proline Repeat-Rich Proteins: Potential Role of eIF5A in Eukaryotic Evolution. PLOS ONE *9*, e111800.

Mathews, M.B., and Hershey, J.W.B. (2015). The translation factor eIF5A and human cancer. Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms *1849*, 836-844.

Munoz, J., Low, T.Y., Kok, Y.J., Chin, A., Frese, C.K., Ding, V., Choo, A., and Heck, A.J. (2011). The quantitative proteomes of human-induced pluripotent stem cells and embryonic stem cells. Molecular systems biology *7*, 550.

Oedem Paulo de, A., Jr., Thais Regina, T., Danuza, R., Daniella de Barros, R., Tatiana Faria, W., Fabio Carrilho, G., Alexandra Ivo, M., Cleslei Fernando, Z., and Sandro Roberto, V. (2014). Hypusine Modification of the Ribosome-binding Protein eIF5A, a Target for New Anti-Inflammatory Drugs: Understanding the Action of the Inhibitor GC7 on a Murine Macrophage Cell Line. Current Pharmaceutical Design *20*, 284-292.

Oliverio, S., Corazzari, M., Sestito, C., Piredda, L., Ippolito, G., and Piacentini, M. (2014). The spermidine analogue GC7 (N1-guanyl-1,7-diamineoheptane) induces autophagy through a mechanism not involving the hypusination of eIF5A. Amino Acids *46*, 2767-2776.

Park, J.H., Aravind, L., Wolff, E.C., Kaevel, J., Kim, Y.S., and Park, M.H. (2006). Molecular cloning, expression, and structural prediction of deoxyhypusine hydroxylase: a HEAT-repeat-containing metalloenzyme. Proc Natl Acad Sci U S A *103*, 51-56.

Park, M.H., and Wolff, E.C. (2018). Hypusine, a polyamine-derived amino acid critical for eukaryotic translation. Journal of Biological Chemistry *293*, 18710-18718.

Pelechano, V., and Alepuz, P. (2017). eIF5A facilitates translation termination globally and promotes the elongation of many non polyproline-specific tripeptide sequences. Nucleic Acids Research 45, 7326-7338.

Pietrocola, F., Lachkar, S., Enot, D.P., Niso-Santano, M., Bravo-San Pedro, J.M., Sica, V., Izzo, V., Maiuri, M.C., Madeo, F., Mariño, G.*, et al.* (2015). Spermidine induces autophagy by inhibiting the acetyltransferase EP300. Cell Death Differ *22*, 509-516.

Po, J.L.Z., Gardner, E.M., Anaraki, F., Katsikis, P.D., and Murasko, D.M. (2002). Age-associated decrease in virus-specific CD8+ T lymphocytes during primary influenza infection. Mechanisms of Ageing and Development *123*, 1167-1181.

Puleston, D.J., Buck, M.D., Klein Geltink, R.I., Kyle, R.L., Caputa, G., O'Sullivan, D., Cameron, A.M., Castoldi, A., Musa, Y., Kabat, A.M.*, et al.* (2019). Polyamines and eIF5A Hypusination Modulate Mitochondrial Respiration and Macrophage Activation. Cell Metabolism *30*, 352-363.e358.

Puleston, D.J., Zhang, H., Powell, T.J., Lipina, E., Sims, S., Panse, I., Watson, A.S., Cerundolo, V., Townsend, A.R.M., Klenerman, P.*, et al.* (2014). Autophagy is a critical regulator of memory CD8+ T cell formation. eLife *3*, e03706.

Raczkowski, F., Ritter, J., Heesch, K., Schumacher, V., Guralnik, A., Höcker, L., Raifer, H., Klein, M., Bopp, T., Harb, H.*, et al.* (2013). The transcription factor Interferon Regulatory Factor 4 is required for the generation of protective effector CD8+ T cells. Proceedings of the National Academy of Sciences *110*, 15019-15024.

Richter, Joel D., and Coller, J. (2015). Pausing on Polyribosomes: Make Way for Elongation in Translational Control. Cell *163*, 292-300.

Robertson, N., Shchepachev, V., Wright, D., Turowski, T.W., Spanos, C., Helwak, A., Zamoyska, R., and Tollervey, D. (2021). A disease-linked lncRNA mutation in RNase MRP inhibits ribosome synthesis. bioRxiv, 2021.2003.2029.437572.

Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics *26*, 139-140.

Salerno, F., Turner, M., and Wolkers, M.C. (2020). Dynamic Post-Transcriptional Events Governing CD8+ T Cell Homeostasis and Effector Function. Trends in Immunology *41*, 240-254.

Santamaría, D., Barrière, C., Cerqueira, A., Hunt, S., Tardy, C., Newton, K., Cáceres, J.F., Dubus, P., Malumbres, M., and Barbacid, M. (2007). Cdk1 is sufficient to drive the mammalian cell cycle. Nature *448*, 811-815.

Schmidt, C., Becker, T., Heuer, A., Braunger, K., Shanmuganathan, V., Pech, M., Berninghausen, O., Wilson, D.N., and Beckmann, R. (2016). Structure of the hypusinylated eukaryotic translation factor eIF-5A bound to the ribosome. Nucleic Acids Research *44*, 1944-1951.

Schuller, A.P., Wu, C.C.-C., Dever, T.E., Buskirk, A.R., and Green, R. (2017). eIF5A Functions Globally in Translation Elongation and Termination. Molecular Cell *66*, 194-205.e195.

Schultz, C.R., Geerts, D., Mooney, M., El-Khawaja, R., Koster, J., and Bachmann, A.S. (2018). Synergistic drug combination GC7/DFMO suppresses hypusine/spermidine-dependent eIF5A activation and induces apoptotic cell death in neuroblastoma. The Biochemical Journal *475*, 531-545.

Schwanhäusser, B., Busse, D., Li, N., Dittmar, G., Schuchhardt, J., Wolf, J., Chen, W., and Selbach, M. (2011). Global quantification of mammalian gene expression control. Nature *473*, 337-342.

Segev, N., and Gerst, J.E. (2018). Specialized ribosomes and specific ribosomal protein paralogs control translation of mitochondrial proteins. J Cell Biol *217*, 117-126.

Shi, Z., Fujii, K., Kovary, K.M., Genuth, N.R., Röst, H.L., Teruel, M.N., and Barna, M. (2017). Heterogeneous Ribosomes Preferentially Translate Distinct Subpools of mRNAs Genome-wide. Mol Cell *67*, 71-83.e77.

Shirai, Y.-T., Mizutani, A., Nishijima, S., Horie, M., Kikuguchi, C., Elisseeva, O., and Yamamoto, T. (2019). CNOT3 targets negative cell cycle regulators in non-small cell lung cancer development. Oncogene *38*, 2580-2594.

Sievert, H., Pällmann, N., Miller, K.K., Hermans-Borgmeyer, I., Venz, S., Sendoel, A., Preukschas, M., Schweizer, M., Boettcher, S., Janiesch, P.C.*, et al.* (2014). A novel mouse model for inhibition of DOHH-mediated hypusine modification reveals a crucial function in embryonic development, proliferation and oncogenic transformation. Dis Model Mech *7*, 963-976.

Sonenberg, N., and Dever, T.E. (2003). Eukaryotic translation initiation factors and regulators. Current Opinion in Structural Biology *13*, 56-63.

Szabo, S.J., Sullivan, B.M., Stemmann, C., Satoskar, A.R., Sleckman, B.P., and Glimcher, L.H. (2002). Distinct effects of T-bet in TH1 lineage commitment and IFN-gamma production in CD4 and CD8 T cells. Science (New York, N.Y.) *295*, 338-342.

Tan, T.C.J., Knight, J., Sbarrato, T., Dudek, K., Willis, A.E., and Zamoyska, R. (2017). Suboptimal T-cell receptor signaling compromises protein translation, ribosome biogenesis, and proliferation of mouse CD8 T cells. Proceedings of the National Academy of Sciences *114*, E6117-E6126.

Wu, L., Wang, Y., Liu, Y., Yu, S., Xie, H., Shi, X., Qin, S., Ma, F., Tan, T.Z., Thiery, J.P.*, et al.* (2014). A central role for TRPS1 in the control of cell cycle and cancer development. Oncotarget *5*, 7677-7690.

Yoon, H., Legge, K.L., Sung, S.S., and Braciale, T.J. (2007). Sequential activation of CD8+ T cells in the draining lymph nodes in response to pulmonary virus infection. Journal of immunology *179*, 391-399.

Zhang, H., Alsaleh, G., Feltham, J., Sun, Y., Napolitano, G., Riffelmacher, T., Charles, P., Frau, L., Hublitz, P., Yu, Z.*, et al.* (2019a). Polyamines Control eIF5A Hypusination, TFEB Translation, and Autophagy to Reverse B Cell Senescence. Molecular Cell *76,* 110-125.e9.

Zhang, X., Wang, W., Zhu, W., Dong, J., Cheng, Y., Yin, Z., and Shen, F. (2019b). Mechanisms and Functions of Long Non-Coding RNAs at Multiple Regulatory Levels. Int J Mol Sci *20*, 5573.

Graphical summary

Fig 1. Production and hypusination of eIF5a is stimulated by T cell activation. Naïve OT-1 CD8 T cells were harvested *ex vivo* and stimulated with agonist peptide SIINFEKL for the times indicated. (A) Total cell lysates were Western blotted with Abs for DHPS, hypusine, eIF5a and ZAP70 as loading control, equal cell numbers were loaded in each lane. (B) Fold up-regulation was calculated relative to the 0h time point. Representation of three independent experiments is shown.

Fig 2. CRISPR-Cas9 and pharmacological targeting of eIF5a and its maturation pathway in activated CD8 T cells impaired proliferation, protein synthesis and cytokine productions in a selective manner. (A) Two-day stimulated OT-1 T cells were either treated with 20μM GC7, or targeted to knockout eIF5a, DHPS or DOHH and analysed for expression of eIF5a or Hypusine after two days (GC7), three days (eIF5a) and four days (DHPS and DOHH) of culture in IL-2 to determine the knockout population (KO, red) and the WT (blue). Cell cycle profiles of the models were obtained by Hoechst 33342 staining. Percentage of cells in the accumulated cell cycle phase (S phase in GC7-treated cells, and G1 phase in others), and number of viable cells recovered from the experiments, were plotted at the right (n ≥ 4, *. P < 0.05, **. P < 0.01, paired T test). Thy1: cells targeted to knockout Thy1 which did not affect expression or function of eIF5a. (B) Cells were transfected and cultured as previously described, and then re-stimulated using agonist peptide for 4 hours. Speed of puromycin incorporation, as well as expression of interferon γ (IFNγ) and tumour necrosis factor (TNFα), in KO and WT populations were measured using flow cytometry. Lower panel: Mean fluorescent intensity (MFI) of puromycin and TNFα, and percentage of IFNγ-positive cells from over 6 experiments were plotted (*. P < 0.05, **. P < 0.01, paired T test). (C) RNA levels of IFNγ and TNFα were confirmed with RT-qPCR in fresh bulk transfected cells (Non-restimulated control, Thy1, DHPS, and DOHH, n= 4, *. P < 0.05, unpaired T test) or fixed sorted cells for eIF5a KO (n ≥ 3, *. P < 0.05, paired T test).

Fig 3. Systematic quantitation of GC7- or eIF5a KO-regulated nascent proteomes. (A) GC7- and Mock-treated cells were labelled with AHA for 2 hours prior to RNA and protein extraction from fresh cells. eIF5a CRISPR cells were similarly AHA-labelled, followed by fixation in 2% formaldehyde, permeabilisation in 90% methanol, intracellular staining with anti-eIF5a antibody and secondary AlexaFluor 647 antibody, cell sorting,

and then protein extraction. AHA-labelled proteins were isolated by Click-chemistry with alkyne agarose beads, and then released from the beads using trypsin digestion after washing. In total 5481 proteins (left) were significantly more abundant in these samples compared to background noise (non-AHA-labelled Mock-treated cells). (Right) iBAQ values from each group were normalised by the mean summary of each sample within the group, and then by the input cell number. Log2-transformed values in each sample after normalisation were plotted as violin plots. (B) 3D-MDS plot was used to visualise the difference between samples (left). The percentage of variance represented by each dimension is plotted on the right. (C) Average normalised protein iBAQ values for all detected transcription factors, ribosomal proteins, and translation factors were compared between KO and WT cells, and between GC7- and Mock-treated cells. A selection of proteins is annotated. Black font: significantly changed (P value or FDR < 0.05, Absolute FC > 1.5); Grey font: proteins not deemed significant by the statistical test, but may still be biologically relevant.

Fig 4. Identification of translationally regulated genes affected by GC7 treatment and eIF5a KO. (A) 1635 genes that are translationally down regulated in GC7-treated cells were defined as those not down regulated in their mRNAs (FDR>0.05, edgeR, n = 4) but decreased in their nascent proteins (FDR<0.05, FC<-1.5, unpaired T test with Benjamini-Hochberg correction, n=4). These genes were further overlaid with those reduced in their nascent proteins in eIF5a KO cells (P<0.05, FC<-1.5, paired T test, n = 4) as an extra

criterion of selection for proteins translationally regulated by eIF5a (119 proteins). Lists of genes in these Venn groups are in Table S3. (B) Top ten enriched GO terms in molecular function were listed for genes translationally affected by GC7-treatment or by eIF5a KO. (C) Frequencies of previously studied eIF5a-controlled sequence motifs in the eIF5a-regulated gene set, normalised by the total length of these proteins. Mean and standard deviation of normalised counts of 50 randomly generated 3-mer sequences is plotted at the right (**. P<0.01, one-sided Fisher's exact test). (D) Genes translationally regulated by eIF5a are enriched in poly-proline in their CDS. Percentage of PPP or PPG-containing genes in 2900 randomly selected genes in the transcriptome (repeated 1000 times), all expressed proteins, and translationally down-regulated genes in GC7-treated or eIF5a KO cells in 4 biological samples are plotted. (E) Systematically identified, then hand chosen eIF5a-regulated proteins involved in regulation of cell cycle progression and cytokine production. (F) Normalised iBAQ values for nascent proteins listed in (E) in eIF5a KO and WT cells (*. P<0.05, FC<-1.5, paired T test, n = 4).

Fig 5. Flow cytometry validation of the systematic datasets. (A) Intracellular staining using antibodies against IRF4, TBET, and CDK1 in *eIF5a*, *Dhps*, or *Dohh* CRISPR cells (n ≥ 3, *. P < 0.05, **. P < 0.01, paired T test). (B) RNA levels of these genes were confirmed with RT-qPCR in fresh bulk transfected cells (*Thy1*, *Dhps*, and *Dohh*) or fixed sorted cells for eIF5a KO (n ≥ 3, *. P < 0.05, **. P < 0.01, unpaired T test).

Fig 6. Restricting availability of polyamine spermidine to cells decreased eIF5a hypusination and some but not all eIF5a-regulated genes. (A-B) *Ex vivo* OT-1 cells were stimulated with SIINFEKL peptide for 2 days, expanded in IL-2 culture for another 2 days, and then treated with IL-2 only, 0.5mM DFMO, DFMO + 10μM spermidine, or DFMO + 10μM GC7 in fresh IL-2 culture for 2 days before re-stimulated with the same peptide, puromycin labelling, and then harvested for intracellular staining and flow cytometry. (A) Inhibition

of spermidine biosynthesis by DFMO reduced hypusination. The defect could be rescued by exogenous supplement of spermidine, or further impaired by treatment with GC7. MFIs of hypusine from 5 independent experiments were plotted on the right (**. P < 0.01, unpaired T test). (B) The same samples were stained for cytokines (IFNγ and TNFα), transcription factors (IRF4 and TBET), cell cycle regulators (CDK1 and CDC45), CD25 (IL2 receptor α), and puromycin. MFIs from ≥4 independent experiments were shown in the lower panel. Statistic tests were performed against cells treated with IL-2 only (*. P < 0.05, **. P < 0.01, unpaired T test).

A GC7 relative to Mock

Down-regulated: 3044 proteins	Up-regulated: 31 proteins
Top 10 enriched GO terms in Molecular Function	
mRNA binding	DNA-binding transcription activator activity, RNA polymerase II-specific
translation factor activity, RNA binding	peptidase regulator activity
rRNA binding	protein C-terminus binding
threonine-type peptidase activity	cytokine receptor activity
ubiquitin-like protein ligase binding	protein binding involved in protein folding
ligase activity	MHC protein binding
actin binding	cysteine-type endopeptidase regulator activity involved in apoptotic process
structural constituent of ribosome	cytokine binding
lyase activity	pre-mRNA binding
oxidoreductase activity, acting on CH-OH group of donors	glycosaminoglycan binding

B *eIF5a* KO relative to WT

Down-regulated: 369 proteins	Up-regulated: 149 proteins
Top 10 enriched GO terms in Molecular Function	
actinin binding	structural constituent of ribosome
coenzyme binding	rRNA binding
ligase activity	transferase activity, transferring glycosyl groups
DNA-binding transcription repressor activity, RNA polymerase II-specific	receptor regulator activity
amide binding	translation regulator activity
protein kinase C binding	mRNA binding
SH2 domain binding	oxidoreductase activity, acting on the aldehyde or oxo group of donors
exopeptidase activity	oxidoreductase activity, acting on NAD(P)H
actin binding	isomerase activity
hormone binding	dynein light chain binding

Fig S1. Analyses of nascent proteins up- or down-regulated following (A) GC7-treatment or (B) eIF5a KO. Volcano plots were plotted using the significance criterion described in methods. Heat maps were constructed using the top 2500 significantly changed proteins (Babicki et al., 2016), relative protein

abundance is graded from low (blue), medium (white) to high (red) to allow comparisons between samples. Top 10 enriched GO terms in molecular function were generated using WebGestalt (Liao et al., 2019).

Fig S2. eIF5a KO and GC7 treatment have different impacts on expression of individual proteins (A) MFIs for IRF4 and TBET staining were taken from Fig 3 and 4. To compare across experiments the ratio of treated to control (GC7 vs untreated or KO versus WT cells) were calculated and plotted (n≥4, *p < 0.05, **p < 0.01, unpaired T test). (B) An example of CD25 expression versus eIF5a measured by flow cytometry at Day 3 after CRIPSR-Cas9 transfection is shown in the dot plots and histogram overlays. The fold reduction of MFIs in KO from 6 independent experiments were compared to those of GC7 treated cells in Fig 4 (**p < 0.01, unpaired T test).