Persistence of SARS CoV-2 S1 Protein in CD16+ Monocytes in Post-Acute Sequelae of COVID-19 (PASC) Up to 15 Months Post-Infection

Summary: SARS CoV-2 S1 protein in CD16+ monocytes in the absence of full-length RNA in patients with PASC up to 15 months post-infection

Bruce K. Patterson[1], Edgar B. Francisco[1], Ram Yogendra[2], Emily Long[1], Amruta Pise[1], Hallison Rodrigues[1], Eric Hall[3], Monica Herrara[3], Purvi Parikh[4], Jose Guevara-Coto[5,6], Timothy J. Triche[7], Paul Scott[7], Saboor Hekmati[7], Dennis Maglinte[7], Xaiolan Chang[8], Rodrigo A Mora-Rodríguez[5], Javier Mora[5]

[1]IncellDx Inc, San Carlos, CA
[2]Lawrence General Hospital, Lawrence, MA
[3]Bio-Rad Laboratories, Hercules, CA
[4]NYU Langone Health, New York, NY
[5]Lab of Tumor Chemosensitivity, CIET / DC Lab, Faculty of Microbiology, Universidad de Costa Rica
[6]Department of Computer Science and Informatics (ECCI), Universidad de Costa Rica, San Jose, Costa Rica
[7]Avrok Laboratories, Inc., Azusa, CA
[8] Vaccine & Gene Therapy Institute and Oregon National Primate Research Center, Oregon   Health & Science University, Portland, OR, USA

Summary: SARS CoV-2 S1 Protein in CD16+ Monocytes In PASC

Corresponding author:
Bruce K. Patterson MD
1541 Industrial Road
San Carlos, CA 94070
Tel: +1.650.777.7630
Fax: +1.650.587.1528
Email: brucep@incelldx.com

Key words:

COVID-19, PASC, SARS CoV-2 S1 protein, non-classical monocytes, CCR5, fractalkine

ABSTRACT

The recent COVID-19 pandemic is a treatment challenge in the acute infection stage but the recognition of chronic COVID-19 symptoms termed post-acute sequelae SARS-CoV-2 infection (PASC) may affect up to 30% of all infected individuals. The underlying mechanism and source of this distinct immunologic condition three months or more after initial infection remains elusive. Here, we investigated the presence of SARS-CoV-2 S1 protein in 46 individuals. We analyzed T-cell, B-cell, and monocytic subsets in both severe COVID-19 patients and in patients with post-acute sequelae of COVID-19 (PASC). The levels of both intermediate (CD14+, CD16+) and non-classical monocyte (CD14Lo, CD16+) were significantly elevated in PASC patients up to 15 months post-acute infection compared to healthy controls (P=0.002 and P=0.01, respectively). A statistically significant number of non-classical monocytes contained SARS-CoV-2 S1 protein in both severe (P=0.004) and PASC patients (P=0.02) out to 15 months post-infection. Non-classical monocytes were sorted from PASC patients using flow cytometric sorting and the SARS-CoV-2 S1 protein was confirmed by mass spectrometry. Cells from 4 out of 11 severe COVID-19 patients and 1 out of 26 PASC patients contained ddPCR+ peripheral blood mononuclear cells, however, only fragmented SARS-CoV-2 RNA was found in PASC patients. No full length sequences were identified, and no sequences that could account for the observed S1 protein were identified in any patient. Non-classical monocytes are capable of causing inflammation throughout the body in response to fractalkine/CX3CL1 and RANTES/CCR5.

INTRODUCTION

Post-acute sequelae SARS-CoV-2 infection (PASC) is a disabling and sometimes debilitating

condition that occurs in 10%-30% of individuals infected by SARS-CoV-2 and has recently been

proposed to cause neurologic symptoms in 30% of those infected (*1*). The number and extent of

symptoms is extremely heterogeneous with some reports suggesting >200 different symptoms

(*2*). The underlying cause of PASC symptoms has remained a mystery though some data has

pointed to tissue reservoirs of persistent SARS-CoV-2 as a potential mechanism (*3,4*). We

recently reported a machine learning approach that identified the unique immunologic signature

of individuals with PASC (*5*). In the same report, we also identified characteristic immune cell

subset abnormalities that accompanied the unique cytokine/chemokine profile. The predominant

immune cell abnormality was elevations in monocyte subsets. Monocyte subpopulations are

divided into 3 phenotypic and functionally distinct types. Classical monocytes exhibit the

CD14++, CD16- phenotype, intermediate monocytes exhibit a CD14+, CD16+ phenotype, and

the non-classical monocytes express CD14lo, CD16+ (*6,7*). Further they express very different

cell surface markers as previously described. In particular, classical monocytes express high

levels of the ACE-2 receptor, the putative receptor for SARS-CoV-2 (*8*). Intermediate and non-

classical monocytes express very little ACE-2 receptor. Similarly, classical monocytes express

low levels of the chemokine receptors CX3R1 and CCR5. Intermediate monocytes express high

levels of CCR5 while non-classical monocytes express high levels of CX3R1. Here, we report

kinetic differences in the proportions of monocyte subsets in severe cases and PASC, as well as

the presence of SARS-CoV-2 protein unaccompanied by corresponding viral RNA in CD14lo,

CD16+ monocytes in PASC patients up to 15 months post-acute SARS-CoV-2 infection.

RESULTS

Similar to other inflammatory and infectious conditions such as sepsis, lupus erythematosis, and

rheumatoid arthritis among others (*9*), we detected statistically significant increases (P<0.002) of

intermediate CD14+, CD16+ monocytes in individuals with PASC compared to healthy controls.

In addition, CD14lo, CD16+ non-classical monocytes were also significantly elevated in PASC

(P=0.01). Neither intermediate nor non-classical monocytes were elevated in severe COVID-19

(Figure 1).

Since the reports by our group and others found that monocyte subsets can be infected by HIV,

HCV, Zika virus and Dengue fever virus (*10-12*), we screened peripheral blood mononuclear

cells (PBMCs) from PASC individuals, as well as acute severe COVID-19 as controls, for

SARS-CoV-2 RNA (Table 1). Using the highly sensitive, quantitative digital droplet PCR

(ddPCR), we found that 36% (4 of 11) of severe COVID-19 patients' PBMCs contained SARS-

CoV-2 RNA compared to 4% (1/26) of PASC patients' PBMCs. The one PASC patient that was

RNA positive was 15 months post infection.

To further establish the exact reservoir contributing to the positive signal detected using ddPCR,

we performed high parameter flow cytometry with antibodies that define B cell, T-cell, and

monocytic subsets in addition to simultaneous staining of these cells with an antibody for the

SARS-CoV-2 S1 protein. As demonstrated in Figure 2, we found distinct subpopulations of

SARS-CoV-2 containing cells in the CD14lo, CD16+ monocytic subset for 73% (19 out of 26)

of PASC patients and 91% (10 out of 11) of severe COVID-19 patients. As demonstrated in

Figure 3, the quantity of SARS-CoV-2 S1 containing cells were statistically significant in both

the severe patients (P=0.004) and in the PASC patients (P=0.02). Neither classical monocytes

nor intermediate monocytes expressed the SARS-CoV-2 S1 protein.

To confirm the presence of SARS-CoV-2 S1 protein, we sorted CD14lo, CD16+ monocytes and

performed Ultra High-Performance Liquid Chromatography (UHPLC). Following

immunoprecipitation, the elution fractions were dried down *in vacuo*, resuspended in ddH$_2$O and

purified by to remove any non-crosslinked SARS-CoV-2 S1 antibody as well as any detergents

from the commercial immunoprecipitation buffers. The UHPLC collected fractions were dried *in*

vacuo, resuspended in 100 mM HEPES (pH 8.0, 20% Acetonitrile), and subjected to cistern:

reduction and alkylation with chloroacetamide. The samples were then digested with AspN and

LysC endopeptidases for 16h at 37ºC. The digested peptides were analyzed on an Agilent 6550

IonFunnel QTOF and 1290 UHPLC by comparing patient samples to identical digests performed

on commercially available SARS-CoV-2 S1 subunit. S1 subunit peptides from patient samples

were mapped to a peptide database generated using commercial S1 subunit digests. Peptide

identification consisted of matches in exact mass, isotope distribution, peptide charge state, and

UHPLC retention time. As shown in Figure 4, the retention time of the representative peptide

NLREFVFK in the digested commercial S1 subunit and Sample LH1-6 matched. Additionally,

the Mass Spectra in Figure 4 show identical mass, isotope distribution, and charge states for the

representative peptide NLREFVFK in the representative LH1 sample and commercial S1 subunit

(also observed in LH 2-6, not shown). Using these metrics, up to 44% of the S1 subunit peptides

could be identified in patient samples LH1-LH6 (Supplementary Table 1), providing

complementary evidence to flow cytometry experiments that demonstrate the presence of S1

subunit protein in these patient cells.

To determine whether the observed S1 spike protein was a product of persistent viral infection,

whole viral genome sequencing was performed on monocytes from five patients. Coverage

analysis of the human control amplicons revealed adequate coverage to positively identify

human genomic content. This is consistent with extraction of viral genomic content from a

human host. Human controls also included targeted amplicons for amelogenin (*AMELX* and

AMELY). The ratio of *AMELX* and *AMELY* reads is consistent with the known genders of each

sample.

The sequencing coverage for the five samples was consistent with low viral titer samples or

samples with high Ct values. Average coverage was between 24.17-592.87x and percent bases

covered at 10x and 20x was between 10.81-19.18% and 7.69-15.24% respectively (Table 2).

This is well below the expected threshold to eliminate stretches of Ns > 99 for consensus

sequence submission to GenBank and > 90% genome coverage at 10x for accurate lineage

determination and sequence submission to GISAID (www.gisaid.org). Evaluation of the reads

revealed predominantly short reads (<100bp). To address poor quality reads, primer-dimers or

reads that could possibly map to multiple loci, reads < MAPQ 10 were filtered resulting in the

removal of 3.63-18.99% of total reads per sample.

Lineage determination of the five samples from high quality mutations in the callable regions

yielded lineages of *B* and *B.1* and were non-specific due to inadequate coverage across the

genome. Mutations were identified in *ORF1ab* in all but sample LH5. LH5 had mutations in *N*,

S, and *ORF3b*. (Figure 5).

DISCUSSION

Here, we report the discovery of persistent SARS-CoV-2 protein in CD14lo, CD16+ monocytes

out to 15 months in some individuals and discuss the implications for the pathogenesis of PASC

and severe cases of COVID-19. The three subtypes of circulating monocytes (classical,

intermediate, non-classical) express very different cell surface molecules and serve very different

functions in the immune system. Generally, classical' monocytes exhibit phagocytic activity,

produce higher levels of ROS and secrete proinflammatory molecules such as IL-6, IL-8, CCL2,

CCL3 and CCL5. Intermediate monocytes express the highest levels of CCR5 and are

characterized by their antigen presentation capabilities, as well as the secretion of TNF-α, IL-1β,

IL-6, and CCL3 upon TLR stimulations. Non-classical monocytes expressing high levels of

CX3CR1 are involved in complement and Fc gamma-mediated phagocytosis and anti-viral

responses (*6*).

After maturation, human monocytes are released from bone marrow into the circulation as

classical monocytes. Currently, strong evidence supports the concept that intermediate and non-

classical monocytes emerge sequentially from the pool of classical monocytes (*13*). This is

supported by transcriptome analysis showing that CD16+ monocytes have a more mature

phenotype (*14*). In humans, 85% of the circulating monocyte pool are classical monocytes,

whereas the remaining 15% consist of intermediate and nonclassical monocytes (*13*). Classical

monocytes have a circulating lifespan of approximately one day before they either migrate into

tissues, die, or turn into intermediate and subsequently nonclassical monocytes (*6,13*).

During pathologic conditions mediated by infectious/inflammatory reactions, the proportions of

monocyte subsets vary according to the functionality of each specific subpopulation (*6,13,15*).

Our previous results show that during early stages of the disease, PASC group have reduced

classical monocyte and increased intermediate monocyte percentages compared with healthy

controls (*5*). We find an increase in nonclassical monocytes in PASC group 6-15 months post

infection, and higher percentages of intermediate and nonclassical monocytes at day 0 in severe

cases, suggesting augmented classical-intermediate-nonclassical monocyte transition in both

groups but with different kinetics.

The clinical relevance of monocyte activation in COVID-19 patients and the significance of

these cells as viral protein reservoir in PASC is supported by our data reporting the presence of

S1 protein within nonclassical monocytes. Viral particles and/or viral proteins can enter

monocyte subpopulations in distinct ways, and this appears to be regulated differently in

individuals that will develop severe disease or PASC. Classical monocytes are primarily

phagocytes and express high levels of the ACE-2 receptor (*8*). Therefore, they could either

phagocyte viral particles and apoptotic virally infected cells or be potential targets for SARS-

CoV-2 infection. Considering their short circulating lifespan, viral protein-containing classic

monocytes turn into intermediate and nonclassical monocytes. According to our results, this

process happens faster in the severe group than in the PASC group. Indeed, at early stages of the

disease the severe group show increased nonclassical monocytes whereas in PASC both the

intermediate monocytes and non-classical monocytes are elevated. Additionally, CD14+CD16+

monocytes express intermediate levels of ACE-2 receptors and could as well serve as an

infectious target of SARS-CoV-2 as it has been proved to be an infectious target of HIV-1 and

HCV[11]. Nonclassical monocytes have been proposed to act as custodians of vasculature by

patrolling endothelial cell integrity (*16*), thus pre-existing CD14lo CD16+ cells could ingest

virally infected apoptotic endothelial cells augmenting the proportion of nonclassical monocytes

containing S1 protein. This mechanism is more likely to take place in the PASC group where the

S1 protein was detected 12-15 months post infection than in the severe group. Furthermore,

nonclassical monocytes are associated with FcR-mediated phagocytosis (*17,18*), which might be

related with the ingestion of opsonized viral particles after antibody production at later stages of

the disease in PASC.

Previous reports indicate that the numbers of classical monocytes decrease, but the numbers of

intermediate and non-classical monocytes increase in COVID-19 patients (*19*). Thus, the

presence of S1 protein in nonclassical monocytes in both severe and PASC, might be associated

with clinical characteristics and outcome of these groups. Previously, we found that individuals

with severe COVID-19 have high systemic levels of IL-6, IL-10, VEGF and sCD40L (*5*).

Consistent with our data, other studies showed association of increased production of IL-6,

VEGF and IL-10 by nonclassical monocytes with disease severity (*20-22*).

In the case of PASC, the persistence of circulating S1-containing nonclassical monocytes up to

15 months post infection, independently of the different possible mechanisms of viral proteins

internalization discussed above, indicates that certain conditions are required to maintain this cell

population. It has been shown in both humans and mice that nonclassical monocytes require

fractalkine (CX3CL1) and TNF to inhibit apoptosis and promote cell survival (*22*). Our previous

data show high IFN-γ levels in PASC individuals (*5*), which can induce TNF-α production (*23*).

Further, TNF-α and IFN-γ induce CX3CL1/Fractalkine production by vascular endothelial cells[24]

creating the conditions to promote survival of nonclassical monocytes. Another important aspect

is the permanency of S1-containing cells in the circulation, intermediate monocytes express high

levels of CCR5 and extravasation of these cells can occur in response to CCL4 gradients. We

showed that PASC individuals have low levels of CCL4 (*5*) maintaining these cells in circulation

until they turn into nonclassical monocytes. Moreover, IFN-γ induced CX3CL1/Fractalkine

production by endothelial cells (*23*) creates a gradient within the vascular compartment

preserving nonclassical monocytes expressing CX3CR1 in the circulation.

Nonclassical monocytes are usually referred as anti-inflammatory cells (*22*), nevertheless it was

recently shown that this subset can acquire a proinflammatory phenotype (*25*). Nonclassical

monocytes acquire hallmarks of cellular senescence, which promote long term survival of these

cells in circulation as explained above. Additionally, this induces an inflammatory state of the

non-classical monocytes that could be a manifestation of the senescence-associated secretory

phenotype (SASP), characterized by a high basal NF-κB activity and production of pro-

inflammatory cytokines such as IL-1α, TNF-α and IL-8 (*25*).

The hallmark of PASC is the heterogeneity of symptoms arising in a variety of tissues and

organs. These symptoms are likely associated with the inflammatory phenotype of these

senescent nonclassical monocytes. The CD14lo, CD16+, S1 protein+ monocytes could be

preferentially recruited into anatomic sites expressing fractalkine and contribute to vascular and

tissue injury during pathological conditions in which this monocyte subset is expanded as

previously demonstrated in non-classical monocytes without S1 protein. Previously, CD16+

monocytes were demonstrated to migrate into the brain of AIDS patients expressing high levels

of CX3CL1 (fractalkine) and SDF-1 (*26*), and mediate blood-brain barrier damage and neuronal

injury in HIV-associated dementia via their release of proinflammatory cytokines and neurotoxic

factors. These sequelae are very common in PASC and these data could represent the underlying

mechanism for the symptoms. Interestingly, a number of papers have been written discussing the

increased mobilization of CD14lo, CD16+ monocytes with exercise (*27*). These data support the

reports of worsening PASC symptoms in individuals resuming pre-COVID exercise regimens. In

summary, the mechanism of PASC discussed in this report suggests that intermediate monocytes

remain in circulation due to low CCL4 levels extending their time to differentiate leading to an

accumulation of non-classical monocytes. The utility of using CCR5 antagonists in preventing

migration of intermediate and non-classical monocytes due to the elevated levels of

CCL5/RANTES in PASC (*5*). Further, our data suggests that interruption of the

CX3CR1/fractalkine pathway would be a potential therapeutic target to reduce the survival of

S1-containing non-classical monocytes and the associated vascular inflammation previously

discussed (*5*) and presented here.

It is important to note that the S1 protein detected in these patients appears to be retained from

prior infection or phagocytosis of infected cells undergoing apoptosis and is not the result of

persistent viral replication. Full length sequencing of the five cases submitted for genomic

analysis failed to identify any full-length sequence in the spike protein gene, or any other gene,

that could account for the observed spike protein detected by proteomic analysis. In contrast,

fragmented SARS-CoV-2 sequence was identified in all five of the cases. We have observed a

pattern of high Ct value or negativity by PCR, accompanied by scant, fragmented viral sequence

identified by whole viral genome sequencing over the past several months, a major shift from the

low Ct value, full length viral sequences identified throughout most of 2020. The reasons for this

shift are unclear, but as seen in these cases, it is unlikely these patients are producing any

replication competent viral genomes, and are thus incapable of transmitting the infection. In

contrast, the patients reported here appear to have developed an immune response to retained

viral antigens, specifically the S1 fragment of the spike protein, which continues to be presented

by CD16+ monocytes, eliciting an innate immune response characterized by elevated

inflammatory markers including interferon γ, IL-6, IL-10, and IL-2, among others. The body of

evidence reported here would not support continued viral replication. Instead, it implicates

dysregulation of innate immunity inflammatory mediators in response to persistent viral protein

presentation by CD16+ monocytes.

MATERIAL/METHODS

Patients

Following informed consent, whole blood was collected in a 10 mL EDTA tube and a 10 mL plasma

preparation tube (PPT). A total of 144 individuals were enrolled in the study consisting of 29 normal

individuals, 26 mild-moderate COVID-19 patients, 25 severe COVID-19 patients and 64 chronic COVID

(long hauler-LH) individuals. Long Haulers symptoms are listed in Figure 1. Study subjects were

stratified according to the following criteria.

Mild

1. Fever, cough, sore throat, malaise, headache, myalgia, nausea, diarrhea, loss of taste and small

2. No sign of pneumonia on chest imaging (CXR or CT Chest)

3. No shortness of breath or dyspnea

Moderate:

1. Radiological findings of pneumonia fever and respiratory symptoms

2. Saturation of oxygen (SpO2) ≥ 94% on room air at sea level

Severe

1. Saturation of oxygen (SpO2) < 94% on room air at sea level

2. Arterial partial pressure of oxygen (PaO2)/ fraction of inspired oxygen (FiO2) < 300mmHG

3. Lung infiltrate > 50% within 24 to 48 hours

4. HR ≥ 125 bpm

5. Respiratory rate ≥ 30 breaths per minute

Critical

1.   Respiratory failure and requiring mechanical ventilation, ECMO, high-flow nasal cannula oxygen

supplementation, noninvasive positive pressure ventilation (BiPAP, CPAP)

2.   Septic Shock- Systolic blood pressure < 90mmHg or Diastolic blood pressure < 60 mmHg or

requiring vasopressors (levophed, vasopressin, epinephrine

3.   Multiple organ dysfunction (cardiac, hepatic, renal, CNS, thrombotic disease)

<u>Post-acute COVID-19 (Long COVID)</u>

1.   Extending beyond 3 weeks from the initial onset of first symptoms

<u>Chronic COVID-19</u>

1.   Extending beyond 12 weeks from the initial onset of first symptoms (Table 1S)

High Parameter Immune Profiling/Flow Cytometry

Peripheral blood mononuclear cells were isolated from peripheral blood using Lymphoprep density

gradient (STEMCELL Technologies, Vancouver, Canada). Aliquots 200 of cells were frozen in media

that contained 90% fetal bovine serum (HyClone, Logan, UT) and 10% dimethyl sulfoxide (Sigma-

Aldrich, St. Louis, MO) and stored at -70°C. Cells were stained and analyzed using a 17-color antibody

cocktail including a PE-labeled SARS-CoV-2 S1 antibody (BioTechne, Minneapolis MN).

Digital Droplet PCR

A QIAamp Viral Mini Kit (Qiagen, Catalog #52906) was used to extract nucleic acids from 300 to 400

mL of plasma sample according to the manufacturer's instructions and eluted in 50 mL of AVE buffer

(RNase-free water with 0.04% sodium azide). The purified nucleic acids were tested immediately with a

Bio-Rad SARS-CoV-2 ddPCR Kit (Bio-Rad, Hercules, CA, USA). The panel was designed for

specifically detecting 2019-nCoV (two primer/probe sets). An additional primer/probe set was used to

detect the human RNase P gene in control samples and clinical specimens. RNA isolated and purified

from the plasma samples (5.5 mL) was added to a master mix comprising 1.1 mL of 2019-nCoV triplex

assay, 2.2 mL of reverse transcriptase, 5.5 mL of supermix, 1.1 mL of dithiothreitol, and 6.6 mL of

nuclease-free water.

The mixtures were then fractionated into up to 20,000 nanoliter-sized droplets in the form of a water-in-

oil emulsion in a QX200 Automated Droplet Generator (Bio-Rad, Hercules, CA). The 96-well real-time-

digital droplet polymerase chain reaction (RT-ddPCR) ready plate containing droplets was sealed with

foil using a plate sealer and thermocycled to reverse transcribe the RNA, before PCR amplification of

cDNA in a C1000 Touch thermocycler (Bio-Rad, Hercules, CA, USA). After PCR, the plate was loaded

into a QX200 Droplet Reader (Bio-Rad, Hercules, CA, USA) and the fluorescence intensity of each

droplet was measured in two channels (FAM and HEX). The fluorescence data were then analyzed with

QuantaSoft 1.7 and QuantaSoft Analysis Pro 1.0 Software (Bio-Rad, Hercules, CA, USA).

Flow Cytometric Cell Sorting

Cryopreserved PBMCs were quick-thawed, centrifuged, and washed in 2% BSA solution in D-

PBS. Cells were blocked for 5 min. in 2% BSA and then incubated at room temperature for 30

min. with Alexa Fluor® 488 Anti-CD45 antibody (IncellDx, 1/100 dilution), 2.5 ug of Alexa

Fluor® 647 Anti-CD16 antibody (BD, Cat. # 55710), and 1 ug of PerCP/Cy5.5 Anti-human

CD14 antibody (Biolegend, Cat. #325622). Cells were washed twice with 2% BSA/D-PBS,

filtered, and kept on ice for the duration of the cell sort. Data was acquired on a Sony SH800,

and only CD45+ cells staining positive for both CD14+ and CD16+ were sorted into test tubes

with 100 uL 2% BSA solution. Sort purity of control PBMCs was confirmed to be >99% by re-

analyzing sorted PBMCs using the same template and gating strategy.

Single Cell Protein Identification

Patient cells were sorted based on phenotypic markers (as above) and frozen at -80° C. Six

patient samples with positive flow cytometry signal and sufficient cell counts were chosen for

LCMS confirmation. Frozen cells were lysed with the IP Lysis/Wash Buffer from the kit

according to the manufacturer's protocol. 10 ug of anti-S1 mAb were used to immunoprecipitate

the S1 Spike protein from cell lysate of each patient. After overnight incubation with end-over-

end rotation at 4°C and then three washes with IP Lysis/Wash Buffer, bound S1 Spike protein

was eluted with the elution buffer from the kit.

IP elution fractions were dried *in vacuo*, resuspended in 20 uL of water, pooled, and purified by

Agilent 1290 UPLC Infinity II on a Discovery C8 (3cm x 2.1 mm, 5 µm, Sigma-Aldrich, room

temperature) using mobile phase solvents of 0.1% trifluoroacetic acid (TFA) in water or

acetonitrile. The gradient is as follows: 5-75% acetonitrile (0.1% TFA) in 4.5 min (0.8 mL/min),

with an initial hold at 5% acetonitrile (0.1% TFA) for 0.5 min (0.8 mL/min). The purified protein

was dried *in vacuo* and resuspended in 50 µL of 100 mM HEPES, pH 8.0 (20% Acetonitrile). 1

µL of TCEP (100 mM) was added and the samples were incubated at 37ºC for 30 min. 1 µL of

chloroacetamide (500 mM) was added to the samples and incubated at room temperature for 30

min. 1 µL rAspN (Promega 0.5 µg/µL) and 1 µL of LysC (Pierce, 1 µg/µL) were added and the

samples incubated at 37ºC for 16 h, prior to LCMS analysis.

LC-MS analysis

Digested recombinant SARS-CoV-2 Spike S1 protein was analyzed by a high mass accuracy

mass spectrometer to generate a list of detectable peptides with retention time and accurate

masses. An Agilent 1290 Infinity II high pressure liquid chromatography (HPLC) system and an

AdvanceBio Peptide Mapping column (2.1 × 150 mm, 2.7 μm) were used for peptide separation

prior to mass analysis. The mobile phase used for peptide separation consists of a solvent A

(0.1% formic acid in H_2O) and a solvent B (0.1% formic acid in 90% CH_3CN). The gradient was

as follows: 0–1 min, 3% B; 1– 30 min, to 40% B; 30–33 min, to 90% B; 33-35 min, 90% B; 37-

39 min, 3% B. Eluted peptides were electrosprayed using a Dual JetStream ESI source coupled

with the Agilent 6550 iFunnel time-of-flight MS analyzer. Data was acquired using the MS

method in 2 GHz (extended dynamic range) mode over a mass/charge range of 50–1700 Daltons

and an auto MS/MS method. Acquired data were saved in both centroid and profile mode using

Agilent Masshunter Workstation B09 Data acquisition Software. The same analytical method

was applied to immunoprecipitated samples from sorted patient cells except no ms/ms was

acquired.

Viral Genome Detection by PCR and Whole Viral Genome Sequencing

Ct Determination with TaqPath Assay

Five RNA samples were subjected to the TaqPath COVID-19 Combo Kit Assay (Thermo Fisher

Scientific Catalog no. A47814) to assess the cycle of threshold. TaqPath COVID-19 Combo Kit assay

was performed according to recommendations of the EUA, using the Applied BioSystems QuantStudio 7

Flex (Thermo Fisher Scientific Catalog no. 4485701).

Whole Genome Sequencing of Samples with Ion AmpliSeq

Five RNA samples were subjected to AmpliSeq library preparation using the Ion AmpliSeq Library Kit

2.0 (Thermo Fisher Scientific Catalog no. 4480441) and the Thermo Fisher Scientific Insight panel,

which consists of 238 amplicons in a two pool design against SARS-CoV-2 and seven amplicons as

human controls. Libraries were prepared following manufacturer's recommendations. Final libraries were

amplified using 5 cycles of amplification and libraries were cleaned up using 0.5X right sided cleanup

and 1.2X left sided cleanup using Kapa Pure Beads (Roche Catalog no l7983298001). Final libraries were

quantified using Ion Library TaqMan Quantitation Kit (ThermoFisher Catalog no. 4468802). Samples

were pooled in an equimolar distribution and loaded on to the Ion Chef Instrument (ThermoFisher

Catalog no. 4484177) for Templating onto a 510 chip. The prepared chip was then loaded onto a

GeneStudio S5 Prime (ThermoFisher Catalog no. A38196) for sequencing.

Genome Assembly, Quality Control, and Sequencing Analysis

Sequencing reads were aligned to the SARS-CoV-2 genome (build NC_045512.2) and human

transcriptome (build GRCh37) using the Thermo Fisher Scientific TMAP aligner. Default parameters

were used except for the *--context* flag.

Coverage analysis was performed by the coverage Analysis plugin in Thermo Fisher Scientific Torrent

Suite software. Reads in the human controls were evaluated for quality control. Per-base coverage,

average coverage, and percent genome covered at various depth thresholds were assessed using custom

software. Read length distribution versus read quality (MAPQ score) were further evaluated.

Variant calling was performed on SARS-CoV-2 using the variantCaller plugin. Callable regions were

identified as regions with read depth >= 20 after filtering reads with MAPQ < 10. Variants were filtered

for quality by removing mutations with allele frequency (AF) < 0.5 in the callable regions. Lineage

determination was made with pangoLEARN v1.2.13 using filtered-in mutations.

REFERENCES

1. R. Rubin. As Their Numbers Grow, COVID-19 "Long Haulers" Stump Experts. *JAMA*. **324,** 1381–1383 (2020). doi:10.1001/jama.2020.17709
2. https://www.cdc.gov/coronavirus/2019-ncov/hcp/clinical-care/post-covid-conditions.html (April 8, 2021).
3. X.H. Yao, Z.C. He, T.Y. Li, H. R. Zhang, Y. Wang, H. Mou, Q. Guo, S.C. Yu, Y. Ding, X. Liu, Y.F. Ping, X.W. Bian. Pathological evidence for residual SARS-CoV-2 in pulmonary tissues of a ready-for-discharge patient. *Cell Res*. **30**, 541-543 (2020). doi: 10.1038/s41422-020-0318-5. Epub 2020 Apr 28. PMID: 32346074; PMCID: PMC7186763.
4. R. Nienhold, Y. Ciani, V.H. Koelzer, A. Tzankov, J.D. Haslbauer, T. Menter, N. Schwab, M. Henkel, A. Frank, V. Zsikla, N. Willi, W. Kempf, T. Hoyler, M. Barbareschi, H. Moch, M. Tolnay, G. Cathomas, F. Demichelis, T. Junt, K.D. Mertz. Two distinct immunopathological profiles in autopsy lungs of COVID-19. *Nat Commun.* **11**, 5086 (2020). doi: 10.1038/s41467-020-18854-2.
5. B.K. Patterson, J. Guevarra-Coto, R. Yogendra, E. B. Francisco, E. Long, A. Pise, H. Rodrigues, P. Parikh, J. Mora. R.A. Mora-Rodrıguez. Immune-based prediction of COVID-19 severity and chronicity decoded using machine learning. *Front Immunol* (2021) https://doi.org/10.3389/fimmu.2021.700782.
6. T.S. Kapellos, L. Bonaguro, I. Gemünd, N. Reusch, A. Saglam, E. R Hinkley, J. L. Schultze. Human monocyte subsets and phenotypes in major chronic inflammatory diseases. *Front Immunol* **10,** 1–13 (2020) https://doi.org/10.3389/fimmu.2019.02035.
7. L. Ziegler-Heitbrock. The CD14+ CD16+ blood monocytes: their role in infection and inflammation. *J Leuk Biol* **81,** 584–592 (2007). https://doi.org/10.1189/jlb.0806510.
8. M. Rutkowska-Zapała, M. Suski, R. Szatanek, M. Lenart, K. Węglarczyk, R. Olszanecki, T. Grodzicki, M. Strach, J. Gąsowski, M. Siedlar. Human monocyte subsets exhibit divergent angiotensin I-converting activity *Clin Exp Immunol* **181,** 126–132 (2015).
9. R. Mukherjee, P. Kanti Barman, P. Kumar Thatoi, R. Tripathy, B. Kumar Das, B. Ravindran. Non-Classical monocytes display inflammatory features: Validation in Sepsis and Systemic Lupus Erythematous. *Sci Rep*. **5**, 13886 (2015). doi: 10.1038/srep13886.
10. D. Michlmayr, P. Andrade, K. Gonzalez, A. Balmaseda, E. Harris. CD14+CD16+ monocytes are the main target of Zika virus infection in peripheral blood mononuclear cells in a paediatric study in Nicaragua. *Nature Microbiology*, **2,**1462–1470 (2017). https://doi.org/10.1038/s41564-017-0035-0
11. Coquillard G, Patterson BK. Determination of hepatitis C virus-infected, monocyte lineage reservoirs in individuals with or without HIV coinfection. *J Infect Dis*. **200,** 947-954 (2009). doi: 10.1086/605476. PMID: 19678757.
12. P. Ancuta, K.J.Kunstman, P. Autissier, T. Zaman, D. Stone, S.M. Wolinsky, D.Gabuzda. CD16+ monocytes exposed to HIV promote highly efficient viral replication upon differentiation into macrophages and interaction with T cells. *Virology* **344,** 267–276 (2006). https://doi.org/10.1016/j.virol.2005.10.027

13. A.A. Patel, Y. Zhang, J.N.Fullerton, L. Boelen,, A. Rongvaux, A.A. Maini, V.Bigley, R.A. Flavell, D.W. Gilroy, B. Asquith, D. Macallan, S. Yona. The fate and lifespan of human monocyte subsets in steady state and systemic inflammation. *J Exp Med.* **214,** 1913-1923 (2017). doi: 10.1084/jem.20170355. Epub 2017 Jun 12. PMID: 28606987; PMCID: PMC5502436.

14. P. Ancuta, K.Y. Liu KY, V. Misra, V.S. Wacleche, A. Gosselin, X. Zhou, D. Gabuzda. Transcriptional profiling reveals developmental relationship and distinct biological functions of CD16+ and CD16- monocyte subsets. *BMC Genomics* **10**, 403 (2009). doi: 10.1186/1471-2164-10-403. PMID: 19712453; PMCID: PMC2741492.

15. T. Tak, R. van Groenendael, P. Pickkers, L. Koenderman. Monocyte subsets are differentially lost from the circulation during acute inflammation induced by human experimental endotoxemia. *J Innate Immun.* **9,** 464-474 (2017). doi: 10.1159/000475665. Epub 2017 Jun 23. PMID: 28641299; PMCID: PMC6738874.

16. C. Auffray, D. Fogg, M. Garfa, G. Elain, O. Join-Lambert, S. Kayal, S. Sarnacki, A. Cumano, G. Lauvau, F. Geissmann. Monitoring of blood vessels and tissues by a population of monocytes with patrolling behavior. *Science* **317,** 666-670 (2007). doi: 10.1126/science.1142883. PMID: 17673663.

17. S.T. Gren, T.B. Rasmussen, S. Janciauskiene. A single-cell gene-expression profile reveals inter-cellular heterogeneity within human monocyte subsets. *PLoS One* **10,** e0144351 (2015). doi: 10.1371/journal.pone.0144351. PMID: 26650546; PMCID: PMC4674153.

18. K.L. Wong, J.J. Tai, W.C. Wong, H. Han, X. Sem, W.H. Yeap, P. Kourilsky, S.C. Wong. Gene expression profiling reveals the defining features of the classical, intermediate, and nonclassical human monocyte subsets. *Blood.* **118,** e16-31 (2011). doi: 10.1182/blood-2010-12-326355. Epub 2011 Jun 7. PMID: 21653326.

19. A. Jafarzadeh, P. Chauhan, B. Saha, S. Jafarzadeh, M. Nemati. Contribution of monocytes and macrophages to the local tissue inflammation and cytokine storm in COVID-19: Lessons from SARS and MERS, and potential therapeutic interventions. *Life Sci.* **257**, 118102 (2020). doi: 10.1016/j.lfs.2020.118102. Epub 2020 Jul 18. PMID: 32687918; PMCID: PMC7367812.

20. Y. Zhou, B. Fu, X. Zheng. Pathogenic T-cells and inflammatory monocytes incite inflammatory storms in severe COVID-19 patients, *National Science Review* **7,** 998–1002 (2020).

21. C.E. Olingy, C.L. San Emeterio, M.E. Ogle, J.R. Krieger, A.C. Bruce, D.D. Pfau, B.T. Jordan, S.M. Peirce, E.A. Botchwey. Non-classical monocytes are biased progenitors of wound healing macrophages during soft tissue injury. *Sci Rep.* **7,** 447 (2017). doi: 10.1038/s41598-017-00477-1. PMID: 28348370; PMCID: PMC5428475.

22. P.B. Narasimhan, P. Marcovecchio, A.A.J. Anouk, C.C. Hedrick. Nonclassical monocytes in health and disease. *Ann Rev Immunol* **37,** 439-456 (2019).

23. V. Vila-del Sol, C. Punzón, M. Fresno. IFN-gamma-induced TNF-alpha expression is regulated by interferon regulatory factors 1 and 8 in mouse macrophages. *J Immunol* **181**:4461-4470 (2008). doi: 10.4049/jimmunol.181.7.4461. PMID: 18802049.

24. T. Matsumiya, K. Ota, T. Imaizumi, H. Yoshida, H. Kimura, K. Satoh. Characterization of Synergistic Induction of CX3CL1/Fractalkine by TNF-α and IFN-γ in vascular endothelial cells: an essential role for TNF-α in post-transcriptional regulation of CX3CL1. *J Immunol* **184,** 4205-4214 (2010). DOI: 10.4049/jimmunol.0903212.

25. S.M. Ong, E. Hadadi, T.M. Dang, W.H. Yeap, C.T. Tan, T.P. Ng, A. Larbi, S.C.Wong. The pro-inflammatory phenotype of the human non-classical monocyte subset is attributed to senescence. *Cell Death Dis* **9,** 266 (2008). https://doi.org/10.1038/s41419-018-0327-1.

26. C.F. Pereira, J. Middel, G. Jansen, H.S. Nottet. Enhanced expression of fractalkine in HIV-1 associated dementia. *J Neuroimmunol* **115,** 168-115 (2001). doi: 10.1016/s0165-5728(01)00262-4. PMID: 11282167.

27. A.R. Jajtner, J.R. Townsend, K.S. Beyer, A.N. Varanoske, D.D. Church, L.P. Oliveira, K.A. Herrlinger, S. Radom-Aizik, D.H. Fukuda, J.R. Stout, J.R. Hoffman. Resistance exercise selectively mobilizes monocyte subsets: role of polyphenols. *Med Sci Sports Exerc.* **50,** 2231-2241 (2018). doi: 10.1249/MSS.0000000000001703. PMID: 29957728.

Ethics
Informed consent was obtained from all participants.

Data and materials availability:
All requests for materials and raw data should be addressed to the corresponding author

Competing interests:
B.K.P, A.P., H.R., E.L, and EBF. are employees of IncellDx, Inc
TJT, PS, SH, DM are employees of Avrok Laboratories, Inc

Author contributions:
R.Y. and P.P. organized the clinical study and actively recruited patients.
B.K.P, A.P., H.R., X.E, E.L., J.B.S., TJT, PS, SH, DM performed experiments and analyzed the data.
J.G-C., R.A.M., J.M., X.C. performed the statistics and bioinformatics
B.K.P., J.M., EBF, J.G-C., R.A.M. wrote the draft of the manuscript and all authors contributed to revising the manuscript prior to submission.

Funding: None

573
574
575 **TABLE and FIGURE LEGENDS**
576
577 Table 1. Molecular analysis of study participants.

	Sars-CoV-2 RNA+		
COVID-19 Status	NS	PBMCs	Months Post-Infection
HC 1	-	-	n/a
HC 2	-	-	n/a
HC 3	-	-	n/a
HC 4	-	-	n/a
HC 5	-	-	n/a
HC 6	-	-	n/a
HC 7	-	-	n/a
HC 8	-	-	n/a
Asymptomatic	+	+	n/a
Severe 1	+	-	n/a
Severe 2	+	+	n/a
Severe 3	+	-	n/a
Severe 4	+	-	n/a
Severe 5	+	-	n/a
Severe 6	+	-	n/a
Severe 7	+	+	n/a
Severe 8	+	-	n/a
Severe 9	+	-	n/a
Severe 10	+	+	n/a
Severe 11	+	+	n/a
LH 1	+	-	13
LH 2	+	-	14
LH 3	+	-	6
LH 4	+	-	11
LH 5	+	+	15
LH 6	+	-	13
LH 7	+	-	12
LH 8	+	-	7
LH 9	+	-	14
LH 10	+	-	13
LH 11	+	-	12
LH 12	+	-	12
LH 13	+	-	6
LH 14	+	-	14
LH 15	+	-	13
LH 16	+	-	9

LH 17	+	-	11
LH 18	+	-	7
LH 19	+	-	14
LH 20	+	-	11
LH 21	+	-	13
LH 22	+	-	10
LH 23	+	-	8
LH 24	+	-	7
LH 25	+	-	12
LH 26	+	-	15

578 **Table 2: Average Coverage and Percent Bases Covered at 20x** While the percent of bases
579 covered varied across patients, all were less than 20% at 10X, and less at 20X coverage. In no
580 case was full length viral genome RNA detected, consistent with a lack of replication competent
581 viral infection.
582

Sample	Average Coverage	Percent Bases Covered at 10x	Percent Bases Covered at 20x
02-03_20210625	171.64	19.18	15.24
ABA-2_20210625	59.67	14.04	10.42
BGI-2_20210625	24.17	10.81	7.69
CST-2_20210625	40.29	11.71	7.79
RG_20210625	592.87	12.6743	10.16

603
604
605 **Figure 1**. Quantification of classical, intermediate and non-classical monocytes in PASC (LH).
606 Non-classical monocytes were significantly elevated in severe COVID-19 and in PASC.
607
608 **Figure 2.** High parameter flow cytometric quantification of SARS-CoV-2 S1 protein in
609 monocytic subsets. Cells were gated on CD45 then analyzed for CD14 and CD16 expression.
610 Classical monocytes are green, intermediate monocytes are red and non-classical monocytes are
611 blue.
612
613 **Figure 3.** Quantification of SARS-CoV-2 S1 protein in monocyte subsets isolated from healthy
614 controls (HC), severe COVID-19 (severe), and PASC patients (LH). SARS-CoV-2 S1 protein
615 was expressed in non-classical monocytes in both severe and PASC individuals. The amount of
616 expression was statistically significant.
617
Figure 4. LCMS confirmation of the presence of S1 subunit in samples LH1-6. A. Extracted ion
chromatogram (EIC) displaying the NLREFVFK peptide. The retention time matches that of the
NLREFVFK peptide in the commercial S1 standard. B. Mass Spectra of the NLREFVFK from
both the commercial standard and patient LH1. The Spectra show the same mass and isotope
distribution.

Figure 5: High Quality Mutations in the Callable Regions. Only fragmented viral RNA was
identified in the five patients, but multiple mutations throughout the viral genome were
identified, the vast majority of which were unique to each patient. Overall coverage was less than
20%, and no complete sequence in any portion of the viral genome was detected, including in the
spike gene encoding the S1 subunit identified by protein analysis in these patients.

A

B

Pure S1 protein
Peptide sequence: NLREFVFK
Calc. *m/z* (M+2H)$^{2+}$: 526.7980
Obsv. *m/z* (M+2H)$^{2+}$: 526.7958
Error (ppm) : 4.2

LH1 Sample
Peptide sequence: NLREFVFK
Calc. *m/z* (M+2H)$^{2+}$: 526.7980
Obsv. *m/z* (M+2H)$^{2+}$: 526.7954
Error (ppm) : 4.9