1

Functional divergence of the *bag of marbles* gene in the *Drosophila*

melanogaster species group

Jaclyn E. Bubnell[1*], Cynthia K.S. Ulbing[1], Paula Fernandez Begne[1], and Charles F.

Aquadro[1*]

[1] Department of Molecular Biology and Genetics, Cornell University, Ithaca, NY, USA

*Correspondence: Jaclyn Bubnell, jeb486@cornell.edu and Charles Aquadro,

cfa1@cornell.edu

Authors performed the following: JEB - conceptualization, investigation, validation,

formal analysis, and writing. CKSU and PFB - investigation, and validation, CFA -

conceptualization, funding acquisition, supervision, and writing.

Abstract

In *Drosophila melanogaster,* a key germline stem cell (GSC) differentiation factor, *bag of marbles* (*bam*) shows rapid bursts of amino acid fixations between sibling species *D. melanogaster* and *D. simulans*, but not in the outgroup species *D. ananassae*. Here, we test the null hypothesis that *bam's* differentiation function is conserved between *D. melanogaster* and four additional *Drosophila* species in the *melanogaster* species group spanning approximately 30 million years of divergence. Surprisingly, we demonstrate that *bam* is not necessary for oogenesis or spermatogenesis in *D. teissieri* nor is *bam* necessary for spermatogenesis in *D. ananassae*. Remarkably *bam* function may change on a relatively short time scale. We further report tests of neutral sequence evolution at *bam* in additional species of *Drosophila* and find a positive, but not perfect, correlation between evidence for positive selection at *bam* and its essential role in GSC regulation and fertility for both males and females. Further characterization of *bam* function in more divergent lineages will be necessary to distinguish between *bam's* critical gametogenesis role being newly derived in *D. melanogaster*, *D. simulans*, *D. yakuba,* and *D. ananassae* females or it being basal to the genus and subsequently lost in numerous lineages.

Introduction

In most sexually reproducing animals, egg and sperm production begins with the differentiation of germline stem cells. These germline stem cells (GSCs) divide both to self-renew to maintain the germline and to produce daughter cells that differentiate to produce gametes (Kahney et al. 2019). The cellular processes of daughter cell differentiation and stem cell self-renewal are the most fundamental functions of stem cells and are therefore highly regulated and presumed to be conserved. In the germline, the mis-regulation of either of these processes leads to sterility (Gleason et al. 2018). However, many of the genes that control GSC maintenance and GSC daughter differentiation in *D. melanogaster* are rapidly evolving due to positive selection for amino acid diversification (Civetta et al. 2006; Bauer DuMont et al. 2007; Choi and Aquadro 2015; Flores, DuMont, et al. 2015).

One such gene is *bag-of-marbles* (*bam*) which exhibits a rapid burst of amino acid substitutions between the sister species *D. melanogaster* and *D. simulans*, acts as the key switch for GSC daughter cell differentiation in *D. melanogaster* females, and is necessary for terminal differentiation of spermatogonia in males (McKearin and Spradling 1990; Civetta et al. 2006; Bauer DuMont et al. 2007; Insco et al. 2009). Despite 60 fixed amino acid differences between *D. melanogaster* and *D. simulans,* a *D. simulans bam* transgene is sufficient for GSC daughter cell differentiation in *D. melanogaster* females and males. The intriguing observations that *bam* is both rapidly evolving due to positive selection and functions as a switch for GSC daughter differentiation in *D. melanogaster* has led us to investigate what could be driving its positive selection between species (Flores, Bubnell, et al. 2015).

In females, *bam* is repressed in GSCs, and upon its expression promotes differentiation by binding to the protein product of *benign gonial cell neoplasm* (*bgcn*) as well as a series of other protein partners to repress the translation of self-renewal factors *nanos* and *eIF4a* (Li et al. 2009; Ting 2013)*.* The GSC daughter cell differentiates into a cystoblast, which then goes through four rounds of mitotic divisions while *bam* localizes

to the fusome, a cellular structure that interconnects the resulting dividing cells that

make up the cyst. In these cells, Bam functions with Bgcn to regulate mitotic synchrony

(Chen and McKearin 2003). Before meiosis begins, *bam* expression is repressed. In

males, *bam* is expressed in GSCs and its expression increases as differentiation

progresses (Insco et al. 2009). In early dividing spermatogonia, Bam protein must reach

a threshold to bind with partners Bgcn and *tumorous testis* (*tut*) to repress *mei-P26* and

switch the cellular program from proliferation to terminal differentiation to begin meiosis

(Chen et al. 2014). After this switch, *bam* expression is rapidly repressed. Loss of *bam*

function in either sex results in the over-proliferation of either GSCs in females or

spermatogonia in males causing tumors and sterility (McKearin and Spradling 1990).

One possible explanation for the observed signal of positive selection at *bam* is a

change of function. As *bam* regulates gametogenesis, it is possible that life history or

environmental factors drive positive selection at *bam*. It is even possible that the genetic

interaction between the endosymbiotic bacteria *Wolbachia pipientis* and *bam* has led to

such changes in *bam*'s function in different lineages, as *Wolbachia* infects the GSC

niche and is propagated through the germline (Flores, Bubnell, et al. 2015). While

functional divergence of *bam* seems less likely due to its essential role in GSC

maintenance and differentiation in *D. melanogaster, bam* sequence is highly divergent

among species. It is even difficult to confidently align *bam* sequence between *D.*

melanogaster and *D. ananassae,* both members of the *melanogaster* species group

(subgenus Sophophora), let alone the many more divergent *Drosophila* species (Bauer

DuMont et al. 2007). Additionally, *bam* appears to be a novel gene in the genus

Drosophila as no orthologues are reported in any non-*Drosophila* genomes using

orthoDB v10 (Kriventseva et al. 2019) and Ensembl Metazoa (Howe et al. 2020) (both

accessed 11/4/2021).

Here, we investigate the possibility that the observed signals of positive selection in *D.*

melanogaster and *D. simulans* at *bam* are due to selection for a novel function for *bam*

in gametogenesis in these lineages. To do this, we ask if *bam* is necessary for

gametogenesis in diverse species within the *Drosophila* genus. We generate *bam* loss-

of-function alleles in five *Drosophila* species in the *melanogaster* species group (*D.*

melanogaster, *D. simulans*, *D. yakuba*, *D. teissieri*, and *D. ananassae*). Surprisingly, we

find that *bam* is not necessary for gametogenesis in one or both sexes in two species

within the *melanogaster* species group. We also analyze additional population samples

of species mainly within the *melanogaster* species group and a few more distant

lineages for nucleotide sequence variation at *bam.* Within the *melanogaster* species

subgroup (*D. melanogaster*, *D. simulans*, *D. yakuba*, *D. santomea*, and *D. teissieri*), we

find evidence of lineage-specific signals of adaptation for three species, but not for *D.*

teissieri or *D. santomea*. Outside of the *melanogaster* species subgroup, we find similar

patterns of lineage-specific signals of positive selection. Together with our *bam* loss-of-

function results, these observations raise the possibility that the critical role for *bam* in

GSC differentiation is newly derived in the *melanogaster* species group and the

signatures of positive selection we observe in some of those species is associated with

species-specific refinements of *bam*'s GSC differentiation function or species-specific

adaptive pressures acting on *bam*'s GSC differentiation function.

Results

Generating *bam* null alleles for non-*melanogaster* species

To ask if *bam's* GSC function is conserved in species outside of *D. melanogaster*, we

designed and generated a series of *bam* null alleles in five species in the *melanogaster*

species group to assay for GSC daughter differentiation (Fig 1A). To do this, we used

CRISPR/Cas9 to introduce a 3xP3-YFP or a 3xP3-DsRed gene cassette into the first

exon of *bam*, thereby disrupting the *bam* coding sequence and introducing a premature

termination codon (Fig 1B). This also generates an allele that is trackable by eye color,

which is necessary for the non-*melanogaster* species of which we do not have balancer

chromosomes to maintain alleles that cause sterility. To generate the homozygous *bam*

null genotype, we cross the 3xP3-YFP line to the 3xP3-DsRed line and select flies with

both DsRed and YFP positive eyes. This scheme also allows us to use the same cross

to assay the heterozygous and wildtype *bam* siblings (Table 1).

We used *D. melanogaster* as a control to ensure our method would generate a loss of

function *bam* phenotype, as the *bam* null phenotype has been well characterized in this

species (Table 1). We also generated *bam* nulls in four other species within the

melanogaster species group *D. simulans*, *D. yakuba*, *D. teissieri,* and *D. ananassae*

(Fig 1A).

bam is necessary for female and male fertility in *D. melanogaster*, *D. simulans*,

and *D. yakuba*, but only in *D. ananassae* females, and in neither *D. teissieri* males

nor females.

In *D. melanogaster*, both male and female *bam* nulls are sterile as early germline cells

fail to differentiate and proceed through gametogenesis. In classic null alleles of *D.*

melanogaster bam (*bam*^{Δ59} and *bam*^{Δ86}), one copy of functional *bam* is sufficient to

rescue the *bam* null sterility phenotype. We first asked if our novel *bam* null alleles

behaved this way in *D. melanogaster*. Heterozygotes of each 3xP3-DsRed or 3xP3-YFP

disruption line showed similar fertility to wildtype *bam* in both males and females (mean

difference of adult progeny, not significant, permutation test) and transheterozygous

males and females carrying both the 3xP3-DsRed and the 3xP3-YFP disruptions (which

we will term the *bam* null genotype) were completely sterile (mean difference of adult

progeny, P < 0.0001, permutation test) (Fig 1 , File S2). We observed this same pattern

for *D. simulans* and *D. yakuba* (Fig 1, File S2). We observed some sterile *D. simulans*

females and males across all *bam* genotypes, which we attribute to the strain and

unlikely to be related to *bam* (Fig 1, File S2). In contrast, in *D. teissieri* we found that

there was no difference in fertility between any of the *bam* genotypes tested for females

or males (mean difference of adult progeny, not significant, permutation test) (Fig 1, File

S2). The *bam* null genotype did not result in sterility, or even reduced fertility. In *D.*

ananassae, *bam* null females were sterile (mean difference of adult progeny, P <

0.0001, permutation test), and heterozygotes of each 3xP3-DsRed or 3xP3-YFP

disruption did not significantly reduce fertility (mean difference of adult progeny, not

significant, permutation test). However, in *D. ananassae* males, *bam* nulls were fertile,

with no significant mean difference in progeny across the tested *bam* genotypes (Fig 1,

File S2). Therefore, *bam* is not necessary for fertility in all species in the *melanogaster*

species group, or even all species within the *melanogaster* species subgroup.

Consistent with the fertility results, *bam* is also necessary for germ cell

differentiation in *D. simulans* and *D. yakuba* females and males, but not in *D.*

***teissieri* males or females, and only *D. ananassae* females.**

The ovaries of *D. melanogaster bam* null females feature over-proliferating germline

stem cells that fail to differentiate into cystoblasts (McKearin and Spradling 1990; Ting

2013). This results in the germarium and developing downstream cysts accumulating

small, GSC-like cells resembling tumors (Fig 3A). In the testes of *D. melanogaster bam*

null males spermatogonia differentiation is blocked. The resulting testes feature over-

proliferating spermatogonia that fail to develop into terminally differentiated

spermatocytes (Fig 3A) (Insco et al. 2009; Insco et al. 2012).

We imaged ovaries from 2-5 day old *D. melanogaster* females from each *bam*

genotype. We assayed for the characteristic *bam* null tumor phenotype using antibodies

to Vasa to visualize the morphology of the germline and Hts-1B1 to visualize the

fusome, which fails to fully form in *bam* mutants (Lavoie et al. 1999; Hinnant et al.

2020). We found that the *bam* null genotype resulted in the characteristic *bag of*

marbles phenotype consistent with classic *bam* null alleles, with tumorous ovaries filled

with undifferentiated GSC-like cells (Fig 3B). One copy of wildtype *bam* was sufficient to

rescue the *bam* null phenotype, as has been previously observed for other *bam* null

alleles (Fig 3B) (Lavoie et al. 1999; Flores, Bubnell, et al. 2015). Therefore, our 3xP3-

DsRed and 3xP3-YFP disruptions successfully recapitulate the *bam* null phenotype in

females. We imaged testes from 3-5 day old *D. melanogaster* males and also

immunostained with antibodies to Vasa to label the germline and Hts-1B1 to label the

fusome. We found that the *bam* null genotype resulted in tumorous testes filled with

small over-proliferating cells that fail to differentiate into spermatocytes (Fig 3B). We

found that one copy of wildtype *bam* was sufficient to rescue the *bam* null tumor

phenotype, as has been previously reported for classic *bam* null alleles (Gönczy et al.

1997; Shivdasani and Ingham 2003)(Fig 3B). Therefore, our *bam* null alleles also

recapitulate the previously described *bam* null phenotype in *D. melanogaster* males.

We repeated the same analysis for the other species in the study. We found for *D.*

simulans and *D. yakuba* both female and male homozygous *bam* null genotypes

exhibited the characteristic ovarian and testes tumors as described in *D. melanogaster*

(Fig 3B). In these two species for both males and females, one copy of *bam* was also

sufficient to rescue the *bam* null phenotype as observed in *D. melanogaster* (Fig 3B).

However, in the *D. teissieri bam* null genotype, we observed no evidence of GSC over-

proliferation in females or spermatogonia over-proliferation in males. We observe the

same phenotype for the heterozygous alleles in both males and females, as well as

wildtype (Fig 3B). This indicates that *bam* is not necessary for GSC daughter

differentiation in *D. teissieri* females nor is *bam* necessary for spermatogonia

differentiation in males. In *D. ananassae bam* null females, we observed the

characteristic *bam* null phenotype observed in *D. melanogaster* with over-proliferating

GSC-like cells in the germarium and cysts and undeveloped fusomes (Fig 3B). One

copy of *bam* was sufficient to rescue the *bam* null tumorous phenotype (Fig 3B).

Therefore the *D. ananassae* female *bam* null phenotype is consistent with that of *D.*

melanogaster, *D. simulans*, and *D. yakuba*. In contrast, in *D. ananassae* males of the

bam null genotype we did not observe over-proliferating spermatogonia (Fig 3B). The

homozygous *bam* null genotype resulted in the same phenotype as the heterozygous

bam null genotypes and wildtype indicating that *bam* is not necessary for the switch

from proliferating spermatogonia to differentiated spermatocytes in *D. ananassae*

males. Therefore, these cytological data are fully consistent with our fertility assay

observations.

Confirming the loss-of-function status of *bam* in the *bam* null disruption design

The only available Bam antibody is weakly cross-reactive in *D. melanogaster* and *D.*

simulans thus we cannot directly demonstrate the absence of Bam protein in all the

species tested (Flores 2013; Flores, Bubnell, et al. 2015). Although we used the same

design to generate the *bam* null alleles across this study and therefore expect the 3xP3-

DsRed and 3xP3-YFP disruptions to have the same effect on *bam* in all five species, we

further confirmed the loss-of-function status of the *bam* disruption alleles in *D.*

ananassae and *D. teissieri* given their functional divergence*.*

First, we asked if *bam* had undergone duplication events in *D. ananassae* and *D.*

teissieri using publicly available sequence data and found no evidence of *bam*

duplication in either species (details described in S1 File, S8 Table). Next, since the

3xP3-DsRed and 3xP3-YFP insertions introduce a premature termination codon, we

expect the mRNA of the disrupted *bam* allele to be degraded through nonsense

mediated mRNA decay (NMD), but if the premature termination codon induced exon

skipping, the disrupted *bam* exon could be alternatively spliced out resulting in

expression of a truncated *bam* allele, which may be sufficient for function (Brogna and

Wen 2009; Mou et al. 2017; Chen et al. 2018; Sui et al. 2018). We used RT-qPCR to

ask if patterns of expression in the *bam* null alleles in *D. ananassae* males and *D.*

teissieri females and males were consistent with expectations for NMD. We found that

for *D. ananassae* and *D. teissieri*, the expression of the *bam* null disruption alleles were

reduced relative to the *bam* wildtype allele to levels consistent with NMD (~1-50%,

details in S1 File, Fig S1. Fig S2) (Pereverzev et al. 2015). Therefore, the mRNA of the

bam disruption alleles are likely targeted for degradation. Additionally, since both *bam*

null *D. teissieri* males and females show wildtype phenotypes we looked for evidence of

exon skipping. We asked if the exon disrupted by 3xP3-YFP or 3xP3-DsRed in the *bam*

null genotypes was expressed in proportion to the non-disrupted exons relative to the

wildtype genotype. We found that the ratio of exon 1 to exon 2 expression was not

significantly different between *D. teissieri* nulls and *D. teissieri* wildtype genotypes for

males and females (details in S1 File, Fig S3).

Although *bam* null *D. ananassae* males and *D. teissieri* males and females exhibited

wildtype phenotypes, together the genomic and RT-qPCR analyses indicate that the

bam null alleles are loss-of-function alleles. This evidence therefore supports our

conclusion that *bam* is not required for spermatogenesis in *D. ananassae* and *D.*

teissieri nor oogenesis in *D. teissieri.*

D. ananassae bam null males show evidence of germline defects

In _D. melanogaster_, Fas3 expression in the testes is limited to the hub, the testes
microenvironment that houses the somatic and germline stem cell populations at the
apical tip of the testes. As we found that _bam_ was necessary for oogenesis in _D.
ananassae_ but not for spermatogenesis, we assayed for more subtle evidence of
germline defects in _D. ananassae_ males. We immunostained testes with an antibody to
Fas3 to look for evidence of ectopic expression of Fas3 as in Gonzalez et al. (Gonzalez
et al. 2015). In _D. melanogaster bam_ null males, Fas3 is ectopically expressed, as the
spermatogonia over proliferate (Fig 4A&B). In _D. ananassae bam_ null males, we also
observed ectopic expression of Fas3 in cells outside of the apical tip of the testes,
although not nearly as severe as in _D. melanogaster bam_ null males (Fig 4D&E). We
then asked if there was any evidence of ectopic Fas3 expression in _D. teissieri bam_ null
males, as we also found that _bam_ was not necessary for spermatogenesis in _D.
teissieri._ In contrast to _D. ananassae_ and _D. melanogaster,_ we did not observe ectopic
expression of Fas3 in _D. teissieri bam_ null males (Fig 4B&C). This indicates that while
bam is not necessary for differentiation in _D. ananassae_ males, it may be playing a role
in regulating the early differentiating germ cell population.

**_bam_ shows lineage-specific bursts of positive selection throughout the
Drosophila genus.**

Previous studies have reported that _bam_ is evolving under strong positive selection
detected by the McDonald-Kreitman test in _D. melanogaster_ and _D. simulans_, but not in
D. ananassae (Civetta et al. 2006; Bauer DuMont et al. 2007; Choi and Aquadro 2014).
As these three species span about 30 million years of divergence, we wanted to know if
there was evidence of positive selection in lineages more closely related to _D.
melanogaster_ and _D. simulans_, or if this signal was unique to the individual _D.
melanogaster_ and _D. simulans_ lineages. We expanded the species sampled to include
three closely related species from the _yakuba_ complex: _D. yakuba, D. santomea,_ and _D.
teissieri_. We also took advantage of published population genomic data from samples of
inbred lines to expand our assessment of _bam_ variation in additional population

samples for *D. melanogaster* (Lack et al. 2015) and *D. simulans* and to newly assess

bam variation in population samples in the *montium* species group (*D. serrata, D.*

bunnanda, D. birchii, and *D. jambulina*) and the *ananassae* species subgroup (*D.*

bipectinata, D. pseudoananassae, and *D. pandora*) (Li et al. 2021).

In addition to further sampling closely related species to *D. melanogaster* and *D.*

simulans, we also wanted to ask if there was any evidence of adaptive evolution at *bam*

in lineages outside of the *melanogaster* group. We resequenced *bam* from population

samples of three more distantly related *Drosophila* species: *D. pseudoobscura*, *D.*

affinis, and *D. mojavensis* which span approximately 70 million years of divergence (Li

et al. 2021)*.* We also used published population genomic data from samples of inbred

lines to assess *bam* variation in *D. immigrans* and *D. rubida*, both members of the

immigrans species group (Li et al. 2021).

To ask if any population samples showed departures from selective neutrality consistent

with positive selection at *bam* similar to what we previously observed for *D.*

melanogaster and *D. simulans*, we used a lineage-specific McDonald-Kreitman test

(MKT) (Siddiq et al. 2017). The null hypothesis of the MKT is that the ratio of

nonsynonymous to synonymous polymorphism within a species is equal to the ratio of

nonsynonymous to synonymous fixed differences between species (or in our case, a

predicted common ancestral sequence). A previous study reported signatures of

positive selection at *bam* as an excess of nonsynonymous fixations in the *D.*

melanogaster and *D. simulans* lineages (Bauer DuMont et al. 2007). As *bam* appears to

be experiencing episodic signals of adaptive evolution restricted to individual lineages,

we measured lineage-specific divergence. We used codeml to generate reconstructed

ancestral sequences at the node of interest by maximum likelihood (Yang 1997; Yang

2007). We then used this sequence to calculate nonsynonymous and synonymous

divergences for each population sample. We excluded polymorphisms below 12% as

low-frequency mutations are likely to be slightly deleterious and reduce the power of the

MKT to identify positive selection (Fay et al. 2002; Charlesworth and Eyre-Walker

2008).

For *D. melanogaster*, we utilized the Zambia population sample from the *Drosophila* genome Nexus to measure *bam* variation (Lack et al. 2015). The Zambia sample contains the largest sample size of intact *bam* sequence (few missing calls), and represents variation found in the ancestral range of *D. melanogaster*. We used the predicted common ancestor of *D. melanogaster* and *D. simulans* and found similar patterns of polymorphism and divergence to the smaller samples of *D. melanogaster* from the USA and Zimbabwe previously reported by Bauer DuMont et al. (Bauer DuMont et al. 2007) (Fig 5). The *D. melanogaster* lineage shows a strong signal of positive selection (alpha = 0.943, P=0.001) (Fig 5).

For *D. simulans bam*, we utilized the Burnley, Victoria population sample from Li et al. (Li et al. 2021). We used the same common ancestral sequence as we did for the *D. melanogaster* analysis to calculate divergence. The *D. simulans* lineage shows a signal of positive selection for *bam* (alpha=0.722, P=0.036) are consistent with those previously reported by Bauer DuMont et al. (Bauer DuMont et al. 2007) (Fig 5). We also assessed *bam* variation in the *D. simulans* CA, USA population sample from Signor et al. (Signor et al. 2018), but due to elevated levels of genome-wide intermediate frequency polymorphisms caused by a recent population bottleneck, this sample violates key assumptions for the McDonald-Kreitman test (Fay 2011; Messer and Petrov 2013). For completeness, we report our findings for the CA sample in Supplementary File S1.

We used the predicted common ancestor of the *yakuba* complex (*D. yakuba*, *D. santomea*, *D. teissieri*) to estimate divergence for *bam* in each of the population samples. For *D. yakuba* we also observed a significant signal of positive selection for *bam* (alpha=0.893, P=0.002) (Fig 5). In contrast, we could not reject neutrality for *D. teissieri bam* or for *D. santomea bam.* The estimate of alpha is very low for *D. teissieri* (alpha=-0.011, P=0.983), but for *D. santomea* the alpha value was high, although not significant (alpha=0.76, P=0.186), which suggests it may be informative to sample *bam* from additional *D. santomea* populations with larger sample sizes (n=11 here) (Fig 5).

In the *montium* species subgroup, we used the predicted common ancestor of *D. serrata, D. bunnanda, D. birchii,* and *D. jambulina* to calculate divergence for *bam.* We found a significant signal of positive selection for *bam* in *D. serrata* (alpha=0.799, P=0.003)*, D. birchii* (alpha=0.675, P=0.003)*,* and *D. jambulina* (alpha=0.618, P=0.023) (Fig 5)*.* However, we could not reject neutrality for *D. bunnanda* (alpha=-1.014, P=0.542) (Fig 5).

In the *ananassae* species group, we used the predicted common ancestor of *D. bipectinata, D. pseudoananassae, D. pandora,* and *D. ananassae* to calculate divergence for *bam.* We reassessed lineage-specific divergence at *bam* for the *D. ananassae* population sample previously reported by Choi et al (Choi and Aquadro 2014) as they reported divergence between *D. ananassae* and *D. atripex*. We did not reject neutrality at *bam* for *D. bipectinata* (alpha=0.355, P=0.357)*, D. pseudoananassae* (alpha=-0.866, P=0.613)*,* or *D. pandora* (alpha=0.666, P=0.11) (Fig 5)*.* However, we found a significant signal of positive selection for *bam* in *D. ananassae* (alpha=1, P=0.045) (Fig 5). Therefore, the signals of positive selection at *bam* previously reported for *D. melanogaster* and *D. simulans* are not unique to the *melanogaster* species group.

In the *obscura* species group, we calculated divergence for *bam* from population samples of *D. pseudoobscura* and *D. affinis* from their predicted common ancestor and observed similar patterns of polymorphism and divergence (*D. pseudoobscura*: alpha=0.487, P=0.448), (*D. affinis*: alpha=-1.064, P=0.349), neither of which were close to rejecting neutral expectations (Fig 5). We used the predicted common ancestor of *D. mojavensis* and *D. arizonae bam* to calculate divergence in the *D. mojavensis* lineage and again did not observe a trend towards an excess of nonsynonymous divergence (alpha=0.285, P=0.771) (Fig 5). For the *immigrans* species group, we calculated divergence for *bam* from the predicted common ancestor of *D. immigrans* and *D. rubida,* We did not reject neutrality for *bam* in *D. immigrans* (alpha=0.116, P=0.882), but we did find a strong signal of positive selection for *bam* in *D. rubida* (alpha=1, P<0.000) (Fig 5).

Therefore, the signals of positive selection at *bam* are heterogeneously distributed
throughout the *Drosophila* genus.

Discussion

Bam's role in germline stem cell maintenance and differentiation shows
remarkable variation in the *Drosophila melanogaster* species group.

Our comparative assessment of *bam* requirement in gametogenesis across diverse
Drosophila species indicates that the functional evolution of *bam* is unexpectedly
complex. We discovered that *D. teissieri* and *D. ananassae* do not require *bam* for
gametogenesis in one or both sexes. A single orthologous copy of *bam* is found in all
Drosophila species examined to date, but is absent in outgroups for example, house fly
(Musca) and the sheep blow fly (Lucinia). These results raise the question as to whether
bam's critical role in GSC differentiation originally characterized in *D. melanogaster* is in
fact a phylogenetically recent gain of function, or whether *bam*'s critical role is basal to
the genus and that *bam*'s role has been recently (and possibly repeatedly) lost in
specific lineages. For example, assuming *bam*'s role in GSC differentiation is basal,
then the most parsimonious explanation of our functional data is that *bam* was required
for oogenesis and spermatogenesis in the common ancestor of the species tested (Fig
1A), then *bam* function was lost in *D. ananassae* males, and then lost again
independently in *D. teissieri* males and females. These alternative scenarios can be
distinguished by future follow-up analyses of *bam* function in more divergent *Drosophila*
species.

Evidence of positive selection for protein diversification at Bam is strikingly
heterogeneous across divergent *Drosophila* lineages.

We analyzed additional population samples of species mainly within the *melanogaster*
species group and a few more distant lineages for nucleotide sequence variation at
bam. We found new evidence of lineage-specific signals positive selection for
accelerated protein sequence evolution at *bam* in three of the *melanogaster* subgroup
species (*D. melanogaster, D. simulans*, and *D. yakuba*)*,* three of the *montium* subgroup

species (*D. serrata, D. birchii,* and *D. jambulina*), and one of the *ananassae* subgroup

species (*D. ananassae*). Additionally, we found evidence of positive selection in the *D.*

rubida lineage, indicating that the protein diversification at *bam* is repeatedly occurring

in species spanning over 70 million years of divergence.

While we could not reject neutrality for *bam* for the remaining lineages, It is important to

note as well that the MKT has limited power statistically when applied to single genes

(Akashi 1999; Zhai et al. 2009) and thus a failure to reject selective neutrality does not

prove that positive selection is not acting or has not acted on this gene.

Are stem cell genes hotspots for adaptive evolution?

The results of past studies of GSC evolution (Bauer DuMont et al. 2007; Choi and

Aquadro 2014; Choi and Aquadro 2015; Flores, DuMont, et al. 2015) and our population

genetic and functional results in this study lead to the question of why GSC

developmental functions are frequent targets of adaptation. It has been documented

that selfish transposable elements are upregulated in the absence of piRNAs, implying

that piRNAs act to protect the germline genome leading to an evolutionary conflict that

may drive positive selection at piRNA genes (Aravin et al. 2007; Simkin et al. 2013).

There is natural variation in the gene *Bruno* that provides tolerance to deleterious P-

element expression in the genome (Kelleher et al. 2018). In a scRNA-seq study of male

D. melanogaster testes, Witt et al. (Witt et al. 2019) report many fixed de novo genes

expressed in GSCs and early spermatogonia (where *bam* is expressed) at similar levels

to testis-specific genes. In contrast, segregating de novo genes show lower expression

in GSCs and early spermatogonia implying that cell type expression may impact the

likelihood for a de novo gene to become fixed. As de novo gene evolution is

increasingly recognized as a critical mode of adaptation, especially in reproduction,

GSCs/early spermatogonia may be generating necessary variation for adaptation (Witt

et al. 2019). A recent finding that the insect gene *oskar,* which is critical for germ cell

formation and oogenesis, arose *de novo* through a horizontal gene transfer event from a

prokaryote, resulting in a fusion between prokaryotic and eukaryotic sequence

illustrates yet another mechanism by which novel genes can take on key developmental

roles in reproduction (Jenny et al. 2006; Blondel et al. 2020).

The *bam* sequence itself has many properties of novel genes: it does not contain any

known DNA or RNA binding motifs, it binds with multiple protein partners, contains

structurally disordered regions, and is expressed in a tight spatiotemporal manner

(Chen et al. 2010; Jiang and Assis 2017; Wilson et al. 2017; Klasberg et al. 2018;

Vakirlis and McLysaght 2019). Novel genes that become fixed often gain critical roles in

conserved gene networks. Their new role may lead to selection to fine tune their critical

role thereby providing additional functional flexibility for these conserved processes

(Chen et al. 2010). Such a process could represent what has been called

developmental systems drift, a model to explain how developmental systems remain

phenotypically conserved while the genetic pathways that determine them diverge (True

and Haag 2001).

It is possible that *bam* was a gene novel to the common ancestor of *Drosophila* (and

possibly some other dipterans) and has functionally diversified across the *Drosophila*

genus, only gaining its role in germline stem cell differentiation in the lineage leading to

the *melanogaster* species group. Additionally, it is possible that *bam* quickly evolved a

role in GSC daughter differentiation in *Drosophila,* but that function has been shaped by

germline conflicts in specific lineages. Conflict with germline agents (such as *W.*

pipientis) may even drive the fixation of novel genes in new roles that successfully

evade their control. As *W. pipientis* must recognize host cues during oogenesis to

successfully infect the developing oocyte, it is possible that novel host genes that

regulate differentiation might then be under selection for their new function if it evades

unfavorable manipulation by *W. pipientis*.

It certainly seems plausible that genetic conflict is one way for new genes to evolve

required functions in conserved processes such as GSC daughter differentiation. The *D.*

melanogaster testes and the accessory gland show enriched expression of novel

genes, many of which show signals of adaptive evolution, and some of which are

predicted to be driven by sexual conflict (Levine et al. 2006; Zhou et al. 2008; Findlay et

al. 2009; Kaessmann 2010; Gubala et al. 2017; Witt et al. 2019). While the

developmental processes of gametogenesis and reproduction must be conserved, our

results demonstrate that the individual genes that function in that process may vary

among even closely related species.

Materials and Methods

Fly stocks and rearing:

We raised fly stocks on standard cornmeal molasses food at room temperature. We

used yeast glucose food for fertility assays. The *D. melanogaster* w^{1118} isogenic line and

w^{1118} balancer lines were gifts from Luis Teixeira, and the *D. simulans* w^{501} line (stock

14021-0251.011), the *D. yakuba* white eyed line (14021-226.03) and the *D. ananassae*

reference genome (14024-0371.14) lines are from the Drosophila species stock center

(http://blogs.cornell.edu/drosophila/). The *W. pipientis*-free *D. ananassae* line Nou83

was a gift from Artyom Kopp. The *D. teissieri* line (*teissieri* syn) was a gift from Daniel

Matute (Turissini et al. 2015).

We tested all lines for *W. pipientis* infection status by qPCR (NEB Luna Universal qPCR

kit) using primer sets for *wsp* and *arm* (S1 Table). Lines that tested positive (*D. teissieri*,

D. ananassae, *D. simulans*) were treated with tetracycline (200 µg/mL) for three

generations to remove *W. pipientis* (Glover et al. 1990). We allowed these lines to

recover for at least three generations before using them in any analyses. We were

unable to fully clear the *D. ananassae* reference line of *W. pipientis* with tetracycline, so

to obtain a *W. pipientis* free stock, we crossed males of our *bam* null lines to females

from the Nou83 line, which we confirmed to be *W. pipientis*-free by qPCR.

Cloning for bam null constructs:

We used Flybase to obtain the nucleotide sequence information for designing

constructs in *D. melanogaster* (Dmel 5.57)*, D. simulans* (dsim r2.01), *D. yakuba* (dyak

r1.04), and *D. ananassae* (dana r1.04) (Thurmond et al. 2019). We obtained sequence

information for the *D. teissieri* sequence from Turissini et al. 2017 (accession
SAMN07407364-SAMN07407376) (Turissini and Matute 2017). We used Geneious for
all cloning design.

We used the NEB Q5 High Fidelity 2X master mix to generate all PCR products. We gel
extracted and purified PCR products using the NEB Monarch DNA gel extraction kit or
Qiagen MinElute gel extraction kit. IDT primers were used for PCR, sequencing, and
cloning. We generated the donor plasmids for the 3xP3-DsRed and 3xP3-YFP bam
disruption lines using the NEB HiFi Assembly Cloning kit into the pHD-attP-DsRed
vector from flyCRISPR (Gratz et al. 2014) as follows: we amplified 1.5 kb homology
arms from genomic DNA of the appropriate species stock flanking the insertion site for
3xP3-DsRed or 3xP3-YFP. 3xP3-DsRed was amplified from the pHD-attP-DsRed
plasmid and 3xP3-YFP was amplified from the *D. simulans* nos-Cas9 line, a gift from
David Stern. We then gel extracted the two homology arms and the appropriate 3xP3
marker, purified them, and assembled them into the pHD vector backbone using the
manufacturer's protocol (S3 Table).

We used the U6:3 plasmid from CRISPRflydesign (Port et al. 2014) to express gRNAs
for generating the *D. yakuba* lines. The gRNA target sequence was generated by
annealing primers with the gRNA sequence and BbsI site overhangs, and ligating into
the BbsI ligated U6:3 vector (T4 ligase NEB, BbsI NEB) (S3 Table). We used NEBalpha
competent cells for transformations.

We prepped and purified all plasmids for embryo injections with the Qiagen plasmid
plus midi-prep kit. We additionally purified plasmids for injections with phenol-chloroform
extraction to remove residual RNases. We confirmed plasmid sequences with Sanger
sequencing (Cornell BRC Genomics Facility).

CRISPR/Cas9:
gRNA selection:

We chose gRNAs using the flyCRISPR target finder (Gratz et al. 2014) for *D.*

melanogaster, *D. simulans*, *D. yakuba*, and *D. ananassae*. For *D. teissieri*, we used one

completely conserved gRNA from *D. yakuba.* We also chose an additional gRNA from

D. yakuba that had two divergent sites between *D. yakuba* and *D. teissieri,* thus we

modified them to the *D. teissieri* sites. We selected gRNAs that had no predicted off-

targets in the reference genome (S4 Table & S5 Table). For *D. yakuba*, we used the

U6:3 gRNA plasmid from CRISPRfly design as described above. For the remaining

species (*D. melanogaster*, *D. simulans*, *D. teissieri*, and *D. ananassae*), we used

synthetic gRNAs (sgRNAs) from Synthego. We used 1-3 gRNAs per injection to

increase the chances of a successful CRISPR event. We used multiple gRNAs if there

were suitable options for that site to increase efficiency (multiple gRNAs within 50 bps

with 0 predicted off-targets).

<u>Injections</u>:

All CRISPR/Cas9 injections were carried out by Genetivision. Lines were injected with

the plasmid donor, gRNAs (synthetic or plasmid), Cas9 protein (Synthego), and an

siRNA for Lig4 (IDT DNA). The same siRNA was used for *D. melanogaster*, *D.*

simulans, *D. yakuba*, and *D. teissieri* (pers communication with Daniel Barbash and

David Stern S6 Table). We created a *D. ananassae*-specific siRNA due to sequence

divergence (S6 Table).

For the *bam* disruption lines, we screened for eye color in F1s in-house using a

Nightsea fluorescence system with YFP (cyan) and DsRed (green) filters. We

backcrossed all CRISPR/Cas9 mutants to the stock line we used for injections for three

generations. All mutants are maintained as heterozygous stocks. We confirmed all

CRISPR insertions by Sanger sequencing (Cornell BRC Genomics Core Facility).

<u>Genotypes for generating bam disruption homozygotes</u>:

We generated two *bam* disruption lines in each species in order to create the genotypes

for the comparative *bam* null analyses. We generated one by disrupting *bam* with 3xP3-

DsRed and the other with 3xP3-YFP. We selected DsRed or YFP positive flies and

crossed them to generate homozygous *bam* disruption flies (*bam^{3xp3-DsRed}/bam^{3xP3-YFP}*).

This cross also results in DsRed or YFP positive heterozygous *bam* null progeny as well

as *bam* wildtype progeny without any fluorescent eye marker.

We targeted the 1st exon of *bam* in all species in order to disrupt the *bam* coding

sequence and introduce an early stop codon. While we were successful in generating

1st exon disruptions in each species, in *D. simulans* we were only able to obtain a 3xP3-

YFP 1st exon disruption line. However, we did find that a *D. simulans* line with a 3xP3-

DsRed insertion in its 2nd intron resulted in homozygous sterile females and males with

tumorous ovaries and testes, while heterozygous females and males were fertile with

wildtype ovary and testes phenotypes. As this edit was specific to the *bam* locus, we

presumed it was an mRNA null allele and we used this line for our analyses. Thus, if the

phenotype of the 3xP3-DsRed 2nd intron allele over the 3xP3-YFP 1st exon allele is the

same as the homozygous 3xP3-DsRed 2nd intron allele, this is consistent with

classification as a genetic null.

Fertility assays:

<u>Female fertility</u>:

All female fertility assays were performed following the approach of Flores et al. (Flores

et al. 2015). Specifically, we collected virgin females and aged them based on their time

to sexual maturity 2-3 days (*D. melanogaster*, *D. simulans*, *D. yakuba,* and *D. teissieri*)

or 5-7 days (*D. ananassae*) (Markow and O'Grady 2006). We collected all genotypes

from each bottle to control for bottle effects. We collected virgin males of the wildtype

genotype for each species and aged them for the same timeframe as the females of

that species. We evenly distributed males collected from different bottles across the

female genotypes to control for any bottle effects. We crossed single females collected

within 48 hours of each other to two virgin males. We allowed the trio to mate for 9

days, and then flipped them to new vials. Except for *D. ananassae* and *D. yakuba*, as

fertility was very low, we cleared the trio after day 9 and did not continue further. For all

other species crosses, after an additional 9 days, we cleared the trio. Then, we counted

the progeny for each trio every two to three days to get the total adult progeny per

female.

Male fertility:

All male fertility assays were performed as described for the female fertility assays

except that one tester male was crossed to two virgin females.

Food and conditions for fertility assays:

For *D. melanogaster*, *D. simulans*, *D. ananassae*, and *D. teissieri*, we performed all

fertility assays on yeast-glucose food. For *D. yakuba*, we performed fertility assays on

cornmeal-molasses food, as this line did not thrive on the yeast-glucose food. We kept

all crosses and fertility experiments in an incubator at 25°C with a 12-hour light-dark

cycle.

Statistics:

For the fertility assays we used estimation statistics to assess the mean difference

(effect size) of adult progeny between the wildtype *bam* genotype and each individual

additional *bam* genotype. For generating the estimation statistics and plots (Ho et al.

2019), we used the dabest package in Python (v. 0.3.1) with 5,000 bootstrap resamples.

Estimation statistics provide a non-parametric alternative to other statistical tests of the

difference of the mean (for example, ANOVA), and allow us to understand the

magnitude of the effect of *bam* genotype on the fertility phenotype. In text, we report

significance as a mean difference (effect size) outside the 95% bootstrap confidence

interval. While one benefit of these statistics is that they are less biased than reporting

traditional P-values, we also report P-values for approximate permutation tests with

5,000 permutations.

Immunostaining:

We used the following primary antibodies, anti-vasa antibody from Santa Cruz

biologicals (anti rabbit 1:200), anti-Hts-1B1 from Developmental Studies Hybridoma

bank (anti-mouse 1:4 for serum, 1:40 for concentrate item), and anti-Fas3 from DSHB

(anti-mouse 1:50). We used the following secondary antibodies: Alexaflour 488, 568

(Invitrogen) at 1:500.

We performed immunostaining as described in Aruna et al. and Flores et al. (Aruna et

al. 2009; Flores, Bubnell, et al. 2015). Briefly, we dissected ovaries and testes in ice-

cold 1X PBS and pipetted ovaries up and down to improve antibody permeability. We

fixed tissues in 4% paraformaldehyde, washed with PBST, blocked in PBTA (Alfa

Aesar, with 0.1% Triton X added), and then incubated in the appropriate primary

antibody in PBTA overnight. We then washed (PBST), blocked (PBTA), and incubated

the tissue in the appropriate secondary antibody for 2 hours, then washed (PBST) and

mounted in mounting media with DAPI (Vecta shield or Prolong Glass with NucBlue) for

imaging.

Confocal Microscopy:

We imaged ovaries and testes on a Zeiss i880 confocal microscope with 405nm, 488

nm, and 568 nm laser lines at 40X (Plan-Apochromat 1.4 NA, oil) (Cornell BRC Imaging

Core Facility). We analyzed and edited images using Fiji (ImageJ).

RNA extraction:

For RT-qPCR experiments we dissected ovaries from 10 one-day old females per

biological replicate or testes from 15 one-day old males in ice cold 1x PBS. The tissue

was immediately placed in NEB DNA/RNA stabilization buffer, homogenized with a

bead homogenizer, and then stored at -80°C until RNA extraction. We used the NEB

Total RNA mini-prep kit for all RNA extractions according to the manufacturer's

protocol and included the on column DNAse treatment. We assessed RNA

concentration and quality with a Qubit 4 fluorometer and a NanoDrop.

RT-qPCR:

For all RT-qPCR analyses, we used the Luna One Step RT-qPCR kit from NEB

according to the manufacturer's instructions at 10µl total volume in 384 well plates. We

used a Viia7 qPCR machine with the appropriate cycle settings as described in the

Luna One Step RT-qPCR kit.

For the analysis of *bam* expression in *D. ananassae*, we used primers to *D. ananassae*

bam with *D. ananassae Rp49* as a control (S1 Table). To assess the possibility that the

bam disruption allele induced exon skipping of the disrupted exon 1 in *D. teissieri*, we

designed a primer pair that spanned exon 1 to exon 2 and another pair that spanned

exon 2 to exon 3 (S1 Table). We used the exon 2 - exon 3 target as the control. For

both the *D. teissieri* and *D. ananassae* assays described above, we included a standard

curve for each target on every plate at a 1:2 dilution and then used the standard curve

method to calculate the relative quantity of *bam* (to either *Rp49* for *D. ananassae* or

bam exon 2-exon 3 for *D. teissieri*) with the QuantStudio software using the wildtype

bam genotype as a control for each species.

For the allele specific analysis in *D. teissieri,* we designed one primer pair specific to

the *bam* 3xP3-DsRed and 3xP3-YFP insertion alleles and another primer pair specific

to the wildtype allele. We then used these two targets to measure the relative

expression of the insertion allele to the wildtype allele in a single heterozygous sample.

The primer pair for the *bam* insertion alleles target the sequence between the SV40-

polyA tail of the 3xP3-DsRed and 3xP3-YFP insertion and the *bam* sequence

downstream of the insertion (S1 Table). The primer pair for the *bam* wildtype allele

target *bam* upstream and downstream of the 3xP3-DsRed and 3xP3-YFP insertion,

and therefore will not amplify the large insertion allele under RT-qPCR conditions (S1

Table). We included a standard curve for each target on every plate at a 1:2 dilution

and then used the standard curve method in the QuantStudio software to calculate the

Ct and Ct standard error for each target and each sample. We report the relative

quantity of the 3xP3-DsRed or 3xP3-YFP allele using the wildtype allele in the same

sample as a control by calculating $2^{-(nullCt-wildtypeCt)}$.

Genomic analysis of *bam* copy number in *D. teissieri:*

While *bam* is reported as a 1:1 syntenic ortholog on FlyBase for *D. melanogaster, D.*

simulans, D. yakuba, and *D. ananassae,* data for *D. teissieri* is not available. To ask if

bam has been recently duplicated in *D. teissieri,* we generated a blast database (blastn

v 2.9.0) from a new de novo assembled *D. teissieri* reference genome (accession

GCA_016746235.1) and blasted *D. teissieri bam* from the resequencing described

above.

Additionally, to assess read coverage for *bam* compared to the surrounding loci and the

entire chromosome 3R, we used publicly available short read WGS data for *D. teissieri*

(SRR13202235) and used bwa-mem (v 0.7.17) to map reads to a new de novo

assembled *D. yakuba* reference genome (accession GCA_016746365.1). We visualized

the alignment and assessed read coverage using Geneious.

Population genetic analyses:

The *D. yakuba*, *D. santomea*, and *D. teissieri* population sample lines were gifted by

Brandon Cooper and Daniel Matute (Cooper et al. 2017). The *D. yakuba* and *D. teissieri*

lines were collected in Bioko in 2013, the *D. santomea* sample was collected in Sao

Tome in 2015. The *D. affinis* population sample lines were gifted by Robert Unckless

(S2 Table). The *D. mojavensis* population sample is from Organ Pipe National

Monument, Arizona and DNA from this sample was gifted by Erin Kelleher and Luciano

Matzkin. The *D. pseudoobscura* population sample is from Apple Hill, California, as

described in (Schaeffer and Miller 1991). We used previously prepared DNA from the *D.*

pseudoobscura population sample as described in (Schaeffer and Miller 1991).

We prepared genomic DNA from *D. affinis, D. yakuba*, *D. santomea*, and *D. teissieri*

samples using the Qiagen PureGene kit. We used Primer3 to generate primers to

amplify *bam* from *D. affinis, D. mojavensis, D. pseudoobscura, D. yakuba*, *D. santomea*,

and *D. teissieri* population samples (S7 Table). We used existing polymorphism data

(Daniel Matute, personal communication) to generate primers for *D. yakuba*, *D.*

santomea, and *D. teissieri*. We resequenced *bam* using Sanger sequencing from

5'UTR-3'UTR for *D. yakuba*, *D. santomea*, and *D. teissieri* and *bam* exons and introns

for *D. affinis* (Cornell BRC Genomics Facility). We used Geneious for aligning the

sequences and calling polymorphisms. We generated fasta files for each individual. We

used DnaSP to phase individuals with heterozygous sites, and then randomly picked

one allele for further analyses (Bauer DuMont et al. 2007). We will deposit all

resequencing data described here in Genbank.

We used polymorphism data for *D. melanogaster* from the Zambia population sample

from the Drosophila genome Nexus (Lack et al. 2015), as this is the largest population

sample with intact *bam* sequence and represents a population in the *D. melanogaster*

ancestral range. For the *D. melanogaster* Zambia population sample, we removed a

single individual that contained missing nucleotide calls in the *bam* coding sequence

(ZI117). As we filtered polymorphisms at < 12% frequency, omitting this single individual

would not affect our downstream analyses.

Polymorphism for *D. simulans*, *D. serrata*, *D. bunnanda*, *D. birchii*, *D. jambulina*, *D.*

bipectinata, *D. pseudoananassae, D. pandora*, *D. immigrans*, and *D. rubida* as well as

the single *bam* sequence for *Z. bogoriensis* that we used as an outgroup

are from Li et al 2021(accession PRJNA736147) (Li et al. 2021).

D. ananassae bam polymorphism, and single *D. atripex bam* sequences are from Choi

et al. (Choi and Aquadro 2014). We obtained single *bam* sequences for species used

for outgroups from FlyBase (*D. erecta*), and Genbank (*D. eugracilis, D. arizonae, D.*

navojoa, D. miranda, D. subobscura).

We performed all multiple sequence alignments for the *bam* coding sequences using

PRANK (v.170427) with the -codon and -F parameters and using the PRANK generated

guide tree. As *bam* is highly divergent across the *Drosophila* genus, we generated

multiple sequence alignments only among closely related clusters of species to ensure

alignment quality. We generated a multiple sequence alignment of the *melanogaster*

subgroup species (*D. melanogaster, D. simulans, D. yakuba, D. santomea, D. teissieri,*

D. erecta, D. eugracilis) and *montium* subgroup species (*D. serrata*, *D. bunnanda*, *D.*

birchii, *D. jambulina*), a multiple sequence alignment of *D. ananassae, D. bipectinata, D.*

pseudoananassae, D. pandora, and *D. atripex* , a multiple sequence alignment of *D.*

mojavensis, D. arizonae, and D. navajo, a multiple sequence alignment of *D. affinis, D.*

pseudoobscura, D. miranda, and *D. subobscura,* and a multiple sequence alignment of

D.immigrans, D. rubida, and *Z. bogoriensis*.

We used only unique sequences for input to codeml. We used the codeml package from

PAML (v. 4.9) (Yang 1997; Yang 2007) to generate the predicted common ancestor

sequences to calculate lineage-specific divergence for *bam* with the McDonald-

Kreitman test. We used the PRANK alignments and their corresponding gene trees as

input to codeml with the following control file parameters (noisy = 9, verbose = 2,

runmode = 0, seqtype = 1, CodonFreq = 2, clock = 0, aaDist = 0, model = 0, NSsites =

0, icode = 0, getSE = 0, RateAncestor = 1, Small_Diff = .5e-6, cleandata = 0, method =

1).

We used the http://mkt.uab.cat/mkt/mkt.asp webtool to perform the standard McDonald-

Kreitman test (MKT) comparing nonsynonymous vs. synonymous changes (Egea et al.

2008). We excluded polymorphic sites at < 12% frequency, as these are likely slightly

deleterious alleles that have not yet been purged by purifying selection (Charlesworth

and Eyre-Walker 2008). For each species where we had polymorphism data, we

performed the MKT using the specified predicted common ancestral sequence to

calculate lineage-specific divergence. We report the values of the contingency table, the

P-value of the Chi-Square test, as well as alpha, the proportion of fixations predicted to

be due to positive selection (Eyre-Walker 2006).

Summary of accession numbers for genomes used in this analysis:

For read coverage analysis of *bam*: *D. teissieri*: SRR13202235, GCA_016746235.1. *D.*

yakuba: GCA_016746365.1. For cloning *D. teissieri bam* null constructs:

SAMN07407364-SAMN07407376. For *bam* polymorphism from *D. simulans, D. serrata,*

D. bunnanda, D. birchii, D. jambulina, D. bipectinata, D. pseudoananassae, D. pandora,

D. immigrans, and D. rubida: PRJNA736147.

Acknowledgements:

We thank Santiago Herrera Alvarez for resequencing *bam* from *D. affinis* and Jae
Young Choi for resequencing *bam* from *D. pseudoobscura* and *D. mojavensis.* We
thank Daniel Matute, Brandon Cooper, Robert Unckless, Erin Kelleher, and Luciano
Matzkin for the population samples they provided for this study. We thank Fang Li for
working to make the variant call files from (Li et al. 2021) quickly available for this study.
Additionally, we thank David L. Stern for his advice and guidance in designing the *bam*
null alleles and for an extremely thoughtful and critical review of an earlier draft of the
manuscript. We are grateful to Mariana F. Wolfner, Andrew G. Clark, Miwa Wenzel and
Catherine Kagemann for valuable input and discussion of these experiments and results
and helpful comments on this manuscript. Lastly, we thank the three anonymous
reviewers for their thoughtful and insightful feedback for improving this manuscript.

Data Availability:

Sequence data generated in this study will be deposited in GenBank prior to publication.
The remaining data underlying this article are available in the article and in its online
supplementary material. Fly lines, CRISPR/Cas9 reagents, and plasmids used in this
study are available upon request.

Funding:

This work was supported by National Institutes of Health grant R01-GM095793 to
C.F.A. and by the National Science Foundation Graduate Research Fellowship Program
DGE-114415 to J.E.B.

REFERENCES

Akashi H. 1999. Inferring the fitness effects of DNA mutations from polymorphism and

divergence data: Statistical power to detect directional selection under stationarity

and free recombination. Genetics. 151(1). doi:10.1093/genetics/151.1.221.

Aravin AA, Hannon GJ, Brennecke J. 2007. The Piwi-piRNA pathway provides an

adaptive defense in the transposon arms race. Science (80-). 318(5851).

doi:10.1126/science.1146484.

Aruna S, Flores HA, Barbash DA. 2009. Reduced fertility of Drosophila melanogaster

Hybrid male rescue (Hmr) mutant females is partially complemented by Hmr

orthologs from sibling species. Genetics. doi:10.1534/genetics.108.100057.

Bauer DuMont VL, Flores HA, Wright MH, Aquadro CF. 2007. Recurrent positive

selection at Bgcn, a key determinant of germ line differentiation, does not appear to

be driven by simple coevolution with its partner protein bam. Mol Biol Evol.

24(1):182–191. doi:10.1093/molbev/msl141.

Blondel L, Jones TEM, Extavour CG. 2020. Bacterial contribution to genesis of the

novel germ line determinant Oskar. Elife. doi:10.7554/eLife.45539.

Brogna S, Wen J. 2009. Nonsense-mediated mRNA decay (NMD) mechanisms. Nat

Struct Mol Biol. doi:10.1038/nsmb.1550.

Charlesworth J, Eyre-Walker A. 2008. The McDonald-Kreitman test and slightly

deleterious mutations. Mol Biol Evol. doi:10.1093/molbev/msn005.

Chen D, McKearin D. 2003. Dpp Signaling Silences bam Transcription Directly to

Establish Asymmetric Divisions of Germline Stem Cells. Curr Biol. 13(20):1786–

1791. doi:10.1016/j.cub.2003.09.033.

Chen D, Tang JX, Li B, Hou L, Wang X, Kang L. 2018. CRISPR/Cas9-mediated

genome editing induces exon skipping by complete or stochastic altering splicing in

the migratory locust. BMC Biotechnol. doi:10.1186/s12896-018-0465-7.

Chen D, Wu C, Zhao S, Geng Q, Gao Y, Li X, Zhang Y, Wang Z. 2014. Three RNA

Binding Proteins Form a Complex to Promote Differentiation of Germline Stem Cell

Lineage in Drosophila. PLoS Genet. 10(11):e1004797.

doi:10.1371/journal.pgen.1004797.

Chen S, Zhang YE, Long M. 2010. New Genes in Drosophila Quickly Become

Essential. Science (80-). 330(6011):1682–1685. doi:10.1126/science.1196380.

Choi JY, Aquadro CF. 2014. The coevolutionary period of Wolbachia pipientis infecting

Drosophila ananassae and its impact on the evolution of the host germline stem

cell regulating genes. Mol Biol Evol. doi:10.1093/molbev/msu204.

Choi JY, Aquadro CF. 2015. Molecular evolution of drosophila germline stem cell and

neural stem cell regulating genes. Genome Biol Evol. 7(11):3097–3114.

doi:10.1093/gbe/evv207.

Civetta A, Rajakumar SA, Brouwers B, Bacik JP. 2006. Rapid evolution and gene-

specific patterns of selection for three genes of spermatogenesis in Drosophila. Mol

Biol Evol. 23(3):655–662. doi:10.1093/molbev/msj074.

Cooper BS, Ginsberg PS, Turelli M, Matute DR. 2017. Wolbachia in the Drosophila

yakuba complex: Pervasive frequency variation and weak cytoplasmic

incompatibility, but no apparent effect on reproductive isolation. Genetics.

205(1):333–351. doi:10.1534/genetics.116.196238.

Egea R, Casillas S, Barbadilla A. 2008. Standard and generalized McDonald-Kreitman

test: a website to detect selection by comparing different classes of DNA sites.

Nucleic Acids Res. doi:10.1093/nar/gkn337.

Eyre-Walker A. 2006. The genomic rate of adaptive evolution. Trends Ecol Evol.

doi:10.1016/j.tree.2006.06.015.

Fay JC. 2011. Weighing the evidence for adaptation at the molecular level. Trends

Genet. 27(9). doi:10.1016/j.tig.2011.06.003.

Fay JC, Wyckoff GJ, Wu CI. 2002. Testing the neutral theory of molecular evolution with

genomic data from Drosophila. Nature. 415(6875). doi:10.1038/4151024a.

Findlay GD, MacCoss MJ, Swanson WJ. 2009. Proteomic discovery of previously

unannotated, rapidly evolving seminal fluid genes in Drosophila. Genome Res.

19(5):886–896. doi:10.1101/gr.089391.108.

Flores HA. 2013. Evolutionary and Functional Analysis of the Drosophila Bag of Marbles

Gene. Cornell University.

Flores HA, Bubnell JE, Aquadro CF, Barbash DA. 2015. The Drosophila bag of marbles

Gene Interacts Genetically with Wolbachia and Shows Female-Specific Effects of

Divergence. PLoS Genet. 11(8). doi:10.1371/journal.pgen.1005453.

Flores HA, DuMont VLB, Fatoo A, Hubbard D, Hijji M, Barbash DA, Aquadro CF. 2015.
Adaptive evolution of genes involved in the regulation of germline stem cells in
Drosophila melanogaster and D. simulans. G3 Genes, Genomes, Genet. 5(4):583–
592. doi:10.1534/g3.114.015875.

Gleason RJ, Anand A, Kai T, Chen X. 2018. Protecting and diversifying the germline.
Genetics. 208(2):435–471. doi:10.1534/genetics.117.300208.

Glover DM, Raff J, Karr TL, O'Neill SL, Lin H, Wolfner MF. 1990. Parasites in
Drosophila embryos. Nature. 348:117. doi:10.1038/348117a0.

Gönczy P, Matunis E, DiNardo S. 1997. bag-of-marbles and benign genial cell
neoplasm act in the germline to restrict proliferation during Drosophila
spermatogenesis. Development. 124(21):4361–4371.
doi:10.1242/dev.124.21.4361.

Gonzalez J, Qi H, Liu N, Lin H. 2015. Piwi Is a Key Regulator of Both Somatic and
Germline Stem Cells in the Drosophila Testis. Cell Rep. 12(1):150–161.
doi:10.1016/J.CELREP.2015.06.004.

Gratz SJ, Ukken FP, Rubinstein CD, Thiede G, Donohue LK, Cummings AM, Oconnor-
Giles KM. 2014. Highly specific and efficient CRISPR/Cas9-catalyzed homology-
directed repair in Drosophila. Genetics. 196(4):961–971.
doi:10.1534/genetics.113.160713.

Gubala AM, Schmitz JF, Kearns MJ, Vinh TT, Bornberg-Bauer E, Wolfner MF, Findlay
GD. 2017. The Goddard and Saturn Genes Are Essential for Drosophila Male
Fertility and May Have Arisen de Novo. Mol Biol Evol. doi:10.1093/molbev/msx057.

Hinnant TD, Merkle JA, Ables ET. 2020. Coordinating Proliferation, Polarity, and Cell
Fate in the Drosophila Female Germline. Front Cell Dev Biol. 8:19.
doi:10.3389/fcell.2020.00019.

Ho J, Tumkaya T, Aryal S, Choi H, Claridge-Chang A. 2019. Moving beyond P values:
data analysis with estimation graphics. Nat Methods. 16(7):565–566.
doi:10.1038/s41592-019-0470-3.

Howe KL, Contreras-Moreira B, De Silva N, Maslen G, Akanni W, Allen J, Alvarez-
Jarreta J, Barba M, Bolser DM, Cambell L, et al. 2020. Ensembl Genomes 2020-
enabling non-vertebrate genomic research. Nucleic Acids Res. 48(D1).

doi:10.1093/nar/gkz890.

Insco ML, Bailey AS, Kim J, Olivares GH, Wapinski OL, Tam CH, Fuller MT. 2012. A

self-limiting switch based on translational control regulates the transition from

proliferation to differentiation in an adult stem cell lineage. Cell Stem Cell.

11(5):689–700. doi:10.1016/j.stem.2012.08.012.

Insco ML, Leon A, Tam CH, McKearin DM, Fuller MT. 2009. Accumulation of a

differentiation regulator specifies transit amplifying division number in an adult stem

cell lineage. Proc Natl Acad Sci. 106(52):22311–22316.

doi:10.1073/PNAS.0912454106.

Jenny A, Hachet O, Závorszky P, Cyrklaff A, Weston MDJ, St Johnston D, Erdélyi M,

Ephrussi A. 2006. A translation-independent role of oskar RNA in early Drosophila

oogenesis. Development. 133(15):2827–2833. doi:10.1242/dev.02456.

Jiang X, Assis R. 2017. Natural selection drives rapid functional evolution of young

drosophila duplicate genes. Mol Biol Evol. 34(12):3089–3098.

doi:10.1093/molbev/msx230.

Kaessmann H. 2010. Origins, evolution, and phenotypic impact of new genes. Genome

Res. 20(10):1313–1326. doi:10.1101/gr.101386.109.

Kahney EW, Snedeker JC, Chen X. 2019. Regulation of Drosophila germline stem cells.

Curr Opin Cell Biol. 60:27–35. doi:10.1016/j.ceb.2019.03.008.

Kelleher ES, Jaweria J, Akoma U, Ortega L, Tang W. 2018. QTL mapping of natural

variation reveals that the developmental regulator bruno reduces tolerance to P-

element transposition in the Drosophila female germline. PLoS Biol. 16(10).

doi:10.1371/journal.pbio.2006040.

Klasberg S, Bitard-Feildel T, Callebaut I, Bornberg-Bauer E. 2018. Origins and

structural properties of novel and *de novo* protein domains during insect evolution.

FEBS J. 285(14):2605–2625. doi:10.1111/febs.14504.

Kriventseva E V., Kuznetsov D, Tegenfeldt F, Manni M, Dias R, Simão FA, Zdobnov

EM. 2019. OrthoDB v10: Sampling the diversity of animal, plant, fungal, protist,

bacterial and viral genomes for evolutionary and functional annotations of

orthologs. Nucleic Acids Res. 47(D1). doi:10.1093/nar/gky1053.

Lack JB, Cardeno CM, Crepeau MW, Taylor W, Corbett-Detig RB, Stevens KA, Langley

CH, Pool JE. 2015. The drosophila genome nexus: A population genomic resource

of 623 Drosophila melanogaster genomes, including 197 from a single ancestral

range population. Genetics. doi:10.1534/genetics.115.174664.

Lavoie CA, Ohlstein B, McKearin DM. 1999. Localization and function of bam protein

require the benign gonial cell neoplasm gene product. Dev Biol.

doi:10.1006/dbio.1999.9346.

Levine MT, Jones CD, Kern AD, Lindfors HA, Begun DJ. 2006. Novel genes derived

from noncoding DNA in Drosophila melanogaster are frequently X-linked and

exhibit testis-biased expression. National Academy of Sciences.

Li F, Rane R V, Luria | Victor, Xiong Z, Chen J, Li Z, Catullo RA, Griffin PC, Schiffer M,

Pearce S, et al. 2021. Phylogenomic analyses of the genus Drosophila reveals

genomic signals of climate adaptation. Mol Ecol Resour. 00:1–23.

doi:10.1111/1755-0998.13561.

Li Y, Minor NT, Park JK, McKearin DM, Maines JZ. 2009. Bam and Bgcn antagonize

Nanos-dependent germ-line stem cell maintenance. Proc Natl Acad Sci U S A.

106(23):9304–9309. doi:10.1073/pnas.0901452106.

Markow TA, O'Grady P. 2006. Drosophila: A Guide to Species Identification and Use.

Elsevier Science.

McKearin DM, Spradling AC. 1990. Bag-of-marbles: A Drosophila gene required to

initiate both male and female gametogenesis. Genes Dev. 4(12 B):2242–2251.

doi:10.1101/gad.4.12b.2242.

Messer PW, Petrov DA. 2013. Frequent adaptation and the McDonald-Kreitman test.

Proc Natl Acad Sci U S A. 110(21):8615–8620. doi:10.1073/pnas.1220835110.

Mou H, Smith JL, Peng L, Yin H, Moore J, Zhang XO, Song CQ, Sheel A, Wu Q, Ozata

DM, et al. 2017. CRISPR/Cas9-mediated genome editing induces exon skipping by

alternative splicing or exon deletion. Genome Biol. 18(1):108. doi:10.1186/s13059-

017-1237-8.

Pereverzev AP, Gurskaya NG, Ermakova G V., Kudryavtseva EI, Markina NM, Kotlobay

AA, Lukyanov SA, Zaraisky AG, Lukyanov KA. 2015. Method for quantitative

analysis of nonsense-mediated mRNA decay at the single cell level. Sci Rep.

5(1):1–10. doi:10.1038/srep07729.

Port F, Chen HM, Lee T, Bullock SL. 2014. Optimized CRISPR/Cas tools for efficient
germline and somatic genome engineering in Drosophila. Proc Natl Acad Sci U S
A. 111(29). doi:10.1073/pnas.1405500111.

Schaeffer SW, Miller EL. 1991. Nucleotide sequence analysis of Adh genes estimates
the time of geographic isolation of the Bogota population of Drosophila
pseudoobscura. Proc Natl Acad Sci U S A. 88(14):6097–6101.
doi:10.1073/pnas.88.14.6097.

Shivdasani AA, Ingham PW. 2003. Regulation of Stem Cell Maintenance and Transit
Amplifying Cell Proliferation by TGF-β Signaling in Drosophila Spermatogenesis.
Curr Biol. doi:10.1016/j.cub.2003.10.063.

Siddiq MA, Loehlin DW, Montooth KL, Thornton JW. 2017. Experimental test and
refutation of a classic case of molecular adaptation in Drosophila melanogaster.
Nat Ecol Evol. 1(2). doi:10.1038/s41559-016-0025.

Signor SA, New FN, Nuzhdin S. 2018. A large panel of drosophila simulans reveals an
abundance of common variants. Genome Biol Evol. doi:10.1093/gbe/evx262.

Simkin A, Wong A, Poh YP, Theurkauf WE, Jensen JD. 2013. Recurrent and recent
selective sweeps in the piRNA pathway. Evolution (N Y). 67(4):1081–1090.
doi:10.1111/evo.12011.

Sui T, Song Y, Liu Z, Chen M, Deng J, Xu Y, Lai L, Li Z. 2018. CRISPR-induced exon
skipping is dependent on premature termination codon mutations. Genome Biol.
doi:10.1186/s13059-018-1532-z.

Thurmond J, Goodman JL, Strelets VB, Attrill H, Gramates LS, Marygold SJ, Matthews
BB, Millburn G, Antonazzo G, Trovisco V, et al. 2019. FlyBase 2.0: the next
generation. Nucleic Acids Res. 47(D1):D759–D765. doi:10.1093/NAR/GKY1003.

Ting X. 2013. Control of germline stem cell self-renewal and differentiation in the
Drosophila ovary: Concerted actions of niche signals and intrinsic factors. Wiley
Interdiscip Rev Dev Biol. 2(2):261–273. doi:10.1002/wdev.60.

True JR, Haag ES. 2001. Developmental system drift and flexibility in evolutionary
trajectories. Evol Dev. 3(2):109–119. doi:10.1046/j.1525-142X.2001.003002109.x.

Turissini DA, Liu G, David JR, Matute DR. 2015. The evolution of reproductive isolation
in the Drosophila yakuba complex of species. J Evol Biol. 28(3):557–575.

doi:10.1111/jeb.12588.

Turissini DA, Matute DR. 2017. Fine scale mapping of genomic introgressions within the

Drosophila yakuba clade. PLoS Genet. 13(9). doi:10.1371/journal.pgen.1006971.

Vakirlis N, McLysaght A. 2019. Computational Prediction of De Novo Emerged Protein-

Coding Genes. In: Methods in Molecular Biology. Vol. 1851. Humana Press Inc. p.

63–81.

Wilson BA, Foy SG, Neme R, Masel J. 2017. Young genes are highly disordered as

predicted by the preadaptation hypothesis of de novo gene birth. Nat Ecol Evol.

1(6). doi:10.1038/s41559-017-0146.

Witt E, Benjamin S, Svetec N, Zhao L. 2019. Testis single-cell RNA-seq reveals the

dynamics of de novo gene transcription and germline mutational bias in drosophila.

Elife. doi:10.7554/eLife.47138.

Yang Z. 1997. Paml: A program package for phylogenetic analysis by maximum

likelihood. Bioinformatics. doi:10.1093/bioinformatics/13.5.555.

Yang Z. 2007. PAML 4: Phylogenetic analysis by maximum likelihood. Mol Biol Evol.

doi:10.1093/molbev/msm088.

Zhai W, Nielsen R, Slatkin M. 2009. An investigation of the statistical power of neutrality

tests based on comparative and population genetic data. Mol Biol Evol. 26(2).

doi:10.1093/molbev/msn231.

Zhou Q, Zhang G, Zhang Y, Xu S, Zhao R, Zhan Z, Li X, Ding Y, Yang S, Wang W.

2008. On the origin of new genes in Drosophila. Genome Res. 18(9):1446–1455.

doi:10.1101/gr.076588.108.

1033

Fig 1. Species chosen to generate *bam* null alleles using CRISPR/Cas9 to disrupt *bam* with a trackable eye marker cassette. (A) The five species we chose to generate *bam* null alleles span approximately 30 million years of divergence. (B) Schematic for generating null alleles with trackable eye markers using CRISPR/Cas9. We generate two null alleles using different color eye markers and cross the heterozygotes together to generate an easily trackable null genotype.

Fig 2. *Bam* is necessary for female and male fertility in *D. melanogaster, D. simulans, and D. yakuba,* but not for *D. ananassae* males nor both *D. teissieri* males and females. Swarm plots showing the total progeny per female or per male for each *bam* genotype and Cumming estimation plots showing the 95% confidence interval effect size. Dotted boxes show genotypes where the phenotype did not match that of *D. melanogaster.* For *D. melanogaster, D. simulans, and D. yakuba*, the *bam* null DsRed/YFP females and males were sterile (P < 0.0001, permutation test), and the heterozygous genotypes did not have a significant effect on fertility (P > 0.05, permutation test). For *D. teissieri,* the *bam* null DsRed/YFP females and males did not have significantly different fertility from wildtype, nor did the heterozygotes. For *D. ananassae*, the *bam* null DsRed/YFP females were sterile (P < 0.0001, permutation test), and the heterozygotes did not affect fertility. However, the *bam* null DsRed/YFP males were fertile and there was no significant difference between the null and heterozygotes and wildtype fertility (P>0.05, permutation test).

Fig 3. *Bam* is necessary for female and male germline stem cell differentiation in *D. melanogaster, D. simulans, and D. yakuba,* but not for *D. ananassae* males nor *D. teissieri* males and females. (A) Schematics of ovaries and testes showing the *bam* wildtype and null cytological phenotypes in *D. melanogaster*. Wildtype *bam* ovaries exhibit developing egg chambers with differentiated nurse cells and testes exhibit large, differentiated spermatocytes. Null *bam* ovaries exhibit egg chambers filled with small GSC-like tumors that contain undeveloped fusomes and testes exhibit cysts filled with small over-proliferating spermatogonia. (B) Ovaries and testes from all five species of

wildtype (*bam⁺/bam⁺*), heterozygous (YFP/+ or DsRed/+), and null (DsRed/YFP) *bam* genotypes. All tissue is labeled with antibodies to Vasa (germline) and Hts-1B1 (fusome). Dashed boxes indicate genotypes and tissue that are not consistent with the *D. melanogaster* phenotype. Ovaries and testes from all five species exhibit developing nurse cells or spermatocytes (respectively) for wildtype and heterozygous *bam* genotypes. We did not observe any tumorous cysts (arrowheads). Ovaries and testes from *bam* null *D. melanogaster, D. simulans, and D. yakuba* all exhibited cysts filled with GSC-like tumors (ovaries) or over-proliferating spermatogonia (testes) (arrows). The *bam* null ovaries exhibit undeveloped fusomes. *D. ananassae bam* null ovaries exhibited the tumorous egg chamber phenotype (arrow), however testes exhibited large developing spermatocytes (arrowhead) and we did not observe any tumors. The *bam* null *D. teissieri* ovaries and testes exhibited nurse cell positive egg chambers and developing spermatocytes (respectively) (arrowheads) and we did not observe any tumors.

Fig 4. Loss of *bam* function in *D. ananassae* males shows ectopic expression of Fas3, a marker expressed in the hub, but not in *D. teissieri*. All testes tissue is labeled with antibodies for Vasa (germline) and Fas3 (hub, arrowheads). Significance was assessed with a Fisher's exact test comparing counts of testes with ectopic Fas3 expression between wildtype and *bam* null testes. (A) *D. melanogaster bam* wildtype testes with Fas3 expression limited to the anterior tip of the testes hub (inset) and featuring large differentiating spermatogonia (Fas3 ectopic expression n=0/13). (B) *D. melanogaster bam* null testes feature ectopic Fas3 expression and small over-proliferating spermatogonia (Fas3 ectopic expression n=8/11, P<0.000). (C) *D. teissieri bam* wildtype testes with Fas3 expression limited to the anterior tip of the testes (hub) and featuring large differentiating spermatogonia (Fas3 ectopic expression n=0/20). (D) *D. teissieri bam* null testes with Fas3 expression limited to the anterior tip of the testes (hub) and large differentiating spermatogonia (Fas3 ectopic expression n=0/16, P=1). (E) *D. ananassae bam* wildtype testes with Fas3 expression limited to the hub and featuring large differentiating spermatogonia (Fas3 ectopic expression n=31). (F) *D. ananassae bam* null

testes with ectopic Fas3 expression and large differentiating spermatogonia (Fas3 ectopic expression n=19/27, P<0.000).

Fig 5. *Bam* shows lineage-specific signals of positive selection scattered across the *Drosophila* genus. Cladogram of *Drosophila* species we sampled in this study. Pink highlighted lineages represent lineages where we have evidence of positive selection at *bam* with the McDonald Kreitman test. Blue highlighted lineages are lineages of which we could not reject neutrality with the McDonald Kreitman test. The 2X2 table from the McDonald Kreitman test is shown on the right (N = nonsynonymous, S = synonymous). Alpha is the proportion of nonsynonymous fixations predicted to be due to positive selection. P-value is from the Chi-square test, n is sample size.

[1]Polymorphism data from Choi and Aquadro 2014.

Fig S1. RT-qPCR experiment to assess *bam* mRNA in *D. ananassae bam* null genotypes reported as the relative quantity (RQ) of *bam* expression to the housekeeping control RP49. *bam* expression in testes of the null (DsRed/YFP) genotype is significantly reduced (standard error) relative to *bam* expression in testes of the wildtype genotype at approximately 10%, consistent with expectations for NMD. The heterozygous genotypes DsRed/+ and YFP/+ do not show reduced *bam* expression relative to wildtype.

Fig S2. RT-qPCR experiment to detect patterns consistent with nonsense mediated mRNA decay of the *bam* disruption allele in *D. teissieri* reported as the relative quantity (RQ) of mRNA from the *bam* disruption allele to the *bam* wildtype allele in heterozygotes. Significance reported as standard error (A) Expression of the *bam* null allele relative to the *bam* wildtype allele in ovaries. Both heterozygous genotypes DsRed/+ and YFP/+ show less than 10% of expression of the *bam* null allele

relative to the wildtype allele, consistent with expectations for NMD. The wildtype negative control shows little to no expression of the null allele as expected. (B) Expression of the *bam* null allele relative to the *bam* wildtype allele in testes. Both heterozygous genotypes DsRed/+ and YFP/+ show less than 50% of expression of the *bam* null allele relative to the wildtype allele, consistent with expectations for NMD. The wildtype negative control shows little to no expression of the null allele as expected.

Fig S3. RT-qPCR experiment to detect evidence of exon skipping of the exon containing the *bam* disruption reported as the relative quantity (RQ) of *bam* exon 1 to exon 2. Significance reported as standard error (A) Expression of *bam* exon 1 relative to exon 2 in ovaries. The RQ of exon 1 to exon 2 in the *bam* null DsRed/YFP genotype is not significantly different than the RQ of exon 1 to exon 2 in the *bam* wildtype genotype, indicating that expression of the exon containing the disruption is comparable to the unaffected exon in both the null and wildtype genotype, and therefore exon skipping is not likely occurring in females. (B) Expression of *bam* exon 1 relative to exon 2 in testes. The RQ of exon 1 to exon 2 in the *bam* null DsRed/YFP genotype is not significantly different than the RQ of exon 1 to exon 2 in the *bam* wildtype genotype, indicating that expression of the exon containing the disruption is comparable to the unaffected exon in both the null and wildtype genotype, and therefore exon skipping is not likely occurring in males.

Table 1. Abbreviations for *bam* genotypes for all species and the expected phenotypes as observed in *D. melanogaster*.

Genotype	Abbreviation	*D. melanogaster* Phenotype Expectations		
		Eye Color	Fertility	Cytological
bam⁺/bam⁺	wildtype	white	wildtype	wildtype
bam^{3xP3-DsRed}/bam⁺	DsRed/+	DsRed	wildtype	wildtype
bam^{3xP3-YFP}/bam⁺	YFP/+	YFP	wildtype	wildtype
bam^{3xP3-DsRed}/bam^{3xP3-YFP}	DsRed/YFP	DsRed & YFP	sterile	tumorous ovaries and testes

Fig1

A.

B.

Fig2

Adult progeny per fly for each *bam* genotype

female   male

melanogaster

simulans

yakuba

teissieri

ananassae

Fig3

A. *D. melanogaster bam* cytological phenotypes

ovaries — wildtype / null
females — fusome/spectrosome

B.

bam disruption genotype

wildtype   heterozygous   null

melanogaster

simulans

yakuba

teissieri

ananassae

Vasa

Fig4

bam disruption genotype

wildtype   null

Vasa   Fas3

A. *melanogaster*   B.

C. *teissieri*   D.

E. *ananassae*   F.

Fig5

	Dn	Ds	Pn	Ps	alpha	P-value	n
D. melanogaster	33	13	1	7	0.943	*0.001*	187
D. simulans	34	17	5	9	0.722	*0.036*	20
D. yakuba	26	18	2	13	0.893	*0.002*	14
D. santomea	24	23	1	4	0.76	0.186	11
D. teissieri	11	13	3	1	-0.181	0.789	11
D. serrata	61	52	4	7	0.799	*0.003*	18
D. bunnanda	67	45	3	1	-1.014	0.542	20
D. birchii	40	39	13	39	0.675	*0.003*	20
D. jambulina	30	28	2	13	0.618	*0.023*	19
D. bipectinata	23	14	18	17	0.355	0.357	20
D. pseudoanananassae	30	22	3	13	-0.866	0.613	20
D. pandora	27	30	3	10	0.666	0.11	17
D. ananassae†	28	33	0	5	1	*0.045*	14
D. pseudoobscura	24	40	2	4	0.487	0.448	15
D. affinis	31	32	2	1	-1.064	0.349	14
D. mojavensis	2	5	2	7	0.285	0.771	22
D. immigrans	60	53	3	3	0.116	0.882	20
D. rubida	75	73	0	14	1	*0.000*	19

Predicted ancestral sequence used to calculate divergence

melanogaster species subgroup

montium species subgroup

ananassae species subgroup

melanogaster species group

obscura species group

repleta group

immigrans species group

Fig S1

Fig S2

A. *D. teissieri* ovaries

B. *D. teissieri* testes

Fig S3

A. *D. teissieri* ovaries

B. *D. teissieri* testes