Use of untargeted magnetic beads to capture *Mycobacterium smegmatis* and *Mycobacterium*

***avium paratuberculosis* prior detection by mycobacteriophage D29 and Real-Time-PCR**

Gabriel Rojas-Ponce[a*], Dominic Sauvageau[b], Roger Zemp[c], Herman W. Barkema[d], and Stephane Evoy[a]

[a]Department of Electrical and Computer Engineering, University of Alberta, Edmonton,

AB, Canada

[b]Department of Chemical and Materials Engineering, University of Alberta, Edmonton, AB, Canada.

[c]Department of Electrical and Computer Engineering, Biomedical Engineering, University of Alberta, Edmonton,

AB, Canada.

[d]Department of Production Animal Health, Faculty of Veterinary Medicine, University of Calgary, Calgary, AB,

Canada

*Corresponding Author: gabrielr_p@yahoo.com

Abstract

Dynabeads® M-280 Tosylactivated (untargeted magnetic beads) were evaluated to capture

Mycobacterium smegmatis and *Mycobacterium avium* subspecies *paratuberculosis* (MAP) from spiked

feces, milk, and urine. Untargeted magnetic beads added to the spiked samples were slightly mixed for 1

hour and separated in a magnetic rack for further detection. Beads recovered more *M. smegmatis* cells

from PBS suspension that the centrifugation method; these results were confirmed by the recovery of

96.31% of 1.68 x 10^4 CFU/mL viable *M. smegmatis* by beads and 0% by centrifugation. Likewise, the F57-

qPCR detection of MAP cells, after being recovered by beads and centrifugation, were different;

cycle threshold (Ct) was lower ($p<0.05$) for the detection of MAP cells recovered by beads than

centrifugation. Magnetic separation of MAP cells from milk, urine, and feces specimens were detected

by amplifying F57 and IS900 sequences. Ct values demonstrated that beads captured no less than 10^9

CFU/mL from feces and no less than 10^4 CFU/mL of MAP cells from milk and urine suspensions. Milk

proteins were denatured by Proteinase k before capturing MAP cells by magnetic beads. *M. smegmatis*

coupled to magnetic beads were infected by mycobacteriophage D29; plaque former units were

observed clearly from urine containing 2 x 10^5 and 2 x 10^3 CFU/mL *M. smegmatis* in 24 hours. The results

of this study encourage further effort to rule out the use of untargeted beads as a simple tool for

diagnosis of Johne's disease and other mycobacterial diseases such as tuberculosis.

Introduction

The genus *Mycobacterium* is grouped in *Mycobacterium tuberculosis* complex (MTBC) and non-tuberculosis mycobacteria (NTM) (1) and can cause disease in humans and animals (2,3). *M. tuberculosis* (MTB), the etiological agent of tuberculosis (TB) in humans cause millions of deaths mainly in developing countries (4), can be isolated from sputum (5), lavage fluid (6,7), gastric lavage (8,9), urine (10–12), and stool (13,14). *M. avium* subspecies *paratuberculosis* (MAP), a NTM that cause Johne's disease (JD) or paratuberculosis, has been isolated from intestinal samples of domestic (e.g., cattle, sheep, goats, and farmed deer), wildlife animals (e.g., rabbit, hare, fox, stoat, weasel, crow, rook); and human beings (15)(16)(17).

Definitive diagnostic of TB and JD is based on the isolation of MTB and MAP from human and animal's specimens on culture media. For that, the specimen, which can be contaminated with normal flora organisms, should be decontaminated and centrifuged (18). Classical decontamination methods to isolate MTB and MAP on culture method are based on the use *N*-acetyl-L-cysteine–sodium hydroxide (NALC-NaOH) (18) and Hexadecylpyridinium chloride (HPC) (19), respectively, and incorporate the concentration of the specimen at 3000 *x g* at 4°C for 15-20 min (18–20). Given that these chemical decontaminants can kill a proportion of mycobacteria present in the specimen also (19–22), and the availability of a refrigerated centrifuge with sealed safety cups and aerosol tight rotors may be challenging in laboratories from low- and middle-income (23), additional effort should be focused on the development of new and simple methods that enable to separate MTB and MAP without affecting their viability to growth on culture media.

Magnetic-separation methods, already established as routine methods to detect *Listeria monocytogenes* (24), *Salmonella* spp. (25), and *Escherichia coli* O157:H7 (26) (27) in food and veterinary clinical samples are promising alternates to selectively separate and concentrate MTB (28–32) and MAP (33,34). Magnetic bead characteristics (composition, size, concentration, and surface modification) and the nature of the coating ligand (antibodies, peptides, chemicals) are important to concentrate the target bacteria and increase the performance of further complementary detection methods such as culture, PCR, microscopy, antigen detection immunoassay, or phage assay (33,35,36).

Tosyl-activated magnetic beads (e.g., Dynabeads M-280 Tosylactivated) can bind covalently to primary amine and sulfhydryl groups in proteins and peptides (37), and as some surface proteins have been identified on mycobacteria (38–40). As per the characteristics of Tosyl-activated magnetic beads, we postulate that Tosyl-activated magnetic beads may interact with these surface proteins for capturing

M. smegmatis (Ms) and MAP cells in suspension. Because the use of untargeted beads to capture MAP

cells would diminish laboratory work, costs, and would increase its affordability in countries with a high

burden of TB and JD, this study determined the potential of Tosyl-activated Dynabeads M-280

(untargeted magnetic beads) to capture Ms and MAP from spiked urine, feces, and milk samples prior

detection by D29 phage and real-time PCR (qPCR).

Material and Methods

Bacterial cultures

MAP ATCC 19698 was cultured in Middlebrook 7H9 broth (BD Biosciences, San Jose, CA, USA)

supplemented with an oleic acid–albumin–dextrose–catalase mixture (BD Biosciences) and mycobactin J

(Allied Monitor, Fayette, MO, USA) at 37^{o}C with constant shaking at 200 rpm for 60 d. Ms mc2155 cells

were cultured under the same conditions but without mycobactin and for 7 d. *Escherichia coli* BL21

(DE3) was cultured in Luria–Bertani (LB) broth at 37^{o}C for 24 h with shaking. Stock Ms and *E. coli* cells

suspensions were adjusted to OD$_{600}$ measurement of 1. Then 4 serial 10-fold dilutions of the stock

suspension were made for CFU counting.

Magnetic beads assay

Tosyl-activated Dynabeads™ M-280 (Invitrogen)(37) were washed twice with sterile PBS (pH 7.4)

before adding into the suspension. Briefly, 21 µl of beads was transferred to one sterile micro centrifuge

tubes of 2 ml capacity (microtube) and washed twice with 1 ml of PBS (pH 7.4) for 1 min. Microtube was

placed on a magnet (Magna grip rack, Millipore) for removing supernatant. Washed beads were used for

magnetic beads assay. Microtubes containing PBS, milk, urine, and feces suspensions spiked with Ms or

MAP cells and beads were placed in a microtube rotator (Thermo Scientific™ Labquake™, Edmonton, AB,

Canada) at room temperature for 1 h. Then, the microtubes were placed on a magnetic rack for 1 min to

carefully remove the supernatant to another microtube. Magnetic beads coupled to Ms and MAP were

washed twice with 1 mL PBS before detecting by D29 phage and qPCR.

Recovery of viable *E. coli* and *M. smegmatis* by magnetic beads

M. smegmatis

From a stock of Ms suspension (adjusted to OD$_{600}$ ~1, microtube#1), 3 serial 10-fold dilutions

(microtube#2, microtube#3, and microtube#4) in PBS were made to evaluate the concentration of cells

by centrifugation and beads. Determination of viable Ms recovered by beads and centrifugation was

estimated as per the number of cells isolated from the supernatant due to the cells sedimented may

form clumps and underestimate the count of Ms isolated on LB plates. To make feasible the count of all

cells remained in the entire volume of supernatant, the experiments were optimized in microtube (2 mL

capacity). Two aliquots (1 mL) of each microtube were transferred to 2 microtubes for centrifugation

(control) and beads methods. Microtube#1 (control) was centrifuged at 3000 $x\,g$ at 4°C for 16 min, and

the total volume of supernatant was cultured on LB agar plates for counting cells that were not

sedimented. Another microtube#1 was assayed for magnetic capture, and the total volume of

supernatant was carefully removed and transferred to one plastic cuvette for measuring turbidity via OD

at 600 nm and for colony counts method on LB-agar plate. Same procedure, except measuring turbidity,

was applied for microtube#2 and microtube#3. From microtube#4, Ms cells recovered by beads and

centrifugation were resuspended in 1 mL, vortexed vigorously and plated on LB agar plates. Likewise,

the total supernatant volume was plated on LB agar for counting cells. The experiment was repeated 3

times.

E. coli. From a stock of *E. coli* suspension (adjusted to OD_{600} ~1), 2 aliquots (1 mL) were transferred to 2

microtubes to evaluate the concentration of cells by centrifugation (2000 $x\,g$, 4°C, 16 min) and capture

by magnetic beads. Recovery of cells by centrifugation (control) was done following the procedure

described for the microtube#1 with Ms. The experiment was repeated 3 times.

Detection of MAP cells by qPCR

This experiment was done to evaluate the recovery of MAP cells by beads and centrifugation

through the detection of DNA of the total, viable and non-viable, MAP cells sedimented. Efficiency of

beads and centrifugation to capture and sediment cells, respectively, was determined by using 2

referential MAP (5 x 10^7 CFU/ml and 5 x 10^5 CFU/mL) concentrations. Briefly, 2 serial 10-fold dilutions

were prepared from the stock (5 x 10^7 CFU/mL) of MAP suspension in PBS. Two aliquots (1 mL) of the

stock (5 x 10^7 CFU/ml of MAP cells) were transferred to 2 microtubes to evaluate the recovery of MAP

by centrifugation (control) and magnetic beads. Recovered MAP cells by beads and centrifugation (3000

$x\,g$, 4°C, 16 min) were analyzed by F57 qPCR. Same procedure was followed from the second serial

dilution containing (5 x 10^5 CFU/mL MAP). The experiment was repeated three times.

Magnetic bead capture MAP from milk, urine and feces

From milk and urine suspension. Three sterile microtubes containing 0.9 mL of PBS (pH 7.4), 0.9 mL of

urine from an apparently healthy human and 0.9 ml of skim milk (Dairyland®) were spiked with 0.1 mL of

10^8 CFU/mL MAP cells. Proteinase K (Pk) was added to the microtube containing milk and incubated at

70°C for 10 min before magnetic beads assay. MAP cells recovered using beads were analyzed by qPCR.

Same procedure was applicable for other microtubes containing less ($10^6, 10^5, 10^4$ CFU/mL) MAP cells.

The experiment was run in triplicate and repeated twice.

From feces suspension. Cow's feces of a known MAP shedding calf (41) was provided by the University

of Calgary. Feces suspension was prepared by mixing 3 g of apparently healthy cattle feces and 6 ml of

PBS in a 15-ml centrifuge microtube. After keeping the microtube in an upright position for 20 min, 0.9

mL of supernatant was transferred to another microtube and spiked with 0.1 mL of 10^8 CFU/mL MAP cell

suspended in PBS. One microtube with 0.9 mL PBS was spiked with 0.1 mL of 10^8 CFU/mL MAP cell. Both

microtubes were used for magnetic beads assay and analyzed by F57 qPCR. The experiment was run in

triplicate and repeated twice.

Quantification of number of MAP cells recovered by beads

MAP cells were counted qPCR based upon the single copy fragment F57 (42). Briefly, DNA

calibration curves were prepared for each experiment by preparing serial 10-fold dilutions of DNA

sample obtained MAP cell (adjusted to OD_{600} ~1). The initial DNA concentration of the first serial dilution

was measured through optical absorbance at 260 nm using a NanoDrop ND 1000 spectrophotometer

(NanoDrop Technologies Inc.). DNA molecule copy number (CFU/mL) was determined using the formula:

(ng DNA × 6,023 × 10^{23})/(4,829,781 × 10^9 × 660)(43). Number of F57 (CFU) detected in each MAP

sediment recovered by beads and control was determined by using the respective threshold cycle (Ct)

values.

DNA extraction

DNA extraction was performed using a GenoLyse® kit (Hain Lifescience, Nehren, BW, Germany)

as per manufacturer's instructions with some modifications. Briefly, MAP cells concentrated by

centrifugation or magnetic beads were resuspended in 100 µl of A-Lys, vortexed, centrifuged at 10,000 *x*

g for 1 min, and incubated for 5 min at 95°C in a water bath. Then, 100 µl of A-NB buffer was added to

the lysate, vortexed, and centrifuged at 10,000 *x g* for 5 min. Supernatant was transferred to sterile 1.5

ml microtubes. Concentration of the isolated DNA was determined by measuring the sample absorbance

at 260 nm using a NanoDrop ND-1000 spectrophotometer (NanoDrop Technologies Inc., Wilmington,

DE, USA).

Quantitative qPCR assay

This procedure was done as per Singh et al. (36). Briefly, qPCR amplification targeting IS900 and

F57 genes was done using Taqman specific probes and StepOnePlus Real Time PCR System (Applied

Biosystems). The 10 µL PCR mixture comprised 5 µL of TaqMan® Universal Master Mix II (Applied

Biosystems) with the uracil-N-glycosylase (UNG), 200 nM of each primer and 250 nM of fluorogenic

probe specific to IS900 and F57. From each template DNA sample, 4 µL was added in the reaction

mixture. qPCR conditions were as follows: 50°C for 2 min for UNG treatment, initial denaturation at 95°C

for 10 min, 40 cycles at 95°C for 15 sec, and at 60°C for 1 min. Each qPCR amplification targeting IS900

and F57 genes was run in triplicate with negative and positive controls. Obtained data were analyzed in

the form of threshold cycle (Ct) values.

Magnetic separation of *M. smegmatis* from urine for mycobacteriophage D29 infection

The experimental procedure was adjusted as per infection of MTB by D29 phage (44) and

doubling time (3 to 4 hours) of Ms (45). Two urine aliquots of 1 ml of 2.0×10^5 and 2.0×10^3 CFU/mL Ms

were transferred to 2 sterile microtubes. Ms captured by beads from urine were detected by D29 phage

(44,46) with some modifications. Briefly, Ms cells captured by beads were resuspended in 150 µL of LB

with 1 mM $CaCl_2$ (LB-S) and infected with 50 µL of 10^8 PFU/mL D29 phage at 37°C. Extracellular D29

phage was eliminated with 0.03 M ferrous ammonium sulfate (FAS) for 10 min; then 0.5 mL of LB-$CaCl_2$

was added to the microtube for stopping FAS virucide activity. The total volume suspension was mixed

with 4 mL of sterile molten agar (0.8%) and 1 mL of 10^8 CFU/mL Ms suspension before pouring onto

Petri dishes with LB agar. Petri dishes were incubated for 24 h at 37°C and the number of infection

plaques was counted. The experiment was repeated 3 times.

Statistical analyses

Number of F57 and IS900 copies, determined in triplicate per each sediment recovered by u-

TDbeads and control, were plotted and analyzed by the Kruskal-Wallis test (GraphPad Instat, version 8;

GraphPad Prism, La Jolla, CA). *P*-values < 0.05 were considered statistically significant.

Results

Capture of *E. coli* and *M. smegmatis* by magnetic beads

Magnetic beads captured Ms cells in PBS with greater capture efficiency compared to *E. coli* cells (*P* <0.05). This is evident by the lack of cells in supernatant of cuvette B2, unlike cuvette A2, where beads efficiently removed cells from suspension (Fig. 1). The number of viable Ms cells isolated from the supernatant of each suspension showed higher (> 90%) recovery of viable cells by beads than centrifugation. *E. coli* cells were not recovered by beads; approximately 94% of cells from suspension were isolated from the supernatant (Table 1). The effect of the centrifugation and magnetic beads methods on the viability of Ms cells was confirmed from the suspension 1.68×10^4 CFU/mL (Fig. 2). As per one experiment results, 17 CFU of Ms isolated from the supernatant and 337 CFU-beads recovered from the sediment of the suspension (1.68×10^4 CFU/mL), 337 CFU compressed 1326 CFU of Ms as consequence of the formation of clumps during the magnetic separation (Fig. 2). Thus, the magnetic beads recovered more (96%) viable Ms from the suspension (1.68×10^4 CFU/mL) than the centrifugation (0%). An electron microscopy image of the capture of Ms cells by beads is presented in Fig. 3.

Detection of MAP cells by qPCR

The initial concentration (100%) of MAP cells in the first and second PBS suspensions was estimated by detecting F57 sequence from all MAP cells collected by centrifugation. Since the centrifugation of PBS suspensions was done at 3,000 *x g* at 4°C for 16 min to concentrate 95% mycobacteria (18), we estimated that the initial concentration (100%) of MAP cells in the first and second PBS were 2.93×10^9 and 1.43×10^7 CFU/mL. Using F57 qPCR, there was a difference (p=0.04) in recovery of MAP cells: from the first suspension 2.8×10^9 CFU/mL (Ct = 17.89) were recovered by magnetic beads and 3.2×10^9 CFU/mL (Ct 17.68) by centrifugation, whereas 3.9×10^7 CFU/mL (Ct = 23.94) were recovered by magnetic beads and 1.4×10^7 CFU/mL (Ct = 25.43) by centrifugation ((p<0.0001); Table 2).

Milk, urine, and feces suspension. Ct values demonstrated that magnetic beads captured no less than 10^9 CFU/mL from feces and not less than 10^4 CFU/mL of MAP cells from milk and urine suspensions (Fig. 4). From milk suspensions the recovery rates, confirmed by F57 sequences, were 6.11% of 8.92×10^9 CFUL/mL, 5.96% of 6.25×10^7 CFU/mL, and 0.62% of 3.80×10^4 CFU/mL. Pk was incorporated in milk suspensions as our previous experiments (data not shown) showed that its activity increased the capture of MAP cells by beads. Recovery rates of MAP cells from urine suspensions using F57 was

30.93% of 8.92×10^9, 43.25% of 6.25×10^7, 44.56% of 6.11×10^6 CFU/mL, and 8.60% of 4.92×10^5

CFU/mL. IS900, unlike F57, was more sensitive to detect lowest (10^4 CFU/mL) MAP cells concentration

from milk and urine (Fig. 4 and Table 3). From feces, 0.66% of 8.92×10^9 CFU/mL MAP cells were

captured and detected by beads and F57-qPCR (IS900 was not done). Other feces suspensions

containing lower concentrations were not detected.

Magnetic separation of *M. smegmatis* from urine for mycobacteriophage D29 infection

Magnetic beads-D29 phage detected Ms spiked in urine in 24 h (Fig. 5). Although diminished

number of plaques were observed from urine spiked with 2×10^5 CFU/mL and 2×10^3 CFU/mL of Ms

(Table 4), the number of PFU detected from urine suspension with 2×10^5 CFU/mL of Ms clearly prove

the presence of Ms in the suspension in 24 h.

Discussion

Untargeted Tosyl-activated Dynabeads™ M-280 recovered more (>90%) viable Ms from spiked PBS

suspension with 10^7, 10^6, 10^5 and 10^4 CFU/mL Ms. Capture efficiency was estimated by counting

unbound Ms cells (supernatant) that were not captured nor sedimented by beads. Counting cells

sedimented by beads, and even, after resuspending them in 1 mL before plating (47), could

underestimate the real number of cells recovered by beads due to formation of Ms and MAP clumps

during the sedimentation process. With regards to the effect of magnetic beads and centrifugation on

the viability of Ms cells, untargeted magnetic beads did not affect the viability of Ms to growth on LB

agar plate and to be infected by D29 phage. Contrary, centrifugation had a negative effect on the

viability of Ms cells sedimented from all four 4 suspensions spiked with 4 different Ms concentrations.

Moreover, the qPCR detection of MAP cells, after sedimentation by centrifugation and magnetic beads,

indicate that the use of magnetic beads would be the best method to precipitate cells from PBS

suspension. Although the centrifugation method (3000 *x g* at 4°C for 16 min) has been established as a

referential method to recovery 95% of cells from suspension (18,48), we agree with others (49–51) that

centrifugation may not be an appropriate method to concentrate specimens containing small numbers

of organisms. The inefficient centrifuge method may be related to the growth rate of bacteria; low

recovery of MAP cells from a stationary culture is expected as the majority of cells have a lower buoyant

density due to the presence of lipid bodies (52). Likewise, as others have reported the effect of

centrifugation process for other bacteria (53), the low number of Ms growth on LB agar may be

explained by the damage of some bacterial cells, and even DNA, during the centrifugation.

Since detection of MAP cells from milk improved after treating with Pk, we postulate that Tosyl

group can be blocked by milk proteins and, as consequence, affect the capture of MAP cells. The effect

of Pk on capture of MAP cells from milk was not observed in urine nor feces, perhaps, due to the higher

protein concentration (1.87%) of milk (54) compared to urine (55) and feces. With regards to the

magnetic separation of MAP cells from urine and feces, we postulate that these differences in the

detection sensitivity of qPCR to detect MAP cells from milk, urine and feces can be explained by its own

inhibitors, and the number of copies of F57 (1 copy) (56,57) and IS900 (14-20 copies)(57,58) in the

genome of MAP. Urea amine group, and other inhibitors such as polysaccharides originated from

vegetable material in the diet (59), gut flora, bile salts, and glucolipids (60), may react with Tosyl groups

and decrease the capture of cells by beads. Likewise, these inhibitors of each specimen could be

transferred with the beads coupled to MAP cells and reduce the amplification efficiency of F57 and

IS900 sequences.

Magnetic beads detected viable Ms from urine suspension containing at least 10^3 CFU/mL by

D29 phage. Since that our experiments have shown that the centrifugation process can affect the

viability of all Ms cells from a suspension with 10^6 CFU/mL bacilli, and considering that the

decontamination of sample may damage the mycobacteria and reduce the sensitivity of phage-based

assays for detecting mycobacteria cells (61), our method magnetic bead-based D29 phage could also be

useful to detect low amount of MAP cells from cow's milk, and even MTB from urine samples, in 24 h.

However, since the number of plaques (02 PFU) from urine suspension spiked with 2 x 10^3 CFU/mL of Ms

may raise questionable results, the addition of an incubation step of recovered Ms at 37°C for 24 hours,

before adding phages, could be necessary to increase the number of cells to be infected by phages, and

as consequence, the increase of plaques units on the indicator plate. Despite that the final

concentration of phages (2.5 x 10^7 PFU/mL) in the PBS and urine suspensions should ensure an infection

rate of 1% (maximum infection rate) (44), the presence of beads in PBS and, as well, the urine urea

would explain the low infection rate (<1%) of cells by phage. Beads may interfere the interaction

between cells and phages while the urine urea (amino group) may interact with Tosyl groups and reduce

the number of cells captured by beads.

Another variable that positively contributed to the capture of mycobacteria by beads was

related to the technical skills to take aliquots of beads in stock suspension, remove the supernatant after

capturing beads coupled to mycobacteria and washing beads. Before taking an aliquot of the stock

beads suspension, it was homogenized by pipetting 3 – 4 times carefully. The supernatant was removed

without taking out the microtube from the magnetic rack to avoid removing beads coupled to

mycobacteria. PBS washing process was done carefully despite that, due to the characteristic of beads

for interacting with proteins, there would be a strong interaction between beads and mycobacteria

surface proteins. PBS was added slowly and carefully through the inner microtube's wall and not directly

to the beads. The microtube should be slightly inverted 5 times to ensure that all beads distributed

homogenously and are not making visible clumps that can carry on some debris present in the sample.

The use of microcentrifuge tube of 2 mL with attached flat top cap ensures a) homogenous mix of

suspension for increasing the attraction between beads and cells, b) removal of supernatant while the

tube is placed in the magnetic rack, and c) addition of PBS to the tube for washing cells coupled to

beads.

Notwithstanding others (62) demonstrated that the use of Tosyl-activated Dynabeads™ M-280

coupled to Gp10, a recombinant lysin from the mycobacteriophage L5, in combination with qPCR, the

study does not show data about the capture efficiency of uncovered magnetic beads. Another study (47)

showed that other types of magnetic beads coupled to some peptides and antibodies, and

complemented with phages, detected around 0.03% and 0.003% of 10^3 to 10^4 CFU/mL MAP cells from

milk, respectively. However, unlike the method of this study, which recovered MAP cells directly from

milk, the MAP cells were sedimented from milk by centrifugation and then were transferred to one

microtube containing 7H9-OADC media for magnetic separation of MAP cells. Moreover, the authors

(47) showed that other types of uncoated beads recovered a moderate percentage (around 10%) of

nontarget mycobacteria.

A limitation of the study was that MAP cells recovered by beads were not cultured. Although the

2 log10 lower counts of MAP cells between theoretical concentration determined at OD600 and F57-

qPCR can be explained by the clumps present in the suspension and not detected at OD600, and our

results are in line with another study (42) that have also reported a difference of two log10 counts

between colony counts on culture media and F57-qPCR, counting these recovered cells before qPCR

could have been useful to infer the presence of death and alive cells recovered by beads and

centrifugation. Other variable that could not be controlled in the study were the effect of calcium

chelating agent in milk. Removing calcium from milk could reduce the formation of MAP cells clumps

and keep a homogenous distribution of cells in the suspension to be captured by beads. This study did

not evaluate the effect of less amount of feces (1 g (19,63), 2g (64,65)) on the capture of MAP cells; even

though less amount of feces can provide less amount of inhibitors, the reduction of feces would reduce

the sensitivity to recovery MAP cells. The evaluation of these variables in further studies would define

the potential to use untargeted magnetic beads to capture MAP cells from milk and feces.

The results of this study encourage further effort to evaluate the utility of Dynabeads® M-280

Tosylactivated, without specific ligand, in real animals' samples with a presumptive diagnosis

of paratuberculosis. Likewise, the use of these beads, in combination with a microbiological (e.g., DS6A

phage, culture media, colorimetric method) or a molecular test (e.g., HAIN-MTBDRplus, HAIN-MTBDRsl,

GeneXpert), would have high potential to be used as a simple and rapid tool for diagnosis of TB disease.

References

1. Baliga S, Murphy C, Sharon L, Shenoy S, Biranthabail D, Weltman H, et al. Rapid method for

detecting and differentiating Mycobacterium tuberculosis complex and non-tuberculous

mycobacteria in sputum by fluorescence in situ hybridization with DNA probes. Int J Infect Dis.

2018;75:1–7.

2. Palmer M V., Welsh MD, Hostetter JM. Mycobacterial diseases of animals. Vet Med Int.

2011;2011:292469.

3. Li H, Turhan V, Chokhani L, Stratton CW, Dunbar SA, Tang YW. Identification and differentiation

of clinically relevant Mycobacterium species directly from acid-fast Bacillus-positive culture

broth. J Clin Microbiol. 2009;47:3814–20.

4. WHO. Global tuberculosis report 2020. 2020. Available from:

https://www.who.int/publications/i/item/9789240013131

5. Narvaiz de Kantor, Isabel, Kim, Sang Jae, Frieden, Thomas R, Laszlo, Adalbert, Luelmo F et al.

Laboratory services in tuberculosis control culture. Part III: Culture. World Health Organization.

1998. Available from:

https://apps.who.int/iris/bitstream/handle/10665/65942/WHO_TB_98.258_%28part3%29.pdf?

sequence=3&isAllowed=y.

6. Nikbakhsh N, Bayani M, Siadati S. The value of bronchoalveolar lavage in the diagnosis of

sputum smear-negative pulmonary tuberculosis. Iran J Pathol. 2015;10:35–40.

7. Baughman RP, Dohn MN, Loudon RG, Frame PT. Bronchoscopy with bronchoalveolar lavage in

tuberculosis and fungal infections. Chest. 1991;99:92–7.

8. DiNardo AR, Detjen A, Ustero P, Ngo K, Bacha J, Mandalakas AM. Culture is an imperfect and

heterogeneous reference standard in pediatric tuberculosis. Tuberculosis. 2016;101:S105–8.

9.   Ruiz Jiménez M, Guillén Martín S, Prieto Tato LM, Cacho Calvo JB, Álvarez García A, Soto Sánchez
B, et al. Induced sputum versus gastric lavage for the diagnosis of pulmonary tuberculosis in
children. BMC Infect Dis. 2013;13:222.

10. Gopinath K, Singh S. Urine as an adjunct specimen for the diagnosis of active pulmonary
tuberculosis. Int J Infect Dis. 2009;13:374–9.

11. Peter J, Green C, Hoelscher M, Mwaba P, Zumla A, Dheda K. Urine for the diagnosis of
tuberculosis: Current approaches, clinical applicability, and new developments. Curr Opin Pulm
Med. 2010;16:262-70.

12. Turbawaty DK, Sugianli AK, Soeroto AY, Setiabudiawan B, Parwati I. Comparison of the
performance of urinary Mycobacterium tuberculosis antigens cocktail (ESAT6, CFP10, and
MPT64) with culture and microscopy in pulmonary tuberculosis patients. Int J Microbiol.
2017;2017:3259329.

13. Walters E, Demers AM, Van Der Zalm MM, Whitelaw A, Palmer M, Bosch C, et al. Stool culture
for diagnosis of pulmonary tuberculosis in children. J Clin Microbiol. 2017;55:3355–65.

14. Oramasionwu GE, Heilig CM, Udomsantisuk N, Kimerling ME, Eng B, Nguyen HD, et al. The utility
of stool cultures for diagnosing tuberculosis in people living with the human immunodeficiency
virus. Int J Tuberc Lung Dis. 2013;17:1023–8.

15. Harris NB, Barletta RG. Mycobacterium avium subsp. paratuberculosis in veterinary medicine.
Clin Microbiol Rev. 2001;14:489–512.

16. de Juan L, Alvarez J, Romero B, Bezos J, Castellanos E, Aranaz A, et al. Comparison of four
different culture media for isolation and growth of type II and type I/III Mycobacterium avium
subsp. paratuberculosis strains isolated from cattle and goats. Appl Environ Microbiol.
2006;72:5927–32.

17. Rangel SJ, Paré J, Doré E, Arango JC, Côté G, Buczinski S, et al. A systematic review of risk factors
associated with the introduction of Mycobacterium avium spp. paratuberculosis (MAP) into
dairy herds. Can Vet J. 2015;56:169-77.

18. Kent P T KGP. Public health mycobacteriology: a guide for the level III laboratory. US Department
of Public Health and Human Services, Public Health Service, Centers for Disease Control, Atlanta,
GA,. 1985. p. 31-67.

19. Sattar A, Zakaria Z, Abu J, Aziz SA, Gabriel RP. Evaluation of six decontamination procedures for
isolation of Mycobacterium avium complex from avian feces. PLoS One. 2018;13:e0202034.

20. Ganoza CA, Ricaldi JN, Chauca J, Rojas G, Munayco C, Agapito J, et al. Novel hypertonic saline-
sodium hydroxide (HS-SH) method for decontamination and concentration of sputum samples
for Mycobacterium tuberculosis microscopy and culture. J Med Microbiol. 2008;57:1094–8.

21. Chatterjee M, Bhattacharya S, Karak K, Dastidar SG. Effects of different methods of
decontamination for successful cultivation of Mycobacterium tuberculosis. Indian J Med Res.
2013;138:541–8.

22. Addise D, Bitew A, Yaregal Z, Yenew B, Mollalign H, Diriba G, et al. Effect of 1.5% sodium
hydroxide final concentration on recovery rate of mycobacterial species and decontamination of
other bacterial and fungal contaminants on sputum. Ethiop J public Heal Nutr. 2016;1:57.

23. Parsons LM, Somoskövi Á, Gutierrez C, Lee E, Paramasivan CN, Abimiku A, et al. Laboratory
diagnosis of tuberculosis in resource-poor countries: Challenges and opportunities. Clin
Microbiol Rev. 2011;24:314-50.

24. Hsih H-Y, Tsen H-Y. Combination of immunomagnetic separation and polymerase chain reaction
for the simultaneous detection of Listeria monocytogenes and Salmonella spp. in food samples.
J Food Prot. 2001;64:1744–50.

25. Rijpens N, Herman L, Vereecken F, Jannes G, De Smedt J, De Zutter L. Rapid detection of stressed
Salmonella spp. in dairy and egg products using immunomagnetic separation and PCR. Int J Food
Microbiol. 1999;46:37–44.

26. Karch H, Janetzki-Mittmann C, Aleksic S, Datz M. Isolation of enterohemorrhagic Escherichia coli
O157 strains from patients with hemolytic-uremic syndrome by using immunomagnetic
separation, DNA-based methods, and direct culture. J Clin Microbiol. 1996;34:516–9.

27. Possé B, De Zutter L, Heyndrickx M, Herman L. Quantitative isolation efficiency of O26, O103,
O111, O145 and O157 STEC serotypes from artificially contaminated food and cattle faeces
samples using a new isolation protocol. J Appl Microbiol. 2008;105:227–35.

28. Ghodbane R, Drancourt M. Magnetic bead protocol for culturing mycobacterium tuberculosis
from sputum specimens. J Clin Microbiol. 2013;51:1578–9.

29. Intorasoot S, Tharinjaroen CS, Phunpae P, Butr-Indr B, Anukool U, Intachai K, et al. Novel
potential diagnostic test for Mycobacterium tuberculosis complex using combined
immunomagnetic separation (IMS) and PCR-CTPP. J Appl Microbiol. 2016;121:528–38.

30. Yoshimatsu S, Kato-Matsumaru T, Aono A, Chikamatsu K, Yamada H, Mitarai S. Factors
contribute to efficiency of specimen concentration of Mycobacterium tuberculosis by
centrifugation and magnetic beads. Int J Mycobacteriology. 2015;4:245–9.

31. Albert H, Ademun PJ, Lukyamuzi G, Nyesiga B, Manabe Y, Joloba M, et al. Feasibility of magnetic
bead technology for concentration of mycobacteria in sputum prior to fluorescence microscopy.
BMC Infect Dis. 2011;11:1–9.

32. Saad J, Loukil A, Drancourt M. Bead-captured Mycobacterium tuberculosis for next-generation
sequencing diagnosis of uncultured tuberculosis. Eur J Clin Microbiol Infect Dis. 2020;39:205-7.

33. Foddai A, Elliott CT, Grant IR. Maximizing capture efficiency and specificity of magnetic
separation for Mycobacterium avium subsp. paratuberculosis cells. Appl Environ Microbiol.
2010;76:7550–8.

34. Husakova M, Dziedzinska R, Slana I. Magnetic separation methods for the detection of
Mycobacterium avium subsp. paratuberculosis in various types of matrices: A review. Biomed
Res Int. 2017;2017:5869854.

35. Ekong MO. A review on laboratory diagnosis of Mycobacterium tuberculosis. Asian J Res Infect
Dis. 2019;2:1–7.

36. Singh U, Arutyunov D, Basu U, Santos Seckler H Dos, Szymanski CM, Evoy S. Mycobacteriophage
lysin-mediated capture of cells for the PCR detection of Mycobacterium avium subspecies
paratuberculosis. Anal Methods. 2014;6:5682–9.

37. Invitrogen. Magnetic Separation Technology Surface-activated Dynabeads ® A wide product
portfolio for flexible molecular separations [Internet]. [cited 2019 Apr 26]. Available from:
https://assets.thermofisher.com/TFS-Assets/LSG/brochures/Surface_Activated_Dynabeads.PDF

38. He Z, De Buck J. Localization of proteins in the cell wall of Mycobacterium avium subsp.
paratuberculosis K10 by proteomic analysis. Proteome Sci. 2010;8:21.

39. Beatty WL, Russell DG. Identification of mycobacterial surface proteins released into subcellular
compartments of infected macrophages. Infect Immun. 2000;68:6997–7002.

40. He Z, De Buck J. Cell wall proteome analysis of Mycobacterium smegmatis strain MC2 155. BMC
Microbiol. 2010;10:121.

41. Mortier RAR, Barkema HW, Bystrom JM, Illanes O, Orsel K, Wolf R, et al. Evaluation of age-
dependent susceptibility in calves infected with two doses of Mycobacterium avium subspecies
paratuberculosis using pathology and tissue culture. Vet Res. 2013;44:1–9.

42. Kralik P, Beran V, Pavlik I. Enumeration of Mycobacterium avium subsp. paratuberculosis by
quantitative real-time PCR, culture on solid media and optical densitometry. BMC Res Notes.
2012;5:114.

43. Sevilla IA, Molina E, Elguezabal N, Pérez V, Garrido JM, Juste RA. Detection of mycobacteria,

Mycobacterium avium subspecies, and Mycobacterium tuberculosis complex by a novel

tetraplex real-time PCR assay. J Clin Microbiol. 2015;53:930–40.

44. McNerney R, Kambashi BS, Kinkese J, Tembwe R, Godfrey-Faussett P. Development of a

bacteriophage phage replication assay for diagnosis of pulmonary tuberculosis. J Clin Microbiol.

2004;42:2115–20.

45. Klann AG, Belanger AE, Abanes-de Mello A, Lee JY, Hatfull GF. Characterization of the dnaG locus

in Mycobacterium smegmatis reveals linkage of DNA replication and cell division. J Bacteriol.

1998;180:65–72.

46. Chauca JA, Palomino JC, Guerra H. Evaluation of rifampicin and isoniazid susceptibility testing of

Mycobacterium tuberculosis by a mycobacteriophage D29-based assay. J Med Microbiol.

2007;56:360–4.

47. Foddai A, Elliott CT, Grant IR. Maximizing capture efficiency and specificity of magnetic

separation for Mycobacterium avium subsp. paratuberculosis. Cells. 2010;76:7550–8.

48. Ratnam S, March SB. Effect of relative centrifugal force and centrifugation time on

sedimentation of mycobacteria in clinical specimens. J Clin Microbiol. 1986;23:582–5.

49. WAYNE LG. The use of millipore filters in clinical laboratories. Am J Clin Pathol. 1957;28:565–7.

50. Smithwick RW, Stratigos CB. Acid-fast microscopy on polycarbonate membrane filter sputum

sediments. J Clin Microbiol. 1981;13:1109–13.

51. Ratnam S, March SB. Effect of relative centrifugal force and centrifugation time on

sedimentation of mycobacteria in clinical specimens. J Clin Microbiol. 1986;23:582–5.

52. Kennedy JA, Baron VO, Hammond RJH, Sloan DJ, Gillespie SH. Centrifugation and

decontamination procedures selectively impair recovery of important populations in

Mycobacterium smegmatis. Tuberculosis. 2018;112:79–82.

53. Peterson BW, Sharma PK, van der Mei HC, Busscher HJ. Bacterial cell surface damage due to

centrifugal compaction. Appl Environ Microbiol. 2012;78:120–5.

54. Górska-Warsewicz H, Rejman K, Laskowski W, Czeczotko M. Milk and dairy products and their

nutritional contribution to the average polish diet. Nutrients. 2019;11:1771.

55. Black CE, Berg RL, Urquhart AC. 24-Hour urinary calcium in primary hyperparathyroidism. Clin

Med Res. 2013;11:219–25.

56. Tasara T, Stephan R. Development of an F57 Sequence-based real-time PCR assay for detection

of Mycobacterium avium subsp. paratuberculosis in milk. Appl Environ Microbiol.

2005;71:5957–68.

57. Donaghy JA, Johnston J, Rowe MT. Detection of Mycobacterium avium ssp. paratuberculosis in

cheese, milk powder and milk using IS900 and F57-based qPCR assays. J Appl Microbiol.

2011;110:479–89.

58. Cunha MV., Rosalino LM, Leão C, Bandeira V, Fonseca C, Botelho A, et al. Ecological drivers of

Mycobacterium avium subsp. paratuberculosis detection in mongoose (Herpestes ichneumon)

using IS900 as proxy. Sci Rep. 2020;10:1–14.

59. Monteiro L, Bonnemaison D, Vekris A, Petry KG, Bonnet J, Vidal R, et al. Complex

polysaccharides as PCR inhibitors in feces: Helicobacter pylori model. J Clin Microbiol.

1997;35:995–8.

60. Schrader C, Schielke A, Ellerbroek L, Johne R. PCR inhibitors - occurrence, properties and

removal J Appl Microbiol. 2012;113:1014-26.

61. Kalantri S, Pai M, Pascopella L, Riley L, Reingold A. Bacteriophage- based tests for the detection

of Mycobacterium tuberculosis in clinical specimens: A systematic review and meta-analysis.

BMC Infect Dis. 2005;5:59.

62. Singh U, Arutyunov D, Basu U, Dos H, Seckler S, Szymanski CM, et al. Mycobacteriophage lysin-

mediated capture of cells for the PCR detection of Mycobacterium avium subspecies

paratuberculosis. Anal Meth 2014; 6:5682-9.

63. Zhang MZ, Zhang S. An efficient DNA extraction method for polymerase chain reaction-based

detection of Mycobacterium avium subspecies paratuberculosis in bovine fecal samples. J Vet

Diagn Invest. 2011;23:41–8.

64. Keller SM, Stephan R, Kuenzler R, Meylan M, Wittenbrink MM. Comparison of fecal culture and

F57 real-time polymerase chain reaction for the detection of Mycobacterium avium subspecies

paratuberculosis in Swiss cattle herds with a history of paratuberculosis. Acta Vet Scand.

2014;56:68.

65. Bauman CA, Jones-Bitton A, Jansen J, Kelton D, Menzies P. Evaluation of fecal culture and fecal

RT-PCR to detect Mycobacterium avium ssp. paratuberculosis fecal shedding in dairy goats and

dairy sheep using latent class Bayesian modeling. BMC Vet Res. 2016;12:212.

Table 1: *Escherichia coli* and *Mycobacterium smegmatis* remained in supernatant after centrifugation and magnetic beads methods.

Method	% E. coli (mean, ±SD) in supernatant from (CFU/mL)	% M. smegmatis (mean, ±SD) in supernatant from (CFU/mL)			
	$1{,}8 \times 10^7$	1.68×10^7	1.68×10^6	1.68×10^5	1.68×10^4
Centrifugation	0	0.66 $(1.12 \times 10^5, \pm 7.64 \times 10^3)$	0	0	0
Magnetic beads	94.63 $(1.7 \times 10^7, \pm 5.75 \times 10^4)$	4.04 $(6.79 \times 10^5, \pm 2.16 \times 10^4)$	3.07 $(5.17 \times 10^4, \pm 2.96 \times 10^3)$	6.71 $(1.13 \times 10^4, \pm 9.57 \times 10^2)$	3.69 $(6.21 \times 10^2, \pm 1.22 \times 10^2)$

Table 2: Recovery and detection of MAP cells by magnetic beads and F57-qPCR

PBS suspension	OD$_{600}$, 1.0 (theoretical)[a] CFU/mL ($\pm$SD)	Initial MAP cells concentration CFU/mL[b]	MAP cells recovered by centrifugation* and quantified by F57-qPCR		MAP cells recovered by beads and quantified by F57-qPCR	
			CFU/mL ($\pm$SD)	Ct	CFU/mL ($\pm$SD)	Ct
First	3.17×10^7 ($\pm 2.02 \times 10^7$)	2.93×10^9	2.8×10^9 ($\pm 4.1 \times 10^8$)	17.89[c]	3.2×10^9 ($\pm 4.5 \times 10^8$)	17.68[c]
Second	3.17×10^5 ($\pm 2.02 \times 10^5$)	1.43×10^7	1.4×10^7 ($\pm 1.8 \times 10^6$)	25.43[d]	3.9×10^7 ($\pm 5.5 \times 10^6$)	23.94[d]

[a] Approximation of CFU in each suspension was assessed quickly by OD600 measurements.

[b] Initial concentration (100%) of MAP cells in the first and second PBS suspensions was estimated by detecting F57 sequence from MAP cells recovered (in theory 95%) by centrifugation*.

[c] F57-qPCR showed that there was a statistically significantly difference (p=0.04) in the recovery of MAP cells by magnetic beads and centrifugation.

[d] F57-qPCR showed that there was a statistically significantly difference (p<0.0001) in the recovery of MAP cells by magnetic beads and centrifugation.

Table 3: Recovery rate of *M. avium paratuberculosis* from milk, urine, and feces sample

Initial concentration (CFU/mL)	Milk				Urine				Feces		
	F57-qPCR			IS900-qPCR	F57-qPCR			IS900-qPCR	F57-qPCR		
	CFU/mL (mean, ±SD)	Recovery (%)	Ct (mean; ±SD)	Ct (mean; SD)	CFU/mL (mean, ±SD)	Recovery (%)	Ct (mean; ±SD)	Ct (mean; ±SD)	CFU/mL (mean, ±SD)	Recovery (%)	Ct (mean; ±SD)
8.92×10^9	5.45×10^8; $\pm 8.31 \times 10^7$	6.11	20.45; ± 0.22	16.30; ± 0.18	2.76×10^9; $\pm 4.46 \times 10^8$	30.93	18.12; ± 0.23	13.84; ± 0.08	$5.93 \times 10^7 \pm 1.54 \times 10^7$	0.66	28.27; ± 0.35
6.25×10^7	3.73×10^6; $\pm 1.01 \times 10^6$	5.96	27.61; ± 0.38	23.89; ± 0.13	2.70×10^7; $\pm 6.85 \times 10^6$	43.25	24.77; ± 0.34	20.53; ± 0.06	UND	UND	UND
6.11×10^6	3.80×10^4; $\pm 2.44 \times 10^4$	0.62	36.44; ± 1.11	32.14; ± 1.74	2.72×10^6; $\pm 2.81 \times 10^6$	44.56	31.72; ± 2.71	26.63; ± 2.93	UND	UND	UND
4.92×10^5	UND	NA	NA	35.45; ± 0.78	4.23×10^4; $\pm 2.70 \times 10^4$	8.60	36.24; ± 0.82	32.42; ± 2.01	UND	UND	UND
5.19×10^4	UND	NA	NA	36.59; NA	UND	NA	NA	35.74; ± 1.18	UND	UND	UND

Table 4: Detection of *M. smegmatis* from PBS and urine by beads and D29 phage

	PFU (mean* ±SD) detected from spiked M. smegmatis (CFU/mL)	
	2×10^5 CFU/mL	2×10^3 CFU/mL
PBS (control)	259.33 ±34.08	12.33 ±2.52
Urine	56 ±8.54	2 ±1

*Mean of three experiments.

FIG 1 Capture of *E. coli* and *M. smegmatis* cells in PBS suspension by magnetic beads.

A. Capturing *E. coli* in PBS by beads. Cuvette A1: Concentration (1,8 x 10^7 CFU/ml) of *E. coli* before capture; cuvette A2: Concentration (1,7 x 10^7 CFU/ml) of *E. coli* suspension after beads capture. Microtube A3: Magnetic beads bound to *E. coli* cells and separated in a magnetic rack; B. Capturing *M. smegmatis* in PBS by beads. Cuvette B1: Concentration (1,68 x 10^7 CFU/ml) of *M. smegmatis* suspension before beads capture; cuvette B2: Concentration (6.56 x 10^5 CFU/ml) of *M. smegmatis* suspension after beads capture; microtube B3: Magnetic beads bound to *M. smegmatis* cells and separated in a magnetic rack.

FIG 2 *M. smegmatis* cells recovered by magnetic beads and centrifugation. From a suspension of 1.68 x 10^4 CFU/mL of *M. smegmatis*, A and B show cells captured (precipitated) and not captured (supernatant) by beads, respectively. As per the recovery (3.69%) of *M. smegmatis* cells from the supernatant, the estimated number of cells recovered was 96.31%. C and D show that there were not cells recovered nor remained in the supernatant after centrifugation process.

FIG 3 Electron microscopy of *M. smegmatis* cell captured by magnetic beads. Capturing *M. smegmatis* cell in PBS by magnetic beads.

FIG 4 Detection of the F57 sequence by qPCR from MAP cells recovered by magnetic beads.

Ct values showed that magnetic beads captured MAP cells from milk, urine, and feces. From feces, milk, and urine, F57 sequences were detected up to 8.92×10^9, 6.11×10^6, and 4.92×10^5 CFU/mL, respectively. IS900, unlike F57, showed better sensitivity to detect lowest (10^4 CFU/mL) MAP cells concentration from milk and urine. Other feces suspensions containing lower concentrations were not detected by qPCR. Detection of IS sequence from feces sample was not done.

FIG 5 Detection of *M. smegmatis* from urine suspensions by mycobacteriophage D29.

Detection of *M. smegmatis* captured from PBS suspensions (A and B) and urine suspensions (C and D) by mycobacteriophage D29 in 24 hours.