Genetic basis and dual adaptive role of floral pigmentation in sunflowers

Marco Todesco[1]*, Natalia Bercovich[1], Amy Kim[1], Ivana Imerovski[1], Gregory L. Owens[1,2],

Óscar Dorado Ruiz[1], Srinidhi V. Holalu[3], Lufiani L. Madilao[4], Mojtaba Jahani[1], Jean-Sébastien

Légaré[1], Benjamin K. Blackman[3], Loren H. Rieseberg[1]*

[1] Department of Botany and Biodiversity Research Centre, University of British Columbia,

Vancouver, British Columbia, Canada.

[2] Department of Biology, University of Victoria, Victoria, British Columbia, Canada.

[3] Department of Plant and Microbial Biology, University of California, Berkeley, Berkeley,

California, USA.

[4]Michael Smith Laboratory and Wine Research Centre, University of British Columbia,

Vancouver, British Columbia, Canada.

*Corresponding authors: mtodesco@biodiversity.ubc.ca (M.T.), lriesebe@mail.ubc.ca (L.H.R.).

Abstract

Variation in floral displays, both between and within species, has been long known to be shaped

by the mutualistic interactions that plants establish with their pollinators. However, increasing

evidence suggests that abiotic selection pressures influence floral diversity as well. Here we

analyze the genetic and environmental factors that underlie patterns of floral pigmentation in

wild sunflowers. While sunflower inflorescences appear invariably yellow to the human eye,

they display extreme diversity for patterns of ultraviolet pigmentation, which are visible to most

pollinators. We show that this diversity is largely controlled by cis-regulatory variation at a

single MYB transcription factor, HaMYB111, through accumulation of UV-absorbing flavonol

glycosides. As expected, different patterns of ultraviolet pigments in flowers have a strong effect

on pollinator preferences. However, variation for floral ultraviolet patterns is also associated with

environmental variables, especially relative humidity, across populations of wild sunflowers.

Larger ultraviolet patterns, which are found in drier environments, limit transpiration, therefore

reducing water loss. The dual role of floral UV patterns in pollination attraction and abiotic

responses reveals the complex adaptive balance underlying the evolution of floral traits.

Introduction

The diversity in colour and colour patterns found in flowers is one of the most extraordinary

examples of adaptive variation in the plant world. As remarkable as the variation that we can

observe is, even more of it lays just outside our perception. Many species accumulate pigments

that absorb ultraviolet (UV) radiation in their flowers; while these patterns are invisible to the

human eye, they can be perceived by pollinators, most of which can see in the near UV (Chittka

et al., 1994, Tovée, 1995). UV patterns have been shown to increase floral visibility, and to have

a major influence on pollinator visitation and preference (Brock et al., 2016, Horth et al., 2014,

Rae and Vamosi, 2013, Sheehan et al., 2016). Besides their role in pollinator attraction, patterns

of UV-absorbing pigments in flowers have also been linked, directly or indirectly, to other biotic

and abiotic factors (Gronquist et al., 2001, Koski and Ashman, 2015, Koski and Ashman, 2016).

Sunflowers have come to enjoy an iconic status in popular culture (as testified by the, arguably

dubious, honour of being one of the only five flower species with a dedicated emoji

(Unicode.org, 2020)). This is despite being apparently largely immune from the aforementioned

diversification for flower colour; all 50 species of wild sunflowers have ligules (the enlarged

modified petals of the outermost whorl of florets in the sunflower inflorescences) that appear of

the same bright yellow colour to the human eye. However, ligules also accumulate UV-

absorbing pigments at their base, while their tip reflects UV radiation (Harborne and Smith,

1978, Wojtaszek and Maier, 2014). Across the whole inflorescence, this results in a bullseye

pattern with an external UV-reflecting ring and an internal UV-absorbing ring. Besides their

well-described role in pollinator attraction, UV bullseyes have been proposed to act as nectar

guides, helping pollinators orient towards nectar rewards once they land on the petal, although

recent experiments have challenged this hypothesis (Koski and Ashman, 2014). Considerable

variation in the size of UV bullseye patterns has been observed between and within plant species

(Koski and Ashman, 2013, Koski and Ashman, 2016); however, little is known about the

ecological factors that drive this variation, or the genetic determinants that control it.

Results and Discussion

Floral UV patterns in wild sunflowers

A preliminary screening of 15 species of wild sunflowers, as well as cultivated sunflower,

suggested that UV bullseye patterns are common across sunflower species (Figure 1 – figure

supplement 1). We also observed substantial within-species variation for the size of UV floral

patterns. Variation for UV bullseye size was previously reported in the silverleaf sunflower

Helianthus argophyllus; however, genetic mapping resolution was insufficient to identify

individual causal genes (Moyers et al., 2017). To better understand the function and genetic

regulation of variation for floral UV pigmentation, we focused on two widespread species of

annual sunflowers, *H. annuus* (the progenitor of the domesticated sunflower) and *H. petiolaris*;

both have broad distributions across North America, but the latter prefers sandier soils (Heiser

and Smith, 1969, Todesco et al., 2020). Over two growing seasons, we measured UV floral

patterns (as the proportion of the ligule that absorbs UV radiation, henceforth "Ligule UV

proportion" or "LUVp") in 1589 *H. annuus* individuals derived from 110 distinct natural

populations, and 351 *H. petiolaris* individuals from 40 populations, grown in common garden

experiments in Vancouver, Canada (Todesco et al., 2020) (Figure 1a,b; Figure 1 – source data 1).

While extensive variation was observed within both species, it was particularly striking for *H.*

annuus, which displayed a phenotypic continuum from ligules with almost no UV pigmentation

to ligules that were entirely UV-absorbing (Figure 1c-e; Figure 1 – source data 2). A relatively

high proportion of *H. annuus* individuals (~13%) had completely UV-absorbing ligules and

therefore lacked UV "nectar guides", suggesting that pollinator orientation is not a necessary

function of floral UV pigmentation in sunflower.

Figure 1: Diversity for floral UV pigmentation patterns in wild sunflowers.

a, Geographic distribution of sampled populations for *H. annuus* and **b,** *H. petiolaris***.** Yellow/orange dots represent

different populations, overlaid grey dots size is proportional to the population mean LUVp. **c,** Range of variation for

floral UV pigmentation patterns in the two species. Scale bar = 1 cm. **d,** LUVp values distribution for *H. annuus* and

e, *H. petiolaris* subspecies.

Genetic control of floral UV patterning

To identify the loci controlling variation for floral UV patterning, we performed a genome-wide

association study (GWAS). We used a subset of the phenotyped plants (563 of the *H. annuus* and

all 351 *H. petiolaris* individuals) for which we previously generated genotypic data at >4.6M

high-quality single-nucleotide polymorphisms (SNPs) (Todesco et al., 2020). Given their

relatively high level of genetic differentiation, analyses were performed separately for the

petiolaris and *fallax* subspecies of *H. petiolaris* (Todesco et al., 2020). We detected several

genomic regions significantly associated with UV patterning in *H. petiolaris petiolaris*, and a

particularly strong association ($P = 5.81e^{-25}$) on chromosome 15 in *H. annuus* (Figure 2a,b;

Figure 2 – figure supplement 1). The chromosome 15 SNP with the strongest association with

ligule UV pigmentation patterns in *H. annuus* (henceforth "Chr15_LUVp SNP") explained 62%

of the observed variation, and allelic distributions at this SNP closely matched that of floral UV

patterns (Figure 2c, compare to Figure 1a; Figure 1 – source data 2).

Genotype at the Chr15_LUVp SNP had a remarkably strong effect on the size of UV bullseyes in

inflorescences. Individuals homozygous for the "large" (L) allele had a mean LUVp of 0.78

(st.dev ±0.16), meaning that ~3/4 of the ligule was UV-absorbing, while individuals homozygous

for the "small" (S) allele had a mean LUVp of 0.33 (st.dev. ±0.15), meaning that only the basal

~1/3 of the ligule absorbed UV radiation. Consistent with the trimodal LUVp distribution

observed for *H. annuus* (Figure 1d), alleles at this locus showed additive effects, with

heterozygous individuals having intermediate phenotypes (LUVp = 0.59 ± 0.18; Figure 2d). The

association between floral UV patterns and the Chr15_LUVp SNP was confirmed in the F_2

progeny of crosses between plants homozygous for the L allele (with completely UV-absorbing

ligules; LUVp = 1) and for the S allele (with a small UV-absorbing patch at the ligule base;

LUVp < 0.18; Figure 2e; Figure 2 – figure supplement 2). Average LUVp values were lower,

and their range smaller, when these populations were grown in a greenhouse rather than in a

field. Plants in the greenhouse experienced relatively uniform temperatures and humidity, and

were shielded from most UV radiation. This suggests that while floral UV patterns have a strong

genetic basis (consistent with previous observations (Koski and Ashman, 2013)), their

expression is also affected by the environment.

Figure 2: A single locus explains most of the variation in floral UV patterning in *H. annuus*.

a, LUVp GWAS. **b,** Zoomed-in Manhattan plot for the chromosome 15 LUVp peak in *H. annuus*. Red lines

represent 5% Bonferroni-corrected significance. GWAs were calculated using two-sided mixed models. Number of

individuals: n = 563 individuals (*H. annuus*); n = 159 individuals (*H. petiolaris petiolaris*). Only positions with -

log10 p-value >2 are plotted. **c,** Geographic distribution of Chr15_LUVp SNP allele frequencies in *H. annuus*. L =

Large and S = Small allele. **d,** LUVp associated with different genotypes at Chr15_LUVp SNP in natural

populations of *H. annuus* grown in a common garden. All pairwise comparisons are significant for $P < 10^{-16}$ (one-

way ANOVA with post-hoc Tukey HSD test, F = 438, df = 2; n = 563 individuals). LUVp values and genotype data

for Chr15_LUVp SNP are reported in Figure 1 – source data 2. **e,** LUVp associated with different genotypes at

Chr15_LUVp SNP in *H. annuus* F_2 populations grown in the field or in a greenhouse (GH). Measurements for the

parental generations are shown: squares = grandparents (field-grown); empty circles = F_1 parents (greenhouse-

grown; figure 2 – figure supplement 2). Boxplots show the median, box edges represent the 25^{th} and 75^{th} percentiles,

whiskers represent the maximum/minimum data points within 1.5x interquartile range outside box edges.

Differences between genotypic groups are significant for P = 0.0057 (Pop. 1 Field, one-way ANOVA, F = 5.73, df =

2; n = 54 individuals); P = 0.0021 (Pop. 2 Field, one-way ANOVA, F = 7.02, df = 2; n = 50 individuals); P =

0.00015 (Pop. 1 GH, one-way ANOVA, F = 11.13, df = 2; n = 42 individuals); P = 0.054 (Pop. 2 GH, one-way

ANOVA, F = 3.17, df = 2; n = 38 individuals). P-values for pairwise comparisons for panels d and e are reported in

the source data for this figure.

***HaMYB111* regulates UV pigment production**

While no obvious candidate genes were found for the GWA peaks for floral UV pigmentation in

H. petiolaris petiolaris, the *H. annuus* chromosome 15 peak is ~5 kbp upstream of *HaMYB111*, a

sunflower homolog of the *Arabidopsis thaliana AtMYB111* gene (Figure 2b). Together with

AtMYB11 and AtMYB12, AtMYB111 is part of a small family of transcription factors (also

called PRODUCTION OF FLAVONOL GLYCOSIDES, PFG) that controls the expression of

genes involved in the production of flavonol glycosides in Arabidopsis (Stracke et al., 2007).

Flavonol glycosides are a subgroup of flavonoids known to fulfill a variety of functions in plants,

including protection against abiotic and biotic stresses (e.g. UV radiation, cold, drought,

herbivory) (Pollastri and Tattini, 2011). Crucially, they absorb strongly in the near UV range

(300-400 nm), and are the pigments responsible for floral UV patterns in several plant species

(Brock et al., 2016, Rieseberg and Schilling, 1985, Sheehan et al., 2016, Thompson et al., 1972).

For instance, alleles of a homolog of *AtMYB111* are responsible for the evolutionary gain and

subsequent loss of flavonol accumulation and UV absorption in flowers of *Petunia* species,

associated with two successive switches in pollinator preferences (from bees, to hawkmoths, to

hummingbirds (Sheehan et al., 2016)). A homolog of *AtMYB12* has also been associated with

variation in floral UV patterns in *Brassica rapa* (Brock et al., 2016). Consistent with this, we

found flavonol glycosides to be the main UV-absorbing pigments in sunflower ligules,

accumulating at much higher levels at their base, and in ligules of plants with larger LUVp

(Figure 3a,b).

AtMYB12 and *AtMYB111* are known to have the strongest effect on flavonol glycoside

accumulation in Arabidopsis (Stracke et al., 2007, Stracke et al., 2010). We noticed, from

existing RNAseq data, that *AtMYB111* expression levels are particularly high in petals

(Klepikova et al., 2016) (Figure 3c), and found that Arabidopsis petals, while uniformly white in

the visible spectrum, absorb strongly in the UV (Figure 3d; Figure 3 – figure supplement 1). To

our knowledge, this is the first report of floral UV pigmentation in Arabidopsis, a highly selfing

species which is seldom insect-pollinated (Hoffmann et al., 2003). Accumulation of flavonol

glycosides in petals is strongly reduced, and UV pigmentation is almost completely absent, in

petals of mutants for *AtMYB111* (*myb111*; Figure 3d,e). UV absorbance is further reduced in

petals of double mutants for *AtMYB12* and *AtMYB111* (*myb12/111*). However, petals of the

single mutant for *AtMYB12* (*myb12*), which is expressed at low levels throughout the plant

(Klepikova et al., 2016), are indistinguishable from wild-type plants (Figure 2 – figure

supplement 1). This shows that flavonol glycosides are responsible for floral UV pigmentation

also in Arabidopsis, and that *AtMYB111* plays a fundamental role in controlling their

accumulation in petals.

To confirm that sunflower *HaMYB111* is functionally equivalent to its Arabidopsis homolog, we

introduced it into *myb111* plants. Expression of *HaMYB111*, either under the control of a

constitutive promoter or of the endogenous *AtMYB111* promoter, restored petal UV pigmentation

and induced accumulation of flavonol glycosides (Figure 3d,e). *HaMYB111* coding sequences

obtained from wild sunflowers with large or small LUVp were equally effective at

complementing the *myb111* mutant. Together with the observation that the strongest GWAS

association with LUVp fell in the promoter region of *HaMYB111*, these results suggest that

differences in the effect of the "small" and "large" alleles of this gene on floral UV pigmentation

are not due to differences in protein function, but rather to differences in gene expression.

Analysis of *HaMYB111* expression in cultivated sunflower revealed that, consistent with a role in

floral UV pigmentation and similar to its Arabidopsis counterpart, it is expressed specifically in

ligules, and it is almost undetectable in other tissues (Badouin et al., 2017) (Figure 3f). Similar to

observations in *Rudbeckia hirta*, another member of the *Heliantheae* tribe (Schlangen et al.,

2009), UV pigmentation is established early in ligule development in both *H. annuus* and *H.*

petiolaris, as their visible colour turns from green to yellow before the inflorescence opens (R4

developmental stage (Schneiter and Miller, 1981); Figure 3g; Figure 3 – figure supplement 2).

HaMYB111 is highly expressed in the part of the ligule that accumulates UV-absorbing

pigments, and especially in developing ligules, consistent with a role in establishing

pigmentation patterns (Figure 3h). We also observed a matching expression pattern for *HaFLS1*,

the sunflower homolog of a gene encoding one of the main enzymes controlling flavonol

biosynthesis in Arabidopsis (*FLAVONOL SYNTHASE 1, AtFLS1*), whose expression is regulated

directly by *AtMYB111* (Stracke et al., 2007) (Figure 3i). Finally, we compared *HaMYB111*

expression levels in a set of 46 field-grown individuals with contrasting LUVp values,

representing 21 different wild populations. *HaMYB111* expression levels differed significantly

between the two groups ($P = 0.009$; Figure 3j). Variation in expression levels within phenotypic

classes was quite large; this is likely due in part to the strong dependence of *HaMYB111*

expression on developmental stage (Figure 3g), and the difficulty of accurately establishing

matching ligule developmental stages across diverse wild sunflowers.

These expression analyses further point to *cis*-regulatory rather than coding sequence differences

between *HaMYB111* alleles being responsible for LUVp variation. Accordingly, direct

sequencing of the *HaMYB111* locus from multiple wild *H. annuus* individuals, using a

combination of Sanger sequencing and long PacBio HiFi reads, identified no coding sequence

variants associated with differences in floral UV patterns, or with alleles at the Chr15_LUVp

SNP. However, we observed extensive variation in the promoter region of *HaMYB111*,

differentiating wild *H. annuus* alleles from each other and from the reference assembly for

cultivated sunflower. Relaxing quality filters to include less well-supported SNPs in our LUVp

GWAS did not identify additional variants with stronger associations than Chr15_LUVp SNP

(Figure 2 – figure supplement 2). However, many of the polymorphisms we identified by direct

sequencing were either larger insertions/deletions (indels) or fell in regions that were too

repetitive to allow accurate mapping of short reads, and would not be included even in the

expanded SNP dataset. While several of these variants in the promoter region of *HaMYB111*

appeared to be associated with the Chr15_LUVp SNP, further studies will be required to confirm

this, and to identify their eventual effects on *HaMYB111* activity.

Interestingly, when we sequenced the promoter region of *HaMYB111* in several *H. argophyllus*

and *H. petiolaris* individuals, we found that they all carried the S allele at the Chr15_LUVp SNP.

Similarly, in a set of previously re-sequenced wild sunflowers, we found the S allele to be fixed

in several perennial (*H. decapetalus*, *H. divaricatus* and *H. grosseserratus*) and annual sunflower

species (*H. argophyllus*, *H. niveus*, *H. debilis*), and to be at >0.98 frequency in *H. petiolaris*,

suggesting that the "small" haplotype is ancestral. Conversely, the L allele at Chr15_LUVp SNP

was almost fixed (>0.98 frequency) in a set of 285 cultivated sunflower lines (Mandel et al.,

2013). Consistent with these patterns, UV bullseyes are considerably smaller in *H. argophyllus*

(mean LUVp ± st.dev. = 0.27 ± 0.09), *H. niveus* (0.15 ± 0.09), and *H. petiolaris* (0.27 ± 0.12;

Figure 1e) than in cultivated sunflower lines (0.62 ± 0.23). Additionally, while 50 of the

cultivated sunflower lines had completely or almost completely UV-absorbing ligules (LUVp >

0.8), no such case was observed in the other three species (Figure 1 – figure supplement 2).

Figure 3: *MYB111* is associated with floral UV pigmentation patterns and flavonol accumulation in sunflower

and Arabidopsis.

 a, UV chromatograms (350 nm) for methanolic extracts of the upper and lower third of ligules with intermediate

UV patterns, and **b,** of ligules with large and small floral UV patterns. Peaks corresponding to flavonols are labelled

(Figure 3 – source data 1). **c,** Expression levels of *AtMYB111* in Arabidopsis. RPKM = Reads Per Kilobase of

transcript per Million mapped reads. **d,** UV pictures of Arabidopsis petals. *HaMYB111* from *H. annuus* plants with

small or large LUVp was introduced into the Arabidopsis *myb111* mutant under the control of a constitutive

promoter (*p35SCaMV*) or of the promoter of the Arabidopsis homolog (*pAtMYB111*). All petals are white in the

visible range (Figure 3 – figure supplement 1). Scale bar = 1mm. **e,** UV chromatograms (350 nm) for methanolic

extracts of petals of Arabidopsis lines. Upper panel: wild-type Col-0 and mutants. Bottom panel:

p35SCaMV::HaMYB111 lines in *myb111* background. Peaks corresponding to flavonols are labelled (Figure 3 –

source data 1). **f,** Expression levels of *HaMYB111* in cultivated sunflower. **g,** Pigmentation patterns in ligules of wild

H. annuus at different developmental stages: R3 = closed inflorescence bud; R4 = inflorescence bud opening; R5 =

inflorescence fully opened. **h,** Expression levels in the UV-absorbing (grey) and UV-reflecting (yellow) portion of

mature (mat) and developing (dev) ligules for *HaMYB111* and **i,** *HaFLS1*, one of its putative targets. Each bar

represents a biological replicate (different inflorescence from a same individual). **j,** *HaMYB111* expression levels in

field-grown wild *H. annuus* with different floral UV pigmentation patterns. The difference between the two groups

is significant for $P = 0.009$ (Welch t-test, $t = 2.81$, df = 27.32, two-sided; $n = 46$ individuals). Boxplots show the

median, box edges represent the 25^{th} and 75^{th} percentiles, whiskers represent the maximum/minimum data points

within 1.5x interquartile range outside box edges.

A dual role for floral UV pigmentation

Although our results show that *HaMYB111* explains most of the variation in floral UV

pigmentation patterns in wild *H. annuus*, why such variation exists in the first place is less clear.

Several hypotheses have been advanced to explain the presence of floral UV patterns and their

variability. Like their visible counterparts, UV pigments play a fundamental role in pollinator

attraction (Horth et al., 2014, Koski and Ashman, 2014, Rae and Vamosi, 2013, Sheehan et al.,

2016). For example, in *Rudbeckia* species, artificially increasing the size of bullseye patterns to

up to 90% of the petal surface resulted in rates of pollinator visitation equal to or higher than

wild-type flowers (which have on average 40-60% of the petal being UV-absorbing).

Conversely, reducing the size of the UV bullseye had a strong negative effect on pollinator

visitation (Horth et al., 2014). To test whether the relative size of UV bullseye patterns affected

pollination, we compared insect visitation rates for wild *H. annuus* lines with contrasting UV

bullseye patterns. An initial experiment, carried out in Vancouver (Canada), found that flowers

with large UV patterns received significantly more visits (Figure 4a). Vancouver is outside the

natural range of *H. annuus*, suggesting that our results are unlikely to be affected by learned

preferences (i.e. pollinators preferring UV patterns they are familiar with in sunflower). While

this experiment revealed a clear pattern of pollinator preferences, it involved plants from only

two different populations, and effects of other unmeasured factors unrelated to UV pigmentation

on visitation patterns cannot be excluded. Therefore, we monitored pollinator visitation in plants

grown in a field including 1484 individuals from 106 *H. annuus* populations, spanning the entire

range of the species. Within this field, we selected 82 plants, from 49 populations, which

flowered at roughly the same time and had comparable numbers of flowers. We divided these

plants into three categories, based on the species-wide distribution of LUVp values in *H. annuus*

(Figure 1d): small (0-0.3); intermediate (0.5-0.8) and large (>0.95) LUVp. Plants with

intermediate UV patterns had the highest visitation rates (Figure 4b; Figure 4 – figure

supplement 1). Visitation to plants with small or large UV patterns was less frequent, and

particularly low for plants with very small LUVp values (<0.15). A strong reduction in

pollination would be expected to result in lower fitness, and to be negatively selected;

accordingly, plants with such small LUVp values were rare (~1.5% of the individuals grown).

These results confirm that floral UV patterns play a major role in pollinator attraction, as has

been already extensively reported (Horth et al., 2014, Koski and Ashman, 2014, Rae and

Vamosi, 2013, Sheehan et al., 2016). They also agree with previous findings in other plant

species suggesting that intermediate-to-large UV bullseyes are preferred by pollinators, and only

very small UV bullseyes are maladaptive (Horth et al., 2014, Koski and Ashman, 2014). While

we cannot exclude that smaller UV bullseyes would be preferred by pollinators in some parts of

the *H. annuus* range, this does not seem likely; the most common pollinators of sunflower are

ubiquitous across the range of *H. annuus*, and many bee species known to pollinate sunflower

are found in both regions where *H. annuus* populations have large LUVp and regions where they

have small LUVp (Hurd Jr et al., 1980). While acting as visual cues for pollinators is therefore

clearly a major function of floral UV bullseyes, it is unlikely to (fully) explain the patterns of

variation that we observe for this trait.

In recent years, the importance of non-pollinator factors in driving selection for floral traits has

been increasingly recognized (Strauss and Whittall, 2006). Additionally, flavonol glycosides, the

pigments responsible for floral UV patterns in sunflower, are known to be involved in responses

to several abiotic stressors (Korn et al., 2008, Nakabayashi et al., 2014b, Pollastri and Tattini,

2011, Schulz et al., 2015). Therefore, we explored whether some of these stressors could drive

diversification in floral UV pigmentation. An intuitively strong candidate is UV radiation, which

can be harmful to plant cells (Stapleton, 1992). Variation in the size of UV bullseye patterns

across the range of *Argentina anserina* (a member of the *Rosaceae* family) has been shown to

correlate positively with intensity of UV radiation. Flowers of this species are bowl-shaped, and

larger UV-absorbing regions have been proposed to protect pollen from UV damage by

absorbing UV radiation that would otherwise be reflected toward the anthers (Koski and

Ashman, 2015). However, sunflower inflorescences are much flatter than *A. anserina* flowers,

making it unlikely that any significant amount of UV radiation would be reflected from the

ligules towards the disk flowers. Studies in another plant with non-bowl-shaped flowers (*Clarkia*

unguiculata) have found no evidence of an effect of floral UV patterns in protecting pollen from

UV damage (Peach et al., 2020). Consistent with this, the associations between the intensity of

UV radiation at our collection sites and floral UV patterns in *H. annuus* was weak (*H. annuus*: R^2

$= 0.01$, $P = 0.12$; Figure 4c; Figure 4 – figure supplement 2).

Across the *Potentillae* tribe (*Rosaceae*), floral UV bullseye size is also weakly associated with

UV radiation, but is more strongly correlated with temperature, with lower temperatures being

associated with larger UV bullseyes (Koski and Ashman, 2016). We found a similar, strong

correlation with temperature in our dataset, with average summer temperatures explaining a large

fraction of the variation in LUVp in *H. annuus* ($R^2 = 0.44$, $P = 2.4 \times 10^{-15}$; Figure 4d; Figure 4 –

figure supplement 2). It has been suggested that the radiation absorbed by floral UV pigments

could contribute to increasing the temperature of the flower, similar to what has been observed

for visible pigments (Koski et al., 2020). This possibility is particularly intriguing for sunflower,

in which flower temperature plays an important role in pollinator attraction; inflorescences of

cultivated sunflowers consistently face East so that they warm up faster in the morning, making

them more attractive to pollinators (Atamian et al., 2016). Larger UV bullseyes could therefore

contribute to increasing temperature of the sunflower inflorescences, and their attractiveness of

sunflowers to pollinators, in cold climates. However, different levels of UV pigmentation had no

effect on the temperature of inflorescences or individual ligules exposed to sunlight (Figure 4e-g;

Figure 4 – figure supplement 3). This is perhaps not surprising, given that UV wavelengths

represents only a small fraction (3-7%) of the solar radiation reaching the Earth surface

(compared to >50% for visible wavelengths), and are therefore unlikely to provide sufficient

energy to significantly warm up the ligules (Nunez et al., 1994).

While several geoclimatic variables are correlated across the range of wild *H. annuus*, the single

variable explaining the largest proportion of the variation in floral UV patterns in this species

was summer relative humidity (RH; $R^2 = 0.51$, $P = 1.4 \times 10^{-18}$; Figure 4h; Figure 4 – figure

supplement 2), with lower humidity being associated with larger LUVp values. Lower relative

humidity is generally associated with higher transpiration rates in plants, leading to increased

water loss. Flavonol glycosides are known to play an important role in responses to drought

stress (Nakabayashi et al., 2014a); in particular, Arabidopsis lines that accumulate higher

concentrations of flavonol glycosides due to over-expression of *AtMYB12* lose water and

desiccate at slower rates than wild-type plants (Nakabayashi et al., 2014b). Similarly, we found

that completely UV-absorbing ligules desiccate significantly slower than largely UV-reflecting

ligules (Figure 4i). This difference is not due to general differences in transpiration rates between

genotypes, since we observed no comparable trend for rates of leaf desiccation in the same set of

sunflower lines (Figure 4j). Transpiration from flowers can be a major source of water loss for

plants, and this is known to drive, within species, the evolution of smaller flowers in populations

living in dry locations (Galen, 2000, Herrera, 2005, Lambrecht, 2013, Lambrecht and Dawson,

2007). Thus, variation in floral UV pigmentation in sunflowers is likely similarly driven by the

role of flavonol glycosides in reducing water loss from ligules, with larger floral UV patterns

helping prevent drought stress in drier environments.

One of the main roles of transpiration in plants is facilitating heat dispersion at higher

temperatures through evaporative cooling (Burke and Upchurch, 1989, Drake et al., 2018),

which could explain the strong correlation between LUVp and temperature across the range of

H. annuus (Figure 4d). Consistent with this, summer relative humidity and summer temperatures

together explain a considerably larger fraction of the variation for LUVp in *H. annuus* than either

variable alone ($R^2 = 0.63$; $P = 0.0017$; Figure 1 – source data 1), with smaller floral UV patterns

being associated with higher relative humidity and higher temperatures (Figure 4 – figure

supplement 2). Despite a more limited range of variation for LUVp, the same trend is present

also in *H. petiolaris* (Figure 4 – figure supplement 4). Consistent with a role of floral UV

pigmentation in the plant's response to variation in both humidity and temperature, we found

strong associations (dB > 10) between SNPs in the *HaMYB111* region and these variables in

genotype-environment association (GEA) analyses (Figure 4k; Figure 4 – source data 3).

Figure 4: Accumulation of UV pigments in flowers affects pollinator visits and transpiration rates.

a, Rates of pollinator visitation measured in Vancouver in 2017 ($P = 0.017$; Mann-Whitney U-tests, W = 150, two-

sided; $n = 143$ pollinator visits) and **b,** in 2019 (differences between LUVp categories are significant for $P = 0.0058$,

Kruskal-Wallis test, $\chi^2 = 14.54$, df = 4; $n = 1390$ pollinator visits. Letters identify groups that are significantly

different for $P < 0.05$ in pairwise comparisons, Wilcoxon rank sum test. Exact p-values are reported in the source

data for this figure). Boxplots show the median, box edges represent the 25^{th} and 75^{th} percentiles, whiskers represent

the maximum/minimum data points within 1.5x interquartile range outside box edges. **c,** Correlation between

average LUVp for different populations of *H. annuus* and summer UV radiation ($R^2 = 0.01$, $P = 0.12$) or **d,** summer

average temperature ($R^2 = 0.44$, $P = 2.4 \times 10^{-15}$). Grey areas represent 95% confidence intervals. **e,** Sunflower

inflorescences pictured in the visible, UV and infrared (IR) range. In the IR picture, a bumblebee is visible in the

inflorescence with large LUVp (right). The higher temperature in the centre (disk) of the inflorescence with small

LUVp does not depend on ligule UV patterns (Figure 4 – support figure 3). **f,** *H. annuus* ligules after having been

exposed to sunlight for 15 minutes. **g,** Pairs of ligules from different sunflower lines were exposed to sunlight for 15

minutes, and their average temperature was measured from IR pictures. **h,** Correlation between average LUVp in *H.*

annuus populations and summer relative humidity ($R^2 = 0.51$, $P = 1.4 \times 10^{-18}$). The grey area represents the 95%

confidence interval. **i,** Rate of water loss from ligules and **j,** leaves of wild *H. annuus* plants with large or small

LUVp. Values reported are means ± standard error of the mean. $n = 16$ (ligules) or 15 (leaves). Ligules and leaves

were left to air-dry and weighed every hour for five hours, after they were left to air-dry overnight (o.n.), and after

they were incubated in an oven to remove any residual humidity (oven-dry). Asterisks denote significant differences

($p < 0.05$, two-sided Welch t-test; exact p-values are reported in the source data for this figure). **k,** GEA for summer

average temperature (Av. T) and summer relative humidity (RH) in the *HaMYB111* region. The dashed orange line

represents Bayes Factor (BF_{is}) = 10 deciban (dB). GEAs were calculated using two-sided XtX statistics. $n = 71$

populations.

Conclusions

Connecting adaptive variation to its genetic basis is one of the main goals of evolutionary

biology. Here, we show that regulatory variation at a single major gene, the transcription factor

HaMYB111, underlies most of the variation for floral UV patterns in wild *H. annuus*, and that

these UV patterns not only have a strong effect on pollinator visits, but they also co-vary with

geoclimatic variables (especially relative humidity and temperature) and affect desiccation rates

in ligules. By reducing the amount of transpiration in environments with lower relative humidity,

UV-absorbing pigments in the ligule help prevent excessive water loss and maintain ligule

turgidity; in humid, hot environments (e.g. Southern Texas), lower accumulation of flavonol

glycosides would instead promote transpiration from ligules, keeping them cool and avoiding

over-heating. The presence of UV pigmentation in the petals of Arabidopsis (also controlled by

the Arabidopsis homolog of *MYB111*) further points to a more general protective role of these

pigments in flowers, since pollinator attraction is likely not critical for fertilization in this largely

selfing species. Additionally, a role in reducing water loss from petals is consistent with the

overall trend in increased size of floral UV patterns over the past 80 years that has been observed

in herbarium specimens (Koski et al., 2020); due to changing climates, relative humidity over

land has been decreasing in recent decades, which could result in higher transpiration rates

(Byrne and O'Gorman, 2018). Further studies will be required to confirm the existence of this

trend and assess its strength.

More generally, our study highlights the complex nature of adaptive variation, with selection

pressures from both biotic and abiotic factors shaping the patterns of diversity that we observe

across natural populations. Floral diversity in particular has long been attributed to the actions of

animal pollinators. Our work adds to a growing literature demonstrating the contributions of

abiotic factors, most notably drought and heat stress, to this diversity. In sum, it is not all about

sex, even for flowers.

Methods

Plant material and growth conditions

Sunflower lines used in this paper were grown from seeds collected from wild populations

(Todesco et al., 2020), or obtained from the North Central Regional Plant Introduction Station in

Ames, Iowa, USA. Sunflower seeds were surface sterilized by immersion for 10 minutes in a

1.5% sodium hypochlorite solution. Seeds were then rinsed twice in distilled water and treated

for at least one hour in a solution of 1% PPM (Plant Cell Technologies, Washington, DC, USA),

a broad-spectrum biocide/fungicide, to minimize contamination, and 0.05 mM gibberellic acid

(Sigma-Aldrich, St. Louis, MO, USA). They were then scarified, de-hulled, and kept for two

weeks at 4 ºC in the dark on filter paper moistened with a 1% PPM solution. Following this,

seeds were kept in the dark at room temperature until they germinated. For common garden

experiments, the seedlings were then transplanted in peat pots, grown in a greenhouse for two

weeks, then moved to an open-sided greenhouse for a week for acclimation, and finally

transplanted in the field. For all other experiments seedlings were transplanted in 2-gallon pots

filled with Sunshine #1 growing mix (Sun Gro Horticulture Canada, Abbotsford, BC, Canada).

For the wild sunflower species shown in Figure 1 – figure supplement 1b, following sterilization,

seeds were scarified and then dipped in fuscicosin solution (1.45 µM) for 15 minutes, dehulled,

germinated in the dark for at least 8-10 days, and then grown in pots for three weeks before

transplanting into 2-gallon pots filled with a blend of sandy loam, organic compost and mulch.

Those plants were grown at the UC Davis field experiment station (California, USA) from July

to October 2017. A complete list of sunflower accessions and their populations of origin is

reported in Figure 1 - source data 1 and Figure 1 - source data 2.

Seeds from the following Arabidopsis lines were obtained from the Arabidopsis Biological

Resource Center: Col-0 (CS28167), *myb111* (CS9813), *myb12* (CS9602) and *myb12/myb111*

(CS9980). Seeds were stratified in 0.1% agar at 4 ºC in the dark for four days, and then sown in

pots containing Sunshine #1 growing mix. Plants were grown in growth chambers at 23 ºC in

long-day conditions (16 h light, 8 h dark).

Common garden

Two common garden experiments were performed, in 2016 and 2019. After germination and

acclimation, plants were transplanted at the Totem Plant Science Field Station of the University

of British Columbia (Vancouver, Canada). In the 2016 common garden experiment, each

sunflower species was grown in a separate field. Pairs of plants from the same population were

randomly distributed within each field. In the 2019 common garden experiment, plants were

sown using a completely randomized design.

In the summer of 2016, ten plants from each of 151 selected populations of wild *H. annuus*, *H.*

petiolaris, *H. argophyllus* and *H. niveus* were grown. Plants were transplanted in the field on the

25th of May (*H. argophyllus*), 2nd of June (*H. petiolaris* and *H. niveus*) and 7th of June 2016 (*H.*

annuus). Up to four inflorescences from each plant were collected for visible and UV

photography. In the summer of 2019, fourteen plants from each of 106 populations of wild *H.*

annuus were transplanted in the field on 6th of June. These included 65 of the populations grown

in the previous common garden experiment, and 41 additional populations that were selected to

complement their geographic distribution. At least two ligules from different inflorescences for

each plant were collected for visible and UV photography.

Sample size for the common garden experiments was determined by the available growing space

and resources. Ten-to-fourteen individuals were grown for each population because this would

provide a good representation of the variation present in each population, while maximizing the

number of populations that could be surveyed.

Researcher were not blinded as to the identity of individual samples. However, information about

their populations of origin and/or LUVp phenotypes were not attached to the samples during data

acquisition.

Ultraviolet and infrared photography

Ultraviolet patterns were imaged in whole flowerheads or detached ligules using a Nikon D70s

digital camera, fitted with a Noflexar 35-mm lens and a reverse-mounted 2-inch Baader U-Filter

(Baader Planetarium, Mammendorf, Germany), which only allows the transmission of light

between 320 and 380 nm. Wild sunflower species shown in Figure 1 – figure supplement 1b

were imaged using a Canon DSLR camera in which the internal hot mirror filter had been

replaced with a UV bandpass filter (LifePixel, Mukilteo, WA). The length of the whole ligule

(L_L) and the length of the UV-absorbing part at the base of the ligule (L_{UV-abs}) were measured

using Fiji (Schindelin et al., 2012, Schneider et al., 2012). Ligule UV proportion was measured

as the ratio between the two ($LUVp = L_{UV-abs}/L_L$). In some *H. annuus* individuals, the upper,

"UV-reflecting" portion of the ligules (L_{UV-ref}) also displayed a lower level of UV-absorption; in

those cases, these regions were weighted at 50% of fully UV-absorbing regions, using the

formula $LUVp = (L_{UV-abs}/L_L) + \frac{1}{2}(L_{UV-ref}/L_L)$. For pictures of whole inflorescences, LUVp values

were averaged from three different ligules. For each individual, LUVp values were averaged

between all the inflorescences or detached ligules measurements.

Infrared pictures were taken using a Fluke TiX560 thermal imager (Fluke Corporation, Everett,

WA, USA) and analyzed using the Fluke Connect software (v1.1.536.0). For time series

experiments, plants were germinated as above (see "Common Garden"), grown in 2-gallon pots

in a greenhouse until they produced four true leaves, and then moved to the field. On three

separate days in August 2017, pairs of inflorescences with opposite floral UV patterns at similar

developmental stages were selected and made to face East. Infrared images were taken just

before sunrise, ~5 minutes after sunrise, and then at 0.5, 1, 2, 3 and 4 hours after sunrise.

For infrared pictures of detached ligules, plants were grown in a greenhouse. Flowerheads were

collected and kept overnight in a room with constant temperature of 21 ºC, with their stems

immersed in a beaker containing distilled water. The following day, pairs of ligules with

contrasting LUVp were removed and arranged on a sheet of white paper. Infrared pictures were

taken immediately before exposing the ligules to the sun, and again 15 minutes after that.

Library preparation, sequencing and SNP calling

Whole-genome shotgun (WGS) sequencing library preparation and sequencing, as well as SNP

calling and variant filtering, for the *H. annuus* and *H. petiolaris* individuals used for GWA

analyses in this paper were previously described (Todesco et al., 2020). Briefly, DNA was

extracted from leaf tissue using a modified CTAB protocol (Murray and Thompson, 1980, Zeng

et al., 2002), the DNeasy Plant Mini Kit or a DNeasy 96 Plant Kit (Qiagen, Hilden, Germany).

Genomic DNA was sheared to an average fragment size of 400 bp using a Covaris M220

ultrasonicator (Covaris, Woburn, MA, USA). Libraries were prepared using a protocol largely

based on (Rowan et al., 2015), the TruSeq DNA Sample Preparation Guide from Illumina

(Illumina, San Diego, CA, USA) and (Rohland and Reich, 2012), with the addition of an

enzymatic repeats depletion step using a Duplex-Specific Nuclease (DSN; Evrogen, Moscow,

Russia) (Matvienko et al., 2013, Shagina et al., 2010, Todesco et al., 2020). All libraries were

sequenced at the McGill University and Génome Québec Innovation Center on HiSeq2500,

HiSeq4000 and HiSeqX instruments (Illumina) to produce paired end, 150 bp reads.

Sequences were trimmed for low quality using Trimmomatic (v0.36) (Bolger et al., 2014) and

aligned to the *H. annuus* XRQv1 genome (Badouin et al., 2017) using NextGenMap (v0.5.3)

(Sedlazeck et al., 2013). We followed the best practices recommendations of The Genome

Analysis ToolKit (GATK) (Poplin et al., 2017) and executed steps documented in GATK's

germline short variant discovery pipeline (for GATK 4.0.1.2). During genotyping, to reduce

computational time and improve variant quality, genomic regions containing transposable

elements were excluded (Badouin et al., 2017). We then used GATK's VariantRecalibrator

(v4.0.1.2) to select high quality variants. SNP data were then filtered for minor allele frequency

(MAF) $\geq$ 0.01, genotype rate $\geq$ 90%, and to keep only bi-allelic SNPs.

Filtered SNPs were then re-mapped to the improved reference assembly HA412-HOv2 (Staton

and Lázaro-Guevara, 2020) using BWA (v0.7.17) (Li, 2013). These re-mapped SNPs were used

for all analyses, excluding the GWA for the region surrounding the *HaMYB111* locus that used

un-filtered variants based on the XRQv1 assembly (Figure 2 – figure supplement 3).

The SNP dataset used to determine the genotype at the Chr15_LUVp SNP in other species (*H.*

argophyllus, *H. niveus*, *H. debilis*, *H. decapetalus*, *H. divaricatus* and *H. grosseserratus*) was

based on WGS data generated for (Todesco et al., 2020) and is described in (Owens et al., 2021).

Sequence data for the Sunflower Association Mapping population was reported in (Hubner et al.,

2019).

Genome-wide association mapping

Genome-wide association analyses for LUVp were performed for *H. annuus, H. petiolaris petiolaris* and *H. petiolaris fallax,* using two-sided mixed models implemented in EMMAX (v07Mar2010) (Kang et al., 2010) or in the EMMAX module in EasyGWAS (Grimm et al., 2017). For all runs, the first three principal components (PCs) were included as covariates, as well as a kinship matrix. Only SNPs with MAF >5% were included in the analyses. Sample size was estimated to be sufficient to provide an 85% probability of detecting loci explaining 5% or more of the phenotypic variance in *H. annuus*, 8% of variance in *H. petiolaris*.

F$_2$ populations and genotyping

Individuals from population ANN_03 from California, USA (large LUVp) and ANN_55 from Texas, USA (small LUVp) were grown in 2-gallon pots in a field. When the plants reached maturity, they were moved to a greenhouse, where several inflorescences were bagged and crossed. The resulting F$_1$ seeds were germinated and grown in a greenhouse, and pairs of siblings were crossed (wild sunflowers are overwhelmingly self-incompatible). The resulting F$_2$ populations were grown both in a greenhouse in the winter of 2019 ($n = 42$ individuals for population 1, 38 individuals for population 2) and in a field as part of the 2019 common garden experiments ($n = 54$ individuals for population 1, 50 individuals for population 2). DNA was extracted from young leaf tissue as described above. All plants were genotyped for the Chr15_LUVp SNPs using a custom TaqMan SNP genotyping assay (Thermo Fisher Scientific, Walthman, MA, USA) on a Viia 7 Real-Time PCR system (Thermo Fisher Scientific).

Metabolite analyses

Methanolic extractions were performed following (Stracke et al., 2007). Sunflower ligules (or

portions of them) and Arabidopsis petals were collected and flash-frozen in liquid nitrogen. The

tissue was ground to a fine powder by adding 10-15 zirconia beads (1 mm diameter) and using a

TissueLyser (Qiagen) for sunflower ligules, or using a plastic pestle in a 1.5 ml tube for

Arabidopsis petals. 0.5 ml of 80% methanol were added, and the samples were further

homogenized and incubated at 70 ºC for 15 minutes. They were then centrifuged at 15.000g for

10 minutes, and the supernatant was dried in a SpeedVac (Thermo Fisher Scientifics) at 60 ºC.

Samples were then resuspended in 1 µl (sunflower) or 2.5 µl (Arabidopsis) of 80% methanol for

every mg of starting tissue.

The extracts were analyzed by LC/MS/MS using an Agilent 1290 UHPLC system (Agilent

Technologies, Santa Clara, CA, USA) coupled with an Agilent 6530 Quadrupole Time of Flight

mass spectrometer. The chromatographic separation was performed on Atlantis T3- C18

reversed-phase (50 mm × 2.1 mm, 3 µm) analytical columns (Waters Corp, Milford, MA, USA).

The column temperature was set at 40 °C. The elution gradient consisted of mobile phase A

(water and 0.2% formic acid) and mobile phase B (acetonitrile and 0.2% formic acid). The

gradient program was started with 3% B, increased to 25% B in 10 min, then increased to 40% B

in 13 min, increased to 90% B in 17 min, held for 1 min and equilibrated back to 3% B in 20

min. The flow rate was set at 0.4 mL/min and injection volume was 1 µl. A PDA (photo diode

array) detector was used for detection of UV-absorption in the range of 190-600 nm.

MS and MS/MS detection were performed using an Agilent 6530 accurate mass Quadrupole

Time of Flight mass spectrometer equipped with an ESI (electrospray) source operating in both

positive and negative ionization modes. Accurate positive ESI LC/MS and LC/MS/MS data were

processed using the Agilent MassHunter software to identify the analytes. The ESI conditions

were as follows: nebulizing gas (nitrogen) pressure and temperature were 30 psi and 325 °C;

sheath gas (nitrogen) flow and temperature were 12 L/min, 325 °C; dry gas (nitrogen) was 7

l/min. Full scan mass range was 50-1700 m/z. Stepwise fragmentation analysis (MS/MS) was

carried out with different collision energies depending on the compound class.

Transgenes and expression assays

Total RNA was isolated from mature and developing ligules, or part of ligules, using TRIzol

(Thermo Fisher Scientific) and cDNA was synthesized using the RevertAid First Strand cDNA

Synthesis kit (Thermo Fisher Scientific). Genomic DNA was extracted from leaves of

Arabidopsis using CTAB (Murray and Thompson, 1980). A 1959 bp-long fragment

(*pAtMYB111*) from the promoter region of *AtMYB111* (*At5g49330*), including the 5'-UTR of the

gene, was amplified using Phusion High-Fidelity DNA polymerase (New England Biolabs,

Ipswich, MA, USA) and introduced in pFK206 derived from pGREEN (Hellens et al., 2000).

Alleles of *HaMYB111* were amplified from cDNA from ligules, and placed under the control of

pAtMYB111 (in the plasmid describe above) or of the constitutive *CaMV 35S* promoter (in

pFK210, derived as well from pGREEN (Hellens et al., 2000)). Constructs were introduced into

A. thaliana plants by *Agrobacterium tumefaciens*-mediated transformation (strain GV3101)

(Weigel and Glazebrook, 2002). At least five independent transgenic lines with levels of UV

pigmentation comparable to the ones shown in Figure 3d were recovered for each construct. For

expression analyses, qRT-PCRs were performed on cDNA from ligules using the SsoFast

EvaGreen Supermix (Bio-Rad, Hercules, CA, USA) on a CFX96 Real-Time PCR Detection

System (Bio-Rad). Expression levels were normalized against *HaEF1α*. For the expression

analyses shown in Figure 3h,i, portions of ligules were collected at different developmental

stages from three separate inflorescences from one individuals for each species (biological

replicates). Three qRT-PCRs were run for each sample (technical replicates). For the expression

analysis shown in Fig. 3j, samples were collected from wild *H. annuus* individuals grown as part

of the 2019 common garden experiment. Ligules were collected from developing inflorescences

of 24 individuals with large LUVp (from 10 populations) and 22 individuals with small LUVp

(from 11 populations). Three technical replicates were performed for each individual. Sample

size for this experiment was determined by the number of available plants with opposite LUVp

phenotypes and at the appropriate developmental stage. Primers used for cloning and qRT-PCR

are given in Table 1.

Sanger and PacBio sequencing

Fragments ranging in size from 1.5 to 5.5 kbp were amplified from genomic DNA of 20

individuals that had been previously re-sequenced (Todesco et al., 2020), and whose genotype at

the Chr15_LUVp SNP was therefore known, using Phusion High-Fidelity DNA polymerase

(New England Biolabs). Fragments were then cloned in either pBluescript or pJET (Thermo

Fisher Scientific) and sequenced on a 3730S DNA analyzer using BigDye Terminator v3.1

sequencing chemistry (Applied Biosystem, Foster City, CA, USA).

For long reads sequencing, seed from wild *H. annuus* populations known to be homozygous for

different alleles at the Chr15_LUVp SNP were germinated and grown in a greenhouse. After

confirming that they had the expected LUVp phenotype, branches from each plant were covered

with dark cloth for several days, and young, etiolated leaves were collected and immediately

frozen in liquid nitrogen. High molecular weight (HMW) DNA was extracted using a modified

CTAB protocol (Stoffel et al., 2012). All individuals were genotyped for the Chr15_LUVp SNP

using a custom TaqMan SNP genotyping assay (Thermo Fisher Scientific, see above) on a

CFX96 Real-Time PCR Detection System (Bio-Rad). Two individuals, one with large and one

with small LUVp, were selected. HiFi library preparation and sequencing were performed at the

McGill University and Génome Québec Innovation Center on a Sequel II instrument (PacBio,

Menlo Park, CA, USA). Each individual was sequenced on an individual SMRT cell 8M,

resulting in average genome-wide sequencing coverage of 6-8X.

Pollinator preferences assays

In September 2017, pollinator visits were recorded in individual inflorescences of pairs of plants

with large and small LUVp grown in pots in a field adjacent the Nursery South Campus

greenhouses of the University of British Columbia. Pairs of inflorescences were filmed using a

Bushnell Trophy Cam HD (Bushnell, Overland Park, KS, USA) in 12-minute intervals.

Visitation rates were averaged over 14 such movies (Figure 4 – source data 1).

In summer 2019, pollinator visits were scored in a common garden experiment consisting of

1484 *H. annuus* plants at the Totem Plant Science Field Station of the University of British

Columbia. Over five days, between the 29th of July and the 7th of August, pollinator visits on

individual plants were counted over five-minute intervals, for a total of 435 series of

measurements on 111 plants from 51 different populations (Figure 4 – source data 1). To

generate a more homogenous and comparable dataset, measurements for plants with too few (1)

or too many (>10) flowers were excluded from the final analysis.

Correlations with environmental variables and genotype-environment association analyses

Twenty topo-climatic factors were extracted from climate data collected over a 30-year period

(1961-1990) for the geographic coordinates of the population collection sites, using the software

package Climate NA (Wang et al., 2016) (Figure 1 – source data 1). Additionally, UV radiation

data were extracted from the glUV dataset (Beckmann et al., 2014) using the R package "raster"

(Hijmans, 2020, R Core Team, 2020). Correlations between individual environmental variables

and LUVp were calculated using the "lm" function implemented in R; a correlation matrix

between all environmental variables and LUVp was calculated using the "cor" function in R and

plotted using the "heatmap.2" function in the "gplots" package (Warnes et al., 2009).

GEAs were analyzed using BayPass (Gautier, 2015) version 2.1. Population structure was

estimated by choosing 10,000 putatively neutral random SNPs under the BayPass core model.

The Bayes factor (denoted BF_{is} as in (Gautier, 2015)) was then calculated under the standard

covariate mode. For each SNP, BF_{is} was expressed in deciban units [dB, $10 \log_{10} (BF_{is})$].

Significance was determined following (Gautier, 2015), and employing Jeffreys' rule (Jeffreys,

1961), quantifying the strength of associations between SNPs and variables as "strong" (10 dB $\leq$

$BF_{is} <$ 15 dB), "very strong" (15 dB $\leq BF_{is} <$ 20 dB) and decisive ($BF_{is} \geq$ 20 dB).

Desiccation assays

Inflorescence and leaves were collected from well-watered, greenhouse-grown plants, and

brought to an environment kept at 21ºC. They were left overnight with their stems or petioles

immersed in a beaker containing distilled water. The following morning 1-2 leaves from each

plant, and three ligules from each inflorescence were individually weighed and hanged to air dry

at room temperature (21 ºC). Their weight was measured at one-hour intervals for five hours, and

then again the following morning. Leaves and ligules were then incubated for 48 hours at 65 ºC

in an oven to determine their dry weight. Total water content was measured as the difference

between the initial fresh weight (W_0) and dry weight (W_d). Water loss was expressed as a

fraction of the total water content of each organ, using the formula $[(W_i-W_d)/(W_0-W_d)] \times 100$,

where W_i is the weight of a sample at a time i. The assay was performed on samples from 16

(ligules) or 15 (leaves) individuals belonging to 7 (ligules) or 8 (leaves) different populations of

H. annuus.

Data availability

All raw sequenced data are stored in the Sequence Read Archive (SRA) under BioProjects

PRJNA532579, PRJNA398560 and PRJNA736734. Filtered SNP datasets are available at

https://rieseberglab.github.io/ubc-sunflower-genome/. Raw sequencing data and SNP datasets

have been previously described in (Todesco et al., 2020). The sequences of individual alleles at

the *HaMYB111* locus and of *HaMYB111* coding sequences have been deposited at GenBank

under accession numbers XXX-XXX and MZA410295- MZA410296, respectively. All other

data are available in the main text or in the source data provided with the manuscript.

Table 1: Oligonucleotides used in this study.

Target	Forward	Reverse	Product size
HaMYB111 CDS	ATGGGAAGGACCCCGTGTT	TTAAGACTGAAACCATGCATCTACC	885 bp
AtMYB111 promoter	CCTGTGCTTTAAGGCTCGAC	TGCTTCTCGGTCTCTTCTGT	1959 bp
HaMYB111 qPCR	ATGGGAAGGACCCCGTGTT	GCAACTCTTTCCGCATCTCA	169 bp
HaFLS1 qPCR	AAACTACTACCCACCATGCC	TCCTTGTTCACTGTTGTTCTGT	243 bp
EF1alpha qPCR	GTGTGTGATGTCGTTCTCCA	ATTCCACCCAAAGCTTGCTC	167 bp

Acknowledgments

Acknowledgments

This research was conducted in the ancestral and unceded territory of the xʷməθkʷəy̓əm

(Musqueam) People. We thank Andrea Todesco, Daniela Rodeghiero, Emma Borger, Quinn

Anderson, Jennifer Lipka, Jasmine Lai, Hafsa Ahmed, Dominique Skonieczny, Ana Parra,

Cassandra Konecny, Kelsie Morioka and Daniel Yang for assistance with field work and data

acquisition, Melina Byron and Glen Healy at UBC and the UC Davis Plant Sciences Field

Station personnel for assistance with greenhouse and field experiments, Elizabeth Elle and Tyler

Kelly for help planning the pollinator preferences experiments, Laura Marek and the USDA-

ARS in Ames, IA, USA, for providing sunflower seeds, and Chase Mason for providing cuttings

for *Phoebantus tenuifolius*. Maps were realized using tiles from Stamen Design

(https://stamen.com), under CC BY 3.0, from data by OpenStreetMaps contributors

(https://openstreetmap.org), under ODbL. Funding was provided by Genome Canada and

Genome BC (LSARP2014-223SUN), the NSF Plant Genome Program (DBI-1444522 and DBI-

1817628), the University of California, Berkeley, and an HFSP long-term postdoctoral

fellowship to M.T. (LT000780/2013).

Competing interests

The authors declare no competing interests.

References

ATAMIAN, H. S., CREUX, N. M., BROWN, E. A., GARNER, A. G., BLACKMAN, B. K. &

HARMER, S. L. 2016. Circadian regulation of sunflower heliotropism, floral orientation,

and pollinator visits. *Science,* 353**,** 587-90.

BADOUIN, H., GOUZY, J., GRASSA, C. J., MURAT, F., STATON, S. E., COTTRET, L.,

CRESPI, M., MANGIN, B., BURKE, J. M., SALSE, J., MUNOS, S., VINCOURT, P.,

RIESEBERG, L. H. & LANGLADE, N. B. 2017. The sunflower genome provides insights

into oil metabolism, flowering and Asterid evolution. *Nature,* 546**,** 148-152.

BECKMANN, M., VÁCLAVÍK, T., MANCEUR, A. M., ŠPRTOVÁ, L., VON WEHRDEN, H.,

WELK, E. & CORD, A. F. 2014. gl UV: a global UV-B radiation data set for

macroecological studies. *Methods in Ecology and Evolution,* 5**,** 372-383.

BOLGER, A. M., LOHSE, M. & USADEL, B. 2014. Trimmomatic: a flexible trimmer for

Illumina sequence data. *Bioinformatics,* 30**,** 2114-20.

BROCK, M. T., LUCAS, L. K., ANDERSON, N. A., RUBIN, M. J., CODY MARKELZ, R.,

COVINGTON, M. F., DEVISETTY, U. K., CHAPPLE, C., MALOOF, J. N. & WEINIG, C.

2016. Genetic architecture, biochemical underpinnings and ecological impact of floral UV

patterning. *Molecular ecology,* 25**,** 1122-1140.

BURKE, J. J. & UPCHURCH, D. R. 1989. Leaf temperature and transpirational control in

cotton. *Environmental and experimental botany,* 29**,** 487-492.

BYRNE, M. P. & O'GORMAN, P. A. 2018. Trends in continental temperature and humidity

directly linked to ocean warming. *Proc Natl Acad Sci U S A,* 115**,** 4863-4868.

CHITTKA, L., SHMIDA, A., TROJE, N. & MENZEL, R. 1994. Ultraviolet as a component of

flower reflections, and the colour perception of Hymenoptera. *Vision research,* 34**,** 1489-

1508.

DRAKE, J. E., TJOELKER, M. G., VÅRHAMMAR, A., MEDLYN, B. E., REICH, P. B.,

LEIGH, A., PFAUTSCH, S., BLACKMAN, C. J., LÓPEZ, R. & ASPINWALL, M. J. 2018.

Trees tolerate an extreme heatwave via sustained transpirational cooling and increased leaf

thermal tolerance. *Global change biology,* 24**,** 2390-2402.

GALEN, C. 2000. High and dry: drought stress, sex-allocation trade-offs, and selection on

flower size in the alpine wildflower *Polemonium viscosum* (Polemoniaceae). *Am Nat,* 156**,**

72-83.

GAUTIER, M. 2015. Genome-wide scan for adaptive divergence and association with

population-specific covariates. *Genetics,* 201**,** 1555-79.

GRIMM, D. G., ROQUEIRO, D., SALOME, P. A., KLEEBERGER, S., GRESHAKE, B., ZHU,

W., LIU, C., LIPPERT, C., STEGLE, O., SCHOLKOPF, B., WEIGEL, D. &

BORGWARDT, K. M. 2017. easyGWAS: a cloud-based platform for comparing the results

of genome-wide association studies. *Plant Cell,* 29**,** 5-19.

GRONQUIST, M., BEZZERIDES, A., ATTYGALLE, A., MEINWALD, J., EISNER, M. &

EISNER, T. 2001. Attractive and defensive functions of the ultraviolet pigments of a flower

(*Hypericum calycinum*). *Proceedings of the National Academy of Sciences,* 98**,** 13745-

13750.

HARBORNE, J. B. & SMITH, D. M. 1978. Anthochlors and other flavonoids as honey guides in

the Compositae. *Biochemical Systematics and Ecology,* 6**,** 287-291.

HEISER, C. B. & SMITH, D. M. 1969. *The North American sunflowers (Helianthus),* Durham,

N.C., Published for the Club by the Seeman Printery, Durham, N.C., USA.

HELLENS, R. P., EDWARDS, E. A., LEYLAND, N. R., BEAN, S. & MULLINEAUX, P. M.

2000. pGreen: a versatile and flexible binary Ti vector for Agrobacterium-mediated plant

transformation. *Plant Mol. Biol.,* 42**,** 819-32.

HERRERA, J. 2005. Flower size variation in *Rosmarinus officinalis*: individuals, populations

and habitats. *Annals of Botany,* 95**,** 431-437.

HIJMANS, R. J. 2020. Geographic data analysis and modeling. Available:

https://rspatial.org/raster/index.html

HOFFMANN, M., BREMER, M., SCHNEIDER, K., BURGER, F., STOLLE, E. & MORITZ,

G. 2003. Flower visitors in a natural population of *Arabidopsis thaliana*. *Plant Biology,* 5**,**

491-494.

HORTH, L., CAMPBELL, L. & BRAY, R. 2014. Wild bees preferentially visit *Rudbeckia*

flower heads with exaggerated ultraviolet absorbing floral guides. *Biology open,* 3**,** 221-230.

HUBNER, S., BERCOVICH, N., TODESCO, M., MANDEL, J. R., ODENHEIMER, J.,

ZIEGLER, E., LEE, J. S., BAUTE, G. J., OWENS, G. L., GRASSA, C. J., EBERT, D. P.,

OSTEVIK, K. L., MOYERS, B. T., YAKIMOWSKI, S., MASALIA, R. R., GAO, L.,

CALIC, I., BOWERS, J. E., KANE, N. C., SWANEVELDER, D. Z. H., KUBACH, T.,

MUNOS, S., LANGLADE, N. B., BURKE, J. M. & RIESEBERG, L. H. 2019. Sunflower

pan-genome analysis shows that hybridization altered gene content and disease resistance.

Nat. Plants, 5**,** 54-62.

HURD JR, P. D., LEBERGE, W. E. & LINSLEY, E. G. 1980. Principal sunflower bees of North

America with emphasis on the southwestern United States (Hymenoptera, Apoidea).

Smithsonian contributions to zoology.

JEFFREYS, H. 1961. *Theory of Probability*, Oxford, Clarendon Press.

KANG, H. M., SUL, J. H., SERVICE, S. K., ZAITLEN, N. A., KONG, S. Y., FREIMER, N. B.,

SABATTI, C. & ESKIN, E. 2010. Variance component model to account for sample

structure in genome-wide association studies. *Nat. Genet.,* 42**,** 348-54.

KLEPIKOVA, A. V., KASIANOV, A. S., GERASIMOV, E. S., LOGACHEVA, M. D. &

PENIN, A. A. 2016. A high resolution map of the *Arabidopsis thaliana* developmental

transcriptome based on RNA-seq profiling. *The Plant Journal,* 88**,** 1058-1070.

KORN, M., PETEREK, S., MOCK, H. P., HEYER, A. G. & HINCHA, D. K. 2008. Heterosis in

the freezing tolerance, and sugar and flavonoid contents of crosses between *Arabidopsis*

thaliana accessions of widely varying freezing tolerance. *Plant, cell & environment,* 31**,**

813-827.

KOSKI, M. H. & ASHMAN, T.-L. 2013. Quantitative variation, heritability, and trait

correlations for ultraviolet floral traits in *Argentina anserina* (Rosaceae): Implications for

floral evolution. *International Journal of Plant Sciences,* 174**,** 1109-1120.

KOSKI, M. H. & ASHMAN, T.-L. 2015. Floral pigmentation patterns provide an example of

Gloger's rule in plants. *Nature Plants,* 1**,** 1-5.

KOSKI, M. H. & ASHMAN, T. L. 2014. Dissecting pollinator responses to a ubiquitous

ultraviolet floral pattern in the wild. *Functional Ecology,* 28**,** 868-877.

KOSKI, M. H. & ASHMAN, T. L. 2016. Macroevolutionary patterns of ultraviolet floral

pigmentation explained by geography and associated bioclimatic factors. *New Phytologist,*

211**,** 708-718.

KOSKI, M. H., MACQUEEN, D. & ASHMAN, T. L. 2020. Floral pigmentation has responded

rapidly to global change in ozone and temperature. *Curr Biol,* 30**,** 4425-4431 e3.

LAMBRECHT, S. C. 2013. Floral water costs and size variation in the highly selfing

Leptosiphon bicolor (Polemoniaceae). *International Journal of Plant Sciences,* 174**,** 74-84.

LAMBRECHT, S. C. & DAWSON, T. E. 2007. Correlated variation of floral and leaf traits

along a moisture availability gradient. *Oecologia,* 151**,** 574-583.

LI, H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

*ArXiv***,** arXiv:1303.3997v2.

MANDEL, J. R., NAMBEESAN, S., BOWERS, J. E., MAREK, L. F., EBERT, D.,

RIESEBERG, L. H., KNAPP, S. J. & BURKE, J. M. 2013. Association mapping and the

genomic consequences of selection in sunflower. *PLoS Genet,* 9**,** e1003378.

MATVIENKO, M., KOZIK, A., FROENICKE, L., LAVELLE, D., MARTINEAU, B.,

PERROUD, B. & MICHELMORE, R. 2013. Consequences of normalizing transcriptomic

and genomic libraries of plant genomes using a duplex-specific nuclease and

tetramethylammonium chloride. *PLoS One,* 8**,** e55913.

MOYERS, B. T., OWENS, G. L., BAUTE, G. J. & RIESEBERG, L. H. 2017. The genetic

architecture of UV floral patterning in sunflower. *Annals of Botany,* 120**,** 39-50.

MURRAY, M. G. & THOMPSON, W. F. 1980. Rapid isolation of high molecular weight plant

DNA. *Nucleic Acids Res.,* 8**,** 4321-5.

NAKABAYASHI, R., MORI, T. & SAITO, K. 2014a. Alternation of flavonoid accumulation

under drought stress in *Arabidopsis thaliana*. *Plant Signal Behav,* 9**,** e29518.

NAKABAYASHI, R., YONEKURA-SAKAKIBARA, K., URANO, K., SUZUKI, M.,

YAMADA, Y., NISHIZAWA, T., MATSUDA, F., KOJIMA, M., SAKAKIBARA, H.,

SHINOZAKI, K., MICHAEL, A. J., TOHGE, T., YAMAZAKI, M. & SAITO, K. 2014b.

Enhancement of oxidative and drought tolerance in Arabidopsis by overaccumulation of

antioxidant flavonoids. *Plant J,* 77**,** 367-79.

NUNEZ, M., FORGAN, B. & ROY, C. 1994. Estimating ultraviolet radiation at the earth's

surface. *International Journal of Biometeorology,* 38**,** 5-17.

OWENS, G. L., TODESCO, M., BERCOVICH, N., LÉGARÉ, J. S., MITCHELL, N.,

WHITNEY, K. D. & RIESEBERG, L. H. 2021. Standing variation rather than recent

adaptive introgression likely underlies differentiation of the *texanus* subspecies of

Helianthus annuus. *Molecular Ecology*.

PEACH, K., LIU, J. W. & MAZER, S. J. 2020. Climate predicts UV floral pattern size,

anthocyanin concentration, and pollen performance in *Clarkia unguiculata*. *Front Plant Sci,*

11**,** 847.

POLLASTRI, S. & TATTINI, M. 2011. Flavonols: old compounds for old roles. *Annals of*

Botany, 108**,** 1225-1233.

POPLIN, R., RUANO-RUBIO, V., DEPRISTO, M. A., FENNELL, T. J., CARNEIRO, M. O.,

VAN DER AUWERA, G. A., KLING, D. E., GAUTHIER, L. D., LEVY-MOONSHINE,

A., ROAZEN, D., SHAKIR, K., THIBAULT, J., CHANDRAN, S., WHELAN, C., LEK,

M., GABRIEL, S., DALY, M. J., NEALE, B., MACARTHUR, D. G. & BANKS, E. 2017.

Scaling accurate genetic variant discovery to tens of thousands of samples. *BioRxiv*.

R CORE TEAM. 2020. R: a language and environment for statistical computing. Available:

https://www.r-project.org/.

RAE, J. M. & VAMOSI, J. C. 2013. Ultraviolet reflectance mediates pollinator visitation in

Mimulus guttatus. *Plant Species Biology,* 28**,** 177-184.

RIESEBERG, L. H. & SCHILLING, E. E. 1985. Floral flavonoids and ultraviolet patterns in

Viguiera (Compositae). *American journal of botany,* 72**,** 999-1004.

ROHLAND, N. & REICH, D. 2012. Cost-effective, high-throughput DNA sequencing libraries

for multiplexed target capture. *Genome Res.,* 22**,** 939-46.

ROWAN, B. A., PATEL, V., WEIGEL, D. & SCHNEEBERGER, K. 2015. Rapid and

inexpensive whole-genome genotyping-by-sequencing for crossover localization and fine-

scale genetic mapping. *G3 (Bethesda),* 5**,** 385-98.

SCHINDELIN, J., ARGANDA-CARRERAS, I., FRISE, E., KAYNIG, V., LONGAIR, M.,

PIETZSCH, T., PREIBISCH, S., RUEDEN, C., SAALFELD, S. & SCHMID, B. 2012. Fiji:

an open-source platform for biological-image analysis. *Nat. Methods,* 9**,** 676.

SCHLANGEN, K., MIOSIC, S., CASTRO, A., FREUDMANN, K., LUCZKIEWICZ, M.,

VITZTHUM, F., SCHWAB, W., GAMSJÄGER, S., MUSSO, M. & HALBWIRTH, H.

2009. Formation of UV-honey guides in *Rudbeckia hirta*. *Phytochemistry,* 70**,** 889-898.

SCHNEIDER, C. A., RASBAND, W. S. & ELICEIRI, K. W. 2012. NIH Image to ImageJ: 25

years of image analysis. *Nat. Methods,* 9**,** 671.

SCHNEITER, A. & MILLER, J. 1981. Description of sunflower growth stages. *Crop Science,*

21**,** 901-903.

SCHULZ, E., TOHGE, T., ZUTHER, E., FERNIE, A. R. & HINCHA, D. K. 2015. Natural

variation in flavonol and anthocyanin metabolism during cold acclimation in *Arabidopsis*

thaliana accessions. *Plant, cell & environment,* 38**,** 1658-1672.

SEDLAZECK, F. J., RESCHENEDER, P. & VON HAESELER, A. 2013. NextGenMap: fast

and accurate read mapping in highly polymorphic genomes. *Bioinformatics,* 29**,** 2790-1.

SHAGINA, I., BOGDANOVA, E., MAMEDOV, I. Z., LEBEDEV, Y., LUKYANOV, S. &

SHAGIN, D. 2010. Normalization of genomic DNA using duplex-specific nuclease.

Biotechniques, 48**,** 455-459.

SHEEHAN, H., MOSER, M., KLAHRE, U., ESFELD, K., DELL'OLIVO, A., MANDEL, T.,

METZGER, S., VANDENBUSSCHE, M., FREITAS, L. & KUHLEMEIER, C. 2016.

MYB-FL controls gain and loss of floral UV absorbance, a key trait affecting pollinator

preference and reproductive isolation. *Nature genetics,* 48**,** 159.

STAPLETON, A. E. 1992. Ultraviolet radiation and plants: burning questions. *The Plant Cell,* 4**,**

1353.

STATON, S. E. & LÁZARO-GUEVARA, J. M. 2020. *Sunflower genome database* [Online].

Available: https://sunflowergenome.org/annotations/ [Accessed].

STEPHENS, J. D., ROGERS, W. L., MASON, C. M., DONOVAN, L. A. & MALMBERG, R.

L. 2015. Species tree estimation of diploid *Helianthus* (Asteraceae) using target enrichment.

American Journal of Botany, 102**,** 910-920.

STOFFEL, K., VAN LEEUWEN, H., KOZIK, A., CALDWELL, D., ASHRAFI, H., CUI, X.,

TAN, X., HILL, T., REYES-CHIN-WO, S. & TRUCO, M.-J. 2012. Development and

application of a 6.5 million feature Affymetrix Genechip® for massively parallel discovery

of single position polymorphisms in lettuce (*Lactuca* spp.). *BMC genomics,* 13**,** 1-17.

STRACKE, R., ISHIHARA, H., HUEP, G., BARSCH, A., MEHRTENS, F., NIEHAUS, K. &

WEISSHAAR, B. 2007. Differential regulation of closely related R2R3-MYB transcription

factors controls flavonol accumulation in different parts of the *Arabidopsis thaliana*

seedling. *Plant J,* 50**,** 660-77.

STRACKE, R., JAHNS, O., KECK, M., TOHGE, T., NIEHAUS, K., FERNIE, A. R. &

WEISSHAAR, B. 2010. Analysis of PRODUCTION OF FLAVONOL GLYCOSIDES-

dependent flavonol glycoside accumulation in *Arabidopsis thaliana* plants reveals MYB11-,

MYB12- and MYB111-independent flavonol glycoside accumulation. *New Phytol,* 188**,**

985-1000.

STRAUSS, S. Y. & WHITTALL, J. B. 2006. Non-pollinator agents of selection on floral traits.

*Ecology and evolution of flowers***,** 120-138.

THOMPSON, W., MEINWALD, J., ANESHANSLEY, D. & EISNER, T. 1972. Flavonols:

pigments responsible for ultraviolet absorption in nectar guide of flower. *Science,* 177**,** 528-

530.

TODESCO, M., OWENS, G. L., BERCOVICH, N., LEGARE, J. S., SOUDI, S., BURGE, D. O.,

HUANG, K., OSTEVIK, K. L., DRUMMOND, E. B. M., IMEROVSKI, I., LANDE, K.,

PASCUAL-ROBLES, M. A., NANAVATI, M., JAHANI, M., CHEUNG, W., STATON, S.

E., MUNOS, S., NIELSEN, R., DONOVAN, L. A., BURKE, J. M., YEAMAN, S. &

RIESEBERG, L. H. 2020. Massive haplotypes underlie ecotypic differentiation in

sunflowers. *Nature,* 584**,** 602-607.

TOVÉE, M. J. 1995. Ultra-violet photoreceptors in the animal kingdom: their distribution and

function. *Trends in ecology & evolution,* 10**,** 455-460.

UNICODE.ORG. 2020. *Full emoji list, v13.0* [Online]. Available:

https://unicode.org/emoji/charts/full-emoji-list.html [Accessed].

WANG, T., HAMANN, A., SPITTLEHOUSE, D. & CARROLL, C. 2016. Locally downscaled

and spatially customizable climate data for historical and future periods for North America.

PloS one, 11**,** e0156720.

WARNES, G. R., BOLKER, B., BONEBAKKER, L., GENTLEMAN, R., HUBER, W., LIAW,

A., LUMLEY, T., MAECHLER, M., MAGNUSSON, A. & MOELLER, S. 2009. gplots:

various R programming tools for plotting data. Available: https://CRAN.R-

project.org/package=gplots

WEIGEL, D. & GLAZEBROOK, J. 2002. *Arabidopsis: A laboratory manual,* Cold Spring

Harbor, N.Y., USA, CSHL Press.

WOJTASZEK, J. & MAIER, C. 2014. A microscopic review of the sunflower and honeybee

mutualistic relationship. *Int. J. AgriSci,* 4**,** 272-282.

ZENG, J., ZOU, Y., BAI, J. & ZHENG, H. 2002. Preparation of total DNA from recalcitrant

plant taxa. *Acta Bot. Sin.,* 44**,** 694-697.

Figure supplements

Figure 1 – figure supplement 1: Floral UV patterns in wild sunflower species and cultivated

sunflower.

a, Visible and UV images of inflorescences from five wild sunflower species, and ligules of six

cultivated sunflower lines. Variation in floral UV patterns was found within all these species.

Scale bar = 1 cm. **b,** UV images of inflorescences from twelve wild sunflower species and for the

outgroup *Phoebanthus tenuifolius*. Images are not to scale. **c,** Species tree for 31 sunflower

species and *P. tenuifolius*, modified from (Stephens et al., 2015). The size of the black dots to the

right of each species name is proportional to the average size of bullseye patterns measured for

that species or subspecies (Figure 1 – source data 1). For the species in **a**, bullseye values are

averages for ≥42 individuals (see also Figure 1 – figure supplement 2). For the species in **b**,

bullseye values are for single individuals or averages for up to three individuals. Two taxa in the

original species tree, *H. petiolaris* and *H. neglectus*, were renamed to *H. petiolaris ssp. petiolaris*

and *H. petiolaris ssp. fallax* to reflect the current understanding of their identities.

Figure 1 – figure supplement 2: LUVp variation in wild sunflower species and cultivated

sunflower.

LUVp values for individuals of four wild sunflower species and of the cultivated sunflower

association mapping (SAM) population, and allele frequencies at the Chr15_LUVp SNP. ANN =

wild *H. annuus* (n = 1589 from 110 populations), PET = *H. petiolaris* (n = 351 individuals from

40 populations), ARG = *H. argophyllus* (n = 105 individuals from 27 populations), NIV = *H.*

niveus (n = 42 individuals from 9 populations), SAM = cultivated *H. annuus* (n = 275

individuals). Letters identify groups that are significantly different for $p < 0.001$ (one-way

ANOVA with post-hoc Tukey HSD test, F = 247, df = 4). Exact p-values for pairwise

comparisons are reported in Figure 1 – source data 2.

Figure 2 – figure supplement 1: LUVp GWAS in *H. petiolaris fallax*.

LUVp GWAS in *H. petiolaris fallax* (n = 193 individuals). The red line represents 5%

Bonferroni-corrected significance. GWAs were calculated using two-sided mixed models. Only

positions with -log10 p-value > 2 are plotted.

Figure 2 – figure supplement 2: Floral UV patterns in the parental lines of F$_2$ populations.

UV images of ligules of the parental lines for the F$_2$ populations shown in Figure 2e, and their F$_1$

progeny. A pair of F$_1$ plants was selected and crossed for each population to generate the F$_2$

progeny.

Figure 2 – figure supplement 3: LUVp GWAS in unfiltered *H. annuus* datasets.

LUVp GWAS *H. annuus* using an un-filtered variants dataset in a 100 kbp region surrounding

HaMYB111 (*n* = 563 individuals). Relaxing variant filtering parameters, to capture more of the

polymorphisms at the *HaMYB111* locus, resulted in an almost 50-fold increase in the number of

variants in this region, from 142 to 6949. Regions in which no SNPs are reported contain highly

repetitive sequences, and were masked before reads mapping. As re-mapping to the improved

HA412v2 reference assembly of the complete *H annuus* set of >222M unfiltered variants would

have been computationally intensive, positions are shown based on the original XRQv1

reference assembly. The red line represents 5% Bonferroni-corrected significance. GWAs were

calculated using two-sided mixed models.

Figure 3 – figure supplement 1: Floral UV patterns in Arabidopsis lines.

Visible and UV images of Arabidopsis petals. Col-0 = wild type Arabidopsis. Scale bar = 1 mm.

Figure 3 – figure supplement 2: Stages of ligule development in *H. petiolaris*.

Figure 4 – figure supplement 1: Mean pollinator visits in the 2019 field experiment divided

by category of pollinators.

"Bees" were exclusively honey bees; "Bumblebees" included several *Bombus* species; "Others"

were mostly *Megachile* bees (n = 1390 pollinator visits). Pollinators were overwhelmingly

bumblebees in the 2017 field experiment.

Figure 4 – figure supplement 2: Spearman correlation heatmap for LUVp and environmental variables in wild *H. annuus* populations.

Source data for this figure are reported in Figure 1 – source data 1.

Figure 4 – figure supplement 3: Inflorescence temperature time series.

Infrared images of East-facing inflorescences of sunflowers with large (LUVp = 1) or small

(LUVp < 0.15) floral UV patterns taken in the summer of 2017. No additional difference was

observed in pictures taken more than three hours after sunrise. While no difference in

temperature was observed in ligules, the centre (disk) of the inflorescence was consistently

warmer in plants with small LUVp. However, this effect is independent of ligule UV patterns,

since it persists in inflorescences in which ligules were removed (right-most column). Pollinator

visits were severely reduced for inflorescences with ligules removed. Bumblebees can be seen on

the disk of inflorescences with large LUVp in the leftmost column of pictures, at 5-10 minutes

and 2 hours. Temperatures values outside of the 10-40 ℃ interval are shown in beige (<10 ℃) or

green (>40 ℃).

Figure 4 – figure supplement 4: Correlations between LUVp and environmental variables

in *H. petiolaris*.

a, Correlation between average LUVp for different populations of *H. petiolaris* and summer UV

radiation ($R^2 = 0.11$, $P = 0.02$), **b,** summer average temperature ($R^2 = 0.69$, $P = 10^{-11}$), or **c,**

summer relative humidity ($R^2 = 0.47$, $P = 4.4 \times 10^{-7}$). Grey areas represent the 95% confidence

interval. **d,** Spearman correlation heatmap for LUVp and environmental variables. Source data

for this figure are reported in Figure 1 – source data 1.

Source data

Figure 1 - source data 1 (Figure 1 - source data 1.xlsx): Population used in this study, average

LUVp values and environmental variables.

Figure 1 - source data 2 (Figure 1 - source data 2.xlsx): Individuals used in this study, LUVp

values and Chr15_LUVp SNP genotypes.

Figure 2 - source data 1 (Figure 2 - source data 1.xlsx): LUVp values and Chr15_LUVp SNP

genotypes for F_2.

Figure 3 - source data 1 (Figure 3 - source data 1.xlsx): Flavonols in methanolic extractions of

sunflower ligules and Arabidopsis petals.

Figure 4 - source data 1 (Figure 4 - source data 1.xlsx): Pollinator experiment data.

1000

1001   **Figure 4 - source data 2 (Figure 4 - source data 2.xlsx):** Ligules and leaves desiccation

1002   experiment data.

1003

1004   **Figure 4 - source data 3 (Figure 4 - source data 3.xlsx):** GEA results for the HaMYB111

1005   region.