Title

Microbiota responses to different prebiotics are conserved within individuals and associated with

habitual fiber intake

Authors

Zachary C. Holmes[1]: MSRB3, 3 Genome Court, Durham NC 27705; zachary.holmes@duke.edu

Max M. Villa[1,2]: MSRB3, 3 Genome Court, Durham NC 27705; max.villa.m@gmail.com

Heather K. Durand[1,2]: MSRB3, 3 Genome Court, Durham NC 27705; heather.durand@duke.edu

Sharon Jiang[1,2]: MSRB3, 3 Genome Court, Durham NC 27705; sharon.jiang@duke.edu

Eric P. Dallow[1,2]: MSRB3, 3 Genome Court, Durham NC 27705; dalloweric@gmail.com

Brianna L. Petrone[1,3]: MSRB3, 3 Genome Court, Durham NC 27705; brianna.petrone@duke.edu

Justin D. Silverman[4,5,6]: Westgate Bldg, University Park, PA 16802; JustinSilverman@psu.edu

Pao-Hwa Lin[7,8]: Stedman Nutrition Ctr, 3475 Erwin Rd, Durham NC 27705;
pao.hwa.lin@duke.edu

Lawrence A. David[1,2,9,*]: MSRB3, 3 Genome Court, Durham NC 27705;
lawrence.david@duke.edu

Affiliations

[1]Department of Molecular Genetics and Microbiology, Duke University, Durham, NC

[2]Center for Genomic and Computational Biology, Duke University, Durham, NC

[3]Medical Scientist Training Program, Duke University, Durham, NC

[4]College of Information Science and Technology, Penn State University, University Park, PA

[5]Department of Medicine, Penn State University, Hershey, PA

[6]Institute for Computational and Data Science, Penn State University, University Park, PA

[7]Duke Molecular Physiology Institute, Duke University, Durham, NC

[8]Department of Medicine, Duke University Medical Center, Durham, NC

[9]Program in Computational Biology and Bioinformatics, Duke University, Durham, NC

[*]Corresponding author

Abstract

Background: Short-chain fatty acids (SCFAs) derived from gut bacteria are associated with

protective roles in diseases ranging from obesity to colorectal cancers. Intake of microbially

accessible dietary fibers (prebiotics) lead to varying effects on SCFA production in human

studies, and gut microbial responses to nutritional interventions vary by individual. It is therefore

possible that prebiotic therapies will require customizing to individuals.

Results: Here, we explored prebiotic personalization by conducting a three-way crossover study

of three prebiotic treatments in healthy adults. We found that within individuals, metabolic

responses were correlated across the three prebiotics. Individual identity, rather than prebiotic

choice, was also the major determinant of SCFA response. Across individuals, prebiotic response

was inversely related to basal fecal SCFA concentration, which, in turn, was associated with

habitual fiber intake. Experimental measures of gut microbial SCFA production for each

participant also negatively correlated with fiber consumption, supporting a model in which

individuals' gut microbiota are limited in their overall capacity to produce fecal SCFAs from

fiber.

Conclusions: Our findings support developing personalized prebiotic regimens that focus on

selecting individuals who stand to benefit, and that such individuals are likely to be deficient in

fiber intake.

Keywords

Microbiome; prebiotic; short-chain fatty acids; diet; personalized nutrition; fiber

Background

A key beneficial role of the human gut microbiota is producing the short-chain fatty acids (SCFAs) acetate, propionate, and butyrate, which are metabolic end products of anaerobic fermentation (1, 2). These SCFA, most notably among them, butyrate, have been shown to promote gastrointestinal (GI) barrier integrity, lower inflammation, increase resistance to pathogen invasion, and contribute positively to enteroendocrine signaling and energy homeostasis (3-9). Augmenting these functions may explain the protective role of SCFAs in diseases ranging from obesity to inflammatory bowel diseases (3, 10-20).

The substrates for SCFA production in the human gut are predominately fibers, which are dietary carbohydrates inaccessible to human enzymes and transporters (21). On average, American adults consume only 21 to 38% of the fiber that is recommended by the USDA (22). This lack of fiber is implicated in the etiology of Western diseases (15, 23, 24), which may in part be tied to the effects of low fiber consumption on microbial SCFA production and SCFA signaling (15, 25-29). To address fiber deficiencies, dietary supplements known as prebiotics have been formulated using specific dietary fibers that are microbially fermentable but host-inaccessible. Prebiotics can increase production of SCFAs by the gut microbiota (30-33), and have emerged as a promising microbiota-targeting approach for promoting health.

A key challenge of using prebiotics to promote SCFA production is heterogeneity in treatment response. Many distinct prebiotic formulations exist (34, 35), and different prebiotics have been associated with diverging patterns of SCFA production (30, 32). Prebiotic trials have also often revealed an increase in SCFA at the population level following treatment, but inconsistent results at the level of individual (30-33). Still, prebiotic trials tend not to examine how responses to different prebiotics vary within individuals, and have to date often profiled

80 single prebiotics (31, 33, 36, 37), or tested different prebiotics in separate cohorts (30, 32, 38).

81 More rare crossover studies have taken place, but have either not investigated individual

82 variation in prebiotic response (39), or solely utilized *in vitro* assays of prebiotic response (40,

83 41). If consistent evidence does emerge that responses to different prebiotics vary within people,

84 then effective prebiotic therapies will face the challenging task of personalizing prebiotic choices

85 for each recipient. Alternatively, if responses to different prebiotics are conserved within

86 individuals, effective prebiotic therapy could primarily focus on selecting individuals for

87 treatment. Evidence supporting this latter scenario include studies showing that individual

88 identity is a stronger predictor of SCFA concentrations than dietary intervention (42), and that

89 individual characteristics like habitual food choice can predict changes to microbiota community

90 composition following prebiotic supplementation (37).

91  To directly investigate the interaction of prebiotic and individual variation in prebiotic

92 response, we report here the results of a randomized, fully balanced, crossover trial where a

93 single set of individuals was treated with multiple prebiotics. Our primary lines of inquiry were

94 to assess the relative contribution of individual and prebiotic differences to SCFA responses to

95 supplementation, with an emphasis on butyrate production. Secondary analyses included

96 determining what host and microbiological factors shaped variation in prebiotic response.

98 **Methods**

99 Participant recruitment:

100  This study was conducted at Duke University in Durham, NC, USA, and was registered

101 with ClinicalTrials.gov (Identifier: NCT03595306). All recruitment and study procedures

102 involving human subjects were approved by the Duke University Institutional Review Board

(Pro00087214). Study participants were recruited by flyers, email list-servs, and word-of-mouth

at Duke University Campus. Individuals with a self-reported history of irritable bowel syndrome,

inflammatory bowel disease, type-2 diabetes, kidney disease, intestinal obstruction, or colorectal

cancer were excluded. Participants were also excluded if they had taken oral antibiotics within

the past 6 months, had any known food allergies, reported dietary intolerances of any kind, or

were pregnant or breastfeeding. If individuals identified as female, their urine was subjected to a

pregnancy test (Quidel QuickView One-Step hCG Urine Test) to minimize the risk that prebiotic

treatment would impact newborn or infant health. Participants who tested positive were deemed

ineligible for participation in the study. Inclusion criteria were that participants were between

the ages of 18 and 70, and able to provide stool samples at no risk to themselves. Fiber

consumption, prebiotic/probiotic usage, and dietary habits were not used as inclusion/exclusion

criteria, but participants were asked to report via surveys as part of the study requirements.

During this period, 81 individuals set up a phone interview to determine eligibility and interest to

participate in the study. Of these, 41 completed electronic informed consent to the study and

were subsequently enrolled. Ultimately, 8 subjects were lost to follow-up, 2 withdrew, and 3

were omitted from some or all analyses due to inability to collect sufficient usable sample,

resulting in 28 subjects included in all final data analyses.

Study design:

The three-period crossover longitudinal study was conducted from May of 2018 to March

of 2019. The crossover design allowed investigation of three prebiotic interventions while also

allowing participants to serve as their own controls (Figure 1). There were six arms of the study,

to cover all possible orders of prebiotic consumption and to eliminate the need to account for

uneven carry-over effects. The study consisted of an initial study visit, followed by a 6-week

study period with alternating weeks of baseline (no prebiotic consumption) and intervention

(prebiotic consumption). Participants were added sequentially to an arm group as they enrolled;

meaning the first participant was enrolled in Arm 1, the second to Arm 2 and so on. Participants

were blinded to the prebiotics that they were consuming until the end of the study. The prebiotics

were taken twice a day, Monday to Friday during the intervention weeks. Monday was a half-

dose, to help allow participants to adjust to the increase in fiber. Following consent, participants

had an initial study visit, to have their baseline physiology assessed by height, weight, and blood

pressure measurement, as well as provide a urine sample and receive the study kit and materials.

Participants were asked to self-collect stool at home three times a week each week during the

study period. Participants also completed 4 diet surveys throughout the study; one instance of the

Diet History Questionnaire Version 3 (DHQ3) (43), covering the one month period prior to the

study, and three instances of the Automated Self-Administered 24-hour (ASA24) Dietary

Assessment Tool, version 2017, developed by the National Cancer Institute, Bethesda, MD (44).

Two ASA24's were completed during the week, and one on a weekend. The DHQ3 was

completed prior to the 6-week study period. Additionally, participants completed online-surveys

about their experiences with the prebiotics following the intervention weeks.

Prebiotic supplementation:

Three prebiotics were tested in this study: inulin (manufactured by Now Foods,

Bloomingdale, IL), wheat dextrin (Benefiber, manufactured by GSK Consumer Healthcare,

Warren, NJ), and galactooligosaccharides (GOS) (Bimuno, manufactured by Clasado Limited,

Shinfield, UK). In a clean non-laboratory space, prebiotics were weighed into individual doses

and packaged into pre-labeled food-safe sealable plastic bags. Each participant received all three

prebiotics throughout the study; inulin at 9g / day, wheat dextrin at 9g / day, and

galactooligosaccharides (GOS) at 3.6g / day. These doses correspond to the recommended daily

intake on the product label. In order to further minimize the possibility of discomfort associated

with consuming prebiotics, the first day of dosing for each prebiotic was a half dose to allow

acclimation (4.5g for inulin and dextrin, 1.8g for GOS). Participants consumed one of the three

prebiotics at a time, twice a day, for a duration of 5 days, followed by a week of washout. The

order was determined by their assigned study arm. Participants were instructed to consume the

prebiotics by mixing the powders in any kind of beverage (such as water) or food (such as

applesauce), so long as it was cold or room temperature.

Stool collection:

At the pre-study visit, participants were provided with a standard urine specimen collection cup

and shown to a bathroom to collect the sample. Urine samples were also sent to LabCorp in a BD

Vacutainer UA tube to measure metabolic variables, including routine urinalysis, sodium,

potassium, urea nitrogen, and creatinine. Consented participants were asked to self-collect stool

three times a week at home using sampling kits provided to them. Participants were asked to

immediately store samples in their personal freezers (-20C). Participants were asked to bring

stored samples to the David Lab freezers on a weekly basis. Stool samples were transferred and

stored at -80C until thawed for DNA extraction and other sample processing.

In vitro fermentation:

In vitro fermentation of stool microbiota was conducted following protocols previously

published (40). In brief, a 10% w/v fecal slurry was made using thawed week-1 stool and

anaerobic 1X PBS (pH 7.0 ± 0.1). This slurry was filtered through 0.33mm pore size mesh, then

mixed 1:1 with a 1% w/v solution of prebiotic and incubated in duplicate in a 24-well cell culture

plate for 24hr. Three prebiotic treatments were used, corresponding to the three prebiotics used

in the *in vivo* portion of this study: inulin (Now Foods Inulin Powder, part #2944),

galactooligosaccharides (GOS; Bimuno Powder), and wheat dextrin (Benefiber Original). The

resulting fermentation conditions where therefore 5% fecal slurry with 0.5% prebiotic (w/v) and

occurred at 37°C for 24 hours. Following fermentation, a 1mL aliquot was taken from each well

for SCFA quantification.

Quantification of SCFA:

Quantification of SCFA by gas chromatograph flame ionization detection (GCFID) was

performed following protocols previously published (40). In brief, a 1mL aliquot of either 10%

fecal slurry in PBS or the fermentation vessel contents was obtained. To this, 50 μL of 6N HCl

was added to acidify the solution to a pH below 3. The mixture was vortexed, centrifuged at

14,000rcf for 5 minutes at 4°C to remove particles. Avoiding the pellet, 750 μL of this supernatant

was passed through a 0.22μm spin column filter. The resulting filtrate was then transferred to a

glass autosampler vial (VWR part #66009-882).

Filtrates were analyzed on an Agilent 7890b gas chromatograph (GC) equipped with a

flame-ionization detector (FID) and an Agilent HP-FFAP free fatty-acid column (25m x .2mm id

x .3μm film). Acetate, propionate, isobutyrate, butyrate, isovalerate, and valerate were identified

and quantified in each sample by comparing to an 8-point standard curve that encompassed the

sample concentration range. Standards contained 0.1mM, 0.2mM, 0.5mM, 1mM, 2mM, 4mM,

8mM, and 16mM concentrations of each SCFA.

DNA extraction, PCR amplification, and sequencing:

We performed 16S rRNA gene amplicon sequencing on human stool samples to determine

microbiota community composition following protocols previously published (40). DNA was

extracted from frozen fecal samples with the Qiagen DNeasy PowerSoil DNA extraction kit (ID

12888-100). Amplicon sequencing was performed using custom barcoded primers targeting the

V4 region of the 16S gene (45), using published protocols (45-47). The sequencing library was

diluted to a 5pM concentration and sequenced using an Illumina MiniSeq and a MiniSeq Mid

Output Kit (FC420-1004) with paired-end 150bp reads.

Identifying sequence variant and taxonomy assignment:

We used an analysis pipeline with DADA2 (48) to identify and quantify sequence

variants, as previously published (40, 49). To prepare data for denoising with DADA2, 16S

rRNA primer sequences were trimmed from paired sequencing reads using Trimmomatic v0.36

without quality filtering (50). Barcodes corresponding to reads that were dropped during

trimming were removed using a custom python script. Reads were demultiplexed without quality

filtering using python scripts provided with Qiime v1.9 (51). Bases between positions 10 and 150

were retained for the forward reads and between positions 0 and 140 were retained for the

reverse reads. This trimming, as well as minimal quality filtering of the demultiplexed reads was

performed using the function fastqPairedFilter provided with the DADA2 R package (v1.8.0).

Sequence variants were inferred by DADA2 independently for the forward and reverse reads of

each of the two sequencing runs using error profiles learned from all 20 samples. Forward and

reverse reads were merged. Bimeras were removed using the function removeBimeraDenovo

with default settings. Taxonomy was assigned using the function assignTaxonomy from

DADA2, trained using version 123 of the Silva database.

Results

Prebiotic supplementation affected SCFA production at the cohort-level. We enrolled

participants into a six-week, three-period prebiotic intervention study (Figure 1). We used a

crossover design to compare multiple prebiotics across a subject pool. We selected inulin, wheat

dextrin, and galactooligosaccharides (GOS) as our prebiotics, as these are commonly consumed

(52), commercially available dietary supplements, and are monomerically distinct. Our crossover

was standardized in series and period, with each prebiotic occurring the same number of times

within each series (once) and study period (twice). The design was also balanced with respect to

first-order carry-over effects, meaning we shuffled the order in which prebiotics were

administered so as to minimize the interactive effects between any two pairs of consecutive

prebiotics. Each study period consisted of an intervention-free baseline week and a prebiotic

intervention week. Stool samples were collected on Wednesday, Thursday, and Friday of each of

the six study weeks (baselines and intervention). Of the 41 participants who initially enrolled, 2

withdrew, 3 were unable to collect at least one usable sample (i.e., insufficient volume or

inadequate labeling) per week, and 8 were lost to follow-up. We focused our subsequent analyses

on the remaining 28 participants who completed the study. These participants in total collected

413 stool samples (mean=2.46 stool samples per person per week).

We first sought to examine differences in microbiome response during each of the

different prebiotic treatments. Analyses of gut microbial community composition did not reveal

associations with inulin, GOS, or wheat dextrin treatment (p > 0.05, PERMANOVA using

unweighted UniFrac distance at species level). In addition, no species were statistically

differentially abundant during any prebiotic intervention and its corresponding prebiotic-free

baseline (p > 0.05, Benjamini-Hochberg corrected Wilcoxon Rank Sum tests using the R

package ALDEx2). While previous studies have observed changes in community composition

following prebiotic administration, such effects have been observed in a subset of tested

prebiotics (30, 32), and with different dosing schedules (30-32, 37).

Still, in contrast to the lack of a coherent 16S rRNA shifts among participants, we found

the three prebiotics varied according to resulting fecal SCFA profiles relative to baseline. GOS

was associated with decreased total stool SCFA concentrations (Table 1, Figure S1C, effect size

= -0.071, p = 0.038, GLMM), an outcome that has been previously observed (53). Differences

within individual SCFAs were also apparent when analyzing absolute fecal SCFA concentrations

(Table 1). Dextrin increased propionate concentration (effect size = 0.074, p = 0.038, GLMM),

and GOS decreased both propionate and butyrate concentrations (effect size = -0.087, p = 0.015;

effect size = -0.14, p = 0.002, GLMMs, respectively). Still, changes in total and individual SCFA

concentrations may be caused by effects of stool hydration, which can vary up to three-fold and

directly impacts measures of SCFA concentration in feces (54, 55). To assess the relative

production of each SCFA, we therefore normalized levels of individual SCFAs to account for

differences in stool hydration. Our resulting analysis found that inulin supplementation increased

proportional butyrate concentrations (butyrate concentration / total SCFA concentration) in

participants' stool (effect size = 0.082, p = 0.007, GLMM; Figure 2, Table 1). These findings are

consistent with prior reports that inulin supplementation may increase butyrogenesis (56-58).

Our analyses also suggested GOS and wheat dextrin decreased proportional butyrate

concentration in participants' stool (GOS; effect size = -0.074, p = 0.012. Dextrin; effect size = -

0.071, p = 0.015, GLMM; Figure 2, Table 1). These findings align with prior trials that found

both GOS (53, 59, 60) and dextrin (61, 62) increases acetate production over more reduced

SCFAs, including butyrate and propionate, potentially by shunting nutrients away from butyrate

production. No prebiotic altered proportional acetate or proportional propionate concentration at

the cohort level (p > 0.05, GLMM; Figures S1A and S1B, Table 1).

Butyrogenic responses are more strongly predicted by individual identity than by prebiotic.

We next investigated how prebiotic responses varied among individuals. We focused our

analysis on prebiotic impacts on butyrate production because it was the sole SCFA that was

altered at the cohort-level, and it is also the microbially derived SCFA most commonly

implicated in enteric health (2, 6-8, 63-66). We also focused on proportional butyrate

concentrations to avoid the effects of stool hydration on our analysis and to be consistent with

previous individual-level analyses of fecal SCFAs (32, 42). In line with prior studies of prebiotic

response, (40) statistical modeling yielded evidence for individual-specific butyrogenic

responses to each of the prebiotics (that is, an interaction between individual and prebiotic terms

in a GLMM). Nonetheless, further analyses suggested that the strongest determinant of

proportional butyrate responses was shaped by individual identity: individual-specific effects

(pseudo-R^2 = 0.39, p < 0.0001; GLM F-test) were much stronger than prebiotic-specific effects

(pseudo-R^2 = 0.05, p = 0.001; GLM F-test). Additionally, individuals who exhibited the largest

response in proportional stool butyrate concentration on a given prebiotic were more likely to

exhibit larger responses on the other two prebiotics ($\rho = 0.49$, 0.37, and 0.28, $p = 0.008$, 0.05,

and 0.15, respectively for inulin vs dextrin, dextrin vs GOS, and inulin vs GOS, Spearman

correlations; Figure 3). This pattern was apparent in spite of correcting for baseline differences in

SCFA production between individuals, which were also consistent over time and between

baseline and treatment weeks.

Individuals consistently varied according to gut metabolic state, microbiota community

composition, and metabolic potential. Why might individual-specific effects be stronger than

prebiotic-specific effects in our cohort? We observed that individuals were idiosyncratic in their

SCFA profiles over time. Both proportional butyrate concentration and butyrate concentration

were correlated within individuals between the first and last baseline weeks ($\rho = 0.49$, $p < 0.007$;

and $\rho = 0.63$, $p = 0.0003$, respectively, Spearman correlations; Figure S2). Variation between

individuals' SCFA profiles could also be linked to inter-individual microbiota variation, a well-

known factor in prebiotic response (42). When we divided participants into 'high' (> median)

and 'low' (<= median) fecal butyrate groups, we observed a significant effect of this grouping on

microbiota community composition ($R^2 = 0.075$, $p = 0.0022$, PERMANOVA using unweighted

UniFrac distance; Figure S3). We identified three specific genera that were differentially

abundant between the groups: *Anaerofilum* and *Ruminiclostridium_5* were decreased in the high

butyrate group, and *Lachnospira* was increased in the high butyrate group ($p = 0.009$, .043, and

0.030, respectively; Benjamini-Hochberg corrected Wilcoxon Rank Sum tests; Figure S3). We

additionally observed a relationship between baseline microbiota community structure and

average response to all three prebiotics (Δ%butyrate). After dividing participants into

'responders' and 'non-responders', based on whether average Δ%butyrate was positive or

negative, we found a significant effect of this grouping on microbiota community composition

($R^2 = 0.059$, p = 0.028, PERMANOVA using unweighted UniFrac distance; Figure S4). Thus,

individuals' starting microbiome composition was associated with prebiotic treatment outcome.

Intake patterns of dietary fiber negatively correlated with response to prebiotic

intervention.

Next, we sought to understand what host factors might influence individuals' microbiomes and

ultimately shape their butyrogenic responses across the tested prebiotics. We focused on fiber

intake, which has been shown to shape microbiota community composition and metabolic output

(67, 68). We measured fiber consumption using a dietary questionnaire spanning the month

preceding the study (43), and we compared fiber intakes to each individual's average response to

the three prebiotic interventions (average Δ%butyrate). Despite prior studies suggesting that

plant or fiber intake would augment gut microbial metabolic capacity (37, 68-70), our

comparison revealed that dietary fiber consumption was negatively correlated with average

Δ%butyrate (ρ = -0.40, p = 0.046; Spearman correlation; Figure 4A). Thus, higher habitual fiber

intake was associated with a lower average Δ%butyrate during the prebiotic intervention. To

corroborate these findings using an objective biomarker, we also compared average Δ%butyrate

to spot urinary potassium, a known marker for fruit and vegetable intake (71), as well as for

general healthy eating (72, 73). We validated that spot urinary potassium was correlated with

dietary fiber intake in our study (ρ = 0.47, p = 0.031; Spearman correlation; Figure S5A) and

found evidence for a negative correlation between spot urinary potassium and average

Δ%butyrate (ρ = -0.38, p = 0.071; Spearman correlation; Figure S5B).

Observational evidence that baseline metabolic state and dietary history limits prebiotic

response. To understand why habitual fiber intake would be negatively associated with prebiotic

responses, we considered recent ecological studies suggesting that nutrient limitation is a key

factor shaping the metabolism of mammalian gut microbiota (74, 75). Even if excess fiber is

available to polysaccharide utilizing microbes, fixed caps in the availability of other nutrients

involved in carbohydrate fermentation would ultimately constrain the total amount of SCFAs

that can be produced from dietary fibers. Individuals who naturally consume high levels of fiber

and already activate SCFA production pathways would therefore exhibit reduced capacity to

increase SCFA production following prebiotic treatment.

We identified three findings that were consistent with a threshold model of metabolic

capacity limitation. First, we observed that individuals with higher starting proportional

concentrations of butyrate exhibited the weakest butyrogenic prebiotic responses. ($p = 0.008$, ρ

$= -0.49$, Spearman correlation; Figure 4B). No participant ultimately exhibited butyrate

concentrations in excess of 29mM following prebiotic treatment.

A second finding linking background nutrient intake and metabolic limitation involved

baseline proportional butyrate concentrations and habitual diet. Dietary fiber consumption was

positively correlated with proportional butyrate concentration at baseline week 1, prior to any

prebiotic intervention (1g fiber/2000 kilocalorie diet led to an increase in proportional butyrate

concentration of 0.010, $p = 0.012$, beta regression; Figure S6) as has been previously suggested

(33, 36, 76). Furthermore, using an ecological technique for identifying relationships between

multivariate datasets known as co-inertia analysis (77), we found a significant interaction

between baseline proportional SCFA concentrations and food choices($p = 0.031$, coinertia

analysis). This analysis associated intake of several fiber-rich food groups with baseline

proportional butyrate concentrations including legumes, berry fruits, and dark-green vegetables

(Figure 5).

Culture-based experimental evidence links fiber intake to limits on microbiome metabolic

response. A third line of experimental evidence arose linking habitual nutrient intake and

microbial SCFA production capacity. We hypothesized that if baseline fiber consumption

reduced the capacity of gut microbiota to produce SCFAs, the metabolic activity of fecal

microbiota would exhibit a negative relationship with fiber intake. To test this hypothesis, we

applied a recently developed an *ex vivo* fermentation model (40) to baseline stool samples

collected from each study participant. This model experimentally tests the metabolic effects of

supplementing a microbial community using a given prebiotic as the sole additional carbon

source, independent of potentially confounding host factors like SCFA absorption. In support of

the model's capability to produce SCFAs following prebiotic administration, we observed an

increase in overall SCFA production relative to control ($p < 0.0001$, t-test; average of 4.41-fold

increase across participants, calculated using the average total SCFA concentration of 3 separate

prebiotic treatments) when the *in vitro* assays were supplemented with prebiotics as the sole

carbon source. When we compared SCFA production to fiber intake, we observed as expected a

negative correlation between fiber consumption and SCFA production ($p = 0.021$, $\rho = -0.46$,

Spearman correlation using average of three prebiotic treatments; Figure 4C). We also observed

a negative correlation between fiber consumption and SCFA production in prebiotic-free control

treatments ($p = 0.019$, $\rho = -0.47$, Spearman correlation; Figure S7. Thus, individuals who tended

to eat more fiber in our cohort also harbored gut microbiota that produced less SCFA when

exposed to prebiotics.

Consistent growth of specific microbial genera supports individual-specific prebiotic response. Notably, our *ex vivo* fermentations also associated specific gut bacterial taxa with prebiotic response. Following 24hr fermentation, microbial community composition was assayed from each prebiotic-supplemented culture and compared to prebiotic-free control cultures, for each of the 28 participants, using ALDEx2. This analysis revealed multiple genera that were over or under-represented in each prebiotic treatment (Figure 6).

 Additionally, most genera that were differentially abundant in one prebiotic treatment exhibited similar differential abundance in the other prebiotic treatments. All genera exhibiting conserved higher abundance in two or three prebiotic treatments compared to control (*Collinsella*, *Faecalibacterium*, *Lachnospiraceae*, *Marvinbryantia*, and *Ruminiclostridium*) are known degraders of common dietary fibers with broad specificity (78-80). By contrast, of the genera exhibiting conserved lower abundance in two or three prebiotic treatments compared to control (*Alistipes*, *Anaerofilum*, *Flavonifractor*, *Holdemania*, and *Parabacteroides*), we are not aware of reports linking them to degradation of inulin, dextrin, or GOS, and only *Parabacteroides* has been shown to degrade any dietary fiber (78, 80). Overall, the consistent response of select bacterial taxa to prebiotic addition _in vitro_ further supports the conclusion that responses across a range of prebiotics will be linked to individual-specific factors like gut microbial community composition.

Discussion

In this study, we administered the prebiotics inulin, GOS, and dextrin to a set of individuals using a fully balanced crossover design. We found that these prebiotics varied in

their impact on fecal SCFA concentrations when data was agglomerated at the population scale,

with inulin increasing proportional butyrate and both GOS and wheat dextrin decreasing

proportional butyrate. Yet, our analysis also suggested that individual identity exerted a much

stronger effect on prebiotic response, and we observed that responses to each prebiotic were

correlated within individuals. Individual response variation was associated with differences in

baseline microbiological and dietary factors. In particular, we found that the general butyrogenic

response to prebiotics was anti-correlated with habitual fiber intake. This negative relationship

may be explained by limitations in the overall SCFA production capacity of human gut

microbiota.

From a microbial ecological perspective, our findings support a model where microbial

metabolic repertoires within individuals' microbiota have the capacity to utilize a diverse array

of chemically distinct fiber types. Evidence in favor of this model include *in vitro* prebiotic

screens suggesting that individuals uniformly harbor bacteria capable of degrading a wide variety

of carbohydrate species (81). Such metabolic plasticity may benefit gut microbiota by enhancing

community stability and invasion resistance (82-84), and could be enabled by substrate-

switching at the level of strains (84, 85) or cross-feeding interactions (86). Still, differences in

the levels of strains underlying these metabolic interactions may vary between individuals (81,

87), which could in turn lead to inter-individual variation in the capacity of individuals'

microbiota to ferment prebiotics.

The nutritional implications of our work were, in our view, more surprising. A positive

relationship between dietary fiber consumption and microbiota community structure (88), and

metabolic function (68, 89-91) is well established, with higher fiber intake linked to greater

SCFA and, specifically, butyrate production. Thus, we expected that individuals habituated to a

high-fiber diet would have a greater capacity to utilize incoming novel fiber, not a lower capacity

as we observed. Still, prior studies reporting links between diet patterns and microbiota response

have specifically associated changes to microbial community composition with baseline fiber

intake (37). Associations between fiber consumption and SCFA production, by contrast, have

either not been tested for in prebiotic trials (92, 93) or not revealed a positive relationship (37).

Additionally, while our analyses confirmed an association between baseline proportional

butyrate levels and intake of plant foods groups linked to dietary fiber (e.g. legumes and dark

green vegetables), it also linked food groups typically associated with a Mediterranean diets (e.g.

seafood, dairy, fruit, and poultry (94)) with higher baseline levels of proportional acetate. Future

prebiotic trials are needed to determine the generality of our observed association between

habitual diet and SCFA production

Conclusions

Our study provides evidence that the responsiveness of an individual to prebiotic

treatment may be predictable from diet and baseline concentrations of SCFA in stool. We

emphasize that we do not believe these predictions should be used to modify the diets of

individuals who normally consume high levels of fiber or have higher concentrations of SCFA in

stool. Ample evidence supports the benefits of both a high fiber diet and adequate microbial

SCFA production (70, 95, 96). Rather, we argue our findings support strategies for personalizing

microbiota therapy that focus on individual selection. Ongoing prioritization efforts are being

used to identify patients most likely to respond to microbiota-directed and microbiota-dependent

disease therapies (97-100), and are also being explored to enhance prebiotic therapy (30, 69, 90,

91). Our study here suggests that such efforts to personalize prebiotic supplementation can rely

on dietary history and fecal metabolite levels as informative and non-invasive biomarkers of

treatment outcome.

Ethics approval and consent to participate

All recruitment and study procedures involving human subjects were approved by the

Duke University Institutional Review Board (Pro00087214). All subjects provided electronic

informed consent prior to enrollment.

Competing interests

The authors declare that they have no competing interests

Funding

This study was supported by funding from the Beckman Young Investigator program, the

Damon Runyon Cancer Research Foundation, the NASA Translational Research

Institute through Cooperative Agreement NNX16AO69A, and NIH 1R01DK116187-01. This

work used a high-performance computing facility partially supported by grant 2016-IDG-1013

(HARDAC+: Reproducible HPC for Next-generation Genomics"). ZH was funded by the

National Defense Science and Engineering Graduate fellowship. MV was funded by a

Postdoctoral Enrichment Program Award from the Burroughs Wellcome Fund. JS and BP were

funded by the Duke School of Medicine Medical Scientist Training Program. BP was

additionally funded by the Integrative Bioinformatics for Investigating and Engineering

Microbiomes (IBIEM) program. No funding body had any role in the design of the study nor the

collection, analysis, or interpretation of the data nor in writing the manuscript.

Authors' contributions

ZH, MV, JS, LD, and HD conceptualized and designed the study. HD, ZH, SJ, and LD coordinated recruitment and engaged with participants. ZH was blinded to participant identity and study arm assignment where possible. ZH, SJ, ED, and HD processed samples, ran *in vitro* fermentation assays, and analyzed samples via gas-chromatography and sequencing. PHL and BP supported collection and analysis of nutrition data. ZH analyzed all data. ZH and LD wrote the manuscript. All authors read and approved the manuscript.

Acknowledgements

The authors acknowledge the Stedman Nutrition Center at Duke University for lending kitchen facilities for food-safe repackaging of prebiotics, and Jeffrey Letourneau, Rachael Bloom, and Savita Gupta for assistance with study coordination and logistics.

Figures

Figure 1: Prebiotic trial design. A three-period crossover design was standardized in series and
period, with each prebiotic occurring the same number of times within each arm (once) and study
period (twice). The design was also balanced with respect to first-order carry-over effects,
meaning prebiotics were administered in all possible orders. Each of the three periods consisted
of a prebiotic-free week followed by a prebiotic-supplemented week. Six study arms
encompassed all possible orders of prebiotic supplementation, and subjects were assigned to
arms sequentially by time of enrollment, resulting in approximately equal numbers of individuals
in each arm. A total of 41 participants were enrolled and 28 were considered to have completed
the study.

504
Figure 2: Effects of prebiotic supplementation on proportional butyrate concentration of
stool. Plotted are the average proportional butyrate concentrations for each study participant
during each prebiotic intervention period (up to 3 samples per person per week), and the average
of all three baseline weeks (up to 9 samples per person). Inulin was associated with significantly
increased proportional butyrate concentration of stool averaged across individuals, while both
dextrin and GOS were associated with decreased proportional butyrate relative to control (dashed
line) (generalized linear mixed model).

	Control	Inulin		GOS		Dextrin	
	value	value	p	value	p	value	p
CONCENTRATION							
Acetate (mM)	48.1	46.4	ns	47.8	ns	51.7	ns
Propionate (mM)	13.8	13.7	ns	13.3	**0.015**	15.7	**0.038**
Butyrate (mM)	12.4	13.7	ns	11.2	**0.002**	12.2	ns
Total SCFA (mM)	74.5	73.9	ns	72.4	**0.038**	79.6	ns
PROPORTION							
Acetate (%)	64.3	62.5	ns	65.3	ns	64.7	ns
Propionate (%)	19.2	19.5	ns	19.1	ns	20	ns
Butyrate (%)	16.5	18.0	**0.009**	15.6	**0.027**	15.3	**0.006**

513 **Table 1: Concentrations and proportions of SCFA in stool during prebiotic and control**
514 **weeks, averaged across individuals**. Significance was determined by generalized linear mixed
515 models.

516
517
Figure 3: Individual changes in proportional butyrate concentration (Δ%butyrate)
compared across multiple prebiotics. Changes in Δ%butyrate were positively correlated when
comparing responses to dextrin and inulin ($\rho = 0.49$, $p = 0.008$; Spearman correlation), as well as
responses to dextrin and GOS ($\rho = 0.37$, $p = 0.05$). The relationship between inulin and GOS
response trended towards a positive association ($\rho = 0.28$, $p = 0.15$).

525
526
Figure 4: Relationships between habitual diet, baseline proportional butyrate
concentration, and metrics of SCFA production. (A) Average change in proportional butyrate
concentration of stool during intervention (Δ%butyrate) was negatively correlated with habitual
dietary fiber consumption (ρ = -0.40, p = 0.046; Spearman correlation). (B) Average Δ%butyrate
was also negatively correlated with baseline proportional butyrate concentration (ρ = -0.49, p =
0.008; Spearman correlation). (C) This relationship between habitual fiber consumption and
SCFA production from prebiotics was also observed *in vitro* (ρ = -0.46, p = 0.021; Spearman
correlation).

Figure 5: Relationship between baseline SCFA profiles and habitual intake of food groups.
(**A**) Baseline SCFA profiles and average food intakes among participants. Participants (rows)
clustered based on proportional SCFA profiles. Cells values represent z-scores within each
column. Loadings shown of proportional SCFA concentrations (**B**) and consumed food groups
(**C**) along canonical axes 1 and 2 resulting from co-inertia analysis of SCFA and diet data (p =
0.031). Each axis is represented identically in each plot, so that eigenvectors (direction of
arrows) correspond between the SCFA and food group plots and the magnitude of eigenvalues
(length of arrows) is related to the strength of that component in the axis loading.

Figure 6: Gut bacterial taxa whose abundance changes during prebiotic-supplemented *ex vivo* fermentation. Bacterial abundances were compared to prebiotic-free control fermentation. Colors represent the number of prebiotic conditions in which a genus was differentially abundant (Benjamini-Hochberg corrected q value < 0.2, visualized with horizontal dashed line). Effect size estimate as reported by ALDEx2 is the median difference between groups (prebiotic and control) divided by the maximal difference within either group, in clr-transformed coordinates.

553 <u>Supplementary figures and tables</u>

554
555

557
558
559 **Figure S1: Proportional Acetate (A), proportional propionate (B), and total SCFA (C)**
560 **concentrations during baseline and prebiotic interventions.** All significant differences
561 between prebiotic treatments and control are designated by displayed p values, calculated using a
562 series of generalized linear mixed models.

563
Figure S2: Average proportional butyrate concentration (A) and average butyrate concentration (B) in participant stool during the first and last week of the study. Both proportional concentration and raw concentration were correlated between the first and last weeks ($\rho = 0.49$, $p < 0.007$; and $\rho = 0.63$, $p = 0.0003$, respectively, Spearman correlations).
568

569
Figure S3: Biplot of stool 16S community composition showing samples (dots) plotted on two principal component axes along with corresponding taxonomic variables (arrows), agglomerated to genus level. Samples are colored by group (high and low; > median baseline %butyrate and <= median baseline %butyrate, respectively) and ellipses represent 95% confidence intervals of the centroid for each group. Centroids are significantly different (p = 0.0022; PERMANOVA using unweighted UniFrac distance). Only significantly differentially abundant genera (Benjamini-Hochberg corrected q < 0.05, ALDEx2) are represented by arrows and labels.

580
Figure S4: PCA of stool 16S community composition showing samples plotted on two
principal component axes. No genera were significantly differentially abundant (Benjamini-
Hochberg corrected q < 0.05, ALDEx2), so no variables are plotted**.** Samples are colored by
group (high and low; > 0 average butyrogenic response to prebiotics and < 0 average butyrogenic
response to prebiotics, respectively) and ellipses represent 95% confidence intervals of the
centroid for each group. Centroids are significantly different (p = 0.028; PERMANOVA using
unweighted UniFrac distance).

588
Figure S5: Relationships between urinary potassium and (A) habitual dietary fiber
consumption, and (B) average change in proportional butyrate concentration during
prebiotic interventions. Urinary potassium was positively correlated with habitual fiber
consumption (ρ = 0.47, p = 0.031; Spearman correlation; A) and showed evidence for a negative
correlation with average change in proportional butyrate concertation during prebiotic
interventions (ρ = -0.38, p = 0.071; Spearman correlation; B).

596
597   **Figure S6: Relationship between habitual dietary fiber consumption and baseline**
598   **proportional butyrate concentration at week 1 (prior to study intervention).** Fiber
599   consumption was positively associated with baseline proportional butyrate concentration, with
600   1g fiber/2000 kilocalorie diet associated with an increase in proportional butyrate concentration
601   of 0.010 (p = 0.012, beta regression).

Figure S7: Relationship between *in vitro* SCFA production in prebiotic-free control treatments and habitual dietary fiber consumption. *In vitro* SCFA production was negatively correlated with habitual fiber consumption (ρ = -0.43, p = 0.031; Spearman correlation).

Citations

1.   Flint HJ, Duncan SH, Scott KP, Louis P. 2015. Links between diet, gut microbiota
composition and gut metabolism. Proceedings of the Nutrition Society 74:13-22.
2.   Morrison DJ, Preston T. 2016. Formation of short chain fatty acids by the gut microbiota
and their impact on human metabolism. Gut microbes 7:189-200.
3.   Samuel BS, Shaito A, Motoike T, Rey FE, Backhed F, Manchester JK, Hammer RE,
Williams SC, Crowley J, Yanagisawa M, Gordon JI. 2008. Effects of the gut microbiota
on host adiposity are modulated by the short-chain fatty-acid binding G protein-coupled
receptor, Gpr41. Proc Natl Acad Sci U S A 105:16767-72.
4.   Smith PM, Howitt MR, Panikov N, Michaud M, Gallini CA, Bohlooly YM, Glickman
JN, Garrett WS. 2013. The microbial metabolites, short-chain fatty acids, regulate colonic
Treg cell homeostasis. Science 341:569-73.
5.   Lee ES, Lee DS, Pandeya PR, Kim YC, Kang DG, Lee HS, Oh BC, Lee DH. 2017.
Sodium butyrate has context-dependent actions on dipeptidyl peptidase-4 and other
metabolic parameters. Korean J Physiol Pharmacol 21:519-529.
6.   Lee C, Kim BG, Kim JH, Chun J, Im JP, Kim JS. 2017. Sodium butyrate inhibits the NF-
kappa B signaling pathway and histone deacetylation, and attenuates experimental colitis
in an IL-10 independent manner. Int Immunopharmacol 51:47-56.
7.   Yan H, Ajuwon KM. 2017. Butyrate modifies intestinal barrier function in IPEC-J2 cells
through a selective upregulation of tight junction proteins and activation of the Akt
signaling pathway. PLoS One 12:e0179586.
8.   Hatayama H, Iwashita J, Kuwajima A, Abe T. 2007. The short chain fatty acid, butyrate,
stimulates MUC2 mucin production in the human colon cancer cell line, LS174T.
Biochem Biophys Res Commun 356:599-603.
9.   Tolhurst G, Heffron H, Lam YS, Parker HE, Habib AM, Diakogiannaki E, Cameron J,
Grosse J, Reimann F, Gribble FM. 2012. Short-chain fatty acids stimulate glucagon-like
peptide-1 secretion via the G-protein-coupled receptor FFAR2. Diabetes 61:364-71.
10.   Brown AJ, Goldsworthy SM, Barnes AA, Eilert MM, Tcheang L, Daniels D, Muir AI,
Wigglesworth MJ, Kinghorn I, Fraser NJ, Pike NB, Strum JC, Steplewski KM, Murdock
PR, Holder JC, Marshall FH, Szekeres PG, Wilson S, Ignar DM, Foord SM, Wise A,
Dowell SJ. 2003. The Orphan G protein-coupled receptors GPR41 and GPR43 are
activated by propionate and other short chain carboxylic acids. J Biol Chem 278:11312-9.
11.   Sonnenburg JL, Bäckhed F. 2016. Diet–microbiota interactions as moderators of human
metabolism. Nature 535:56.
12.   Gao Z, Yin J, Zhang J, Ward RE, Martin RJ, Lefevre M, Cefalu WT, Ye J. 2009.
Butyrate improves insulin sensitivity and increases energy expenditure in mice. Diabetes
58:1509-1517.
13.   Lu Y, Fan C, Li P, Lu Y, Chang X, Qi K. 2016. Short Chain Fatty Acids Prevent High-
fat-diet-induced Obesity in Mice by Regulating G Protein-coupled Receptors and Gut
Microbiota. Sci Rep 6:37589.
14.   Frost G, Sleeth ML, Sahuri-Arisoylu M, Lizarbe B, Cerdan S, Brody L, Anastasovska J,
Ghourab S, Hankir M, Zhang S, Carling D, Swann JR, Gibson G, Viardot A, Morrison D,
Louise Thomas E, Bell JD. 2014. The short-chain fatty acid acetate reduces appetite via a
central homeostatic mechanism. Nat Commun 5:3611.
15.   Thorburn AN, Macia L, Mackay CR. 2014. Diet, metabolites, and "western-lifestyle"
inflammatory diseases. Immunity 40:833-42.

16.   Nie Y, Lin Q, Luo F. 2017. Effects of Non-Starch Polysaccharides on Inflammatory
Bowel Disease. Int J Mol Sci 18.
17.   Pascal V, Pozuelo M, Borruel N, Casellas F, Campos D, Santiago A, Martinez X, Varela
E, Sarrabayrouse G, Machiels K, Vermeire S, Sokol H, Guarner F, Manichanh C. 2017.
A microbial signature for Crohn's disease. Gut 66:813-822.
18.   Thibault R, De Coppet P, Daly K, Bourreille A, Cuff M, Bonnet C, Mosnier JF,
Galmiche JP, Shirazi-Beechey S, Segain JP. 2007. Down-regulation of the
monocarboxylate transporter 1 is involved in butyrate deficiency during intestinal
inflammation. Gastroenterology 133:1916-27.
19.   De Preter V, Arijs I, Windey K, Vanhove W, Vermeire S, Schuit F, Rutgeerts P, Verbeke
K. 2012. Impaired butyrate oxidation in ulcerative colitis is due to decreased butyrate
uptake and a defect in the oxidation pathway. Inflamm Bowel Dis 18:1127-36.
20.   Palmieri O, Creanza TM, Bossa F, Palumbo O, Maglietta R, Ancona N, Corritore G,
Latiano T, Martino G, Biscaglia G, Scimeca D, De Petris MP, Carella M, Annese V,
Andriulli A, Latiano A. 2015. Genome-wide Pathway Analysis Using Gene Expression
Data of Colonic Mucosa in Patients with Inflammatory Bowel Disease. Inflamm Bowel
Dis 21:1260-8.
21.   Institute of Medicine. 2001. Dietary Reference Intakes: Proposed Definition of Dietary
Fiber National Academies Press, Washington DC.
22.   USDA Agricultural Research Service. 2021. Usual Nutrient Intake from Food and
Beverages, by Gender and Age, What We Eat in America, NHANES 2015-2018
23.   Park Y, Subar AF, Hollenbeck A, Schatzkin A. 2011. Dietary fiber intake and mortality
in the NIH-AARP diet and health study. Archives of internal medicine 171:1061-1068.
24.   Ou J, Carbonero F, Zoetendal EG, DeLany JP, Wang M, Newton K, Gaskins HR,
O'Keefe SJD. 2013. Diet, microbiota, and microbial metabolites in colon cancer risk in
rural Africans and African Americans. The American journal of clinical nutrition 98:111-
120.
25.   Maslowski KM, Mackay CR. 2011. Diet, gut microbiota and immune responses. Nature
immunology 12:5-9.
26.   Atarashi K, Tanoue T, Shima T, Imaoka A, Kuwahara T, Momose Y, Cheng G,
Yamasaki S, Saito T, Ohba Y. 2011. Induction of colonic regulatory T cells by
indigenous Clostridium species. Science 331:337-341.
27.   Atarashi K, Tanoue T, Oshima K, Suda W, Nagano Y, Nishikawa H, Fukuda S, Saito T,
Narushima S, Hase K, Kim S, Fritz JV, Wilmes P, Ueha S, Matsushima K, Ohno H, Olle
B, Sakaguchi S, Taniguchi T, Morita H, Hattori M, Honda K. 2013. Treg induction by a
rationally selected mixture of Clostridia strains from the human microbiota. Nature
500:232-236.
28.   Geuking MB, Cahenzli J, Lawson MAE, Ng DCK, Slack E, Hapfelmeier S, McCoy KD,
Macpherson AJ. 2011. Intestinal bacterial colonization induces mutualistic regulatory T
cell responses. Immunity 34:794-806.
29.   Smith PM, Howitt MR, Panikov N, Michaud M, Gallini CA, Bohlooly-y M, Glickman
JN, Garrett WS. 2013. The microbial metabolites, short-chain fatty acids, regulate colonic
Treg cell homeostasis. Science 341:569-573.
30.   Baxter NT, Schmidt AW, Venkataraman A, Kim KS, Waldron C, Schmidt TM. 2019.
Dynamics of Human Gut Microbiota and Short-Chain Fatty Acids in Response to Dietary
Interventions with Three Fermentable Fibers. mBio 10.

31. Venkataraman A, Sieber JR, Schmidt AW, Waldron C, Theis KR, Schmidt TM. 2016.
Variable responses of human microbiomes to dietary supplementation with resistant
starch. Microbiome 4:9.
32. Deehan EC, Yang C, Perez-Munoz ME, Nguyen NK, Cheng CC, Triador L, Zhang Z,
Bakal JA, Walter J. 2020. Precision Microbiome Modulation with Discrete Dietary Fiber
Structures Directs Short-Chain Fatty Acid Production. Cell Host Microbe 27:389-404 e6.
33. McOrist AL, Miller RB, Bird AR, Keogh JB, Noakes M, Topping DL, Conlon MA.
2011. Fecal butyrate levels vary widely among individuals but are usually increased by a
diet high in resistant starch. J Nutr 141:883-9.
34. Davani-Davari D, Negahdaripour M, Karimzadeh I, Seifan M, Mohkam M, Masoumi SJ,
Berenjian A, Ghasemi Y. 2019. Prebiotics: Definition, Types, Sources, Mechanisms, and
Clinical Applications. Foods 8.
35. Cerdó T, García-Santos JA, G. Bermúdez M, Campoy C. 2019. The Role of Probiotics
and Prebiotics in the Prevention and Treatment of Obesity. Nutrients 11.
36. Holscher HD, Bauer LL, Gourineni V, Pelkman CL, Fahey GC, Jr., Swanson KS. 2015.
Agave Inulin Supplementation Affects the Fecal Microbiota of Healthy Adults
Participating in a Randomized, Double-Blind, Placebo-Controlled, Crossover Trial. J
Nutr 145:2025-32.
37. Healey G, Murphy R, Butts C, Brough L, Whelan K, Coad J. 2018. Habitual dietary fibre
intake influences gut microbiota response to an inulin-type fructan prebiotic: a
randomised, double-blind, placebo-controlled, cross-over, human intervention study. Br J
Nutr 119:176-189.
38. Kleessen B, Schwarz S, Boehm A, Fuhrmann H, Richter A, Henle T, Krueger M. 2007.
Jerusalem artichoke and chicory inulin in bakery products affect faecal microbiota of
healthy volunteers. British Journal of Nutrition 98:540-549.
39. Vester Boler BM, Rossoni Serao MC, Bauer LL, Staeger MA, Boileau TW, Swanson KS,
Fahey GC. 2011. Digestive physiological outcomes related to polydextrose and soluble
maize fibre consumption by healthy adult men. British Journal of Nutrition 106:1864-
1871.
40. Holmes ZC, Silverman JD, Dressman HK, Wei Z, Dallow EP, Armstrong SC, Seed PC,
Rawls JF, David LA. 2020. Short-Chain Fatty Acid Production by Gut Microbiota from
Children with Obesity Differs According to Prebiotic Choice and Bacterial Community
Composition. mBio 11.
41. Gamage H, Tetu SG, Chong RWW, Bucio-Noble D, Rosewarne CP, Kautto L, Ball MS,
Molloy MP, Packer NH, Paulsen IT. 2018. Fiber Supplements Derived From Sugarcane
Stem, Wheat Dextrin and Psyllium Husk Have Different In Vitro Effects on the Human
Gut Microbiota. Front Microbiol 9:1618.
42. Salonen A, Lahti L, Salojarvi J, Holtrop G, Korpela K, Duncan SH, Date P, Farquharson
F, Johnstone AM, Lobley GE, Louis P, Flint HJ, de Vos WM. 2014. Impact of diet and
individual variation on intestinal microbiota composition and fermentation products in
obese men. ISME J 8:2218-30.
43. National Institutes of Health ARP, National Cancer Institute. 2018. Diet History
Questionnaire, Version 3.0.
44. Subar AF, Kirkpatrick SI, Mittl B, Zimmerman TP, Thompson FE, Bingley C, Willis G,
Islam NG, Baranowski T, McNutt S, Potischman N. 2012. The Automated Self-

Administered 24-hour dietary recall (ASA24): a resource for researchers, clinicians, and
educators from the National Cancer Institute. J Acad Nutr Diet 112:1134-7.

45.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ,
Fierer N, Knight R. 2011. Global patterns of 16S rRNA diversity at a depth of millions of
sequences per sample. Proceedings of the National Academy of Sciences 108:4516.

46.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, Owens SM,
Betley J, Fraser L, Bauer M, Gormley N, Gilbert JA, Smith G, Knight R. 2012. Ultra-
high-throughput microbial community analysis on the Illumina HiSeq and MiSeq
platforms. The Isme Journal 6:1621.

47.   Maurice Corinne F, Haiser Henry J, Turnbaugh Peter J. 2013. Xenobiotics Shape the
Physiology and Gene Expression of the Active Human Gut Microbiome. Cell 152:39-50.

48.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes SP. 2016.
DADA2: High-resolution sample inference from Illumina amplicon data. Nat Methods
13:581-3.

49.   Silverman JD, Durand HK, Bloom RJ, Mukherjee S, David LA. 2018. Dynamic linear
models guide design and analysis of microbiota studies within artificial human guts.
Microbiome 6:202.

50.   Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer
N, Pena AG, Goodrich JK, Gordon JI, Huttley GA, Kelley ST, Knights D, Koenig JE,
Ley RE, Lozupone CA, McDonald D, Muegge BD, Pirrung M, Reeder J, Sevinsky JR,
Turnbaugh PJ, Walters WA, Widmann J, Yatsunenko T, Zaneveld J, Knight R. 2010.
QIIME allows analysis of high-throughput community sequencing data. Nat Methods
7:335-6.

51.   McMurdie PJ, Holmes S. 2013. phyloseq: an R package for reproducible interactive
analysis and graphics of microbiome census data. PLoS One 8:e61217.

52.   Davani-Davari D, Negahdaripour M, Karimzadeh I, Seifan M, Mohkam M, Masoumi SJ,
Berenjian A, Ghasemi Y. 2019. Prebiotics: Definition, Types, Sources, Mechanisms, and
Clinical Applications. Foods (Basel, Switzerland) 8:92.

53.   Liu F, Li P, Chen M, Luo Y, Prabhakar M, Zheng H, He Y, Qi Q, Long H, Zhang Y,
Sheng H, Zhou H. 2017. Fructooligosaccharide (FOS) and Galactooligosaccharide (GOS)
Increase Bifidobacterium but Reduce Butyrate Producing Bacteria with Adverse
Glycemic Metabolism in healthy young population. Scientific Reports 7:11789.

54.   Rose C, Parker A, Jefferson B, Cartmell E. 2015. The Characterization of Feces and
Urine: A Review of the Literature to Inform Advanced Treatment Technology. Crit Rev
Environ Sci Technol 45:1827-1879.

55.   Liebisch G, Ecker J, Roth S, Schweizer S, Öttl V, Schött H-F, Yoon H, Haller D, Holler
E, Burkhardt R, Matysik S. 2019. Quantification of Fecal Short Chain Fatty Acids by
Liquid Chromatography Tandem Mass Spectrometry-Investigation of Pre-Analytic
Stability. Biomolecules 9:121.

56.   Clarke ST, Green-Johnson JM, Brooks SPJ, Ramdath DD, Bercik P, Avila C, Inglis GD,
Green J, Yanke LJ, Selinger LB, Kalmokoff M. 2016. beta 2-1 Fructan supplementation
alters host immune responses in a manner consistent with increased exposure to microbial
components: results from a double-blinded, randomised, cross-over study in healthy
adults. British Journal of Nutrition 115:1748-1759.

57.   Falony G, Verschaeren A, De Bruycker F, De Preter V, Verbeke K, Leroy F, De Vuyst L.
2009. In Vitro Kinetics of Prebiotic Inulin-Type Fructan Fermentation by Butyrate-

Producing Colon Bacteria: Implementation of Online Gas Chromatography for
Quantitative Analysis of Carbon Dioxide and Hydrogen Gas Production. Applied and
Environmental Microbiology 75:5884-5892.
58.   De Vuyst L, Leroy F. 2011. Cross-feeding between bifidobacteria and butyrate-producing
colon bacteria explains bifdobacterial competitiveness, butyrate production, and gas
production. Int J Food Microbiol 149:73-80.
59.   Kim MG, Jo K, Chang YB, Suh HJ, Hong KB. 2020. Changes in the Gut Microbiome
after Galacto-Oligosaccharide Administration in Loperamide-Induced Constipation. J
Pers Med 10.
60.   O'Connor KM, Lucking EF, Bastiaanssen TFS, Peterson VL, Crispie F, Cotter PD,
Clarke G, Cryan JF, O'Halloran KD. 2020. Prebiotic administration modulates gut
microbiota and faecal short-chain fatty acid concentrations but does not prevent chronic
intermittent hypoxia-induced apnoea and hypertension in adult rats. EBioMedicine
59:102968-102968.
61.   Barczynska R, Jurgoński A, Slizewska K, Juśkiewicz J, Kapusniak J. 2017. Effects of
potato dextrin on the composition and metabolism of the gut microbiota in rats fed
standard and high-fat diets. Journal of Functional Foods 34:398-407.
62.   Barczynska R, Jurgoński A, Slizewska K, Juśkiewicz J, Kapusniak J. 2019. Corn starch
dextrin changes intestinal microbiota and its metabolic activity in rats fed a basal and
high-fat diet. British Food Journal 121:2219-2232.
63.   Thangaraju M, Cresci GA, Liu K, Ananth S, Gnanaprakasam JP, Browning DD,
Mellinger JD, Smith SB, Digby GJ, Lambert NA, Prasad PD, Ganapathy V. 2009.
GPR109A is a G-protein-coupled receptor for the bacterial fermentation product butyrate
and functions as a tumor suppressor in colon. Cancer Res 69:2826-32.
64.   Di Sabatino A, Morera R, Ciccocioppo R, Cazzola P, Gotti S, Tinozzi FP, Tinozzi S,
Corazza GR. 2005. Oral butyrate for mildly to moderately active Crohn's disease.
Aliment Pharmacol Ther 22:789-94.
65.   Hamer HM, Jonkers D, Venema K, Vanhoutvin S, Troost FJ, Brummer RJ. 2008. Review
article: the role of butyrate on colonic function. Aliment Pharmacol Ther 27:104-19.
66.   Mathewson ND, Jenq R, Mathew AV, Koenigsknecht M, Hanash A, Toubai T, Oravecz-
Wilson K, Wu S-R, Sun Y, Rossi C, Fujiwara H, Byun J, Shono Y, Lindemans C,
Calafiore M, Schmidt TM, Honda K, Young VB, Pennathur S, van den Brink M, Reddy
P. 2016. Gut microbiome–derived metabolites modulate intestinal epithelial cell damage
and mitigate graft-versus-host disease. Nature Immunology 17:505.
67.   McDonald D, Hyde E, Debelius JW, Morton JT, Gonzalez A, Ackermann G, Aksenov
AA, Behsaz B, Brennan C, Chen Y, DeRight Goldasich L, Dorrestein PC, Dunn RR,
Fahimipour AK, Gaffney J, Gilbert JA, Gogul G, Green JL, Hugenholtz P, Humphrey G,
Huttenhower C, Jackson MA, Janssen S, Jeste DV, Jiang L, Kelley ST, Knights D,
Kosciolek T, Ladau J, Leach J, Marotz C, Meleshko D, Melnik AV, Metcalf JL,
Mohimani H, Montassier E, Navas-Molina J, Nguyen TT, Peddada S, Pevzner P, Pollard
KS, Rahnavard G, Robbins-Pianka A, Sangwan N, Shorenstein J, Smarr L, Song SJ,
Spector T, Swafford AD, Thackray VG, et al. 2018. American Gut: an Open Platform for
Citizen Science Microbiome Research. mSystems 3.
68.   David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, Ling AV,
Devlin AS, Varma Y, Fischbach MA, Biddinger SB, Dutton RJ, Turnbaugh PJ. 2014.
Diet rapidly and reproducibly alters the human gut microbiome. Nature 505:559-63.

69. Healey GR, Murphy R, Brough L, Butts CA, Coad J. 2017. Interindividual variability in
gut microbiota and host response to dietary interventions. Nutr Rev 75:1059-1080.
70. Tap J, Furet JP, Bensaada M, Philippe C, Roth H, Rabot S, Lakhdari O, Lombard V,
Henrissat B, Corthier G, Fontaine E, Dore J, Leclerc M. 2015. Gut microbiota richness
promotes its stability upon increased dietary fibre intake in healthy adults. Environ
Microbiol 17:4954-64.
71. Appel LJ, Moore TJ, Obarzanek E, Vollmer WM, Svetkey LP, Sacks FM, Bray GA,
Vogt TM, Cutler JA, Windhauser MM, Lin P-H, Karanja N, Simons-Morton D,
McCullough M, Swain J, Steele P, Evans MA, Miller ER, Harsha DW. 1997. A Clinical
Trial of the Effects of Dietary Patterns on Blood Pressure. New England Journal of
Medicine 336:1117-1124.
72. Posma JM, Garcia-Perez I, Frost G, Aljuraiban GS, Chan Q, Van Horn L, Daviglus M,
Stamler J, Holmes E, Elliott P, Nicholson JK. 2020. Nutriome-metabolome relationships
provide insights into dietary intake and metabolism. Nature food 1:426-436.
73. Mente A, Irvine EJ, Honey RJDA, Logan AG. 2009. Urinary Potassium Is a Clinically
Useful Test to Detect a Poor Quality Diet. The Journal of Nutrition 139:743-749.
74. Reese AT, Pereira FC, Schintlmeister A, Berry D, Wagner M, Hale LP, Wu A, Jiang S,
Durand HK, Zhou X, Premont RT, Diehl AM, O'Connell TM, Alberts SC, Kartzinel TR,
Pringle RM, Dunn RR, Wright JP, David LA. 2018. Microbial nitrogen limitation in the
mammalian large intestine. Nat Microbiol 3:1441-1450.
75. Macfarlane S, Macfarlane GT. 2003. Regulation of short-chain fatty acid production.
Proc Nutr Soc 62:67-72.
76. Holscher HD. 2017. Dietary fiber and prebiotics and the gastrointestinal microbiota. Gut
Microbes 8:172-184.
77. Dolédec S, Chessel D. 1994. Co-inertia analysis: an alternative method for studying
species–environment relationships. Freshwater Biology 31:277-294.
78. Magnúsdóttir S, Heinken A, Kutt L, Ravcheev DA, Bauer E, Noronha A, Greenhalgh K,
Jäger C, Baginska J, Wilmes P, Fleming RMT, Thiele I. 2017. Generation of genome-
scale metabolic reconstructions for 773 members of the human gut microbiota. Nature
Biotechnology 35:81-89.
79. Giallo J, Gaudin C, Belaich JP, Petitdemange E, Caillet-Mangin F. 1983. Metabolism of
glucose and cellobiose by cellulolytic mesophilic Clostridium sp. strain H10. Applied and
Environmental Microbiology 45:843.
80. Bergey DH, Garrity GM, Boone DR, Castenholz RW, Brenner DJ, Krieg NR, Staley JT.
2001. Bergey's manual of systematic bacteriology. New York : Springer, 2001-, New
York.
81. Villa MM, Bloom RJ, Silverman JD, Durand HK, Jiang S, Wu A, Dallow EP, Huang S,
You L, David LA. 2020. Interindividual Variation in Dietary Carbohydrate Metabolism
by Gut Bacteria Revealed with Droplet Microfluidic Culture. mSystems 5.
82. Maldonado-Gomez MX, Martinez I, Bottacini F, O'Callaghan A, Ventura M, van
Sinderen D, Hillmann B, Vangay P, Knights D, Hutkins RW, Walter J. 2016. Stable
Engraftment of Bifidobacterium longum AH1206 in the Human Gut Depends on
Individualized Features of the Resident Microbiome. Cell Host Microbe 20:515-526.
83. Deriu E, Liu JZ, Pezeshki M, Edwards RA, Ochoa RJ, Contreras H, Libby SJ, Fang FC,
Raffatellu M. 2013. Probiotic bacteria reduce salmonella typhimurium intestinal
colonization by competing for iron. Cell Host Microbe 14:26-37.

84.   Pereira FC, Berry D. 2017. Microbial nutrient niches in the gut. Environ Microbiol
19:1366-1378.

85.   Steeb B, Claudi B, Burton NA, Tienz P, Schmidt A, Farhan H, Maze A, Bumann D.
2013. Parallel exploitation of diverse host nutrients enhances Salmonella virulence. PLoS
Pathog 9:e1003301.

86.   Rakoff-Nahoum S, Foster KR, Comstock LE. 2016. The evolution of cooperation within
the gut microbiota. Nature 533:255-9.

87.   Samuel BS, Gordon JI. 2006. A humanized gnotobiotic mouse model of host-archaeal-
bacterial mutualism. Proc Natl Acad Sci U S A 103:10011-6.

88.   Wu GD, Chen J, Hoffmann C, Bittinger K, Chen YY, Keilbaugh SA, Bewtra M, Knights
D, Walters WA, Knight R, Sinha R, Gilroy E, Gupta K, Baldassano R, Nessel L, Li H,
Bushman FD, Lewis JD. 2011. Linking long-term dietary patterns with gut microbial
enterotypes. Science 334:105-8.

89.   Duncan SH, Belenguer A, Holtrop G, Johnstone AM, Flint HJ, Lobley GE. 2007.
Reduced dietary intake of carbohydrates by obese subjects results in decreased
concentrations of butyrate and butyrate-producing bacteria in feces. Appl Environ
Microbiol 73:1073-8.

90.   Kolodziejczyk AA, Zheng D, Elinav E. 2019. Diet-microbiota interactions and
personalized nutrition. Nat Rev Microbiol 17:742-753.

91.   Leshem A, Segal E, Elinav E. 2020. The Gut Microbiome and Individual-Specific
Responses to Diet. mSystems 5.

92.   Eid N, Osmanova H, Natchez C, Walton G, Costabile A, Gibson G, Rowland I, Spencer
JP. 2015. Impact of palm date consumption on microbiota growth and large intestinal
health: a randomised, controlled, cross-over, human intervention study. Br J Nutr
114:1226-36.

93.   Griffin NW, Ahern PP, Cheng J, Heath AC, Ilkayeva O, Newgard CB, Fontana L,
Gordon JI. 2017. Prior Dietary Practices and Connections to a Human Gut Microbial
Metacommunity Alter Responses to Diet Interventions. Cell Host Microbe 21:84-96.

94.   Davis C, Bryan J, Hodgson J, Murphy K. 2015. Definition of the Mediterranean Diet; a
Literature Review. Nutrients 7:9139-9153.

95.   Pryde SE, Duncan SH, Hold GL, Stewart CS, Flint HJ. 2002. The microbiology of
butyrate formation in the human colon. FEMS Microbiology Letters 217:133-139.

96.   Burkitt DP, Walker AR, Painter NS. 1972. Effect of dietary fibre on stools and the
transit-times, and its role in the causation of disease. Lancet 2:1408-12.

97.   Chumpitazi BP, Cope JL, Hollister EB, Tsai CM, McMeans AR, Luna RA, Versalovic J,
Shulman RJ. 2015. Randomised clinical trial: gut microbiome biomarkers are associated
with clinical response to a low FODMAP diet in children with the irritable bowel
syndrome. Alimentary Pharmacology & Therapeutics 42:418-427.

98.   Chumpitazi BP, Hollister EB, Oezguen N, Tsai CM, McMeans AR, Luna RA, Savidge
TC, Versalovic J, Shulman RJ. 2014. Gut microbiota influences low fermentable
substrate diet efficacy in children with irritable bowel syndrome. Gut Microbes 5:165-
175.

99.   Davar D, Dzutsev AK, McCulloch JA, Rodrigues RR, Chauvin J-M, Morrison RM,
Deblasio RN, Menna C, Ding Q, Pagliano O, Zidi B, Zhang S, Badger JH, Vetizou M,
Cole AM, Fernandes MR, Prescott S, Costa RGF, Balaji AK, Morgun A, Vujkovic-Cvijin
I, Wang H, Borhani AA, Schwartz MB, Dubner HM, Ernst SJ, Rose A, Najjar YG,

Belkaid Y, Kirkwood JM, Trinchieri G, Zarour HM. 2021. Fecal microbiota transplant overcomes resistance to anti–PD-1 therapy in melanoma patients. Science 371:595.

100. Gopalakrishnan V, Spencer CN, Nezi L, Reuben A, Andrews MC, Karpinets TV, Prieto PA, Vicente D, Hoffman K, Wei SC. 2018. Gut microbiome modulates response to anti–PD-1 immunotherapy in melanoma patients. Science 359:97-103.