Investigating the reliability of molecular estimates of evolutionary time when substitution rates and speciation rates vary

Andrew M. Ritchie[1][*], Xia Hua[1][2], Lindell Bromham[1]

1. *Research School of Biological Sciences, Australian National University, Canberra, ACT 2600, Australia*
2. *Mathematical Sciences Institute, Australian National University, Canberra, ACT 2600, Australia*

* Corresponding Author:

Andrew M. Ritchie

Research School of Biological Sciences, Robertson Building, 134 Linnaeus Way

Australian National University, Canberra, ACT 2600, Australia

Ph: +61 2 6125 5111

Email: Andrew.Ritchie@anu.edu.au

Acknowledgements

We would like to extend our gratitude to Marcel Cardillo and Alex Skeels for helpful conversations in the preparation of this manuscript.

Funding

This work was supported by the Australian Research Council grant DP190101916.

Abstract

<u>Background</u>

An accurate timescale of evolutionary history is essential to testing hypotheses

about the influence of historical events and processes, and the timescale for evolution is

increasingly derived from analysis of DNA sequences. But variation in the rate of molecular

evolution complicates the inference of time from DNA. Evidence is growing for numerous

factors, such as life history and habitat, that are linked both to the molecular processes of

mutation and fixation and to rates of macroevolutionary diversification. However, the most

widely used models of molecular rate variation, such as the uncorrelated and autocorrelated

lognormal clocks, rely on idealised models of rate variation and molecular dating methods

are rarely tested against complex models of rate change. One relationship that is not

accounted for in molecular dating is the potential for interaction between molecular

substitution rates and speciation, a relationship that has been supported by empirical studies

in a growing number of taxa. If these relationships are as widespread as evidence indicates,

they may have a significant influence on molecular dates.

<u>Results</u>

We simulate phylogenies and molecular sequences under three different realistic rate

variation models – one in which speciation rates and substitution rates both vary but are

unlinked, one in which they covary continuously and one punctuated model in which

molecular change is concentrated in speciation events, using empirical case studies to

parameterise realistic simulations. We test two commonly used "relaxed clock" molecular

dating methods against these realistic simulations to explore the degree of error in molecular

dates under each model. We find average divergence time inference errors ranging from 12%

of node age for the unlinked model when reconstructed under an uncorrelated rate prior, to up

to 93% when punctuated simulations are reconstructed under an autocorrelated prior.

Conclusions

We demonstrate the potential for substantial errors in molecular dates when both

speciation rates and substitution rates vary between lineages. This study highlights the need

for tests of molecular dating methods against realistic models of rate variation generated from

empirical parameters and known relationships.

1.Background

Understanding the timescale of evolution is critical to researching the processes that

generate and shape the diversity of life on Earth. Analysis of DNA sequences offers the

possibility of investigating the tempo and mode of evolution of all extant lineages, not just

those that have a detailed fossil record. The earliest molecular dating analyses concluded that

the average rate of molecular evolution was relatively constant over time in some proteins

because of an apparently linear accumulation of amino acid differences when compared

between species (1, 2). However, it has since become apparent that the average rate of

molecular evolution can be influenced by a wide range of species traits, such as generation

time, body size and longevity (3, 4), by macroevolutionary processes such as net

diversification rate (5-10) and potentially also by environmental features like temperature (9-

11). If the average rate of molecular evolution varies substantially and consistently between

lineages, then we ideally need to incorporate rate variation in models of molecular evolution

in order to accurately infer divergence times among species (12).

A wide variety of models have been developed to account for different patterns of

among-lineage molecular rate variation (13-16). In the first instance, these models differ in

how many different evolutionary rate parameters are estimated for different parts of the

phylogeny. For example, a 'strict clock' model estimates only one rate parameter that applies

to every branch in the tree, while a 'local clock' model estimates additional rate parameters

for subsets of branches (17-19), and a 'relaxed clock' provides for a separate rate on every

branch (20). Models also differ in how the rates on different branches are related to one

another. Some models, notably in the penalized-likelihood framework, aim to minimize the

overall variation of branch-specific rates or the differences between adjacent branches (21,

22). For Bayesian phylogenetic methods (23, 24), branch-specific rates are drawn from a

prior probability distribution that determines which rates and patterns of relative rate

variation are considered *a priori* more probable. We refer to this prior as the rate prior.

Common variants include uncorrelated models, in which each branch-specific rate is an

independent draw from a distribution with a common mean (20); and autocorrelated models,

in which the mean of the distribution for each branch-specific rate depends on the rate drawn

for its parent branch (25, 26).

Testing over the last two decades has overall supported the robustness of divergence

time estimates under the Bayesian methodology to the choice of rate prior, provided that

enough separate rate parameters are allowed and sufficient calibrating information is

provided (27-33); but see (34). However, these studies rely heavily on simulated data or on

specific and usually well-calibrated empirical examples, whereas simulations have indicated

that different rate priors may give different results when calibrating information is poorer

(30). Changing the rate prior in these circumstances could lead to differences in empirical

dates and associated causal hypotheses. Examples of empirical studies where the choice of

rate prior appears to be a factor in differing age estimates include influenza viruses (35), the

age of grasstrees (36) and the origins of Metazoa (37).

Currently used priors on rate variation are generally either uncorrelated, so that each

lineage-specific rate is drawn independently from a common distribution, or autocorrelated

so that lineage-specific rates depend evolve from the rates of their parent lineages according

to a stochastic, random walk. However, the factors shaping variation in rate of molecular

evolution are complex: in addition to random variation in rates between lineages, there is

clearly also a systematic component influenced by many factors including species traits,

population dynamics, environment and evolutionary history (38). Evolutionary rates among

eukaryotes appear to be faster at lower elevations and latitudes (9, 39, 40) but slower in arid

climates (41). Life history covaries with molecular rates across multiple axes, including body

size (42-45), fecundity (46) generation time (47).and longevity (48, 49). Associations have

also been observed between substitution rates and more complex species traits, such as

parasitism (50), flightlessness (51) and sexual competition (52). Furthermore, substitution

rates are the result of underlying microevolutionary processes, and may therefore be affected

by population size (53-56).

In addition to covarying with other evolving traits, there is evidence that variation in

molecular rates can be associated with macroevolutionary patterns. Relationships between the

relative size of sister clades, representing their net diversification rates, and their molecular

rates have been detected in several studies, include in reptiles (8, 57), birds (7, 58), flowering

plants (6, 59, 60) and across Metazoa (61). Several studies have also noted relationships

between the genetic divergence of species in a phylogeny and the number of intervening

speciation events, which may also represent examples of this phenomenon (62-64).

The mechanism behind observed correlations between diversification and molecular

evolution remains a matter of debate. One line of argument proposes that higher molecular

rates could promote speciation by speeding the development of reproductive isolation (65). In

this hypothesis, subpopulations become sporadically isolated over time and may accumulate

alleles that may be effectively neutral on their own but deleterious in combination (66, 67). In

this case, greater rates of molecular evolution could lead to faster development of

reproductive isolation during periods of isolation, leading to a greater average rate of

speciation (68, 69). Alternatively, higher molecular rates could reduce extinction rates by

accumulating more standing genetic diversity which provides a buffer against changing

environment by allowing more rapid selective response (70). Speciation rates and molecular

rates could also be related through a common driving factor, such as climate or

environmental productivity (5, 10). In the other causal direction, speciation itself could

increase average molecular rates by producing rapid bursts of evolution (62, 63, 71).

Suggested mechanisms for an increase in substitution rates during speciation involve an

increase in fixation rates of mildly deleterious alleles and linked neutral mutations due to

drift, or large-scale genomic divergence caused by an increase in the frequency of disruptive

genomic events such as polyploidy or chromosomal rearrangements (71).

An association between substitution rates and diversification rates could affect the

performance of existing divergence time estimation methods (e.g. 72). In particular, Bayesian

methods divide their specification of prior beliefs about the evolutionary process between the

rate prior, the prior on divergence times and tree topology (sometimes referred to as the tree

prior or node time prior), and calibrating distributions on the ages of individual speciation

events (23, 73). These priors are essential to estimating divergence time, because genetic

distances between sequences are the product of rate and time (the total number of character

differences between related sequences). This means that even if genetic distances are

correctly inferred on a phylogeny, rate and time are unidentifiable without additional

constraints on joint values they can hold (74, 75). The rate prior, node time prior and

calibrating distributions place either soft or hard constraints on these values, allowing

researchers to infer evolutionary history on an absolute timescale. However, this means that

inferred divergence times are always subject to the choice of rate, time and calibration priors

even with large amounts of sequence data (76, 77). Most commonly used Bayesian molecular

dating methods do not propose *a priori* relationships between the rate prior and the tree prior,

so the joint distribution implied by these priors will assign relatively higher probability to

long-lived, fast-evolving lineages and short-lived, slow-evolving lineages than would be

expected when rates and times are related. It is therefore possible that an unmodelled

relationship between rates and times could produce distributions of molecular rates and

speciation times that differ substantially from those implied by common Bayesian priors,

leading to widespread error in estimates of evolutionary dates.

Here, we investigate the potential influence of speciation-related rates of molecular

evolution on the reliability of molecular dating studies. Rather than assuming only one

specific model of the association between diversification rates and rates of molecular, we

model a range of possible causal links. We include three alternative models of the association

between rates of molecular evolution and speciation rates. In the first, both speciation rates

and substitution rates vary over the phylogeny, but neither influences the other. In the second,

we model continuous variation in a linked manner, which might represent either direct

influence of one on the other (65) or an indirect association between speciation rates and

substitution rates, for example if they are both influenced by environmental differences (5,

10). The third model is a punctuated model in which bursts of substitution are associated with

speciation events (62, 63, 71, 78). The proportion of all substitutions associated with such

bursts has recently been shown to impact Bayesian inference of clade crown ages under an

uncorrelated lognormal rate prior (72).

In this study, we simulate the evolution of molecular sequences under all three models

of the link between speciation and rate of molecular evolution – unlinked, continuous

coevolution and a punctuated model - allowing for independent variation in molecular and

speciation rates, and observe the average effects on errors in Bayesian divergence time

estimates. In order to make our test have real-world significance, we also base our

simulations, as far as possible, on empirically determined relationships between rates of

molecular evolution and diversification rate. We parameterize our models using values taken

from empirical inferences on bird data (79, 80) or which reproduce established empirical

relationships between rates of molecular evolution and diversification rate in birds (81). Since

the choice of rate prior may affect the results of phylogeny reconstruction (34), we also

examine the effect of divergence time reconstruction under uncorrelated and autocorrelated

rate priors. We reconstruct divergence times on these simulated data sets and examine the

effects of different forms of correlated molecular and speciation rates on the performance of

molecular divergence time reconstruction.

2. Results

2.1 Simulation design

We refer throughout to simulated phylogenies, which are known without error, as the

'simulated trees', and to the phylogenies inferred from analyzing the sequences evolved

along each of these simulated phylogenies as the 'reconstructed trees'. Comparison of the

reconstructed trees to the simulated trees allows us to evaluate the degree to which

phylogenetic inference methods correctly infer the known evolutionary history of the

simulated data.

Our overall design is summarised in Figure 1. For our simulation study, we first

generate phylogenetic trees in units of absolute time with evolving rates of speciation and

molecular evolution. We then simulate DNA sequences along these phylogenies. Finally, we

infer dated trees from the alignment of these sequences using standard methods in Bayesian

phylogenetics and compare the inferred tree to the simulated tree from the known

evolutionary history from each simulation.

We simulate sets of 50 phylogenetic trees and nucleotide sequence alignments under

each of three different models describing the association between speciation rates and rates of

molecular evolution: one "Unlinked" model in which molecular and diversification rates each

vary independently, as well as two alternative mechanisms for an association between rates

of molecular evolution and speciation rates ("Punctuated" and "Continuous": see Table 1).

We do not consider models in which molecular rates affect rates of extinction, so in our

simulations, extinction rates remain constant over the tree. For all models, we first simulate

the phylogenetic tree in continuous time, together with speciation and substitution rates that

vary along the tree. From this, we derive branch lengths in expected genetic divergence

(substitutions/site). Finally, we simulate the evolution of nucleotide sequences along these

phylogenies.

Table 1. Simulation design.

Simulation type	Speciation rates covary with substitution rates?	Instantaneous covariance	Punctuated burst of substitutions at nodes?
Unlinked	*No*	0	*no*
Continuous	*Yes*	0.0044	*no*
Punctuated	*No*	0	*yes*

234 We lay out the conditions for each set of simulated trees. Each set consisted of 50 trees.
235 Lineage-specific speciation rates and substitution rates evolve continuously through time via
236 Brownian motion in all three studies. The three simulations differ in whether the covariance
237 between speciation and substitution rates is greater than zero in the Brownian motion and
238 whether an additional burst of substitutions occurs at nodes of the tree (speciation events).
239

In the Unlinked model, we simulate continuous variation in both substitution rate and

speciation rate through time, but neither affects the other. Substitution and speciation rates

evolve independently by Brownian motion, where each lineage continuously adds a separate

random increment to its substitution rate and speciation rate in each of many small time

intervals. The Unlinked model serves as a baseline to determine the level of uncertainty in

molecular dates caused by the random variation in substitution and diversification rate when

there is no mechanistic link between the two. In the Continuous model, speciation rates and

substitution rates coevolve with each other continuously over time. The coevolution is

simulated as a correlated Brownian motion, where the random increments in speciation and

molecular rates added over time are dependent on one another. Consequently, substitution

rate and speciation rate tend to increase or decrease together. For this model, we also ensured

realism simulated many data sets with different covariance parameters, and selected a set of

50 trees that reproduced the empirical correlation of (7) between clade size and branch length

(Supplementary Methods S1.1). In the Punctuated model, the genetic divergence along a

branch is the sum of a gradual accumulation of substitutions at a rate that evolves as in the

Unlinked model, and a burst component at each speciation event. A speciation event occurs

each time a new lineage is born, whether any of its descendants are observed in the present or

not. The amount of additional branch length due to the burst is independently and randomly

drawn for each speciation event. For this model, we condition our simulations to reproduce

the relationship observed in (62). There it was observed that the total number of substitutions

occurring along root-to-tip paths in their sample of trees correlates with the number of nodes

along the path. Although this relationship could also be produced by a mechanism similar to

the Continuous model (82-84), we model the mechanism proposed in (62) that the node-

associated substitutions are produced in bursts associated with speciation events.

We derived plausible speciation, extinction and substitution rates for our simulations

using the avian phylogeny in (85). We do not intend to make any inferences about the actual

evolution of birds or the accuracy of inferred divergence times. Birds are chosen only

because the relationships among traits, molecular rates and diversification rates are well

studied, so that we can use these to give us an empirical ground for choosing parameter

values. Branch-specific diversification rates have been estimated for many of the lineages in

this tree in (80), while substitution rates have been inferred in (79) assuming covariation of

molecular rates and life history. Speciation rate and molecular rate values taken from avian

data have been previously used to inform realistic simulations on ecological assemblies for

the purpose of estimating the effect of dating error on phylogenetic diversity estimates (86),

and we make use of similar procedures here. A relationship between molecular rates and net

diversification rate has also been established for birds (7, 8). We use this fact to select our

Continuous data set by simulating multiple data sets and varying the covariance parameter

until the relationship appears. Full details of how we used these studies to determine realistic

parameterisations for our simulations are available as Supplementary Methods S1.1.

2.2 Validation of tree simulations

The simulation procedure produced trees with average characteristics similar to the

avian clades used to parameterize the study (80, supplementary data, 85). Across all data sets,

the mean root date of simulated trees was 28.4 million years, compared to a range of 23.8-

35.6 million years for bird clades with 70-80 taxa. Simulated lineage-specific speciation rates

at the tips were on average slightly lower than the speciation rate of similar clades in the bird

tree (80). The range of simulated speciation rates on log scale was 0.009-0.92 compared to a

range of 0.034-2.25 average log lineages/million years for bird clades with 70-80 taxa,

similar taxon number to our simulated trees. The range of speciation rates within each tree

was also similar to the range across the avian clades with similar taxon number, with an

average magnitude of 0.44 log lineages/million years compared to 0.32-2.21 for avian clades

with 70-80 taxa.

2.3 Error in reconstructed trees

We used three metrics (3.2), the gamma statistic (3.3) and the sister pair analysis (3.4)

to assess the accuracy of the reconstruction of trees simulated from the Unlinked model, the

Continuous model, and the Punctuated model, under uncorrelated lognormal (UCLN) and

autocorrelated lognormal (ACLN) models of among-lineage rate variation (Table 2).

Table 2. Summary of results.

Rate prior	Simulation	Node age error (%)	Topo. error	HPD width (%)	Gamma Statistic	Sister pair analysis
UCLN	*Unlinked*	6.3–30.9	0–14	43–29	-1.46–1.09	Null
UCLN	*Continuous*	6.7–30.3	0–16	35–605	-1.52–0.20	Null
UCLN	*Punctuated*	8.8–80.4	0-8	42–154	-2.16–3.92	Negative
ACLN	*Unlinked*	11–417	Fixed	84.9–217.8	-7.13–5.62	Positive
ACLN	*Continuous*	9.4–170	Fixed	73.5–319.4	-4.92–3.32	Positive
ACLN	*Punctuated*	13.7–397.8	Fixed	76.7 - 144.1	-4.95–2.88	Positive

Summary of reconstructed tree characteristics. Trees simulated under three models: Unlinked , Continuous, and Punctuated. Molecular dates were then reconstructed under two methods, the uncorrelated lognormal model (UCLN) and the autocorrelated lognormal model (ACLN). We calculated three metrics for each tree and summarised these over each simulation type. The metrics were median absolute % error of node ages, topological error (Topo. Error) by number of true bipartitions not reconstructed, and uncertainty by mean highest posterior density interval width as a proportion of reconstructed node age. We compared the gamma statistics of the reconstructed trees to the simulated trees in order to detect any systematic bias in node age error. We also performed sister pair analysis to test if we can detect the positive correlation between speciation rate and substitution rate from the reconstructed trees.

First, we examined the median absolute percentage error (MAPE) in the ages of all nodes within each tree whose topology were correctly reconstructed (Figure 2). For trees reconstructed under UCLN, the average (min–max) MAPE was 12.2% for Unlinked simulations (6.3–30.9%). Errors for the Continuous simulation were similar at 14.2% (6.7–30.3%). Average MAPE for the Punctuated simulation was higher than for Unlinked trees at 20.3% (8.8–80.4%). For trees reconstructed under the ACLN in PAML, MAPE values were higher and range of errors were larger than for the UCLN in BEAST. Average MAPE for Unlinked trees was 93.3% (11–417%), compared to 43.4% (9.4%–170%) for Continuous trees and 84% (13.7–397.8%) for Punctuated trees.

Second, for trees reconstructed under the UCLN model in BEAST, we also

investigated the degree of topological error via the Robinson-Foulds distance between the

true and estimated tree (87). The median background topological error for the Unlinked

simulations was 6 bipartitions (0–14), compared to 5 (0–16) for the Continuous model

(Figure 3). Error was lower for the Punctuated model with a median distance of 2

bipartitions (0-8).

Third, we calculated the uncertainty of node age reconstruction as reported by the

method via highest posterior density intervals. Under UCLN (Figure 5), the average HPD

interval for the Unlinked trees had a width 60% of the reconstructed age of its associated

node (43 – 29%). For the Continuous trees this was 108% (35 – 605%), and for the

Punctuated trees it was 70% of node age (42 – 154%). Under ACLN, the average HPD width

for Unlinked trees was 120.7% of node age (84.9 – 217.8%); for Continuous trees, 112.3%

(73.5% – 319.4%); and for Punctuated trees, 107.0% (76.7 – 144.1%).

2.4 Distribution of reconstructed vs simulated node ages

If the difference in gamma values between reconstructed and true trees is negative,

the reconstructed tree has shorter internal branches and longer terminal branches than the true

tree, while if it is positive the reconstructed tree has longer internal branches and shorter

terminal branches than the true tree. Under the UCLN, Gamma for the reconstructed

Unlinked trees was distributed evenly around the gamma of the simulated tree with a mean

value of +0.11 (-1.46 – + 1.09; Figure 4). This distribution was skewed negative (shorter tips)

relative to the simulated tree for Continuous trees with a mean of -0.44 (-1.52 – +0.20). This

negative skew was more pronounced for the Punctuated trees at -0.94 (-2.16 – +3.92). Under

ACLN, the relative change in gamma was universally negative. For the Unlinked trees it was

-2.87 (-7.13 – +5.62), while for the Continuous trees it was less negative at -1.33 (-4.92 –

+3.32), and the Punctuated trees were again skewed negative with a mean of -2.74 (-4.95 –

+2.88).

2.5 Efficacy of sister pair correlation on reconstructed branch lengths

In order to determine whether the simulated relationship between molecular rates and

diversification rates could be recovered after tree reconstruction, we regressed log clade size

contrasts against contrasts in log median branch length estimates in the data set of sister pair

analysis generated from reconstructed trees under UCLN or ACLN for each simulation

models (see 2.8). We present the p-value and sign of slope for each regression (Figure 6).

Unlinked simulated trees show no significant relationship. Significant positive relationships

are found for the Continuous and Punctuated simulations; note that this was a condition for

selection of the Continuous data set (Supplementary Methods). When reconstructed under

UCLN, the Unlinked data set and the Continuous data set show no significant relationship,

while the Punctuated data set giving a significant negative relationship, opposite to what we

expect. When reconstructed under ACLN, all three of the Unlinked, Continuous and

Punctuated data sets have significant positive relationships between clade sizes and median

branch length estimates.

3.Discussion

3.1 Errors associated with variation in speciation and substitution rate

The addition of rate-variable 'relaxed clock' models for use as rate priors represents a

significant extension to the sophistication of modern molecular dating methods (15). While

there is a wide variety of rate-variable molecular dating methods, all of them to date rely on

stochastic models of rate variation, such that rates of molecular evolution are expected to

evolve randomly over the phylogeny. However, it is clear that real substitution rates do not

always vary randomly. Instead, rates of molecular evolution are influenced by species life

history, niche and environment and macroevolutionary processes. It is important to examine

whether the stochastic models of rate variation can adequately model realistic biological

patterns of rate variation (3, 62, 65, 88). In the case of substitution rates, this means that we

must test molecular dating methods using rate simulations that are empirically based and

incorporate the effects of other biological processes. Here we have focused on testing the

impact of widely reported correlations between molecular rates and diversification on the

accuracy of molecular date reconstruction (5-8).

We used three models, Unlinked, Continuous and Punctuated, to reflect different

views on the possible links between substitution rates and diversification rates – either

varying independently; covarying continuously, linked directly by common mechanisms or

indirect through shared influences on rate variation; or linked by punctuated bursts of

increased substitutions related to speciation events. Rather than choosing parameters

arbitrarily, we parameterised our models using realistic values from the literature, or found

plausible values by tuning the models to produce data sets that reproduce empirical findings

(Supplementary Methods; 7, 79, 80). We used three different metrics (Table 2) to compare

the accuracy of phylogenetic inference using two population 'relaxed clock' molecular dating

methods, BEAST (using an uncorrelated rates model) and PAML (using an autocorrelated

rates model).

3.2 Unlinked model

The Unlinked model describes the error that can be expected from independent

variation in speciation and molecular rates. Inference on trees simulated under this model

produced median absolute errors in node age inference of about 12% on average under the

UCLN and a very high value of 93% under the ACLN (Figure 2). The largest errors were up

to 30% for UCLN and the worst instances for ACLN had errors of over 400%. The Unlinked

model produced lower errors under UCLN than the other two models, with an average error

of 12% (Figure 2). Gamma differences for the Unlinked model are distributed evenly around

zero under UCLN, so there is no overall tendency for nodes to be over- or under-estimated

(Figure 4). The sources of error in the Unlinked model are most likely to be violation of the

assumptions of rate priors and node time (tree) priors. Variation in speciation rates violates

the assumptions of the birth-death node time priors used in both BEAST 2 and PAML, which

assume universally constant rates of speciation and extinction (89). Speciation rate variation

among lineages also induces more imbalanced trees on average (90), which can lead to bias

and loss of precision in node age inferences (91). While the degree and scale on which

speciation and extinction rates vary remains contested (92-95), there is increasing evidence

that radiations and other forms of variation are a factor in the evolution of many taxa, so that

errors of this magnitude may be widespread (96-99). Substitution rates that evolve in an

autocorrelated fashion also violate the assumptions of the UCLN, which assumes that the

evolutionary rate of each lineage is independent (20), and this may also induce errors in

inference (30, 34).

The causes of the extremely high errors of the ACLN-based phylogenetic inference in

PAML for the Unlinked trees (and throughout the study for the other models) are unclear.

Examination of the output trees suggests that the most extreme errors are in trees with

shallower calibrations, and that the ages of large uncalibrated clades are sometimes

overestimated severely. The inferences in PAML and BEAST are not directly comparable

due to numerous technical differences, such as the tree topology being fixed in PAML and

BEAST placing prior distributions on speciation and extinction parameters, which are fixed

in PAML. Nevertheless it is surprising that the performance of the ACLN in PAML was

worse than UCLN in BEAST even though the underlying rate variation in these simulations

is indeed autocorrelated. There are some indications that ACLN can be difficult to fit to data

even when it is the correct model (91, 100). Another possibility is that the sensitivity of the

prior to model violation in PAML was enhanced by our use of the approximate likelihood

function (101). This is a second-order Taylor approximation to the likelihood evaluated at the

maximum likelihood estimates of the branch lengths and other parameters. As discussed by

dos Reis and Yang (101), while the approximation is very accurate under normal

circumstances when the branch lengths proposed during Bayesian inference are very far from

the maximum likelihood values the approximation becomes much less accurate. This can

happen when the rate or node time (tree) priors are misspecified. Nevertheless, absolute error

rates in other simulation studies have rarely been greater than 20% even with a deliberately

misspecified rate prior and more limited calibrating information (30), so the extreme error

rates we encountered could be down to analytical factors and not solely due to the rate prior

itself.

3.3 Continuous model

The Continuous model has substitution rates and speciation rates that coevolve

continuously rather than varying independently. Notably, we chose the degree of covariance

that reproduced two well-established empirical phenomena – the correlation of clade size and

branch length contrasts detected in a sister pairs analysis by (7), and the widespread

relationship between branch lengths and tree nodes discovered by (62). This model produced

absolute node time inference errors of 14% under UCLN, slightly higher than the Unlinked

model, but the worst errors were up to 30% (Figure 2). Negatively-skewed gamma

differences indicate that error generally leads to node ages being overestimated, pushing

reconstructed nodes back towards the root (Figure 4). In addition to the error, which is the

difference between reconstructed time estimates and the truth, we also examine how the

reconstruction methods report their uncertainty about the truth through 95% highest posterior

density intervals. Although not greatly different in terms of absolute error, the reported

uncertainty of node time estimates was very high for this model under UCLN, with average

HPD interval widths more than 107% of node height (Figure 5). This average is driven partly

by some outlying trees with very high uncertainty, up to 600% of node height; the median is

75% of node height. Wide HPD intervals can be a sign of conflict between the prior and data,

so it may be that covariance between diversification rates and substitution rates tends to

produce patterns of branch lengths that are not well described by independent rate and time

priors. The practical impact of this level of uncertainty could depend on how it is distributed

within the tree and the nature of any downstream inferences; if the question at hand is the

dating of major clades then high uncertainty in the age of tips may be tolerable, while the

same level of uncertainty in internal branches would render the tree unusable. However, any

macroevolutionary inference that relies on the distribution of nodes throughout the tree would

likely be affected (59). Under ACLN, error rates were very high (mean of 43.5%; Figure 2)

but were lower for the Continuous simulations than they were for the Unlinked simulations.

A reason for this could be that when substitution rate is positively correlated with speciation

rate it is also negatively correlated with branch lengths in units of time. Then short terminal

branches will tend to have faster rates and build up more substitutions. This in turn could

enable lower variance in estimates of rate variation across the tips of the tree, which would

aid in inferring the degree of autocorrelation.

3.4 Punctuated model

Levels of absolute error were highest under UCLN when speciation events were

associated with large punctuational bursts of molecular evolution (mean of 20.3%, up to

80.4%; Figure 2). This implies that, if real evolutionary processes were to commonly involve

such bursts of substitutions associated with speciation, or produce a similar distribution of

evolutionary histories and rate variation through some other mechanism, median node age

estimates reconstructed under the UCLN rate prior would be generally incorrect. Error rates

were also extreme for ACLN (mean 80.4%, max 397.7%; Figure 2). The finding of high

error rates in the Punctuated model corroborates a recent analysis (72), which showed that

greater proportions of branch length associated with speciation events lead to greater crown

age errors. Here, we show that the effect is strong enough to impact the median absolute error

of all node ages, even when the crown node is calibrated. The very negative skew of Gamma

away from the simulated value shows that these internal node ages are severely

overestimated, leading to very distorted trees with long tips (Figure 4).

Despite having the highest degree of branch length error under UCLN, the Punctuated

model had the lowest degree of topological error (a median of 2 bipartitions not correctly

reconstructed; Figure 3), comparing to the Unlinked and Continuous models (median of 5

and 6 respectively, Figure 3). Intuitively, this could be because rapidly diversifying clades

will normally have many short branches, making their internal relationships more uncertain.

The Punctuated model adds more substitutions to every branch regardless of its duration in

time, and so helps resolve these rapidly diversifying clades. The precise impact of topological

error is difficult to interpret, since our investigation does not distinguish among topological

errors at different levels and we do not designate any particular nodes as important. But the

practical upshot of systematically lower topological error in the simulation with the most

severe node age errors under UCLN is that these errors will be difficult to spot. Major

topological errors are easier to diagnose by checking against classical taxonomies or other

information, while little is usually known about the dates of uncalibrated nodes ahead of

time.

As an alternative method of diagnosis, we tried conducting a sister pairs analysis on

each set of simulated and reconstructed trees to see if an empirical relationship could be

recovered (Figure 6). However, tree reconstruction under either the UCLN or ACLN rate

prior disrupts the relationships between branch lengths and clade sizes displayed by the

underlying simulated trees (Figure 4). Under the UCLN they are even inverted, giving a

negative correlation between clade sizes and branch lengths. This could be related to the fact

that lineages with many speciation events have higher variance in branch lengths under

UCLN (102) which means their total length could be more strongly constrained by the node

time (tree) prior. Conversely, under ACLN we found significant positive relationships

between inferred branch lengths and clade size for all three models, even the Unlinked model

where speciation rates and substitution rates are not positively associated. A possible cause

of this inference of a positive assertion between diversification rates and substitution rate,

even in the absence of an underlying association between the two, is the node-density effect,

which results from systematic underestimation of branch lengths with high uncertainty (103).

The influence of the node density effect should be reduced by our procedure of selecting an

even number of samples in each sister clade, but may not be eliminated by it due to the

extremely high error rates under ACLN. The failure to recover the empirical correlation from

branch lengths reconstructed as part of a Bayesian dating analysis suggests that this is not a

useful way to test for the presence of interactions between substitution rates and speciation

rates. For some data sets it may be practical to check for such correlations by reconstructing

branch lengths in units of genetic divergence using a fast branch length reconstruction

method that does not impose a model of timing or rate variation (e.g. 104, 105-107).

However, even if such correlations can be diagnosed, there is currently no available method

for correcting any associated bias. This should be an area of active research.

The true role of punctuational bursts or coevolution of speciation and substitution

rates in macroevolution is currently a matter of debate. The primary empirical evidence is the

analysis of a large selection of empirical phylogenies in (63) and (62). These studies have

found that the length of root-to-tip paths in the phylogenies in substitutions per site is

frequently higher when there are more speciation events on the path. The relationship

remains even after removing the node-density effect that can generate spurious correlations

between branch lengths and node densities across clades (103, 108). Regardless of the

mechanisms, we have shown that if there were punctuated busts of substitutions associated

with speciation events, we could expect it to impact the accuracy of molecular date estimates,

since any mechanism that leads to similar patterns of rate variation and apparent correlation

between branch lengths and speciation events would produce similar effects.

3.5 Molecular dating methodology

Our study contrasts two common reconstruction methods, rather than varying aspects

of data choice. Many aspects of data choice and analytical method could exacerbate or reduce

the errors found in this analysis. For example, our alignments are shorter than many used in

actual Bayesian phylogenetic analyses but also likely more informative due to being

simulated by a simple method without invariant sites or many real sources of substitution

model misspecification. Shorter or less informative alignments could lead to greater or more

widespread error. Perhaps the most important element that we did not explore was the

influence of calibrations. Calibration choice and positioning, as well as the shape of the

calibrating distribution used, is often the most influential factor influencing divergence time

reconstruction (e.g. 32, 109) and is the source of many disagreements in published molecular

dating studies, including high-profile dating studies in birds (110-112) mammals (113, 114),

and insects (115-117). Simulation studies have suggested that calibrations can rescue

analyses strongly affected by rate prior misspecification (30), although others have found that

sufficiently misspecified rate priors cannot be rescued by more calibrations (34). The largest

and most data-rich molecular dating studies may have many more calibrations than we

implement (e.g. 139, 118). However our trees are smaller than in these studies, and our

calibrations are guaranteed to be correctly placed, have hard constraints, and are centered on

the true value. We also ensure the root is calibrated, which has been shown to have the largest

effect on accuracy in several studies (30, 34, 119). We therefore believe that our method

overall represents a balance between realism and conservatism in determining the amount of

calibrating information available to inform divergence time estimates. However, any

extension of the analyses undertaken in this study would certainly indicate investigation of

the effects of both alignment length and calibration numbers and positioning.

4. Conclusions

Our results here show the value of testing molecular phylogenetic and dating methods

against complex simulation models informed by empirically determined patterns of rate

variation. We have demonstrated that commonly applied Bayesian divergence time

estimation methods can experience average errors of 12% in node dates under a model in

which speciation and molecular rates vary independently, 14% when substitution rates are

linked to speciation rates and 20% under a punctuational model when analysed using an

ucorrelated rates prior (UCLN) in BEAST 2. Errors were much higher under all models when

trees were analysed with an autocorrelated rate prior (ACLN) in PAML. As demonstrated by

the negative skewness of gamma statistics, for trees with links between speciation and

molecular evolution these errors could lead to systematic overestimation of node ages.

Regardless of the mechanism generating the association between speciation rates and

substitution rates, we show that the potential for divergence time estimation error associated

with known empirical relationships between molecular evolution (manifest in substitution

rate) and macroevolution (manifest in the topology and branch lengths of phylogenetic trees)

can be significant and should be taken into account. Future work should see a broader range

of sophisticated divergence time estimation methods be tested against a wider variety of more

empirically realistic simulated data, including geographic and environmental biases,

concerted changes in population size, and directional changes in life history characteristics.

Ultimately the goal should be to produce new models that relax the assumption of

independence between priors on divergence times and priors on substitution rates. These will

lead to greater confidence in applications that depend on robust estimates from molecular

dating procedures.

5.Methods

5.2 Simulation trees and alignments

To simulate the phylogeny, we adopt a forward approach, starting at the root of the

phylogeny and moving forward in time, as lineages divide into two daughter lineages, go

extinct, and evolve in speciation and substitution rate as a continuous time Markov process.

In each step, we draw the time until the next event from an exponential distribution with rate

equal to the sum of the rates of all events in all lineages. Then, for every lineage l that is

present at time t, one of the following three events may take place:

(i) The lineage gives rise to two daughter lineages with rate $\lambda(l, t)$, the current speciation

rate of lineage l at time t. Both daughter lineages initially retain the same speciation rate

and trait values as their parent lineages.

(ii) The lineage becomes extinct with rate μ, which is constant over the tree.

(iii) The lineage updates its speciation rate and substitution rate with rate q, which is

constant throughout the tree. The update process adds an increment to the current

speciation rate, which is a random draw from a bivariate normal distribution with mean 0

and an instantaneous variance-covariance matrix Σ multiplied by the time since last jump.

Until one of these events happens, the lineage continues with the same speciation and

molecular rates. High values of q make for longer simulations but produce a better

approximation to Brownian motion. We found that q had no influence on observed

speciation or substitution rate variance or average tree age for $q > 10$. For our final

628   simulations, we selected $q = 50$ to allow a significant margin of safety.

630   Specifically, the instantaneous variance-covariance matrix takes the form:

631
$$\Sigma = \begin{bmatrix} \sigma_\lambda^2 & \text{Cov}(\lambda, r) \\ \text{Cov}(\lambda, r) & \sigma_r^2 \end{bmatrix}$$

where σ_λ^2 describes how fast the speciation rate evolves and σ_r^2 describes how fast the

substitution rate evolves. $\text{Cov}(\lambda, r) = \rho\sigma_\lambda\sigma_r$, where ρ is the correlation coefficient between

speciation rate and substitution rate, so $\rho = 0$ in the Unlinked Model and the Punctuated

model. We employ the Generalised Sampling Approach (120) to ensure a correct distribution

of trees with a given number of tips, which is set to 75, the approximate mean size of the

avian clades taken from (80). Details of the approach is in Supplementary Methods S1.2.

In the Punctuated model, we simulate an initial tree as per the Unlinked sample, but

without removing extinct taxa. We generate a burst of substitutions following each speciation

event. We generate Punctuated trees to match the formula found by (62), in which $16 \pm 5.4\%$

of the tree length (sum of all branch lengths) results from node-associated bursts

(Supplementary Methods S1.3).

Once we have simulated a tree, we simulate the evolution of nucleotide sequences

along the tree, given that branch lengths in the tree represent the expected number of

substitutions per site along the branch. The branch lengths in this tree represent the expected

number of substitutions per site along the branch. We simulate alignments of 2000 bases,

which is equivalent to a 6000-base coding alignment, since most substitutions in a coding

alignment occur in the third codon sites. A 6000-base coding alignment is about the size of

the concatenated coding sequences of a mitochondrial genome. Previous literature reviews

have shown this to be a common length for phylogenetic data sets in published ecological

studies, and a tractable length for large Bayesian simulation studies (86). We parameterize

our sequence simulations using realistic among-site rate variation values from birds

(Supplementary Methods S1.4).

5.3 Calibrations

In addition to the trees and sequence alignments, we need calibrations to provide

information on absolute time to allow substitution rates to be identifiable from absolute ages

of nodes. For simulated trees of similar size to ours, diminishing accuracy returns for adding

new calibrations are achieved between 1 and 5 calibrations, especially if one is at the root

(30). Since our calibration dates are known with more accuracy than those in a real analysis,

and we remove the possibility of incorrectly assigned calibrations, we choose to assign two

calibrations per tree. For each tree, we choose calibrations at the root node and at one

randomly chosen internal node. To ensure that the calibration is randomly chosen across the

nodes of the simulated trees, we first decide the level of the calibrated node as a random

integer between 1 and the largest number of nodes along any root-to-tip path in the tree. Then

we randomly pick the calibrated node from all the nodes at that level.

For analyses in both BEAST 2 and PAML, we apply a hard-bounded uniform prior

for the age of each calibrated node. Both calibrations have bounds at the true age ± 15%.

Note that this is a conservative test of molecular dating because in the simulated datasets we

have accurate knowledge of the true date of the speciation event. In reality, calibrations may

be misplaced or incorrectly dated, and may only provide maximum or minimum bounds.

5.4 Phylogenetic inference

We generate a set of 50 trees under each of the Unlinked, Continuous and Punctuated

simulation types. After sequence simulation, this gives us 50 sequence alignments for each of

the three models.We infer divergence times from the simulated sequences using common

Bayesian molecular dating procedures. First, we generate posterior distributions of tree

topologies and divergence times using two common molecular dating methods: the

uncorrelated lognormal rate prior (UCLN), implemented in the software package 'BEAST 2'

(BEAST 2; 121), in which rates for each branch are independent draws from a common

lognormal distribution, and the autocorrelated lognormal rate prior (ACLN), implemented in

the 'mcmctree' program in the PAML software suite (105), in which the rate of each branch

is lognormally distributed with a mean equal to the rate of its parent branch. To speed

analysis, we used the approximate likelihood algorithm implemented in mcmctree. The

PAML analyses differ from the BEAST analyses in that PAML requires a fixed tree

topology, which in this case was the topology of the true simulated tree. Therefore only

BEAST incorporates the possibility of topological inference errors. However, the three

simulation models can be compared within each reconstruction method. Details of the

inference method are available as Supplementary Methods (S1.5).

5.5 Error in reconstructed trees

For each reconstructed tree, we calculated accuracy in the estimation of node times

using the median absolute percentage error (MAPE) of inferred node times for clades that are

present in both true and reconstructed trees. The median was chosen rather than the mean to

reduce the impact of extremely high values for the estimates of shallow nodes. Since these

are only calculated for correctly reconstructed branches and do not capture errors in

reconstructing topology, we also calculate the Robinson-Foulds distance or number of non-

shared bipartitions, which only reflects typographical error (87).

Model misspecification in Bayesian molecular dating methods can sometimes be

associated with higher uncertainty in the reconstructed node ages, as measured by the width

of Bayesian credible intervals. This is undesirable because the reconstructed history is less

useful as an explanatory tool and less informative in downstream analyses, but it can also be

a simple way to diagnose problems with the specified model. We examined the uncertainty

with which node ages were reconstructed by the two Bayesian molecular dating methods. We

scored uncertainty by averaging the width of the reconstructed 95% highest posterior density

(HPD) intervals as a proportion of reconstructed node height across each tree, with a larger

score indicating less precise estimates.

5.6 Distribution of reconstructed vs simulated node ages

We examine the direction of error by calculating the gamma statistic for each tree and

comparing to the true tree (122). The gamma statistic is a description of how the times

between branching events in a tree whose tips are all extant taxa change as they become

closer to the present. It summarises information relating to the pattern of diversification,

following a standard normal distribution for a tree with a constant rate of speciation and no

extinction. Trees with higher values have a relatively greater proportion of their length in

older branches, while those with lower values have relatively more of their length in younger

branches. The statistic therefore tells us whether the ages of internal nodes within each tree

tend to be older or younger on average than the true times while accounting for the different

scale of this bias for older and younger nodes.

5.7 Efficacy of sister pair correlation on reconstructed branch lengths

We checked to see if we can use the reconstructed branch lengths, which are the

median of posterior branch lengths calculated by BEAST or mcmctee, to detect the

correlation between branch lengths and clade sizes found by sister pair analysis on the

simulated trees (see *Simulating realistic rate variation* above). This could be used as a rapid

post-hoc check to see whether published dated trees could be affected by reconstruction error

that is associated with substitution and speciation rate covariation. Data sets for the sister pair

analysis are generated from the 50 simulated trees under each of the three simulation models,

where each tree gives a pair of sister clades and the sister clades are the two clades that split

at the root. So, under each simulation model, we have a data set of 50 sister pairs. We then

regress contrasts in log clade sizes against contrasts in the log of the median branch length

estimates from BEAST or mcmctree.

References

1.   Zuckerkandl E, Pauling L, Bryson V, Vogel H. Evolving genes and proteins. 1965.
2.   Dickerson RE. The structure of cytochrome c and the rates of molecular evolution. J Mol Evol. 1971;1(1):26-45.
3.   Bromham L. Why do species vary in their rate of molecular evolution? Biol Lett. 2009;5:401-4.
4.   Bromham L. The genome as a life-history character: why rate of molecular evolution varies between mammal species. Phil Trans Roy Soc B. 2011;366(1577):2503-13.
5.   Barraclough TG, Savolainen V. Evolutionary rates and species diversity in flowering plants. Evolution. 2001;55:677-83.
6.   Bromham L, Hua X, Lanfear R, Cowman P. Exploring the relationships between mutation rates, life history, genome size, environment and species richness in flowering plants. Am Nat. 2015;185:507-24.
7.   Lanfear R, Ho SYW, Love D, Bromham L. Mutation rate is linked to diversification in birds. Proc Natl Acad Sci U S A. 2010;107(47):20423-8.
8.   Eo SH, DeWoody JA. Evolutionary rates of mitochondrial genomes correspond to diversification rates and to contemporary species richness in birds and reptiles. Proc R Soc B. 2010 277:3587-92.

9.   Gillman LN, Keeling DJ, Gardner RC, Wright SD. Faster evolution of highly
conserved DNA in tropical plants. J Evol Biol. 2010;23(6):1327-30.
10.   Gillman LN, Wright SD. Species richness and evolutionary speed: the influence of
temperature, water and area. J Biogeogr. 2014;41(1):39-51.
11.   Orton MG, May JA, Ly W, Lee DJ, Adamowicz SJ. Is molecular evolution faster in
the tropics? Heredity (Edinb). 2019;122(5):513-24.
12.   Ho SYW. The changing face of the molecular evolutionary clock. Trends Ecol Evol.
2014;29(9):496-503.
13.   Kumar S, Hedges SB. Advances in time estimation methods for molecular data. Mol
Biol Evol. 2016;33(4):863-9.
14.   Kumar S. Molecular clocks: four decades of evolution. Nat Rev Genet.
2005;6(8):654-62.
15.   dos Reis M, Donoghue PC, Yang Z. Bayesian molecular clock dating of species
divergences in the genomics era. Nat Rev Genet. 2016;17(2):71-80.
16.   Ho SYW, Duchêne S. Molecular-clock methods for estimating evolutionary rates and
timescales. Mol Ecol. 2014;23(24):5947-65.
17.   Heath TA, Holder MT, Huelsenbeck JP. A Dirichlet process prior for estimating
lineage-specific substitution rates. Mol Biol Evol. 2012;29(3):939-55.
18.   Yoder AD, Yang Z. Estimation of primate speciation dates using local molecular
clocks. Mol Biol Evol. 2000;17(7):1081-90.
19.   Drummond AJ, Suchard MA. Bayesian random local clocks, or one rate to rule them
all. BMC Biol. 2010;8(1):114.
20.   Drummond AJ, Ho SYW, Phillips MJ, Rambaut A. Relaxed phylogenetics and dating
with confidence. PLOS Biol. 2006;4(5):e88.
21.   Sanderson MJ. r8s: inferring absolute rates of molecular evolution and divergence
times in the absence of a molecular clock. Bioinformatics. 2003;19(2):301-2.
22.   Smith SA, O'Meara BC. treePL: divergence time estimation using penalized
likelihood for large phylogenies. Bioinformatics. 2012;28(20):2689-90.
23.   Sauquet H. A practical guide to molecular dating. Comptes Rendus Palevol.
2013;12(6):355-67.
24.   Wilke T, Schultheiß R, Albrecht C. As time goes by: a simple fool's guide to
molecular clock approaches in invertebrates. Am Malac B. 2009;27(1/2):25-45.
25.   Thorne JL, Kishino H, Painter IS. Estimating the rate of evolution of the rate of
molecular evolution. Mol Biol Evol. 1998;15(12):1647-57.
26.   Kishino H, Thorne JL, Bruno WJ. Performance of a divergence time estimation
method under a probabilistic model of rate evolution. Mol Biol Evol. 2001;18(3):352-61.
27.   Aris-Brosou S, Yang Z. Effects of models of rate evolution on estimation of
divergence dates with special reference to the metazoan 18S ribosomal RNA phylogeny. Syst
Biol. 2002;51(5):703-14.
28.   Lepage T, Lawi S, Tupper P, Bryant D. Continuous and tractable models for the
variation of evolutionary rates. Math Biosci. 2006;199(2):216-33.
29.   Sarver BAJ, Pennell MW, Brown JW, Keeble S, Hardwick KM, Sullivan J, et al. The
choice of tree prior and molecular clock does not substantially affect phylogenetic inferences
of diversification rates. PeerJ. 2019;7.
30.   Duchêne S, Lanfear R, Ho SYW. The impact of calibration and clock-model choice
on molecular estimates of divergence times. Mol Phylogenet Evol. 2014;78:277-89.
31.   Lepage T, Bryant D, Philippe H, Lartillot N. A general comparison of relaxed
molecular clock models. Mol Biol Evol. 2007;24(12):2669-80.

32.   Foster CSP, Sauquet H, van der Merwe M, McPherson H, Rossetto M, Ho SYW.
Evaluating the impact of genomic data and priors on Bayesian estimates of the angiosperm
evolutionary timescale. Syst Biol. 2017;66(3):338-51.
33.   Smith AB, Pisani D, Mackenzie-Dodds JA, Stockley B, Webster BL, Littlewood DTJ.
Testing the molecular clock: molecular and paleontological estimates of divergence times in
the Echinoidea (Echinodermata). Mol Biol Evol. 2006;23(10):1832-51.
34.   Battistuzzi FU, Filipski A, Hedges SB, Kumar S. Performance of relaxed-clock
methods in estimating evolutionary divergence times and their credibility intervals. Mol Biol
Evol. 2010;27(6):1289-300.
35.   Worobey M, Han G-Z, Rambaut A. A synchronized global sweep of the internal
genes of modern avian influenza virus. Nature. 2014;508(7495):254-7.
36.   Crisp MD, Hardy NB, Cook LG. Clock model makes a large difference to age
estimates of long-stemmed clades with no internal calibration: a test using Australian
grasstrees. BMC Evol Biol. 2014;14:263-.
37.   dos Reis M, Thawornwattana Y, Angelis K, Telford Maximilian J, Donoghue
Philip CJ, Yang Z. Uncertainty in the timing of origin of animals and the limits of precision
in molecular timescales. Curr Biol. 2015;25(22):2939-50.
38.   Welch JJ, Bromham L. Molecular dating when rates vary. Trends in Ecology &
Evolution. 2005;20(6):320-7.
39.   Lin G, Huang Z, Wang L, Chen Z, Zhang T, Gillman LN, et al. Evolutionary rates of
bumblebee genomes are faster at lower elevations. Mol Biol Evol. 2019;36(6):1215-9.
40.   Lourenço JM, Glémin S, Chiari Y, Galtier N. The determinants of the molecular
substitution process in turtles. J Evol Biol. 2013;26(1):38-50.
41.   Goldie X, Gillman L, Crisp M, Wright S. Evolutionary speed limited by water in arid
Australia. Proc R Soc B. 2010;277(1694):2645-53.
42.   Berv JS, Field DJ. Genomic signature of an avian Lilliput effect across the K-Pg
extinction. Syst Biol. 2018;67(1):1-13.
43.   Barrera-Redondo J, Ramirez-Barahona S, Eguiarte LE. Rates of molecular evolution
in tree ferns are associated with body size, environmental temperature, and biological
productivity. Evolution. 2018;72(5):1050-62.
44.   Qiu F, Kitchen A, Burleigh JG, Miyamoto MM. Scombroid fishes provide novel
insights into the trait/rate associations of molecular evolution. J Mol Evol. 2014;78(6):338-
48.
45.   May JA, Feng Z, Orton MG, Adamowicz SJ. The effects of ecological traits on the
rate of molecular evolution in ray-finned fishes: a multivariable approach. J Mol Evol.
2020;88(8-9):689-702.
46.   Welch JJ, Bininda-Emonds ORP, Bromham L. Correlates of substitution rate
variation in mammalian protein-coding sequences. BMC Evol Biol. 2008;8(1):53.
47.   Thomas JA, Welch JJ, Lanfear R, Bromham L. A generation time effect on the rate of
molecular evolution in invertebrates. Mol Biol Evol. 2010;27(5):1173-80.
48.   Hua X, Cowman P, Warren D, Bromham L. Longevity is linked to mitochondrial
mutation rates in rockfish: a test using Poisson regression. Mol Biol Evol. 2015;32(10):2633-
45.
49.   Galtier N, Blier PU, Nabholz B. Inverse relationship between longevity and
evolutionary rate of mitochondrial proteins in mammals and birds. Mitochondrion.
2009;9(1):51-7.
50.   Bromham L, Cowman PF, Lanfear R. Parasitic plants have increased rates of
molecular evolution across all three genomes. BMC Evol Biol. 2013;13(1):126.
51.   Mitterboeck TF, Adamowicz SJ. Flight loss linked to faster molecular evolution in
insects. Proc R Soc B. 2013;280(1767):20131128.

52.   Wong A. Covariance between testes size and substitution rates in primates. Mol Biol Evol. 2014;31(6):1432-6.

53.   Woolfit M, Bromham L. Population size and molecular evolution on islands. Proc Biol Sci. 2005;272(1578):2277-82.

54.   Lynch M, Ackerman MS, Gout J-F, Long H, Sung W, Thomas WK, et al. Genetic drift, selection and the evolution of the mutation rate. Nat Rev Genet. 2016;17(11):704-14.

55.   Gossmann TI, Keightley PD, Eyre-Walker A. The effect of variation in the effective population size on the rate of adaptive molecular evolution in eukaryotes. Genome Biol Evol. 2012;4(5):658-67.

56.   Charlesworth B. Fundamental concepts in genetics: effective population size and patterns of molecular evolution and variation. Nat Rev Genet. 2009;10(3):195-205.

57.   Mindell DP, Sites Jr JW, Graur D. Speciational evolution: a phylogenetic test with allozymes in Sceloporus (Reptilia). Cladistics. 1989;5(1):49-61.

58.   Iglesias-Carrasco M, Jennions MD, Ho SYW, Duchene DA. Sexual selection, body mass and molecular evolution interact to predict diversification in birds. Proc R Soc B. 2019;286(1899).

59.   Duchêne DA, Hua X, Bromham L. Phylogenetic estimates of diversification rate are affected by molecular rate variation. J Evol Biol. 2017;30(10):1884–97.

60.   Lancaster LT. Molecular evolutionary rates predict both extinction and speciation in temperate angiosperm lineages. BMC Evol Biol. 2010;10(1):162.

61.   Fontanillas E, Welch JJ, Thomas JA, Bromham L. The influence of body size and net diversification rate on molecular evolution during the radiation of animal phyla. BMC Ecol Evol. 2007;7:95.

62.   Pagel M, Venditti C, Meade A. Large punctuational contribution of speciation to evolutionary divergence at the molecular level. Science. 2006;314(5796):119.

63.   Webster AJ, Payne RJH, Pagel M. Molecular phylogenies link rates of evolution and speciation. Science. 2003;301:478.

64.   Ezard THG, Thomas GH, Purvis A. Inclusion of a near-complete fossil record reveals speciation-related molecular evolution. Methods Ecol Evol. 2013;4(8):745-53.

65.   Hua X, Bromham L. Darwinism for the genomic age: connecting mutation to diversification. Front Genet. 2017;8:12.

66.   Dobzhansky T. Genetics and the origin of species. New York: Columbia University Press; 1937.

67.   Muller H, editor Isolating mechanisms, evolution, and temperature. Biol Symp; 1942.

68.   Gavrilets S. Fitness landscapes and the origin of species (MPB-41): Princeton University Press; 2004.

69.   Orr HA. The population genetics of speciation: the evolution of hybrid incompatibilities. Genetics. 1995;139(4):1805-13.

70.   Frankham R. Genetics and extinction. Biol Cons. 2005;126(2):131-40.

71.   Venditti C, Pagel M. Speciation as an active force in promoting genetic evolution. Trends Ecol Evol. 2010;25(1):14-20.

72.   Janzen T, Bokma F, Etienne RS. Nucleotide substitutions during speciation may explain substitution rate variation. bioRxiv. 2021:2020.08.19.256891.

73.   Bromham L, Duchêne S, Hua X, Ritchie AM, Duchêne DA, Ho SYW. Bayesian molecular dating: opening up the black box. Biol Rev. 2018;93(2):1165-91.

74.   Warnock RCM, Parham JF, Joyce WG, Lyson TR, Donoghue PCJ. Calibration uncertainty in molecular dating analyses: there is no substitute for the prior evaluation of time priors. Proc Biol Sci. 2015;282(1798):20141013-.

75.   Yang Z, Rannala B. Bayesian estimation of species divergence times under a molecular clock using multiple fossil calibrations with soft bounds. Mol Biol Evol. 2005;23(1):212-26.

76.   Zhu T, Dos Reis M, Yang Z. Characterization of the uncertainty of divergence time estimation under relaxed molecular clock models using multiple loci. Syst Biol. 2015;64(2):267-80.

77.   dos Reis M, Yang Z. The unbearable uncertainty of Bayesian divergence time estimation. J Systemat Evol. 2013;51(1):30-43.

78.   Manceau M, Marin J, Morlon H, Lambert A. Model-based inference of punctuated molecular evolution. Mol Biol Evol. 2020;37(11):3308-23.

79.   Nabholz B, Lanfear R, Fuchs J. Body mass-corrected molecular rate for bird mitochondrial DNA. Mol Ecol. 2016;25(18):4438-49.

80.   Maliet O, Hartig F, Morlon H. A model with many small shifts for estimating species-specific diversification rates. Nat Ecol Evol. 2019;3(7):1086-92.

81.   Lanfear R, Ho SYW, Love D, Bromham L. Mutation rate influences diversification rate in birds. Proc Natl Acad Sci U S A. 2010;107(47):20423-8.

82.   Rabosky DL. Positive correlation between diversification rates and phenotypic evolvability can mimic punctuated equilibrium on molecular phylogenies. Evolution. 2012;66(8):2622-7.

83.   Pennell MW, Harmon LJ, Uyeda JC. Is there room for punctuated equilibrium in macroevolution? Trends Ecol Evol. 2014;29(1):23-32.

84.   Pennell MW, Harmon LJ, Uyeda JC. Speciation is unlikely to drive divergence rates. Trends Ecol Evol. 2014;29(2):72-3.

85.   Jetz W, Thomas GH, Joy JB, Hartmann K, Mooers AO. The global diversity of birds in space and time. Nature. 2012;491:444.

86.   Ritchie AM, Hua X, Cardillo M, Yaxley KJ, Dinnage R, Bromham L. Phylogenetic diversity metrics from molecular phylogenies: modelling expected degree of error under realistic rate variation. Divers Distrib. 2020;n/a(n/a).

87.   Robinson DF, Foulds LR. Comparison of phylogenetic trees. Math Biosci. 1981;53(1):131-47.

88.   Bromham L. The genome as a life-history character: why rate of molecular evolution varies between mammal species. Phil Trans Roy Soc B. 2011;366(1577):2503-13.

89.   Gernhard T. The conditioned reconstructed process. J Theor Biol. 2008;253(4):769-78.

90.   Stadler T. Recovering speciation and extinction dynamics based on phylogenies. J Evol Biol. 2013;26(6):1203-19.

91.   Duchêne D, Duchêne S, Ho SYW. Tree imbalance causes a bias in phylogenetic estimation of evolutionary timescales using heterochronous sequences. Mol Ecol Res. 2015;15(4):785-94.

92.   Hedges SBK, Sudhir The Timetree of Life. New York: Oxford University Press; 2009.

93.   Hedges SB, Marin J, Suleski M, Paymer M, Kumar S. Tree of life reveals clock-like speciation and diversification. Mol Biol Evol. 2015;32(4):835-45.

94.   McPeek MA, Brown JM. Clade age and not diversification rate explains species richness among animal taxa. Am Nat. 2007;169(4):E97-106.

95.   Marin J, Rapacciuolo G, Costa GC, Graham CH, Brooks TM, Young BE, et al. Evolutionary time drives global tetrapod diversity. Proc R Soc B. 2018;285(1872):20172378.

96.   Cardillo M. Latitude and rates of diversification in birds and butterflies. Proc R Soc B. 1999;266(1425):1221-5.

97.   Alfaro ME, Santini F, Brock C, Alamillo H, Dornburg A, Rabosky DL, et al. Nine exceptional radiations plus high turnover explain species diversity in jawed vertebrates. Proc Natl Acad Sci U S A. 2009;106(32):13410-4.

98.   Kozak KH, Wiens JJ. Testing the relationships between diversification, species richness, and trait evolution. Syst Biol. 2016;65(6):975-88.

99.   Scholl JP, Wiens JJ. Diversification rates and species richness across the Tree of Life. Proc R Soc B. 2016;283(1838):20161334.

100.   Ho SY, Duchêne S, Duchêne D. Simulating and detecting autocorrelation of molecular evolutionary rates among lineages. Mol Ecol Resour. 2015;15(4):688-96.

101.   dos Reis M, Yang Z. Approximate likelihood calculation on a phylogeny for Bayesian estimation of divergence times. Mol Biol Evol. 2011;28(7):2161-72.

102.   Guindon S. Rates and rocks: strengths and weaknesses of molecular dating methods. Front Genet. 2020;11:526-.

103.   Hugall AF, Lee MS. The likelihood node density effect and consequences for evolutionary studies of molecular rates. Evolution. 2007;61(10):2293-307.

104.   Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics. 2014;30(9):1312-3.

105.   Yang ZH. PAML 4: Phylogenetic analysis by maximum likelihood. Mol Biol Evol. 2007;24(8):1586-91.

106.   Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol. 2014;32(1):268-74.

107.   Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. Syst Biol. 2010;59(3):307-21.

108.   Venditti C, Meade A, Pagel M. Detecting the node-density artifact in phylogeny reconstruction. Syst Biol. 2006;55(4):637-43.

109.   Sauquet H, Ho SYW, Gandolfo MA, Jordan GJ, Wilf P, Cantrill DJ, et al. Testing the impact of calibration on molecular divergence times using a fossil-rich group: the case of Nothofagus (Fagales). Syst Biol. 2011;61(2):289-313.

110.   Jarvis ED, Mirarab S, Aberer AJ, Li B, Houde P, Li C, et al. Whole-genome analyses resolve early branches in the tree of life of modern birds. Science. 2014;346(6215):1320-31.

111.   Mitchell KJ, Cooper A, Phillips MJ. Comment on "Whole-genome analyses resolve early branches in the tree of life of modern birds". Science. 2015;349(6255):1460-.

112.   Cracraft J, Houde P, Ho SYW, Mindell DP, Fjeldså J, Lindow B, et al. Response to Comment on "Whole-genome analyses resolve early branches in the tree of life of modern birds". Science. 2015;349(6255):1460-.

113.   Meredith RW, Janečka JE, Gatesy J, Ryder OA, Fisher CA, Teeling EC, et al. Impacts of the Cretaceous terrestrial revolution and KPg extinction on mammal diversification. Science. 2011;334(6055):521-4.

114.   Phillips MJ. Geomolecular dating and the origin of placental mammals. Syst Biol. 2016;65(3):546-57.

115.   Misof B, Liu S, Meusemann K, Peters RS, Donath A, Mayer C, et al. Phylogenomics resolves the timing and pattern of insect evolution. Science. 2014;346(6210):763-7.

116.   Tong KJ, Duchêne S, Ho SYW, Lo N. Comment on "Phylogenomics resolves the timing and pattern of insect evolution". Science. 2015;349(6247):487-.

117.   Kjer KM, Ware JL, Rust J, Wappler T, Lanfear R, Jermiin LS, et al. Response to Comment on "Phylogenomics resolves the timing and pattern of insect evolution". Science. 2015;349(6247):487-.

118.   Rabosky DL, Chang J, Title PO, Cowman PF, Sallan L, Friedman M, et al. An
inverse latitudinal gradient in speciation rate for marine fishes. Nature. 2018;559(7714):392-
5.
119.   Wertheim JO, Sanderson MJ. Estimating diversification rates: how useful are
divergence times? Evolution. 2011;65(2):309-20.
120.   Hartmann K, Wong D, Stadler T. Sampling trees from evolutionary models. Syst
Biol. 2010;59(4):465-76.
121.   Bouckaert R, Vaughan TG, Barido-Sottani J, Duchêne S, Fourment M, Gavryushkina
A, et al. BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis.
PLOS Comp Biol. 2019;15(4):e1006650.
122.   Pybus OG, Harvey PH. Testing macro-evolutionary models using incomplete
molecular phylogenies. Proc Biol Sci. 2000;267(1459):2267-72.
1018
1019

1020

1021

1022

1023

1024

1025

1026

1027

1028

1029

1030

1031

1032

1033
1034
1035

Figure 1. Flowchart of study design. Trees with branch lengths in units of time are

simulated under three simulation conditions. For the Unlinked model, birth rates (λ)and

substitution rates (sub) vary independently, while for the Continuous model they are

correlated. For the Punctuated model, trees are simulated as for Unlinked, then some of the

branch length is redistributed to form bursts of substitutions occurring upon speciation. Trees

and branch lengths are then used to simulate molecular sequence alignments. Time trees are

then reconstructed under two rate priors, UCLN and ACLN, and compared to the original,

simulated trees to determine the error (Medain abs % error), uncertainty (HPD width) and

bias towards over- or under-estimation (Gamma).

Figure 2. Median absolute percentage error of node times (on log scale) reconstructed from data sets simulated under three simulation types with different relationships between molecular evolution and speciation. The mean error is calculated only for branches whose bipartitions are shared between the true and inferred trees, and therefore ignores errors in topology which may occur in the BEAST analyses. The three simulation models are Unlinked (instantaneous covariance of molecular rates and speciation rates = 0), Continuous (instantaneous covariance = 0.0044), and Punctuated (instantaneous covariance = 0, bursts of substitutions added at speciation events). Topologies and node times were reconstructed using two different analytical methods, the uncorrelated lognormal 'relaxed clock' rate prior (UCLN) in BEAST 2, and the autocorrelated lognormal rate prior (ACLN) in PAML.

1058
1059
Figure 3. Topological error of trees reconstructed from data sets simulated under three

simulation types with different relationships between molecular evolution and speciation.

Topological error is given by the Robinson-Foulds distance (number of non-shared

bipartitions). This distance incorporates both topological and branch time errors. The three

simulation models are Unlinked (instantaneous covariance of molecular rates and speciation

rates = 0), Continuous (instantaneous covariance = 0.0044), and Punctuated (instantaneous

covariance = 0, bursts of substitutions added at speciation events). Topologies and node times

were reconstructed using the uncorrelated lognormal 'relaxed clock' rate prior (UCLN) in

BEAST 2. No results are available for the autocorrelated lognormal rate prior in PAML

because the topology was fixed for the analysis.

1070

1071
1072

Figure 4. Difference in gamma statistic values between reconstructed and simulated trees,

where simulated trees were generated under three models with different relationships

between molecular evolution and speciation. Negative scores indicate that reconstructed

trees have a greater proportion of their branch times in tips vs internal branches ('tippier')

than the simulated tree. The three simulation models are Unlinked (instantaneous covariance

of molecular rates and speciation rates = 0), Continuous (instantaneous covariance = 0.0044),

and Punctuated (instantaneous covariance = 0, bursts of substitutions added at speciation

events). Topologies and node times were reconstructed using two different analytical

methods, the uncorrelated lognormal 'relaxed clock' rate prior (UCLN) in BEAST 2, and the

autocorrelated lognormal rate prior (ACLN) in PAML.

Figure 5. Uncertainty of trees reconstructed from phylogenies simulated under three models

with different relationships between molecular evolution and speciation, in width as a

percentage of reconstructed node age. Precision is measured by averaging the width of 95%

highest posterior density (HPD) intervals as a proportion of reconstructed node height across

each tree, with a larger score indicating less precise estimates. The three simulation models

are Unlinked (instantaneous covariance of molecular rates and speciation rates = 0),

Continuous (instantaneous covariance = 0.0044), and Punctuated (instantaneous covariance =

0, bursts of substitutions added at speciation events). Topologies and node times were

reconstructed using two different analytical methods, the uncorrelated lognormal 'relaxed

clock' rate prior (UCLN) in BEAST 2, and the autocorrelated lognormal rate prior (ACLN)

in PAML.

Figure 6. Slope and significance of mock sister pair analyses of clade size contrasts against

contrasts in simulated branch lengths (SIM) or median branch length estimates reconstructed

from simulated trees. Results are shown for branch lengths reconstructed under the

uncorrelated lognormal clock (UCLN) or autocorrelated lognormal clock (ACLN). The three

simulation models are Unlinked (instantaneous covariance of molecular rates and speciation

rates = 0), Continuous (instantaneous covariance = 0.0044), and Punctuated (instantaneous

covariance = 0, bursts of substitutions added at speciation events). Black triangles represent

positive regression coefficients, while white inverted triangles represent negative coefficients.

The red line indicates significance at the 0.05 level.