Investigating the reliability of molecular estimates of evolutionary time when

substitution rates and speciation rates vary

ANDREW M. RITCHIE[1][*], XIA HUA[1][2], LINDELL BROMHAM[1]

1. *Research School of Biological Sciences, Australian National University, Canberra,*

ACT 2600, Australia

2. *Mathematical Sciences Institute, Australian National University, Canberra, ACT*

2600, Australia

* Corresponding Author:

Andrew M. Ritchie

Research School of Biological Sciences, Robertson Building, 134 Linnaeus Way

Australian National University, Canberra, ACT 2600, Australia

Ph: +61 2 6125 5111

Email: Andrew.Ritchie@anu.edu.au

17
18
19
20
Abstract

<u>Background</u>

An accurate timescale of evolutionary history is essential to testing hypotheses

about the influence of historical events and processes, and the timescale for evolution is

increasingly derived from analysis of DNA sequences. But variation in the rate of molecular

evolution complicates the inference of time from DNA. Evidence is growing for numerous

factors, such as life history and habitat, that are linked both to the molecular processes of

mutation and fixation and to rates of macroevolutionary diversification. However, the most

widely used methods rely on idealised models of rate variation, such as the uncorrelated and

autocorrelated clocks, and molecular dating methods are rarely tested against complex

models of rate change. One relationship that is not accounted for in molecular dating is the

potential for interaction between molecular substitution rates and speciation, a relationship

that has been supported by empirical studies in a growing number of taxa. If these

relationships are as widespread as current evidence suggests, they may have a significant

influence on molecular dates.

Results

We simulate phylogenies and molecular sequences under three different realistic rate

variation models – one in which speciation rates and substitution rates both vary but are

unlinked, one in which they covary continuously and one punctuated model in which

molecular change is concentrated in speciation events, using empirical case studies to

parameterise realistic simulations. We test three commonly used "relaxed clock" molecular

dating methods against these realistic simulations to explore the degree of error in molecular

dates under each model. We find average divergence time inference errors ranging from 12%

of node age for the unlinked model when reconstructed under an uncorrelated rate prior using

BEAST 2, to up to 91% when sequences evolved under the punctuated model are

reconstructed under an autocorrelated prior using PAML.

Conclusions

We demonstrate the potential for substantial errors in molecular dates when both

speciation rates and substitution rates vary between lineages. This study highlights the need

for tests of molecular dating methods against realistic models of rate variation generated from

empirical parameters and known relationships.

Keywords

Molecular dating, substitution rates, diversification rates, Bayesian phylogenetics

1.Background

Understanding the timescale of evolution is critical to researching the processes that

generate and shape the diversity of life on Earth. Analysis of DNA sequences offers the

possibility of investigating the tempo and mode of evolution of all extant lineages, not just

those that have a detailed fossil record. The earliest molecular dating analyses concluded that

the average rate of molecular evolution was relatively constant over time in some proteins

because of an apparently linear accumulation of amino acid differences when compared

between species (1, 2). However, it has since become apparent that the average rate of

molecular evolution can be influenced by a wide range of species traits, such as generation

time, body size and longevity (3, 4), by macroevolutionary processes such as net

diversification rate (5-10) and potentially also by environmental features like temperature (9-

11). If the average rate of molecular evolution varies substantially and consistently between

lineages, then we may need to incorporate this rate variation into models of molecular

evolution in order to accurately infer divergence times among species (12).

A wide variety of models have been developed to account for different patterns of

among-lineage molecular rate variation (13-16). In the first instance, these models differ in

how many different evolutionary rate parameters are estimated for different parts of the

phylogeny. For example, a 'strict clock' model estimates only one rate parameter that applies

to every branch in the tree, while a 'local clock' model estimates additional rate parameters

for subsets of branches (17-19), and a 'relaxed clock' provides for a separate rate on every

branch (20). Models also differ in how the rates on different branches are related to one

another. Some models, notably in the penalized-likelihood framework, aim to minimize the

overall variation of branch-specific rates or the differences between adjacent branches (21,

22). For Bayesian phylogenetic methods (23, 24), branch-specific rates are drawn from a

prior probability distribution that determines which rates and patterns of relative rate

variation are considered *a priori* more probable (we refer to this prior as the rate prior).

Common variants include uncorrelated models, in which each branch-specific rate is an

independent draw from a distribution with a common mean (20), and autocorrelated models,

in which the mean of the distribution for each branch-specific rate depends on the rate drawn

for its parent branch (25, 26).

Testing over the last two decades has generally supported the robustness of

divergence time estimates under the Bayesian methodology to the choice of rate prior,

provided that enough separate rate parameters are allowed and sufficient calibrating

information is provided (27-33); but see (34). However, these studies rely heavily on

simulated data or on specific and usually well-calibrated empirical examples, whereas

simulations have indicated that different rate priors may give different results when

calibrating information is poorer (30). Changing the rate prior in these circumstances could

lead to differences in empirical dates and associated causal hypotheses. Examples of

empirical studies where the choice of rate prior appears to be a factor in differing age

estimates include influenza viruses (35), the age of grasstrees (36) and the origins of Metazoa

(37).

Currently used models of rate variation are generally either uncorrelated, so that each

lineage-specific rate is drawn independently from a common distribution, or autocorrelated

so that lineage-specific rates evolve from the rates of their parent lineages according to a

stochastic, random walk. However, the factors shaping variation in rate of molecular

evolution are complex: in addition to random variation in rates between lineages, there is

clearly also systematic variation influenced by many factors including species traits,

population dynamics, environment and evolutionary history (38). Evolutionary rates can be

faster at lower elevations and latitudes (9, 39, 40) but slower in arid climates (41). Life

history can covary with molecular rates across multiple axes, including body size (42-45),

fecundity (46), generation time (47) and longevity (48, 49). Associations have also been

observed between substitution rates and more complex species traits, such as parasitism (50),

flightlessness (51) and sexual competition (52). Furthermore, substitution rates are the result

of underlying microevolutionary processes, and may therefore be affected by population size

(53-56).

In addition to covarying with other evolving traits, there is evidence that variation in

molecular rates can be associated with macroevolutionary patterns. Relationships between the

relative size of sister clades, representing their net diversification rates, and their molecular

rates have been detected in several studies, include in reptiles (8, 57), birds (7, 58), flowering

plants (6, 59, 60) and across Metazoa (61). Several studies have also noted relationships

between the genetic divergence of species in a phylogeny and the number of intervening

speciation events, which may also represent examples of this phenomenon (62-64).

The mechanism behind observed correlations between diversification and molecular

evolution remains a matter of debate. One line of argument proposes that higher molecular

rates could promote speciation by speeding the development of reproductive isolation (65). In

this hypothesis, subpopulations become sporadically isolated over time and may accumulate

alleles that may be effectively neutral on their own but deleterious in combination (66, 67). In

this case, greater rates of molecular evolution could lead to faster development of

reproductive isolation during periods of isolation, leading to a greater average rate of

speciation (68, 69). Alternatively, higher substitution rates could reduce extinction rates by

accumulating more standing genetic diversity which provides a buffer against changing

environment by allowing more rapid selective response (70). Speciation rates and molecular

rates could also be related through a common driving factor, such as climate or

environmental productivity (5, 10). In the other causal direction, speciation itself could

increase average rates of molecular evolution by producing rapid bursts of evolution (62, 63,

71). Suggested mechanisms for an increase in substitution rates during speciation include an

increase in fixation rates of mildly deleterious alleles and linked neutral mutations due to

drift, or large-scale genomic divergence caused by an increase in the frequency of disruptive

genomic events such as polyploidy or chromosomal rearrangements (71).

An association between substitution rates and diversification rates could affect the

performance of existing divergence time estimation methods (e.g. 72). In particular, Bayesian

methods divide their specification of prior beliefs about the evolutionary process between the

rate prior, the prior on divergence times and tree topology (sometimes referred to as the tree

prior or node time prior), and calibrating distributions on the ages of individual speciation

events (23, 73). These priors are essential to estimating divergence time, because genetic

distances between sequences are the product of rate and time (the total number of character

differences between related sequences). This means that even if genetic distances are

correctly inferred on a phylogeny, rate and time are unidentifiable without additional

constraints on joint values they can hold (74, 75). The rate prior, node time prior and

calibrating distributions place either soft or hard constraints on these values, allowing

researchers to infer evolutionary history on an absolute timescale. However, this means that

inferred divergence times are always subject to the choice of rate, time and calibration priors

even with large amounts of sequence data (76, 77). Most commonly used Bayesian molecular

dating methods do not propose *a priori* relationships between the rate prior and the tree prior,

so the joint distribution implied by these priors will assign relatively higher probability to l

long branches with high substitution rates and short branches with low substitution rates than

would be expected when rates and times are related. It is therefore possible that an

unmodelled relationship between rates and times could produce distributions of molecular

rates and speciation times that differ substantially from those implied by common Bayesian

priors, leading to widespread error in estimates of evolutionary dates.

Here, we investigate the potential influence of speciation-related rates of molecular

evolution on the reliability of molecular dating studies. Rather than assuming only one

specific model of the association between diversification rates and rates of molecular

evolution, we model a range of possible causal links. We include three alternative models of

the association between rates of molecular evolution and speciation rates. In the first, both

speciation rates and substitution rates vary over the phylogeny, but neither influences the

other. In the second, we model continuous variation in a linked manner, which might

represent either direct influence of one on the other (65) or an indirect association between

speciation rates and substitution rates, for example if they are both influenced by

environmental factors (5, 10). The third model is a punctuated model in which bursts of

substitution are associated with speciation events (62, 63, 71, 78). The proportion of all

substitutions associated with such bursts has recently been shown to impact Bayesian

inference of clade crown ages under an uncorrelated lognormal rate prior (72).

In this study, we simulate the evolution of molecular sequences under all three models

of the link between speciation and rate of molecular evolution – unlinked, continuous

coevolution and punctuated models – and observe the average effects on errors in Bayesian

divergence time estimates. Most simulation studies use arbitrarily chosen values to

parameterize the simulations. In order to make our test have real-world significance, we base

our simulations, as far as possible, on empirically determined relationships between rates of

molecular evolution and diversification rate. As a convenient well-studied case study, we

parameterize our models using values taken from empirical inferences on bird data (79, 80),

or by using parameter values which reproduce established empirical relationships between

rates of molecular evolution and diversification rate in birds (81). We have chosen birds as a

convenient exemplar since an association between rates of molecular evolution and rates of

diversification has been reported for birds (58), and there are few if any other taxa whose

rates of molecular evolution and diversification have been as well-described. We test the

performance of three commonly used "relaxed clock" molecular dating methods: since the

choice of rate prior and methodological details may affect the results of phylogeny

reconstruction (34), we have tested uncorrelated and autocorrelated rate models in two

different molecular dating programs. We reconstruct divergence times on these simulated

data sets and examine the effects of different forms of correlated molecular and speciation

rates on the performance of molecular divergence time reconstruction.

2. Results

2.1 Validation of tree simulations

In order to ensure our simulations are as close as possible to real-world data, we chose

to parameterize the simulations using empirical estimates of parameters from bird data,

because birds are one of the most well-studied groups for both substitution rate variation and

diversification rate. While these parameters might not exactly capture processes in other taxa,

we believe that basing the simulations on a real case study lends a degree of realism that

would be lacking if we arbitrarily chose convenient parameter values. We checked that the

simulation procedure produced trees with average characteristics similar to the avian clades

used to parameterize the study (80 [supplementary data], 82). Across all data sets, the mean

root date of simulated trees was 28.4 million years, compared to a range of 23.8-35.6 million

years for bird clades with 70-80 taxa. Simulated lineage-specific speciation rates at the tips

were on average slightly lower than the speciation rate of similar clades in the bird tree (80).

The range of simulated speciation rates on log scale was 0.009-0.92 compared to a range of

0.034-2.25 average log lineages/million years for bird clades with 70-80 taxa, similar taxon

numbers to our simulated trees. The range of speciation rates within each tree was also

similar to the range across the avian clades with similar taxon numbers, with an average

magnitude of 0.44 log lineages/million years compared to 0.32-2.21 for avian clades with 70-

80 taxa. These observations suggest that the simulations produce realistic phylogenies,

similar to those associated with real taxa.

2.2 Error in reconstructed trees

Trees were simulated under three models: Unlinked (speciation rates and molecular

rates vary independently), Continuous (speciation rates and molecular rates covary), and

Punctuated (molecular change occurs in bursts associated with speciation events as well as

continuously within lineages) (Table 1). We used three different metrics, plus the gamma

statistic and a sister pair analysis, to assess the accuracy of the reconstruction of trees

simulated from the Unlinked model, the Continuous model, and the Punctuated model, under

uncorrelated lognormal (UCLN) and autocorrelated lognormal (ACLN) models of among-

lineage rate variation (Table 2).

Table 1. Simulation design.
238

Simulation type	Speciation rates covary with substitution rates?	Brownian motion covariance	Punctuated burst of substitutions at nodes?
Unlinked	*No*	0	*no*
Continuous	*Yes*	0.0044	*no*
Punctuated	*No*	0	*yes*

239
240   We lay out the conditions for each set of simulated trees. Each set consisted of 50 trees.
241   Lineage-specific speciation rates and substitution rates evolve continuously through time via
242   Brownian motion in all three studies. The three simulations differ in whether the covariance
243   between speciation and substitution rates is greater than zero in the Brownian motion and
244   whether an additional burst of substitutions occurs at nodes of the tree (speciation events).
245

Node dates were then inferred using three different reconstruction methods: BEAST 2 (83,

84) using the uncorrelated lognormal rate prior (UCLN); PAML (85) using the

autocorrelated lognormal rate prior (ACLN); and PAML using UCLN. To compare the

reconstructed molecular dates to the true dates from the simulated data, we calculated three

metrics for each tree and summarised these over each simulation type (Table 2). The metrics

were median absolute percentage error of node ages (MAPE), topological error (Topo. Error)

by number of true bipartitions not reconstructed, and uncertainty by mean highest posterior

density interval width as a percentage of reconstructed node age (HPD width %). We

compared the gamma statistics of the reconstructed trees to the simulated trees in order to

detect any systematic bias in node age error. As a way of evaluating the degree to which the

modelled relationship between speciation and substitution rates is reflected in the phylogenies

reconstructed from the simulated data, we also performed sister pair analysis to test if we can

detect the positive correlation between speciation rate and substitution rate from the

reconstructed trees.

Table 2. Summary of results.

Recon. method	Simulation	MAPE (%)	Topo. error	HPD width (%)	Gamma Statistic	Sister pair analysis
BEAST 2 (UCLN)	*Unlinked*	6.3–30.9	0–14	43–29	-1.46–1.09	None
	Continuous	6.7–30.3	0–16	35–605	-1.52–0.20	None
	Punctuated	8.8–80.4	0-8	42–154	-2.16–3.92	Negative
PAML (ACLN)	*Unlinked*	11.4–192.1	Fixed	62–178	-5.32–9.77	Positive
	Continuous	9.4–188.0	Fixed	57–151	-5.15–6.25	Positive
	Punctuated	15.4–438.9	Fixed	73 - 327	-8.4–2.99	Positive
PAML (UCLN)	*Unlinked*	22.7–245.2	Fixed	49–263	-6.53–7.51	Positive
	Continuous	26.2–227.5	Fixed	49–379	-6.246.65	Positive
	Punctuated	18.1–535.4	Fixed	102–301	-8.43–4.83	Positive

Summary of reconstructed tree characteristics. Trees simulated under three models:
Unlinked, Continuous, and Punctuated. Molecular dates were then reconstructed using three

different methods: BEAST 2 (uncorrelated lognormal model, UCLN), PAML (autocorrelated
lognormal model, ACLN), and PAML (UCLN). For each reconstructed tree we calculated
median absolute % error of node ages (MAPE %), topological error (Topo. Error) by number
of true bipartitions not reconstructed, and uncertainty by mean highest posterior density
interval width as a percentage of reconstructed node age (HPD width %). We compared the
gamma statistics of the reconstructed trees to the simulated trees and used sister pair analysis
to test if we can detect the positive correlation between speciation rate in the reconstructed
trees (None = no significant correlation).
274

First, we examined the median absolute percentage error (MAPE) in the ages of all

nodes within each tree whose topology was correctly reconstructed (Figure 2). For trees

reconstructed using BEAST 2 (UCLN), the average (min–max) MAPE was 12.2% for

Unlinked simulations (6.3–30.9%). Errors for the Continuous simulation were similar at

14.2% (6.7–30.3%). Average MAPE for the Punctuated simulation was higher than for

Unlinked trees at 20.3% (8.8–80.4%). For trees reconstructed using PAML (ACLN), MAPE

values were higher and range of errors were larger than for BEAST 2 (UCLN). Average

MAPE for Unlinked trees was 66.0% (11–192%), compared to 60.6% (9.4%–188%) for

Continuous trees and 91.1% (15.4–438.9%) for Punctuated trees. For reconstructions with

PAML (UCLN), average MAPE was even higher than PAML (ACLN) at 77.0% (22.7–

245.2) for Unlinked trees, for Continuous trees 69.6% (26.2–227.5), but lower for Punctuated

trees at 74.9% (18.1-535.3). Because these errors can only be calculated for nodes that are

shared between the true and reconstructed trees, we also examined an alternative error metric

using the branch score (Kühner-Felsenstein) distance, which is based on the sum of squared

branch length errors, includes all nodes, and treats branches that are incorrectly reconstructed

as length zero, thus accounting for topology errors (86). This metric led to similar patterns

and is included as online Supplementary Information (Figure S1).

Figure 1. Median absolute percentage error of node times (on log scale) reconstructed from

data sets simulated under three simulation types with different relationships between

molecular evolution and speciation. The mean error is calculated only for branches whose

bipartitions are shared between the true and inferred trees, and therefore ignores errors in

topology which may occur in the BEAST 2 analyses. The three simulation models are

Unlinked (instantaneous covariance of molecular rates and speciation rates = 0), Continuous

(instantaneous covariance = 0.0044), and Punctuated (instantaneous covariance = 0, bursts of

substitutions added at speciation events). Topologies and node times were reconstructed

using three different analytical methods, with an uncorrelated lognormal 'relaxed clock' rate

prior (UCLN) in BEAST 2, the autocorrelated lognormal rate prior (ACLN) in PAML, and

the UCLN in PAML.

Second, for trees reconstructed using BEAST 2 (UCLN), we also investigated the

degree of topological error via the Robinson-Foulds distance between the true and estimated

308   tree (87). This measure can only be used for BEAST 2 because the 'mcmctree' program in

309   PAML does not infer topology. The median background topological error for the Unlinked

310   simulations was 6 out of 74 bipartitions (0–14), compared to 5 (0–16) for the Continuous

311   model (Figure 3). Error was lower for the Punctuated model with a median distance of 2

312   bipartitions (0-8).

314
315
Figure 2. Topological error of trees reconstructed from data sets simulated under three

simulation types with different relationships between molecular evolution and speciation (see

Table 1). Topological error is given by the Robinson-Foulds distance (number of non-shared

bipartitions). The three simulation models are Unlinked (instantaneous covariance of

molecular rates and speciation rates = 0), Continuous (instantaneous covariance = 0.0044),

and Punctuated (instantaneous covariance = 0, bursts of substitutions added at speciation

events). Topologies and node times were reconstructed using the uncorrelated lognormal

'relaxed clock' rate prior in BEAST 2 (UCLN). No results are available for PAML because

the topology was fixed for these analyses.

Third, we calculated the uncertainty of node age reconstruction as reported by the

method via the width of 95% highest posterior density intervals as a proportion of node age.

Division by node age is required because the infinite-sites model of molecular evolution

predicts that the relationship between posterior mean node age estimates and 95% credible

intervals should approach a straight line with sufficient molecular data (75, 77). Using

BEAST 2 (UCLN) (Figure 4), the average HPD width for the Unlinked trees had a width

60% of the reconstructed age of its associated node (43 – 29%). For the Continuous trees this

was 108% (35 – 605%), and for the Punctuated trees it was 70% of node age (42 – 154%).

Using PAML (ACLN), the average HPD width for Unlinked trees was 107.0% of node age

(62.2 – 178.4%); for Continuous trees, 100.4% (57.2 – 151.5%); and for Punctuated trees,

112.4% (73.4 – 327.0%). Using PAML (UCLN), the average HPD width was greater than

for PAML (ACLN) for all three simulation models, at 160% (49.5–263.4%) of node age for

Unlinked trees; 158% (48.7–379.5%) for Continuous trees; and 165% (102.4–301.2%) for

Punctuated trees.

Figure 3. Uncertainty of trees reconstructed from phylogenies simulated under three models

with different relationships between molecular evolution and speciation, in width as a

percentage of reconstructed node age. Precision is measured by averaging the width of 95%

highest posterior density intervals as a proportion of reconstructed node height across each

tree, with a larger score indicating less precise estimates (HPD width). The three simulation

models are Unlinked (instantaneous covariance of molecular rates and speciation rates = 0),

Continuous (instantaneous covariance = 0.0044), and Punctuated (instantaneous covariance =

0, bursts of substitutions added at speciation events). Topologies and node times were

reconstructed using three different analytical methods, the uncorrelated lognormal 'relaxed

clock' (UCLN) rate prior in BEAST 2, the autocorrelated lognormal rate (ACLN) prior in

PAML, and the UCLN in PAML.

2.4 Distribution of reconstructed vs simulated node ages

If the difference in gamma values between reconstructed and true trees is negative,

the reconstructed tree has nodes distributed more towards the root, while if it is positive the

reconstructed tree has nodes distributed more towards the tips than the true tree. Using

BEAST 2 (UCLN), gamma values for the reconstructed Unlinked trees were distributed

evenly around the gamma value of the simulated tree with a mean shift of +0.11 (-1.46 – +

1.09; Figure 5). This distribution was skewed negative (towards the root) relative to the

simulated tree for Continuous trees with a mean of -0.44 (-1.52 – +0.20). This negative skew

was more pronounced for the Punctuated trees at -0.94 (-2.16 – +3.92). Using

PAML(ACLN), the relative change in gamma was on average negative across all three

simulation models. For the Unlinked trees it was -0.28 (-5.32 – +9.77), while for the

Continuous trees it was -0.76 (-5.15 – +6.25), and the Punctuated trees were again skewed

negative with a mean of -2.87 (-8.40 – +2.99). In contrast when using PAML (UCLN), the

relative change in gamma was on average positive for the Unlinked trees at +0.57 (-6.53 –

+7.51) and for the Continuous trees at +1.07 (-6.24 – +6.65). It was slightly negative for the

Punctuated trees at -0.16 (-8.43 – +4.83).

Figure 4. Difference in gamma statistic values between reconstructed and simulated trees,

where simulated trees were generated under three models with different relationships

between rates of molecular evolution and speciation rates. Negative scores indicate that

reconstructed trees have nodes distributed more towards the root ('tippier') than the

simulated tree. The three simulation models are Unlinked (instantaneous covariance of

molecular rates and speciation rates = 0), Continuous (instantaneous covariance = 0.0044),

and Punctuated (instantaneous covariance = 0, bursts of substitutions added at speciation

events). Topologies and node times were reconstructed using three different methods, an

uncorrelated lognormal (UCLN) rate prior in BEAST 2, an autocorrelated lognormal rate

(ACLN) prior in PAML, and the UCLN in PAML.

2.5 Detection of rate differences on reconstructed phylogenies

Sister pairs analysis have been used to detect an association between diversification

rates and rates of molecular evolution in a wide range of datasets (6-8, 48, 88). In order to

determine whether the simulated relationship between molecular rates and diversification

rates could be detected in reconstructed trees, we regressed log clade size contrasts against

contrasts in log median branch length estimates in the data set of sister pair analysis

generated from reconstructed trees. We present the p-value and sign of slope for each

regression (Figure 6). First, we asked whether the sister pairs analysis suggest a relationship

between rates of molecular evolution and speciation rate in the simulated data (that is, in the

true trees). As expected, Unlinked simulated trees show no significant relationship between

clade size and branch length. Significant positive relationships are found for the Continuous

and Punctuated simulations; note that this was a condition for selection of the Continuous

data set (Supplementary Methods). Second, we asked whether the relationship between rates

of molecular evolution and diversification rate could be detected in the reconstructed

phylogenies. When reconstructed using BEAST 2 (UCLN), the Unlinked data set and the

Continuous data set show no significant relationship, while the Punctuated data set gives a

significant negative relationship, the opposite to what we expect. When reconstructed under

PAML(ACLN) and PAML (UCLN), all three of the Unlinked, Continuous and Punctuated

data sets have significant positive relationships between clade sizes and median branch length

estimates.

Figure 5. Slope and significance of mock sister pair analyses of clade size contrasts against

contrasts in simulated branch lengths (SIM) or median branch length estimates reconstructed

from simulated trees. Results are shown for branch lengths reconstructed using an

uncorrelated lognormal (UCLN) rate prior in BEAST 2, an autocorrelated lognormal (ACLN)

rate prior in PAML, or the UCLN in PAML. The three simulation models are Unlinked

(instantaneous covariance of molecular rates and speciation rates = 0), Continuous

(instantaneous covariance = 0.0044), and Punctuated (instantaneous covariance = 0, bursts of

substitutions added at speciation events). Black triangles represent positive regression

coefficients, while white inverted triangles represent negative coefficients. The red line

indicates significance at the 0.05 level.

3.Discussion

3.1 Dating errors associated with variation in speciation and substitution rate

The addition of rate-variable 'relaxed clock' models represents a significant extension to the sophistication of modern molecular dating methods (15). While there is a wide variety of rate-variable molecular dating methods, all of them to date rely on stochastic models of rate variation, such that rates of molecular evolution are expected to evolve randomly over the phylogeny. However, it is clear that real substitution rates do not always vary randomly. Instead, rates of molecular evolution can be influenced by species life history, niche and environment and macroevolutionary processes. It is important to examine whether the stochastic models of rate variation can adequately model realistic biological patterns of rate variation (3, 62, 65, 89). In the case of substitution rates, this means that we must test molecular dating methods using rate simulations that are empirically based and incorporate the effects of other biological processes. Here we have focused on testing the impact of widely reported correlations between molecular rates and diversification on the accuracy of molecular date reconstruction (5-8).

We used three models, Unlinked, Continuous and Punctuated, to reflect different possible links between substitution rates and diversification rates – either varying independently; covarying continuously, linked directly by common mechanisms or indirect through shared influences on rate variation; or linked by punctuated bursts of increased substitutions related to speciation events. Rather than choosing parameters arbitrarily, we parameterised our models using realistic values from the literature, or found plausible values

by tuning the models to produce data sets that reproduce empirical findings (Supplementary

Methods; 7, 79, 80). To make sure that we had realistic parameters for simulations, we based

our parameter values on empirically derived values for birds. This approach produces more

realistic simulations than the common approach choosing arbitrary values for convenience.

Although these values may not represent all taxa, they are still more likely to be a realistic

representation than random values, and here are used only to explore whether molecular

dating methods are accurate given simulated datasets that mimic, as much as possible, a real

case study. Exploring the performance of methods on simulations modelled on other case

studies would be desirable, if appropriate parameters were available. Modelling population

genetic processes and multiple loci could also add real-world error sources such as gene tree

discordance, coalescent error and incomplete lineage sorting, many of which would be

affected by variation in substitution rates and branch times. Our framework focuses on the

most common use cases for macroevolutionary studies, which generally involve analyses of

few loci at broad taxonomic scales for which these problems may be less prevalent. However,

this would be a useful path for future research.

Similarly, we use three examples of 'relaxed clock' molecular dating methods:

BEAST 2 (using an uncorrelated rates model, UCLN) and PAML (using either the

autocorrelated rates model, ACLN, or the UCLN). Clearly, these methods and models do not

represent all possible approaches to inferring molecular dates using a rate-variable method,

but they were chosen because they are widely used and differ in some key respects, and

therefore present two useful test cases for the accuracy of "relaxed clock" methods in the face

of systematic rate variation.

3.2 Unlinked model

The Unlinked model describes the error that can be expected from independent

variation in speciation and molecular rates. Inference on trees simulated under this model

produced median absolute errors in node age inference of about 12% on average using

BEAST 2 (UCLN), and high values of 66% using the PAML (ACLN) and 77% using PAML

(UCLN) (Figure 2). The largest errors were up to 30% for BEAST 2 (UCLN),while the worst

instances for PAML (ACLN) had errors up to 192% and the worst MAPE for any individual

tree was in PAML (UCLN) at over 500%. The Unlinked model produced lower errors using

BEAST 2 (UCLN) than the other two models(Figure 2). Gamma differences for the Unlinked

model are distributed evenly around zero using BEAST 2 (UCLN), so there is no overall

tendency for nodes to be over- or under-estimated (Figure 4). The sources of error in the

Unlinked model are most likely to be violation of the assumptions of rate priors and node

time (tree) priors. Variation in speciation rates violates the assumptions of the birth-death

node time priors used in both BEAST 2 and PAML, which assume universally constant rates

of speciation and extinction (90). Speciation rate variation among lineages also induces more

imbalanced trees on average (91), which can lead to bias and loss of precision in node age

inferences (92). While the degree and scale on which speciation and extinction rates vary

remains contested (93-96), there is increasing evidence that radiations and other forms of

variation are a factor in the evolution of many taxa, so that errors of this magnitude may be

widespread (97-100). Substitution rates that evolve in an autocorrelated fashion also violate

the assumptions of the UCLN, which assumes that the evolutionary rate of each lineage is

independent (20), and this may also induce errors in inference (30, 34).

The causes of the extremely high errors of the PAML(ACLN) and PAML (UCLN)

phylogenetic inference for the Unlinked trees (and throughout the study for the other models)

are unclear. Examination of the output trees suggests that the most extreme errors are in trees

with shallower calibrations, and that the ages of large uncalibrated clades are sometimes

overestimated severely. The inferences from PAML and BEAST 2 are not directly

comparable due to numerous technical differences: for example, tree topology is fixed in

PAML but estimated in BEAST 2; and BEAST 2 has prior distributions on speciation and

extinction parameters, which are fixed in PAML. Nevertheless it is surprising that the

performance of the ACLN model in PAML was worse than the UCLN model in BEAST 2

even though the underlying rate variation in these simulations is indeed autocorrelated. The

fact that even higher errors and greater HPD widths are found when using PAML (UCLN)

than using PAML (ACLN) (Figures 1, 3) suggests that the choice of rate prior can have some

ameliorating influence, but the differences between the methods are not wholly due to the

rates prior, because PAML (UCLN) still performs significantly worse than BEAST2 UCLN,

indicating that there must be additional differences in methodology impacting the accuracy of

inference. One possibility is that the sensitivity of the prior to model violation in PAML

(ACLN) was enhanced by our use of the approximate likelihood function (101). This is a

second-order Taylor approximation to the likelihood evaluated at the maximum likelihood

estimates of the branch lengths and other parameters. As discussed by dos Reis and Yang

(101), while the approximation is very accurate under normal circumstances, when the

branch lengths proposed during Bayesian inference are very far from the maximum

likelihood values the approximation becomes much less accurate. This can happen when the

rate or node time (tree) priors are misspecified. PAML also requires some parameters of the

prior to be specified exactly, such as the speciation and extinction rates, whereas in BEAST 2

these values are usually given distributions. This could mean that PAML is not as robust to

violation of the assumption of constant diversification rates. Nevertheless, absolute error rates

in other simulation studies have been less than 20% even with a deliberately misspecified rate

prior and more limited calibrating information (30), and the high error rates may be

indicating that patterns of rate variation likely to exist in real data may reveal weaknesses in

methods that were not obvious from simulating simpler, stochastic patterns of rate variation.

3.3 Continuous model

The Continuous model has substitution rates and speciation rates that coevolve

continuously rather than varying independently. Notably, we chose the degree of covariance

that reproduced two well-established empirical phenomena – the correlation of clade size and

branch length contrasts detected in a sister pairs analysis by Lanfear et al. (7), and the

widespread relationship between branch lengths and tree nodes reported by Pagel et al. (62).

On average, this model produced absolute node time inference errors of 14% using BEAST 2

(UCLN), slightly higher than the Unlinked model, but the worst errors were up to 30%

(Figure 2). Negatively-skewed gamma differences indicate that error generally leads to node

ages being overestimated, pushing reconstructed nodes back towards the root (Figure 4). In

addition to the error, which is the difference between reconstructed time estimates and the

truth, we also examine how the reconstruction methods report their uncertainty about the

truth through 95% highest posterior density intervals. Although not greatly different in terms

of absolute error, the reported uncertainty of node time estimates was very high for this

model using BEAST 2 (UCLN), with average HPD widths more than 107% of node height

(Figure 5). This average is driven partly by some outlying trees with very high uncertainty,

up to 600% of node height; the median is 75% of node height. Wide HPD intervals can be a

sign of conflict between the prior and data, so it may be that covariance between

diversification rates and substitution rates tends to produce patterns of branch lengths that are

not well described by independent rate and time priors. The practical impact of this level of

uncertainty could depend on how it is distributed within the tree and the nature of any

downstream inferences; if the question at hand is the dating of major clades then high

uncertainty in the age of tips may be tolerable, while the same level of uncertainty in internal

branches would render the tree unusable. However, any macroevolutionary inference that

relies on the distribution of nodes throughout the tree would likely be affected (59). Error

rates for the Continuous simulations were similar to those for the Unlinked simulations using

PAML (ACLN) (mean of 60.6%; Figure 2) and PAML (UCLN) (mean of 69.6%). Similar

error rates suggest that the accuracy of PAML analysis, while having low precision under all

simulation models, was not substantially affected by the degree of correlation between

substitution rates and diversification rates.

3.4 Punctuated model

Levels of absolute error were highest under BEAST 2 (UCLN) when speciation

events were associated with large punctuational bursts of molecular evolution (mean of

20.3%, up to 80.4%; Figure 2). This implies that, if real evolutionary processes were to

commonly involve such bursts of substitutions associated with speciation, or produce a

similar distribution of evolutionary histories and rate variation through some other

mechanism, median node age estimates reconstructed under the UCLN rate prior would be

generally incorrect. Error rates were also extreme for PAML (ACLN) (mean 91.4%, max

438.9%; Figure 2) and for PAML (UCLN) (mean 74.9%, max 535%). These rates were

slightly higher on average than the rates for Unlinked and Continuous models, though also

depending more on the specific tree. The finding of high error rates in the Punctuated model

corroborates a recent analysis (72), which showed that greater proportions of branch length

associated with speciation events lead to greater errors in the age of the root. Here, we show

that the effect is strong enough to impact the median absolute error of all node ages, even

when the crown node is calibrated. The very negative skew of Gamma away from the

simulated value shows that these internal node ages are severely overestimated, leading to

very distorted trees with long tips (Figure 4).

Despite having the highest degree of branch length error using BEAST 2 (UCLN), the

Punctuated model had the lowest degree of topological error (a median of 2 bipartitions not

correctly reconstructed; Figure 3), comparing to the Unlinked and Continuous models

(median of 5 and 6 respectively, Figure 3). Intuitively, this could be because rapidly

diversifying clades will normally have many short branches, making their internal

relationships more uncertain. The Punctuated model adds more substitutions to every branch

regardless of its duration in time, and so helps resolve these rapidly diversifying clades. The

precise impact of topological error is difficult to interpret, since our investigation does not

distinguish among topological errors at different timescales and we do not designate any

particular nodes as important. But the practical upshot of systematically lower topological

error in the simulation with the most severe node age errors using BEAST 2 (UCLN) is that

these errors will be difficult to diagnose. Major topological errors are easier to diagnose by

checking against classical taxonomies or other information, while little is usually known

about the dates of uncalibrated nodes ahead of time.

Given the errors in inference of phylogenetic branch lengths and topology under these

rate models, and the aforementioned difficulty in diagnosing these errors from inferred

phylogenies, would we expect to be able to detect an association between diversification rates

and substitution rates in these reconstructed phylogenies? We asked if the underlying

relationship could be detected using a standard analytical approach with sister pairs (Figure

6). Although the association between speciation rates and substitution rates was evident in the

simulated trees, tree reconstruction using PAML(ACLN), PAML (UCLN) or BEAST 2

(UCLN) disrupts the relationship between clade size and branch lengths and clade sizes

displayed by the underlying simulated trees (Figure 4). In the BEAST 2 (UCLN) phylogenies

the relationship between speciation and molecular rates is inverted, giving a negative

correlation between clade sizes and branch lengths. This could be related to the fact that

lineages with many speciation events have higher variance in branch lengths in the BEAST 2

(UCLN) reconstruction (102) which means their total length could be more strongly

constrained by the node time (tree) prior. Conversely, for the PAML (ACLN) and PAML

(UCLN) reconstructed trees, we found significant positive relationships between inferred

branch lengths and clade size for all three models, even the Unlinked model where speciation

rates and substitution rates are not positively associated. A possible cause of this inference of

a positive association between diversification rates and substitution rate, even in the absence

of an underlying association between the two, is the node-density effect, which results from

systematic underestimation of branch lengths with high uncertainty (103). The influence of

the node density effect should be reduced by our procedure of selecting an even number of

samples in each sister clade, but may not be eliminated by it due to the extremely high error

rates in the PAML (ACLN) or PAML (UCLN) reconstructions. The node-density effect may

not be observed in our BEAST 2 (UCLN) because the aforementioned artefactual negative

correlation overcomes any positive correlation due to node density. However the reasons why

this does not occur using PAML (UCLN) are unclear and cannot be distinguished using our

data. The failure to recover the empirical correlation from branch lengths reconstructed as

part of a Bayesian dating analysis suggests that this is not a useful way to test for the

presence of interactions between substitution rates and speciation rates. For some data sets it

may be practical to check for such correlations by reconstructing branch lengths in units of

genetic divergence using a fast branch length reconstruction method that does not impose a

model of timing or rate variation (e.g. 85, 104, 105, 106). However, even if such correlations

can be diagnosed, there is currently no available method for correcting any associated bias.

This should be an area of active research.

The true role of punctuational bursts or coevolution of speciation and substitution

rates in macroevolution is currently a matter of debate. The primary empirical evidence is the

analysis of a large selection of empirical phylogenies in Webster et al. (63) and Pagel et al.

(62). These studies have found that the length of root-to-tip paths in the phylogenies in

substitutions per site is frequently higher when there are more nodes on the path. The

relationship remains even after removing the node-density effect that can generate spurious

correlations between branch lengths and node densities across clades (103, 107). Regardless

of the mechanisms, we have shown that if there were punctuated busts of substitutions

associated with speciation events, we could expect it to impact the accuracy of molecular date

estimates, since any mechanism that leads to similar patterns of rate variation and apparent

correlation between branch lengths and speciation events would produce similar effects.

3.5 Molecular dating methodology

Our study contrasts two common reconstruction methods, rather than varying aspects

of data choice. Many aspects of data choice and analytical method could exacerbate or reduce

the errors found in this analysis. For example, our alignments are shorter than many used in

actual Bayesian phylogenetic analyses but also likely more informative due to being

simulated without invariant sites or other more complex patterns of substitution. Shorter or

less informative alignments could lead to greater or more widespread error. Perhaps the most

important element that we did not explore was the influence of calibrations. Calibration

choice and positioning, as well as the shape of the calibrating distribution used, is often the

most influential factor influencing divergence time reconstruction (e.g. 32, 108) and is the

source of many disagreements in published molecular dating studies, including high-profile

dating studies in birds (109-111) mammals (112, 113), and insects (114-116). Simulation

studies have suggested that calibrations can rescue analyses strongly affected by rate prior

misspecification (30), although others have found that sufficiently misspecified rate priors

cannot be rescued by more calibrations (34). The largest and most data-rich molecular dating

studies may have many more calibrations than we implement (e.g. 139, 117). However, our

trees are smaller than in these studies, and, unlike empirical studies, our calibrations are

guaranteed to be correctly placed, have hard constraints, and are centered on the true value.

We also ensure the root is calibrated, which has been shown to have the largest effect on

accuracy in several studies (30, 34, 118). We therefore believe that our method overall

represents a balance between realism and conservatism in determining the amount of

calibrating information available to inform divergence time estimates. However, a usefl

extension of the analyses undertaken in this study would be investigation of the effects of

both alignment length and calibration numbers and positioning.

4. Conclusions

Our results here show the value of testing molecular phylogenetic and dating methods

against complex simulation models informed by empirically determined patterns of rate

variation. We have demonstrated that commonly applied Bayesian divergence time

estimation methods can experience average errors of 12% in node dates under a model in

which speciation and molecular rates vary independently, 14% when substitution rates are

linked to speciation rates and 20% under a punctuational model when analysed using an

uncorrelated rates prior in BEAST (UCLN). Errors were much higher under all models when

trees were analysed with an autocorrelated rate prior in PAML (ACLN), and still higher with

the uncorrelated lognormal rate prior in PAML (UCLN). As demonstrated by the negative

skewness of gamma statistics, for trees with links between speciation and molecular

evolution these errors could lead to systematic overestimation of node ages. Regardless of

the mechanism generating the association between speciation rates and substitution rates, we

show that the potential for divergence time estimation error associated with known empirical

relationships between molecular evolution (manifest in substitution rate) and macroevolution

(manifest in the topology and branch lengths of phylogenetic trees) can be significant and

should be taken into account. Future work should see a broader range of sophisticated

divergence time estimation methods be tested against a wider variety of more empirically

realistic simulated data, including geographic and environmental biases, concerted changes in

population size, and directional changes in life history characteristics. Ultimately the goal

should be to produce new models that relax the assumption of independence between priors

on divergence times and priors on substitution rates. These will lead to greater confidence in

applications that depend on robust estimates from molecular dating procedures.

5.Methods

5.1 Simulation design

We refer throughout to simulated phylogenies, which are known without error, as the

'simulated trees', and to the phylogenies inferred from analyzing the sequences evolved

along each of these simulated phylogenies as the 'reconstructed trees'. Comparison of the

reconstructed trees to the simulated trees allows us to evaluate the degree to which

phylogenetic inference methods correctly infer the known evolutionary history of the

simulated data.

Our overall design is summarised in Figure 1. For our simulation study, we first

generate phylogenetic trees in units of absolute time with evolving rates of speciation and

molecular evolution. We then simulate DNA sequences along these phylogenies. Finally, we

infer dated trees from the alignment of these sequences using standard methods in Bayesian

phylogenetics and compare the inferred tree to the simulated tree from the known

evolutionary history from each simulation.

We simulate sets of 50 phylogenetic trees and nucleotide sequence alignments under

each of three different models describing the association between speciation rates and rates of

molecular evolution: one "Unlinked" model in which molecular and diversification rates each

vary independently, as well as two alternative mechanisms for an association between rates

of molecular evolution and speciation rates ("Punctuated" and "Continuous": see Table 1).

We do not consider models in which molecular rates affect rates of extinction, so in our

simulations, extinction rates remain constant over the tree. For all models, we first simulate

the phylogenetic tree in continuous time, together with speciation and substitution rates that

vary along the tree. From this, we derive branch lengths in expected genetic divergence

(substitutions/site). Finally, we simulate the evolution of nucleotide sequences along these

phylogenies.

Figure 6. Flowchart of study design. Trees with branch lengths in units of time are

simulated under three simulation conditions. For the Unlinked model, birth rates (λ)and

substitution rates (sub) vary independently, while for the Continuous model they are

correlated. For the Punctuated model, trees are simulated as for Unlinked, then some of the

branch length is redistributed to form bursts of substitutions occurring upon speciation. Trees

and branch lengths are then used to simulate molecular sequence alignments. Time trees are

then reconstructed using three different methods with two rate priors: theuncorrelated

lognormal in BEAST 2 (UCLN), the autocorrelated lognormal in PAML (ACLN), and the

uncorrelated lognormal in PAML (UCLN). Reconstructed trees are then compared to the

original, simulated trees to determine the error (Median absolute % error, MAPE),

uncertainty (HPD width) and bias towards over- or under-estimation (Gamma).

In the Unlinked model, we simulate continuous variation in both substitution rate and

speciation rate through time, but neither affects the other. Substitution and speciation rates

evolve independently by Brownian motion, where each lineage continuously adds a separate

random increment to its substitution rate and speciation rate in each of many small time

intervals. The Unlinked model serves as a baseline to determine the level of uncertainty in

molecular dates caused by the random variation in substitution and diversification rate when

there is no mechanistic link between the two. In the Continuous model, speciation rates and

substitution rates coevolve with each other continuously over time. The coevolution is

simulated as a correlated Brownian motion, where the random increments in speciation and

molecular rates added over time are dependent on one another. Consequently, substitution

rate and speciation rate tend to increase or decrease together. For this model, we also ensured

realism simulated many data sets with different covariance parameters, and selected a set of

50 trees that reproduced the empirical correlation of (7) between clade size and branch length

(Supplementary Methods S1.1). In the Punctuated model, the genetic divergence along a

branch is the sum of a gradual accumulation of substitutions at a rate that evolves as in the

Unlinked model, and a burst component at each speciation event. A speciation event occurs

each time a new lineage is born, whether any of its descendants are observed in the present or

not. The amount of additional branch length due to the burst is independently and randomly

drawn for each speciation event. For this model, we condition our simulations to reproduce

the relationship observed in (62). There it was observed that the total number of substitutions

occurring along root-to-tip paths in their sample of trees correlates with the number of nodes

along the path. Although this relationship could also be produced by a mechanism similar to

the Continuous model (119-121), we model the mechanism proposed in (62) that the node-

associated substitutions are produced in bursts associated with speciation events.

We derived plausible speciation, extinction and substitution rates for our simulations

using the avian phylogeny in (82). We do not intend to make any inferences about the actual

evolution of birds or the accuracy of inferred divergence times. Birds are chosen only

because the relationships among traits, molecular rates and diversification rates are well

studied, so that we can use these to give us an empirical ground for choosing parameter

values. Branch-specific diversification rates have been estimated for many of the lineages in

this tree in (80), while substitution rates have been inferred in (79) assuming covariation of

molecular rates and life history. Speciation rate and molecular rate values taken from avian

data have been previously used to inform realistic simulations on ecological assemblies for

the purpose of estimating the effect of dating error on phylogenetic diversity estimates (122),

and we make use of similar procedures here. A relationship between molecular rates and net

diversification rate has also been established for birds (7, 8). We use this fact to select our

Continuous data set by simulating multiple data sets and varying the covariance parameter

until the relationship appears. Full details of how we used these studies to determine realistic

parameterisations for our simulations are available as Supplementary Methods S1.1.

5.2 Simulation trees and alignments

To simulate the phylogeny, we adopt a forward approach, starting at the root of the

phylogeny and moving forward in time, as lineages divide into two daughter lineages, go

extinct, and evolve in speciation and substitution rate as a continuous time Markov process.

In each step, we draw the time until the next event from an exponential distribution with rate

equal to the sum of the rates of all events in all lineages. Then, for every lineage l that is

present at time *t*, one of the following three events may take place:

(i) The lineage gives rise to two daughter lineages with rate $\lambda(l, t)$, the current speciation

rate of lineage *l* at time *t*. Both daughter lineages initially retain the same speciation rate

and trait values as their parent lineages.

(ii) The lineage becomes extinct with rate μ, which is constant over the tree.

(iii) The lineage updates its speciation rate and substitution rate with rate *q*, which is

constant throughout the tree. The update process adds an increment to the current

speciation rate, which is a random draw from a bivariate normal distribution with mean 0

and an instantaneous variance-covariance matrix Σ multiplied by the time since last jump.

Until one of these events happens, the lineage continues with the same speciation and

molecular rates. High values of *q* make for longer simulations but produce a better

approximation to Brownian motion. We found that *q* had no influence on observed

speciation or substitution rate variance or average tree age for *q* > 10. For our final

simulations, we selected *q* = 50 to allow a significant margin of safety.

Specifically, the instantaneous variance-covariance matrix takes the form:

808
$$\Sigma = \begin{bmatrix} \sigma_\lambda^2 & \mathrm{Cov}(\lambda, r) \\ \mathrm{Cov}(\lambda, r) & \sigma_r^2 \end{bmatrix}$$

where σ_λ^2 describes how fast the speciation rate evolves and σ_r^2 describes how fast the

substitution rate evolves. $\mathrm{Cov}(\lambda, r) = \rho\sigma_\lambda\sigma_r$, where ρ is the correlation coefficient between

speciation rate and substitution rate, so $\rho = 0$ in the Unlinked Model and the Punctuated

model. We employ the Generalised Sampling Approach (123) to ensure a correct distribution

of trees with a given number of tips, which is set to 75, the approximate mean size of the

avian clades taken from (80). Details of the approach is in Supplementary Methods S1.2.

In the Punctuated model, we simulate an initial tree as per the Unlinked sample, but

without removing extinct taxa. We generate a burst of substitutions following each speciation

event. We generate Punctuated trees to match the formula found by (62), in which $16 \pm 5.4\%$

of the tree length (sum of all branch lengths) results from node-associated bursts

(Supplementary Methods S1.3).

Once we have simulated a tree, we simulate the evolution of nucleotide sequences

along the tree, given that branch lengths in the tree represent the expected number of

substitutions per site along the branch. The branch lengths in this tree represent the expected

number of substitutions per site along the branch. We simulate alignments of 2000 bases,

which is roughly equivalent to a 6000-base coding alignment, since most substitutions in a

coding alignment occur in the third codon sites. A 6000-base coding alignment is about the

size of the concatenated coding sequences of a mitochondrial genome. Previous literature

reviews have shown this to be a common length for phylogenetic data sets in published

ecological studies, and a tractable length for large Bayesian simulation studies (122). We

parameterize our sequence simulations using realistic among-site rate variation values from

birds (Supplementary Methods S1.4).

5.3 Calibrations

In addition to the trees and sequence alignments, we need calibrations to provide

information on absolute time to allow substitution rates to be identifiable from absolute ages

of nodes. For simulated trees of similar size to ours, diminishing accuracy returns for adding

new calibrations are achieved between 1 and 5 calibrations, especially if one is at the root

(30). Since our calibration dates are known with more accuracy than those in a real analysis,

and we remove the possibility of incorrectly assigned calibrations, we choose to assign two

calibrations per tree. For each tree, we choose calibrations at the root node and at one

randomly chosen internal node. To ensure that the calibration is randomly chosen across the

nodes of the simulated trees, we first decide the level of the calibrated node as a random

integer between 1 and the largest number of nodes along any root-to-tip path in the tree. Then

we randomly pick the calibrated node from all the nodes at that level.

For analyses in both BEAST 2 and PAML, we apply a hard-bounded uniform prior

for the age of each calibrated node. Both calibrations have bounds at the true age $\pm$ 15%.

Note that this is a conservative test of molecular dating because in the simulated datasets we

have accurate knowledge of the true date of the speciation event. In reality, calibrations may

be misplaced or incorrectly dated, and may only provide maximum or minimum bounds.

5.4 Phylogenetic inference

We generate a set of 50 trees under each of the Unlinked, Continuous and Punctuated

simulation types. After sequence simulation, this gives us 50 sequence alignments for each of

the three models.We infer divergence times from the simulated sequences using common

Bayesian molecular dating procedures. First, we generate posterior distributions of tree

topologies and divergence times using three common molecular dating methods: the

uncorrelated lognormal rate prior (UCLN), as implemented in the software package 'BEAST

2' (BEAST 2; 83), in which rates for each branch are independent draws from a common

lognormal distribution;the autocorrelated lognormal rate prior (ACLN), implemented in the

'mcmctree' program in the PAML software suite (85), in which the rate of each branch is

lognormally distributed with a mean equal to the rate of its parent branch; and the UCLN as

implemented in 'mcmctree' in PAML. To speed analysis when using PAML, we used the

approximate likelihood algorithm implemented in mcmctree. The PAML analyses also differ

from the BEAST 2 analyses in that some model parameters are provided as fixed numbers

rather than distributions, and that PAML requires a fixed tree topology. In this case the fixed

topology was the topology of the true simulated tree. Therefore only BEAST 2 (UCLN)

incorporates the possibility of topological inference errors. However, the three simulation

models can be compared within each reconstruction method. Details of the inference method

are available as Supplementary Methods (S1.5).

5.5 Error in reconstructed trees

For each reconstructed tree, we calculated accuracy in the estimation of node times

using the median absolute percentage error (MAPE) of inferred node times for clades that are

present in both true and reconstructed trees. The median was chosen rather than the mean to

reduce the impact of extremely high values for the estimates of shallow nodes. Since these

are only calculated for correctly reconstructed branches and do not capture errors in

reconstructing topology, we also calculate the Robinson-Foulds distance or number of non-

shared bipartitions, which only reflects topological error (87).

Model misspecification in Bayesian molecular dating methods can sometimes be

associated with higher uncertainty in the reconstructed node ages, as measured by the width

of Bayesian credible intervals. This is undesirable because the reconstructed history is less

useful as an explanatory tool and less informative in downstream analyses, but it can also be

a simple way to diagnose problems with the specified model. We examined the uncertainty

with which node ages were reconstructed by the two Bayesian molecular dating methods. We

scored uncertainty by averaging the width of the reconstructed 95% highest posterior density

intervals as a proportion of reconstructed node height across each tree (HPD width), with a

larger score indicating less precise estimates.

5.6 Distribution of reconstructed vs simulated node ages

We examine the direction of error by calculating the gamma statistic for each tree and

comparing to the true tree (124). The gamma statistic is a description of how the times

between branching events in a tree whose tips are all extant taxa change as they become

closer to the present. It summarises information relating to the pattern of diversification,

following a standard normal distribution for a tree with a constant rate of speciation and no

extinction. Trees with higher values have a relatively greater proportion of their length in

older branches, while those with lower values have relatively more of their length in younger

branches. The statistic therefore tells us whether the ages of internal nodes within each tree

tend to be older or younger on average than the true times while accounting for the different

scale of this bias for older and younger nodes.

5.7 Efficacy of sister pair correlation on reconstructed branch lengths

We checked to see if we can use the reconstructed branch lengths, which are the

median of posterior branch lengths calculated by BEAST 2 (UCLN), PAML (ACLN) or

PAML (UCLN), to detect the correlation between branch lengths and clade sizes found by

sister pair analysis on the simulated trees (see Supplemental Methods available as online

supplementary information). This could be used as a rapid post-hoc check to see whether

published dated trees could be affected by reconstruction error that is associated with

substitution and speciation rate covariation. Data sets for the sister pair analysis are generated

from the 50 simulated trees under each of the three simulation models, where each tree gives

a pair of sister clades and the sister clades are the two clades that split at the root. So, under

each simulation model, we have a data set of 50 sister pairs. We then regress contrasts in log

clade sizes against contrasts in the log of the median branch length estimates from BEAST 2

(UCLN), PAML (ACLN), and PAML (UCLN).

921
922 **Abbreviations**
923
924 UCLN – Uncorrelated Lognormal
925
926 ACLN – Autocorrelated Lognormal
927
928 HPD – Highest Posterior Density
929
930 MAPE – Mean Absolute Percentage Error
931
932
933 **Declarations**
934
Ethics approval

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

All data generated during this study are included in this published article as Supplementary

Information. The datasets generated are also available online at

https://github.com/amritchie/Sim_Evolving_Rates_Molecular_Dates.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the Australian Research Council grant DP190101916.

Authors' contributions

The study was conceived by L.B. A.M.R. wrote simulation code and conducted the analysis.

X.H. reviewed simulation code and assisted in the design of the study. A.R. and L.B. wrote

the manuscript. All authors reviewed the manuscript.

Acknowledgements

We would like to extend our gratitude to Marcel Cardillo and Alex Skeels for helpful

conversations in the preparation of this manuscript.

References

1.   Zuckerkandl E, Pauling L, Bryson V, Vogel H. Evolving genes and proteins. 1965.
2.   Dickerson RE. The structure of cytochrome c and the rates of molecular evolution. J
Mol Evol. 1971;1(1):26-45.
3.   Bromham L. Why do species vary in their rate of molecular evolution? Biol Lett.
2009;5:401-4.
4.   Bromham L. The genome as a life-history character: why rate of molecular evolution
varies between mammal species. Phil Trans Roy Soc B. 2011;366(1577):2503-13.
5.   Barraclough TG, Savolainen V. Evolutionary rates and species diversity in flowering
plants. Evolution. 2001;55:677-83.
6.   Bromham L, Hua X, Lanfear R, Cowman P. Exploring the relationships between
mutation rates, life history, genome size, environment and species richness in flowering
plants. Am Nat. 2015;185:507-24.
7.   Lanfear R, Ho SYW, Love D, Bromham L. Mutation rate is linked to diversification
in birds. Proc Natl Acad Sci U S A. 2010;107(47):20423-8.
8.   Eo SH, DeWoody JA. Evolutionary rates of mitochondrial genomes correspond to
diversification rates and to contemporary species richness in birds and reptiles. Proc R Soc B.
2010 277:3587-92.

9.   Gillman LN, Keeling DJ, Gardner RC, Wright SD. Faster evolution of highly conserved DNA in tropical plants. J Evol Biol. 2010;23(6):1327-30.

10.   Gillman LN, Wright SD. Species richness and evolutionary speed: the influence of temperature, water and area. J Biogeogr. 2014;41(1):39-51.

11.   Orton MG, May JA, Ly W, Lee DJ, Adamowicz SJ. Is molecular evolution faster in the tropics? Heredity (Edinb). 2019;122(5):513-24.

12.   Ho SYW. The changing face of the molecular evolutionary clock. Trends Ecol Evol. 2014;29(9):496-503.

13.   Kumar S, Hedges SB. Advances in time estimation methods for molecular data. Mol Biol Evol. 2016;33(4):863-9.

14.   Kumar S. Molecular clocks: four decades of evolution. Nat Rev Genet. 2005;6(8):654-62.

15.   dos Reis M, Donoghue PC, Yang Z. Bayesian molecular clock dating of species divergences in the genomics era. Nat Rev Genet. 2016;17(2):71-80.

16.   Ho SYW, Duchêne S. Molecular-clock methods for estimating evolutionary rates and timescales. Mol Ecol. 2014;23(24):5947-65.

17.   Heath TA, Holder MT, Huelsenbeck JP. A Dirichlet process prior for estimating lineage-specific substitution rates. Mol Biol Evol. 2012;29(3):939-55.

18.   Yoder AD, Yang Z. Estimation of primate speciation dates using local molecular clocks. Mol Biol Evol. 2000;17(7):1081-90.

19.   Drummond AJ, Suchard MA. Bayesian random local clocks, or one rate to rule them all. BMC Biol. 2010;8(1):114.

20.   Drummond AJ, Ho SYW, Phillips MJ, Rambaut A. Relaxed phylogenetics and dating with confidence. PLOS Biol. 2006;4(5):e88.

21.   Sanderson MJ. r8s: inferring absolute rates of molecular evolution and divergence times in the absence of a molecular clock. Bioinformatics. 2003;19(2):301-2.

22.   Smith SA, O'Meara BC. treePL: divergence time estimation using penalized likelihood for large phylogenies. Bioinformatics. 2012;28(20):2689-90.

23.   Sauquet H. A practical guide to molecular dating. Comptes Rendus Palevol. 2013;12(6):355-67.

24.   Wilke T, Schultheiß R, Albrecht C. As time goes by: a simple fool's guide to molecular clock approaches in invertebrates. Am Malac B. 2009;27(1/2):25-45.

25.   Thorne JL, Kishino H, Painter IS. Estimating the rate of evolution of the rate of molecular evolution. Mol Biol Evol. 1998;15(12):1647-57.

26.   Kishino H, Thorne JL, Bruno WJ. Performance of a divergence time estimation method under a probabilistic model of rate evolution. Mol Biol Evol. 2001;18(3):352-61.

27.   Aris-Brosou S, Yang Z. Effects of models of rate evolution on estimation of divergence dates with special reference to the metazoan 18S ribosomal RNA phylogeny. Syst Biol. 2002;51(5):703-14.

28.   Lepage T, Lawi S, Tupper P, Bryant D. Continuous and tractable models for the variation of evolutionary rates. Math Biosci. 2006;199(2):216-33.

29.   Sarver BAJ, Pennell MW, Brown JW, Keeble S, Hardwick KM, Sullivan J, et al. The choice of tree prior and molecular clock does not substantially affect phylogenetic inferences of diversification rates. PeerJ. 2019;7.

30.   Duchêne S, Lanfear R, Ho SYW. The impact of calibration and clock-model choice on molecular estimates of divergence times. Mol Phylogenet Evol. 2014;78:277-89.

31.   Lepage T, Bryant D, Philippe H, Lartillot N. A general comparison of relaxed molecular clock models. Mol Biol Evol. 2007;24(12):2669-80.

32.   Foster CSP, Sauquet H, van der Merwe M, McPherson H, Rossetto M, Ho SYW.
Evaluating the impact of genomic data and priors on Bayesian estimates of the angiosperm
evolutionary timescale. Syst Biol. 2017;66(3):338-51.
33.   Smith AB, Pisani D, Mackenzie-Dodds JA, Stockley B, Webster BL, Littlewood DTJ.
Testing the molecular clock: molecular and paleontological estimates of divergence times in
the Echinoidea (Echinodermata). Mol Biol Evol. 2006;23(10):1832-51.
34.   Battistuzzi FU, Filipski A, Hedges SB, Kumar S. Performance of relaxed-clock
methods in estimating evolutionary divergence times and their credibility intervals. Mol Biol
Evol. 2010;27(6):1289-300.
35.   Worobey M, Han G-Z, Rambaut A. A synchronized global sweep of the internal
genes of modern avian influenza virus. Nature. 2014;508(7495):254-7.
36.   Crisp MD, Hardy NB, Cook LG. Clock model makes a large difference to age
estimates of long-stemmed clades with no internal calibration: a test using Australian
grasstrees. BMC Evol Biol. 2014;14:263-.
37.   dos Reis M, Thawornwattana Y, Angelis K, Telford Maximilian J, Donoghue
Philip CJ, Yang Z. Uncertainty in the timing of origin of animals and the limits of precision
in molecular timescales. Curr Biol. 2015;25(22):2939-50.
38.   Welch JJ, Bromham L. Molecular dating when rates vary. Trends in Ecology &
Evolution. 2005;20(6):320-7.
39.   Lin G, Huang Z, Wang L, Chen Z, Zhang T, Gillman LN, et al. Evolutionary rates of
bumblebee genomes are faster at lower elevations. Mol Biol Evol. 2019;36(6):1215-9.
40.   Lourenço JM, Glémin S, Chiari Y, Galtier N. The determinants of the molecular
substitution process in turtles. J Evol Biol. 2013;26(1):38-50.
41.   Goldie X, Gillman L, Crisp M, Wright S. Evolutionary speed limited by water in arid
Australia. Proc R Soc B. 2010;277(1694):2645-53.
42.   Berv JS, Field DJ. Genomic signature of an avian Lilliput effect across the K-Pg
extinction. Syst Biol. 2018;67(1):1-13.
43.   Barrera-Redondo J, Ramirez-Barahona S, Eguiarte LE. Rates of molecular evolution
in tree ferns are associated with body size, environmental temperature, and biological
productivity. Evolution. 2018;72(5):1050-62.
44.   Qiu F, Kitchen A, Burleigh JG, Miyamoto MM. Scombroid fishes provide novel
insights into the trait/rate associations of molecular evolution. J Mol Evol. 2014;78(6):338-
48.
45.   May JA, Feng Z, Orton MG, Adamowicz SJ. The effects of ecological traits on the
rate of molecular evolution in ray-finned fishes: a multivariable approach. J Mol Evol.
2020;88(8-9):689-702.
46.   Welch JJ, Bininda-Emonds ORP, Bromham L. Correlates of substitution rate
variation in mammalian protein-coding sequences. BMC Evol Biol. 2008;8(1):53.
47.   Thomas JA, Welch JJ, Lanfear R, Bromham L. A generation time effect on the rate of
molecular evolution in invertebrates. Mol Biol Evol. 2010;27(5):1173-80.
48.   Hua X, Cowman P, Warren D, Bromham L. Longevity is linked to mitochondrial
mutation rates in rockfish: a test using Poisson regression. Mol Biol Evol. 2015;32(10):2633-
45.
49.   Galtier N, Blier PU, Nabholz B. Inverse relationship between longevity and
evolutionary rate of mitochondrial proteins in mammals and birds. Mitochondrion.
2009;9(1):51-7.
50.   Bromham L, Cowman PF, Lanfear R. Parasitic plants have increased rates of
molecular evolution across all three genomes. BMC Evol Biol. 2013;13(1):126.
51.   Mitterboeck TF, Adamowicz SJ. Flight loss linked to faster molecular evolution in
insects. Proc R Soc B. 2013;280(1767):20131128.

52.   Wong A. Covariance between testes size and substitution rates in primates. Mol Biol Evol. 2014;31(6):1432-6.

53.   Woolfit M, Bromham L. Population size and molecular evolution on islands. Proc Biol Sci. 2005;272(1578):2277-82.

54.   Lynch M, Ackerman MS, Gout J-F, Long H, Sung W, Thomas WK, et al. Genetic drift, selection and the evolution of the mutation rate. Nat Rev Genet. 2016;17(11):704-14.

55.   Gossmann TI, Keightley PD, Eyre-Walker A. The effect of variation in the effective population size on the rate of adaptive molecular evolution in eukaryotes. Genome Biol Evol. 2012;4(5):658-67.

56.   Charlesworth B. Fundamental concepts in genetics: effective population size and patterns of molecular evolution and variation. Nat Rev Genet. 2009;10(3):195-205.

57.   Mindell DP, Sites Jr JW, Graur D. Speciational evolution: a phylogenetic test with allozymes in Sceloporus (Reptilia). Cladistics. 1989;5(1):49-61.

58.   Iglesias-Carrasco M, Jennions MD, Ho SYW, Duchene DA. Sexual selection, body mass and molecular evolution interact to predict diversification in birds. Proc R Soc B. 2019;286(1899).

59.   Duchêne DA, Hua X, Bromham L. Phylogenetic estimates of diversification rate are affected by molecular rate variation. J Evol Biol. 2017;30(10):1884–97.

60.   Lancaster LT. Molecular evolutionary rates predict both extinction and speciation in temperate angiosperm lineages. BMC Evol Biol. 2010;10(1):162.

61.   Fontanillas E, Welch JJ, Thomas JA, Bromham L. The influence of body size and net diversification rate on molecular evolution during the radiation of animal phyla. BMC Ecol Evol. 2007;7:95.

62.   Pagel M, Venditti C, Meade A. Large punctuational contribution of speciation to evolutionary divergence at the molecular level. Science. 2006;314(5796):119.

63.   Webster AJ, Payne RJH, Pagel M. Molecular phylogenies link rates of evolution and speciation. Science. 2003;301:478.

64.   Ezard THG, Thomas GH, Purvis A. Inclusion of a near-complete fossil record reveals speciation-related molecular evolution. Methods Ecol Evol. 2013;4(8):745-53.

65.   Hua X, Bromham L. Darwinism for the genomic age: connecting mutation to diversification. Front Genet. 2017;8:12.

66.   Dobzhansky T. Genetics and the origin of species. New York: Columbia University Press; 1937.

67.   Muller H, editor Isolating mechanisms, evolution, and temperature. Biol Symp; 1942.

68.   Gavrilets S. Fitness landscapes and the origin of species (MPB-41): Princeton University Press; 2004.

69.   Orr HA. The population genetics of speciation: the evolution of hybrid incompatibilities. Genetics. 1995;139(4):1805-13.

70.   Frankham R. Genetics and extinction. Biol Cons. 2005;126(2):131-40.

71.   Venditti C, Pagel M. Speciation as an active force in promoting genetic evolution. Trends Ecol Evol. 2010;25(1):14-20.

72.   Janzen T, Bokma F, Etienne RS. Nucleotide substitutions during speciation may explain substitution rate variation. bioRxiv. 2021:2020.08.19.256891.

73.   Bromham L, Duchêne S, Hua X, Ritchie AM, Duchêne DA, Ho SYW. Bayesian molecular dating: opening up the black box. Biol Rev. 2018;93(2):1165-91.

74.   Warnock RCM, Parham JF, Joyce WG, Lyson TR, Donoghue PCJ. Calibration uncertainty in molecular dating analyses: there is no substitute for the prior evaluation of time priors. Proc Biol Sci. 2015;282(1798):20141013-.

75.   Yang Z, Rannala B. Bayesian estimation of species divergence times under a molecular clock using multiple fossil calibrations with soft bounds. Mol Biol Evol. 2005;23(1):212-26.

76.   Zhu T, Dos Reis M, Yang Z. Characterization of the uncertainty of divergence time estimation under relaxed molecular clock models using multiple loci. Syst Biol. 2015;64(2):267-80.

77.   dos Reis M, Yang Z. The unbearable uncertainty of Bayesian divergence time estimation. J Systemat Evol. 2013;51(1):30-43.

78.   Manceau M, Marin J, Morlon H, Lambert A. Model-based inference of punctuated molecular evolution. Mol Biol Evol. 2020;37(11):3308-23.

79.   Nabholz B, Lanfear R, Fuchs J. Body mass-corrected molecular rate for bird mitochondrial DNA. Mol Ecol. 2016;25(18):4438-49.

80.   Maliet O, Hartig F, Morlon H. A model with many small shifts for estimating species-specific diversification rates. Nat Ecol Evol. 2019;3(7):1086-92.

81.   Lanfear R, Ho SYW, Love D, Bromham L. Mutation rate influences diversification rate in birds. Proc Natl Acad Sci U S A. 2010;107(47):20423-8.

82.   Jetz W, Thomas GH, Joy JB, Hartmann K, Mooers AO. The global diversity of birds in space and time. Nature. 2012;491:444.

83.   Bouckaert R, Vaughan TG, Barido-Sottani J, Duchêne S, Fourment M, Gavryushkina A, et al. BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis. PLOS Comp Biol. 2019;15(4):e1006650.

84.   Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu C-H, Xie D, et al. BEAST 2: a software platform for Bayesian evolutionary analysis. PLOS Comp Biol. 2014;10(4):e1003537.

85.   Yang ZH. PAML 4: Phylogenetic analysis by maximum likelihood. Mol Biol Evol. 2007;24(8):1586-91.

86.   Kuhner MK, Felsenstein J. A simulation comparison of phylogeny algorithms under equal and unequal evolutionary rates. Mol Biol Evol. 1994;11(3):459-68.

87.   Robinson DF, Foulds LR. Comparison of phylogenetic trees. Math Biosci. 1981;53(1):131-47.

88.   Goldie X, Lanfear R, Bromham L. Diversification and the rate of molecular evolution: no evidence of a link in mammals. BMC Evol Biol. 2011;11(1):286.

89.   Bromham L. The genome as a life-history character: why rate of molecular evolution varies between mammal species. Phil Trans Roy Soc B. 2011;366(1577):2503-13.

90.   Gernhard T. The conditioned reconstructed process. J Theor Biol. 2008;253(4):769-78.

91.   Stadler T. Recovering speciation and extinction dynamics based on phylogenies. J Evol Biol. 2013;26(6):1203-19.

92.   Duchêne D, Duchêne S, Ho SYW. Tree imbalance causes a bias in phylogenetic estimation of evolutionary timescales using heterochronous sequences. Mol Ecol Res. 2015;15(4):785-94.

93.   Hedges SBK, Sudhir The Timetree of Life. New York: Oxford University Press; 2009.

94.   Hedges SB, Marin J, Suleski M, Paymer M, Kumar S. Tree of life reveals clock-like speciation and diversification. Mol Biol Evol. 2015;32(4):835-45.

95.   McPeek MA, Brown JM. Clade age and not diversification rate explains species richness among animal taxa. Am Nat. 2007;169(4):E97-106.

96.   Marin J, Rapacciuolo G, Costa GC, Graham CH, Brooks TM, Young BE, et al. Evolutionary time drives global tetrapod diversity. Proc R Soc B. 2018;285(1872):20172378.

97. Cardillo M. Latitude and rates of diversification in birds and butterflies. Proc R Soc B. 1999;266(1425):1221-5.

98. Alfaro ME, Santini F, Brock C, Alamillo H, Dornburg A, Rabosky DL, et al. Nine exceptional radiations plus high turnover explain species diversity in jawed vertebrates. Proc Natl Acad Sci U S A. 2009;106(32):13410-4.

99. Kozak KH, Wiens JJ. Testing the relationships between diversification, species richness, and trait evolution. Syst Biol. 2016;65(6):975-88.

100. Scholl JP, Wiens JJ. Diversification rates and species richness across the Tree of Life. Proc R Soc B. 2016;283(1838):20161334.

101. dos Reis M, Yang Z. Approximate likelihood calculation on a phylogeny for Bayesian estimation of divergence times. Mol Biol Evol. 2011;28(7):2161-72.

102. Guindon S. Rates and rocks: strengths and weaknesses of molecular dating methods. Front Genet. 2020;11:526-.

103. Hugall AF, Lee MS. The likelihood node density effect and consequences for evolutionary studies of molecular rates. Evolution. 2007;61(10):2293-307.

104. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics. 2014;30(9):1312-3.

105. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol. 2014;32(1):268-74.

106. Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. Syst Biol. 2010;59(3):307-21.

107. Venditti C, Meade A, Pagel M. Detecting the node-density artifact in phylogeny reconstruction. Syst Biol. 2006;55(4):637-43.

108. Sauquet H, Ho SYW, Gandolfo MA, Jordan GJ, Wilf P, Cantrill DJ, et al. Testing the impact of calibration on molecular divergence times using a fossil-rich group: the case of Nothofagus (Fagales). Syst Biol. 2011;61(2):289-313.

109. Jarvis ED, Mirarab S, Aberer AJ, Li B, Houde P, Li C, et al. Whole-genome analyses resolve early branches in the tree of life of modern birds. Science. 2014;346(6215):1320-31.

110. Mitchell KJ, Cooper A, Phillips MJ. Comment on "Whole-genome analyses resolve early branches in the tree of life of modern birds". Science. 2015;349(6255):1460-.

111. Cracraft J, Houde P, Ho SYW, Mindell DP, Fjeldså J, Lindow B, et al. Response to Comment on "Whole-genome analyses resolve early branches in the tree of life of modern birds". Science. 2015;349(6255):1460-.

112. Meredith RW, Janečka JE, Gatesy J, Ryder OA, Fisher CA, Teeling EC, et al. Impacts of the Cretaceous terrestrial revolution and KPg extinction on mammal diversification. Science. 2011;334(6055):521-4.

113. Phillips MJ. Geomolecular dating and the origin of placental mammals. Syst Biol. 2016;65(3):546-57.

114. Misof B, Liu S, Meusemann K, Peters RS, Donath A, Mayer C, et al. Phylogenomics resolves the timing and pattern of insect evolution. Science. 2014;346(6210):763-7.

115. Tong KJ, Duchêne S, Ho SYW, Lo N. Comment on "Phylogenomics resolves the timing and pattern of insect evolution". Science. 2015;349(6247):487-.

116. Kjer KM, Ware JL, Rust J, Wappler T, Lanfear R, Jermiin LS, et al. Response to Comment on "Phylogenomics resolves the timing and pattern of insect evolution". Science. 2015;349(6247):487-.

117. Rabosky DL, Chang J, Title PO, Cowman PF, Sallan L, Friedman M, et al. An inverse latitudinal gradient in speciation rate for marine fishes. Nature. 2018;559(7714):392-5.

118. Wertheim JO, Sanderson MJ. Estimating diversification rates: how useful are
divergence times? Evolution. 2011;65(2):309-20.
119. Rabosky DL. Positive correlation between diversification rates and phenotypic
evolvability can mimic punctuated equilibrium on molecular phylogenies. Evolution.
2012;66(8):2622-7.
120. Pennell MW, Harmon LJ, Uyeda JC. Is there room for punctuated equilibrium in
macroevolution? Trends Ecol Evol. 2014;29(1):23-32.
121. Pennell MW, Harmon LJ, Uyeda JC. Speciation is unlikely to drive divergence rates.
Trends Ecol Evol. 2014;29(2):72-3.
122. Ritchie AM, Hua X, Cardillo M, Yaxley KJ, Dinnage R, Bromham L. Phylogenetic
diversity metrics from molecular phylogenies: modelling expected degree of error under
realistic rate variation. Divers Distrib. 2020;n/a(n/a).
123. Hartmann K, Wong D, Stadler T. Sampling trees from evolutionary models. Syst
Biol. 2010;59(4):465-76.
124. Pybus OG, Harvey PH. Testing macro-evolutionary models using incomplete
molecular phylogenies. Proc Biol Sci. 2000;267(1459):2267-72.
1245
1246