Gut microbiota of ring-tailed lemurs (*Lemur catta*) vary across natural and captive

populations and correlate with environmental microbiota

Sally L. Bornbusch[1†], Lydia K. Greene[2], Sylvia Rahobilalaina[3], Samantha Calkins[4], Ryan S.

Rothman[5], Tara A. Clarke[6], Marni LaFleur[7], Christine M. Drea[1]

[1]Department of Evolutionary Anthropology, Duke University, Durham NC

SLB: sally.bornbusch@gmail.com[†]

CMD: cdrea@duke.edu

[2]Duke Lemur Center, Durham NC

LKG: lydiakgreene@gmail.com

[3]Faculty of Sciences, University of Antananarivo , Antananarivo, Madagascar

SR: hobilalainasylvia@gmail.com

[4]Department of Psychology, Program in Animal Behavior and Conservation, Hunter College,

New York, NY

SC: samanthacalkins92@gmail.com

[5]Institute for the Conservation of Tropical Environments, Stony Brook University, Stony Brook,

NY

RSR: ryan.rothman@stonybrook.edu

[6]Department of Sociology and Anthropology, North Carolina State University, Raleigh, NC

TAC: taclarke@ncsu.edu

[7]Department of Anthropology, University of San Diego, 5998 Alcala Park, San Diego, CA

ML: marni.lafleur@gmail.com

† - corresponding author

Abstract

Background: Inter-population variation in host-associated microbiota reflects differences in the

hosts' environments, but this characterization is typically based on studies comparing few

populations. The diversity of natural habitats and captivity conditions occupied by any given host

species has not been captured in these comparisons. Moreover, intraspecific variation in gut

microbiota, generally attributed to diet, may also stem from differential acquisition of

environmental microbes – an understudied mechanism by which host microbiomes are directly

shaped by environmental microbes. To more comprehensively characterize gut microbiota in an

ecologically flexible host, the ring-tailed lemur (*Lemur catta*; n = 209), while also investigating

the role of environmental acquisition, we used 16S rRNA sequencing of lemur gut and soil

microbiota sampled from up to 13 settings, eight in the wilderness of Madagascar and five in

captivity in Madagascar or the U.S. Based on matched fecal and soil samples, we used source-

sink ecological theory to examine covariation between the two types of consortia.

Results: The diversity of lemur gut microbes varied markedly within and between settings.

Microbial diversity was not consistently greater in wild than in captive lemurs, indicating that

this metric is not necessarily an indicator of host habitat or condition. Variation in microbial

composition was inconsistent with either a single, representative gut community for wild

conspecifics or a universal 'signal of captivity' that homogenizes the gut consortia of captive

animals. Despite the similar, commercial diets of captive lemurs on both continents, lemurs

within Madagascar were compositionally most similar, suggesting that non-dietary factors

govern some of the variability. In particular, soil microbial communities were most distinct

between the two continents, and there was significant and context-specific covariation between

lemur gut and soil microbiota.

Conclusions: As one of the broadest, single-species investigations of primate microbiota, our

study highlights that gut consortia are sensitive to multiple scales of environmental differences.

This finding begs a reevaluation of the simple 'captive vs. wild' dichotomy. Beyond the

important implications for animal care, health, and conservation, our finding that environmental

acquisition may mediate aspects of host-associated consortia further expands the framework for

how host-associated and environmental microbes interact across different microbial landscapes.

Introduction

The structure of gut microbial communities within vertebrates is influenced in part by

endogenous host factors, such as genotype and physiology[1–3], and in part by exogenous factors,

such as sociality, seasonality, habitat quality, and diet[4–6]. These exogenous factors can influence

which microbial taxa in a gut community thrive or become depauperate, as amply demonstrated

in dietary studies[7–10], or they can provide opportunities for more direct routes of microbial

acquisition[11–14]. For example, the transmission of microbes between hosts, including horizontal

pathogen transfer[15–17] or vertical transmission during the birthing process and nursing[18,19], are

significant drivers of host health and development. There is, likewise, the potential for horizontal

acquisition of microbes via exposure to environmental consortia on natural (e.g., soil) and man-

made surfaces, plus on food and in water[12,20–23]; however, this latter route to shaping host-

associated communities, hereafter referred to as 'environmental acquisition,' remains

understudied. Here, we match-sampled ring-tailed lemur (*Lemur catta*) feces and soil from 13

settings, representing both a large portion of the lemurs' natural habitat range in Madagascar and

a range of captive housing conditions in Madagascar and the U.S. (Table 1), to (a) characterize

variation in host gut microbiota, (b) characterize variation in soil microbiota, and (c) test for any

covariation between host and soil communities. Examining environmental microbes alongside

host-associated communities is a first step to understanding the role of environmental acquisition

in population-level differences between host microbiomes.

Previous studies of intraspecific variation in gut microbiota, often framed using a 'wild vs.

captive' comparison, have provided valuable descriptions of differences in presumed extremes[24–

26]. For example, researchers often report a 'signal of captivity,' whereby the gut microbiota of

captive hosts differ significantly from those of wild conspecifics, converging on a perturbed or

'humanized' composition[25,27,28]. Perturbations of this nature are generally attributed to

commercial diets that include manufactured chow and cultivated produce[27,29,30]; nevertheless,

studies of captive populations have been focused on accredited zoos or rescue facilities that may

not represent the range of captive conditions or may be confounded by within-species

comparisons across continents[26,29,31]. Even comparative field studies have been limited in the

number of populations per species studied, typically to a few populations that differ on a given

metric of interest (e.g. season, habitat type or quality[32–34]). Because hosts experience a wider

range of environmental settings than is typically encompassed within wild vs. captive

comparisons, a broader comparative approach is necessary to provide a more comprehensive and

nuanced understanding of gut microbiota variation.

As noted, differential exposure to environmental microbes provides potential for horizontal

transmission and environmental acquisition[20,22,23,35–37], with the ingestion of specific microbes

being linked to novel digestive functions of the gut microbiota[38–40]. Under certain conditions,

environmental acquisition has been shown to outweigh vertical transmission as the main mode of

microbial colonization[41,42]. Although environmental acquisition may promote heterogeneity

within and between hosts[43], its role rarely has been considered a differentiating factor between

wild and captive hosts. Husbandry practices and veterinary care, for example, introduce cleaning

products and antibiotics to the microbial environment of captive animals[44,45], further

differentiating it from natural habitats[46], with potentially critical consequences to microbiome

structure and function.

Ring-tailed lemurs are semi-terrestrial, omnivorous strepsirrhine primates[47,48] that occupy

various habitats across southern Madagascar[49] and also survive well in captivity[50]. Their

ecological flexibility, coupled with existing knowledge about their gut microbiome[26,51–53],

motivates broader comparative study of intraspecific variation that takes environmental

acquisition of microbes into consideration. Here, we combine 16S rRNA sequencing and

statistical tools based on source-sink ecological theory[54] to analyze covariation between lemur

gut microbiota (e.g., sink communities) and soil microbiota (e.g., source communities). Given

the variability of environmental factors across our multiple settings, we expect the diversity,

membership, and composition of lemur gut microbiota and soil microbiota to differ within and

between three, broad 'environmental' conditions: wilderness, captivity in Madagascar, and

captivity in the U.S (Table 1).

If diet or habitat quality were the main driver of gut microbiota composition, we would

expect (a) wild lemurs to show variation between their natural settings, (b) captive lemurs,

regardless of continent, to show similar gut microbiota between their settings (reflecting

commercial diets and perturbed habitats), and (c) wild and captive lemurs to differ most

drastically from one another, in line with prior studies[27]. If, however, environmental acquisition

were to play a major role in shaping lemur gut microbiota, we would again expect (a) wild

lemurs to show variation between their natural settings (reflecting the soil microbiota of the

lemurs' habitat), but we would expect (b) Malagasy lemurs (wild and captive) to share certain

soil-derived microbiota, differing most drastically from lemurs in captivity in the U.S., and (c)

differential access to soil within captive conditions to correlate with differential soil-associated

microbes present in hosts. With regard to the latter, for example, we might expect greater

proportions of soil-associated microbes in captive lemurs that gain access to natural enclosures

compared to their counterparts that are housed indoors.

Results

Lemur gut microbiota: Variation in diversity, membership, and composition

Alpha diversity. Across the gut microbiota of all ring-tailed lemurs sampled in this study, metrics of alpha diversity differed significantly by environmental condition (Generalized Linear Models or GLM; Shannon: $F = 23.773$, $p < 0.001$; Faith's phylogenetic: $F = 4.415$, $p = 0.013$; Figures 1a, b) and by setting (GLM; Shannon: $F = 13.157$, $p < 0.001$; Faith's phylogenetic: $F = 5.628$, $p < 0.001$; Figures 1c, d; Supplementary Material 1). The gut microbiota of wild lemurs and captive lemurs in the U.S. (hereafter, 'captive$_{U.S.}$ lemurs') were similarly diverse overall (pairwise Wilcoxon test; Shannon: $p = 0.635$; Faith's phylogenetic: $p = 0.056$; Figures 1a, b), whereas those of captive lemurs in Madagascar (hereafter, 'captive$_M$ lemurs') were significantly less diverse (pairwise Wilcoxon test; Shannon, wild vs. captive$_M$ lemurs: $p < 0.001$; wild vs. captive$_{U.S.}$ lemurs: $p < 0.001$; Faith's phylogenetic, wild vs. captive$_M$ lemurs: $p = 0.022$; wild vs. captive$_{U.S.}$ lemurs: $p = 0.021$; Figures 1a, b). Within environmental condition, however, both metrics of alpha diversity varied widely between the different settings (Figures 1c, d; Supplementary Material 1). For example, among wild lemurs, setting was a significant predictor of both metrics of alpha diversity (GLM; Shannon diversity: $F = 20.768$, $p < 0.001$; Faith's phylogenetic: $F = 11.104$, $p < 0.001$). Sex was not a significant predictor in any models of either alpha diversity metric (Supplementary Material 1).

Community membership. The membership of lemur gut microbiota included 64 abundant taxa (i.e., those that accounted for >1% of sequences). Of these 64 taxa, only four (6.2%) were shared across lemurs from all settings: the genera *Bacteroides* (phylum Bacteroidetes), *Rikenellaceae RC9 gut group* (Bacteroidetes), *Erysipelotrichaceae* UCG-004 (Firmicutes), and *Treponema 2* (Spirochaetes). Within condition, five (7.8%) taxa were shared by all wild lemurs,

whereas 10 (15.6%) and six (9.4%) taxa were shared by captive$_M$ and captive$_{U.S.}$ lemurs,

respectively (Figure 2). Using Analysis of Compositions of Microbiomes (ANCOM), we

identified 801 amplicon sequence variants (ASVs) that were differentially abundant across the

three conditions. For example, members of the Erysipelotrichaceae family characterized the

microbiota of wild lemurs, whereas taxa from the Spirochaetaceae and Prevotellaceae families

were more abundant in the gut microbiota of captive lemurs from both continents.

Erysipelotrichaceae UCG-004 and *Treponema 2*, for example, were abundant in all lemurs

(Figure 2), but the log ratios of the two genera distinguished lemur gut microbiota by the three

environmental conditions and, in particular, differentiated wild lemurs from captive$_{U.S.}$ lemurs

(Figure 3).

Beta diversity. The composition of lemur gut microbial communities was significantly

distinct across the three environmental conditions, as revealed by beta diversity (Permutational

Multivariate Analysis of Variance or PERMANOVA; wild vs. captive$_M$ lemurs: pseudo-F =

30.169, $p < 0.001$; wild vs. captive$_{U.S.}$ lemurs: pseudo-F = 97.912, $p < 0.001$; captive$_M$ vs.

captive$_{U.S.}$ lemurs: pseudo-F = 20.808, $p < 0.001$). Across all subjects, gut microbiota

composition clustered distinctly by condition (principal coordinate analysis of unweighted

UniFrac distances; Figures 4a, b). One notable exception, however, owed to a single pet lemur:

Unlike its in-country peers (i.e., other captive$_M$ lemurs), its microbial community structure

matched those of wild lemurs (see arrows in Figures 4a, b).

Across the three environmental conditions, Random Forest Analysis accurately assigned 208

of the 209 gut microbial profiles to the correct condition, with a low (0.48%) out-of-bag (OOB)

error rate. Based on its gut microbiota, only the previously mentioned pet lemur (see arrows in

Figure 2a, b) was misclassified as a wild lemur. Across the 13 settings, Random Forest Analysis

accurately classified 189 of the 209 microbial profiles (OOB error = 9.57%). The gut microbial

communities of wild and captive lemurs in Madagascar were misclassified at rates of 7.9% and

7.3%, respectively, whereas those of captive lemurs in the U.S. were misclassified at a rate of

20.6%.

With respect to uniformity within environmental condition, the composition of gut microbial

communities were least dissimilar between wild lemurs and most dissimilar between captive$_M$

lemurs (Kruskal-Wallis test; main effect of condition on beta diversity: $\chi^2 = 27487$, $p < 0.0001$;

pairwise Wilcoxon test; within wild vs. within captive$_M$ lemurs: $p < 0.001$; within wild vs. within

captive$_{U.S.}$ lemurs: $p < 0.0001$; Figure 4c). Between conditions, the microbiota of wild and

captive$_M$ lemurs were the least dissimilar, whereas the microbiota of wild vs. captive$_{U.S.}$ lemurs

were the most dissimilar (pairwise Wilcoxon test: 'wild vs. captive$_M$' vs. 'wild vs. captive$_{U.S.}$', p

< 0.0001; 'wild vs. captive$_M$' vs. 'captive$_M$ vs. captive$_{U.S.}$', $p < 0.0001$; Figure 4c). Considering

wild lemurs only, microbiota composition clustered by setting (Figure 4d). Although there was

some overlap between settings, the patterns are suggestive of microbial 'signatures' across

different settings.

Soil microbiota: Variation in diversity, membership, and composition

<u>Alpha diversity</u>. Across the eight settings for which we sampled soil, the alpha diversity of

soil microbiota did not vary significantly between conditions (Kruskal-Wallis test; Shannon

diversity: $\chi^2 = 3.3457$, $p = 0.187$; Faith's phylogenetic: $\chi^2 = 3.433$, $p = 0.179$; Figure 5) nor

between settings (Kruskal-Wallis test; Shannon diversity: $\chi^2 = 7.496$, $p = 0.379$; Faith's

phylogenetic: $\chi^2 = 8.936$, $p = 0.257$; Figure 5). These null findings may owe to small sample

sizes.

195  Community membership. The membership of soil communities included 77 abundant taxa,

196 of which none were shared across all settings (Figure 6). Of the identified soil microbiota,

197 78.12% were unique to the soil samples and were not found in any lemur fecal samples. For the

198 five wild populations for which we sampled soil, only five abundant taxa were shared: the genera

199 *Bacillus* (phylum Firmicutes), *Steroidobacter* (Proteobacteria), *Bryobacter* (Acidobacteria), and

200 *RB41* (Acidobacteria), and an unidentified member of the class Subgroup 6 (Acidobacteria).

201 ANCOM identified nine ASVs that were differentially abundant across all soil samples, five of

202 which (55.6%) belonged to the Balneolaceae family. In addition, compared to soil from

203 Madagascar, the soil communities from the U.S. (hereafter, captivity$_{U.S.}$ soil) were differentially

204 enriched for the genus *Bacillus*. By contrast, members of the family Nitrososphaeraceae

205 (Thaumarchaeota) and the genus *Acinetobacter* (Proteobacteria) characterized soil from the

206 wilderness and captivity settings in Madagascar (hereafter, wilderness and captivity$_M$ soils,

207 respectively; Figure S1).

208  Beta diversity. The beta diversity of the soil microbiota varied between conditions (Figure 7),

209 but only significantly so between wilderness and captivity$_M$ soils (PERMANOVA; wilderness

210 vs. captivity$_M$ soils: pseudo-F = 1.337, p = 0.202; wilderness vs. captivity$_{U.S}$ soils: pseudo-F =

211 3.897, p = 0.012; captivity$_M$ vs. captivity$_{U.S}$ soils: pseudo-F = 7.752, p = 0.329). Variation in soil

212 communities within a condition was not significantly different between wilderness soils and

213 captivity$_{U.S.}$ soils (pairwise Wilcoxon test, p = 0.130; Figure 7c). Between conditions, wilderness

214 and captivity$_M$ soils had the lowest dissimilarities (pairwise Wilcoxon test; 'wilderness vs.

215 captivity$_M$' vs. 'wilderness vs. captivity$_{U.S}$' soils: p < 0.001; 'wilderness vs. captivity$_M$' vs.

216 'captivity$_M$ vs. captivity$_{U.S}$': p = 0.016; 'wild vs. captivity$_{U.S}$' vs. 'captivity$_M$ vs. captivity$_{U.S}$': p =

217 0.338 Figure 7c).

Covariation of lemur gut and soil microbiota

There were 191 ASVs shared between lemur gut and soil microbiota. These were dominated

by members of the Firmicutes (75 ASVs or 39.3%), Proteobacteria (49 ASVs or 25.6%), and

Bacteroidetes (38 ASVs or 19.9%) phyla. Although many of the shared taxa were abundant

(>1%) in either lemur gut or soil microbiota, only one genus, *Acinetobacter* (Proteobacteria), was

abundant in both lemur gut and soil microbiota.

As would be predicted if environmental acquisition impacts host microbial communities,

there was a significant correlation between the abundances of microbes in lemur feces and soil

samples (Mantel test; $r = 0.494$, $p < 0.001$). The proportion of 'soil-associated' microbes found

in lemur gut microbiota varied significantly across conditions (Kruskal-Wallis test; $\chi^2 = 73.862$,

$p < 0.001$; Figure 8a) and settings (Kruskal-Wallis test; $\chi^2 = 112.69$, $p < 0.001$; Figure 8b).

Overall, the gut microbiota of wild lemurs had significantly greater proportions of soil-associated

microbes compared to those of all captive lemurs (pairwise Wilcoxon test, $p < 0.001$; Figure 8).

In addition, captive$_M$ lemurs had significantly greater proportions of soil-associated microbes in

their gut microbiota compared to captive$_{U.S.}$ lemurs (pairwise Wilcoxon test; $p < 0.001$; Figure

8). For lemurs housed at the DLC, those that semi-free-ranged in outdoor, natural habitat

enclosures had significantly greater proportions of soil-associated microbes in their gut

microbiota compared to lemurs that did not have access to forested enclosures (Kruskal-Wallis

test; $\chi^2 = 4.641$, $p = 0.031$; Figure 8c).

Soil from within a lemur's setting accounted for, on average, significantly greater proportions

of the lemur's gut microbiota than did soil communities from other settings (Figure 9,

Supplementary Material 2). Overall, the greatest proportion of soil-associated microbes within

lemur gut microbiota occurred when the lemurs and soil were both from the wilderness (Figure

9; Supplementary Material 2). The proportion of soil-associated microbes in lemur gut

microbiota dropped to near zero when comparing the lemur gut and soil microbiota between

samples from the wilderness and those from captivity in the U.S. (Figure 9; Supplementary

Material 2).

Discussion

Through extensive fecal and soil sampling from multiple settings representing the ring-

tailed lemurs' natural range in Madagascar and in captivity on two continents, we have

highlighted (1) the wide and often underrepresented variety of gut microbiota present within a

single host species, (2) the lack of a universal 'signal of captivity' that uniformly decreases

microbial diversity, (3) aspects of microbiota membership and composition that differ markedly

between wild and captive populations, and (4) covariance between lemur gut and soil microbiota,

which points to a key role of environmental microbes. Researchers have reported host 'group

signatures' in microbiota, often attributed to the social transmission of microbes[5,55–58]; our results

expand this concept to 'population signatures' and draw attention to the potential role of

environmental acquisition of microbes in mediating significant inter-population variation.

Across populations of wild lemurs, we first observed substantial variation in gut microbiota

diversity, membership, and composition, indicating that there is not a single 'representative' gut

community for wild ring-tailed lemurs, as is likely the case for most host species. Nonetheless,

the pattern of natural variation observed did not always meet expectations. For example, lemurs

living in what is considered a relatively 'pristine' site, IVO – a recently discovered humid forest

patch that is relatively undisturbed by human activity – unexpectedly had the second-lowest

diversity of gut microbes. To the extent that lack of disturbance is a proxy of habitat quality, this

pattern would be inconsistent with previous reports that greater habitat quality promotes more

diverse gut microbiota[59,60]. In prior studies, the gut microbiota of ring-tailed lemurs were

relatively unaffected by habitat degradation[52]. Therefore, either pristine habitats can be of low

quality or the ecological and dietary flexibility of this species may dampen the impact of

variation in habitat quality and type, relative to more specialized primates (e.g., folivores)[26,61–63].

That we found significant, natural, inter-population variation in a relatively hardy and robust

species[49,64] suggests that hosts with greater sensitivity to environmental variation, including

habitat quality and type, would likely show even greater variation than that described herein. If

so, studies constrained to single or few host populations are likely to underrepresent the wide-

scale, natural variation in host gut microbiota.

We next observed gut microbiota to be compositionally distinct across populations of captive

lemurs. Contrary to many previous studies[65–68], but consistent with others[69–72], the gut microbiota

of captive lemurs were not consistently less diverse than those of wild lemurs nor were they

compositionally homogenized by the similar commercial diets provided to captive subjects[67,73].

Heterogeneous gut microbiota could reflect slight differences in the diets provided (as the

produce and browse available differ between captivity settings), but such minor dietary variation

is unlikely to be the sole driver of such marked microbial differences, particularly in an

omnivorous host. Non-dietary factors must have contributed to distinguishing the gut

communities of captive lemurs. Indeed, the gut microbiota of captive lemurs in Madagascar were

compositionally more similar to those of their wild counterparts than to those of captive lemurs

in the U.S. Based on this observation, we suggest that the effect of a commercial diet is not

necessarily the strongest differentiator of gut consortia and that the effects of captivity cannot be

standardized across populations. Our results raise questions about the commonly held view that

greater alpha diversity is both (a) a hallmark of wild individuals and (b) a proxy for a healthier

gut community[74–78]. Although we did not assess gut health, pet lemurs fed diets of rice and fruit,

living in close contact with people and domestic animals, often housed solitarily indoors, are

prone to disease[79–81]; yet, their gut consortia were as diverse as those of wild lemurs living at the

relatively pristine site, IVO. Moreover, captive lemurs from the DLC and NCZ in the U.S. had

some of the most diverse gut consortia, equaling the greatest diversity seen in wild lemurs (e.g.,

in BEZ lemurs). These results add to the mounting evidence[61,82,83] that alpha diversity alone,

without the context of host ecology and other microbiome data, should not be used to extrapolate

about the health state of gut consortia or the host's environment.

We also found that, between wild and captive lemurs, the membership and composition of

gut microbiota was indicative of the environmental condition. There was little evidence of a

diverse 'core' microbiome, as only four taxa were found to be abundant across all lemur

populations. Two of those core taxa, *Erysipelotrichaceae UCG-004* and *Treponema 2*, were

differentially abundant between the three conditions. Despite links between members of

Erysipelotrichaceae and high-fat, commercial diets in humans[84], *Erysipelotrichaceae* microbes

were reported to be enriched in wild compared to captive chimpanzees[85], mirroring our findings

in lemurs. Furthermore, the genus *Erysipelotrichaceae UCG-004* was more abundant in the gut

microbiota of chimpanzees, relative to humans[86], and in folivorous woolly lemurs compared to

other lemur species[87]. The functionally diverse members of the *Treponema* genus were more

abundant in the gut microbiota of captive vs. wild hosts in other species[85,88]. *Treponema*

members break down pectin[89,90], a complex plant polysaccharide enriched in ripe fruits, such as

those commonly provided to captive ring-tailed lemurs[91,92]. Compositionally, the gut microbiota

of wild lemurs were markedly less varied than those of lemurs in all captivity settings. These

findings support the "Anna Karenina" principle[93,94], which posits that perturbations of microbiota

result in unstable communities and, thus, 'unperturbed' hosts have less variation in their

microbiota than do 'disturbed' hosts. A single exception to the gut microbiota clustering

according to the hosts' conditions was a pet lemur with gut microbiota that resembled that of

wild lemurs. Although we can only speculate about this individual's history, if recently taken

from their natural habitat, the gut microbiota could still reflect the wild origins of this animal,

potentially indicative of gradual change in response to environmental shifts[95,96].

Lastly, we observed that patterns in lemur gut microbiota were somewhat mirrored in the

diversity and composition of soil microbiota, suggesting that environmental conditions other than

diet, including exposure to external microbes in soils, may influence gut microbiomes[97].

Madagascar's geographical isolation for ~88 million years accounts for high levels of floral and

faunal endemism[98,99]. The same is true of microbes, as evidenced by the numerous, unique

pathogenic microorganisms found on the island[100–103]. Unsurprisingly, therefore, soil microbiota

in Madagascar, whether originating in wilderness or captivity settings, were similar in

composition and significantly divergent from soils in the U.S.[104]. Given the disparate geographic

distributions of many wild vs. captive animals, environmental acquisition that reflects local

microbial endemism may be particularly relevant for distinguishing gut microbiota between wild

and captive conspecifics. For example, the natural ranges of most primates occur in the

tropics[105,106], yet most accredited zoos and captive facilities that house primates are found outside

of tropical regions (in e.g., Europe and North America)[107,108]; the distinct environmental

consortia surrounding wild and captive conspecifics should reflect their geographic or

continental divides.

Regarding the exposure to environmental microbes, soil-associated microbes were more prevalent in lemurs that had greater exposure to natural environments and the acquired soil microbes were specific to the lemurs' environment, reflecting active environmental acquisition. This observation expands on findings that abiotic soil properties mediate primate gut microbiota[97]. Wild and captive ring-tailed lemurs perform geophagy (i.e., earth-eating), a behavior that is linked to nutrient and microbial supplementation[109,110] and is a potential vector for the incorporation of environmental microbes[39]. Similarly, dietary items may act as vessels of soil or environmental microbes[40]; dietary variation across wild and captive lemurs may influence gut microbiomes by simultaneously offering different nutrients and different microbes. Akin to most cross-sectional studies of microbiomes, we were unable to assess the persistence or viability of the soil-associated microbes in lemur gut communities. It is, therefore, possible that the soil-associated microbes in lemur guts were ephemeral or non-viable; however, our results indicate setting-specific, environmental acquisition, supporting that these patterns are not random and that the acquired microbes may be subject to filters that enable the incorporation of only specific microbes[20,111,112]. Furthermore, we analyzed these data from the perspective that environmental consortia act as sources of microbes for host-associated communities, but we expect consistent, bidirectional transmission of microbes between hosts and their environments, a relationship that warrants further investigation.

While expanding our understanding of the factors that shape host-microbe relationships, these results also have significant potential to inform animal care and conservation strategies. Perturbed microbiota are increasingly recognized as culprits of obesity, gastrointestinal distress, and even associated mortality in captive animals[113–115,75]. Given that lemurs are among the most endangered mammals on the planet[116], maintaining populations of healthy animals in captivity is

an important 'safety net' that augments *in-vivo* conservation efforts[117,118]. We suggest that

environmental acquisition may be a key component of 'rewilding' or 'bioaugmenting' captive

animal gut microbiota, a process by which gut consortia can be reshaped to better promote host-

microbe symbiosis[26,117,119]. Identifying what comprises healthy gut microbiota is a complex and

ongoing area of research; nonetheless, we show that environmental acquisition is a potential

driver of microbial communities and thus should be considered as a component of animal health.

Conclusions

Even in a relatively robust, omnivorous host, gut microbiota are distinct across populations.

This variation reflects environmental variability that is underrepresented by a simple wild vs.

captive dichotomy. Moreover, concurrent analysis of lemur gut and soil microbiota supports the

premise that environmental acquisition contributes to shaping host-associated microbiota; hosts

and their associated microbes are components of a larger landscape that includes interactions

with environmental microbes. Together, these results expand our understanding of intraspecific

host-microbe dynamics under varying environmental conditions and reinforce the value of

broad-scale, comparative investigations of microbial variation within a single host species.

Methods

Study sites

Our research sites included 13 settings (one per 'population'), grouped under the following

three environmental conditions: wilderness in Madagascar (8 settings), captivity in Madagascar

(2 settings), and captivity in the U.S. (3 settings; Table 1). The wilderness settings occurred in

protected areas (e.g., national parks, community-managed reserves) that varied in habitat type

(Table 1). The captivity settings in Madagascar included the Lemur Rescue Center (LRC;

Toliara, Madagascar), where the animals were socially housed, and various townships that were

home to individual pets. Lastly, the captivity settings in the U.S. included the North Carolina

Zoo (NCZ; Asheboro, NC), the Duke Lemur Center (DLC; Durham, NC), and the National

Zoological Park (NZP; Washington, DC). These facilities were comparable to one another, all

with socially housed lemurs.

Subjects

Across all research sites, our subjects included 215 adult, ring-tailed lemurs (82 male, 81

female, 52 of unknown sex; Table 1). The wilderness sites were each occupied by multiple lemur

troops, ranging in size from 5-24 individuals. Excluding the pets, all captive settings included

groups of 2-7 lemurs that had access to indoor and outdoor enclosures, and were provided

facility-standardized diets (i.e., fresh produce and commercial chow, freely available water).

Certain animals at the LRC and the DLC also had access to natural habitat enclosures that,

respectively, consisted of dry and spiny forest (LRC) or North American deciduous and pine

hardwood forest (DLC). The pets were kept in human dwellings (i.e., houses or hotels) and were

fed fruit, rice, and other foods intended for human consumption.

Sample collection

During a span of four years (2016-2020), we collected 'matched' fecal and soil samples from

our subjects and study sites, respectively. Within 8 weeks of fecal or soil collection, the samples

were transported to the U.S., where they were stored at -80 °C, until analysis.

For feces, we opportunistically collected fresh samples, upon the lemur's observed voiding.

In Madagascar, collections occurred during the dry season (May-October) and, in the U.S.,

collections occurred end of summer through fall (August-November). To avoid soil

contamination of the fecal sample, we removed the outer layer of each fecal pellet. We then

placed the sample in an Omnigene tube that contained a stabilizing buffer that preserved

microbial communities at room temperature for 8 weeks (Omnigene.Gut tube, DNAgenotek,

Ontario, Canada[120,121]). All settings were represented by fecal samples from minimally two

lemurs (the maximum number of individuals represented was 33).

When collecting soil in nature, we avoided high-defecation areas (e.g., under sleeping trees)

while identifying core areas where lemurs most commonly spent time on the ground. Within

these core areas, we demarcated a 2-3 m^2 area and collected soil from each of five evenly spaced

locations, using a clean, individually wrapped, sterile plastic spatula. For each area, the five

aliquots of topsoil (top 2-3 cm of soil) were pooled in a single Omnigene tube to create a

representative soil sample. Because multiple lemur troops inhabited each of the wilderness

settings, in some cases with overlapping core areas, we prioritized collecting soil samples from

areas of maximal use. In some cases, we were unable to collect soil samples for every troop that

provided fecal samples. At the LRC and DLC, we used the same collection methods to collect

soil samples from areas in the natural habitat enclosures where lemurs semi-free-ranged. Because

it is illegal to own pet lemurs in Madagascar, we minimized owner concern by collecting only

fecal samples for this group. Because of other logistical and analytical constraints (see below),

only eight of the 13 settings were represented by usable, pooled soil samples.

Microbial DNA extraction and sequencing

Following the manufacturer's protocols for the DNeasy Powersoil kit (QIAGAN, Frederick,

MD), we extracted bacterial genomic DNA from fecal and soil samples. We quantified DNA

using a Fluorometer (broad-spectrum kit, Qubit 4, Thermo Fisher Scientific, Waltham, MA).

Aliquots of extracted DNA were sent to Argonne National Laboratory's Environmental

Sequencing facility (Lemont, IL) for library preparation and amplicon sequencing of the 16S

rRNA gene. After amplification of the V4 region with region-specific primers and sample-

specific 12-base barcodes, samples were pooled and amplicon libraries were cleaned using

AMPure XP Beads. Amplicons were then sequenced on a 151 x 151 base pair Illumina MiSeq

run[122].

Bioinformatics and statistics

We processed the raw sequence data using a previously published bioinformatics pipeline

generated in QIIME2[123]. In brief, we used the pipeline to join forward and reverse reads,

demultiplex and quality filter the joined reads (DADA2), generate a phylogenetic tree, and

assign taxonomy based on 99% sequence similarity (SILVA database [124,125], ver. 138.1). After

quality filtering, samples with fewer than 10,000 sequences were removed from downstream

analyses, resulting in 209 fecal samples and 25 soil samples with over 11 million combined reads

and an average of ~50,000 reads per sample. To visually represent rare taxa that had relative

abundances < 1% of the total sequences, we combined them into the conglomerate "Other"

category (Figures 1 and 6). Using tables of amplicon sequence variants (ASVs), we calculated

metrics of alpha diversity (Shannon and Faith's Phylogenetic diversity metric) and beta diversity

(UniFrac distances).

To test for differences in alpha diversity between the gut microbiota of lemurs under the

three environmental conditions and in the 13 settings, we first used generalized linear models

(GLMs; glm in R, ver, 4.0.2) with condition or setting and sex as fixed effects. To further test for

variation in lemur gut microbiota and soil microbiota alpha diversity, we used nonparametric

statistics (e.g., Kruskal-Wallis tests, and pairwise Wilcoxon rank sum tests with Benjamini-

Hochberg adjustment) to perform pairwise comparisons between the various conditions and

settings. To identify and test for effects of condition or setting on beta diversity (unweighted

UniFrac distances) in lemur fecal and soil microbiota, we used principal coordinate analysis (i.e.,

to visualize clustering of microbiota composition) and Permutational Multivariate Analysis of

Variance (PERMANOVA) in QIIME2. We then performed Random Forest Analysis[126], which is

a supervised learning technique that uses decision trees to classify data to specific categories and

provides an overall model error rate (out of the bag error or OOB error). To identify microbes

enriched in specific groups of samples, we used differential abundance analyses via Analysis of

Compositions of Microbiomes (ANCOM) and songbird software[127] in QIIME2, paired with

visualization through Qurro[128].

For the eight settings where we obtained matched fecal and soil samples (Table 1), we

analyzed covariation between lemur gut microbiota and the associated soil communities by

performing a Mantel test on microbial abundance matrices of lemur gut and soil microbiota.

Because multiple lemur fecal samples were associated with each soil sample, we created

comparable matrices for the Mantel test by averaging the microbial abundances across the fecal

samples of lemurs directly associated with a given soil sample, resulting in a single, mean lemur

gut community associated with each soil community. For this process, we omitted fecal samples

from troops not represented by a soil sample or for which troop identity was unknown.

To test if soil-associated microbes were present in lemur gut microbiota, we used FEAST, a

tool for fast expectation-maximization microbial source tracking[129]. FEAST assumes each 'sink'

sample is a convex combination of known and unknown 'sources' and uses multinomial

distributions and machine-learning classification to model the microbial source-sink data[129]. For

this analysis, we used the matched lemur gut and soil samples; all soil samples collected in a

given setting were used to represent the potential exposure to environmental microbes

experienced by all sampled lemurs in that same setting, regardless of troop identity. Because we

were testing whether environmental acquisition influences lemur gut microbiota, and because

this analysis requires an assumption of directionality (i.e., from a source to a sink), we

categorized soil samples as 'sources' and lemur fecal samples as 'sinks'; however, we

acknowledge and discuss the potential for bi-directional transmission of microbes between

lemurs and soil. For each lemur fecal sample, we calculated the proportions of microbes that

were identified as stemming from each soil community and from a default 'unknown source' that

accounts for microbes not relevant to soil microbiota. Lastly, we used FEAST to test for

differences in the proportion of soil microbes in the gut microbiota of lemurs at the DLC that

were either semi-free-ranging or sequestered to indoor enclosures.

Declarations

Ethics approval and consent to participate

Sampling in Madagascar occurred with approval from Madagascar National Parks and

appropriate governmental agencies (Ministry of Environment, Ecology, and Forests; permit #s

147/18/MEEF/SG/DGF/DSAP/SCB.Re, 152/19/MEDD/SG/DGEF/DGRNE,

159/16/MEEF/SG/DGF/DSAP/SCB.Re, 154/17/ MEEF/SG/DGF/DSAP/SCB.Re,

492 156/19/MEEF/SG/DGF/DSAP/SCB.Re). Sampling at the DLC, NCZ, and NZP occurred with

493 approval from the appropriate Animal Care and Use Committees (Duke University's Institutional

494 Animal Care and Use Committee: protocol #A111-16-05; North Carolina Zoo Animal Care and

495 Use Committee: approved without protocol number; Smithsonian National Zoological Park

496 Research Animal Care and Use Committee: approved without protocol number).

497 *Consent for publication*

498 Not applicable. This study does not contain any individual person's data in any form.

499 *Availability of data and material*

500 Sequencing reads are available in the National Center for Biotechnology Information's

501 Sequence Read Archive (BioProject ID #TBD, BioSample accession #s TBD). Additional

502 datasets generated and/or analyzed during the current study are available from the corresponding

503 author upon reasonable request.

504 *Competing interests*

505 We attest that no author has financial or non-financial competing interests.

506 *Funding*

507 Funding was provided by awards from the National Science Foundation (BCS 1749465 to

508 CMD), the Triangle Center for Evolutionary Medicine (Graduate Student Research Award to

509 SLB), the Kenan Institute for Ethics at Duke University (Anthropocene Graduate Research Grant

510 to SLB). During collections, ML was funded by the Margot Marsh Biodiversity Fund.

511 *Authors' contributions*

512 SLB and CMD conceived of the study, with input from LKG. SLB, LKG, SR, SC, RSR,

513 TAC, and ML collected samples, documented metadata, and transported materials/samples. SLB

performed the bioinformatic and statistical analyses. SLB and CMD wrote the manuscript and all

authors read and approved the submitted version.

Acknowledgements

For their assistance with sample collection in the wilderness sites, we are deeply grateful to

Laurent 'Raleso' Randrianasolo, Remi Rakotovao, Georges René Rakotonirina, Soatata Honore

Reseva, Chelsea Southworth, Melina Nolas, and Lauren Petronaci. We thank Dr. Patricia Wright

and the Centre ValBio for providing accessibility to the IVO collection site and assistance in

sample transportation and storage. We further thank the current and past staff members of the

LRC, DLC, NCZ, and NZP for their assistance with sample collection in captivity settings. We

are grateful to Sarah Owens at Argonne National Laboratory for providing guidance and

sequencing services. This is DLC publication number (##TBD).

References

1   Hansen J, Gulati A, Sartor RB. The role of mucosal immunity and host genetics in

defining intestinal commensal bacteria. *Curr Opin Gastroenterol* 2010; **26**: 564.

2   Amato KR, Sanders JG, Song SJ, Nute M, Metcalf JL, Thompson LR, Morton JT, Amir

A, McKenzie VJ, Humphrey G. Evolutionary trends in host physiology outweigh dietary

niche in structuring primate gut microbiomes. *ISME J* 2019; : 1.

3   Milani C, Alessandri G, Mancabelli L, Mangifesta M, Lugli GA, Viappiani A, Longhi G,

Anzalone R, Duranti S, Turroni F. Deciphering the impact of diet and host physiology on

the mammalian gut microbiome by multi-omics approaches. *Appl Environ Microbiol*

2020.

4   Rothschild D, Weissbrod O, Barkan E, Kurilshikov A, Korem T, Zeevi D, Costea PI,

Godneva A, Kalka IN, Bar N. Environment dominates over host genetics in shaping

human gut microbiota. *Nature* 2018; **555**: 210–215.

5   Tung J, Barreiro LB, Burns MB, Grenier JC, Lynch J, Grieneisen LE, Altmann J, Alberts

SC, Blekhman R, Archie EA. Social networks predict gut microbiome composition in

wild baboons. *Elife* 2015; **2015**. doi:10.7554/eLife.05224.

6   Tasnim N, Abulizi N, Pither J, Hart MM, Gibson DL. Linking the gut microbial

ecosystem with the environment: does gut health depend on where we live? *Front

Microbiol* 2017; **8**: 1935.

7   David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, Ling A V,

Devlin AS, Varma Y, Fischbach MA. Diet rapidly and reproducibly alters the human gut

microbiome. *Nature* 2014; **505**: 559.

8   Kartzinel TR, Hsing JC, Musili PM, Brown BRP, Pringle RM. Covariation of diet and gut

microbiome in African megafauna. *Proc Natl Acad Sci* 2019; **116**: 23588–23593.

9   Youngblut ND, Reischer GH, Walters W, Schuster N, Walzer C, Stalder G, Ley RE,

Farnleitner AH. Host diet and evolutionary history explain different aspects of gut

microbiome diversity among vertebrate clades. *Nat Commun* 2019; **10**: 1–15.

10   Greene LK, McKenney EA, O'Connell TM, Drea CM. The critical role of dietary foliage

in maintaining the gut microbiome and metabolome of folivorous sifakas. *Sci Rep* 2018; **8**:

14482.

11   Peccia J, Kwan SE. Buildings, beneficial microbes, and health. *Trends Microbiol* 2016;

24: 595–597.

12   Hyde ER, Navas-Molina JA, Song SJ, Kueneman JG, Ackermann G, Cardona C,

Humphrey G, Boyer D, Weaver T, Mendelson JR. The oral and skin microbiomes of

captive komodo dragons are significantly shared with their habitat. *MSystems* 2016; **1**:

e00046-16.

13 Cardona C, Lax S, Larsen P, Stephens B, Hampton-Marcell J, Edwardson CF, Henry C,

Van Bonn B, Gilbert JA. Environmental sources of bacteria differentially influence host-

associated microbial dynamics. *MSystems* 2018; **3**.

14 Kuthyar S, Manus MB, Amato KR. Leveraging non-human primates for exploring the

social transmission of microbes. *Curr Opin Microbiol* 2019; **50**: 8–14.

15 Durrer S, Schmid-Hempel P. Shared use of flowers leads to horizontal pathogen

transmission. *Proc R Soc London Ser B Biol Sci* 1994; **258**: 299–302.

16 Kulkarni S, Heeb P. Social and sexual behaviours aid transmission of bacteria in birds.

Behav Processes 2007; **74**: 88–92.

17 Pedersen AB, Davies TJ. Cross-species pathogen transmission and disease emergence in

primates. *Ecohealth* 2009; **6**: 496–508.

18 Arora N, Sadovsky Y, Dermody TS, Coyne CB. Microbial vertical transmission during

human pregnancy. *Cell Host Microbe* 2017; **21**: 561–567.

19 Funkhouser LJ, Bordenstein SR. Mom knows best: the universality of maternal microbial

transmission. *PLoS Biol* 2013; **11**: e1001631.

20 Walke JB, Becker MH, Loftus SC, House LL, Cormier G, Jensen R V, Belden LK.

Amphibian skin may select for rare environmental microbes. *ISME J* 2014; **8**: 2207.

21 Seedorf H, Griffin NW, Ridaura VK, Reyes A, Cheng J, Rey FE, Smith MI, Simon GM,

Scheffrahn RH, Woebken D. Bacteria from diverse habitats colonize and compete in the

mouse gut. *Cell* 2014; **159**: 253–266.

22 Smith CCR, Snowberg LK, Caporaso JG, Knight R, Bolnick DI. Dietary input of

microbes and host genetic variation shape among-population differences in stickleback gut

microbiota. *ISME J* 2015; **9**: 2515–2526.

23   Selway CA, Mills JG, Weinstein P, Skelly C, Yadav S, Lowe A, Breed MF, Weyrich LS.

Transfer of environmental microbes to the skin and respiratory tract of humans after urban

green space exposure. *Environ Int* 2020; **145**: 106084.

24   Schmidt E, Mykytczuk N, Schulte-Hostedde AI. Effects of the captive and wild

environment on diversity of the gut microbiome of deer mice (Peromyscus maniculatus).

ISME J 2019; **13**: 1293–1305.

25   McKenzie VJ, Song SJ, Delsuc F, Prest TL, Oliverio AM, Korpita TM, Alexiev A, Amato

KR, Metcalf JL, Kowalewski M. The Effects of Captivity on the Mammalian Gut

Microbiome. *Integr Comp Biol* 2017; **57**: 690–704.

26   Greene LK, Bornbusch SL, McKenney EA, Harris RL, Gorvetzian SR, Yoder AD, Drea

CM. The importance of scale in comparative microbiome research: New insights from the

gut and glands of captive and wild lemurs. *Am J Primatol* 2019.

27   Clayton JB, Vangay P, Huang H, Ward T, Hillmann BM, Al-Ghalith GA, Travis DA,

Long HT, Tuan B Van, Minh V Van *et al.* Captivity humanizes the primate microbiome.

Proc Natl Acad Sci 2016; **113**: 10376–10381.

28   Yao R, Xu L, Hu T, Chen H, Qi D, Gu X, Yang X, Yang Z, Zhu L. The "wildness" of the

giant panda gut microbiome and its relevance to effective translocation. *Glob Ecol*

Conserv 2019; **18**: e00644.

29   Frankel JS, Mallott EK, Hopper LM, Ross SR, Amato KR. The effect of captivity on the

primate gut microbiome varies with host dietary niche. *Am J Primatol* 2019; **81**: e23061.

30   Bornbusch SL, Greene LK, McKenney EA, Volkoff SJ, Midani FS, Joseph G, Gerhard

WA, Iloghalu U, Granek J, Gunsch CK. A comparative study of gut microbiomes in captive nocturnal strepsirrhines. *Am J Primatol* 2019; **81**: e22986.

Gibson KM, Nguyen BN, Neumann LM, Miller M, Buss P, Daniels S, Ahn MJ, Crandall KA, Pukazhenthi B. Gut microbiome differences between wild and captive black rhinoceros–implications for rhino health. *Sci Rep* 2019; **9**: 1–11.

Ren T, Boutin S, Humphries MM, Dantzer B, Gorrell JC, Coltman DW, McAdam AG, Wu M. Seasonal, spatial, and maternal effects on gut microbiome in wild red squirrels. *Microbiome* 2017; **5**: 1–14.

Hicks AL, Lee KJ, Couto-Rodriguez M, Patel J, Sinha R, Guo C, Olson SH, Seimon A, Seimon TA, Ondzie AU. Gut microbiomes of wild great apes fluctuate seasonally in response to diet. *Nat Commun* 2018; **9**: 1–18.

Baniel A, Amato KR, Beehner JC, Bergman TJ, Mercer A, Perlman RF, Petrullo L, Reitsema L, Sams S, Lu A. Seasonal shifts in the gut microbiome indicate plastic responses to diet in wild geladas. *Microbiome* 2021; **9**: 1–20.

Guarner F, Bourdet-Sicard R, Brandtzaeg P, Gill HS, McGuirk P, Van Eden W, Versalovic J, Weinstock J V, Rook GAW. Mechanisms of disease: the hygiene hypothesis revisited. *Nat Clin Pract Gastroenterol Hepatol* 2006; **3**: 275–284.

Chen C-Y, Chen P-C, Weng FC-H, Shaw GT-W, Wang D. Habitat and indigenous gut microbes contribute to the plasticity of gut microbiome in oriental river prawn during rapid environmental change. *PLoS One* 2017; **12**: e0181427.

Mushegian AA, Arbore R, Walser J-C, Ebert D. Environmental sources of bacteria and genetic variation in behavior influence host-associated microbiota. *Appl Environ Microbiol* 2019; **85**.

38 Sylvain F-É, Derome N. Vertically and horizontally transmitted microbial symbionts

shape the gut microbiota ontogenesis of a skin-mucus feeding discus fish progeny. *Sci Rep*

2017; **7**: 1–14.

39 Borruso L, Checcucci A, Torti V, Correa F, Sandri C, Luise D, Cavani L, Modesto M,

Spiezio C, Mimmo T. I Like the Way You Eat It: Lemur (Indri indri) Gut Mycobiome and

Geophagy. *Microb Ecol* 2021; : 1–9.

40 Li H, Li T, Yao M, Li J, Zhang S, Wirth S, Cao W, Lin Q, Li X. Pika gut may select for

rare but diverse environmental bacteria. *Front Microbiol* 2016; **7**: 1269.

41 Kikuchi Y, Hosokawa T, Fukatsu T. Insect-microbe mutualism without vertical

transmission: a stinkbug acquires a beneficial gut symbiont from the environment every

generation. *Appl Environ Microbiol* 2007; **73**: 4308–4316.

42 Inoue R, Ushida K. Vertical and horizontal transmission of intestinal commensal bacteria

in the rat model. *FEMS Microbiol Ecol* 2003; **46**: 213–219.

43 Leftwich PT, Edgington MP, Chapman T. Transmission efficiency drives host–microbe

associations. *Proc R Soc B* 2020; **287**: 20200820.

44 Maamar S Ben, Hu J, Hartmann EM. Implications of indoor microbial ecology and

evolution on antibiotic resistance. *J Expo Sci Environ Epidemiol* 2020; **30**: 1–15.

45 Hartmann EM, Hickey R, Hsu T, Betancourt Román CM, Chen J, Schwager R, Kline J,

Brown GZ, Halden RU, Huttenhower C. Antimicrobial chemicals are associated with

elevated antibiotic resistance genes in the indoor dust microbiome. *Environ Sci Technol*

2016; **50**: 9807–9815.

46 Thompson LR, Sanders JG, McDonald D, Amir A, Ladau J, Locey KJ, Prill RJ, Tripathi

A, Gibbons SM, Ackermann G. A communal catalogue reveals Earth's multiscale

microbial diversity. *Nature* 2017; **551**: 457–463.

47   Jolly A, Sussman RW, Koyama N, Rasamimanana H. *Ringtailed Lemur Biology: Lemur*

catta in Madagascar (Google eBook).

2006http://books.google.com/books?id=WQ5yWlEJVVYC&pgis=1.

48   Gould L. Lemur catta ecology: what we know and what we need to know. In: *Lemurs*.

Springer, 2006, pp 255–274.

49   Gabriel DN. Habitat Use and Activity Patterns as an Indication of Fragment Quality in a

Strepsirrhine Primate. *Int J Primatol* 2013; **34**: 388–406.

50   Mason GJ. Species differences in responses to captivity: stress, welfare and the

comparative method. *Trends Ecol Evol* 2010; **25**: 713–721.

51   Bornbusch SL, Harris RL, Grebe NM, Dimac-Stohl K, Drea CM. Longitudinal effects of

antibiotics and fecal transplant on lemur gut microbiota structure, associations, and

resistomes. doi:https://doi.org/10.1101/2020.11.11.378349;

52   Bennett G, Malone M, Sauther ML, Cuozzo FP, White B, Nelson KE, Stumpf RM, Knight

R, Leigh SR, Amato KR. Host age, social group, and habitat type influence the gut

microbiota of wild ring-tailed lemurs (Lemur catta). *Am J Primatol* 2016; **78**: 883–892.

53   Fogel AT. The gut microbiome of wild lemurs: a comparison of sympatric Lemur catta

and Propithecus verreauxi. *Folia Primatol* 2015; **86**: 85–95.

54   Dias PC. Sources and sinks in population biology. *Trends Ecol Evol* 1996; **11**: 326–330.

55   Leclaire S, Nielsen JF, Drea CM. Bacterial communities in meerkat anal scent secretions

vary with host sex, age, and group membership. *Behav Ecol* 2014; **25**: 996–1004.

56   Theis KR, Schmidt TM, Holekamp KE. Evidence for a bacterial mechanism for group-

specific social odors among hyenas. *Sci Rep* 2012; **2**: 615.

57   Vernier CL, Chin IM, Adu-Oppong B, Krupp JJ, Levine J, Dantas G, Ben-Shahar Y. The

gut microbiome defines social group membership in honey bee colonies. *Sci Adv* 2020; **6**:

eabd3431.

58   Sarkar A, Harty S, Johnson KV-A, Moeller AH, Archie EA, Schell LD, Carmody RN,

Clutton-Brock TH, Dunbar RIM, Burnet PWJ. Microbial transmission in animal social

networks and the social microbiome. *Nat Ecol Evol* 2020; **4**: 1020–1035.

59   Amato KR, Yeoman CJ, Kent A, Righini N, Carbonero F, Estrada A, Gaskins HR, Stumpf

RM, Yildirim S, Torralba M. Habitat degradation impacts black howler monkey (Alouatta

pigra) gastrointestinal microbiomes. *ISME J* 2013; **7**: 1344.

60   Barelli C, Albanese D, Donati C, Pindo M, Dallago C, Rovero F, Cavalieri D, Tuohy KM,

Hauffe HC, De Filippo C. Habitat fragmentation is associated to gut microbiota diversity

of an endangered primate: implications for conservation. *Sci Rep* 2015; **5**: 14862.

61   Barelli C, Albanese D, Stumpf RM, Asangba A, Donati C, Rovero F, Hauffe HC. The gut

microbiota communities of wild arboreal and ground-feeding tropical primates are

affected differently by habitat disturbance. *Msystems* 2020; **5**.

62   Trosvik P, Rueness EK, de Muinck EJ, Moges A, Mekonnen A. Ecological plasticity in

the gastrointestinal microbiomes of Ethiopian Chlorocebus monkeys. *Sci Rep* 2018; **8**: 1–

10.

63   Kohl KD, Skopec MM, Dearing MD. Captivity results in disparate loss of gut microbial

diversity in closely related hosts. *Conserv Physiol* 2014; **2**: cou009.

64   Cameron A, Gould L. Fragment-Adaptive Behavioural Strategies and Intersite Variation

in the Ring-Tailed Lemur (Lemur catta) in South-Central Madagascar. In: Marsh LK,

Chapman CA (eds). *Primates in Fragments SE - 16*. Springer New York, 2013, pp 227–

243.

65   Chi X, Gao H, Wu G, Qin W, Song P, Wang L, Chen J, Cai Z, Zhang T. Comparison of

gut microbiota diversity between wild and captive bharals (Pseudois nayaur). *BMC Vet*

Res 2019; **15**: 1–8.

66   Hale VL, Tan CL, Niu K, Yang Y, Zhang Q, Knight R, Amato KR. Gut microbiota in

wild and captive Guizhou snub-nosed monkeys, Rhinopithecus brelichi. *Am J Primatol*

2019; **81**: e22989.

67   Clayton JB, Vangay P, Huang H, Ward T, Hillmann BM, Al-Ghalith GA, Travis DA,

Long HT, Van Tuan B, Van Minh V. Captivity humanizes the primate microbiome. *Proc*

Natl Acad Sci 2016; : 201521835.

68   Clayton JB, Al-Ghalith GA, Long HT, Van Tuan B, Cabana F, Huang H, Vangay P, Ward

T, Van Minh V, Tam NA. Associations between nutrition, gut microbiome, and health in a

novel nonhuman primate model. *Sci Rep* 2018; **8**.

69   Nelson TM, Rogers TL, Carlini AR, Brown M V. Diet and phylogeny shape the gut

microbiota of Antarctic seals: a comparison of wild and captive animals. *Environ*

Microbiol 2013; **15**: 1132–1145.

70   Greene L, Blanco MB, Rambeloson E, Graubics K, Fanelli B, Colwell RR, Drea CM. Gut

microbiota of frugo-folivorous sifakas across environments. *Anim Microbiome* 2021.

71   Tsukayama P, Boolchandani M, Patel S, Pehrsson EC, Gibson MK, Chiou KL, Jolly CJ,

Rogers J, Phillips-Conroy JE, Dantas G. Characterization of wild and captive baboon gut

microbiota and their antibiotic resistomes. *Msystems* 2018; **3**.

72   Narat V, Amato KR, Ranger N, Salmona M, Mercier-Delarue S, Rupp S, Ambata P,

Njouom R, Simon F, Giles-Vernick T. A multi-disciplinary comparison of great ape gut

microbiota in a central African forest and European zoo. *Sci Rep* 2020; **10**: 1–15.

73   Xiao Y, Xiao G, Liu H, Zhao X, Sun C, Tan X, Sun K, Liu S, Feng J. Captivity causes

taxonomic and functional convergence of gut microbial communities in bats. *PeerJ* 2019;

7: e6844.

74   Fujimura KE, Slusher NA, Cabana MD, Lynch S V. Role of the gut microbiota in defining

human health. *Expert Rev Anti Infect Ther* 2010; **8**: 435–454.

75   Clayton JB, Gomez A, Amato K, Knights D, Travis DA, Blekhman R, Knight R, Leigh S,

Stumpf R, Wolf T. The gut microbiome of nonhuman primates: Lessons in ecology and

evolution. *Am J Primatol* 2018; : e22867.

76   Cheng Y, Fox S, Pemberton D, Hogg C, Papenfuss AT, Belov K. The Tasmanian devil

microbiome—implications for conservation and management. *Microbiome* 2015; **3**: 1–11.

77   Borbón-García A, Reyes A, Vives-Flórez M, Caballero S. Captivity shapes the gut

microbiota of Andean bears: insights into health surveillance. *Front Microbiol* 2017; **8**:

1316.

78   Ma T, Villot C, Renaud D, Skidmore A, Chevaux E, Steele M. Linking perturbations to

temporal changes in diversity, stability, and compositions of neonatal calf gut microbiota:

prediction of diarrhea. *ISME J* 2020; : 1–13.

79   Chomel BB, Belotto A, Meslin F-X. Wildlife, exotic pets, and emerging zoonoses. *Emerg*

Infect Dis 2007; **13**: 6.

80   LaFleur M, Reuter KE, Hall MB, Rasoanaivo HH, McKernan S, Ranaivomanana P,

Michel A, Rabodoarivelo MS, Iqbal Z, Rakotosamimanana N. Drug-Resistant

Tuberculosis in Pet Ring-Tailed Lemur, Madagascar. *Emerg Infect Dis* 2021; **27**: 977.

81   LaFleur M, Clarke TA, Reuter KE, Schaefer MS. Illegal Trade of Wild-Captured Lemur

catta within Madagascar. *Folia Primatol* 2019; **90**: 199–214.

Tong Q, Cui L-Y, Du X-P, Hu Z-F, Bie J, Xiao J-H, Wang H-B, Zhang J-T. Comparison of gut microbiota diversity and predicted functions between healthy and diseased captive Rana dybowskii. *Front Microbiol* 2020; **11**: 2096.

Watson SE, Hauffe HC, Bull MJ, Atwood TC, McKinney MA, Pindo M, Perkins SE. Global change-driven use of onshore habitat impacts polar bear faecal microbiota. *ISME J* 2019; **13**: 2916–2926.

Kaakoush NO. Insights into the role of Erysipelotrichaceae in the human host. *Front Cell Infect Microbiol* 2015; **5**: 84.

Campbell TP, Sun X, Patel VH, Sanz C, Morgan D, Dantas G. The microbiome and resistome of chimpanzees, gorillas, and humans across host lifestyle and geography. *ISME J* 2020; **14**: 1584–1599.

Nishida AH, Ochman H. A great-ape view of the gut microbiome. *Nat Rev Genet* 2019; **20**: 195–206.

Greene LK, Williams C V, Junge RE, Mahefarisoa KL, Rajaonarivelo T, Rakotondrainibe H, O'Connell TM, Drea CM. A role for gut microbiota in host niche differentiation. *ISME J* 2020; : 1–13.

Sun Y, Sun Y, Shi Z, Liu Z, Zhao C, Lu T, Gao H, Zhu F, Chen R, Zhang J. Gut Microbiota of Wild and Captive Alpine Musk Deer (Moschus chrysogaster). *Front Microbiol* 2020; **10**: 3156.

Liu J, Pu Y-Y, Xie Q, Wang J-K, Liu J-X. Pectin induces an in vitro rumen microbial population shift attributed to the pectinolytic Treponema group. *Curr Microbiol* 2015; **70**: 67–74.

90 Liu J, Wang J-K, Zhu W, Pu Y-Y, Guan L-L, Liu J-X. Monitoring the rumen pectinolytic

bacteria Treponema saccharophilum using real-time PCR. *FEMS Microbiol Ecol* 2014;

87: 576–585.

91 Dishman DL, Thomson DM, Karnovsky NJ. Does simple feeding enrichment raise

activity levels of captive ring-tailed lemurs (Lemur catta)? *Appl Anim Behav Sci* 2009;

116: 88–95.

92 Mowry CB, Campbell JL, Mowry CB, Campbell JL. AZA Nutrition Advisory Group

TAG/SSP Husbandry Notebook Nutrition Section Lemur catta (Ring-tailed lemur). 2001.

93 Zaneveld JR, McMinds R, Thurber RV. Stress and stability: applying the Anna Karenina

principle to animal microbiomes. *Nat Microbiol* 2017; **2**: 1–8.

94 Ahmed HI, Herrera M, Liew YJ, Aranda M. Long-term temperature stress in the coral

model Aiptasia supports the "Anna Karenina principle" for bacterial microbiomes. *Front

Microbiol* 2019; **10**: 975.

95 Kohl KD, Dearing MD. Wild-caught rodents retain a majority of their natural gut

microbiota upon entrance into captivity. *Environ Microbiol Rep* 2014; **6**.

doi:10.1111/1758-2229.12118.

96 Martínez-Mota R, Kohl KD, Orr TJ, Dearing MD. Natural diets promote retention of the

native gut microbiota in captive rodents. *ISME J* 2020; **14**: 67–78.

97 Grieneisen LE, Charpentier MJE, Alberts SC, Blekhman R, Bradburd G, Tung J, Archie

EA. Genes, geology and germs: gut microbiota across a primate hybrid zone are explained

by site soil properties, not host species. *Proc R Soc B* 2019; **286**: 20190431.

98 Goodman SM, Benstead JP. Updated estimates of biotic diversity and endemism for

Madagascar. *Oryx* 2005; **39**: 73–77.

99   Ganzhorn JU, Lowry PP, Schatz GE, Sommer S. The biodiversity of Madagascar: one of

791   the world's hottest hotspots on its way out. *Oryx* 2001; **35**: 346–348.

792   100   Dietrich M, Wilkinson DA, Soarimalala V, Goodman SM, Dellagi K, Tortosa P.

793   Diversification of an emerging pathogen in a biodiversity hotspot: L eptospira in endemic

794   small mammals of M adagascar. *Mol Ecol* 2014; **23**: 2783–2796.

795   101   Jeffries CL, Tantely LM, Raharimalala FN, Hurn E, Boyer S, Walker T. Diverse novel

796   resident Wolbachia strains in Culicine mosquitoes from Madagascar. *Sci Rep* 2018; **8**: 1–

797   15.

798   102   Larsen PA, Hayes CE, Williams C V, Junge RE, Razafindramanana J, Mass V,

799   Rakotondrainibe H, Yoder AD. Blood transcriptomes reveal novel parasitic zoonoses

800   circulating in Madagascar's lemurs. *Biol Lett* 2016; **12**: 20150829.

801   103   Guiyoule A, Rasoamanana B, Buchrieser C, Michel P, Chanteau S, Carniel E. Recent

802   emergence of new variants of Yersinia pestis in Madagascar. *J Clin Microbiol* 1997; **35**:

803   2826–2833.

804   104   Bahram M, Hildebrand F, Forslund SK, Anderson JL, Soudzilovskaia NA, Bodegom PM,

805   Bengtsson-Palme J, Anslan S, Coelho LP, Harend H. Structure and function of the global

806   topsoil microbiome. *Nature* 2018; **560**: 233–237.

807   105   Reed KE, Fleagle JG. Geographic and climatic control of primate diversity. *Proc Natl*

808   *Acad Sci* 1995; **92**: 7874–7876.

809   106   Mittermeier RA. Primate diversity and the tropical forest. *Biodiversity* 1988.

810   107   Melfi V. The appliance of science to zoo-housed primates. *Appl Anim Behav Sci* 2005; **90**:

811   97–106.

812   108   Primates M, Altschul DM, Beran MJ, Bohn M, Call J, DeTroy S, Duguid SJ, Egelkamp

813    CL, Fichtel C, Fischer J. Establishing an infrastructure for collaboration in primate

814    cognition research. *PLoS One* 2019; **14**: e0223675.

815 109 Johns T, Duquette M. Detoxification and mineral supplementation as functions of

816    geophagy. *Am J Clin Nutr* 1991; **53**: 448–456.

817 110 Krishnamani R, Mahaney WC. Geophagy among primates: adaptive significance and

818    ecological consequences. *Anim Behav* 2000; **59**: 899–915.

819 111 Won Y-J, Hallam SJ, O'Mullan GD, Pan IL, Buck KR, Vrijenhoek RC. Environmental

820    acquisition of thiotrophic endosymbionts by deep-sea mussels of the genus

821    Bathymodiolus. *Appl Environ Microbiol* 2003; **69**: 6785–6792.

822 112 Tout J, Astudillo-García C, Taylor MW, Tyson GW, Stocker R, Ralph PJ, Seymour JR,

823    Webster NS. Redefining the sponge-symbiont acquisition paradigm: sponge microbes

824    exhibit chemotaxis towards host-derived compounds. *Environ Microbiol Rep* 2017; **9**:

825    750–755.

826 113 Caravaggi A, Plowman A, Wright DJ, Bishop CM. The composition of captive ruffed

827    lemur (Varecia spp.) diets in UK zoological collections, with reference to the problems of

828    obesity and iron storage disease. *J Zoo Aquarium Res* 2018; **6**: 41–49.

829 114 McPherson FJ. Normal blood parameters, common diseases and parasites affecting

830    captive non-human primates. *J Primatol* 2013; **2**: e112.

831 115 McKenney EA, Greene LK, Drea CM, Yoder AD. Down for the count: Cryptosporidium

832    infection depletes the gut microbiome in Coquerel's sifakas. *Microb Ecol Health Dis*

833    2017; **28**: 1335165.

834 116 Schwitzer C, Mittermeier RA, Johnson SE, Donati G, Irwin M, Peacock H, Ratsimbazafy

835    J, Razafindramanana J, Louis EE, Chikhi L. Averting lemur extinctions amid

Madagascar's political crisis. *Science (80-)* 2014; **343**: 842–843.

117 Trevelline BK, Fontaine SS, Hartup BK, Kohl KD. Conservation biology needs a

microbial renaissance: a call for the consideration of host-associated microbiota in wildlife

management practices. *Proc R Soc B* 2019; **286**: 20182448.

118 Tenhumberg B, Tyre AJ, Shea K, Possingham HP. Linking wild and captive populations

to maximize species persistence: optimal translocation strategies. *Conserv Biol* 2004; **18**:

1304–1314.

119 Mills JG, Weinstein P, Gellie NJC, Weyrich LS, Lowe AJ, Breed MF. Urban habitat

restoration provides a human health benefit through microbiome rewilding: the

Microbiome Rewilding Hypothesis. *Restor Ecol* 2017; **25**: 866–872.

120 Song SJ, Amir A, Metcalf JL, Amato KR, Xu ZZ, Humphrey G, Knight R. Preservation

methods differ in fecal microbiome stability, affecting suitability for field studies.

MSystems 2016; **1**.

121 Choo JM, Leong LE, Rogers GB. Sample storage conditions significantly influence faecal

microbiome profiles. *Sci Rep* 2015; **5**: 16350.

122 Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, Owens SM,

Betley J, Fraser L, Bauer M. Ultra-high-throughput microbial community analysis on the

Illumina HiSeq and MiSeq platforms. *ISME J* 2012; **6**: 1621.

123 Bornbusch SL, Grebe NM, Lunn S, Southworth CA, Dimac-Stohl K, Drea C. Stable and

transient structural variation in lemur vaginal, labial and axillary microbiomes: patterns by

species, body site, ovarian hormones and forest access. *FEMS Microbiol Ecol* 2020; **96**:

fiaa090.

124 Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO. The

SILVA ribosomal RNA gene database project: improved data processing and web-based

tools. *Nucleic Acids Res* 2012; **41**: D590–D596.

125   Yarza P, Yilmaz P, Pruesse E, Glöckner FO, Ludwig W, Schleifer K-H, Whitman WB,

Euzéby J, Amann R, Rosselló-Móra R. Uniting the classification of cultured and

uncultured bacteria and archaea using 16S rRNA gene sequences. *Nat Rev Microbiol*

2014; **12**: 635.

126   Breiman L. Random forests. *Mach Learn* 2001; **45**: 5–32.

127   Morton JT, Marotz C, Washburne A, Silverman J, Zaramela LS, Edlund A, Zengler K,

Knight R. Establishing microbial composition measurement standards with reference

frames. *Nat Commun* 2019; **10**: 1–11.

128   Fedarko MW, Martino C, Morton JT, González A, Rahman G, Marotz CA, Minich JJ,

Allen EE, Knight R. Visualizing'omic feature rankings and log-ratios using Qurro. *NAR*

genomics Bioinforma 2020; **2**: lqaa023.

129   Shenhav L, Thompson M, Joseph TA, Briscoe L, Furman O, Bogumil D, Mizrahi I, Pe'er

I, Halperin E. FEAST: fast expectation-maximization for microbial source tracking. *Nat*

Methods 2019; **16**: 627.

Table 1. Research settings (names, descriptions, and locations) and samples collected under wilderness conditions and under captivity conditions in Madagascar and the U.S. A subset of the samples collected were omitted from analyses owing to low-yield extractions or low-quality sequencing. Soil samples could not always be obtained. Maps show locations of each setting; the gray shaded area shows the natural range of wild ring-tailed lemurs in Madagascar.

Condition	Setting (abbreviation)	Setting description	Samples: collected (analyzed) fecal	soil
wilderness	Amoron'I Onilahy (AMO)	Riverine gallery forest, dry scrub forest	20	4 (3)
	Berenty Reserve (BER)	Semi-arid dry forest, spiny forest	22 (19)	4
	Beza Mahafaly Special Reserve (BEZ)	Riverine gallery and semi-arid spiny forest	26	4 (3)
	Fiheranana (FIH)	Dry forest and spiny forest	2	-
	Isalo National Park (ISO)	Dry deciduous forest	18 (16)	3
	Ivohiboro (IVO)	Humid forest, grassland	15	-
	Ranomay (RAN)	Dry forest	13	2 (1)
	Tsimanampetsotsa National Park (TSI)	Dry forest and spiny forest	29 (28)	8
captivity: Madagascar	Lemur Rescue Center (Toliara, Madagascar; LRC)	Outdoor enclosures in dry and spiny forest	33	2 (1)
	Various towns (pets)	Pets housed in human dwellings	8	-
captivity: U.S.	Duke Lemur Center (Durham, NC; DLC)	Indoor and outdoor enclosures, including free ranging in semi-deciduous forest	22	3 (2)
	National Zoological Park (Washington, DC; NZP)	Indoor and outdoor enclosures, moated island with vegetation	4	-
	North Carolina Zoo (Asheboro, NC; NCZ)	Indoor and outdoor enclosures, moated island with vegetation	3	-
		Total samples:	215 (209)	30 (25)

Figure and legends

Figure 1. Alpha diversity metrics of lemur gut microbiota (a, b) collapsed by environmental
condition, including wilderness (wild lemurs; green), captivity in Madagascar (captive$_M$ lemurs;
pink), and captivity in the U.S. (captive$_{U.S.}$ lemurs; orange), and (c, d) averaged across
individuals for each of the 13 different settings inhabited (reprising the color codes of each
condition, delineated by dashed vertical lines). Shown are both (a, c) Shannon diversity and (b,
d) Faith's phylogenetic diversity. Across the (c, d) settings within a condition (see Table 1 for
names of abbreviated study sites), the data are plotted in descending order of mean Shannon
diversity.

908
909
910 Figure 2. Mean proportion of sequences assigned to microbial taxa across lemurs at each of the 13 different settings, with the three
911 conditions (wilderness, captivity in Madagascar, and captivity in the U.S.) delineated by dashed vertical lines (see Table 1 for names
912 of abbreviated study sites). Taxa are identified by phylum and deepest possible taxonomic level (i.e., genus level or above); those
913 representing < 1% of the microbiomes were combined into the category "Other"

914
915
Figure 3. Differential abundance of *Erysipelotrichaceae UCG-004* and *Treponema* 2 in the gut
microbiota of lemurs. (a) Differential rank plot showing lemur gut microbial taxa (x axis) ranked
by their differentials (y axis; the estimated log-fold changes for taxa abundances across sample
groups) for lemurs in the wilderness vs. captivity settings in the U.S. Those taxa that are more
abundant in the wild lemurs compared to captive lemurs in the U.S. appear on the right side of
the plot whereas those that are less abundant in wild lemurs appear on the left side.
Erysipelotrichaceae UCG-004 and *Treponema 2* differentials are highlighted in red and blue,
respectively, with other taxa represented in gray. (b) Natural log ratios of *Erysipelotrichaceae*
UCG-004 vs. *Treponema 2* in lemurs across all three environmental conditions.
925
926

Figure 4. Beta diversity (unweighted UniFrac distances) of lemur gut microbiota across three environmental conditions – wilderness (wild lemurs; green), captivity in Madagascar (captive$_M$ lemurs; pink), and captivity in the U.S. (captive$_{U.S.}$ lemurs; orange) – that encompass 13 setting (see Table 1 for names of abbreviated study sites). (a, b) Principal coordinate plots, showing axes 1 and 2, or 1 and 3, respectively, of individual gut microbial communities colored by setting and encircled by normal data ellipses reflecting environmental condition. (c) Mean beta diversity distance scores within (single color) and between (two colors) environmental conditions. (d) Principal coordinate plots, showing axes 1 and 2, for the eight settings within the wilderness condition. Kruskal-Wallis test with Benjamini-Hochberg correction; **** $p < 0.0001$.

Figure 5. Alpha diversity metrics of soil microbiota (a, b) collapsed by environmental condition, including wilderness (wilderness soil; green), captivity in Madagascar (captivity$_M$ soil; pink), and captivity in the U.S. (captivity$_{U.S.}$ soil; orange) and (c, d) averaged across individuals for each of the eight different settings (reprising the color codes of each condition, delineated by dashed vertical lines). Shown are both (a, c) Shannon diversity and (b, d) Faith's phylogenetic diversity. Across the (c, d) settings within a condition (see Table 1 for names of abbreviated study sites), the data are plotted in descending order of mean Shannon diversity.

953
954 Figure 6. Mean proportion of sequences assigned to microbial taxa of soil at each of the eight settings sampled, within the three
955 conditions: wilderness (wilderness soil; green), captivity in Madagascar (captivity_M soil; pink), and captivity in the U.S. (captivity_{U.S.}
956 soil; orange), which are delineated by dashed vertical lines (see Table 1 for names of abbreviated study sites). Taxa are identified by
957 phylum and deepest possible taxonomic level (i.e., genus level or above); those representing < 1% of the microbiomes were combined
958 into the category "Other".

960
961
Figure 7. Beta diversity (unweighted UniFrac distances) of soil microbiota across three environmental conditions - wilderness (wilderness soil; green), captivity in Madagascar (captivity$_M$ soil; pink), and captivity in the U.S. (captivity$_{U.S.}$ soil; orange) – that encompass eight setting (see Table 1 for names of abbreviated study sites). (a, b) Principal coordinate plots, showing axes 1 and 2, or 1 and 3, respectively, of soil microbial communities colored by setting and encircled by normal data ellipses reflecting environmental condition. (c) Mean beta diversity distance scores within (single color) and between (two colors) environmental conditions. Kruskal-Wallis test with Benjamini-Hochberg correction; * $p < 0.05$, **** $p < 0.0001$.

970
971

972
973
Figure 8. Mean proportion of total soil-associated microbes in the gut microbiota of lemurs (a)
collapsed by environmental condition: wilderness (wild lemurs; green), captivity in Madagascar
(captive$_M$ lemurs; pink), and captivity in the U.S. (captive$_{U.S.}$ lemurs; orange). Averaged soil-
associated microbes across individuals for (b) each of the eight different settings (reprising the
color codes of each condition, delineated by dashed vertical lines) and (c) at the Duke Lemur
Center (DLC) that were semi-free-ranging in natural habitat enclosures or were housed indoors.
Kruskal-Wallis test with pairwise comparisons and Benjamini-Hochberg correction; * $p < 0.05$,
** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$, ns = nonsignificant.

Figure 9. Mean proportion of soil-associated microbes in the gut microbiota of lemurs within (single color) and between (two colors) the three conditions: wilderness (wild lemurs; green), captivity in Madagascar (captive$_M$ lemurs; pink), and captivity in the U.S. (captive$_{U.S.}$ lemurs; orange) – that encompass eight setting. Within the gut microbiota of lemurs from a given condition (first color = fecal source condition), values show the proportion of soil associated microbes from a given condition (second color = soil source condition). Kruskal-Wallis test with pairwise comparisons and Benjamini-Hochberg correction; ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.