Anoikis resistance in mammary epithelial cells is mediated by semaphorin 7a

Semaphorin-7A and anoikis resistance

Taylor R. Rutherford[1,23] Alan M Elder[1,2,4] Traci R. Lyons[1,2,4,5]

Corresponding author: traci.lyons@cuanscutz.edu

[1] Division of Medical Oncology, Department of Medicine, School of Medicine, University of Colorado Anschutz Medical Campus, Aurora, Colorado.

[2] Young Women's Breast Cancer Translational Program, University of Colorado Anschutz Medical Campus, Aurora, Colorado.

[3] Cell biology, Stem cell, and Development Graduate Training Program, University of Colorado Anschutz Medical Campus, Aurora, Colorado.

[4] Cancer biology Graduate Training Program, University of Colorado Anschutz Medical Campus, Aurora, Colorado.

[5] University of Colorado Cancer Center, Aurora, Colorado.

ABSTRACT

Semaphorin-7a (SEMA7A), best known as a neuroimmune molecule, plays a diverse role in many cellular processes and pathologies. Here, we show that SEMA7A promotes anoikis resistance in cultured mammary epithelial cells through integrins and activation of pro-survival kinase AKT, which led us to investigate a role for SEMA7A during postpartum mammary gland involution—a normal developmental process where cells die by anoikis. Our results reveal that SEMA7A is expressed on live mammary epithelial cells during involution, that SEMA7A expression is primarily observed in $\alpha6$-integrin expressing cells, and that luminal progenitor cells, specifically, are decreased in mammary glands of SEMA7A-/- mice during involution. We further identify a SEMA7A-$\alpha6/\beta1$-integrin dependent mechanism of mammosphere formation and chemoresistance in mammary epithelial cells and suggest that this mechanism is relevant for recurrence in breast cancer patients.

INTRODUCTION

It has been postulated that the functional plasticity of the adult mammary epithelium, which is required for response to hormonal changes during pregnancy and lactation, may make the breast tissue more susceptible to malignancy.Identifying mechanisms of normal mammary epithelial cell (MEC) biology that contribute to tumorigenesis is critically important for furthering our understanding of breast cancer and extending the survival of breast cancer patients. Of particular interest are postpartum breast cancers (PPBCs), defined as breast cancers diagnosed in women within 10 years of most recent childbirth, as these cancers are nearly three times more likely to result in lethal metastasis(1).

The majority of mammary morphogenesis occurs postnatally beginning at puberty and culminating with lactation following pregnancy. As such, differentiation of the mammary

epithelium is not complete until a full pregnancy-lactation cycle. Upon weaning, or after

pregnancy in the absence of nursing, the mammary gland regresses to resemble the pre-

pregnant architecture through a process termed postpartum involution. In mice, postpartum

involution occurs in two primary phases(2). In the first phase, milk proteins accumulate in the

mammary ducts and alveoli resulting in downstream phosphorylation and activation of

signal transducer and activator of transcription (pSTAT3), and subsequent lysosomal-mediated

cell death(3). In mice, if pups are returned within 48 hours of weaning, this process can be

reversed, and lactation can continue. If pups are not returned, however, the irreversible second

phase of involution begins. This phase is characterized by degradation of the mammary

basement membrane by matrix metalloproteinases and anoikis-mediated cell death in the

absence of laminin-mediated activation of integrin alpha-6/beta-1 ($\alpha6\beta1$) and downstream pro-

survival signaling(4-6).

SEMA7A, or CD108w, is a member of the semaphorin protein family, named after the English

verb "semaphore", which means to transmit a signal across a long distance. SEMA7A was first

identified for its expression on lymphocytes(7), but has since been shown to exert diverse roles

in neuronal development(8), cell differentiation(9-11), ECM deposition(12, 13), various

components of the immune system(14-17), and pathologies including fibrosis(12, 18, 19) and

cancer,(13, 16, 20-22). Though elevated SEMA7A is observed in breast cancer, where we have

established a role for it in conferring cell survival, in part, via promotion of anoikis resistance(13,

23), the role of SEMA7A in normal MECs has not previously been investigated. SEMA7A is

known to signal through $\beta1$-integrin in immune cells, neurons, and cancer; $\beta1$-integrin also

mediates MEC survival during development, in part, via phosphorylation and activation of

downstream pro-survival signaling mediated by PI3-kinase and protein kinase B, or Akt1(24).

Though it is known that redundant secretory epithelial cells die during postpartum involution, the

factor(s) that allow some cells to survive and regenerate a lactation-competent gland for future

pregnancies remain unknown. Herein, we have identified enrichment for SEMA7A expressing

$\alpha6\beta1$ positive MECs during the matrix remodeling period of postpartum involution and a role for

SEMA7A in specifically promoting the survival of $\alpha6$-integrin+ luminal progenitor cells. We

provide evidence that exogenous stimulation of immortalized MECs with purified SEMA7A

protein promotes anoikis resistance via activation of $\alpha6$ and $\beta1$-integrins and activation of

phosphorylated AKT (pAKT). Consistent with a role for SEMA7A in promoting anoikis

resistance, SEMA7A knockout mice (SEMA7A[tmalk1/J]) exhibit phenotypes associated with

accelerated involution including early presence of apoptotic cell bodies in mammary lumens,

increased pSTAT3, and premature reappearance of adipocytes. $\alpha6$-integrin also drives

maintenance of tissue resident stem cells and therapy resistant cancer stem cells(25) and our

findings reveal that SEMA7A drives stem cell phenotypes and chemoresistance in cultured

MECs in a manner that is dependent on $\alpha6$ and $\beta1$. Finally, we provide evidence for SEMA7A in

driving transformed phenotypes in normal MECs and reversing transformed phenotypes in

genetically identical, ras-transformed, variants of the same cell line. Our results are the first to

identify a role for SEMA7A in anoikis resistance and promotion of stem and cancer cell

phenotypes via activation of $\alpha6$ and $\beta1$-integrins.

METHODS

Cell culture:

MCF10A/MCF12A cells were obtained from, and cultured, according to ATCC.

MCF10DCIS.com cells were obtained from K. Polyak (University of Harvard, Cambridge, MA).

Cells were validated by the DNA sequencing core at the University of Colorado Anschutz

Medical Campus and found to be pure populations of their respective cell lines. Cells were

regularly tested for mycoplasma throughout studies. Cells were sub-cultured as previously

described (reference), or according to ATCC standards. Exogenous SEMA7A protein utilized in

our studies was purified from media of cells stably transfected with an Fc-tagged version of

SEMA7A by published methods(14, 26). Control plasmid (pcDNA3.1) was obtained from H.

Ford (University of Colorado Anschutz Medical Campus, Denver, CO).

shRNA Knockdown and Overexpression

shRNA plasmids targeting Sema7a, and a negative control shRNA (SABiosciences), were

amplified in E. coli and plasmid DNA was isolated by Plasmid Maxi-Prep (Qiagen).

MCF10DCIS.com-GFP cells were cultured overnight to approximately 80% confluence. 1 μg/μL

of each shRNA was added to Transfectagro Reduced-Serum Medium (Corning) and incubated

for 15 minutes with 4 μL of X-treme Gene HP DNA transfection reagent (Roche). Transfected

cells were selected for by hygromycin resistance. Stable knockdown was confirmed by qPCR as

shown in Black et al(27). Negative control cells were transfected with a scrambled artificial

sequence not matching human, mouse, or rat. Overexpression plasmid (SEMA7A-Fc) was a

generous gift from R. Medzhitov (Yale University, New Haven, CT). All other over-expression

plasmids (p304-V5-Blasticidin and V5-SEMA7A)were obtained from the Functional Genomics

Core at the CUAnschutz Medical Campus and overexpression was confirmed via qPCR and

western blot analysis. KD (27) and overexpression (13) as previously described.

Animal Studies:

All animal procedures were approved by the University of Colorado Anschutz Medical Campus

Institutional Animal Care and Use Committee. Sema7a[tm1Alk/J] mice (a generous gift from Alex

Kolodkin at John's Hopkins University) and C57Bl/6 (Jax) were housed and bred as previously

described.(28) Briefly, C57Bl/6 SEMA7A[tm1Alk] females were bred with WT C57Bl/6 males to

induce pregnancy, lactation, and postpartum involution. Pup numbers were normalized to 6-8

per dam after birth to ensure for adequate lactation. Postpartum involution was initiated by

forced weaning of pups at the peak of lactation (L10-L12). #4 right and left mammary glands

were harvested from age-matched nulliparous animals and postpartum dams at involution days

1,2,4,6,8,10,14, and 28. For flow cytometry, inguinal lymph nodes were removed prior to tissue

digestion. #3 mammary glands were also harvested, formalin fixed, paraffin embedded, and 5

micron sections generated for histologic/immunohistochemical analaysis. See Supplemental

Table 1 for animal study sample sizes.

In vitro survival assays:

MCF10A and MCF12A cells were cultured with 200ng/ml purified SEMA7A or PBS vehicle

control in adherent (tissue culture plastic) or forced suspension conditions (ultra-low attachment

plates (Corning, Corning, NY, USA: #3473)). Cells were seeded at a density of 1000 cells/well in

a 96-well plate plus media only wells as controls. Cell death was measured 24 hours post

seeding via luminescence using the Caspase Glo assay (Promega, Madison, WI, USA:

#G8090). Annexin-V /7AAD staining (Biolegend, San Diego, CA, USA: #640930) and

pAKTS473 (ThermoFisher: Waltham, MA, USA, #17-9715-42) were also used to confirm cell

viability by flow cytometry. See flow cytometry methods for staining protocol. Function blocking

antibodies for b1 (CD29- 9EG7; BD Biosciences: #550531) and a6 integrin (ThermoFisher: #14-

0495-82) were used to disrupt integrin signaling in the presence of SEMA7A or as antibody

alone controls to determine off-target effects. IgG controls were also used. 9EG7 was used at a

concentration of 0.6ug/ml while GoH3 was used at a concentration of 100ug/mL for inhibition

studies consistent with previous reports (29, 30). Cells were cultured with anti-integrin inhibitors

at time of seeding.

Label retention stem cell assay:

MCF10A and MCF12A cells were labeled with CellTrace Violet (ThermoFisher: #C34557) for 20

minutes at 37°C according to manufacturer's instructions. A sample of labeled cells was

analyzed by flow cytometry to determine labeling efficiency and fluorescent intensity at time of

seeding **(Supplemental Figure 1)** Single cells were seeded at a density of 4000 cells/well in

ultra-low attachment 24-well plates to induce mammosphere formation. Cells were cultured in

normal culture media for 7 days and were subsequently counted, dissociated enzymatically in

Accutase, (Stem Cell Technologies, Vancouver, Canada: #07922), and analyzed by flow

cytometry. See flow cytometry methods for detailed information.

Paclitaxel resistance studies: MCF10DCIS.com cells were treated with 100nM paclitaxel

(Millipore Sigma, Burling, MA, USA #T7402-5MG) for 24 hours prior to analysis by flow

cytometry. Dose was determined experimentally based on cell viability and cancer stem cell

enrichment. Drug resistance was measured by label-retention and mammosphere formation as

described above and %CD44+ (Biolegend: #103018) CD24- (Biolegend: #311104)

of singlets was used as a molecular definition of drug-resistant stem cells. See label retention

stem cell assay and flow cytometry methods for additional details.

Flow cytometry:

Single cell suspensions were generated from cultured cells via enzymatic harvest with

Accutase. Mouse mammary tissues were minced with scalpels, digested in Click's media

containing 500 units/mL Collagenase II (Worthington, Columbus, OH, USA; LS004174) and

Collagenase IV (Worthington, LS004186) and 20ug/mL DNAse (Worthington, LS002004) for 1

hour at 37°C with occasional trituration and strained through 70um filters to generate single cell

suspensions. Single cells were stained at 4°C for 30 minutes and washed with PBS+2%FBS.

Cells were filtered through 30um filters, diluted to $<10^6$ cells/ml, and analyzed with the ZE5-YETI

flow cytometer. Fixation/Permeabilization kit (BD Biosciences, Franklin Lakes, NJ, USA:

#554715) was used for intracellular antigen staining. Data were analyzed with Kaluza software.

Single stain, FMO, and unstained controls were used to set gates. Single cells were lineage

depleted using CD45- (BD Biosciences: #564279)/CD31- (BD Biosciences: #745436) gating.

Full gating strategy is included as Supplemental Figure 2.

Immunohistochemistry:

Tissues were formalin-fixed and paraffin-embedded as previously described(31). Hematoxylin

and Eosin staining was used to define morphological features. Antigen retrieval was performed

using target retrieval solution (Dako, Glostrup, Denmark; #S1699) for CC3(Cell Signaling

Technologies, Danvers, MA, USA; #9661) and Perilipin(Cell Signaling: #3470) or (EDTA Dako

cat# S2367) for pSTAT3 (Cell Signaling Technologies; #9145). ImmPRESS polymer anti-rabbit

IgG secondary reagent (Vector Labs, San Francisco, CA, USA t#MP-7401-15) was used for

secondary staining of CC3 and pSTAT3 stained tissues and anti-rabbit secondary (Dako

#K4003) for perilipin stains. DAB (Vector Labs,; #SK-4105) and counterstaining were performed

with hematoxylin (Vector Labs #H-3401). Alveolar area was measured on H&E-stained tissues

in ImageJ on 10 alveoli/field; 5 fields/tissue. 10 fields/tissue were analyzed for positive CC3,

pSTAT3, and perilipin stained tissues using CellSense Dimension software count and measure

feature on regions of interest.

Data Mining:

Analysis of SEMA7A in breast cancer versus normal was performed on

http://ualcan.path.uab.edu/index.html by selecting breast cancer samples from the The Cancer

Genome Atlas (TCGA) (32, 33). 5-year relapse free survival curves for *Sema7a (*Affy ID:

230345_at) and *Itga6 (*Affy ID: 201656_at) were generated in KM plot using default

parameters and no restrictions. Patients split by median survival. For multi-gene

analyses, genes were filtered by median expression(34).

Statistical Analysis:

Unpaired *t*-test and one-way ANOVA were performed in GraphPad Prism. Nonparametric

equivalent analyses were used for samples with uneven distribution or unequal standard

deviations. *p* values of less than 0.05 were considered significant. Error bars represent mean +/-

standard error of the mean. All in vivo studies were replicated in at least 3 mice and in vitro

studies performed in technical and biological triplicate with representative data presented.

RESULTS

Semaphorin-7a promotes anoikis resistance in normal MECs

To identify whether SEMA7A promotes survival of normal MECs we utilized two methods:

transfection of FC-tagged SEMA7A into immortalized human (MCF10A) cells **(Supplemental**

Figure 3) and addition of exogenous SEMA7A, which we purified from FC-tagged SEMA7A

expressing cells, to wild-type MCF10A and MCF12A cells—another non-transformed MEC line.

We found that genetic SEMA7A overexpression resulted in resistance to cell death, as

measured by cleaved caspase 3/7, in a forced suspension assay, but not in attached conditions

suggesting that SEMA7A specifically mediates resistance to cell death by anoikis(**Figure 1A)**.

To determine the contribution of endogenous SEMA7A protein to anoikis resistance, we

measured SEMA7A+ MECs by flow over time in attached and suspended conditions and found

that SEMA7A+ MECs in attached conditions remain unchanged while SEMA7A+ cells become

significantly enriched as early as 24 hours in detached conditions(**Figure 1B**). We also show

that stimulation with exogenous purified SEMA7A protein in both MCF10A and MCF12A cells is

sufficient to significantly decrease cell death and Inhibition of a known SEMA7A receptor, b1-

integrin, significantly abrogated SEMA7A-mediated anoikis resistance **(Figure 1C and**

Supplemental Figures 4 and 5). We independently confirmed decreased cell death in

detached MECs cultured with exogenous SEMA7A by flow cytometry for Annexin-V/7AAD

(**Supplemental Figure 6**). We also examined pAKT in MECs cultured with SEMA7A using

insulin as a positive control as the IGF-1 receptor (and insulin) can activate pAKT via b1-integrin

in the mammary gland(35, 36). We observed percentages of pAkt expressing cells that were

similar to those observed with insulin when cells were exposed to SEMA7A *in vitro* **(Figure 1C)**.

To determine whether SEMA7A inhibits anoikis *in vivo*, we examined MECs harvested from

age-matched nulliparous and force weaned WT animals for SEMA7A expression. We observed

a low percentage of SEMA7A+EpCAM+ cells in the nulliparous mammary gland, which was

enriched approximately 3-fold at involution day 3 when basement membrane degradation and

extracellular matrix remodeling ensues. The percentage of SEMA7A+ epithelial cells decreased

starting at involution day 4 and returned to pre-involution day 3 levels by involution day 6

(Figure 1E). We also compared live epithelial cells by flow cytometry in age matched SEMA7A

knockout mice (Sema7a[tm1Alk/J]) and WT C57Bl/6 mice at days 1, 2, 3, & 6 post-wean and

observed a modest, but significant, reduction in MECs as measured by the percentage of

Aquazombie- EpCAM+ cells **(Figure 1F)**. To test whether the decrease in live cells during

involution in Sema7a[tm1Alk/J] animals was accompanied by changes to gland histology and

programs normally activated during postpartum involution, we performed H&E and

immunohistochemical analysis of whole mammary tissues isolated from Sema7a[tm1Alk/J] dams

and WT controls. We observed accelerated glandular regression, as evidenced by decreased

alveolar area, increased pSTAT3, precocious appearance of cleaved caspase 3 (CC3) positive

cells, and accelerated adipocyte reappearance in glands harvested from Sema7a[tm1Alk/J] animals

(Figure 2 and Figure 3) suggesting that loss of SEMA7A accelerates postpartum involution.

SEMA7A promotes anoikis-resistance via α6-integrin and activates AKT.

MECs can be divided into four primary cell types: luminal, luminal progenitor (LPCs), basal, and

basal progenitor cells (BPCs) and these populations can be resolved by flow cytometry based

largely on their expression of epithelial cell adhesion molecule (EpCAM) and CD49f, also known

as α6-integrin(37, 38). Since the magnitude of the changes observed in Figure 1E was relatively

small, we used flow cytometry to resolve the four primary MEC populations to determine which

cells are specifically affected by loss of SEMA7A during postpartum involution. We observed a

significant decrease in EpCAM^{hi}CD49f+ LPCs at all timepoints examined, which was not

consistently observed in the other cell populations **(Figure 4A,B and Supplemental Figure 7)**.

We also found that more than 80% of the live LPCs express SEMA7A at involution days 2-4

(Figure 4C). Furthermore, SEMA7A+ cells are enriched in each MEC population at some point

during postpartum involution except for mature luminal cells. **(Supplemental Figure 8)**. One

factor that distinguishes the SEMA7A+ MEC populations from the SEMA7A- luminal cells is

their expression of α6-integrin. β1-integrin is a known receptor that mediates SEMA7A cellular

signaling(8), but the preferred alpha binding partner in MECs has not been identified. We

quantified cells that co-express α6 and β1 in WT and SEMA7A$^{tmalk1/J}$ animals and found a

significant reduction during involution **(Figure 4D)**. Furthermore, we observed that over 90% of

the α6 and β1 positive MECs express SEMA7A at the transition to the reversible phase of

involution when basement membrane degradation and anoikis-induced cell death is initiated

(Figure 4E). To determine whether SEMA7A confers anoikis-resistance via α6-integrin

dependent activation of pro-survival signaling, we cultured MCF12A cells with SEMA7A in the

presence of the α6-integrin blocking antibody (GoH3) in forced suspension cultures. Consistent

with our previous results, SEMA7A decreased CC3/7 activity, which was blocked by GoH3.

SEMA7A treatment also resulted in enrichment for pAKT+ cells which was inhibited by α6 β1

blockade **(Figure 4F&G).** Inhibition of both α6 and β1 did not increase CC3/activity or alter

pAKT over vehicle alone in the absence of SEMA7A (**Figure 1C and 4G**).Taken together, our

results suggest SEMA7A promotes anoikis-resistance during involution, in part, through

α6β1and activation of pAKT.

SEMA7A promotes stem cell phenotypes, cellular transformation, and chemoresistance

Anoikis-resistance is rarely observed in normal epithelial cells; rather, it is a characteristic of

stem cells and tumor cells. Given our findings that SEMA7A promotes anoikis-resistance in

normal MECs, we tested whether it also promotes functional features of mammary stem cells

using cultured MECs. Previously identified functional properties of mammary stem cells include

a quiescent or slowly proliferative phenotype and their ability to survive and proliferate in

anchorage-independent conditions where they give rise to clonal spheroids. These spheroids,

also known as "mammospheres", allow for identification of stem cells as they only undergo one

or two rounds of division before reentering quiescence, allowing them to be identified based on

their ability to retain fluorescent dyes that are normally progressively lost by dilution in the

actively proliferating cells(39-41). We used a cell trace violet dye and mammosphere assay to

identify functional mammary stem cell phenotypes *in vitro* and found that SEMA7A expressing

MCF10A cells are enriched for a quiescent, or senescent, cell population **(Figure 5A)**. We also

measured mammosphere formation as a readout for self-renewal in cells cultured with

exogenous SEMA7A in the presence or absence of function blocking antibodies for α6 and β1

and found that increased sphere formation occurred with SEMA7A which was abrogated with

the inhibitors **(Figure 5B&C)**. Secondary sphere formation was also increased in MCF10A and

MCF12A cells cultured with SEMA7A **(Supplemental Figure 9).** We then tested the ability of

SEMA7A to promote cell growth in anchorage-independent conditions using a soft agar assay—

a classical measurement of cellular transformation. We found that overexpression of SEMA7A

in MCF10A cells is sufficient to increase growth in soft agar to levels similar to a Ras-

transformed variant of these cells, MCF10DCIS.com.

To determine a role for SEMA7A in breast cancer, we examined *SEMA7A* expression in

breast cancers in The Cancer Genome Atlas (TCGA) to show that *SEMA7A* is significantly

upregulated in primary tumors **(Figure 6A)**, is higher in all stages **(Figure 6B)**, is upregulated in

breast cancers in women of all ages **(Figure 6C)** and is expressed in every major subtype of

breast cancer with highest expression occurring in the triple-negative class when compared to

normal breast tissues **(Figure 6D and Supplemental Figure 10)**. We also utilied Kaplan-Meier

analysis to examine relapse free survival (RFS) to determine whether coordination between

SEMA7A and *α6-integrin* impacts patient recurrence. We observed that expression of each

molecule individually did not decrease RFS, but co-expression of *SEMA7A* in $α6^{high}$ tumors did

(HR=1.5; 95% CI=1.18-1.9; p=0.00075) **(Figure 6F and Supplemental Figure 11)**. Taken

together, these data suggest a clinically relevant mechanism by which SEMA7A and a6-integrin

may promote cellular phenotypes associated with tumor progression and chemoresistance.

DISCUSSION

Despite published roles for SEMA7A in breast cancer progression, the current study is, to our

knowledge, the first investigation into the role of SEMA7A in normal MEC survival during

postpartum mammary gland involution. In 2004, Clarkson and Watson published a

comprehensive analysis of gene expression across mammary gland development in which they

identified roles for both death receptors and immunomodulation during specific phases of

postpartum development including the first and second phases of involution(43). In this highly

impactful study, they identified seven basic profiles of gene expression across the lactation-

involution cycle. *SEMA7A* was identified in the Inv1 cluster, which was upregulated at the

transition between lactation and involution. In this study, we assessed cell surface SEMA7A

protein expression by flow cytometry in recently weaned mice to show its expression during the

initial phases of postpartum involution. We found that the proportion of MECs expressing

SEMA7A peaks at involution day 3, marking the transition from the reversible to the irreversible

phase of mammary gland remodeling. During the irreversible phase, MECs lose contact with the

basement membrane, primarily composed of laminin, and die by anoikis. Since anoikis is one of

the hallmarks of the irreversible phase of mammary involution, and our lab has published that

SEMA7A overexpression in multiple breast cancer cell lines decreases anoikis and, most

recently, that SEMA7A expression increases over time when cells are detached from matrix, we

predicted that expression of SEMA7A would promote anoikis resistance during the second

phase of irreversible involution(13, 44). We now also show that treating cells with exogenous

SEMA7A can confer resistance to anoikis in cultured human MECs suggesting that SEMA7A

may signal to neighboring cells to promote resistance to cell death. This is consistent with

recent data showing that, in human breast tissues, entire acini are lost during involution rather

than a percentage of cells from each acini(45). We propose that SEMA7A promotes anoikis

resistance, in part, by signaling to detached cells in a manner similar to matrix molecules,

thereby mimicking attachment. Consistent with this hypothesis, we show that exogenous

SEMA7A increases pAKT, and cell survival, in cultured MECs in a manner that is dependent, in

part, on $\alpha6$ and $\beta1$-integrin, which is normally the integrin receptor pair for pro-survival signaling

via ligation by laminin in the mammary gland(6, 24). While $\beta1$-integrin is a known mediator of

SEMA7A signaling in a variety of contexts, few studies have addressed the alpha integrin that

cooperates with $\beta1$ to mediate SEMA7A-activated signaling. Our results presented herein

suggest, for the first time, that SEMA7A may mimic the effects of laminin to promote anoikis

resistance via $\alpha6\beta1$ heterodimers in MECs. One limitation of our study is that we have not

directly assessed the effect of SEMA7A on anoikis-resistance in the presence of an AKT

inhibitor which would directly establish a relationship between SEMA7A and AKT-dependent

anoikis-resistance. However, literature suggests that AKT is important for promoting anoikis

resistance, via inhibition of BIM, and that loss of AKT increases cell death by anoikis. Therefore,

the ability of SEMA7A to enrich for pAKT+ cells suggests it may be sufficient to induce anoikis-

resistance through published AKT-dependent mechanisms, but that SEMA7A is not the only

molecule that can activate AKT to promote cell survival(46-48). Additional investigation is

required to further understand the role of SEMA7A in this pathway.

Anoikis resistance is a hallmark of stem cells and breast cancer cells, particularly those that

result in lethal metastasis as these cells must survive in the absence of extracellular matrix

signaling as they travel to and throughout the vasculature to seed distant tumors(49). Since

tumor cells are known to hijack normal developmental programs, we postulated that sustained

expression of SEMA7A in MECs, particularly in postpartum women, could result in pro-tumor

characteristics that are similar to those induced by Ras—a known oncogene. This hypothesis

was based on our recent publication showing that SEMA7A expression is higher in normal

adjacent breast tissue from postpartum women and our identification of a cut-point of

expression that predicts for recurrence in women with PPBC(50). Our hypothesis is supported

by our findings that SEMA7A mediates hallmarks of cellular transformation and stemness in

non-transformed cultured MECs. While others have shown that SEMA7A is required for Ras

dependent induction of EMT(51), we now show that growth in soft agar in Ras transformed

variants of MCF10A cells is dependent on SEMA7A and that SEMA7A can promote growth of

non-transformed cells in soft agar(26, 27, 30). Therefore, we predict that SEMA7A promotes

cellular transformation and tumor initiation in breast cancer patients and, in particular, in PPBC

patients(22, 30). A question that remains unanswered is whether sustained expression of

SEMA7A is causative for malignancy or whether SEMA7A expression in the tumor influences

normal adjacent cells to express SEMA7A thereby promoting tumor progression and

metastasis. Evidence for the latter comes from our studies showing that SEMA7A can affect

multiple aspects of the tumor microenvironment including collagen and fibronectin deposition,

lymphangiogenesis and macrophage infiltration, as well as neighboring cells such as

fibroblasts(26, 30). Additionally, we provide evidence that SEMA7A expressing cells upregulate

drug resistant cell-surface markers(23) and therefore may not respond to conventional

chemotherapy; this, along with our analysis of RFS in a large patient dataset revealing that co-

expression of *ITGA6* and *SEMA7A* significantly influences probability of relapse suggests that

SEMA7A can not only initiate tumor formation, but also influence clinical outcomes. Future

studies will investigate these important questions to identify how to best prevent and treat PPBC

and all patients with α6-integrin+/SEMA7A+ breast cancers.

Finally, to date, the majority of research into PPBC has focused on how the tissue

microenvironment in the postpartum breast drives metastasis with lesser focus on cellular

mechanisms at play in the MECs that may also support transformation(50). As more than half of

all breast cancers in patients under 40 can be defined as PPBC, which are highly metastatic,

understanding the mechanisms at play in postpartum breast tissue is, thus, critically important

for identifying molecular drivers of PPBC. In this study, we have identified a known tumor-

promotional protein, SEMA7A, as being temporally expressed during the irreversible phase of

postpartum mammary gland involution where it promotes anoikis resistance ,and we have

shown for the first time that SEMA7A can promote properties of transformation, stemness, and

metastasis in non-cancerous cells.

ACKNOWLEDGEMENTS

This work was supported by NIH/NCI R01 CA211696-01A1 (to T.R. Lyons), NIH/NCI R01CA211696-

02S1(to Taylor Rutherford), ACS RSG-16-171-010CSM (to T.R. Lyons), and Department of

Medicine Outstanding Early Career Scholars Award (to T.R. Lyons). The authors acknowledge the

University of Colorado Cancer Center Flow Cytometry Core for use of equipment and technical

assistance and the University of Colorado Cancer Center Support Grant (P30CA046934). The

authors also thank Alexander Stoller for assistance with animal studies and Veronica Wessels for

histological expertise. Contents are the authors' sole responsibility and do not necessarily represent

official NIH views.

The authors declare no potential conflicts of interest.

REFERENCES

1.   Goddard ET, Bassale S, Schedin T, Jindal S, Johnston J, Cabral E, et al. Association Between
Postpartum Breast Cancer Diagnosis and Metastasis and the Clinical Features Underlying Risk. JAMA
Netw Open. 2019;2(1):e186997.
2.   Lund LR, Romer J, Thomasset N, Solberg H, Pyke C, Bissell MJ, et al. Two distinct phases of
apoptosis in mammary gland involution: proteinase-independent and -dependent pathways.
Development. 1996;122(1):181-93.
3.   Schere-Levy C, Buggiano V, Quaglino A, Gattelli A, Cirio MC, Piazzon I, et al. Leukemia Inhibitory
Factor Induces Apoptosis of the Mammary Epithelial Cells and Participates in Mouse Mammary Gland
Involution. Experimental Cell Research. 2003;282(1):35-47.
4.   Gilmore AP. Anoikis. Cell death and differentiation. 2005;12 Suppl 2:1473-7.
5.   Boudreau N, Sympson CJ, Werb Z, Bissell MJ. Suppression of ICE and apoptosis in mammary
epithelial cells by extracellular matrix. Science. 1995;267(5199):891-3.
6.   Pullan S, Wilson J, Metcalfe A, Edwards GM, Goberdhan N, Tilly J, et al. Requirement of
basement membrane for the suppression of programmed cell death in mammary epithelium. Journal of
cell science. 1996;109 (Pt 3):631-42.
7.   Cerny J, Stockinger H, Horejsi V. Noncovalent associations of T lymphocyte surface proteins. Eur
J Immunol. 1996;26(10):2335-43.
8.   Pasterkamp RJ, Peschon JJ, Spriggs MK, Kolodkin AL. Semaphorin 7A promotes axon outgrowth
through integrins and MAPKs. Nature. 2003;424(6947):398-405.
9.   Jaimes Y, Gras C, Goudeva L, Buchholz S, Eiz-Vesper B, Seltsam A, et al. Semaphorin 7A inhibits
platelet production from CD34+ progenitor cells. Journal of thrombosis and haemostasis : JTH.
2012;10(6):1100-8.
10.   Delorme G, Saltel F, Bonnelye E, Jurdic P, Machuca-Gayet I. Expression and function of
semaphorin 7A in bone cells. Biology of the cell / under the auspices of the European Cell Biology
Organization. 2005;97(7):589-97.
11.   Esnault S, Torr EE, Bernau K, Johansson MW, Kelly EA, Sandbo N, et al. Endogenous Semaphorin-
7A Impedes Human Lung Fibroblast Differentiation. PloS one. 2017;12(1):e0170207.
12.   De Minicis S, Rychlicki C, Agostinelli L, Saccomanno S, Trozzi L, Candelaresi C, et al. Semaphorin
7A contributes to TGF-beta-mediated liver fibrogenesis. The American journal of pathology.
2013;183(3):820-30.
13.   Tarullo SE, Hill RC, Hansen K, Behbod F, Borges VF, Nelson AC, et al. Postpartum breast cancer
progression is driven by semaphorin 7a-mediated invasion and survival. Oncogene.
2020;https://doi.org/10.1038/s41388-020-1192-9.

14.   Czopik AK, Bynoe MS, Palm N, Raine CS, Medzhitov R. Semaphorin 7A Is a Negative Regulator of
T Cell Responses. Immunity. 2006;24(5):591-600.
15.   Xie J, Wang H. Semaphorin 7A as a potential immune regulator and promising therapeutic target
in rheumatoid arthritis. Arthritis Res Ther. 2017;19(1):10.
16.   Elder AM, Tamburini BA, Crump LS, Black SA, Wessells VM, Schedin PJ, et al. Semaphorin 7A
promotes macrophage-mediated lymphatic remodeling during postpartum mammary gland involution
and in breast cancer. Cancer research. 2018.
17.   Esnault S, Kelly EA, Johansson MW, Liu LY, Han ST, Akhtar M, et al. Semaphorin 7A is expressed
on airway eosinophils and upregulated by IL-5 family cytokines. Clinical immunology. 2014;150(1):90-
100.
18.   Gan Y, Reilkoff R, Peng X, Russell T, Chen Q, Mathai SK, et al. Role of semaphorin 7a signaling in
transforming growth factor beta1-induced lung fibrosis and scleroderma-related interstitial lung disease.
Arthritis and rheumatism. 2011;63(8):2484-94.
19.   Lee Y, Blount KL, Dai F, Thompson S, Scher JK, Bitterman S, et al. Semaphorin 7A in circulating
regulatory T cells is increased in autosomal-dominant polycystic kidney disease and decreases with
tolvaptan treatment. Clin Exp Nephrol. 2018;22(4):906-16.
20.   Garcia-Areas R, Libreros S, Amat S, Keating P, Carrio R, Robinson P, et al. Semaphorin7A
promotes tumor growth and exerts a pro-angiogenic effect in macrophages of mammary tumor-bearing
mice. Frontiers in physiology. 2014;5:17.
21.   Garcia-Areas R, Libreros S, Simoes M, Castro-Silva C, Gazaniga N, Amat S, et al. Suppression of
tumor-derived Semaphorin 7A and genetic ablation of host-derived Semaphorin 7A impairs tumor
progression in a murine model of advanced breast carcinoma. Int J Oncol. 2017;51(5):1395-404.
22.   Borges VF, Hu J, Young C, Maggard J, Parris HJ, Gao D, et al. Semaphorin 7a is a biomarker for
recurrence in postpartum breast cancer. npj Breast Cancer. 2020;6(1):56.
23.   Crump LS, Wyatt GL, Rutherford TR, Richer JK, Porter WW, Lyons TR. Hormonal Regulation of
Semaphorin 7a in ER(+) Breast Cancer Drives Therapeutic Resistance. Cancer research. 2021;81(1):187-
98.
24.   Gilmore AP, Metcalfe AD, Romer LH, Streuli CH. Integrin-mediated survival signals regulate the
apoptotic function of Bax through its conformation and subcellular localization. The Journal of cell
biology. 2000;149(2):431-46.
25.   Krebsbach PH, Villa-Diaz LG. The Role of Integrin alpha6 (CD49f) in Stem Cells: More than a
Conserved Biomarker. Stem cells and development. 2017;26(15):1090-9.
26.   Tarullo SE, Hill RC, Hansen KC, Behbod F, Borges VF, Nelson AC, et al. Postpartum breast cancer
progression is driven by semaphorin 7a-mediated invasion and survival. Oncogene. 2020;39(13):2772-
85.
27.   Black SA, Nelson AC, Gurule NJ, Futscher BW, Lyons TR. Semaphorin 7a exerts pleiotropic effects
to promote breast tumor progression. Oncogene. 2016;35(39):5170-8.
28.   Lyons TR, Borges VF, Betts CB, Guo Q, Kapoor P, Martinson HA, et al. Cyclooxygenase-2-
dependent lymphangiogenesis promotes nodal metastasis of postpartum breast cancer. The Journal of
clinical investigation. 2014;124(9):3901-12.
29.   Gustafson-Wagner E, Stipp CS. The CD9/CD81 tetraspanin complex and tetraspanin CD151
regulate alpha3beta1 integrin-dependent tumor cell behaviors by overlapping but distinct mechanisms.
PloS one. 2013;8(4):e61834.
30.   Elder AM, Tamburini BAJ, Crump LS, Black SA, Wessells VM, Schedin PJ, et al. Semaphorin 7A
Promotes Macrophage-Mediated Lymphatic Remodeling during Postpartum Mammary Gland Involution
and in Breast Cancer. Cancer research. 2018;78(22):6473-85.

31.   O'Brien J, Lyons T, Monks J, Lucia MS, Wilson RS, Hines L, et al. Alternatively activated
macrophages and collagen remodeling characterize the postpartum involuting mammary gland across
species. The American journal of pathology. 2010;176(3):1241-55.
32.   Chandrashekar DS, Bashel B, Balasubramanya SAH, Creighton CJ, Ponce-Rodriguez I,
Chakravarthi B, et al. UALCAN: A Portal for Facilitating Tumor Subgroup Gene Expression and Survival
Analyses. Neoplasia. 2017;19(8):649-58.
33.   Chen F, Chandrashekar DS, Varambally S, Creighton CJ. Pan-cancer molecular subtypes revealed
by mass-spectrometry-based proteomic characterization of more than 500 human cancers. Nature
communications. 2019;10(1):5679.
34.   Lanczky A, Gyorffy B. Web-Based Survival Analysis Tool Tailored for Medical Research (KMplot):
Development and Implementation. J Med Internet Res. 2021;23(7):e27633.
35.   Lee YJ, Streuli CH. Extracellular matrix selectively modulates the response of mammary epithelial
cells to different soluble signaling ligands. The Journal of biological chemistry. 1999;274(32):22401-8.
36.   Farrelly N, Lee YJ, Oliver J, Dive C, Streuli CH. Extracellular matrix regulates apoptosis in
mammary epithelium through a control on insulin signaling. The Journal of cell biology.
1999;144(6):1337-48.
37.   Shehata M, Teschendorff A, Sharp G, Novcic N, Russell IA, Avril S, et al. Phenotypic and
functional characterisation of the luminal cell hierachy of the mammary gland. Breast cancer research :
BCR. 2012;14(5):R134.
38.   Visvader JE, Stingl J. Mammary stem cells and the differentiation hierarchy: current status and
perspectives. Genes & development. 2014;28(11):1143-58.
39.   Tosoni D, Di Fiore PP, Pece S. Functional purification of human and mouse mammary stem cells.
Methods in molecular biology. 2012;916:59-79.
40.   Dontu G, Abdallah WM, Foley JM, Jackson KW, Clarke MF, Kawamura MJ, et al. In vitro
propagation and transcriptional profiling of human mammary stem/progenitor cells. Genes &
development. 2003;17(10):1253-70.
41.   Cicalese A, Bonizzi G, Pasi CE, Faretta M, Ronzoni S, Giulini B, et al. The tumor suppressor p53
regulates polarity of self-renewing divisions in mammary stem cells. Cell. 2009;138(6):1083-95.
42.   Liu R, Chen Y, Liu G, Li C, Song Y, Cao Z, et al. PI3K/AKT pathway as a key link modulates the
multidrug resistance of cancers. Cell Death Dis. 2020;11(9):797.
43.   Clarkson RW, Wayland MT, Lee J, Freeman T, Watson CJ. Gene expression profiling of mammary
gland development reveals putative roles for death receptors and immune mediators in post-lactational
regression. Breast cancer research : BCR. 2004;6(2):R92-109.
44.   Crump LS, Wyatt G, Rutherford TR, Richer JK, Porter WW, Lyons TR. Hormonal regulation of
Semaphorin 7a in ER+ breast cancer drives therapeutic resistance. Cancer research. 2020.
45.   Jindal S, Gao D, Bell P, Albrektsen G, Edgerton SM, Ambrosone CB, et al. Postpartum breast
involution reveals regression of secretory lobules mediated by tissue-remodeling. Breast cancer research
: BCR. 2014;16(2):R31.
46.   Adeshakin FO, Adeshakin AO, Afolabi LO, Yan D, Zhang G, Wan X. Mechanisms for Modulating
Anoikis Resistance in Cancer and the Relevance of Metabolic Reprogramming. Front Oncol.
2021;11:626577.
47.   de Sousa Mesquita AP, de Araujo Lopes S, Pernambuco Filho PCA, Nader HB, Lopes CC.
Acquisition of anoikis resistance promotes alterations in the Ras/ERK and PI3K/Akt signaling pathways
and matrix remodeling in endothelial cells. Apoptosis. 2017;22(9):1116-37.
48.   Woods NT, Yamaguchi H, Lee FY, Bhalla KN, Wang HG. Anoikis, initiated by Mcl-1 degradation
and Bim induction, is deregulated during oncogenesis. Cancer research. 2007;67(22):10744-52.
49.   Cao Z, Livas T, Kyprianou N. Anoikis and EMT: Lethal "Liaisons" during Cancer Progression.
Critical reviews in oncogenesis. 2016;21(3-4):155-68.

50.   Borges VF, Lyons TR, Germain D, Schedin P. Postpartum Involution and Cancer: An Opportunity
for Targeted Breast Cancer Prevention and Treatments? Cancer research. 2020;80(9):1790-8.
51.   Allegra M, Zaragkoulias A, Vorgia E, Ioannou M, Litos G, Beug H, et al. Semaphorin-7a reverses
the ERF-induced inhibition of EMT in Ras-dependent mouse mammary epithelial cells. Molecular biology
of the cell. 2012;23(19):3873-81.

FIGURE LEGENDS

Figure 1: Semaphorin-7a promotes anoikis resistance in normal MECs. a. Cleaved

caspase 3/7 (CC3/7) expression in MCF10A EV or S7a-OE cells after 24 hours in attached or

forced suspension cultures. **b.** SEMA7A expression by flow cytometry in attached and forced

suspension MCF12A cultures. **c.** CC3/7 in MCF12A cells in forced suspension +/- exogenous

S7a protein and/or ITGB1 blocking antibody, 9eg7. **d**. Flow cytometry analysis of pAKT in

insulin or S7a-treated MCF12A with representative flow plots. **e.** Flow cytometry for %SEMA7A+

mammary epithelial cells from tissues harvested from nulliparous (N) or recently weaned

C57/BL6 mice at involution days 1-28 (I1-I28). **f.** Flow cytometry of live epithelial cells as

measured by Aquazombie-EpCAM+ in mammary tissues harvested from recently weaned WT

or Sema7a[tm1Alk/J] mice. WT (I1: n=5, I2: n=6, I3: n=5, I6: n=4 glands/group) SEMA7A [tm1Alk/J] (I1-

I6: n=6 glands/group). Error bars indicate mean +/- standard error of the mean. Significance

denoted as: $p < 0.05 = *$, $p < 0.01 = **$, $p < 0.001 = ***$, $p < 0.0001 = ****$.

Figure 2: Loss of SEMA7A promotes histologic features of accelerated involution.

Representative 20x images of #3 mammary glands from WT or SEMA7A [tm1Alk/J] mice at 1,2,3,&6

days of forced involution. **a.** Hematoxylin and Eosin, **b.** pSTAT3, **c.** cleaved caspase-3 and **d.**

perilipin staining.

Figure 3: Loss of SEMA7A promotes alveolar collapse through alterations to cell survival

and adipocyte repopulation. Immunohistochemical quantification of mammary glands from

WT (I1: n=5, I2: n=6, I3: n=5, I6: n=4 glands/group) or SEMA7A [tm1Alk/J] (I1-I6: n=6 glands/group)

mice at 1,2,3,&6 days post forced involution. Quantification of **a.** alveolar area **b.** pSTAT3 **c.**

cleaved caspase-3 and **d.** perilipin staining. Error bars indicate mean +/- standard error of the

mean. Significance denoted as: $p<0.05=*$, $p<0.01=**$, $p<0.001=***$, $p<0.0001=****$.

Figure 4: Semaphorin-7a promotes survival via integrins and enriches for pAKT+ cells. a-

e. Flow cytometry analysis of mammary glands from WT (left, I1: n=4, I2: n=6, I3: n=5, I6: n=4

glands/group) and SEMA7A$^{tm1Alk/J}$ (right, I1-I6: n=6 glands/group) mice. **a.** Stratification of MEC

subtypes by EpCAM/CD49f staining. **b.** %EpCAMhiCD49f+ LPCs of EpCAM+ cells. **c.**

%SEMA7A+ of live (Aquazombie-) EpCAMhiCD49f+ LPCs. **d. %**ITGB1+ITGA6+ of EpCAM

cells. **e.** %SEMA7A of ITGB1+ITGA6+ cells. **f.** CC3/7 expression in MCF12A cells +/- S7a

and/or GoH3. **g.** %pAKT in MCF12A cells +/- S7a and/or 9eg7/GoH3, n=3. Error bars indicate

mean +/- standard error of the mean. Significance denoted as: $p<0.05=*$, $p<0.01=**$,

$p<0.001=***$, $p<0.0001=****$.

Figure 5: SEMA7A and ITGB1/ITGA6 promote features of transformation and

chemoresistance. a. %Celltrace violet bright of singlets in MCF10A EV and OE cells. **b,c.**

Number of mammospheres formed by MCF12A cells +/- S7a and GoH3 or 9eg7. **d.** Number of

colonies formed in soft agar assay.

Figure 6: *Sema7a* is upregulated in multiple breast cancers and predicts for decreased

relapse-free survival when expressed with α6-integrin. a-d. TCGA data for *Sema7a* in a.

normal (n=114) vs primary tumor (1097) **b.** normal versus stages I-IV (I: n=183, II: n=615, III:

n=247, IV: n=20) **c.** normal versus cancer stratified by age (21-40: n=97, 41-60: n=505, 61-80:

n=431, 81-100: n=54) , **d.** normal versus breast cancer subclass (luminal: n=566, HER2+: n=37,

TNBC-BL1: n=13, TNBC-BL2: n=11, TNBC-IM: n=20, TNBC-LAR: n=8, TNBC-MSL: n=8,

TNBC-M: n=29, TNBC-UNS: n=27) Error bars indicate ??? Significance denoted as: $p<0.05=*$,

p<0.01=**, p<0.001=***, p<0.0001=****. **e.** Kaplan-Meier 5-year relapse-free survival analysis

for *Sema7a in Itga6* low (n=2032) or high (n=2032) expressing patients.