Anoikis resistance in mammary epithelial cells is mediated by semaphorin 7a

Semaphorin-7A and anoikis resistance

Taylor R. Rutherford[1,23] Alan M Elder[1,2,4] Traci R. Lyons[1,2,4,5]

Corresponding author: traci.lyons@cuanscutz.edu

[1] Division of Medical Oncology, Department of Medicine, School of Medicine, University of Colorado Anschutz Medical Campus, Aurora, Colorado.

[2] Young Women's Breast Cancer Translational Program, University of Colorado Anschutz Medical Campus, Aurora, Colorado.

[3] Cell biology, Stem cell, and Development Graduate Training Program, University of Colorado Anschutz Medical Campus, Aurora, Colorado.

[4] Cancer biology Graduate Training Program, University of Colorado Anschutz Medical Campus, Aurora, Colorado.

[5] University of Colorado Cancer Center, Aurora, Colorado.

ABSTRACT

Semaphorin-7a (SEMA7A), best known as a neuroimmune molecule, plays a diverse role in many cellular processes and pathologies. Here, we show that SEMA7A promotes anoikis resistance in cultured mammary epithelial cells through integrin mediated activation of pro-survival kinase AKT, which led us to investigate a role for SEMA7A during postpartum mammary gland involution—a normal process in development where cells die by anoikis. Our results reveal that SEMA7A is expressed on live mammary epithelial cells during involution, that SEMA7A expression is primarily observed in α6-integrin expressing cells, and that luminal progenitor cells, specifically, are decreased in mammary glands of SEMA7A-/- mice during involution. We further identify a SEMA7A-α6/β1-integrin dependent mechanism of cell survival that promotes anoikis resistance, mammosphere formation, and chemoresistance in mammary epithelial cells and suggest that this mechanism is relevant for recurrence in breast cancer patients.

INTRODUCTION

Identifying mechanisms of normal mammary epithelial cell (MEC) biology that contribute to tumorigenesis is critically important for furthering our understanding of breast cancer and extending the survival of breast cancer patients. It has been postulated that the functional plasticity of the adult mammary epithelium that is required for response to hormonal changes during pregnancy and lactation may make the tissue more susceptible to malignancy. Of particular interest are postpartum breast cancers (PPBCs), defined as breast cancers diagnosed in women within 10 years of most recent childbirth, as these cancers are nearly three times more likely to result in lethal metastasis(1).

The majority of mammary morphogenesis occurs postnatally beginning at puberty and

culminating with lactation following pregnancy. As such, differentiation of the mammary

epithelium is not complete until a full pregnancy-lactation cycle. Upon weaning, or after

pregnancy in the absence of nursing, the mammary gland regresses to resemble the pre-

pregnant architecture through a process termed postpartum involution. Postpartum involution

occurs in two primary phases(2). In the first phase, milk proteins accumulate in the mammary

ducts and alveoli resulting in activation of leukemia inhibitory factor (LIF), phosphorylation of

signal transducer and activator of transcription (pSTAT3), and subsequent lysosomal-mediated

cell death(3). In mice, if pups are returned within 48 hours of weaning, this process can be

reversed, and lactation can continue. If pups are not returned, however, the irreversible second

phase of involution begins. This phase is characterized by degradation of the mammary

basement membrane by matrix metalloproteinases and anoikis-mediated cell death in the

absence of laminin-mediated activation of integrin alpha-6/beta-1 ($\alpha6\beta1$) and downstream pro-

survival signaling(4-6).

SEMA7A, or CD108w, is a member of the semaphorin protein family, named after the English

verb "semaphore", which means to transmit a signal across a long distance. SEMA7A was first

identified for its expression on lymphocytes(7), but has since been shown to exert diverse roles

in embryonic neuronal development(8), cell differentiation(9-11), ECM deposition(12, 13),

various components of the immune system(14-17), and pathologies including fibrosis(12, 18,

19) and cancer·(13, 16, 20-22). Though elevated SEMA7A is observed in breast cancer, where

it has an established role in promoting tumor cell survival(13, 23), the role of SEMA7A in normal

MECs has not previously been investigated. SEMA7A is known to signal through $\beta1$-integrin in

immune cells, neurons, and cancer; $\beta1$-integrin also mediates MEC survival during

development, in part, via phosphorylation and activation of downstream pro-survival signaling

mediated by PI3-kinase and protein kinase B, or Akt(24). Though it is known that redundant

secretory epithelial cells die during postpartum involution, the factor(s) that allow some cells to

survive and regenerate a lactation-competent gland for future pregnancies remain unknown.

Herein, we have identified enrichment for SEMA7A expressing $\alpha6\beta1$ positive MECs during the

matrix remodeling period of postpartum involution and a role for SEMA7A in specifically

promoting the survival of $\alpha6$-integrin+ luminal progenitor cells. We provide evidence that

exogenous stimulation of immortalized MECs with purified SEMA7A protein promotes anoikis

resistance via activation of $\alpha6$ and $\beta1$-integrins and downstream activation of phosphorylated

AKT (pAKT). Consistent with a role for SEMA7A in promoting anoikis resistance, SEMA7A

knockout mice (SEMA7A[tmalk1/J]) exhibit phenotypes associated with accelerated involution

including early presence of apoptotic cell bodies in mammary lumens, increased pSTAT3, and

premature adipocyte repopulation. $\alpha6$-integrin also drives maintenance of tissue resident stem

cells and therapy resistant cancer stem cells(25) and our findings reveal that SEMA7A drives

stem cell phenotypes and chemoresistance in cultured MECs in a manner that is dependent on

$\alpha6\beta1$. Finally, we provide evidence for SEMA7A in driving transformed phenotypes in normal

MECs and reversing transformed phenotypes in genetically identical, ras-transformed, variants

of the same cell line. Our results are the first to identify a role for SEMA7A in anoikis resistance

and promotion of stem and cancer cell phenotypes via activation of $\alpha6$ and $\beta1$-integrins.

METHODS

Cell culture:

MCF10A/MCF12A cells were obtained from, and cultured, according to ATCC.

MCF10DCIS.com cells were obtained from K. Polyak (University of Harvard, Cambridge, MA).

Cells were validated by the DNA sequencing core at the University of Colorado Anschutz

Medical Campus and found to be pure populations of their respective cell lines. Cells were

regularly tested for mycoplasma throughout studies. Cells were sub-cultured as previously

described (reference), or according to ATCC standards. Overexpression plasmid (SEMA7A-Fc)

was a generous gift from R. Medzhitov (Yale University, New Haven, CT). Control plasmid

(pcDNA3.1) was obtained from H. Ford (University of Colorado Anschutz Medical Campus,

Denver, CO).

Animal Studies:

All animal procedures were approved by the University of Colorado Anschutz Medical Campus

Institutional Animal Care and Use Committee. Sema7a[tm1Alk/J] mice (a generous gift from Alex

Kolodkin at John's Hopkins University) and C57Bl/6 (Jax) were housed and bred as previously

described.[26] Briefly, C57Bl/6 SEMA7A[tm1Alk] females were bred with WT C57Bl/6 males to

induce pregnancy, lactation, and postpartum involution. Pup numbers were normalized to 6-8

per dam after birth to ensure for adequate lactation. Postpartum involution was initiated by

forced weaning of pups at the peak of lactation (L10-L12). #4 right and left mammary glands

were harvested from age-matched nulliparous animals and postpartum dams at involution days

1,2,4,6,8,10,14, and 28. For flow cytometry, inguinal lymph nodes were removed prior to tissue

digestion.

In vitro survival assays:

MCF10A and MCF12A cells were cultured with 200ng/ml purified SEMA7A or PBS vehicle

control in adherent (tissue culture plastic) or forced suspension conditions (ultra-low attachment

plates (Corning, Corning, NY, USA: #3473)). Cells were seeded at a density of 1000 cells/well in

a 96-well plate plus media only wells as controls. Cell death was measured 24 hours post

seeding via luminescence using the Caspase Glo assay (Promega, Madison, WI, USA:

#G8090). Annexin-V /7AAD staining (Biolegend, San Diego, CA, USA: #640930) and

pAKTS473 (ThermoFisher: Waltham, MA, USA, #17-9715-42) were also used to confirm cell

viability by flow cytometry. See flow cytometry methods for staining protocol.

Label retention stem cell assay:

MCF10A and MCF12A cells were labeled with CellTrace Violet (ThermoFisher: #C34557) for 20

minutes at 37°C according to manufacturer's instructions. A sample of labeled cells was

analyzed by flow cytometry to determine labeling efficiency and fluorescent intensity at time of

seeding. Single cells were seeded at a density of 4000 cells/well in ultra-low attachment 24-well

plates to induce mammosphere formation. Cells were cultured in normal culture media for 7

days and were subsequently counted, dissociated enzymatically in Accutase, (Stem Cell

Technologies, Vancouver, Canada: #07922), and analyzed by flow cytometry. See flow

cytometry methods for detailed information.

Paclitaxel resistance studies: MCF10DCIS.com cells were treated with 100nM paclitaxel

(Millipore Sigma, Burling, MA, USA #T7402-5MG) for 24 hours prior to analysis by flow

cytometry. Drug resistance was measured by label-retention and mammosphere formation as

described above and %CD44+ (Biolegend: #103018) CD24- (Biolegend: #311104)

of singlets was used as a molecular definition of drug-resistant stem cells.

Flow cytometry:

Single cell suspensions were generated from cultured cells via enzymatic harvest with

Accutase. Mouse mammary tissues were minced with scalpels, digested in Click's media

containing 500 units/mL Collagenase II (Worthington, Columbus, OH, USA; LS004174) and

Collagenase IV (Worthington, LS004186) and 20ug/mL DNAse (Worthington, LS002004) for 1

hour at 37°C with occasional trituration and strained through 70um filters to generate single cell

suspensions. Single cells were stained at 4°C for 30 minutes and washed with PBS+2%FBS.

Cells were filtered through 30um filters, diluted to $<10^6$ cells/ml, and analyzed with the ZE5-YETI

flow cytometer. Fixation/Permeabilization kit (BD Biosciences, Franklin Lakes, NJ, USA:

#554715) was used for intracellular antigen staining. Data were analyzed with Kaluza software.

Single cells were lineage depleted using CD45- (BD Biosciences: #564279)/CD31- (BD

Biosciences: #745436) gating.

Immunohistochemistry:

Tissues were formalin-fixed and paraffin-embedded as previously described(27). Hematoxylin

and Eosin staining was used to define morphological features. Antigen retrieval was performed

using target retrieval solution (Dako, Glostrup, Denmark; #S1699) for CC3(Cell Signaling

Technologies, Danvers, MA, USA; #9661) and Perilipin(Cell Signaling: #3470) or (EDTA Dako

cat# S2367) for pSTAT3 (Cell Signaling Technologies; #9145). ImmPRESS polymer anti-rabbit

IgG secondary reagent (Vector Labs, San Francisco, CA, USA t#MP-7401-15) was used for

secondary staining of CC3 and pSTAT3 stained tissues and anti-rabbit secondary (Dako

#K4003) for perilipin stains. DAB (Vector Labs,; #SK-4105) and counterstaining were performed

with hematoxylin (Vector Labs #H-3401). Alveolar area was measured on H&E-stained tissues

in ImageJ on 10 alveoli/field; 5 fields/tissue. 10 fields/tissue were analyzed for positive CC3,

pSTAT3, and perilipin stained tissues using CellSense Dimension software count and measure

feature on regions of interest.

Statistical Analysis:

Unpaired *t*-test and one-way ANOVA were performed in GraphPad Prism. Nonparametric

equivalent analyses were used for samples with uneven distribution or unequal standard

deviations. *p* values of less than 0.05 were considered significant. Error bars represent mean +/-

standard error of the mean. All in vivo studies were replicated in at least 3 mice and in vitro

studies performed in technical and biological triplicate with representative data presented.

RESULTS

Semaphorin-7a promotes anoikis resistance in normal MECs

To identify whether SEMA7A promotes survival of normal MECs we utilized two methods: transfection of FC-tagged SEMA7A into immortalized human (MCF10A) cells and addition of exogenous SEMA7A to wild-type MCF10A and MCF12A cells—another non-transformed MEC line. We found that genetic SEMA7A overexpression resulted in resistance to cell death, as measured by cleaved caspase 3/7, in a forced suspension assay, but not in attached conditions suggesting that SEMA7A specifically mediates resistance to cell death by anoikis. Therefore, all subsequent analyses of cell death were performed in forced suspension. Stimulation with exogenous purified SEMA7A protein in both MCF10A and MCF12A cells also significantly decreased cell death **(Figure 1A, B)**. We independently confirmed decreased cell death in detached MECs cultured with exogenous SEMA7A by flow cytometry for Annexin-V/7AAD **(SFigure 1)**. Inhibition of a known SEMA7A receptor, β1-integrin, with the function blocking antibody 9EG7 significantly abrogated SEMA7A-mediated anoikis resistance **(Figure 1B).** We also examined pAKT in MECs cultured with SEMA7A using insulin as a positive control as the IGF-1 receptor (and insulin) can activate pAKT via β1-integrin in the mammary gland(28, 29). We observed percentages of pAkt expressing cells similar to those observed with insulin when cells were exposed to SEMA7A *in vitro* **(Figure 1C)**.

To determine whether SEMA7A inhibits anoikis *in vivo*, we examined MECs harvested from age-matched nulliparous and force weaned WT animals for SEMA7A expression. We observed a low percentage of SEMA7A+EpCAM+ cells in the nulliparous mammary gland, which was enriched approximately 3-fold at involution day 3 when basement membrane degradation and extracellular matrix remodeling ensues. The percentage of SEMA7A+ epithelial cells declined sharply starting at involution day 4 and returned to levels comparable to those observed in

nulliparous glands at involution day 6 **(Figure 1D).** We also compared live epithelial cells by flow

cytometry in age matched SEMA7A knockout mice (Sema7a$^{tm1Alk/J}$) and WT C57Bl/6 mice at

days 1, 2, 3, & 6 post-wean and observed a modest, but significant, reduction in MECs as

measured by the percentage of Aquazombie- EpCAM+ cells **(Figure 1E)**. To test whether the

decrease in live cells during involution in Sema7a$^{tm1Alk/J}$ animals was accompanied by changes

to gland histology and programs normally activated during postpartum involution, we performed

H&E and immunohistochemical analysis of whole mammary tissues isolated from Sema7a$^{tm1Alk/J}$

dams and WT controls. We observed accelerated glandular regression, as evidenced by

decreased alveolar area, increased pSTAT3, precocious appearance of cleaved caspase 3

(CC3) positive cells, and increased adipocyte repopulation in glands harvested from

Sema7a$^{tm1Alk/J}$ animals **(Figure 2 and Figure 3)** suggesting that loss of SEMA7A accelerates

postpartum involution.

SEMA7A promotes anoikis-resistance via α6-integrin mediated pAKT activation.

MECs can be divided into four primary cell types: luminal, luminal progenitor (LPCs), basal, and

basal progenitor cells (BPCs) and these populations can be resolved by flow cytometry based

largely on their expression of epithelial cell adhesion molecule (EpCAM) and CD49f, also known

as α6-integrin(30, 31). Since the magnitude of the changes observed in Figure 1E was relatively

small, we used flow cytometry to resolve the four primary MEC populations to determine which

cells are specifically affected by loss of SEMA7A during postpartum involution. We observed a

significant decrease in EpCAM^{hi}CD49f+ LPCs at all timepoints examined, which were not

consistently observed in the other SEMA7A expressing cell populations. Additionally, luminal

cells remained unchanged **(Figure 4A, B and SFigure 2&3)**. We also found that more than

80% of the live LPCs express SEMA7A at involution days 2-4 **(Figure 4C)**. One factor that

distinguishes the SEMA7A+ MEC populations from the SEMA7A- luminal cells is their

expression of CD49f, or α6-integrin. β1-integrin is a known receptor that mediates SEMA7A

cellular signaling(8), but the preferred alpha binding partner has not been identified. We

quantified cells that co-express $\alpha6$ and $\beta1$ in WT and SEMA7A[tmalk1/J] animals and found a

significant reduction during involution **(Figure 4D)**. Furthermore, we observed that over 90% of

the $\alpha6$ and $\beta1$ positive MECs express SEMA7A at the transition to the reversible phase of

involution when basement membrane degradation and anoikis-induced cell death is initiated

(Figure 4E). To determine whether SEMA7A confers anoikis-resistance via $\alpha6$-integrin

dependent activation of pAKT, we cultured MCF10A cells with SEMA7A in the presence of the

$\alpha6$-integrin blocking antibody (GoH3) in forced suspension cultures. We observed increased

CC3/7+ cells in GoH3 conditions as well as a significant decrease in pAKT-expressing cells

(Figure 4F&G). Taken together, our results suggest SEMA7A promotes anoikis-resistance

during involution, in part, through $\alpha6\beta1$-mediated activation of pAKT.

SEMA7A promotes stem cell phenotypes, cellular transformation, and chemoresistance

Anoikis-resistance is rarely observed in normal epithelial cells; rather, it is a characteristic of

stem cells and tumor cells. Given our findings that SEMA7A promotes anoikis-resistance in

normal MECs, we tested whether it also confers features associated with stem cells and

transformation. We used a label-retention and mammosphere assay to identify phenotypes

associated with mammary stem cells *in vitro* and found that SEMA7A overexpression in

MCF10A cells increases the quiescent stem cell population **(Figure 5A & SFigure 4)**. We also

measured mammosphere formation as a readout for self-renewal in cells cultured with

exogenous SEMA7A in the presence or absence of function blocking antibodies for $\alpha6$ and $\beta1$

and found that increased sphere formation occurred with SEMA7A which was abrogated with

either inhibitor **(Figure 5B&C)**. Secondary sphere formation was also increased in MCF10A and

MCF12A cells cultured with SEMA7A **(SFigure 5).** We then tested the ability of SEMA7A to

promote cell growth in anchorage-independent conditions using a soft agar assay—a classical

measurement of cellular transformation. We found that overexpression of SEMA7A in MCF10A

cells is sufficient to increase growth in soft agar to levels similar to a Ras-transformed variant of

these cells, MCF10DCIS.com. Moreover, we found that shRNA-mediated knockdown of

SEMA7A in MCF10DCIS cells(13, 32) significantly decreased anchorage-independent growth,

to levels more similar to the MCF10A non-transformed EV control and this effect was rescued

by addition of exogenous SEMA7A **(Figure 5D).** Cancer stem cells, and tumor cells expressing

pAkt, are also known to be chemoresistant(33). To test whether SEMA7A promotes

chemoresistance in breast cancer cells, we treated SEMA7A-overexpressing and knockdown

MCF10DCIS mammosphere cultures with the chemotherapeutic drug, paclitaxel. We found that

SEMA7A overexpression increases mammosphere formation in treated MCF10DCIS cells,

which can be partially blocked with 9EG7 and completely blocked with GoH3. We then

confirmed an increase in tumor-initiating and drug-resistant cells in these cultures by increased

label-retention and CD44+CD24- expression—markers frequently used to characterize drug

resistant cancer stem cells **(Figure 5E-G).** Since patients with chemotherapy resistant disease

often experience disease relapse, we examined whether co-expression of *SEMA7A* and α6-

integrin (*ITGA6*) was associated with decreased relapse free survival (RFS) by Kaplan Meier

analysis. We observed that expression of each molecule individually did not decrease RFS, but

co-expression of *SEMA7A* in *ITGA6*[high] tumors did (HR=1.5; 95% CI=1.18-1.9; p=0.00075)

(Figure 6). Taken together, these data suggest a clinically relevant mechanism by which

SEMA7A and α6-integrin may promote cellular phenotypes associated with tumor progression

and chemoresistance.

DISCUSSION

Despite published roles for SEMA7A in breast cancer progression, the current study is, to our

knowledge, the first investigation into the role of SEMA7A in normal MEC survival during

postpartum mammary gland involution. In 2004, Clarkson and Watson published a

comprehensive analysis of gene expression across mammary gland development in which they

identified roles for both death receptors and immunomodulation during specific phases of

postpartum development including the first and second phases of involution(34). In this highly

impactful study, they identified seven basic profiles of gene expression across the lactation-

involution cycle. *SEMA7A* was identified in the Inv1 cluster, which was upregulated at the

transition between lactation and involution. In this study, we assessed cell surface SEMA7A

protein expression by flow cytometry in recently weaned mice to show its expression during the

initial phases of postpartum involution. We found that the proportion of MECs expressing

SEMA7A peaks at involution day 3, marking the transition from the reversible to the irreversible

phase of mammary gland remodeling. During the irreversible phase, MECs lose contact with the

basement membrane, primarily laminin, and die by anoikis. Since anoikis is one of the hallmarks

of the irreversible phase of mammary involution, and our lab has published that SEMA7A

overexpression in multiple breast cancer cell lines decreases anoikis and, most recently, that

SEMA7A expression increases over time when cells are detached from matrix, we predicted

that expression of SEMA7A would promote anoikis resistance during the second phase of

irreversible involution(13, 35). We now also show that treating cells with exogenous SEMA7A

can confer resistance to anoikis in cultured human MECs suggesting that SEMA7A may signal

to neighboring cells to promote resistance to cell death. This is consistent with recent data

showing that, in human breast tissues, entire acini are lost during involution rather than a

percentage of cells from each acini(36). We propose that SEMA7A promotes anoikis resistance,

in part, by signaling to detached cells in a manner similar to matrix molecules, thereby

mimicking attachment. Consistent with this hypothesis, we show that exogenous SEMA7A

increases pAKT in cultured MECs in a manner that is dependent on α6 and β1-integrins, which

is normally the receptor for pro-survival signaling via laminin in the mammary gland(6, 24).

While β1-integrin is a known mediator of SEMA7A signaling in a variety of contexts, few studies

have addressed the alpha integrin that cooperates with β1 to mediate SEMA7A-activated

signaling. Our results presented herein identify, for the first time, that SEMA7A may mimic the

effects of laminin to promote anoikis resistance via α6β1 heterodimers in MECs.

Anoikis resistance is a hallmark of stem cells and breast cancer cells, particularly those that

result in lethal metastasis as these cells must survive in the absence of extracellular matrix

signaling as they travel to and throughout the vasculature to seed distant tumors(37). Since

tumor cells are known to hijack normal developmental programs, we postulated that sustained

expression of SEMA7A in MECs, particularly in postpartum women, could result in pro-tumor

characteristics that are similar to those induced by Ras—a known oncogene. This hypothesis

was based on our recent publication showing that SEMA7A expression is higher in normal

adjacent breast tissue from postpartum women and our identification of a cut-point of

expression that predicts for recurrence in women with PPBC(38). Our hypothesis is supported

by our findings that SEMA7A mediates hallmarks of cellular transformation and stemness in

non-transformed cultured MECs. While others have shown that SEMA7A is required for Ras

dependent induction of EMT(39), we now show that growth in soft agar in Ras transformed

variants of MCF10A cells is dependent on SEMA7A and that SEMA7A can promote growth of

non-transformed cells in soft agar(32, 40, 41). Therefore, we predict that SEMA7A promotes

cellular transformation and tumor initiation in breast cancer patients and, in particular, in PPBC

patients(22, 41). A question that remains unanswered is whether sustained expression of

SEMA7A is causative for malignancy or whether SEMA7A expression in the tumor influences

normal adjacent cells to express SEMA7A thereby promoting tumor progression and

metastasis. Evidence for the latter comes from our studies showing that SEMA7A can affect

multiple aspects of the tumor microenvironment including collagen and fibronectin deposition,

lymphangiogenesis and macrophage infiltration, as well as neighboring cells such as

fibroblasts(40, 41). Additionally, we provide evidence that SEMA7A expressing cells upregulate

drug resistant cell-surface markers(23) and therefore may not respond to conventional

chemotherapy; this, along with our analysis of RFS in a large patient dataset revealing that co-

expression of *ITGA6* and *SEMA7A* significantly influences probability of relapse suggests that

SEMA7A can not only initiate tumor formation, but also influence clinical outcomes. Future

studies will investigate these important questions to identify how to best prevent and treat PPBC

and all patients with α6-integrin+SEMA7A+ breast cancers.

To date, the majority of research into PPBC has focused on how the tissue microenvironment in

the postpartum breast drives metastasis with lesser focus on cellular mechanisms at play in the

MECs that may also support transformation(38). As more than half of all breast cancers in

patients under 40 can be defined as PPBC, which are highly metastatic, understanding the

mechanisms at play in postpartum breast tissue is, thus, critically important for identifying

molecular drivers of PPBC. In this study, we have identified a known tumor-promotional protein,

SEMA7A, as being temporally expressed during the irreversible phase of postpartum mammary

gland involution where it promotes anoikis resistance and we have shown for the first time that

SEMA7A can promote properties of transformation, stemness, and metastasis in non-cancerous

cells.

ACKNOWLEDGEMENTS

This work was supported by NIH/NCI R01 CA211696-01A1 (to T.R. Lyons), NIH/NCI R01CA211696-

02S1(to Taylor Rutherford), ACS RSG-16-171-010CSM (to T.R. Lyons), and Department of

Medicine Outstanding Early Career Scholars Award (to T.R. Lyons). The authors acknowledge the

University of Colorado Cancer Center Flow Cytometry Core for use of equipment and technical

assistance and the University of Colorado Cancer Center Support Grant (P30CA046934). The

authors also thank Alexander Stoller for assistance with animal studies. Contents are the authors'

sole responsibility and do not necessarily represent official NIH views.

The authors declare no potential conflicts of interest.

REFERENCES

1.   Goddard ET, Bassale S, Schedin T, Jindal S, Johnston J, Cabral E, et al. Association Between
Postpartum Breast Cancer Diagnosis and Metastasis and the Clinical Features Underlying Risk. JAMA
Netw Open. 2019;2(1):e186997.
2.   Lund LR, Romer J, Thomasset N, Solberg H, Pyke C, Bissell MJ, et al. Two distinct phases of
apoptosis in mammary gland involution: proteinase-independent and -dependent pathways.
Development. 1996;122(1):181-93.
3.   Schere-Levy C, Buggiano V, Quaglino A, Gattelli A, Cirio MC, Piazzon I, et al. Leukemia Inhibitory
Factor Induces Apoptosis of the Mammary Epithelial Cells and Participates in Mouse Mammary Gland
Involution. Experimental Cell Research. 2003;282(1):35-47.
4.   Gilmore AP. Anoikis. Cell death and differentiation. 2005;12 Suppl 2:1473-7.
5.   Boudreau N, Sympson CJ, Werb Z, Bissell MJ. Suppression of ICE and apoptosis in mammary
epithelial cells by extracellular matrix. Science. 1995;267(5199):891-3.
6.   Pullan S, Wilson J, Metcalfe A, Edwards GM, Goberdhan N, Tilly J, et al. Requirement of
basement membrane for the suppression of programmed cell death in mammary epithelium. Journal of
cell science. 1996;109 (Pt 3):631-42.
7.   Cerny J, Stockinger H, Horejsi V. Noncovalent associations of T lymphocyte surface proteins. Eur
J Immunol. 1996;26(10):2335-43.
8.   Pasterkamp RJ, Peschon JJ, Spriggs MK, Kolodkin AL. Semaphorin 7A promotes axon outgrowth
through integrins and MAPKs. Nature. 2003;424(6947):398-405.
9.   Jaimes Y, Gras C, Goudeva L, Buchholz S, Eiz-Vesper B, Seltsam A, et al. Semaphorin 7A inhibits
platelet production from CD34+ progenitor cells. Journal of thrombosis and haemostasis : JTH.
2012;10(6):1100-8.
10.   Delorme G, Saltel F, Bonnelye E, Jurdic P, Machuca-Gayet I. Expression and function of
semaphorin 7A in bone cells. Biology of the cell / under the auspices of the European Cell Biology
Organization. 2005;97(7):589-97.
11.   Esnault S, Torr EE, Bernau K, Johansson MW, Kelly EA, Sandbo N, et al. Endogenous Semaphorin-
7A Impedes Human Lung Fibroblast Differentiation. PloS one. 2017;12(1):e0170207.
12.   De Minicis S, Rychlicki C, Agostinelli L, Saccomanno S, Trozzi L, Candelaresi C, et al. Semaphorin
7A contributes to TGF-beta-mediated liver fibrogenesis. The American journal of pathology.
2013;183(3):820-30.
13.   Tarullo SE, Hill RC, Hansen K, Behbod F, Borges VF, Nelson AC, et al. Postpartum breast cancer
progression is driven by semaphorin 7a-mediated invasion and survival. Oncogene.
2020;https://doi.org/10.1038/s41388-020-1192-9.
14.   Czopik AK, Bynoe MS, Palm N, Raine CS, Medzhitov R. Semaphorin 7A Is a Negative Regulator of
T Cell Responses. Immunity. 2006;24(5):591-600.
15.   Xie J, Wang H. Semaphorin 7A as a potential immune regulator and promising therapeutic target
in rheumatoid arthritis. Arthritis Res Ther. 2017;19(1):10.

371 16. Elder AM, Tamburini BA, Crump LS, Black SA, Wessells VM, Schedin PJ, et al. Semaphorin 7A
372 promotes macrophage-mediated lymphatic remodeling during postpartum mammary gland involution
373 and in breast cancer. Cancer research. 2018.
374 17. Esnault S, Kelly EA, Johansson MW, Liu LY, Han ST, Akhtar M, et al. Semaphorin 7A is expressed
375 on airway eosinophils and upregulated by IL-5 family cytokines. Clinical immunology. 2014;150(1):90-
376 100.
377 18. Gan Y, Reilkoff R, Peng X, Russell T, Chen Q, Mathai SK, et al. Role of semaphorin 7a signaling in
378 transforming growth factor beta1-induced lung fibrosis and scleroderma-related interstitial lung disease.
379 Arthritis and rheumatism. 2011;63(8):2484-94.
380 19. Lee Y, Blount KL, Dai F, Thompson S, Scher JK, Bitterman S, et al. Semaphorin 7A in circulating
381 regulatory T cells is increased in autosomal-dominant polycystic kidney disease and decreases with
382 tolvaptan treatment. Clin Exp Nephrol. 2018;22(4):906-16.
383 20. Garcia-Areas R, Libreros S, Amat S, Keating P, Carrio R, Robinson P, et al. Semaphorin7A
384 promotes tumor growth and exerts a pro-angiogenic effect in macrophages of mammary tumor-bearing
385 mice. Frontiers in physiology. 2014;5:17.
386 21. Garcia-Areas R, Libreros S, Simoes M, Castro-Silva C, Gazaniga N, Amat S, et al. Suppression of
387 tumor-derived Semaphorin 7A and genetic ablation of host-derived Semaphorin 7A impairs tumor
388 progression in a murine model of advanced breast carcinoma. Int J Oncol. 2017;51(5):1395-404.
389 22. Borges VF, Hu J, Young C, Maggard J, Parris HJ, Gao D, et al. Semaphorin 7a is a biomarker for
390 recurrence in postpartum breast cancer. npj Breast Cancer. 2020;6(1):56.
391 23. Crump LS, Wyatt GL, Rutherford TR, Richer JK, Porter WW, Lyons TR. Hormonal Regulation of
392 Semaphorin 7a in ER(+) Breast Cancer Drives Therapeutic Resistance. Cancer research. 2021;81(1):187-
393 98.
394 24. Gilmore AP, Metcalfe AD, Romer LH, Streuli CH. Integrin-mediated survival signals regulate the
395 apoptotic function of Bax through its conformation and subcellular localization. The Journal of cell
396 biology. 2000;149(2):431-46.
397 25. Krebsbach PH, Villa-Diaz LG. The Role of Integrin alpha6 (CD49f) in Stem Cells: More than a
398 Conserved Biomarker. Stem cells and development. 2017;26(15):1090-9.
399 26. Lyons TR, Borges VF, Betts CB, Guo Q, Kapoor P, Martinson HA, et al. Cyclooxygenase-2-
400 dependent lymphangiogenesis promotes nodal metastasis of postpartum breast cancer. The Journal of
401 clinical investigation. 2014;124(9):3901-12.
402 27. O'Brien J, Lyons T, Monks J, Lucia MS, Wilson RS, Hines L, et al. Alternatively activated
403 macrophages and collagen remodeling characterize the postpartum involuting mammary gland across
404 species. The American journal of pathology. 2010;176(3):1241-55.
405 28. Lee YJ, Streuli CH. Extracellular matrix selectively modulates the response of mammary epithelial
406 cells to different soluble signaling ligands. The Journal of biological chemistry. 1999;274(32):22401-8.
407 29. Farrelly N, Lee YJ, Oliver J, Dive C, Streuli CH. Extracellular matrix regulates apoptosis in
408 mammary epithelium through a control on insulin signaling. The Journal of cell biology.
409 1999;144(6):1337-48.
410 30. Shehata M, Teschendorff A, Sharp G, Novcic N, Russell IA, Avril S, et al. Phenotypic and
411 functional characterisation of the luminal cell hierarchy of the mammary gland. Breast cancer research :
412 BCR. 2012;14(5):R134.
413 31. Visvader JE, Stingl J. Mammary stem cells and the differentiation hierarchy: current status and
414 perspectives. Genes & development. 2014;28(11):1143-58.
415 32. Black SA, Nelson AC, Gurule NJ, Futscher BW, Lyons TR. Semaphorin 7a exerts pleiotropic effects
416 to promote breast tumor progression. Oncogene. 2016;35(39):5170-8.
417 33. Liu R, Chen Y, Liu G, Li C, Song Y, Cao Z, et al. PI3K/AKT pathway as a key link modulates the
418 multidrug resistance of cancers. Cell Death Dis. 2020;11(9):797.

34.   Clarkson RW, Wayland MT, Lee J, Freeman T, Watson CJ. Gene expression profiling of mammary
gland development reveals putative roles for death receptors and immune mediators in post-lactational
regression. Breast cancer research : BCR. 2004;6(2):R92-109.
35.   Crump LS, Wyatt G, Rutherford TR, Richer JK, Porter WW, Lyons TR. Hormonal regulation of
Semaphorin 7a in ER+ breast cancer drives therapeutic resistance. Cancer research. 2020.
36.   Jindal S, Gao D, Bell P, Albrektsen G, Edgerton SM, Ambrosone CB, et al. Postpartum breast
involution reveals regression of secretory lobules mediated by tissue-remodeling. Breast cancer research
: BCR. 2014;16(2):R31.
37.   Cao Z, Livas T, Kyprianou N. Anoikis and EMT: Lethal "Liaisons" during Cancer Progression.
Critical reviews in oncogenesis. 2016;21(3-4):155-68.
38.   Borges VF, Lyons TR, Germain D, Schedin P. Postpartum Involution and Cancer: An Opportunity
for Targeted Breast Cancer Prevention and Treatments? Cancer research. 2020;80(9):1790-8.
39.   Allegra M, Zaragkoulias A, Vorgia E, Ioannou M, Litos G, Beug H, et al. Semaphorin-7a reverses
the ERF-induced inhibition of EMT in Ras-dependent mouse mammary epithelial cells. Molecular biology
of the cell. 2012;23(19):3873-81.
40.   Tarullo SE, Hill RC, Hansen KC, Behbod F, Borges VF, Nelson AC, et al. Postpartum breast cancer
progression is driven by semaphorin 7a-mediated invasion and survival. Oncogene. 2020;39(13):2772-
85.
41.   Elder AM, Tamburini BAJ, Crump LS, Black SA, Wessells VM, Schedin PJ, et al. Semaphorin 7A
Promotes Macrophage-Mediated Lymphatic Remodeling during Postpartum Mammary Gland Involution
and in Breast Cancer. Cancer research. 2018;78(22):6473-85.

FIGURE LEGENDS

Figure 1: Semaphorin-7a promotes anoikis resistance in normal MECs. a. Cleaved

caspase 3/7 expression in MCF10A EV or S7a-OE cells in adherent and forced suspension

cultures, n=3 and **b.** in MCF12A cells +/- exogenous S7a protein and/or ITGB1 blocking

antibody, 9eg7, n=3. **c.** Flow cytometry analysis of pAKT in insulin or S7a-treated MCF12A (left)

with representative flow plots and MCF10A cells (right), n=3. **d.** Flow cytometry for %S7a+

mammary epithelial cells from tissues harvested from nulliparous (N) or recently weaned

C57/BL6 mice (N: n=6, I1: n=5, I2: n=5, I3: n=5, I4: n=6, I6: n=4, I10: n=6, I14: n=4, I28: n=6

glands/group). **e.** Flow cytometry analysis of live epithelial cells as measured by Lin-

Aquazombie-EpCAM+ in mammary tissues harvested from recently weaned WT (I1: n=5, I2:

n=6, I3: n=5, I6: n=4 glands/group) or Sema7a$^{tm1Alk/J}$ (I1-I6: n=6 glands/group) mice . Error bars

indicate mean +/- standard error of the mean. Significance denoted as: p<0.05=*, p<0.01=**,

p<0.001=***, p<0.0001=****.

Figure 2: Loss of SEMA7A promotes histologic features of accelerated involution. Representative 20x images of #3 mammary glands from WT or SEMA7A [tm1Alk/J] mice at 1,2,3,&6 days post-wean. **a.** Hematoxylin and Eosin, **b.** pSTAT3, **c.** cleaved caspase-3 and **d.** perilipin staining.

Figure 3: Loss of SEMA7A promotes alveolar collapse through alterations to cell survival and adipocyte repopulation. Immunohistochemical quantification of mammary glands from WT (I1: n=5, I2: n=6, I3: n=5, I6: n=4 glands/group) or SEMA7A [tm1Alk/J] (I1-I6: n=6 glands/group) mice at 1,2,3,&6 days post-wean. Quantification of **a.** alveolar area **b.** pSTAT3 **c.** cleaved caspase-3 and **d.** perilipin staining. Error bars indicate mean +/- standard error of the mean. Significance denoted as: $p<0.05$=*, $p<0.01$=**, $p<0.001$=***, $p<0.0001$=****.

Figure 4: Semaphorin-7a promotes survival via integrin mediated activation of pAKT. a-e. Flow cytometry analysis of Lin- singlets from #4 mammary glands from WT (left, I1: n=4, I2: n=6, I3: n=5, I6: n=4 glands/group) and SEMA7A[tm1Alk/J] (right, I1-I6: n=6 glands/group) mice. **a.** %MEC subtype of total MECs by EpCAM/CD49f **b.** %EpCAMhiCD49f+ LPCs of Lin-EpCAM+. **c.** %SEMA7A+ of live (Aquazombie-) Lin-EpCAMhiCD49f+ LPCs. **d.** %ITGB1+ITGA6+ of Lin-EpCAM cells. **e.** %SEMA7A of ITGB1+ITGA6+ cells. **f.** CC3/7 expression in MCF12A cells +/- S7a and/or GoH3, n=3. **g.** %pAKT in MCF12A cells +/- S7a and/or GoH3, n=3. Error bars indicate mean +/- standard error of the mean. Significance denoted as: $p<0.05$=*, $p<0.01$=**, $p<0.001$=***, $p<0.0001$=****.

Figure 5: SEMA7A promotes features of transformation and chemoresistance via β1/α6-integrins. a. %Celltrace violet bright of singlets in MCF10A EV and OE cells, n=3. **b,c.** Number of mammospheres formed by MCF12A cells +/- S7a and GoH3 or 9eg7, n=4. **d.** Number of colonies formed in soft agar assay, n=3. **e,f,g.** DCIS KD or OE mammospheres cultured in

paclitaxel +/- 9eg7 or GoH3, n=3. **e**. Number of mammospheres. **f.** %CD44+CD24- of singlets

by flow cytometry. **g.** Representative images, 20x magnification. Error bars indicate mean +/-

standard error of the mean. Significance denoted as: p<0.05=*, p<0.01=**, p<0.001=***,

p<0.0001=****.

Figure 6: High expression of SEMA7A and α6-integrin in breast cancer correlates with

decreased relapse-free survival. a-d. Kaplan-Meier survival curves of relapse-free survival

over 5 years in breast cancer patients. **a.** SEMA7A **b.** α6-integrin **c.** SEMA7A in patients with

low α6-integrin **d.** SEMA7A in patients with high α6-integrin.