Caspase-1-driven neutrophil pyroptosis promotes an incomplete NETosis upon *Pseudomonas aeruginosa* infection

Karin Santoni[1], David Pericat[1], Miriam Pinilla[1], Salimata Bagayoko[1], Audrey Hessel[1], Stephen-Adonai Leon-Icaza[1], Elisabeth Bellard[1], Serge Mazères[1], Emilie Doz-Deblauwe[2], Nathalie Winter[2], Christophe Paget[3], Jean-Philippe Girard[1], Céline Cougoule[1], Renaud Poincloux[1], Mohamed Lamkanfi[4], Emma Lefrançais[1], Etienne Meunier[1,#,¶], Rémi Planès[1,#,¶]

[1] Institute of Pharmacology and Structural Biology (IPBS), University of Toulouse, CNRS, Toulouse; France.

[2] INRAE, Université de Tours, ISP, Nouzilly, France.

[3] Institut national de la santé et de la recherche médicale, Centre d'Etude des Pathologies Respiratoires, UMR 1100, Tours, France.

[4] Department of Internal Medicine and Pediatrics, Ghent University, Ghent, Belgium.

[#] Address correspondence to: etienne.meunier@ipbs.fr (EM) and Remi.planes@ipbs.fr (RP)

Present Address: Institute of Pharmacology and Structural Biology (IPBS), University of Toulouse, CNRS, Toulouse; France

[¶] These authors equally supervised this work.

Abstract

Multiple neutrophil death programs contribute to host defense against infections. Although expressing all necessary components, neutrophils specifically fail to undergo pyroptosis, a lytic form of cell death triggered by the activation of the pro-inflammatory complex inflammasome. In the light of the arm race, we hypothesized that intrinsic neutrophil pyroptosis resistance might be bypassed in response to specific microbial species. We show that *Pseudomonas aeruginosa* (*P. aeruginosa*) stimulates Caspase-1-dependent pyroptosis in human and murine neutrophils. Mechanistically, activated NLRC4 inflammasome supports Caspase-1-driven Gasdermin-D (GSDMD) activation, IL-1β cytokine release and neutrophil pyroptosis. Furthermore, GSDMD activates Peptidyl Arginine Deaminase-4 which drives an "incomplete NETosis" where neutrophil DNA fills the cell cytosol but fails crossing plasma membrane. Finally, we show that neutrophil Caspase-1 account for IL-1β production and contributes to various *P. aeruginosa* strains spread in mice. Overall, we demonstrate that neutrophils are fully competent for Caspase-1-dependent pyroptosis, which drives an unsuspected "incomplete NETosis".

Summary

Neutrophils play an essential roles against infections. Although multiple neutrophil death programs contribute to host defense against infections, they fail to undergo pyroptosis, a pro-inflammatory form of cell death. Upon Infections, pyroptosis can be induced in macrophages or epithelial cells upon activation of pro-inflammatory complexes, inflammasomes that trigger Caspase-1-driven Gasdermin dependent plasma membrane lysis. In the light of host-microbe interactions, we hypothesized that yet to find microbial species might hold the capacity to overcome neutrophil resistance to inflammasome-driven pyroptosis. Among several bacterial species, we describe that the bacterium *Pseudomonas aeruginosa* specifically engages the NLRC4 inflammasome, which promotes Caspase-1-dependent Gasdermin-D activation and subsequent neutrophil pyroptosis. Furthermore, inflammasome-driven pyroptosis leads to DNA decondensation and expansion into the host cell cytosol but not to the so called Neutrophil Extracellular Trap (NET) release as DNA fails breaching the plasma membrane. Finally, *in vivo P. aeruginosa* infections highlight that Caspase-1-driven neutrophil pyroptosis is functional and is detrimental upon *P. aeruginosa* infection. Altogether, our results unexpectedly underline neutrophil competence for

57 Caspase-1-dependent pyroptosis, a process that contributes to host susceptibility to

58 *P. aeruginosa* infection.

Introduction

Over the last 30 years, non-apoptotic forms of cell death have emerged as crucial processes driving inflammation, host defense against infections but also (auto) inflammatory pathologies [1].

Unique among all forms of regulated cell necrosis is the capacity of granulocyte neutrophils to undergo the process of NETosis [2]. NETosis is an antimicrobial and pro-inflammatory from of cell death that promotes the formation of extracellular web-like structures called Neutrophil Extracellular Traps (NETs) [2]. Although the importance of NETosis in host immunity to infections has been well established [2–5], NETosis dysregulation also associates to autoimmunity, host tissue damages, aberrant coagulation and thrombus that all contribute to pathology such as sepsis or autoimmune lupus [6–13].

NETosis consists in sequential steps that start with nuclear envelope disintegration, DNA decondensation, cytosolic expansion of DNA and its subsequent expulsion through plasma membrane [14]. Completion of DNA decondensation and expulsion requires various cellular effectors. Among them, neutrophil serine proteases (Neutrophil elastase, Cathepsin G, Proteinase 3) or Caspase-11 can mediate histone cleavage, which relaxes DNA tension [3,13,15–17]. In addition, granulocyte-enriched Protein arginine deaminase 4 (PAD4), citrullinates histone-bound DNA, which neutralizes arginine positive charges, thus helping nuclear DNA relaxation and decondensation [4,18,19]. Then, decondensed DNA is mixed with the neutrophil cytoplasmic granule content such as NE, CathG, PR3 and Myeloperoxidase (MPO) proteins [3,15,18]. Finally, sub-cortical actin network disassembly is required to ensure efficient DNA extrusion through the permeabilized plasma membrane [18,20].

Depending on the initial trigger, various signaling pathways such as calcium fluxes [17,18], necroptosis-associated MLKL phosphorylation [21], ROS-induced Neutrophil protease release [15] or endotoxin-activated Caspase-11 [3,5,22] all bring neutrophil into NETosis. Common to both ROS- and Caspase-11-dependent NETosis is the requirement of the pyroptosis executioner Gasdermin-D (GSDMD) cleavage by both neutrophil serine and Caspase-11 proteases, which triggers neutrophil NETosis [3,16]. Specifically, active GSDMD forms a pore on PIP2-enriched domains of the plasma and

nuclear membrane of neutrophils, which ensures both IL-1-related cytokine release

[23–26] and osmotic imbalance-induced DNA decondensation and expulsion [3,16].

An intriguing feature of neutrophils is that, despite GSDMD activation, they resist

canonical inflammasome-induced Caspase-1-dependent cell pyroptosis upon

Salmonella Typhimurium and *Burkholderia thaïlandensis*-activated NLRC4

inflammasome or upon Nigericin/ATP-mediated NLRP3 inflammasome activation

[3,5,27,28]. However, recent studies indirectly challenged canonical pyroptosis

impairment in neutrophils by showing that sterile activators of the NLRP3

inflammasome also contribute to canonical inflammasome-dependent neutrophil death

and subsequent NETosis [29,30]. If it does exist specific microbial species that can

promote neutrophil canonical pyroptosis has remained an open question so far.

Specifically lung infections triggered by the bacterium *Pseudomonas aeruginosa* (*P.*

aeruginosa) can promote acute and chronic life-threatening infections in

immunocompromised or hospitalized patients [31]. *P. aeruginosa* strains express a

Type-3 Secretion System (T3SS) that allows injecting a specific set of virulence factors

into host target cells, including macrophages and neutrophils [32]. T3SS-expressing

Pseudomonas aeruginosa strains classically segregate into two mutually exclusive

clades. Those expressing the bi-cistronic ADP-rybosylating and GTPase Activating

Protein (GAP) virulence factor ExoS, and those expressing the lytic phospholipase of

the patatin-like family, ExoU [32]. Common to most of *P. aeruginosa* strains is the

expression of two other toxins, ExoY and ExoT whose functions in bacterial infection

remain still unclear. All Exo toxins are injected by the T3SS into host target cells upon

infections. Finally *P. aeruginosa* strains also use their T3SS to inject Flagellin into host

target cells, which promotes the activation of the NAIP5-NLRC4 inflammasome and

subsequent Caspase-1-driven GSDMD-dependent pyroptosis of macrophages [33–

39]. Although numerous studies underlined that neutrophils are targeted by

Pseudomonas aeruginosa virulence factors, which could promote NETosis [12,40–42],

the critical effectors and their host cell targets remain extensively debated. Regarding

this, defect in the Nadph oxidase (Nox2) enzyme expression has also been found to

sensitive murine neutrophils to some degree of Caspase-1-driven neutrophil death

upon infection with *Pseudomonas aeruginosa* [43]. This suggests that under certain

conditions neutrophils might be prone to undergo Caspase-1-dependent pyroptosis.

Whether such process also occurs in WT neutrophils, and its molecular as well as
immune significance remain unknown.

Here, we aimed at determining if *Pseudomonas aeruginosa* but also other bacterial
species could bypass neutrophil resistance to canonical pyroptosis induction as well
as the key host and microbial effectors involved. We describe that the bacterial
pathogen *Pseudomonas aeruginosa* uniquely triggers measurable Caspase-1-
dependent pyroptosis in WT human and murine neutrophils. This requires the
expression of a functional Type-3 Secretion System (T3SS) and Flagellin, but not other
T3SS-derived toxins. Noticeable, deletion of Exotoxins U or S in *P. aeruginosa* entirely
rewires neutrophil death towards a Caspase-1-driven pyroptosis. Specifically, *P.*
aeruginosa activates the neutrophil NLRC4 inflammasome, which ensures Caspase-
1-driven Gasdermin D (GSDMD) cleavage and the formation of GSDMD pores.
GSDMD pores promote neutrophil pyroptosis and subsequent Peptidyl Arginine
Deaminase 4 (PAD4) activation. PAD4 Citrullinates Histones, which leads to an
incomplete NETosis where neutrophil DNA is decondensed and fills the host cell
cytosol but is not expulsed out from the cells. Finally, using intravital microscopy on
MRP8-GFP mice, we show that this incomplete NETosis occurs in neutrophils in
infected lungs and that neutrophil Caspase-1 contributes both to IL-1β production and
to the spread of several *P. aeruginosa* strains in mice. Overall, our results highlight
that Caspase-1-dependent pyroptosis is a functional process in neutrophils which
promotes a unique "incomplete NETosis".

Results

Various *Pseudomonas aeruginosa* strains trigger Caspase-1-dependent and -independent neutrophil lysis

In order to determine whether neutrophils could undergo pyroptosis upon bacterial infections, we first infected WT and *Caspase-1$^{-/-}$* (*Casp1$^{-/-}$*) mouse Bone Marrow Neutrophils (BMNs) with various bacterial strains, know to activate different inflammasomes in macrophages (**Fig. 1A**). We measured neutrophil lysis (LDH release) as well as IL-1β release as a hallmark of inflammasome activation. At the exception of *Staphylococcus aureus*, all bacteria triggered a Caspase-1-dependent IL-1β release, suggesting that an inflammasome-dependent response was induced in neutrophils upon various bacterial challenge (**Fig. 1A**). However, despite inducing significant neutrophil lysis, only *Pseudomonas aeruginosa* infection triggered a robust Caspase-1-dependent lysis (referred as pyroptosis) (**Fig. 1A**). Furthermore, infection of human blood neutrophils with various *Pseudomonas aeruginosa* strains (PAO1 and CHA) also triggered Caspase-1-dependent IL-1B and lysis, thus suggesting that both murine and human neutrophils are competent for Caspase-1-driven pyroptosis in response to specific bacterial species (**Fig. 1B**).

Next, we aimed at determining the means by which *P. aeruginosa* could promote Caspase-1-dependent neutrophil pyroptosis. As *P. aeruginosa* T3SS-mediated injection of toxins (ExoS, U, T, Y) and Flagellin play a major role in virulence, we hypothesized that one or many of those virulence factors might contribute to neutrophil pyroptosis. Hence, we infected murine BMNs with *P. aeruginosa* strains lacking or not expression of each toxin (PAO1^{ExoS-}, PAO1^{ExoT-}, PAO1^{ExoY-}, PAO1^{FliC-}) or deficient for the expression of the T3SS (PAO1^{ExsA-}). We measured the ability of WT and *Casp1$^{-/-}$* neutrophils to undergo lysis (LDH release), to promote IL1β release or to exhibit plasma membrane permeabilization (SYTOX Green incorporation) upon infections with those *Pseudomonas aeruginosa* mutants.

P. aeruginosa strains lacking expression of T3SS or Flagellin were unable to promote a robust Caspase-1-dependent neutrophil lysis, IL-1β release and plasma membrane permeabilization, suggesting that T3SS and Flagellin are major effectors of Caspase-1-driven neutrophil death (**S1A Fig, Fig. 1C, D**).

To the contrary, strains lacking ExoY or T did not influence significantly neutrophil lysis, IL-1β release and plasma membrane permeabilization (**S1A Fig, Fig. 1C, D**). We also

noticed that the infection of neutrophils with *P. aeruginosa* deficient for ExoS triggered
increased lysis, membrane permabilization and IL-1β release of neutrophils (**S1A Fig,**
Fig. 1C, D). In addition, PAO1^{ExoS-} -induced neutrophil lysis was completely abrogated
in Caspase-1 deficient neutrophils, hence suggesting that ExoS expression might
promote a Caspase-1-independent neutrophil death (**S1A Fig, Fig. 1C, D**). Caspase-
1 is a central effector of macrophage lysis through the cleavage of the pyroptosis
executioner Gasdermin D (GSDMD). In this context, over time measures showed that
both *Casp1$^{-/-}$* and *GsdmD$^{-/-}$* neutrophils exhibited reduced LDH and IL-1β release upon
Pseudomonas aeruginosa infection (**Fig. 1E**). Again, we remarked that PAO1^{ExoS-}-
induced neutrophil lysis and IL-1β release were fully CASP1- and GSDMD- dependent
(**Fig. 1E**).
As 30% of *Pseudomonas aeruginosa* strains do not express ExoS but instead express
the extremely lytic phospholipase toxin ExoU, we also challenged our findings by
infecting neutrophils with ExoU-expressing strains (PP34^{ExoU}). We observed no defect
of *Casp1$^{-/-}$* neutrophil lysis capacity upon PP34^{ExoU} infection (**S1B, C Fig.**). However,
infection of neutrophils in presence of MAFP, an inhibitor of ExoU lytic activity or with
PP34^{ExoU-} or strains expressing a catalytically dead mutant of ExoU (PP34^{ExoUS142A})
showed a fully Caspase-1-dependent neutrophil lysis, plasma membrane
permeabilization and IL-1β release both in human and murine BMNs (**S1B, C Fig.**).
This, suggests that in absence of ExoU, *P. aeruginosa* also triggers neutrophil death
in a fully Caspase-1 and GSDMD-dependent manner.
Altogether, our results show the unexpected ability of Caspase-1 to promote GSDMD-
dependent neutrophil lysis upon *Pseudomonas aeruginosa* infection.
249
***P. aeruginosa* infection engages a canonical NLRC4-Caspase-1-Gasdermin D-**
dependent pyroptosis axis in neutrophils.
Next, we investigated the molecular pathways by which *P. aeruginosa* promoted
Caspase-1-dependent neutrophil death. The infection of WT murine neutrophils or
deficient for various inflammasome sensors showed that only *Nlrc4$^{-/-}$* and *ASC$^{-/-}$* BMNs
had resistance to cell lysis, IL-1β release, Caspase-1 (p20) and GSDMD (p30)
processing in response to multiple strains of *P. aeruginosa* (PAO1, PAO1^{ExoS-},
PP34^{ExoUS142A}, CHA, CHA^{ExoS-}, PP34$^{ExoU-,}$ PA14^{ExoU-}) (**Fig. 2A-E, S2A, B Fig.**). Again,
PAO1^{ExoS-} and PP34^{ExoU-} strains triggered exacerbated neutrophil pyroptosis, IL1β

release but also Caspase-1 and GSDMD processing, a process that required NLRC4
expression (**Fig. 2A-E, S2A, B Fig.**).

Neutrophils resist NLRC4/Caspase-1-dependent pyroptosis upon *Salmonella* infection
[28]. Hence, we next infected WT or *Nlrc4*[-/-] BMNs with various bacteria (*Salmonella*
Typhimurium, *Shigella flexnerii*, *Chromobacter violaceum*, *Burkholderia thailandensis*)
known to trigger a NLRC4 inflammasome response and monitored for cell death and
IL-1β release (**Fig. 3E**) [5,28,44–46]. None of the tested bacteria triggered a significant
NLRC4-dependent neutrophil lysis of murine neutrophils although they promoted
NLRC4-dependent IL-1β release and Gasdermin D processing (**S2C Fig.**).

Finally, to determine if *P. aeruginosa*-induced neutrophil NLRC4 inflammasome
activation also occurs *in vivo*, we infected ASC-Citrine mice with low doses (1.10^5
CFUs) of *P.* aeruginosa strains that triggered specific NLRC4-dependent neutrophil
pyroptosis, namely PP34^{ExoU-} or its isogenic mutant PP34$^{ExoU-/FliC-}$, deficient for the
expression of Flagellin (**Fig. 2D, S2D Fig.**). ImageStreamX observation of neutrophils
presenting an active ASC supramolecular speck (ASC speck$^+$/LY6G$^+$ neutrophils)
showed that PP34^{ExoU-} infection triggered inflammasome activation in neutrophils,
which was reduced when mice where infected with PP34$^{ExoU-/FliC-}$ (**Fig. 2D, S2D Fig.**).
Altogether, our results show that the NLRC4/CASP1/GSDMD axis is fully functional to
promote neutrophil pyroptosis in response to *P. aeruginosa* but not to various other
NLRC4-activating bacteria.

Caspase-1-induced neutrophil pyroptosis promotes PAD4-dependent
incomplete NETosis

Next, we sought to determine whether Caspase-1-induced neutrophil pyroptosis could
also lead to NETosis upon *P. aeruginosa* infection. We infected WT, *Casp1*[-/-] or
GsdmD[-/-] murine BMNs with PAO1, PAO1^{ExoS-}, PP34^{ExoU} or PP34^{ExoU-} strains and
monitored for the presence of NETs using Scanning Electron Microscopy (SEM) (**Fig.**
3A). PP34^{ExoU}, and to a lower extend PAO1, induced NETs in WT*, Casp1*[-/-] and
GsdmD[-/-] neutrophils (**Fig. 3A**). However, we observed that neutrophil pyroptosis
induced by PAO1^{ExoS-} and PP34^{ExoU-} strains failed to induce NETs (**Fig. 3A, B, S3A**
Fig.). Rather, SEM and immunofluorescence experiments revealed that the fully
pyroptotic strains of *P. aeruginosa* (PAO1^{ExoS-}, PP34^{ExoU-}, PP34^{ExoUS142A}) triggered
efficient DNA decondensation as well as exit from the nuclear envelop (Lamin-B1
staining) but no or few DNA release from the neutrophil plasma membrane of BMNs

exhibiting an active inflammasome complex (referred as ASC specks, ASC⁺) (**Fig. 3A,**

B, S3A Fig.). Further experiments using time lapse fluorescent microscopy on ASC

Citrine neutrophils infected with the pyroptotic strain PP34^{ExoU-} or the NETotic strain

PP34^{ExoU+} showed that both bacteria triggered efficient neutrophil DNA

decondensation (**Fig. 3C, S1, 2 Movies**). Yet, pyroptotic neutrophils uniquely failed to

complete DNA release out from the plasma membrane (stained with WGA) (**Fig. 3C**).

NETosis drives the release of DNA and bound Histones outside from neutrophils.

Hence, we reasoned that during pyroptosis, neutrophils might keep intracellularly

Histone-bound DNA. Hence, immunoblotting of Histones in various neutrophil fractions

(soluble, insoluble and supernatant) showed that PP34^{ExoU+}-induced NETosis

efficiently promoted Histone release in the extracellular medium (**Fig. 3D**). However,

neither PAO1^{ExoS-} nor PP34^{ExoU-} induced Histone release, although they efficiently

promoted the release of intracellular soluble and insoluble components such as

GAPDH, the nuclear membrane structural component Lamin B1, the nuclear alarmin

HMGB1 or NLRC4 (**Fig. 3D, S3B Fig.**). Importantly such process required NLRC4

expression (**Fig. 3D, S3B Fig.**). This suggests that Caspase-1-induced neutrophil

pyroptosis can specifically promote an "incomplete NETosis" process where both DNA

decondensation and release from the nuclear membrane occur, but in which DNA does

not breach plasma membrane of neutrophils.

Next, we wondered about the mechanisms by which Caspase-1 promotes incomplete

NETosis. Among others, Histone degradation and DNA citullination are two conserved

mechanisms that promote DNA relaxation and delobulation. Neutrophil elastase and

Caspase-11 promote Histone degradation and PAD4 triggers Histone citrullination.

Therefore, we first explored whether Caspase-1-induced neutrophil DNA delobulation

and release required PAD4-dependent citrullination. Immunoblotting and microscopy

analysis of Histone citrullination showed that PP34^{ExoUS142A} and PAO1^{ExoS-} induced a

robust Histone3-Citrullination in a NLRC4-, GSDMD- and CASP-1-dependent manner

(**Fig. 3E, S3C-E Fig.**), a process that was also seen in human blood neutrophils (**S3C**

Fig.). In order to control the specific involvement of NLRC4 at regulating Histone

Citrullination in neutrophils, we used Ionomycin, a known inducer of NETosis, and

observed that Ionomycin triggered NLRC4-independent Histone Citrullination in

neutrophils (**S3E Fig.**). Conversely, ASC-Citrine BMNs revealed that

NLRC4/CASP1/GSDMD-induced DNA decondensation required PAD4 as

pharmacological inhibition of PAD4 (GSK484) abrogated both Histone citrullination as

well as the DNA nuclear release (**Fig. 3E**). In addition, measure of ASC specks (ASC$^+$), cell lysis (LDH release) in ASC-Citrine, WT, *Pad4$^{-/-}$* and *Nlrc4$^{-/-}$* BMNs highlighted that PAD4 was not involved in PP34^{ExoUS142A}-induced NLRC4 inflammasome activation (**Fig. 3E, S3F Fig.**).

Finally, to determine if Caspase-1-induced neutrophil lysis and PAD4-dependent DNA decondensation plays a microbicidal function, we infected WT, *Pad4$^{-/-}$* and *Nlrc4$^{-/-}$* BMNs with various *Pseudomonas* strains and evaluated their cell-autonomous immune capacities. *Nlrc4$^{-/-}$* BMNs had improved ability to restrict PAO1, PAO1^{ExoS-} and PP34^{ExoUS142A} infection than WT and *Pad4$^{-/-}$* neutrophils (**S3F Fig.**), suggesting that neutrophil pyroptosis more than PAD4-driven DNA decondensation promotes neutrophil failure to restrict PAO1, PAO1^{ExoS-} and PP34^{ExoUS142A}.

All in one, our results describe a pathway where NLRC4-induced neutrophil pyroptosis generates PAD4-dependent intracellular but not extracellular DNA structures.

Neutrophil Caspase-1 contributes to IL-1β production and to *Pseudomonas aeruginosa* spread in mice

Our results showed that various *P. aeruginosa* strains trigger a Caspase-1-dependent neutrophil pyroptosis *in vitro* (**Fig. 2A**). In this context, we aimed at understanding the specific role of neutrophil Caspase-1 upon *P. aeruginosa* infection in mice. First, to determine if neutrophils can undergo Caspase-1-dependent incomplete NETosis in mice, we infected MRP8-GFP$^+$ (granulocytes, including neutrophils express GFP) mice and monitored for the granulocyte death features using intravital microscopy (**Fig. 4A, S3 Movie**). Although necrotic granulocytes exhibited NETotic features (e.g. extracellular DNA) upon exposure to PP34^{ExoU}, PP34^{ExoUS142A} infection led to the appearance of swelled-round necrotic granulocytes that exhibited intracellular decondensed DNA, similarly to what we observed *in vitro* (**Fig. 4A, S1, 2 Movies**). This suggests that upon lung infection, Caspase-1-induced neutrophil pyroptosis is well occurring and displays morphological and immunological distinct characteristics to NETs.

Next, we sought to determine the importance of neutrophil Caspase-1 in response to *Pseudomonas aeruginosa* infection. Thus, we infected mice lacking CASP1 expression in the granulocytic compartment (MRP8^{Cre+}*Casp1^{flox}*) and their respective controls (MRP8^{Cre-}*Casp1^{flox}*) mice either intranasally or systemically with ExoS- or ExoU-expressing *Pseudomonas aeruginosa* (respectively PAO1^{ExoS} and PP34^{ExoU}) or

with their isogenic mutants PAO1^{ExoS-} and PP34^{ExoU-}, both triggering a complete Caspase-1-dependent neutrophil death *in vitro*. We observed that, upon lung infections with PP34^{ExoU}, MRP8^{Cre+}*Casp1*^{flox} mice did not show any differences in bacterial elimination, IL1β production, confirming previous work that *ExoU*-expressing *Pseudomonas* promote successful infection independently of the inflammasome pathways (**Fig. 4B-D**). To the contrary, MRP8^{Cre+}*Casp1*^{flox} mice infected with PAO1^{ExoS}, showed a slight but significant improved bacterial elimination in Bronchoalveolar Fluids (BALF) and lungs, a phenotype that was further amplified upon infection with PAO1^{ExoS-} or PP34^{ExoUS142A} pyroptotic strains (**Fig. 4B-G**). Furthermore, IL-1β levels in BALFs were decreased in MRP8^{Cre+}*Casp1*^{flox} mice infected with PAO1^{ExoS}, PAO1^{ExoS-} or PP34^{ExoUS142A}, hence suggesting that neutrophil Caspase-1 is also a contributor of IL-1β production upon PAO1^{ExoS}, PAO1^{ExoS-} or PP34^{ExoUS142A} infections (**Fig. 4D, G**).

Altogether, our results highlight that neutrophil Caspase-1 activity contributes to both IL-1β release and spreading of several *P. aeruginosa* strains.

Discussion

Our study initially aimed at determining if neutrophil resistance to undergo Caspase-1-dependent pyroptosis can be overcome upon bacterial infection. Screening various inflammasome-activating bacteria [5,27,28], we found that *Pseudomonas aeruginosa* successfully trigger murine and human neutrophil pyroptosis through the engagement of the fully competent canonical NLRC4 inflammasome, process that requires T3SS-dependent injection of bacterial Flagellin. Although those results show a clear ability of neutrophils to undergo Caspase-1-dependent pyroptosis upon *Pseudomonas aeruginosa* infection, our investigations performed with mice specifically lacking Caspase-1 in the granulocyte compartment do suggest a minor contribution of Caspase-1 in favoring *Pseudomonas aeruginosa* spread. Related to this, previous studies showed a robust contribution of NLRC4 at promoting *Pseudomonas aeruginosa* spread in various organs, which suggests that macrophages or other NLRC4-expressing cells are stronger contributors of mouse susceptibility to *P. aeruginosa* [47,48].

Although neutrophils show intrinsic resistance to NLRC4-dependent pyroptosis (e.gs. ESCRT machinery, Caspase-1 expression levels, Ragulator pathway) [3,49–51], the unique ability of *P. aeruginosa* strains and isogenic mutants (PAO1, CHA, PAO1^{ExoS-}, PP34^{ExoU-}) to trigger neutrophil pyroptosis suggests that several neutrophil factors could restrict the ability of other bacteria to promote NLRC4-dependent pyroptosis. Seminal study from Zynchlynsky and colleagues found that neutrophil serine proteases could degrade the Type-3 Secretion System and flagellin virulence factors of *Shigella flexneri* [52], hence limiting their ability to hijack the neutrophil autonomous immunity and restraining *Shigella*-induced neutrophil necrosis. Similarly, upon *P. aeruginosa* infection, mouse neutrophils deficient for the NADPH oxidase enzyme Nox2 undergo increased Caspase-1-dependent pyroptosis [43]. Supporting this, Warnatsch et al., could link the extracellular Oxygen Reactive Species (ROS) localization in neutrophils exposed to *Candida albicans* to an exacerbated IL-1β production and Caspase-1 activation whereas intracellular ROS had an inhibitory effect on IL1β production and Caspase-1 activity in neutrophils [53]. Whether this could explain the capacity of the extracellularly-adapted *Pseudomonas aeruginosa* to specifically promote Caspase-1-dependent neutrophil pyroptosis but not intracellular-adapted bacterial pathogens such as *Shigella* or *Salmonella* will require further investigations.

Striking to us was the observation that expression of the key virulence factors ExoS or
ExoU strongly influences neutrophils to go into NETosis whereas strains lacking ExoU
or ExoS induced a complete rewiring of neutrophil death toward a Caspase-1-driven
way. We hypothesize that the potent cytotoxic effect of these toxins towards various
cell types, including neutrophils, may overcome inflammasome detection of
Pseudomonas aeruginosa and triggers other neutrophil death programs [41,42,54–
56]. Another non mutually exclusive guess is that these toxins directly interfere with
the activation of the inflammasome pathway, thus removal of such toxins leads to an
exacerbated inflammasome-response as previously reported for ExoU [39] and ExoS
[57].

Although NLRC4 activation leads to Caspase-1 dependent neutrophil pyroptosis, in a
pathway similar to what was previously reported in macrophage [39,48], neutrophil
pyroptosis exhibits a unique feature characterized by nuclear membrane rupture, DNA
decondensation and expansion within cell cytosol. One key enzyme responsible for
this morphological characteristic of neutrophil pyroptosis is the Protein arginine
deiminase 4 (PAD4). Indeed, similar to the process induced by various NETosis
inducers, Caspase-1 also promoted PAD4-dependent Histone Citrullination, which
stimulated DNA relaxation and release from the nucleus but, surprisingly, not its
extracellular expulsion. Why upon Caspase-11 [3], MLKL [21], NADPH [17] or
NE/CatG/Pr3 [15] stimulation but not upon Caspase-1 activation neutrophils generate
two different types of DNA structures remains yet to be investigated. Recently, Thiam
et al., [18] observed that pharmacological stabilization of F-actin allowed the
development of this "incomplete NETosis" upon Ionomycin-exposure. Interestingly,
neutrophil elastase has also been shown to degrade actin [58], hence ensuring
complete NETosis process. This, suggests that efficient actin degradation and/or
depolymerization may be an essential player of extracellular DNA release, which would
imply that the final step of NETosis might actually be a cell-regulated process involving
various controllers [14,20]. Interestingly, Chen and colleagues recently found that upon
infection with *Yersinia*, murine neutrophils induce a pyroptotic program that involves
virulence-inhibited innate immune sensing, hence promoting RIPK1-induced Caspase
3-dependent Gasdermin E cleavage and activation and pyroptosis [59], a process that
does not trigger NETosis or "incomplete NETosis". This further suggests that multiple
Gasdermins can trigger neutrophil death through multiple molecular pathways and
promote different morphological outcomes of neutrophils.

Regarding the immunological purpose of Caspase-1-induced neutrophil pyroptosis, we hypothesize that the decondensation of DNA but its conservation into the intracellular space might be a physical mean for neutrophils to trap some intracellular DAMPs, hence avoiding their passive release and a too strong exacerbation of the inflammatory response. Supporting this, we observed that DNA-bound Histones mostly remain trapped intracellularly, but not HMGB1 alarmin, both initially located in the nucleus. In the light of the recent discovery from Kayagaki and colleagues on the role for Ninjurin-1 at promoting active cell shrinkage and HMGB1/nucleosome DAMP release downstream of Gasdermin-D pores in macrophages, the use of Ninjurin-1 deficient mice are full of promises [60]. Another theoretical purpose of neutrophil "incomplete NETosis" could be the generation of "Intracellular Traps" following the "Pyroptosis-induced Intracellular Traps" model in macrophages described by Miao and colleagues. Following such speculative model, intracellular pathogens, in addition to intracellular toxic DAMPs (e.g. Histones, DNA) might remain intracellularly, hence limiting both microbial spread and pathological DAMP-dependent sterile inflammation.

All in one, our results unveil an unsuspected ability of neutrophils to undergo Caspase-1-dependent pyroptosis upon *Pseudomonas aeruginosa* infection which drives an intriguing "incomplete NETosis", hence expanding the spectrum of neutrophil death mechanisms.

ACKNOWLEDGEMENTS

$Nlrc4^{-/-}$ mice were provided by Clare E. Bryant [61] and generated by Millenium Pharmaceutical, $GsdmD^{-/-}$ mice [62] came from P. Broz (Univ of Lausanne, Switzerland), $Casp11^{-/-}$ and $Casp1^{-/-}/ Casp11^{-/-}$ came from B. Py (ENS Lyon, France) and Junying Yuan (Harvard Med School, Boston, USA) [63,64]. Virginie Petrilli (ENS Lyon, France) provided $Nlrp3^{-/-}$ mice that were generated by Fabio Martinon [65]. Thomas Henry (CIRI, Lyon, France) provided $ASC^{-/-}$ and $AIM2^{-/-}$ mice upon agreement with Genentech (San Francisco, Roche, USA) and. ASC-Citrine (#030744) and $Pad4^{-/-}$ (#030315) mice came from Jaxson Laboratory (USA) and were generated by Douglas T Golenbock (University of Massachusetts Medical School, USA) and Kerri Mowen (The Scripps Research Institute, USA) respectively. MRP8^{Cre}/$Casp1^{flox}$ mice are provided by Natalie Winter (INRAE Tours Nouzilly, France) and were generated by crossing MRP8^{Cre} (Jackson # 021614) mice with $Caspase1^{flox}$ mice generated by Mohamed Lamkanfi (Univ. of Ghent, Belgium)[66]. MRP8^{CreGFP} and mTmG mice were

obtained from Jackson laboratories and generated respectively by Emmanuelle Passegue (UCSF, USA) and Liqun Luo, (Stanford University, USA). *Pseudomonas aeruginosa* strains were a kind gift of Ina Attrée (CNRS, Grenoble, France) and Julien Buyck (Univ. of Poitiers, France). Authors also acknowledge the animal facility and Cytometry/microscopy platforms of the INFINITY, CBI and IPBS institutes and particularly Valerie Duplan-Eche for Imagestream acquisition and analysis. This project was funded by grants from the Fonds de Recherche en Santé Respiratoire - Fondation du Souffle to EL, ATIP-Avenir program, FRM "Amorçage Jeunes Equipes" (AJE20151034460) and the ERC (StG INFLAME 804249) to EM, the European Society of Clinical Microbiology and Infectious Diseases (ESCMID, 2020) to RP, Invivogen-CIFRE collaborative PhD (to MP) and post-doctoral fellowships (to RP) and from Mali and Campus France cooperative agencies to SB. Funders had not interest and regard in the conduct of the project.

AUTHOR CONTRIBUTIONS

RP and EM designed the experiments and acquired funding. RP, KS and EM wrote the manuscript. RP and KS performed the experiments with the help of DP, MP, SB, PJB, AH, SALI, CC. Specifically RP and RP performed SEM experiments, SM, EB and EL set up and performed intravital mouse experiments, JPG, ML and NW provided essential reagents, tools and inputs for the conduct of the project. EM, RP and KS supervised the entire study.

CONFLICT OF INTEREST

Authors have no conflict of interest to declare.

 Figure legends

Figure 1. Various *Pseudomonas aeruginosa* strains trigger Caspase-1-dependent and -independent neutrophil lysis

A. Measure of cell lysis (release of LDH) and IL-1β release in WT or *Casp1*^{-/-} murine Bone Marrow Neutrophils (BMNs) infected for 3 hours with various bacteria at a multiplicity of infection (MOI) of 10. ***p ≤ 0.001, T-test with Bonferroni correction. Values are expressed as mean ± SEM. Graphs show one experiment representative of three independent experiments.

B. Measure of cell lysis (release of LDH) and IL-1β release in human blood neutrophils infected for 3 hours with *Pseudomonas aeruginosa* strains PAO1 or CHA (MOI 5) in presence/absence of various Caspase inhibitors, Z-VAD (pan Caspase, 20µM), Z-YVAD (Casp1 inhibitor, 40µM), Z-DEVD (Casp3 inhibitor, 40µM), Z-IETD (Casp8 inhibitor, 40µM). ***p ≤ 0.001, T-test with Bonferroni correction. Values are expressed as mean ± SEM. Graphs show one experiment representative of three independent experiments.

C, D. Measure of plasma membrane permeabilization and associated fluorescent microscopy images over time using SYTOX Green incorporation in WT or *Casp1$^{-/-}$* BMNs infected with *Pseudomonas aeruginosa* PAO1 or various isogenic mutants lacking T3SS-derived toxins ExoS, ExoY, ExoT or Flagellin (FliC) (PAO1^{ExoS-}, PAO1^{ExoY-}, PAO1^{ExoT-}, PAO1^{FliC-}). ***p ≤ 0.001, T-test with Bonferroni correction. Values are expressed as mean ± SEM. Graphs show one experiment representative of three independent experiments. Scale Bar: 150µm

E. Measure of cell lysis (release of LDH) and IL-1β release in WT, *Casp1$^{-/-}$* and *GsdmD$^{-/-}$* murine Bone Marrow Neutrophils (BMNs) infected for the indicated times with PAO1 or its isogenic mutant PAO1^{ExoS-} at an MOI of 10. ***p ≤ 0.001, T-test with Bonferroni correction. Values are expressed as mean ± SEM. Graphs show one experiment representative of three independent experiments.

Figure 2. *P. aeruginosa* **infection engages a canonical NLRC4-Caspase-1-**
Gasdermin D-dependent pyroptosis axis in neutrophils.

A. Measure of cell lysis (release of LDH) and IL-1β release in WT or *Nlrc4*[-/-] murine
Bone Marrow Neutrophils (BMNs) infected for 3 hours with *Pseudomonas aeruginosa*
PAO1 or CHA strains and their isogenic mutants lacking T3SS-derived toxins ExoS,
ExoY, ExoT) at a multiplicity of infection (MOI) of 10. ***p ≤ 0.001, T-test with
Bonferroni correction. Values are expressed as mean ± SEM. Graphs show one
experiment representative of three independent experiments.

B. Immunoblotting of GAPDH, pro-forms of Caspase-1 (p45) and Gasdermin-D (p55),
processed Caspase-1 (p20) and Gasdermin D (p30), in WT, *Casp1*[-/-] and *GsdmD*[-/-] in
cell lysates and cell supernatants (SNs) of BMNs infected for 3 hours with PAO1 or its
isogenic mutants lacking T3SS expression (PAO1[ExsA-]) or ExoS (PAO1[ExoS-]) at a

multiplicity of infection (MOI) of 10. Immunoblots show lysates and supernatants from one experiment performed three times.

C. Immunoblotting of preforms of Caspase-1 (p45) and Gasdermin-D (p55), processed Caspase-1 (p20) and Gasdermin-D (p30) and GAPDH in WT and $Nlrc4^{-/-}$ BMNs infected for 3 hours with PP34^{ExoUS142A} (MOI2) or *S.* Typhimurium (*S.*Tm, MOI 10). Immunoblots show combined lysates and supernatants from one experiment performed three times.

D. Measure of cell lysis (release of LDH) and IL-1β release in WT or $Aim2^{-/-}$, $Nlrp3^{-/-}$, $Casp11^{-/-}$, $ASC^{-/-}$, $Nlrc4^{-/-}$, $Casp1^{-/-}$ and $GsdmD^{-/-}$ murine Bone Marrow Neutrophils (BMNs) infected for 3 hours with PP34^{ExoUS142A} at a multiplicity of infection (MOI) of 2. ***$p \leq 0.001$, T-test with Bonferroni correction. Values are expressed as mean ± SEM. Graphs show one experiment representative of three independent experiments.

E. Immunoblotting of Myeloperoxidase (MPO), pro-forms of Caspase-1 (p45) and Gasdermin-D (p55), processed Caspase-1 (p20) and Gasdermin D (p30), in WT, $Nlrc4^{-/-}$ and $GsdmD^{-/-}$ BMNs infected for 3 hours with PP34^{ExoU} or its isogenic mutant lacking ExoU activity (PP34^{ExoUS142A}) at a multiplicity of infection (MOI) of 2. Immunoblots show combined lysates and supernatants from one experiment performed three times.

F. Imagestream experiments and quantifications of for *in vivo* formation of ASC specks in bronchoalveolar (BALs) neutrophils from ASC-Citrine mice intranasally infected with 1.10^5 PP34^{ExoU-} or PP34$^{ExoU-FliC-}$ for 6 hours. The gating strategy used to evaluate inflammasome activation in neutrophils was performed as follow: (i) a gate was set on cells in focus [Cells in Focus] and (ii) a sub-gate was created on single cells [Single Cells]. Then we gated first on (iii) LY6G+ Neutrophils [LY6G+] and second on (iv) ASC-citrine+ and Hoechst+ cells [Hoechst+/ASC-Citrine+] within LY6G+ population. (v) To distinguish cells with active (ASC-speck) versus inactive inflammasome (Diffuse ASC), we plotted the Intensity with the area of ASC-citrine. This strategy allows to distinguish cells with active inflammasome that were visualized and quantified. Values are expressed as mean ± SEM. Graphs show one experiment representative of two independent experiments.

Figure 3. Caspase-1-induced neutrophil pyroptosis promotes PAD4-dependent incomplete NETosis

A. Scanning electron microscopy (SEM) observation of pyroptosis in WT, *Casp1*$^{-/-}$ and *GsdmD*$^{-/-}$ BMNs 3 hours after PAO1, PAO1^{ExoS-}, PP34^{ExoU}, PP34^{ExoUS142A} infection at an MOI of 2. Images are representative of one experiment performed 3 times. Scale bars are directly indicated in the Figure.

B. Confocal microscopy observations of PP34^{ExoU}- or PP34^{ExoUS142A}-infected human blood neutrophils (MOI 2) for 3 hours harboring ASC complexes and decondensed DNA. Nucleus (blue) was stained with Hoescht; ASC is in red (anti-ASC); plasma membrane is in grey (WGA staining). Images are representative of one experiment performed 3 times. Scale bar 10µm.

C. Representative time lapse fluorescence microscopy images of ASC-Citrine murine BMNs infected with PP34^{ExoU}- or PP34^{ExoUS142A} (MOI 2) for 9 hours (540 minutes). Nucleus (blue) was stained with Hoescht; ASC is in yellow (ASC-Citrine); plasma membrane is in green (WGA staining); plasma membrane permeabilization is stained

in red (cell impermanent DNA dye Propidium Iodide, PI). Images are representative of

one movie out of three independent movies. Scale bar 10µm.

D. Immunoblotting observation of Histone 3, HMGB1, Lamin B1, GAPDH, Actin,

Gasdermin D (GSDMD) and NLRC4 in cellular soluble and insoluble fractions as well

as in the supernatant from WT and *Nlrc4$^{-/-}$* murine BMNs infected with PP34^{ExoU}- or

PP34^{ExoUS142A} (MOI 2) for 3 hours. Immunoblots show one experiment performed three

times.

E. Confocal microscopy observations and quantifications of the percentage of cells

harboring ASC complexes, H3Citrullination and nuclear/decondensed DNA in WT-

ASC-Citrine^{+} BMNs infected for 3 hours with PP34^{ExoUS142A} in presence/absence of the

PAD4 inhibitor GSK484 (10µM). Nucleus (blue) was stained with Hoescht; Histone-3

Citrullination is in red (Anti-H3Cit staining); plasma membrane is in grey (WGA

staining). Scale bar 10µm. For quantifications, the percentage of cells with ASC

complexes, nuclear DNA or positives for H3Cit (H3-Cit^{+}) was determined by quantifying

the ratios of cells positives for ASC speckles, nuclear DNA or H3Cit. At least 10 fields

from n=3 independent experiments were analyzed. Values are expressed as mean ±

SEM.

Figure 4. Neutrophil Caspase-1 contributes to IL-1β production and to ***Pseudomonas aeruginosa* spread in mice**

A. Intravital microscopy visualization of granulocyte death in MRP8-GFP⁺ mice infected with $2.5.10^5$ CFUs of PP34^{ExoU} (NETosis-inducing strain) or PP34^{ExoUS142A} (incomplete NETosis-inducing strain) in presence of SYTOX Blue for 10 hours. Granulocyte death was observed in infected lungs by the appearance of SYTOX blue fluorescence. Pseudo colors represent vessels (gray, mTG); Granulocytes (Green, MRP8-GFP⁺); Dead cells (Purple, SYTOX blue). Scale bar: 20µm. Data show one experiment representative of 5 independent mice.

B. Survival of MRP8^{Cre-} *Casp1*^{flox} and MRP8^{Cre+}*Casp1*^{flox} mice intranasally infected with 5.10^5 CFUs of PP34^{ExoU} or PP34^{ExoUS142A} for 24 hours (n=6 animals per condition). Graphs represent one experiment (6 mice/group) out of three independent *in vivo* experiments. Log-rank Cox-Mantel test was used for survival comparisons. ***p ≤ 0.001. NS; Not significant.

C. Bronchoalveolar (BAL) and lung bacterial loads (colony forming units, CFUs) in MRP8^{Cre-} *Casp1*^{flox} and MRP8^{Cre+}*Casp1*^{flox} mice intranasally infected with $2.5.10^5$ CFUs

of PP34^{ExoU} or PP34^{ExoUS142A} for 24 hours. Graphs represent one experiment (7

mice/group) out of three independent *in vivo* experiments; **p ≤ 0.01, Mann-Whitney

analysis test.

D. Determination of IL-1β levels in Bronchoalveolar Fluids (BALFs) MRP8^{Cre-} *Casp1*^{flox}

and MRP8^{Cre+}*Casp1*^{flox} mice at 10 hours after intranasal infection with $5\ 10^5$ CFUs (5

mice/group) of PP34^{ExoU} or PP34^{ExoUS142A}. Graphs represent one experiment (5

mice/group) out of three independent *in vivo* experiments; **p ≤ 0.01, Mann-Whitney

analysis test.

E. Survival of MRP8^{Cre-} *Casp1*^{flox} and MRP8^{Cre+}*Casp1*^{flox} mice intranasally infected with

1.10^7 CFUs of PAO1^{ExoS} or PAO1^{ExoS-} for 24 hours (n=6 animals per condition). Graphs

represent one experiment (6 mice/group) out of three independent *in vivo* experiments.

Log-rank Cox-Mantel test was used for survival comparisons. NS; Not significant.

F. Bronchoalveolar (BAL) and lung bacterial loads (colony forming units, CFUs) in

MRP8^{Cre-} *Casp1*^{flox} and MRP8^{Cre+}*Casp1*^{flox} mice intranasally infected with 5.10^6 CFUs

of PAO1^{ExoS} or PAO1^{ExoS-} for 24 hours. Graphs represent one experiment (7

mice/group) out of three independent *in vivo* experiments; *p ≤ 0.05, **p ≤ 0.01, Mann-

Whitney analysis test.

G. Determination of IL-1β levels in Bronchoalveolar Fluids (BALFs) MRP8^{Cre-} *Casp1*^{flox}

and MRP8^{Cre+}*Casp1*^{flox} mice at 8 hours after intranasal infection with $5.\ 10^6$ CFUs (5

mice/group) of PAO1^{ExoS} or PAO1^{ExoS-}. Graphs represent one experiment (5

mice/group) out of three independent *in vivo* experiments; **p ≤ 0.01, Mann-Whitney

analysis test.

Supplemental Figures

S1 Movie. Intravital microscopy visualization of granulocyte death in MRP8-GFP$^+$ mice

infected with 2.5.10^5 CFUs of PP34^{ExoU} (NETosis-inducing strains) or PP34^{ExoUS142A}

(incomplete NETosis-inducing strain) in presence of SYTOX Blue for 10 hours.

Granulocyte death was observed in infected lungs by the appearance of SYTOX blue

fluorescence. Pseudo colors represent vessels (gray, mTG); Granulocytes (Green,

MRP8-GFP+); Dead cells (Purple, SYTOX blue). Scale bar: 20µm.

S2 Movie. Time Lapse Fluorescence microscopy of ASC-Citrine murine BMNs

infected with PP34^{ExoU} (MOI 2) for 9 hours (540 minutes). Nucleus (blue) was stained

with Hoescht; ASC is in yellow (ASC-Citrine); plasma membrane is in green (WGA

staining); plasma membrane permeabilization is stained in red (cell impermanent DNA

dye Propidium Iodide, PI).

S3 Movie. Time Lapse Fluorescence microscopy of ASC-Citrine murine BMNs

infected with PP34^{ExoUS142} (MOI 2) for 9 hours (540 minutes). Nucleus (blue) was

stained with Hoescht; ASC is in yellow (ASC-Citrine); plasma membrane is in green

(WGA staining); plasma membrane permeabilization is stained in red (cell

impermanent DNA dye Propidium Iodide, PI).

Supplemental Figure 1. Multiple *P. aeruginosa* strains triggers Caspase-1-
dependent pyroptosis in neutrophils.

A. Measure of cell lysis (release of LDH) and IL-1β release in WT or *Casp1⁻/⁻* murine
Bone Marrow Neutrophils (BMNs) infected for 3 hours with *Pseudomonas aeruginosa*
PAO1 and its isogenic mutants lacking T3SS-derived toxins ExoS, ExoY, ExoT or
Flagellin (FliC) (PAO1^{ExoS-}, PAO1^{ExoY-}, PAO1^{ExoT-}, PAO1^{FliC-}) at a multiplicity of infection
(MOI) of 10. ***p ≤ 0.001, T-test with Bonferroni correction. Values are expressed as
mean ± SEM. Graphs show one experiment representative of three independent
experiments.

B. Measure of cell lysis (release of LDH) and IL-1β release in WT, *Casp1⁻/⁻*, *Casp1⁻/⁻*
Casp11⁻/⁻, *Casp11⁻/⁻* and *GsdmD⁻/⁻* murine Bone Marrow Neutrophils (BMNs) infected
for 3 hours with *Pseudomonas aeruginosa* PP34^{ExoU} and its isogenic mutant PP34^{ExoU-}
at a multiplicity of infection (MOI) of 2. ***p ≤ 0.001, T-test with Bonferroni correction.
Values are expressed as mean ± SEM. Graphs show one experiment representative
of three independent experiments.

C. Measure of plasma membrane permeabilization over time using SYTOX Green
incorporation in murine BMNs or in human blood neutrophils infected with PP34^{ExoU}
and its isogenic catalytically inactive mutant PP34^{ExoUS142A} at a multiplicity of infection
(MOI) of 2 in presence/absence of the phospholipase inhibitor MAFP (20µM). ***p ≤
0.001, T-test with Bonferroni correction. Values are expressed as mean ± SEM.
Graphs show one experiment representative of three independent experiments.

D. Measure of plasma membrane permeabilization over time using SYTOX Green incorporation in murine WT and *Casp1*[-/-] BMNs or in human blood neutrophils infected with PP34[ExoU] and its isogenic catalytically inactive mutant PP34[ExoUS142A] at a multiplicity of infection (MOI) of 2 in presence/absence of various Caspase inhibitors, Z-YVAD (Casp1 inhibitor, 20µM), Z-DEVD (Casp3 inhibitor, 20µM), Z-IETD (Casp8 inhibitor, 20µM). ***p ≤ 0.001, T-test with Bonferroni correction. Values are expressed as mean ± SEM. Graphs show one experiment representative of three independent experiments.

Supplemental Figure 2. *P. aeruginosa* **uniquely triggers a NLRC4-dependent**

pyroptosis in neutrophils.

A, B. Measure of cell lysis (release of LDH) in WT or *Nlrc4*[-/-] murine Bone Marrow

Neutrophils (BMNs) infected for 3 hours with various ExoU-expressing *Pseudomonas*

aeruginosa strains (PP34, PA14, PA103) or their isogenic mutants lacking T3SS-

derived ExoU, Flagellin (FliC) or T3SS expression (ExsA) at a multiplicity of infection

(MOI) of 2 (PP34) or 10 (PA14, PA103). ***$p \le 0.001$, T-test with Bonferroni correction.

Values are expressed as mean ± SEM. Graphs show one experiment representative

of three independent experiments.

C. Measure of cell lysis (release of LDH) and IL-1β release in WT and *Nlrc4*[-/-] BMNs

infected for 3 hours with PP34[ExoUS142A] (MOI 2), *Salmonella* Typhimurium (*S*.Tm

SL1344, MOI 10), *Chromobacter violaceum* (*C. violaceum*, MOI 20), *Shigella flexneri*

(*S. flexneri*, MOI 40) or *Burkholderia thailandensis* (*B. thailandensis*, MOI 40). ***$p \le$

0.001, T-test with Bonferroni correction. Values are expressed as mean ± SEM.

Graphs show one experiment representative of three independent experiments.

D. Gating strategy used to evaluate inflammasome activation in neutrophils was performed as follow: (i) a gate was set on cells in focus [Cells in Focus] and (ii) a sub-gate was created on single cells [Single Cells]. Then we gated first on (iii) LY6G⁺ Neutrophils [LY6G⁺] and second on (iv) ASC-citrine⁺ and Hoechst⁺ cells [Hoechst⁺/ASC-Citrine⁺] within LY6G⁺ population. (v) To distinguish cells with active (ASC-speck) versus inactive inflammasome (Diffuse ASC), we plotted the Intensity with the area of ASC-citrine. This strategy allow to distinguish cells with active inflammasome that were visualized and quantified. Values are expressed as mean ± SEM. Graphs show one experiment representative of two independent experiments.

Supplemental Figure 3. NLRC4-dependent neutrophil pyroptosis promotes an incomplete NETosis.

A. Confocal microscopy observations of WT, *Casp1*[-/-] and *GsdmD*[-/-] BMNs infected for 3 hours with PP34[ExoUS142A] and harboring ASC complexes, decondensed DNA and nuclear membrane (LaminB1). Nucleus (blue) was stained with Hoescht; LaminB1 is in red (anti LaminB1); ASC is in Green (anti-ASC); plasma membrane is in grey (WGA staining). Scale bar 10µm. Images are representative of one experiment performed three times with at least 200 neutrophils observed/ experiment.

B. Immunoblotting observation of Histone 3, HMGB1, Lamin B1, GAPDH, Actin, Gasdermin D (GSDMD) and NLRC4 in cellular soluble and insoluble fractions as well as in the supernatant from WT and *Nlrc4*[-/-] murine BMNs infected with PAO1[ExoS-] (MOI 10) for 3 hours. Immunoblots show one experiment performed two times.

C. Confocal microscopy observations of WT, *Nlrc4*[-/-] and *GsdmD*[-/-] BMNs infected for 3 hours with PP34[ExoUS142A] and harboring Citrullinated Histone 3 (H3-Cit), decondensed DNA and nuclear membrane (LaminB1). Nucleus (blue) was stained with Hoescht; Citrullinated Histone-3 is in red (anti H3-Cit); plasma membrane is in grey (WGA staining). Scale bar 10µm. Images are representative of one experiment performed three times with at least 150 neutrophils observed/ experiment.

D. Confocal microscopy observations of human blood neutrophils infected for 3 hours with PAO1[ExoS-] (MOI 10) or PP34[ExoUS142A] (MOI 2) in presence/absence of Caspase-1 inhibitor Z-YVAD (20µM) and harboring Citrullinated Histone 3 (H3-Cit), decondensed DNA and nuclear membrane (LaminB1). Nucleus (blue) was stained with Hoescht; Citrullinated Histone-3 is in red (anti H3-Cit); plasma membrane is in grey (WGA staining). Scale bar 10µm. Images are representative of one experiment performed three times with at least 150 neutrophils observed/ experiment.

E. Immunoblotting of Citrullinated Histone 3 (H3Cit), total Histone 3 and preformed and cleaved Gasdermin-D (p55/p30) in WT, *Nlrc4*[-/-] and *GsdmD*[-/-] BMNs treated with Ionomycin (10µM, 3 hours) or infected for 3 hours with PAO1, PAO1[ExoS-], PAO1[ExsA-] (MOI 10) or with PP34[ExoU] , PP34[ExoU-] (MOI 2). Immunoblots show combined lysates and supernatants from one experiment performed three times.

F. Measure of cell lysis (release of LDH) and bacterial elimination (Colony-forming Units, CFUs) in WT, *Pad4*[-/-] and *Nlrc4*[-/-] BMNs infected for 3 hours with PP34[ExoU] (MOI2), PP34[ExoUS142A] (MOI2), PAO1 (MOI 10) or PAO1[ExoS-] (MOI 10). ***p ≤ 0.001, T-test with Bonferroni correction. NS: Not significant. Values are expressed as mean ± SEM. Graphs show one experiment representative of three independent experiments.

Methods

All reagents, concentrations of use and their references are listed in **Table S1**

Mice

Nlrc4$^{-/-}$ [61], *Nlrp3$^{-/-}$* [65], *ASC$^{-/-}$*, *Casp11$^{-/-}$* [63,64], *Casp1$^{-/-}$Casp11$^{-/-}$* [63,64], *GsdmD$^{-/-}$*, *Aim2$^{-/-}$*, *Pad4$^{-/-}$*, *MRP8^{Cre+}GFP$^+$*, *MRP8^{Cre+}Casp1^{flox}* were generated and described in previous studies. Mice were bred at the IPBS (Toulouse, France) and INRAE (Tours Nouzilly, France) animal facilities in agreement to the EU and French directives on animal welfare (Directive 2010/63/EU). Charles Rivers provided WT C57BL/6 mice. Mice experiments are under legal authorizations APAFIS#8521-2017041008135771 and APAFIS#12812-2018031218075551, according to the local, French and European ethic laws.

MRP8^{Cre}*Casp1*^{flox} mice genotyping

Casp1$^{flox/flox}$ mice were kindly provided by Mo Lamkanfi and were crossed to MRP8^{Cre} mice to generate MRP8^{Cre}*Casp1*^{flox}. *Caspase-1* genotyping was performed using Primer Fw: CGAGGGTTGGAGCTCAAGTTGACC and Primer Rv: CACTTTGACTTCTCTAAGGACAG. *Cre* genotyping was performed using Primers Fw: CGCCGTAAATCAATCGATGAGTTGCTTC and Primers Rv: GATGCCGGTGAACGTGCAAAACAGGCTC.

Bacterial cultures

P. aeruginosa strains (PAO1, CHA, PP34, PA14, PA103) and their isogenic mutants were grown overnight in Luria Broth (LB) medium at 37°C with constant agitation. Bacteria were sub-cultured the next day by diluting overnight culture 1:25 and grew until reaching an optical density (OD) O.D.600 of 0.6 – 0.8. Bacterial strains and their mutants are listed in **Table S1**.

Bacterial KO generation and complementation

The knockout vector pEXG2 was constructed and used based on the protocol described by Rietsch et al. [67] with the following modifications. Briefly, 700-bp sequences of the flanking regions of the selected gene were amplified by PCR with Q5 high fidelity polymerase (New England Biolabs). Then, the flanking regions were gel

purified and inserted into pEXG2 plasmid by Gibson assembly [68]. The assembled

plasmid was directly transformed into competent SM10λpir using Mix&Go competent

cells (Zymo Research Corporation) and plated on selective LB plates containing 50

µg/mL kanamycin and 15 µg/mL gentamicin. The resulting clones were sequenced,

and mating was allowed for 4 h with PAO1 strain at 37°C. The mated strains were

selected for single cross over on plates containing 15 µg/mL gentamicin and 20 µg/mL

Irgasan (removal of E.coli SM10 strains). The next day, some clones were grown in LB

for 4 hours and streaked on 5% sucrose LB plates overnight at 30°C. P. aeruginosa

clones were then checked by PCR for mutations. All primers were designed with

Snapgene software (GSL Biotech LLC).

Mice infections

Age and sex-matched animals (5–8 weeks old) per group were infected intranasally

with 5.10^5 (lethal doses) or $2.5.10^5$ CFUs of PP34^{ExoU}/PP34^{ExoUS142A} or with 1.10^7 CFUs

(lethal doses) or 5.10^6 CFUs of PAO1/PAO1^{ExoS-} strains suspended in 25µL of PBS.

Animals were sacrificed at indicated times after infection and bronchoalveolar fluids

(BALFs) and lungs were recovered. When specified, bacterial loads (CFU plating),

cytokine levels (ELISA) were evaluated. No randomization or blinding were done.

Intravital microscopy experiments

We relied on the previously published lung intravital microscopy method using an

intercoastal thoracic window [69,70], adapted at the IPBS CNRS-University of

Toulouse TRI platform.

MRP8-mTmG mice (8-12 weeks old) were infected intratracheally with 5.10^5 CFUs of

P. aeruginosa ExoU or ExoU^{S142A} strains resuspended in 50µL of PBS and imaged 6

to 8 hours after infection. 50µL of 50µM solution of SYTOX blue (Life Technologies)

was injected both intravenously (retroorbital) and intratracheally just before imaging to

visualize extracellular DNA.

Next, mice were anesthetized with ketamine and xylazine and secured to a microscope

stage. A small tracheal cannula was inserted, sutured and attached to a MiniVent

mouse ventilator (Harvard Apparatus). Mice were ventilated with a tidal volume of 10

µl of compressed air (21% O_2) per gram of mouse weight, a respiratory rate of 130-

140 breaths per minute, and a positive-end expiratory pressure of 2-3 cm H_2O.

Isoflurane was continuously delivered to maintain anesthesia and 300 µl of 0.9% saline

solution were i.p. administered in mice every hour for hydration. Mice were placed in the right lateral decubitus position and a small surgical incision was made to expose the rib cage. A second incision was then made into the intercostal space between ribs 4 and 5, through the parietal pleura, to expose the surface of the left lung lobe. A flanged thoracic window with an 8 mm coverslip was inserted between the ribs and secured to the stage using a set of optical posts and a 90° angle post clamp (Thor Labs). Suction was applied to gently immobilize the lung (Dexter Medical). Mice were placed in 30°C heated box during microscopy acquisition to maintain the body temperature and the 2-photon microscope objective was lowered over the thoracic window. Intravital imaging was performed using a Zeiss 7MP upright multi-photon microscope equipped with a 20×/1.0 objective and a Ti-Sapphire femtosecond laser, Chameleon-Ultra II (Coherent Inc.) tuned to 920 nm. SYTOX Blue, GFP and Tomato emission signals were detected thanks to the respective bandpass filters: Blue (SP485), Green (500-550) and Red (565-610). Images were analyzed using Imaris software (Bitplane) and Zen (Zeiss).

Isolation of primary murine neutrophils

Murine Bone marrow cells were isolated from tibias and femurs, and neutrophils were purified by positive selection using Anti-Ly-6G MicroBead Kit (Miltenyi Biotech) according to manufacturer's instructions. This process routinely yielded >95% of neutrophil population as assessed by flow cytometry of $Ly6G^+/CD11b^+$ cells.

Isolation of primary human neutrophils

Whole blood was collected from healthy donors by the "Ecole française du sang" (EFS, Tolouse Purpan, France) in accordance with relevant guidelines. Written, informed consent was obtained from each donor. Neutrophils were then isolated by negative selection using MACSxpress® Whole Blood Human Neutrophil Isolation Kit (Miltenyi Biotech) according to manufacturer's instructions. Following isolations cells were centrifuged 10 min at 300 g and red blood cells were eliminated using Red blood cells (RBC) Lysis Buffer (BioLegend). This procedure gives >95% of neutrophil population as assessed by flow cytometry of CD15+/CD16+ cells. License to use human samples is under legal agreement with the EFS; contract n° 21PLER2017-0035AV02, according to Decret N° 2007-1220 (articles L1243-4, R1243-61).

Cell plating and treatment of Neutrophils

Following isolation, Neutrophils were centrifugated for 10 min at 300 g and pellet was resuspendent in serum free OPTI-MEM medium. Absolute cell number was determined with automated cell counter Olympus R1 with trypan blue cell death exclusion method (typically living cells represent >70% of cell solution) and cell density was adjusted at 10^6 / mL by adding OPTI-MEM culture medium. Neutrophils were then plated in either 96 well plates, 24 well plates or 6 well plates with 100 μL (10^5 cells), 500 μL (5.10^5 cells) or 2 mL (2.10^6 cells) respectively. When indicated cells were incubated with chemical inhibitors Z-VAD-fmk (20 μM), Y-VAD-fmk (40 μM), Z-DEVD-fmk (40μM), Z-IETD-fmk (40μM), GSK484 (10 μM), as indicated in each experimental setting. Neutrophils were infected with various bacterial strains and multiplicity of infections (M.O.I.) as indicated.

Kinetic analysis of Neutrophil's permeability by SYTOX Green incorporation assay

Cells are plated at density of 1 x 10^5 per well in Black/Clear 96-well Plates in OPTI-MEM culture medium supplemented with SYTOX-Green dye (100ng/mL) and infected/treated as mentioned in figure legend. Green fluorescence are measured in real-time using Clariostar plate reader equipped with a 37°C cell incubator.

ELISA and plasma membrane lysis tests

Cell death was measured by quantification of the lactate dehydrogenase (LDH) release into the cell supernatant using LDH Cytotoxicity Detection Kit (Takara). Briefly, 100 μL cell supernatant were incubated with 100 μL LDH substrate and incubated for 15 min. The enzymatic reaction was stopped by adding 50 μL of stop solution. Maximal cell death was determined with whole cell lysates from unstimulated cells incubated with 1% Triton X-100. Human and mouse IL-1β secretion was quantified by ELISA kits (Thermo Fisher Scientific) according to the manufacturer's instructions.

Preparation of neutrophil lysates and supernatant for immunoblot

At the end of the treatment 5 mM of diisopropylfluorophosphate (DFP) cell permeable serine protease inhibitor was added to cell culture medium. Cell' Supernatant was collected and clarified from non-adherent cells by centrifugation for 10 min at 300 g.

Cell pellet and adherent cells were lysed in 100 µL of RIPA buffer (150 mM NaCl,
50 mM Tris-HCl, 1% Triton X-100, 0.5% Na-deoxycholate) supplemented with 5 mM
diisopropylfluorophosphate (DFP) in addition to the protease inhibitor cocktail (Roche).
Cell scrapper was used to ensure optimal recovery of cell lysate. Collected cell lysate
was homogenized by pipetting up and down ten times and supplemented with laemli
buffer (1X final) before boiling sample for 10 min at 95°C. Soluble proteins from cell
supernatant fraction were precipitated as described previously [71]. Precipitated pellet
was then resuspended in 100 µL of RIPA buffer plus laemli supplemented with 5 mM
diisopropylfluorophosphate (DFP) and protease inhibitor cocktail (Roche) and heat
denaturated for 10 min at 95°C. Cell lysate and cell supernatant fraction were then
analysed by immunoblot either individually or in pooled sample of lysate plus
supernatant (equal vol/vol).

1020

Treatment of Neutrophils for Immunofluorescences
5.10^5 Cells were plated on 1.5 glass coverslips in 24 well plate and infected/treated as
described above. At the end of the assay, cell supernatant was removed and cells were
fixed with a 4% PFA solution for 10 min at 37°C. PFA was then removed and cells were
washed 3 times with HBSS. When desired, plasma membrane was stained with Wheat
Germ Agglutinin, Alexa Fluor™ 633 Conjugate (ThermoFisher Scientifique) at 1/100th
dilution in HBSS, and incubated for 30 min under 100 rpm orbital shaking conditions.
Then cells were washed with HBSS and processed for further staining.
Permeabilization was performed by incubating cells for 10 min in PBS containing 0.1%
Triton X-100. To block unspecific binding of the antibodies cells are incubated in PBS-
T (PBS+ 0.1% Tween 20), containing 2% BSA, 22.52 mg/mL glycine in for 30 min. 3
washes with PBS-T was performed between each steps. Primary antibodies staining
was performed overnight at 4°C in BSA 2% - Tween 0.1% - PBS (PBS-T) solution.
Coverslips were washed three times with PBS-T and incubated with the appropriate
fluor-coupled secondary antibodies for 1 hour at room temperature. DNA was
counterstained with Hoechst. Cells were then washed three times with PBS and
mounted on glass slides using Vectashield (Vectalabs). Coverslips were imaged using
confocal Zeiss LSM 710 (INFINITY, Toulouse) or Olympus Spinning disk (Image core
Facility, IPBS, Toulouse) using a 40x or a 63x oil objective. Unless specified, for each
experiment, 5-10 fields (~50-250 cells) were manually counted using Image J
software.

Scanning Electron Microscopy experiments

For scanning electron microscopy observations, cells were fixed with 2.5% glutaraldehyde in 0.2M cacodylate buffer (pH 7.4). Preparations were then washed three times for 5min in 0.2M cacodylate buffer (pH 7.4) and washed with distilled water. Samples were dehydrated through a graded series (25 to 100%) of ethanol, transferred in acetone and subjected to critical point drying with CO_2 in a Leica EM CPD300. Dried specimens were sputter-coated with 3 nm platinum with a Leica EM MED020 evaporator and were examined and photographed with a FEI Quanta FEG250.

ImageStreamX

Cells isolated from peritoneal washes were pelleted by centrifugation (10 min at 300 g). Neutrophils were stained prior to fixation with anti-Ly6G (APC-Vio770, Miltenyi-Biotec Clone: REA526 | Dilution: 1:50) in MACS buffer (PBS-BSA 0,5%-EDTA 2mM) in presence of FC block (1/100) and Hoechst (1 µM). Then, cells were fixed in 4% PFA. Data were acquired on ImageStreamXMKII (Amnis) device (CPTP Imaging and Cytometry core facility) and analyzed using IDEAS software v2.6 (Amnis). The gating strategy used to evaluate inflammasome activation in neutrophils was performed as follow: (i) a gate was set on cells in focus [Cells in Focus] and (ii) a sub-gate was created on single cells [Single Cells]. Then we gated first on (iii) LY6G+ Neutrophils [LY6G+] and second on (iv) ASC-citrine+ and Hoechst+ cells [Hoechst+/ASC-Citrine+] within LY6G+ population. (v) To distinguish cells with active (ASC-speck) versus inactive inflammasome (Diffuse ASC), we plotted the Intensity with the area of ASC-citrine. This strategy allow to distinguish cells with active inflammasome that were visualized and quantified (**S2 Fig.**).

Statistical tests used

Statistical analysis was performed with Prism 8.0a (GraphPad Software, Inc.). Otherwise specified, data are reported as mean with SEM. T-test with Bonferroni correction was chosen for comparison of two groups. For *in vivo* mice experiments and comparisons we used Mann-Whitney tests and mouse survival analysis were performed using log-rank Cox-Mantel test. P values are shown in figures with the following meaning; NS non-significant and Significance is specified as $*p \leq 0.05$; $**p \leq 0.01$, $***p \leq 0.001$.

 Supplemental 1 Table

1077 Reagents and Tools are available upon request to Etienne.meunier@ipbs.fr or

1078 Remi.planes@ipbs.fr

REAGENT or RESSOURCE	SOURCE	IDENTIFIER
Antibodies		
Anti- mouse Caspase 1, 1: 1000	AdipoGen	AG20B-0042
Anti-Ly6G APC-Vio770, 1:50	Miltenyi-Biotec	**130-119-126**
Anti- mouse Gasdermin D, 1: 1000	Abcam	AB209845
Anti- mouse IL-1beta, 1: 1000	R&D Systems	AF-401-NA
Anti-MPO , 1: 1000	R&D Systems	AF3667
Anti-ASC, IF 1:100	Novus	NBP1-78-977
Anti-HMGB1, 1: 1000	Genetex	GTX-101277
Anti-HMGB1, 1: 1000	abcam	ab18256
Anti- LaminB1, 1: 1000 Blots, 1:250 IF	abcam	AB229025 [EPR22165-121]
Anti-H3 citrullinated, 1: 1000 Blots, 1:250 IF	abcam	AB5103
Anti-H3, 1: 1000	Cell signalling Techonology	3638S
Anti-H4, 1: 1000	Cell signalling Techonology	2935S
Anti-H2A, 1: 1000	Cell signalling Techonology	2578S
Anti-H1, 1: 1000	Abcam	AB134914 [EPR6536]
Anti-PAD4, 1: 1000	Abcam	AB214810 [EPR20706]
Anti-NLRC4 1: 1000	Abcam	ab201792 [EPR19733]
Anti-β-actin 1: 1000	Cell signalling Techonology	4967S
Anti-β-actin 1: 5000	Sigma-Aldrich	A1978
Goat anti-mouse HRP (1/4000)	SouthernBiotech	1034-05
Goat-anti-rabbit IgG (H+L), HRP conjugate (1/4000)	Advansta	R-05072-500
Rabbit anti-Goat IgG (H+L) Secondary Antibody, HRP (1/4000)	Invitrogen	81-1620
Goat anti-rabbit IgG (H&L)Dylight 488 (1/1000)	Immunoreagents	GtxRb-003-D488NHSX
Donkey anti-Rabbit IgG Dylight 550 (1/1000)	Immunoreagents	DkxRb-003-D550NHSX
Goat anti-mouse IgG Dylight 550 (1/1000)	Immunoreagents	GtxMu-003-D550NHSX
Goat anti-mouse IgG Dylight 488 (1/1000)	Immunoreagents	GtxMu-003-D488NHSX
Hoescht	Sigma-Aldrich	H6024
WGA-Alexa 633 (1/100)	Thermo Fisher Scientific	W21404
Alexa Fluor™ 488 Phalloidin	Thermo Fisher Scientific	A12379

RNA, DNA primers/sequences		
MRP8-Cre Fw primers	CGCCGTAAATCAATCGATGAGTTGCTTC	
MRP8-Cre Rv primers	GATGCCGGTGAACGTGCAAAACAGGCTC	
Casp1flox Fw primers	CGAGGGTTGGAGCTCAAGTTGACC	
Casp1flox Rv primers	CACTTTGACTTCTCTAAGGACAG	
Chemicals, peptides and recombinant proteins		
DFP (5mM)	Sigma-Aldrich	D0879
GSK484 (10µM)	Cayman Chemical	17488
ZVAD (20µM)	Invivogen	tlrl-vad
Y-VAD (40µM)	Invivogen	inh-yvad
MAFP (20µM)	Sigma	M2939
SYTOX™ Green Nucleic Acid Stain (100ng/mL)	Thermo Fisher Scientific	S7020
SYTOX™ Blue Nucleic Acid Stain (100ng/mL)	Thermo Fisher Scientific	S11348
PFA (4%)	Sigma-Aldrich	1004969011
cOmplete™ Protease Inhibitor Cocktail	Sigma-Aldrich	11697498001
Lipofectamine-2000	Thermo Fisher Scientific	11668019
Triton X-100	Sigma-Aldrich	X100-500ML
Newborn Calf Serum (FCS)	Thermo Fisher Scientific	16010159
DMEM	Gibco	11574486
HEPES	Invitrogen	15630080
Opti-MEM	Gibco	11524456
RPMI without phenol red	Gibco	11564456
HBSS without Calcium, Magnesium or Phenol Red	Gibco	14-175-095
Clarity Max™ Western ECL Substrate	Bio-Rad	1705062
Tris Base Ultrapure	Euromedex	EU1018-A
SDS Ultrapure 4X	Euromedex	1012-A
Acrylamide / Bisacrylamide 37.5/1 30%	Euromedex	EU0088-B
TEMED	Sigma-Aldrich	T9281
PageRuler™ Prestained Protein Ladder	Thermo Fisher Scientific	11822124
Vectashield	Vectorlabs	H-1000-10
MACSxpress Whole Blood Neutrophil Isolation Kit human	Miltenyi Biotech	130-104-434
Anti-Ly-6G MicroBeads UltraPure, mouse	Miltenyi Biotech	130-120-337
Critical comercial assays		
LDH Cytotoxicity Detection Kit	Takara	MK401
IL-1 beta Mouse Uncoated ELISA Kit	Thermo Fisher Scientific	88-7013-88
Cell death detection ELISA PLUS	Sigma-Aldrich	11774425001
Bacterial strains		
PP34	Ina Attrée	[72]
PP34ExoUS142A	Ina Attrée	[72]
PP34ExoU-	Ina Attrée	[72]
CHA	Ina Attrée	[72]
CHA ExoS-	Ina Attrée	[72]
PAO1	J Buyck	N.A.

PAO1 ExoS-	J Buyck	N.A.
PAO1 ExoY-	J Buyck	N.A.
PAO1 ExoT-	J Buyck	N.A.
PAO1 ExsA-	J Buyck	N.A.
Softwares		
Prism v.8		
FlowJo v.10		
ImageLab		
FiJi		
IDEAS software v2.6 (Amnis)		
Snapgene		

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

1100

1101

1102

1103

1104

1105

References

1. Galluzzi L, Vitale I, Aaronson SA, Abrams JM, Adam D, Agostinis P, et al. Molecular mechanisms of cell death: Recommendations of the Nomenclature Committee on Cell Death 2018. Cell Death and Differentiation. Nature Publishing Group; 2018. pp. 486–541. doi:10.1038/s41418-017-0012-4

2. Brinkmann V, Reichard U, Goosmann C, Fauler B, Uhlemann Y, Weiss DS, et al. Neutrophil Extracellular Traps Kill Bacteria. Science (80-). 2004;303: 1532–1535. doi:10.1126/science.1092385

3. Chen KW, Monteleone M, Boucher D, Sollberger G, Ramnath D, Condon ND, et al. Noncanonical inflammasome signaling elicits gasdermin D–dependent neutrophil extracellular traps. Sci Immunol. 2018;3: 6676. doi:10.1126/sciimmunol.aar6676

4. Li P, Li M, Lindberg MR, Kennett MJ, Xiong N, Wang Y. PAD4 is essential for antibacterial innate immunity mediated by neutrophil extracellular traps. J Exp Med. 2010;207: 1853–1862. doi:10.1084/jem.20100239

5. Kovacs SB, Oh C, Maltez VI, McGlaughon BD, Verma A, Miao EA, et al. Neutrophil Caspase-11 Is Essential to Defend against a Cytosol-Invasive Bacterium. Cell Rep. 2020;32. doi:10.1016/j.celrep.2020.107967

6. Martinod K, Fuchs TA, Zitomersky NL, Wong SL, Demers M, Gallant M, et al. PAD4-deficiency does not affect bacteremia in polymicrobial sepsis and ameliorates endotoxemic shock. Blood. 2015;125: 1948–1956. doi:10.1182/blood-2014-07-587709

7. Biron BM, Chung C-S, Chen Y, Wilson Z, Fallon EA, Reichner JS, et al. PAD4 Deficiency Leads to Decreased Organ Dysfunction and Improved Survival in a Dual Insult Model of Hemorrhagic Shock and Sepsis. J Immunol. 2018;200: ji1700639. doi:10.4049/jimmunol.1700639

8. Kahlenberg JM, Carmona-Rivera C, Smith CK, Kaplan MJ. Neutrophil Extracellular Trap–Associated Protein Activation of the NLRP3 Inflammasome Is Enhanced in Lupus Macrophages. J Immunol. 2013;190: 1217–1226. doi:10.4049/jimmunol.1202388

9. Fuchs TA, Brill A, Duerschmied D, Schatzberg D, Monestier M, Myers DD, et al. Extracellular DNA traps promote thrombosis. Proc Natl Acad Sci U S A. 2010;107: 15880–15885. doi:10.1073/pnas.1005743107

10. Apel F, Andreeva L, Knackstedt LS, Streeck R, Frese CK, Goosmann C, et al.
The cytosolic DNA sensor cGAS recognizes neutrophil extracellular traps. Sci
Signal. 2021;14. doi:10.1126/scisignal.aax7942

11. Kumar SVR, Kulkarni OP, Mulay SR, Darisipudi MN, Romoli S, Thomasova D,
et al. Neutrophil extracellular trap-related extracellular histones cause vascular
necrosis in severe GN. J Am Soc Nephrol. 2015;26: 2399–2413.
doi:10.1681/ASN.2014070673

12. Lefrançais E, Mallavia B, Zhuo H, Calfee CS, Looney MR. Maladaptive role of
neutrophil extracellular traps in pathogen-induced lung injury. JCI insight.
2018;3. doi:10.1172/jci.insight.98178

13. Knackstedt SL, Georgiadou A, Apel F, Abu-Abed U, Moxon CA, Cunnington AJ,
et al. Neutrophil extracellular traps drive inflammatory pathogenesis in malaria.
Sci Immunol. 2019;4: 336. doi:10.1126/SCIIMMUNOL.AAW0336

14. Thiam HR, Wong SL, Wagner DD, Waterman CM. Cellular Mechanisms of
NETosis. Annual Review of Cell and Developmental Biology. Annual Reviews
Inc.; 2020. pp. 191–218. doi:10.1146/annurev-cellbio-020520-111016

15. Papayannopoulos V, Metzler KD, Hakkim A, Zychlinsky A. Neutrophil elastase
and myeloperoxidase regulate the formation of neutrophil extracellular traps. J
Cell Biol. 2010;191: 677–691. doi:10.1083/jcb.201006052

16. Sollberger G, Choidas A, Burn GL, Habenberger P, Lucrezia R Di, Kordes S, et
al. Gasdermin D plays a vital role in the generation of neutrophil extracellular
traps. Sci Immunol. 2018;3. doi:10.1126/sciimmunol.aar6689

17. Kenny EF, Herzig A, Krüger R, Muth A, Mondal S, Thompson PR, et al. Diverse
stimuli engage different neutrophil extracellular trap pathways. Elife. 2017;6.
doi:10.7554/eLife.24437

18. Thiama HR, Wong SL, Qiu R, Kittisopikul M, Vahabikashi A, Goldman AE, et al.
NETosis proceeds by cytoskeleton and endomembrane disassembly and PAD4-
mediated chromatin decondensation and nuclear envelope rupture. Proc Natl
Acad Sci U S A. 2020;117: 7326–7337. doi:10.1073/pnas.1909546117

19. Wang Y, Li M, Stadler S, Correll S, Li P, Wang D, et al. Histone hypercitrullination
mediates chromatin decondensation and neutrophil extracellular trap formation.
J Cell Biol. 2009;184: 205–213. doi:10.1083/jcb.200806072

20. Neubert E, Meyer D, Rocca F, Günay G, Kwaczala-Tessmann A, Grandke J, et
al. Chromatin swelling drives neutrophil extracellular trap release. Nat Commun.

1174    2018;9: 1–13. doi:10.1038/s41467-018-06263-5

1175 21. D'Cruz AA, Speir M, Bliss-Moreau M, Dietrich S, Wang S, Chen AA, et al. The

1176    pseudokinase MLKL activates PAD4-dependent NET formation in necroptotic

1177    neutrophils. Sci Signal. 2018;11: 1716. doi:10.1126/scisignal.aao1716

1178 22. Chen W, Zhao J, Mu D, Wang Z, Liu Q, Zhang Y, et al. Pyroptosis Mediates

1179    Neutrophil Extracellular Trap Formation during Bacterial Infection in Zebrafish. J

1180    Immunol. 2021; ji2001335. doi:10.4049/jimmunol.2001335

1181 23. Xia S, Zhang Z, Magupalli VG, Pablo JL, Dong Y, Vora SM, et al. Gasdermin D

1182    pore structure reveals preferential release of mature interleukin-1. Nature. 2021;

1183    1–5. doi:10.1038/s41586-021-03478-3

1184 24. Tsuchiya K, Hosojima S, Hara H, Kushiyama H, Mahib MR, Kinoshita T, et al.

1185    Gasdermin D mediates the maturation and release of IL-1α downstream of

1186    inflammasomes. Cell Rep. 2021;34: 108887. doi:10.1016/j.celrep.2021.108887

1187 25. Evavold CL, Ruan J, Tan Y, Xia S, Wu H, Kagan JC. The Pore-Forming Protein

1188    Gasdermin D Regulates Interleukin-1 Secretion from Living Macrophages.

1189    Immunity. 2018;48: 35-44.e6. doi:10.1016/j.immuni.2017.11.013

1190 26. Heilig R, Dick MS, Sborgi L, Meunier E, Hiller S, Broz P. The Gasdermin-D pore

1191    acts as a conduit for IL-1β secretion in mice. Eur J Immunol. 2018;48: 584–592.

1192    doi:10.1002/eji.201747404

1193 27. Karmakar M, Minns M, Greenberg EN, Diaz-Aponte J, Pestonjamasp K, Johnson

1194    JL, et al. N-GSDMD trafficking to neutrophil organelles facilitates IL-1β release

1195    independently of plasma membrane pores and pyroptosis. Nat Commun.

1196    2020;11. doi:10.1038/s41467-020-16043-9

1197 28. Chen KW, Groß CJ, Sotomayor FV, Stacey KJ, Tschopp J, Sweet MJ, et al. The

1198    Neutrophil NLRC4 Inflammasome Selectively Promotes IL-1β Maturation without

1199    Pyroptosis during Acute Salmonella Challenge. Cell Rep. 2014;8: 570–582.

1200    doi:10.1016/j.celrep.2014.06.028

1201 29. Münzer P, Negro R, Fukui S, di Meglio L, Aymonnier K, Chu L, et al. NLRP3

1202    Inflammasome Assembly in Neutrophils Is Supported by PAD4 and Promotes

1203    NETosis Under Sterile Conditions. Front Immunol. 2021;12: 1977.

1204    doi:10.3389/FIMMU.2021.683803/BIBTEX

1205 30. Aymonnier K, Ng J, Fredenburgh LE, Zambrano-Vera K, Münzer P, Gutch S, et

1206    al. Inflammasome activation in neutrophils of patients with severe COVID-19.

1207    Blood  Adv.  2022  [cited  26  Jan  2022].

doi:10.1182/BLOODADVANCES.2021005949

31. Lin CK, Kazmierczak BI. Inflammation: A Double-Edged Sword in the Response
to Pseudomonas aeruginosa Infection. J Innate Immun. 2017;9: 250–261.
doi:10.1159/000455857

32. Hauser AR. The Type III Secretion System of Pseudomonas aeruginosa:
Infection   by   Injection.   Nat   Rev   Microbiol.   2009;7:   654.
doi:10.1038/NRMICRO2199

33. Zhao Y, Shi J, Shi X, Wang Y, Wang F, Shao F. Genetic functions of the NAIP
family of inflammasome receptors for bacterial ligands in mice. J Exp Med.
2016;213: 647–656. doi:10.1084/JEM.20160006

34. Zhao Y, Yang J, Shi J, Gong YN, Lu Q, Xu H, et al. The NLRC4 inflammasome
receptors for bacterial flagellin and type III secretion apparatus. Nature.
2011;477: 596–602. doi:10.1038/NATURE10510

35. Rauch I, Tenthorey JL, Nichols RD, Al Moussawi K, Kang JJ, Kang C, et al. NAIP
proteins are required for cytosolic detection of specific bacterial ligands in vivo.
J Exp Med. 2016;213: 657–665. doi:10.1084/JEM.20151809

36. Miao EA, Ernst RK, Dors M, Mao DP, Aderem A. Pseudomonas aeruginosa
activates caspase 1 through Ipaf. Proc Natl Acad Sci U S A. 2008;105: 2562–
2567. doi:10.1073/pnas.0712183105

37. Franchi L, Stoolman J, Kanneganti TD, Verma A, Ramphal R, Núñez G. Critical
role for Ipaf in Pseudomonas aeruginosa-induced caspase-1 activation. Eur J
Immunol. 2007;37: 3030–3039. doi:10.1002/eji.200737532

38. Hardy KS, Tessmer MH, Frank DW, Audia JP. Perspectives on the
Pseudomonas aeruginosa Type III Secretion System Effector ExoU and Its
Subversion of the Host Innate Immune Response to Infection. Toxins (Basel).
2021;13: 880. doi:10.3390/toxins13120880

39. Sutterwala FS, Mijares LA, Li L, Ogura Y, Kazmierczak BI, Flavell RA. Immune
recognition of Pseudomonas aeruginosa mediated by the IPAF/NLRC4
inflammasome. J Exp Med. 2007;204: 3235–3245. doi:10.1084/JEM.20071239

40. Rada B, Jendrysik MA, Pang L, Hayes CP, Yoo DG, Park JJ, et al. Pyocyanin-
Enhanced Neutrophil Extracellular Trap Formation Requires the NADPH
Oxidase. PLoS One. 2013;8: e54205. doi:10.1371/JOURNAL.PONE.0054205

41. Thanabalasuriar A, Scott BNV, Peiseler M, Willson ME, Zeng Z, Warrener P, et
al. Neutrophil Extracellular Traps Confine Pseudomonas aeruginosa Ocular

Biofilms and Restrict Brain Invasion. Cell Host Microbe. 2019;25: 526-536.e4. doi:10.1016/J.CHOM.2019.02.007

42. Skopelja-Gardner S, Theprungsirikul J, Lewis KA, Hammond JH, Carlson KM, Hazlett HF, et al. Regulation of Pseudomonas aeruginosa-Mediated Neutrophil Extracellular Traps. Front Immunol. 2019;10. doi:10.3389/FIMMU.2019.01670/FULL

43. Ryu JC, Kim MJ, Kwon Y, Oh JH, Yoon SS, Shin SJ, et al. Neutrophil pyroptosis mediates pathology of P. aeruginosa lung infection in the absence of the NADPH oxidase NOX2. Mucosal Immunol. 2017;10: 757–774. doi:10.1038/mi.2016.73

44. Zhao Y, Yang J, Shi J, Gong YN, Lu Q, Xu H, et al. The NLRC4 inflammasome receptors for bacterial flagellin and type III secretion apparatus. Nature. 2011;477: 596–602. doi:10.1038/nature10510

45. Maltez VI, Tubbs AL, Cook KD, Aachoui Y, Falcone EL, Holland SM, et al. Inflammasomes Coordinate Pyroptosis and Natural Killer Cell Cytotoxicity to Clear Infection by a Ubiquitous Environmental Bacterium. Immunity. 2015;43: 987–997. doi:10.1016/j.immuni.2015.10.010

46. Kumari P, Russo AJ, Wright SS, Muthupalani S, Correspondence VAR, Rathinam VA. Hierarchical cell-type-specific functions of caspase-11 in LPS shock and antibacterial host defense. 2021 [cited 6 May 2021]. doi:10.1016/j.celrep.2021.109012

47. Faure E, Mear J-B, Faure K, Normand S, Couturier-Maillard A, Grandjean T, et al. *Pseudomonas aeruginosa* Type-3 Secretion System Dampens Host Defense by Exploiting the NLRC4-coupled Inflammasome. Am J Respir Crit Care Med. 2014;189: 799–811. doi:10.1164/rccm.201307-1358OC

48. Cohen TS, Prince AS. Activation of inflammasome signaling mediates pathology of acute P. aeruginosa pneumonia. J Clin Invest. 2013;123: 1630–1637. doi:10.1172/JCI66142

49. Evavold CL, Hafner-Bratkovič I, Kagan JC. Downstream of gasdermin D cleavage, a Ragulator-Rag-mTORC1 pathway promotes pore formation and pyroptosis. bioRxiv. bioRxiv; 2020. p. 2020.11.02.362517. doi:10.1101/2020.11.02.362517

50. Rühl S, Shkarina K, Demarco B, Heilig R, Santos JC, Broz P. ESCRT-dependent membrane repair negatively regulates pyroptosis downstream of GSDMD activation. Science (80-). 2018;362: 956–960. doi:10.1126/science.aar7607

1276   51.   Bjanes E, Sillas RG, Matsuda R, Demarco B, Fettrelet T, Delaney AA, et al. The
1277   Card19 locus of murine chromosome 13 regulates terminal cell lysis downstream
1278   of caspase activation and Gasdermin-D cleavage Card19 locus regulates
1279   caspase-dependent cell lysis. bioRxiv. 2021; 2021.03.19.436207.
1280   doi:10.1101/2021.03.19.436207

1281   52.   Weinrauch Y, Drujan D, Shapiro SD, Weiss J, Zychlinsky A. Neutrophil elastase
1282   targets virulence factors of enterobacteria. Nature. 2002;417: 91–94.
1283   doi:10.1038/417091a

1284   53.   Warnatsch A, Tsourouktsoglou TD, Branzk N, Wang Q, Reincke S, Herbst S, et
1285   al. Reactive Oxygen Species Localization Programs Inflammation to Clear
1286   Microbes of Different Size. Immunity. 2017;46: 421–432.
1287   doi:10.1016/J.IMMUNI.2017.02.013

1288   54.   Kaminski A, Gupta KH, Goldufsky JW, Lee HW, Gupta V, Shafikhani SH.
1289   Pseudomonas aeruginosa ExoS Induces Intrinsic Apoptosis in Target Host Cells
1290   in a Manner That is Dependent on its GAP Domain Activity. Sci Reports 2018
1291   81. 2018;8: 1–15. doi:10.1038/s41598-018-32491-2

1292   55.   Vareechon C, Zmina SE, Karmakar M, Pearlman E, Rietsch A. Pseudomonas
1293   aeruginosa Effector ExoS Inhibits ROS Production in Human Neutrophils. Cell
1294   Host Microbe. 2017;21: 611-618.e5. doi:10.1016/J.CHOM.2017.04.001

1295   56.   Bagayoko S, Leon Icaza S, Pinilla M, Hessel A, Bordignon P-J, Moreau F, et al.
1296   Phospholipid peroxidation fuels ExoU phospholipase-dependent cell necrosis 1
1297   and supports Pseudomonas aeruginosa-driven pathology 2. bioRxiv. 2021;
1298   2021.02.17.431580. doi:10.1101/2021.02.17.431580

1299   57.   Galle M, Schotte P, Haegman M, Wullaert A, Yang HJ, Jin S, et al. The
1300   Pseudomonas aeruginosa Type III secretion system plays a dual role in the
1301   regulation of caspase-1 mediated IL-1β maturation. J Cell Mol Med. 2008;12:
1302   1767–1776. doi:10.1111/J.1582-4934.2007.00190.X

1303   58.   Metzler KD, Goosmann C, Lubojemska A, Zychlinsky A, Papayannopoulos V.
1304   Myeloperoxidase-containing complex regulates neutrophil elastase release and
1305   actin dynamics during NETosis. Cell Rep. 2014;8: 883–896.
1306   doi:10.1016/j.celrep.2014.06.044

1307   59.   Chen KW, Demarco B, Heilig R, Ramos SP, Grayczyk JP, Assenmacher C-A, et
1308   al. RIPK1 activates distinct gasdermins in macrophages and neutrophils upon
1309   pathogen blockade of innate immune signalling. bioRxiv. 2021;

2021.01.20.427379. doi:10.1101/2021.01.20.427379

60. Kayagaki N, Kornfeld OS, Lee BL, Stowe IB, O'Rourke K, Li Q, et al. NINJ1
mediates plasma membrane rupture during lytic cell death. Nature. 2021;591:
131–136. doi:10.1038/s41586-021-03218-7

61. Man SM, Hopkins LJ, Nugent E, Cox S, Glück IM, Tourlomousis P, et al.
Inflammasome activation causes dual recruitment of NLRC4 and NLRP3 to the
same macromolecular complex. Proc Natl Acad Sci U S A. 2014;111: 7403–
7408. doi:10.1073/pnas.1402911111

62. Demarco B, Grayczyk JP, Bjanes E, Roy D Le, Tonnus W, Assenmacher CA, et
al. Caspase-8-dependent gasdermin D cleavage promotes antimicrobial
defense but confers susceptibility to TNF-induced lethality. Sci Adv. 2020;6:
3465–3483. doi:10.1126/sciadv.abc3465

63. Li P, Allen H, Banerjee S, Franklin S, Herzog L, Johnston C, et al. Mice deficient
in IL-1β-converting enzyme are defective in production of mature IL-1β and
resistant to endotoxic shock. Cell. 1995;80: 401–411. doi:10.1016/0092-
8674(95)90490-5

64. Wang S, Miura M, Jung YK, Zhu H, Li E, Yuan J. Murine caspase-11, an ICE-
interacting protease, is essential for the activation of ICE. Cell. 1998;92: 501–
509. doi:10.1016/S0092-8674(00)80943-5

65. Martinon F, Pétrilli V, Mayor A, Tardivel A, Tschopp J. Gout-associated uric acid
crystals activate the NALP3 inflammasome. Nature. 2006;440: 237–241.
doi:10.1038/nature04516

66. Van Gorp H, Saavedra PHV, De Vasconcelos NM, Van Opdenbosch N, Walle L
Vande, Matusiak M, et al. Familial Mediterranean fever mutations lift the
obligatory requirement for microtubules in Pyrin inflammasome activation. Proc
Natl Acad Sci U S A. 2016;113: 14384–14389. doi:10.1073/pnas.1613156113

67. Rietsch A, Vallet-Gely I, Dove SL, Mekalanos JJ. ExsE, a secreted regulator of
type III secretion genes in Pseudomonas aeruginosa. Proc Natl Acad Sci U S A.
2005;102: 8006–8011. doi:10.1073/PNAS.0503005102

68. Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, Smith HO.
Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat
Methods 2009 65. 2009;6: 343–345. doi:10.1038/nmeth.1318

69. Headley MB, Bins A, Nip A, Roberts EW, Looney MR, Gerard A, et al.
Visualization of immediate immune responses to pioneer metastatic cells in the

lung. Nature. 2016;531: 513–517. doi:10.1038/nature16985

70.   Looney MR, Thornton EE, Sen D, Lamm WJ, Glenny RW, Krummel MF.
Stabilized imaging of immune surveillance in the mouse lung. Nat Methods.
2011;8: 91–96. doi:10.1038/nmeth.1543

71.   Eren E, Planès R, Bagayoko S, Bordignon P, Chaoui K, Hessel A, et al. Irgm2
and Gate-16 cooperatively dampen Gram-negative bacteria-induced caspase-
11 response. EMBO Rep. 2020;21: e50829. doi:10.15252/embr.202050829

72.   Deruelle V, Bouillot S, Job V, Taillebourg E, Fauvarque M-O, Attrée I, et al. The
bacterial toxin ExoU requires a host trafficking chaperone for transportation and
to induce necrosis. bioRxiv. 2020; 2020.11.04.367706.
doi:10.1101/2020.11.04.367706

1355