1 **Title**
2
3 **Nanoscopical analysis reveals an orderly arrangement of the presynaptic**
4 **scaffold protein Bassoon at the Golgi-apparatus**
5
6 **Authors**
7

8 Tina Ghelani[1,6], Carolina Montenegro[2,3,4,] Anna Fejtova[2,5] and Thomas Dresbach[1*]
9
1. Institute of Anatomy and Embryology, University Medical Center Göttingen,
Kreuzbergring 36, 37075 Goettingen, Germany
2. Department of Neurochemistry and Molecular Biology, Leibniz Institute for
Neurobiology, Magdeburg, Germany
3. Center of Behavioral Brain Sciences, Magdeburg, Germany
4. Institute for Pharmacology and Toxicology, Medical Faculty, Otto von Guericke
University, Magdeburg, Germany
5. Department of Psychiatry and Psychotherapy, University Hospital Erlangen,
Friedrich-Alexander Universität Erlangen-Nürnberg, Erlangen, Germany
6. Present address: Department of Biology / Chemistry / Pharmacy, Free University
of Berlin, Takustr. 6, D-14195 Berlin
21
22 *** Correspondence:**
23 Thomas Dresbach
24 thomas.dresbach@med.uni-goettingen.de

25
26
27 **Abstract**
28

Bassoon is a core scaffold protein of the presynaptic active zone. In brain synapses,
the C-terminus of Bassoon is oriented toward the plasma membrane and its N-
terminus is oriented towards synaptic vesicles. At the Golgi-apparatus Bassoon is
thought to assemble active zone precursor structures, but whether it is arranged in an
orderly fashion is unknown. Understanding the topology of this large scaffold protein
is important for models of active zone biogenesis.
35
Using stimulated emission depletion nanoscopy in cultured hippocampal neurons, we
found that an N-terminal intramolecular tag of recombinant Bassoon, but not C-terminal
tag, colocalized with markers of the trans-Golgi network. The N-terminus of Bassoon
was located between 48 nm and 69 nm away from TGN38, while its C-terminus was
located between 100 nm and 115 nm away from TGN38. Sequences within the first 95
amino acids of Bassoon were required for this arrangement.
42
43 Our data are consistent with a model, in which the N-terminus of Bassoon binds to the
44 membranes of the trans-Golgi network, while the C-terminus associates with active
45 zone components, thus reflecting the topographic arrangement characteristic of
46 synapses also at the Golgi-apparatus.

48
49
50

Introduction

Scaffold proteins recruit and anchor molecules to subcellular sites. Due to their multi-domain, modular structure, they bind and regulate multiple proteins to coordinate biochemical reactions in space and time. Employing scaffold proteins is a fundamental principle of cell function, operating during protein folding, receptor and signaling molecule clustering, and at cell-cell-junctions (Good et al., 2011).

Synapses are asymmetric cell-cell junctions assembled and regulated by scaffold molecules. On the presynaptic side, a set of synaptic scaffold proteins confines the docking of synaptic vesicles and the exocytotic release of neurotransmitter from these vesicles to specialized sites of the axonal plasma membrane, called active zones. Several families of scaffold proteins operate at active zones, including RIMs, RIM binding proteins, Munc13s, α-liprins and ELKS/CAST/ERC proteins, as well as the particularly large scaffold proteins Bassoon and Piccolo (Sudhof, 2012; Gundelfinger et al., 2016). One way by which the presynaptic machinery acts is through RIMs, which recruit both voltage gated calcium channels and Munc13s, a family of proteins essential for making synaptic vesicles tethered at the active zone fusion competent (Südhof, 2012; Imig et al., 2014; Acuna et al., 2016). Bassoon regulates this core transmitter release machinery, at least at some synapses, by selectively recruiting the P/Q type of voltage gated calcium channels and by speeding up synaptic vesicle reloading to release sites during ongoing activity (Davydova et al., 2014; Hallermann et al., 2010; Mendoza-Schulz et al., 2014). In addition to regulating transmitter release, Bassoon and Piccolo maintain synaptic integrity by reducing the proteasome- and autophagy-mediated degradation of presynaptic molecules (Waites et al., 2013; Okerlund et al., 2017; Hoffmann-Conaway et al., 2020; Montenegro-Venegas et al., 2021). At the electron microscopy level, the multimolecular complex of presynaptic scaffold proteins manifests as a meshwork of filamentous structures termed the presynaptic particle web (Philips et al., 2001) or cytomatrix of active zones, i.e., CAZ (Cases-Langhoff et al., 1996; Garner et al., 2000; Dresbach et al., 2001).

Bassoon is a particularly large CAZ molecule, comprising 3938 amino acids in the rat, and 3926 amino acids in humans (tom Dieck et al., 1998). It shares 10 regions of sequence homology with Piccolo/Aczonin (Fenster et al., 2000; Wang et al., 1999). Light microscopy super-resolution studies and electron microscopy studies have revealed that Bassoon and Piccolo are oriented in a particular way at synapses, with their C-termini closer to the active zone plasma membrane than their N-termini (Dani et al., 2010; Limbach et al., 2011). Thus, Bassoon and Piccolo appear to be extended proteins with a parallel orientation at synapses, consistent with the assumption that they may represent some of the filamentous CAZ structures observed by electron microscopy.

Using recombinant Bassoon constructs (Dresbach et al., 2003) we previously imaged the incorporation of Bassoon into nascent synapses and its turnover at existing synapses (Shapira et al., 2003; Bresler et al., 2004; Tsuriel et al., 2006; Tsuriel et al., 2009). In the course of these studies, we also found that Bassoon– in addition to being a CAZ protein – is associated with the Golgi-apparatus, and that associating with the Golgi-apparatus is a prerequisite for the subsequent trafficking of Bassoon to synapses (Dresbach et al., 2006). Indeed, Bassoon, Piccolo and ELKS/CAST/ERC are all detected at the Golgi apparatus, and appear to exit the Golgi apparatus on transport vesicles that may carry CAZ material to synapses (Zhai et al., 2001; Maas et al. 2012). Unlike at synapses, the nanostructure of Bassoon at its second prominent subcellular localization, i.e., the Golgi-apparatus, has not been investigated. Here, we created a

101 new generation of Bassoon constructs and determined their localization and
102 arrangement at the Golgi-apparatus by stimulated emission depletion (STED)
103 microscopy. We find that Bassoon is an extended molecule at the trans-Golgi-network
104 (TGN) with its N-terminus closer to the TGN than its C-terminus.
105
Results

Characterizing second generation full-length Bassoon constructs

We first aimed at improving three features of recombinant Bassoon:
a) Faithful expression of the full-length protein including its C-terminal tag: when
expressed in neurons, Bassoon-1-3938-EGFP, in addition to producing punctate
synaptic fluorescence signals, also produces diffusely distributed green fluorescence,
presumably resulting from soluble EGFP or a soluble C-terminal fragment of Bassoon
with the EGFP-tag attached (Dresbach et al., 2003). We realized serendipitously that
this diffusely distributed green fluorescence also occurred when the EGFP coding
sequence was attached out of frame to the 3' end of Bassoon, suggesting that a cryptic
ribosomal entry site exists somewhere near the 3'-region of Bassoon or in the linker
located between Bassoon and EGFP. To prevent translation of the C-terminal EGFP-
tag we changed the linker sequence and removed the start ATG from the EGFP coding
sequence.
b) The accessibility of its N-terminus: Bassoon contains a functional consensus
site for N-myristoylation (Dresbach et al., 2003), so N-terminal tags might impair N-
myristoylation. Ideally, a tag designed to locate the N-terminal region of Bassoon
should be placed downstream of the N-myristoylation consensus site. To leave this
consensus site unaffected, we placed either RFP, CFP or GFP as an intramolecular
tag 97 amino acids downstream of the N-terminus of Bassoon, using an endogenous
HindIII site in the rat Bassoon cDNA. We will refer to these tags as "intramolecular N-
terminal" tags, to highlight both of their features, i.e., leaving the very N-terminus intact,
and placed close to it.
c) Its tendency to aggregate: Bassoon may form homodimers, and heterodimers
with Piccolo (Maas et al., 2012). Tags with an inherent capacity to dimerize could cause
aberrant oligomerization and generate non-functional aggregates of Bassoon. To
prevent this, we used monomeric fluorescent proteins, including RFP, CFP and the
A207K variant of EGFP. We use the term "EGFP" when referring to previously
generated constructs, which harbor standard EGFP, and we use the term "GFP" when
referring to new constructs, which harbor the monomeric variant. A schematic synopsis
of these new constructs is presented in Figure 1A.
To test the new constructs, we transfected dissociated rat hippocampal cultures with
the new full-length single-tagged and double-tagged Bassoon constructs on day 3 after
plating (day in vitro 3; DIV 3). We characterized the subcellular localization of these
Bassoon constructs in young (DIV5) and mature (>DIV13) neurons (Fig 1.), by
immunostaining fixed cultures using a single-domain antibody (nanobody) directed
against RFP, a polyclonal antisera directed against GFP, and monoclonal or polyclonal
antibodies directed against markers for subcellular structures. The extended
characterization of all single-tagged second-generation Bassoon constructs can be
found in supplementary Figure 1.
In the soma of young neurons, all constructs were readily detected at the Golgi
apparatus, labeled by the trans-Golgi-network (TGN) transmembrane protein TGN38
(Figure 1B—E). Previous reports observed similar localizations of endogenous and

149
Figure 1. Full-length Bassoon constructs localize at the TGN in young neurons, traffick to synaptic sites, and are
incorporated into the insoluble AZ scaffold of mature neurons. Panel **A** is a schematic diagram of full-length Bassoon
sequence compared to the sequence of full-length double- and single-tagged (either mRFP / mEGFP tagged) Bassoon
constructs where M stands for N-myristoylationsequence, Zn1 and Zn2 are the two zinc finger domains and cc1, cc2, cc3, are
the three predicted coiled-coil regions. Immunostained DIV7 (**B—E**) and DIV14 (**F—M**) hippocampal neurons transfected with
full-length double-and single-tagged Bassoon constructs with GFP, post a DIV3 lipofectamine transfection, are co-stained with
the TGN38 (**B—E**), synaptophysin (**F—I**) and Piccolo (**J—M**) markers. Panels **B, F,** and **J** represent 40X over views of the
transfections and **C, G** and **K** represent the zooms of their white square ROIs, respectively. Neurons in panels **B,C, J** and **K**
were briefly fixed in cold methanol prior to normal fixation, to quench the RFP and GFP autofluorescence. Panels **J—K** are
stained with TGN38 (purple) and a GFP antibody(green). Panels **J—M** are immuno-labeled for GFP antibody (green) and
Piccolo (purple) after a 90 second treatment of 0.1%Triton X-100 and five minute methanol wash. **N** is the colocalization
quantification of anti GFP immunofluorence of Bassoon for panels **F—M**; data are represented as mean ± SD, N=5 cells from
two separate experiments for each quantification. Scale bars 10µm (**B, F,** and **J**), 5µm (**C—E**) and 2µm(**G—I** and **K—M**).

recombinant Bassoon signals in young neurons (Dresbach et al., 2006; Maas et al.,
2012). To test the targeting of these constructs to synapses and their incorporation into
the CAZ matrix, we analyzed their localization in mature neuronal cultures. In DIV14
neurons immunostained for the tags and the synaptic vesicle marker synaptophysin or
the CAZ marker Piccolo, none of the constructs showed the diffusely distributed green
fluorescence associated with the first-generation Bassoon-EGFP (Dresbach et al.,
2003), and all of the constructs accumulated at synaptic sites (Fig. 1F-N). The degree
of colocalization with synaptophysin was 70.25% (± 18.23% SD) for the dually tagged
construct, 66.7% (± 12.9% SD) for mRFP Bassoon, and 62.64% (± 6.37% SD) for Bsn-
mRFP (Figure 1F—I,N and supplementary Figure 1). Likewise, these Bassoon
accumulations colocalized with the core CAZ scaffold protein Piccolo (Figure 1J—N
and supplementary Figure 1), further corroborating their localization to synapses. This
was true for the dually tagged (85.4% ± 18.42% SD) and the single-tagged Bassoon
constructs (69% ± 5.6% SD for mRFP-Bsn and 76.97% ± 10.16% SD Bsn-mRFP).

Active zone proteins become resistant to Triton X-100 extraction once they became
incorporated into the CAZ scaffold (Dresbach et al., 2003). When we applied a
0.1%Triton X-100 extraction to live neurons, followed by fixation and immunostaining
for the tags and for endogenous Piccolo, we found that the synaptic accumulations of
the recombinant proteins were indeed preserved. Colocalization with Piccolo was
85.43% (± 18.42% SD) for the dually tagged Bassoon, 64.85% (± 15.74% SD) for GFP-
Bsn, and 76.97% (± 24.01% SD) for Bsn-GFP). The resistance of the recombinant
Bassoon to the Triton X-100 treatment is indicative of their successful incorporation
into mature active zone scaffolds (Figure 1 J—N).

Visualizing the orientation of full-length Bassoon constructs with nanobodies
and super-resolution imaging

We then employed these constructs to study recombinant Bassoon by STED
nanoscopy. To take full advantage of super-resolution microscopy, we used camelid
antibodies, called nanobodies, to detect the tags. These anti-mRFP and anti-mEGFP
nanobodies are small (1.5 nm x 2.5 nm) single-domain molecules derived from one
heavy chain of an alpaca IgG antibody (Hamers-Casterman et al., 1993). They are
designed to identify a single epitope on the tertiary structure of mRFP and mEGFP
fluorophores (5nm diameters). These nanobodies were pre-coupled to two molecules
of organic ATTO-TEC dyes, each 2—3nm in size. The mRFP nanobody was coupled
to ATTO594, the mEGFP nanobody was coupled to ATTO647. Compared to traditional
primary and secondary antibodies, which create a 30 nm labeling distance from the
epitope site, the nanobody-ATTO dye complex generates a three times smaller label
cloud around the tags (Wildanger et al., 2009).

Our microscopy setup allowed us to record triple-immunofluorescence images,
with two dyes recorded in STED mode at a resolution of 20 nm, and a third dye
recorded in confocal mode. In the first set of experiments, we analyzed cultures
transfected with dually tagged Bassoon, asking three questions: 1) Do the nanobodies
allow for detection of the construct in STED mode? 2) Can we spatially resolve the N-
and the C-terminus of the dually tagged construct? 3) If so, is the recombinant
construct oriented as predicted, i.e., with the C-terminus closer to the active zone than
the N-terminus (Dani et al., 2010)?

Both the RFP-nanobody and the GFP-nanobody produced line-shaped or
crescent-shaped signals (Fig. 2A-D), as expected for the appearance of active zone
associated proteins at super-resolution (Dani et al., 2010). To make sure that what we
analyzed represented synaptic Bassoon, we only analyzed signals fulfilling two criteria:
they had to be line-shaped or crescent-shaped, suggesting that they represent side-
view synapses, and, in addition, they had to colocalize with postsynaptic marker
PSD95 immunofluorescence recorded in the confocal mode, corroborating that these
signals represent synaptic Bassoon (Fig. 2A-D). Comparing the nanobody-signals to
signals produced by a conventional monoclonal antibody directed against amino acids
756-1001 of Bassoon and detected by secondary antibodies revealed that the
nanobody signals and the indirect immunofluorescence signals colocalized when
analyzed by dual color STED microscopy. In addition, the nanobody signals produced
sub-clusters of fluorescence within the lines and crescents (Fig. 2A-D). This is
consistent with the assumption that their small size and their direct coupling to the
fluorescent dyes allows for greater detection precision compared to indirect
immunofluorescence with a conventional antibody. Overall, these data indicate that the
RFP- and GFP-nanobodies can localize the N- and C-terminus of recombinant
Bassoon molecules within the CAZ of mature neurons, and that the spatial precision

of detection at least equals, and may exceed, the spatial precision provided by indirect
immunofluorescence using a conventional antibody.
228

Figure 2. Visualizing the termini of full-length mRFP-Bassoon-mEGFP construct and its orientation with nanobodies, at mature synaptic sites. DIV21 mass hippocampal cultures transfected with the mRFP-Bassoon-mEGFP construct and imaged using two-color STED microscopy. The construct was visualized at synaptic sites with postsynaptic scaffold markers PSD95 and Shank2, using either the RFP-nanobody-Atto594 to visualize the RFP tag and the N-terminus of the Bassoon construct (**A, B,** and **G**) / the GFP-nanobody-Atto594 to visualize the GFP tag and the C-terminus of the Bassoon construct (**C—E**). Panels **A—D** are two-color-STED deconvolved (Deconv.) images of triple color stainings that label the endogenous presynaptic Bassoon signals with a traditional monoclonal antibody, nanoclusters of nanobody signals within the endogenous presynaptic Bassoon signals, and postsynaptic scaffold marker PSD95 (in confocal mode). Panels **E** and **G** show the localization of nanobody labelled C- and N-termini of mRFP-Bassoon-mEGFP and Shank2 in side-view images of its synapses. Distribution of localization points within a 350nm thick line profiles at the center of the synapse (as shown by the area within the dashed lines) were measured, fit with gaussian distributions and are plotted in panels **F** (C-terminus of tagged Bassoon and Shank2) and **H** (N-terminus of tagged Bassoon and Shank2). The distance between centroids of the two Gaussians defines the Bassoon-Shank2 distances. Scale bars 250nm (**A—G**).

The orientation of endogenous Bassoon at synapses in brain sections was previously
analyzed using conventional antibodies imaged by Stochastic Optical Reconstruction
Microscopy (STORM) (Dani et al., 2010, Held et al., 2020). These studies used the
monoclonal antibody to detect amino acids 756-1001 and/or a polyclonal antibody
against the C-terminal 330 amino acids of Bassoon. They showed that Bassoon
molecules possess an extended conformation within the CAZ scaffold and are oriented
with their C-termini closer to the synaptic cleft than their N-termini. Using our dual color
STED setup, we tested whether our constructs adopt a similar orientation in cultured
neurons. We found that they indeed possess a similarly extended conformation, with
the C-terminus 140 nm and the N-terminus 180 nm from the postsynaptic scaffold
marker Shank2 (Fig. 2E-H). These constructs, in combination with nanobodies and
STED microscopy, therefore, form an effective toolbox to visualize the nanoscopic
localization and orientation of recombinant Bassoon in neurons.

Visualizing the orientation of new full-length Bassoon constructs at the Golgi-
apparatus
How Bassoon is arranged at subcellular sites other than synapses has not been
investigated by nanoscopy thus far. Bassoon may form primordial scaffolds at the

level of the Golgi-apparatus (Dresbach et al., 2006; Maas et al., 2012). To arrive at a
more comprehensive understanding of the possible arrangement of Bassoon, we
used STED microscopy to determine the localization and orientation of our new
constructs at the Golgi-apparatus of immature neurons, at DIV7.

Figure 3. Full-length Bassoon localizes to the trans-Golgi network, not the trans-Golgi sub-compartment.
DIV7 hippocampal neurons were transfected with CFP-Golgi (*trans*-Golgi sub-compartment maker), full-length single-tagged mRFP-Bsn construct, and immunostained using GFP and/or RFP nanobodies against tagged constructs and from **A—E** with TGN38 (trans-Golgi network maker). Two-color STED images of both Golgi sub-compartment markers (**A—E**), CFP-Golgi and RFP-Bsn constructs (**F—J**), and RFP-Bsn at TGN38 (a *trans*-Golgi Network maker) (**K—O**). **A, F, K** show wide field overview of transfected construct, **B, G, L** the confocal zooms of the soma, insets **a** reflect the single channels and merged full STED deconvolved (Deconv.) images of **C—E**, **H—J**, and **M—O**, while insets **b** represents the zooms of STED images. Scale bars 4µm (**B, G** and **L**) and 1µm (**E, J** and **O**).

We began by testing which Golgi compartment recombinant Bassoon is located
to. The Golgi compartment is polarized across the flat stacks of cisternae/lamellae that
form sub-compartments at its *cis* side, which receives materials from the endoplasmic
reticulum (ER), and the Trans Golgi *(TG)* side, which sends them forward to their
destinations. Additional vesicular-tubular structures both on the *cis* and *trans* ends of
the Golgi stack, are known to form specialized compartments for cargo sorting at the
entry and exit sides of the Golgi. The *trans*-Golgi network (TGN) is a post-Golgi
compartment, following the TG sub-compartment that is involved in complex cargo
sorting mechanisms (Griffiths andSimons, 1986; Mellman and Simons, 1992). We
identified the TG sub-compartment and its neighboring compartment the TGN
separately and compared the extent of co-localization of Bassoon at these two
substructures in the soma of cultured neurons. To label the TG sub-compartment, we
expressed a CFP-tagged construct (named CFP-Golgi) containing the cytoplasmic
domain of ß-1,4-galactosyltransferase-11, which labels the last Golgi cisterna of the
trans-Golgi compartment (Wittenmayer, 2014). To label the TGN, we immunostained
for the marker TGN38. We detected recombinant constructs using the nanobodies,
and we detected TGN38 using conventional primary and secondary antibodies. STED

imaging revealed that the signals for the TG sub-compartment and TGN can be
resolved and occupy different localizations in the soma (Fig. 3 A-E). mRFP-Bsn
displayed virtually no colocalization with CFP-Golgi labeled TG lamellae (Fig. 3 F-J),
but colocalized extensively with TGN38 (Fig.3 K-O), indicating that this construct is
more closely associated with the TGN lamella than with the TG sub-compartment
lamellae.
We then sought to determine the localization – and possibly the orientation – of
Bassoon at the TGN in more detail. We first imaged the double-tagged Bassoon
construct (RFP-Bsn-GFP) in somas of DIV7 neurons labeled with TGN38. The
intramolecular N-terminal tag of the construct colocalized more extensively with
TGN38 than its C-terminal tag (Fig. 4A-K). On average, 67.5% (±12.5% SD) of the
signals coming from the intramolecular N-terminal tag, but only 29.8% (±7.7% SD) of
the signals coming from the C-terminal tag colocalized with TGN38 (Figure 4T). To
rule out that the difference could be caused by different avidities of the RFP- and the
GFP- nanobodies we transfected neurons either with RFP-Bsn or with Bsn-RFP and
detected the tags of both constructs using the RFP nanobody. Again, the
intramolecular N-terminal tag colocalized more extensively with TGN38 than the C-
terminal tag (Fig. 4L-S). In particular, the intramolecular N-terminal RFP showed 84.8%
(±4.6% SD) colocalization while the C-terminally located RFP showed 44.4% (±2.4%
SD) colocalization (Fig. 4T). Thus, both the dually tagged and the single tagged
constructs reveal a closer apposition of the N-terminal region of Bassoon with TGN38
than its C-terminal region.
Next, we quantified the distribution pattern of tagged Bassoon termini at certain
distances relative to the TGN. To this end, we defined two distance categories, i.e., 0-
100 nm and 101nm—1µm from TGN38 signals. We excluded Bassoon signals farther
than 1µm from TGN38 signals, assuming that these signals were not associated with
the TGN. The intramolecular N-terminal tags of both double- and single-tagged
Bassoon predominantly occupied the 0-100nm distance range. Fraction sizes were
0.67 ±0.1 SD for RFP-Bsn-GFP and 0.84 ±0.04 SD for RFP-Bsn. A smaller fraction of
each construct occupied the 101nm—1µm distance range. Here, fraction sizes were
0.33 ±0.1 SD for RFP-Bsn-GFP and 0.15 ±0.04 SD RFP-Bsn (Fig. 4U). In contrast to
the intramolecular N-terminal tags, the C-terminal tags of all constructs were more
evenly distributed between the two categories, with a tendency towards localization in
the 101nm-1µm range: a slightly larger fraction of RFP-Bsn-GFP (0.63±0.1 SD) and
Bsn-RFP (0.55±0.03 SD) resided within 101nm—1µm, and the remaining smaller
fraction of signals, for double- (0.37±0.1 SD) and single-tagged (0.45 ±0.03 SD)
Bassoon constructs, were present within the 0-100nm distance range (Fig. 4U). Thus,
all constructs were distributed in such a way that the intramolecular N-terminal tag had
a greater likelihood for detection within 100nm of TGN38 than the C-terminal tag.
Overall, both the percentage of colocalization and the distance distribution indicate an
orientation of recombinant Bassoon at the TGN where the N-termini are arranged
closer to the TGN than their C-termini.
Is this arrangement of Bassoon specific for the localization of Bassoon relative
to one particular Golgi-protein, i.e., TGN38, or does it reflect the orientation of Bassoon
relative to the TGN-compartment in general? To assess this, we imaged RFP-Bsn,
Bsn-RFP and RFP-Bsn-EGFP in neurons immunostained for another TGN marker,
i.e., Syntaxin6 (Syn6). As seen before with TGN38, the percentage of colocalization
with Syn6 was higher for the intramolecular N-terminal tags of RFP-Bsn-EGFP (71.8%)
and RFP-Bsn (75.8%) than for the C-terminal tags of RFP-Bsn-EGFP (23.8%) and

Figure 4 Orientation double-and single-tagged Bassoon constructs at the *trans*-Golgi network marker TGN38. DIV7 hippocampal neurons transfected with double-(**A—K**) and single-tagged (**L—S**) full-length Bassoon are immunostained with the *trans*-Golgi network marker TGN38 and an RFP-nanobody-Atto594 to visualize the RFP tag (**A—G, L—O** and **P—S**) / a GFP-nanobody-Atto594 to visualize the GFP tag (**H—K**). The experimental schematic (panels **A—C**) demonstrates the acquisition of two-color STED images wherein the wide field over view of the transfected neuron (**A**), its corresponding confocal over view of the soma, with GFP autofluorescence in blue, (**B**) and a 10μmX10μm inset scanned in STED mode to visualize the constructs' N—terminus (**C, D** and **L**) and C—terminus (**H** and **P**). Merged and single channel views of the zoom images of **E—G**, **I—K**, **M—O,** and **Q—S** are representations of the white ROIs in the STED images **D, H, L** and **P** respectively. **T** represents the colocalization quantification, and **U** represents the allocation quantification at and away from TGN lamella i.e. 0—100nm or 101nm—1μm, respectively. Data are represented as mean ± SD, N=10 cells from two separate experiments, statistically tested with a one-way ANOVA with the Tukey's multiple comparison's post-hoc test ***$p \le 0.001$. **V** is a Schematic representation of the proposed distribution of N- and C-terminal Bassoon immunosignals at the TGN. Scale bars 2μm (**B**) and 1μm (**C—S**).

Bsn-RFP (43.7%; Fig. 5A-Q). The distance analysis also revealed a similar pattern compared to what was seen for TGN38: a significantly larger fraction of intramolecular N-terminal than C-terminal Bassoon signals of single- (0.8±0.03 SD) and double-tagged (0.71±0.02 SD) Bassoon constructs were distributed within the 0-100nm distance range (Fig. 5R). Again, the C-terminal tags of both constructs were more evenly distributed between the two distance categories, and again with a tendency towards localization in the 101nm-1μm range, i.e., a slight majority of C-terminal Bassoon signals of the single- (0.62±0.03 SD) and double-tagged (0.65±0.04 SD) constructs were present in the 101nm—1μm range from the nearest Syn6 signal (Fig.

346 5R). These results show that Bassoon molecules possess similar colocalization and
347 distance distribution relative to both Syn6 and TGN38 markers of the TGN. This
348 suggests that Bassoon molecules are similarly oriented all over the TGN lamellae, with
349 the N-terminus more closely associated with the TGN membrane than the C-terminus.
350

Figure 5. Orientation double-and single-tagged Bassoon constructs at the *trans*-Golgi network marker Syntaxin 6 (Syn6).
Transfected DIV7 hippocampal neurons immunostained for either one or both termini of full-length Bassoon with RFP-
nanobody-Atto594 / GFP-nanobody-Atto647 and the Syn6 marker. Two-color, deconvolved, 10μmX10μm STED images and
their ROI zooms, show double-tagged and single-tagged Bassoon constructs in panels **A—D** and **I—L** (of which the N–termini
of constructs were imaged) and panels **E—H** and **M—P** (of which the C–termini of the constructs were imaged), respectively.
Graph **Q** quantifies the amount of Bassoon colocalization with Syn6 and graph **R** quantifies the Bassoon signal allocations at
and away from TGN38 lamella i.e., 0—100nm or 101nm—1μm, respectively. Data are represented as mean ± SD, N=10 cells
from two separate experiment, statistically tested with a one-way ANOVA with the Tukey's multiple comparison's post-hoc
test **p ≤ 0.01 and ***p ≤ 0.001. Scale bars 1μm (**A—P**).
360

Which domains of Bassoon are involved in orienting the molecule?

Next, we aimed at characterizing the features of Bassoon required for bringing the N-
terminal region of Bassoon into close apposition to the TGN. N-myristoylation is an
obvious possibility for anchoring a protein to membranes. Therefore, we generated a

point mutated version of the dually tagged Bassoon construct, called G2A-Bsn, where
we replaced glycine in position 2 with alanine. Dual color STED images of TGN38 and
the intramolecular N-terminal tag or the C-terminal tag of G2A-Bsn revealed the
previously seen pattern of high colocalization of the intramolecular N-terminal tag
(74.5%) and low colocalization (28.2%) of the C-terminal tag with TGN38 (Fig. 6A-H,
Q). Signals from the intramolecular N-terminal tag were predominantly (fraction size
0.75 ± 0.02 SD) located within 0 - 100nm from the TGN38, while the majority of signals
from the C-terminal tag (fraction size 0.75 ± 0.02 SD) were distributed within 101nm-
1µm from the TGN (Fig. 6R). A functional N-myristoylation site is thus not necessary
for the orientation of recombinant Bassoon at the TGN. These results do not exclude
the possibility that the G2A-mutant recombinant Bassoon dimerizes with endogenous
Bassoon, and that the intact endogenous Bassoon somehow helps orienting the N-
terminus of the mutant protein. However, when we expressed G2A-Bsn in cultures
obtained from mice lacking Bassoon the intramolecular N-terminal tag of G2A-Bsn was
still oriented towards the TGN, further corroborating that a functional N-myristoylation
site is not necessary for the orientation of Bassoon at the TGN (Suppl. Figure 2).
Since an intact N-myristoylation consensus site was not required for the
orientation of Bassoon, we wondered if any other sequences upstream of the
intramolecular N-terminal tag were required for the orientation of Bassoon. To test this,
we expressed the previously generated GFP-Bsn95-3938 construct aka GFP-95-Bsn.
In this construct, the N-terminal 94 amino acids are replaced by EGFP, but it
accumulates at the Golgi-apparatus and at synapses (Dresbach et al., 2003; Dresbach
et al., 2006). The EGFP tag of this construct, detected by our GFP-nanobody,
displayed remarkably low colocalization with TGN38 at the nanoscopical level (Fig. 6
I-L). Its colocalization with TGN38 was 36.2% and the majority of these signals
(0.60±0.04 SD) were located farther than 100 nm from the nearest TGN38 signals (Fig.
6Q,R), suggesting that sequences in the first 94 amino acids contribute to orienting
the N-terminal area of Bassoon towards the TGN.
A central region of Bassoon, comprising amino acids 2088-2563 and termed Bsn-GBR
(for Golgi-binding region), is required for targeting Bassoon to the Golgi-apparatus.
Thus, this region may contribute to bringing parts of Bassoon into close proximity to
the TGN. A construct consisting of these amino acids fused to the C-terminus of EGFP
indeed targets to the Golgi-apparatus, presumably via dimerization with endogenous
Bassoon (Dresbach et al., 2006; Maas et al., 2012). Where is this construct located
relative to TGN38 at the nanoscopical level? Our STED analysis revealed a relatively
low colocalization with TGN38 (41.5%) and predominant signal allocations (0.67±0.03
SD) in the 101nm—1µm distance range from TGN38 signals (Fig. 6. M-P and Q,R).
Thus, this region alone, while harboring Golgi-targeting capacity, cannot account for
the close apposition of the N-terminal Bassoon regions to the TGN.
Together, these results indicate that neither of the two regions of Bassoon equipped
with known Golgi-targeting sequences, i.e., the N-myristoylation consensus site and
amino acids 2088-2563, account for the particular orientation of Bassoon at the TGN.
Instead, the first 94 amino acids of Bassoon contribute to orienting the N-terminal
region of Golgi-associated Bassoon towards the TGN.

Estimating the extension of Bassoon molecules at the TGN

Having analyzed the distribution of N- and C-terminal tags over two distance categories, i.e., within 100 nm of TGN38 signals and between 101 nm and 1 µm from the TGN, we wondered whether we might be able to extract the average distance between a tag and TGN38 from the data sets. Figure 7A shows the distances of signals relative to TGN38 obtained from all Bassoon constructs and tags, displayed as violin plots.

The analysis shows that most signals were located within 220 nm from the TGN, irrespective of the construct and tag. Among these signals, two differences between the constructs were obvious: first, the median of the values for distance from TGN38 was smaller for the intramolecular N-terminal tags, both in the dually tagged construct

and in the single tagged construct, compared to all other constructs; second, wider
sections of the violin plot, representing a higher probability that data in the population
take on a certain value, indicated a clustering of intramolecular N-terminal signals close
to TGN38. In contrast, all other construct showed a more uniform distribution of signals
across the 0-220 nm range (Fig. 7A). We then analyzed the signals from this 0-220 nm
range to extract the average distances for each of the constructs and tags from TGN38
(Fig. 7B). In dually tagged Bassoon, average distances from TGN38 were 69 nm for
the intramolecular N-terminal tag and 115 nm for the C-terminal tag. Thus, amino acid
97 of Bassoon, where the intramolecular N-terminal tag is located, and amino acid
3938, where the C-terminal tag is located, are estimated to be 46 nm from each other.
In single tagged constructs, the intramolecular N-terminal tag was on average 48 nm
away from TGN38, the C-terminal tag was 100 nm away. This yields an estimated
distance of 52 nm between the average location of the two tags. The average distances
from TGN38 of GFP-Bsn95-3938 (106 nm) and Bsn-GBR (115 nm) show that the tags
of these deletion constructs occupy similar locations compared to the C-terminal tags
of full-length Bassoon constructs (Fig. 7B).
In summary, both transfection of a dually tagged construct and separate transfections
of single tagged constructs lead to similar conclusions: first, the intramolecular N-
terminal tag is located closer to TGN38 than the C-terminal tag; second, the average
distance from TGN38 is between 48 nm and 69 nm for the intramolecular N-terminal
tags; the average distance from TG38 is between 100 nm and 115 nm for the C-
terminal tags; third, the distance between the intramolecular N-terminal and the C-
terminal tag is estimated to be between 46 nm and 52 nm. Overall, this indicates an
orderly arrangement of Bassoon molecules with the N-terminus facing the TGN
membrane (Fig. 7C).

455
Figure 7. Distribution and average distances of all full-length and mutant Bassoon signals from the TGN38 reference maker.
Bean plots in panel **A** display all signal distributions of single- and double-tagged full-length and deletion constructs of
Bassoon from their nearest TGN38 marker. The light grey lines within each individual bean plot show individual observations
of Bassoon signals. The blue ROI represents the 220nm range that corresponds to 75% of the total Bassoon signals oriented
around the TGN lamella (**A**). The average distances of the Bassoon tags of all five constructs, within the 200nm distance range,
are plotted in **B**. **C** is a schematic diagram that shows the average distance N and C-termini of full-length Bassoon molecules
possess at the TGN lamella.
463
464 ## Discussion
465
Bassoon is a presynaptic scaffold protein predicted to be up to 80 nm long
(Gundelfinger et al., 2016). At active zones, Bassoon appears to have an extended
conformation with its C-terminus facing the plasma membrane (Dani et al., 2010;
Limbach et al., 2011). At synapses, Bassoon may thus represent one of the
filamentous structures or dense projections characteristic of active zones (Phillips et
al., 2001; Dresbach et al., 2001). Here, we find that recombinant Bassoon expressed
in cultured hippocampal neurons, has an extended conformation at the Golgi-
apparatus. At this subcellular site, the N-terminus of Bassoon faces the TGN

membrane. The fact that Bassoon is an extended protein already at this early stage of
its trafficking path supports the notion that primordial active zone scaffolds assemble
at the Golgi-apparatus (Dresbach et al., 2006). Its orientation relative to membranes -
i.e., with the N-terminus facing the TGN membrane, and the C-terminus facing the
active zone plasma membrane – adds new insights and raises questions regarding the
topology of its trafficking from the Golgi apparatus to active zones.
480
Design of new Bassoon constructs
We performed this study using a new generation of full-length Bassoon constructs.
These constructs were designed to have several advantages. First, the C-terminal tag
of these constructs does not produce diffusely distributed fluorescence anymore. This
allows for better detection of punctate signals because these are not hidden in a cloud
of homogeneous cytoplasmic background fluorescence anymore. We do not know
what caused this background in previous constructs (Dresbach et al., 2003).
Proteolytic cleavage of the C-terminal tag is a possible cause. But here we found that
there is diffusely distributed green fluorescence when the GFP sequence is attached
out of frame to the 3' end of full-length Bassoon, suggesting that a cryptic translation
initiation sequence may be present in the 3' area of the classical construct. By
removing the start codon of GFP we eliminated this possibility. Second, to reduce the
probability of artificial aggregation we used monomeric versions of fluorescent
proteins, i.e., mRFP and the A207K variant of EGFP. Third, all new constructs have a
free N-terminus, thus leaving the N-myristoylation consensus site of Bassoon
(Dresbach et al., 2003) intact.
497
To introduce a tag close to the N-terminus, but outside the N-myristoylation consensus
sequence, we placed mRFP or mGFP downstream of amino acid 97 in the new
Bassoon constructs. In particular, the intramolecular tag is preceded by amino acids
1-97 of Bassoon and followed by amino acids 95-3938 of Bassoon. Originally, this
location was chosen because it contains a conveniently located HindIII cloning site in
rat Bassoon cDNA. Later it became clear that the first 94 amino acids appear to be a
structurally compact unit without persistent folding (Gundelfinger et al., 2016). Glycine
and proline constitute 48 percent of the first 94 amino acids of rat Bassoon (25 glycine
residues and 20 proline residues), while this percentage steeply drops to 18 percent in
amino acids 95-197. Because of the location of the intramolecular tag, the new full-
length constructs are expected to combine the properties of two previously
characterized constructs: Amino acids 95-3938 preceded by EGFP are correctly
targeted to the Golgi-apparatus and to synapses (Dresbach et al., 2003; Bresler et al.,
2004; Dresbach et al., 2006; Tsuriel et al., 2009), and amino acids 1-97 followed by
EGFP retain targeting capacity for the Golgi-apparatus through their N-myristoylation
site (Dresbach et al., 2003; Dresbach et al., 2006). To avoid problems sometimes
associated with co-expression, one of our new constructs is dually tagged, with an N-
terminal intramolecular mRFP and C-terminal mGFP.
516
Validating the constructs
All constructs accumulated at the Golgi-apparatus and at presynaptic terminals, as
expected (Dresbach et al., 2003; Dresbach et al., 2006). Moreover, their resistance to
extraction with Triton X100 indicates their proper incorporation into the CAZ network
at synapses (Dresbach et al., 2003). Colocalization levels with the synapse markers
Piccolo and Synaptophysin was between 62 percent and 85 percent, and is likley
underestimated due to rigorous thresholding. The remaining, non-synaptic puncta are
probably mobile transport units. This is expected for Bassoon, which traffics on mobile

units (Shapira et al., 2003, Bresler et al. 2004) and undergoes activity-induced synapse
recruitment to synapses from kinesin-1 – motor dependent axonal carriers (Cai et al.,
2007). In addition, we verified by conventional and by STED microscopy that the dually
tagged construct was enriched at synapses based on co-localization, in confocal
images, with the postsynaptic markers PSD95 and Shank-2/ProSAP1. STED
microscopy revealed that this construct was oriented with its C-terminus towards the
active zone, as previously shown for endogenous Bassoon (Dani et al., 2010). Thus,
we conclude that the novel constructs undergo proper subcellular targeting, that the
dually tagged construct adopts the predicted orientation at synapses, and that the two
tags can be spatially resolved at subcellular sites where Bassoon has an extended
conformation.
Our use of recombinant protein bears the inherent caveat of overexpression.
However, recombinant Bassoon has been used widely by us and others to monitor
synapse assembly and turnover (Shapira et al., 2003; Cai et al., 2007; Lee et al., 2008;
Tsuriel et al., 2009; Matz et al., 2010). In addition, our recombinant proteins
accumulated at the expected subcellular sites, and the dually tagged construct showed
the expected orientation at the synapse. The design of the constructs also provides
several advantages. Under ideal expression conditions the location of the N-terminal
intramolecular tag (downstream of amino acid 94) and the C-terminal tag (downstream
of amino acid 3938) should report the maximal extension of Bassoon more accurately
compared to the routinely used antibodies, whose epitopes are located between amino
acids 756 and 1001 and between amino acids 3908 and 3938 (e.g., Dani et al., 2010).
Indeed, we observed a distance of 46—52 nm between the tags, compared to 30 nm
distance calculated for the two epitopes in Dani et al., 2010. This is consistent with the
tags being farther apart within recombinant Bassoon than the two epitopes are in
endogenous Bassoon. We emphasize, however, that Dani et al., 2010 performed an
extensive 3D analysis of a large number of synapses in brain sections, while we only
performed a proof-of-principle analysis for our construct, analysing a set of synapses
that appeared to be visible as side views in our cultured neurons.
Using nanobodies to detect the tags provides an additional advantage: because
of their small size and direct coupling to fluorophores, nanobodies bring the fluorescent
dye closer to the epitope compared to indirect immunofluorescence using primary and
secondary antibodies. The subclusters of recombinant Bassoon we detected with
nanobodies are consistent with this increased spatial resolution. Maybe these
nanobodies detect Bassoon arranged in the presynaptic particle web (Phillips et al.,
2001). Overall, we conclude that our analysis of recombinant Bassoon at synapses
yields results consistent with previous observations and shows that our approach
provides at least similar spatial resolution as previous nanoscopical approaches.
Based on these assumptions we conducted our detailed analysis of recombinant
Bassoon at the Golgi-apparatus.
565
Trans Golgi versus trans-Golgi network
STED microscopy revealed that recombinant Bassoon colocalizes with two TGN
markers, i.e., TGN38 and Syntaxin-6, but not with a TG sub-compartment marker. This
supports and extends previous data showing that both endogenous and recombinant
Bassoon colocalize with TGN38 and Syntaxin 6 upon confocal analysis (Dresbach et
al., 2006; Maas et al., 2012). This further corroborates that the recombinant protein
faithfully represents the localization of endogenous Bassoon. In addition, it narrows
down the exact localization of recombinant Bassoon by showing that it is more closely
associated with the TGN than with the sub-compartment. This observation is
particularly insightful because the recombinant sub-compartment marker we have

used here was shown by electron microscopy to selectively label the most "trans"
located lamellae of the Golgi stack, i.e., the one immediately preceding the TGN
(Wittenmayer, 2014).
579

Localization and orientation of full-length Bassoon at the trans-Golgi network
A key finding of our study is that the N-terminal, intramolecular tag of Bassoon was
located closer to the TGN than the C-terminal tag. This was true both for the dually
tagged Bassoon and for the two constructs that carried single mRFP tags. In addition,
it was true both when we used TGN38 and when we used Syntaxin-6 as TGN markers.
Finally, both colocalization analysis and the distance distribution of the Bassoon
constructs indicated this. Thus, recombinant Bassoon is an extended protein located
at the TGN, with the N-terminal area closer to the TGN than the C-terminus, and the
majority of Bassoon molecules have this orientation.
The distance distribution also revealed that the N-terminal, intramolecular, tag
had a higher likelihood of being located within 100 nm nanometers from the TGN than
being located inside the 100 nm – to 1 µm range, whereas the C-terminal tag appeared
to be distributed more evenly between the two distance regimes. On average, the two
tags were located 46 nm away from each other in the dually tagged construct, and 52
nm away from each other when the two single tag constructs were compared. A
distance of approximately 50 nm between the intramolecular tag and the C-terminal
tag is well within the range of the 80 nm maximal extension of Bassoon predicted by
Gundelfinger et al. (2016). It is also consistent with the two tags being farther apart
within the primary structure of Bassoon than the two antibody epitopes used in Dani et
al. (2010), who estimated a distance of 30 nm between those epitopes based on
STORM data. Taken together, our data indicate that recombinant Bassoon is an
extended protein located at the TGN, with its N-terminal area oriented towards the
TGN membrane and the C-terminus farther away. In addition, they suggest an average
distance of 46-52 nm between the intramolecular tag located downstream of amino
acid 97 and the C-terminal tag located downstream of amino acid 3938 of rat Bassoon.
605

Localization and orientation of deletion constructs
Bassoon includes at least two regions with binding capacity for the Golgi-apparatus,
including the myristoylated N-terminus and the region spanning amino acids 2088-
2563 called the Golgi-binding region (GBR). Using conventional epifluorescence
microscopy, we had previously detected constructs encoding either one of these
regions at the TGN (Dresbach et al., 2003; Dresbach et al., 2005). The results obtained
from analyzing mutated constructs of Bassoon provided some novel insights and
raised new questions.
First, the myristoylation-deficient G2A point mutant was still oriented with its N-
terminus towards the TGN, even in Bassoon knockout cultures. Thus, insertion of
myristic acid into the lipid bilayer is not required for the orientation of the N-terminal
region of Bassoon towards the TGN. Through its second coiled-coil domain the G2A-
mutant full-length protein may still bind to endogenous Piccolo (Maas et al., 2012). But
it is unlikely that this helps orienting the N-terminus of Bassoon towards the TGN,
because another construct, GFP-Bsn95-3938, that is also predicted to bind to
endogenous Bassoon and Piccolo, showed an aberrant orientation. Thus, N-
myristoylation appears to be dispensable for the orientation of Bassoon.
Second, as mentioned above, GFP-Bsn95-3938 was less closely associated
with the TGN than the intramolecular tag in the full-length Bassoon constructs. This
shows that the N-terminal 97 amino acids of Bassoon are essential for orienting the N-
terminus of Bassoon towards the TGN. It is likely that the unusually high percentage

of glycine and proline characteristic of this region of Bassoon contributes to this, but
the mechanisms and putative binding partners providing this orientation have yet to be
discovered. Surprisingly, the tag in GFP-Bsn95-3938 was located unexpectedly far
away from the TGN and was rather distributed like the C-terminal tag of the full-length
construct. We do not know if some extensive bending of the N-terminal end of this
construct towards its C-terminus or a completely aberrant localization account for this.
Third, Bsn-GBR showed a similar localization. This construct includes the
second coiled-coil domain of Bassoon, located between amino acids 2246 and 2366
of rat Bassoon (tom Dieck et al., 1998). Bsn-GBR harbors binding capacity for an
unknown target site on the Golgi-apparatus and, in addition, dimerizes or oligomerizes
with endogenous Bassoon through the CC2 domain (Dresbach et al., 2006; Maas et
al., 2012). Therefore, we expected that this construct might bind to the TGN or to a
central region of endogenous Bassoon or both. In particular, we aimed to find out if
Bsn-GBR locates to the same TGN site that the N-terminus of Bassoon locates to.
However, this construct was located farther away from the TGN than the intramolecular
tag of Bassoon and had a similarly widespread distribution like the C-terminal tag of
full-length Bassoon.
What may account for this localization? Bassoon and Piccolo are required for
the biogenesis of Golgi-derived transport vesicles, called gPTVs, that also contain the
scaffold protein ELKS2 (Maas et al., 2012). Bassoon binds to ELKS2 through its CC3
regions, to the Golgi apparatus through at least two regions, and to CTBP1, a protein
involved in the fission of vesicles budding from the Golgi-apparatus. Thus, Bassoon is
endowed with binding capacity for proteins that together could, theoretically, mediate
the generation and fission of gPTVs. Overexpressing Bsn-GBR causes the
accumulation of Bassoon, Piccolo and ELKS2 at the Golgi-apparatus. Bsn-GBR
prevents forward trafficking of gPTVs either by preventing binding of endogenous
Bassoon to the Golgi-apparatus, or by impairing oligomerization of Bassoon and
possibly Piccolo (Maas et al., 2012). Hence, a complex situation may arise where
endogenous proteins gPTVs accumulate and where, in addition, endogenous proteins
may be misplaced, when Bsn-GBR is overexpressed. Therefore, the relatively
widespread distribution of Bsn-GBR at the nanoscopical level may reflect this construct
binding to its target sites in a condition of reduced gPTV exit from the Golgi.
Overall, our results show, that at the nanoscopical level the Bsn-GBR does not
bind to the same TGN-region as the N-terminus of Bassoon. Thus, the two Golgi-
binding regions of Bassoon seem to associate with distinct sites at the Golgi-
apparatus, and the N-terminal 95 amino acids fulfil a special role in orienting the N-
terminus towards the TGN.
664
Perspective: towards a topological scenario
At active zones, the C-terminus of Bassoon is located closer to the plasmamembrane
than the N-terminus (Dani et al., 2010; our study). Assuming that Bassoon travels to
active zones on Golgi-derived Piccolo-Bassoon transport vesicles (gPTVs; Zhai et al.,
2001; Maas et al., 2012) one might predict that the C-terminus of Bassoon is attached
to Golgi-membranes and subsequently to the gPTV membrane; deposition of Bassoon
at synapses, perhaps by exocytotic fusion of the transport vesicle with the presynaptic
plasmamembrane, would then directly place the C-terminus close to the active zone
membrane. This scenario is "simple" because it involves no topological
rearrangements, i.e. the C-terminus of Bassoon is attached to equivalent membranes
all along the trafficking route.
Here, we find that the N-terminus of Bassoon is oriented towards the TGN
membrane, while the C-terminus is located farther away from it. In the context of the

gPTV model (Zhai et al., 2001; Maas et al., 2012), this is consistent with the following scenario, which also involves no topological rearrangements: at the Golgi-apparatus, the N-terminus of Bassoon may be attached to TGN-associated synaptic vesicle precursor membranes, while the C-terminus may become attached – simultaneously or later – to gPTVs. In this way, Bassoon would travel out of the soma on gPTVs and at the same time carry along synaptic vesicle precursors via its N-terminal region. This speculative scenario is consistent with several observations. First, Bassoon constructs lacking N-terminal areas appear at synapses as small spots the size of Piccolo immunosignals, suggesting that these constructs incorporate into the active zone cytomatrix immediately adjacent to the plasmamembrane; in contrast, a construct comprised of the N-terminal 609 amino acids of Bassoon appears at synapses as larger spots similar to the size of entire synaptic vesicle clusters, suggesting that this N-terminal region of Bassoon may bind to synaptic vesicles (Dresbach et al., 2003). Second, clouds of clear-core vesicles, dense-core vesicles and Bassoon were detected by electron immune-electron microscopy in axons (Tao-Cheng, 2007). It is possible that the dense core vesicles may represent gPTVs while the clear-core vesicles represent synaptic vesicle precursors. Third, a substantial fraction of synaptic vesicles precursors, labelled by recombinant Synaptophysin and called synaptic vesicle transport vesicles (STVs) in this study, co-traffics with active zone precursors, labelled by recombinant Bassoon, as revealed by live imaging studies (Bury and Sabo, 2011; Bury and Sabo, 2015). Recently, a type of Golgi-derived precursor vesicle called PLV, for presynaptic lysosome related vesicles, was identified. PLVs are required for the transport of presynaptic material. They carry, in addition to lysosomal proteins, the synaptic vesicle protein VGlut1 and co-traffic with Bassoon (Vukoja et al., 2018). Thus, at the Golgi-apparatus Bassoon could connect to PTVs via its C-terminus and to synaptic vesicle precursors via its N-terminus, and in this way generate already at the Golgi-apparatus a topological scenario that is later encountered at active zones. Whether this scenario holds true will need to be investigated in future studies.

Materials and Methods

Animals
Cells and tissues used in the study were obtained from bassoon gene trap (Bsn^{GT}) (Hallermann et al., 2010); mouse strains backcrossed over more than 10 generations to C57BL/6N. Bsn^{GT} mice were obtained from Omnibank ES cell line OST486029 by Lexicon Pharmaceuticals, Inc. (The Woodlands, TX). All experiments were performed in accordance with the European Committees Council Directive (86/609/EEC) and approved by the local animal care committees (Landesverwaltungsamt Sachsen-Anhalt, Germany, and the State Government of Lower Saxony, Germany.

Antibodies
The following antibodies were used for immunocytochemistry (IC): mouse anti-Bassoon (1:500 ENZO life systems), rabbit anti-Piccolo(1:200 Synaptic systems), mouse anti-TGN38 (1 to 500BD-Transduction Laboratories), mouse anti-Syntaxin 6 (1:300 Abcam), chicken anti-GFP (1: 3000 Abcam), rabbit anti-GFP (1: 1000 Abcam), mouse anti-Synaptophysin (1: 1000 Sigma Aldrich), guinea pig anti-SHANK2 (1 to 1000 Synaptic systems), Nanobodies: RFP-Booster-Atto594 and GFP-Booster-Atto647 (1:300 Chromotek), Mouse AlexaFluor®647 (1:1000 for Epifluorescence microscopy/1:100 for STORM Invitrogen), Chicken Cy5.5 (1:150 for STORM

729 Jackson/Invitrogen), Chicken and Rabbit Alexa Fluor®488 (1:1000 Invitrogen), ATTO-
730 TEC dyes: Mouse AttoKK1212 and Mouse Atto594 (1: 100) and Abberior dyes: Mouse
731 STAR 638, Mouse STAR 635p and Rabbit STAR 639 (1: 100).
732
Full-length Bassoon constructs
Full-length rat Bassoon constructs mRFP-Bsn-mEGFP, mEGFP-Bsn, Bsn-mEGFP,
mRFP-Bsn, Bsn-mRFP, and mutant G2A-mRFP-Bsn-mEGFP were created in the
ampicillin-resistant pCS2$^+$ vector backbone and designed to include either an
intramolecular mRFP/mEGFP tag and/or a mEGFP/mRFP fused to the C-terminus of
Bassoon. The intramolecular tag was created by gene synthesis in a way that its coding
sequence, after insertion into the HindIII site of rat Bassoon, was preceded by amino
acids 1-97 of Bassoon and was followed by amino acids 95-3938 of Bassoon. We
generated the G2A-mRFP-Bsn-mEGFP myristoyl mutant by inserting a point mutant
at the second amino acid position, thereby replacing a glycine amino acid with alanine
in the sequence of the full-length mRFP-Bsn-mEGFP construct. The plasmid called
"CFP-Golgi" (containing amino acids 1–81 of ß-1,4glycosyltransferase) was originally
purchased from Clontech.
Bassoon-mRFP was cloned by replacing mEGFP of the Bassoon-mEGFP construct
with an mRFP tag using the Mlu and SpeI sites. Similarly, mEGFP-Bassoon was
cloned by replacing mRFP-Bassoon by inserting a mEGFP, at the intramolecular 97th
amino acid position, by using the HindIII site. CFP-ß-1,4glycosyltransferase (amino
acids 0—81), also known as CFP-Golgi was generously provided by also Craig Garner.
Bassoon-mRFP was cloned by replacing mEGFP of the Bassoon-mEGFP construct
with an mRFP tag from the Mlu-mRFP-SpeI-PUC vector (Clonetech). Both constructs
were linearized via a Mlu and SpeI digestion, followed by a CIP dephosphorylation and
ligated. The ligation mix was transformed into XL1 Blue competent cell and was grown
on LB–ampicillin plates. Extracted DNA from the colonies grown on the plates were
sequenced with Bsn_FW_1, Bsn_FW_3c, Bsn_FW_4, and Bsn_FW_5 sequencing
primers to confirm colonies the successfully cloned and direction of insert in the
construct. Similarly, we cloned mEGFP-Bassoon by replacing mRFP-Bassoon by
inserting a mEGFP at the intramolecular 97th amino acid position by using the mEGFP-
HindIII-PUC vector (Clonetech) and a HindIII digestion followed by the same protocol
as described for Bassoon-mRFP generation.
762
Bsn_FW_1   CTAATGGGAGGTCTATATAAG
Bsn_FW_2   AGCACTAGCTGGCGGCGGACA
mRFP-FW-3a   GTAATGCAGAAGAAGACCATG
mRFP-Rev-3b   CATGGTCTTCTTCTGCATTAC
Bsn_FW_3c GGGCTTCAAGTGGGAGCG
Bsn_FW_4   GGGCCAGGAGGAGACAGACG
Bsn_FW_5   GCTCCAAACCGGCAGCCAAAG
770
Primary hippocampal neuron cultures
Rat cultures: E19 rat hippocampi were dissected as previously described (Dresbach
et al., 2006). Hippocampi were dissociated by a 20 min of trypsin treatment at 37°C
and trituration. 50,000 dissociated neurons/cm^2 were grown on poly-lysine coated
coverslips (Sigma Aldrich) in Neurobasal medium enhanced with 2% B-27 and 0.5%
L-Glutamine (Life technologies). Primary hippocampal neurons growing on either
12mm or 18mm coverslips, in 24-well or 12-well plates, respectively, were transfected
with the calcium phosphate method at DIV3 and fixed, with 4% paraformaldehyde, for
imaging of mature DIV15—30 transfected cultures. The protocol for which was

performed as described previously[3] (Figures 1—2, Suppl.1). Neurons were prepared
for calcium phosphate transfection by replacing and saving the conditioned medium
with 500μl (for 12mm coverslips) or 750μl (for 18mm coverslips) Optimem (Life
Technologies) at 37°C and incubated for 20—30min. A DNA mix containing 7μg of
DNA and 250mM $CaCl_2$ was vortexed during the dropwise addition of 105μl of
transfection buffer (274mM NaCl, 10mM KCl, 1.4mM Na2HPO4, 15mM glucose,
42mM HEPES, pH 7.06) and left to incubate for 20min at room temperature. 30μl per
well of the DNA-Calcium phosphate mix was applied onto neurons and incubated
further for 75min. We used three washes of pre-warmed Neurobasal medium to
complete the transfection, reinstated the transfected neurons into their original
conditioned medium, and incubated them at 37°C and 5% $CO2$ until the transfected
cultures had matured.
Mouse cultures were prepared as described (Montenegro-Venegaas et al. 2021).
Briefly, P0-P1 wild type mice cortexes were dissected to generate the feeder layer of
a sandwich culture. Hemispheres of cortexes with their meninges removed were
chopped up in 4.5 ml of HBSS and incubated for 15 minutes at 37°C in 2.5% trypsin
(without EDTA). These pieces were then washed in HBSS and dissociated in glia
medium that consisted of 90% plating medium and 10% DNAse (Invitrogen). Both
hemispheres of one brain were dissociated in 1 ml glia medium and plated in 10 ml of
plating medium that was identical to rat primary culture plating medium. The medium
was changed every 4—5 days, and the confluent glia were trypsinated, washed in
HBSS, and 5 ml of the glia were plated on a 6cm dish. Two P1 Bsn-/- knockout mice
and two wildtype Bsn+/+ littermates were prepped into dissociated primary
hippocampal neurons following the same protocol as was used for rat primary culture.
100 ml of 5000 hippocampal cells were plated on coated 18 mm round glass coverslips.
These coverslips were first incubated for 1 hour at 37°C and 5% $CO2$ and then
transferred, neurons facing down, onto dishes containing the feeder layer of glia and
5 ml of culturing medium (94% Neurobasal, 2% Glutamax (Invitrogen), 2% B27, 1%
NaPyr (0.1M), 1% Pen/Strep (0.1M). These coverslips were left to grow at 37°C and
5% $CO2$ and were treated with 2 μl Ara-C (Sigma) on day in vitro 1 (DIV1) and DIV3
to prevent glia overgrowth and were fed once a week with an exchange of 1 ml of fresh
culturing medium to maintain optimal growth of the culture.
812
Transfection
To visualize Golgi association of Bassoon constructs in hippocampal neurons (Figures
1, 3—6), we applied the Lipofectamine transfection method on DIV5/6 neurons and
fixed them on DIV6/7 neurons with 4% paraformaldehyde, as previously described.
Briefly, a conditioned medium of neurons, on 12mm coverslips, was exchanged with
500ml pre-warmed Neurobasal medium, containing 2% B-27 and 1% of 2mM L-
Glutamine, saved and incubated along with the neurons for 20—30min. A
Lipofectamine solution and a DNA solution of 25μl Optimem/well (Life Technologies)
with 1μl of Lipofectamine 2000/well (Invitrogen) and 1μg of plasmid DNA/well were
prepared and mixed after 10min room temperature incubation. The Lipofectamine-
DNA mix was further incubated for 20min at room temperature; 50μl/well of the solution
was dropwise applied on the neurons and incubated for 75min in 37°C and 5% $CO2$
conditions. The transfection was completed after three pre-warmed Neurobasal
washes and reinstating the transfected neurons in their conditioned medium at 37°C.
These neurons were fixed the next day for 20min in cold 4% paraformaldehyde solution
before immunocytochemistry was performed.
829
Immunocytochemistry

Primary cultures of hippocampal neurons were washed multiple times after the
paraformaldehyde fixation, blocked for 20 min with the primary antibody-buffer (10%
FBS, 5% sucrose, 2% albumin, 0.3% Triton X-100 in 1× PBS) at room temperature,
and stained with the primary antibodies, diluted in the primary antibody-buffer,
overnight at 4°C. Following multiple washes, secondary antibody dilutions were
prepared in the secondary antibody-buffer (0.3% Triton X-100, 5% sucrose, and 2%
albumin in 1× PBS) and applied on the coverslips for 1 hour, in darkness, at room
temperature. Three washed of 1xPBS and one of distilled water were performed on
the coverslips before being mounted on slides with DABCO-mowiol (Calbiochem) and
left to dry overnight.
841
842
Epifluorescence microscopy
An inverted Zeiss fluorescence microscope (Observer.Z1) with a Photometrics
CoolSnap HQ2 camera was used to image samples at a magnification of 40X and 63X.
The following filters from AHF were used: F46-000 for DAPI, F46-002 for GFP and
Alexa 488, F46-004 for Atto594 and Alexa 546 dyes, and F46-006 for AttoKK1212 and
Alexa 647. Exposure times of 500ns for F46-002 and F46-004 filters and 1000ns for
the F46-006 filter were applied. The images were processed with the Image J software
(NIH) (imagej.nih.gov/ij/) to generate scale-bar inserted RGB merged TIFF files for
further analysis with Imaris MeasurementPro software. Images in Figure 1 and
Supplementary 1 were adjusted for brightness and contrast in Image J, where needed,
calculated, and stamped with a suitable sized scale bar.
854
Colocalization analysis for conventional epifluorescence images
Merged multi-channel 40X light microscopy images were analyzed using MetaMorph
Offline Version 7.7.0.0 (Molecular Devices, Inc.). A threshold is set for each channel
followed by the generation of a mask for all channels, in three areas of size 25 pixels
long (representing 2 μm in the sample) and 4 pixels wide, per image. These area
masks were then overlaid in the arithmetic tool and divided to generate a third mask
containing only the population of fluorescence signals in the mask that do colocalize.
The amount of bassoon colocalized is represented as a percentage of the bassoon
colocalized population divided by the total bassoon population.
864
STED microscopy
STED images in Figures 2—6 were acquired on a custom-built two-color STED
microscope that includes a 1.4 NA 100× objective (PL APO HCX 100x 1.4-0.7 Oil,
Leica Microsystems, Wetzlar), and a 775nm STED laser (ELP-5-775-DG, IPG
Photonics Corporation, Oxford, MA, USA)[4]. The dyes were excited at wavelengths of
470nm, 595nm, and 640nm, while the fluorescence was detected with avalanche
photodiodes from 500-550nm, 600-640nm, and 660-720nm, respectively.
Corroborative confocal images were acquired using a LED illumination source, a
monochrome filter, and a camera (DMK41 AU02, The Imaging Source). The LED
illumination source for such overview images was manually installed every session,
wherein a 590nm LED was installed with the 700/60 fluorescence filter, and 640RDC
dichroic filter or a 490nm LED was installed (upon requirement) with the 450/60
fluorescence filter in the camera path. Images were taken at 300—700 mW STED
power, 4μW excitation power, dwell time of 30—100ms and a pixel size of 10nm. The
resolution regularly obtained for Atto594 antibodies was 25-40nm and for AttoKK1212
antibodies was 20-35nm (at 300mW and 700 mW STED-power, respectively). Figures
2-6 were acquired on the Abberior QuadSCAN two-color STED microscope. The setup

was equipped with a pulsed 775nm STED lazer and two pulsed excitation lazer-
sources at 594nm and 640nm integrated into an Olympus IX83 microscope. The setup
also included a 100× 1.4NA objective; a 4-color LED illumination source, a gated
avalanche photodiode (APD), and a wide monochrome field. A pixel size of 10 nm,
dwell time of 3ms, and three-line accumulations were applied to the images. All STED
images were acquired with the ImSpector Software[5] (Max Planck Innovation) and were
processed using the Richardson—Lucy deconvolution function. In combination with
the deconvolution processing, a 2D Lorentz function, that fits the full width at half
maximum (FWHM) fitted of the point spread function of each individual image to the
resolution estimate was used. These images were then analyzed using Imaris
MeasurementPro. Images for figures were adjusted for brightness and contrast with
Image J software (NIH).
894
Quantitative analysis of the probability and amount of colocalization and the
distribution of Bassoon constructs at the Golgi
Merged TIFF epifluorescence images or STED images were analyzed using Imaris
MeasurementPro 8.1(Bitplane AG.) software to ascertain the probability of
colocalization (Pearson's correlation coefficient), amount of colocalization, and
distribution pattern of signals within the images.
Soma Images in Figures 3—6 were analyzed within a ROI, generated by a free-hand
drawn mask, to exclude any signals in the image that was in the nucleus, outside the
cell soma, or in a neighboring neuronal process. Images were analyzed for the
probability of colocalization through the ImarisColoc module. The integrated Costes P-
Value approximation plugin[7], within the ImarisColoc module, first generates automated
thresholds for all the channels of all the images, which are subsequently used to
calculate the colocalization Pearson's correlation coefficient constants in the picture.
The amount of colocalization and distribution pattern of signals was calculated after
Imaris Spots generation. Imaris Spots, a built-in spot detection algorithm, was used to
generate objects for every punctuate signal, for all channels, in the images. These spot
objects were defined by the automated intensity threshold value for each channel
(calculated by the Costes P-Value approximation), signal diameter size range (10nm—
160nm for STED images and >200nm for light microscopy images), and an automated
splitting of signal clusters (defined as >120nm for STED images and >400nm for light
microscopy images), for each channel.
The amount of colocalization was calculated using Spots Colocalize, a MATLAB
extension in the Imaris Spots module, at a distance threshold of 0—100nm (for STED
images) and 0—350nm (for light microscopy images) from the spot centers.
The distribution pattern of AZ protein signals at the Golgi was extracted by applying
the distance transformation MATLAB extension[8], from the Imaris XT module to the
spots object information generated via the Imaris Spots module. This distance
transformation extension was applied on the Golgi label channel transforming the Golgi
voxel intensity data into spot coordinates. These spot coordinates were then used to
calculate the shortest distance of each AZ protein spot object to the object border of a
Golgi-label spot. The data for the shortest distances between all the AZ protein signals
and the border coordinates of the Golgi-label, and the total number of AZP signals
within 0—100nm or 101—1000nm distance ranges was extracted from the statistics of
the Imaris Spot module, statistically tested, and graphically represented with GraphPad
Prism.
The distribution pattern of Bassoon constructs in Figure 7A was generated in Python
and graphically visualized as violin plots, with a signal size upper limit of 600nm; to
clearly represent the distribution and permit visualization of the median and

interquartile ranges of the data. The average distance of each cell per set, in Figure 7B, was extracted from the Imaris Spots statistics and limited to the 220nm cut-off were plotted in GraphPad Prism.

Statistical analysis and representation of all resultant data were prepared in GraphPad Prism 5.02. Data are presented as mean ± SEM, and statistical differences were considered significant, strongly significant, and extremely significant at respective p values of $*p < 0.05$, $**p \leq 0.01$, and $***p \leq 0.001$. In a two-step statistical testing protocol, first, a one-way ANOVA test, with a post-hoc Tukey's multiple comparisons test, was performed, and for every significant difference observed between two relevant groups, an additional two-tailed, unpaired Student's t-test, with different variances, was also performed to reveal the same significant difference.

Quantitative analysis of distances between Bassoon's termini and the post synapse

All acquired, or rendered images were processed and visualized using ImageJ (imagej.nih.gov/ij/) and ImSpector software (Max-Planck Innovation) (Figure 2) or Daxviewer software (Mark Bates) (Figure 3). Line profiles were measured with ImageJ software along a 350nm thick line profiles. Inter-peak Bassoon-SHANK2 distances were determined after fitting a Gaussian distribution in GraphPad Prism.

To factor out the effect of the varying number of signals counted per size of hand-drawn mask in each image, the area in μm3 of the mask used, was divided by the signals measured per image.

Acknowledgements

We are thankful to Prof. Stefan W. Hell, Dr. Dirk Kamin and Dr. Fabian Göttfert for the training and guidance on the STED setups a Max Plank institute of Biophysical Chemistry. We thank Irmgard Weiss for technical assistance. This work was supported by the DFG Research Center for Nanoscale Microscopy and Molecular Physiology of the Brain (CNMPB) to TD.

References

Acuna, C., Liu, X., and Südhof, T.C. (2016). How to Make an Active Zone: Unexpected Universal Functional Redundancy between RIMs and RIM-BPs. Neuron *91*, 792–807.

Bresler, T., Shapira, M., Boeckers, T., Dresbach, T., Futter, M., Garner, C.C., Rosenblum, K., Gundelfinger, E.D., and Ziv, N.E. (2004). Postsynaptic density assembly is fundamentally different from presynaptic active zone assembly. J Neurosci *24*, 1507–1520.

Bury, L.A.D., and Sabo, S.L. (2011). Coordinated trafficking of synaptic vesicle and active zone proteins prior to synapse formation. Neural Dev *6*, 24.

Bury, L.A.D., and Sabo, S.L. (2016). Building a Terminal: Mechanisms of Presynaptic Development in the CNS. Neuroscientist *22*, 372–391.

Cai, Q., Pan, P.-Y., and Sheng, Z.-H. (2007). Syntabulin-kinesin-1 family member 5B-mediated axonal transport contributes to activity-dependent presynaptic assembly. J Neurosci *27*, 7284–7296.

Cases-Langhoff, C., Voss, B., Garner, A.M., Appeltauer, U., Takei, K., Kindler, S., Veh,
R.W., De Camilli, P., Gundelfinger, E.D., and Garner, C.C. (1996). Piccolo, a novel
420 kDa protein associated with the presynaptic cytomatrix. Eur J Cell Biol *69*, 214–
223.

Dani, A., Huang, B., Bergan, J., Dulac, C., and Zhuang, X. (2010). Superresolution
imaging of chemical synapses in the brain. Neuron *68*, 843–856.

Davydova, D., Marini, C., King, C., Klueva, J., Bischof, F., Romorini, S., Montenegro-
Venegas, C., Heine, M., Schneider, R., Schröder, M.S., et al. (2014). Bassoon
specifically controls presynaptic P/Q-type Ca(2+) channels via RIM-binding protein.
Neuron *82*, 181–194.

tom Dieck, S., Sanmartí-Vila, L., Langnaese, K., Richter, K., Kindler, S., Soyke, A.,
Wex, H., Smalla, K.H., Kämpf, U., Fränzer, J.T., et al. (1998). Bassoon, a novel zinc-
finger CAG/glutamine-repeat protein selectively localized at the active zone of
presynaptic nerve terminals. J Cell Biol *142*, 499–509.

Dresbach, T., Qualmann, B., Kessels, M.M., Garner, C.C., and Gundelfinger, E.D.
(2001). The presynaptic cytomatrix of brain synapses. Cell Mol Life Sci *58*, 94–116.

Dresbach, T., Hempelmann, A., Spilker, C., tom Dieck, S., Altrock, W.D., Zuschratter,
W., Garner, C.C., and Gundelfinger, E.D. (2003). Functional regions of the presynaptic
cytomatrix protein bassoon: significance for synaptic targeting and cytomatrix
anchoring. Mol Cell Neurosci *23*, 279–291.

Dresbach, T., Torres, V., Wittenmayer, N., Altrock, W.D., Zamorano, P., Zuschratter,
W., Nawrotzki, R., Ziv, N.E., Garner, C.C., and Gundelfinger, E.D. (2006). Assembly
of active zone precursor vesicles: obligatory trafficking of presynaptic cytomatrix
proteins Bassoon and Piccolo via a trans-Golgi compartment. J Biol Chem *281*, 6038–
6047.

Fejtova, A., and Gundelfinger, E.D. (2006). Molecular organization and assembly of
the presynaptic active zone of neurotransmitter release. Results Probl Cell Differ *43*,
49–68.

Fenster, S.D., Chung, W.J., Zhai, R., Cases-Langhoff, C., Voss, B., Garner, A.M.,
Kaempf, U., Kindler, S., Gundelfinger, E.D., and Garner, C.C. (2000). Piccolo, a
presynaptic zinc finger protein structurally related to bassoon. Neuron *25*, 203–214.

Garner, C.C., Kindler, S., and Gundelfinger, E.D. (2000). Molecular determinants of
presynaptic active zones. Curr Opin Neurobiol *10*, 321–327.

Good, M.C., Zalatan, J.G., and Lim, W.A. (2011). Scaffold proteins: hubs for controlling
the flow of cellular information. Science *332*, 680–686.

Griffiths, G., and Simons, K. (1986). The trans golgi network: sorting at the exit site of
the golgi complex. Science 234, 438–443.

Gundelfinger, E.D., Reissner, C., and Garner, C.C. (2015). Role of Bassoon and
Piccolo in Assembly and Molecular Organization of the Active Zone. Front Synaptic
Neurosci *7*, 19.

Hallermann, S., Fejtova, A., Schmidt, H., Weyhersmüller, A., Silver, R.A., Gundelfinger, E.D., and Eilers, J. (2010). Bassoon speeds vesicle reloading at a central excitatory synapse. Neuron *68*, 710–723.

Hamers-Casterman, C., Atarhouch, T., Muyldermans, S., Robinson, G., Hamers, C., Songa, E.B., Bendahman, N., and Hamers, R. (1993). Naturally occurring antibodies devoid of light chains. Nature *363*, 446–448.

Held, R.G., Liu, C., Ma, K., Ramsey, A.M., Tarr, T.B., De Nola, G., Wang, S.S.H., Wang, J., van den Maagdenberg, A.M.J.M., Schneider, T., et al. (2020). Synapse and Active Zone Assembly in the Absence of Presynaptic Ca2+ Channels and Ca2+ Entry. Neuron 107, 667-683.e9.

Hoffmann-Conaway et al., S., Brockmann, M.M., Schneider, K., Annamneedi, A., Rahman, K.A., Bruns, C., Textori-Taube, K., Trimbuch, T., Smalla, K.-H., Rosenmund, C., Gundelfinger, E.D., Garner, C.C., and Montenegro-Venegas, C. (2020). Parkin contributes to synaptic vesicle autophagy in Bassoon-deficient mice. Elife 9: e56590

Imig, C., Min, S.-W., Krinner, S., Arancillo, M., Rosenmund, C., Südhof, T.C., Rhee, J., Brose, N., and Cooper, B.H. (2014). The morphological and molecular nature of synaptic vesicle priming at presynaptic active zones. Neuron *84*, 416–431.

Lee, S.-H., Peng, I.-F., Ng, Y.G., Yanagisawa, M., Bamji, S.X., Elia, L.P., Balsamo, J., Lilien, J., Anastasiadis, P.Z., Ullian, E.M., et al. (2008). Synapses are regulated by the cytoplasmic tyrosine kinase Fer in a pathway mediated by p120catenin, Fer, SHP-2, and beta-catenin. J Cell Biol *183*, 893–908.

Limbach, C., Laue, M.M., Wang, X., Hu, B., Thiede, N., Hultqvist, G., and Kilimann, M.W. (2011). Molecular in situ topology of Aczonin/Piccolo and associated proteins at the mammalian neurotransmitter release site. Proc Natl Acad Sci U S A *108*, E392-401.

Maas, C., Torres, V.I., Altrock, W.D., Leal-Ortiz, S., Wagh, D., Terry-Lorenzo, R.T., Fejtova, A., Gundelfinger, E.D., Ziv, N.E., and Garner, C.C. (2012). Formation of Golgi-derived active zone precursor vesicles. J Neurosci *32*, 11095–11108.

Matz, J., Gilyan, A., Kolar, A., McCarvill, T., and Krueger, S.R. (2010). Rapid structural alterations of the active zone lead to sustained changes in neurotransmitter release. Proc Natl Acad Sci U S A *107*, 8836–8841.

Mellman, I., and Simons, K. (1992). The golgi complex: in vitro veritas. Cell 68, 829–840.

Mendoza Schulz, A., Jing, Z., Sánchez Caro, J.M., Wetzel, F., Dresbach, T., Strenzke, N., Wichmann, C., and Moser, T. (2014). Bassoon-disruption slows vesicle replenishment and induces homeostatic plasticity at a CNS synapse. EMBO J *33*, 512–527.

Montenegro-Venegas, C., Fienko, S., Anni, D., Pina-Fernandez, E., Frischknecht, R., and Fejtova, A. (2021). Bassoon inhibits proteasome activity via interaction with PSMB4. Cell Mol Life Sci 78: 1545-1563.

Okerlund, N.D., Schneider, K., Leal-Ortiz, S., Montenegro-Venegas, C., Kim, S.A.,
Garner, L.C., Waites, C.L., Gundelfinger, E.D., Reimer, R.J., and Garner, C.C. (2017).
Bassoon Controls Presynaptic Autophagy through Atg5. Neuron *93*, 897-913.e7.

Phillips, G.R., Huang, J.K., Wang, Y., Tanaka, H., Shapiro, L., Zhang, W., Shan, W.S.,
Arndt, K., Frank, M., Gordon, R.E., et al. (2001). The presynaptic particle web:
ultrastructure, composition, dissolution, and reconstitution. Neuron *32*, 63–77.

Sanmartí-Vila, L., tom Dieck, S., Richter, K., Altrock, W., Zhang, L., Volknandt, W.,
Zimmermann, H., Garner, C.C., Gundelfinger, E.D., and Dresbach, T. (2000).
Membrane association of presynaptic cytomatrix protein bassoon. Biochem Biophys
Res Commun *275*, 43–46.

Schoch, S., and Gundelfinger, E.D. (2006). Molecular organization of the presynaptic
active zone. Cell Tissue Res *326*, 379–391.

Shapira, M., Zhai, R.G., Dresbach, T., Bresler, T., Torres, V.I., Gundelfinger, E.D., Ziv,
N.E., and Garner, C.C. (2003). Unitary assembly of presynaptic active zones from
Piccolo-Bassoon transport vesicles. Neuron *38*, 237–252.

Südhof, T.C. (2012). The presynaptic active zone. Neuron *75*, 11–25.

Tao-Cheng, J.-H. (2007). Ultrastructural localization of active zone and synaptic
vesicle proteins in a preassembled multi-vesicle transport aggregate. Neuroscience
150, 575–584.

Tsuriel, S., Geva, R., Zamorano, P., Dresbach, T., Boeckers, T., Gundelfinger, E.D.,
Garner, C.C., and Ziv, N.E. (2006). Local sharing as a predominant determinant of
synaptic matrix molecular dynamics. PLoS Biol *4*, e271.

Tsuriel, S., Fisher, A., Wittenmayer, N., Dresbach, T., Garner, C.C., and Ziv, N.E.
(2009). Exchange and redistribution dynamics of the cytoskeleton of the active zone
molecule bassoon. J Neurosci *29*, 351–358.

Vukoja, A., Rey, U., Petzoldt, A.G., Ott, C., Vollweiter, D., Quentin, C., Puchkov, D.,
Reynolds, E., Lehmann, M., Hohensee, S., et al. (2018). Presynaptic Biogenesis
Requires Axonal Transport of Lysosome-Related Vesicles. Neuron *99*, 1216-1232.e7.

Waites, C.L., Leal-Ortiz, S.A., Okerlund, N., Dalke, H., Fejtova, A., Altrock, W.D.,
Gundelfinger, E.D., and Garner, C.C. (2013). Bassoon and Piccolo maintain synapse
integrity by regulating protein ubiquitination and degradation. EMBO J *32*, 954–969.

Wang, X., Kibschull, M., Laue, M.M., Lichte, B., Petrasch-Parwez, E., and Kilimann,
M.W. (1999). Aczonin, a 550-kD putative scaffolding protein of presynaptic active
zones, shares homology regions with Rim and Bassoon and binds profilin. J Cell Biol
147, 151–162.

Wildanger, D., Medda, R., Kastrup, L., and Hell, S.W. (2009a). A compact STED
microscope providing 3D nanoscale resolution. J Microsc *236*, 35–43.

Wildanger, D., Bückers, J., Westphal, V., Hell, S.W., and Kastrup, L. (2009b). A STED
microscope aligned by design. Opt Express *17*, 16100–16110.

Wittenmayer, N. (2014). Photoconversion of CFP to study neuronal tissue with electron
microscopy. Methods Mol Biol *1148*, 77–87.

Zhai, R.G., Vardinon-Friedman, H., Cases-Langhoff, C., Becker, B., Gundelfinger,
E.D., Ziv, N.E., and Garner, C.C. (2001). Assembling the presynaptic active zone: a
characterization of an active one precursor vesicle. Neuron *29*, 131–143.

1104
1105