Circadian modulation of human sleep spindle frequency

Róbert Bódizs[1,2], Csenge G. Horváth[1], Orsolya Szalárdy[1,3], Péter P. Ujma[1,2], Péter Simor[1,4,5], Ferenc Gombos[6,7], Ilona Kovács[6,7], Lisa Genzel[8], Martin Dresler[8]

[1]*Institute of Behavioural Sciences, Semmelweis University, Budapest, Hungary*

[2]*National Institute of Clinical Neurosciences, Budapest, Hungary*

[3]*Institute of Cognitive Neuroscience and Psychology, Research Centre for Natural Sciences, Budapest, Hungary*

[4]*Institute of Psychology, ELTE, Eötvös Loránd University, Budapest, Hungary*

[5]*UR2NF, Neuropsychology and Functional Neuroimaging Research Unit at CRCN – Center for Research in Cognition and Neurosciences and UNI – ULB Neurosciences Institute, Université Libre de Bruxelles (ULB), Brussels, Belgium*

[6]*Department of General Psychology, Pázmány Péter Catholic University, Budapest, Hungary*

[7]*MTA-PPKE Adolescent Development Research Group, Budapest, Hungary*

[8]*Donders Institute for Brain, Behaviour and Cognition, Radboud University, Nijmegen, The Netherlands*

Corresponding author:

Róbert Bódizs, PhD

Address: Institute of Behavioural Sciences, Semmelweis University, Nagyvárad tér 4., H-1089 Budapest, Hungary

E-mail: bodizs.robert@med.semmelweis-univ.hu

Phone: +361 210 2953

Fax: +361 210 2955

Conflict of interests: None declared.

Abstract

Homeostatic and circadian processes play a pivotal role in determining sleep structure, timing and quality. In sharp contrast with the wide accessibility of the EEG index of sleep homeostasis, an electrophysiological measure of the circadian modulation of sleep is still non-available. As a consequence, the circadian component of sleep regulation is assumed, but not assessed in routine polysomnographic records. Available evidence suggests that sleep spindle frequencies are markers of the circadian modulation of sleep. In order to test the above assumption, the Budapest-Munich database of sleep records (N = 251), as well as an afternoon nap sleep record database (N = 112) were analysed by the Individual Adjustment Method of sleep spindle analysis. Slow and fast sleep spindle frequencies were characterized by U-shaped overnight dynamics, with highest values in the first and the fourth-to-fifth sleep cycle and the lowest values in the middle of the sleeping period (cycles 2–3). Age-related attenuation of sleep spindle deceleration was evident. Estimated phases of the nadirs in sleep spindle frequencies were advanced in children as compared to other age groups. Additionally, nap sleep spindles were faster than night sleep spindles (0.57 and 0.39 Hz difference for slow and fast types, respectively). Our findings suggest that the circadian modulation of sleep is measurable by a fine frequency resolution analysis of sleep spindles. Moreover, age-related attenuation of circadian sleep modulation can be measured by assessing the overnight dynamics in sleep spindle frequency. Phase of the minimal sleep spindle frequency is a putative biomarker of chronotype.

Keywords: thalamocortical oscillations, hyperpolarization-rebound sequences, time of day-effects, NREM sleep, electroencephalogram, chronobiology

Introduction

Easily accessible and reliable biomarkers of sleep regulatory processes are of utmost importance in the objective measurement of sleep quality and rest-activity rhythms. The two-process model is one of the most influential and substantive theories of sleep regulation, proposing a linear interaction between sleep homeostasis (process S) and the circadian rhythm (process C) in humans (Borbély 1982). Although the basic tenets and the main predictions of the model are widely accepted and empirically supported (Borbély et al. 2016), investigators conducting common polysomnography studies are often challenged by the task of depicting and characterizing the two sleep regulatory processes. Thanks to widely available digital electroencephalogram (EEG) recording and analysis tools, sleep homeostasis is easily measurable by quantifying EEG slow wave activity (SWA: spectral power in the 0.75–4.5 Hz range) (Achermann 2009). In contrast, the circadian modulation of sleep EEG is either assumed and hypothetically included in the model, without considering the individual differences in phase (Achermann and Borbély 2003; Borbély 1982) or measured in particularly complex chronobiology study settings, like constant routine (Knoblauch et al. 2005) or forced desynchrony protocols (Wei et al. 1999), which are not easy to implement in common clinical and research sleep studies and require notoriously large time and labor investments. Another objective method which is instrumental in the assessment of the circadian modulation of sleep consists of long-term physiological measurements of core body temperature and/or specific endocrine factors, like melatonin or cortisol release (Oster et al. 2017; Reid 2019). Although these more direct ways of assessing the circadian component of sleep regulation seem promising, they are widely non-convenient and expensive, thus rarely included in the routine polysomnography examination protocols. Omitting these measurements leads to a permanent lack of information on the individual differences in chronotype, the latter being defined as the phase of entrainment of the circadian rhythm and the zeitgebers/local time (Roenneberg 2012). Standard polysomnography examinations disregard the fact that subjects with earlier or later chronotypes are characterized by different levels of circadian modulation of their sleep during the same period of recording. Known age effects of chronotype include a progressive phase delay from childhood till the end of adolescence, followed by a slow gradual phase advancement during aging (Roenneberg et al. 2004). As a consequence, a widely accessible EEG index of the circadian processes could significantly improve the insight into sleep regulation and complete our understanding of sleep and chronotype.

Given the intermingling of sleep homeostasis and circadian modulation in natural, night-time sleep, specific EEG-measures of the circadian process would be advantageous. Results of studies implementing forced desynchrony, constant routine, sleep displacement or overnight polysomnography protocols suggest that various aspects of sleep spindles reflect the circadian

modulation of sleep or time-of-day (Aeschbach, Dijk, and Borbély 1997; Knoblauch et al. 2005; Purcell et al. 2017; Wei et al. 1999). Sleep spindles are known as trains of distinct sinusoidal EEG waves with a frequency of 11–16 Hz (most frequently 12–14 Hz) lasting at least 0.5 s and emerging in non-rapid-eye-movement (NREM) sleep stages N2 and N3 (Berry et al. 2018). These oscillatory events were shown to arise from the hyperpolarization-rebound sequences of widely synchronized thalamocortical neurons. The thalamic reticular nucleus is hypothesized to be the main source of hyperpolarization, whereas the T-type Ca^{2+}-channels are sources of the rhythmic recurrence of firing (Fernandez and Lüthi 2020). Based on topography and dominant frequency, two types of sleep spindles are distinguishable. The anterior (frontal) slow spindles are known to consist of waves of roughly 12 Hz (<12.5 Hz), whereas the posterior (centroparietal) fast sleep spindles are oscillations with a typical frequency of 14 Hz (>12.5 Hz) (Gibbs and Gibbs 1951). Thorough analyses of sleep spindles indicate considerable individual differences (Bódizs et al. 2009; Cox et al. 2017), as well as age and sex effects. The latter two consists of unusually low frequency sleep spindles in prepubertal (and pre-schooler) ages (Ujma et al. 2016), as well as a slightly (0.5 Hz) increased sleep spindle frequency and variability in pubertal girls and adult women, as compared to boys and men (Bódizs et al. 2021; Ujma et al. 2014). Increased variability of sleep spindles in women is known to reflect the neural effects of hormonal variation during the menstrual cycles (Ishizuka et al. 1994). That is, the differentiation of slow and fast sleep spindles and the individual adjustment of sleep spindle frequencies is a basic requirement in conducting studies in the field.

Circadian modulation of sleep spindle frequency was evidenced in a forced desynchrony study, which is instrumental in differentiating sleep homeostatic and circadian effects by invoking the non-24-hours (usually 28 hours) days approach. The frequency of sleep spindles was lowest at the nadir of the core body temperature rhythm and peaked at the acrophase of the body temperature rhythm. In addition, circadian modulation of sleep spindle frequency, that is the decrease in oscillatory frequency during the biological night, was attenuated in aged subjects, as compared to young ones (Wei et al. 1999). These findings cohere with outcomes of nap studies performed in constant routine conditions. The nadir in sleep spindle frequency was shown to coincide with the acrophase of salivary melatonin levels in humans in this report (Knoblauch et al. 2005). In addition, experimental manipulation of sleep timing indicates prominent time of day-effects in sleep spindle frequency activities, suggesting a frequency-specific circadian modulation: lower bins (12.25–13 Hz) of the spindle frequency spectral power peak during the middle of the night sleep period (between 2:00 and 5:00 AM), when the highest bins (14.25–15 Hz) reach their nadir (Aeschbach et al. 1997). In addition, daytime recovery sleep after 25 hours of wakefulness was shown to be characterized by increased spindle frequency compared to baseline night sleep records (Rosinvil et al. 2015). This

latter effect was more pronounced in young as compared to middle-aged participants. That is sleep spindles are slower in the middle of the habitual sleep periods, characterized by high melatonin and low core body temperature levels. It has to be noted that brain temperature per se is a direct modulator of sleep spindle frequency: higher temperatures imply faster spindles (Csernai et al. 2019).

The above studies were not designed to differentiate between slow and fast sleep spindles, thus authors could not deduce if the reported post-midnight deceleration of sleep spindles reflects a frequency decrease or a change in the relative predominance of slow over fast sleep spindles (Aeschbach et al. 1997; Wei et al. 1999). Moreover, there is no direct evidence for the detectability of the above described frequency evolution of sleep spindles during habitual (non-displaced) sleep periods. A study examining the overnight evolution of sleep spindle frequencies, but without differentiating slow and fast sleep spindle events, revealed a sleep time-dependent increase in frequency (Himanen et al., 2002). This pattern evidently contrasts the convergent findings reported by studies implementing chronobiology protocols and systematic sleep displacement. The discrepancy might reflect a contamination of two different factors: change in sleep spindle frequencies and relative slow over fast sleep spindle incidence during the course of the night.

Our aim is to test the circadian modulation of sleep spindle frequencies in one-night-records of habitually timed sleep periods in different age groups by analysing its overnight dynamics (change over consecutive sleep cycles) and by comparing night-time sleep spindle frequency with afternoon nap sleep spindle frequency. We also aim to provide a differential analysis of slow and fast sleep spindles and depict the potential difference between oscillatory deceleration of slow and/or fast sleep spindles and alternatively, the change in the relative predominance of slow over fast sleep spindle events. In order to achieve these goals, we use an already established procedure of individualizing slow and fast sleep spindle frequencies with high frequency resolution (Bódizs et al. 2009). Given the already evidenced circadian modulation and the reported melatonin- and/or temperature-dependence of sleep spindle frequency, we hypothesize that:

1. Overnight dynamics in sleep spindle frequency is characterized by a U-shaped distribution (sleep spindles are slower in the middle of the habitual sleep period, as compared to the first and the last sleep cycles)
2. Middle night slowing of sleep spindle frequency is reduced in aged subjects as compared to young participants
3. Night sleep spindles are slower than nap sleep spindles

4. Estimated phase of the nadir in sleep spindle frequency is delayed in teenagers and young adults, as compared to children and middle aged adults.

Methods

Subjects and databases

Multiple, published databases are used in this study. The Munich-Budapest database of sleep records consists of 251 night-timed polysomnography registrations (Bódizs et al. 2017). Subjects' age varies between 4 and 69 years (122 females, mean age: 25.73 years). Participants of the night sleep record dataset slept at their habitual sleeping time in the laboratory (N = 208) or in their homes (recorded by ambulatory polysomnography, N = 43) on two consecutive nights. In order to attenuate the first-night effect (Agnew et al. 1966), only the second night data were used. Caffeine containing, but not alcoholic beverages were allowed in the morning hours in our adult subjects (maximum 2 cups of coffee/subjects before noon), whereas some of the participants who were light smokers (N = 8) were allowed to smoke during the day. The nap sleep records (N = 112) stem from studies on the effects of napping on memory consolidation (Genzel et al. 2012, 2014), as well as from a study analysing the relationship between nap sleep spindles and intelligence (Ujma, Bódizs, et al. 2015). Age of the napping subjects varied between 18 and 30 years (mean: 23.72 years). Women involved in the nap sleep studies (N = 30) were recorded twice: afternoon naps took place during the first and the third week of their menstrual cycle (early-follicular and mid-luteal phases, respectively with hormonal blood tests assuring this assumption). Data of the two nap records of these subjects were averaged before statistical analyses. All subjects were healthy and free of any medications, except contraceptives in some of the women in the reproductive age. Details of the recording procedures are reported in Table 1. All subjects (night sleepers and nap sleepers) were free of acute and chronic medical conditions, including the history of sleep disorders. The research protocols were approved by the Ethical Committee of the Semmelweis University (Budapest, Hungary) or the Medical Faculty of the Ludwig Maximilians University (Munich, Germany) in accordance with the Declaration of Helsinki. All subjects or the parents of the underage participants signed informed consent for the participation in the studies.

EEG processing

Records were scored according to standard criteria of sleep-wake states (Berry et al. 2018), followed by artefact-removal on a 4 seconds basis. Sleep cycle segmentation was based on reported criteria

(Aeschbach and Borbély 1993). EEG signals (mathematically-linked mastoid reference) of the non-artefactual N2 and N3 sleep periods were subjected to the Individual Adjustment Method (IAM) of sleep spindle analysis (Bódizs et al. 2009) in chunks based on successive sleep cycles of night-time records and as a whole in case of nap sleep. Only the frequency outputs of the IAM is reported in the present study.

The IAM consists of a high-resolution (1/16 Hz) frequency adjustment of characteristic slow- and fast sleep spindle frequencies, based on the zero-crossing points of the averaged second order derivatives of the NREM sleep EEG amplitude spectra (magnitude of the 4 s, Hanning tapered mixed-radix Fast Fourier Transforms). That is, amplitude spectra of NREM sleep EEG with 0.25 Hz frequency resolution were subjected to a numerical derivation procedure as follows: a second-degree polynomial curve fitting was performed using all sets of successive bin triplets (0.75 Hz), with an overlap of 2 bins (0.5 Hz) in the 9–16 Hz range. Derivatives were calculated in the middle of the triplets by using the rules of derivation for quadratic formulas, returning practically the slopes of the curves fitted to the triplets. This procedure is repeated on the series composed by the first derivatives in order to obtain the second derivatives of the amplitude spectra. Second derivatives were averaged over the EEG recording locations resulting in one averaged second derivative function per subject and sleep cycle (NREM period). The zero crossing points of these averaged second derivative were refined on the frequency scale of the high-resolution spectra (1/16 Hz) and considered as lower and upper frequency boundaries of slow and fast sleep spindles if

a. they compassed a negative peak (positive peaks are reverted and negative in the second order derivatives)
b. amplitude spectra at the corresponding frequencies were characterized by frontal (average of Fp1, Fp2, F3, F4, Fz, F7, and F8) and centro-parietal (average of C3, C4, Cz, P3, and P4) dominance, respectively (Fig. 1).

In cases in which more than two peaks were present, we based our selection on frontal and centro-parietal dominances of the respective frequencies, relying on the amplitude spectra. That is, the peak with most pronounced frontal and centro-parietal dominance in amplitude spectra, were considered as slow and fast sleep spindle frequencies, respectively. Lack of a definitive slow spindle peak was handled by searching the relative maxima of the frontal minus centro-parietal amplitude spectral values, the slow spindle spectral peaks being defined at the edges of the associated bulge of second-order derivatives (which latter approach but do not reach zero in these cases) (see Ujma et al. 2016). Fast sleep spindle frequencies were unambiguously detected in all subjects and sleep cycles, without relying on the above described heuristic. Last, but not least the spectral peaks of slow and fast sleep

spindles could partially overlap in some cases. In such cases the intersection of slow and fast sleep spindle-related spectral peaks was considered as a border frequency.

Slow and fast sleep spindle frequencies were defined as the middle of the respective frequency bands (arithmetic mean of lower and upper frequency boundaries).

The estimated phases of the nadirs of slow and fast sleep spindle frequencies was assessed as follows:

i.   finding the sleep cycle characterized by the individual minima in sleep spindle frequency

ii.   determining the local time of day of the middle of the respective sleep cycle (NREM-REM period) by recovering the start time of the respective sleep recording

iii.   transforming time of day values (HH:MM:SS) to hours and fractions of hours relative to midnight (that is times of day before midnight became negative, whereas those after midnight emerged as positive values; higher values indicate later phases of the nadir in sleep spindle frequency).

We could recover the start times of the original recording in N = 248 subjects (98.8% of the subjects). In order to provide a composite measure of the assumed circadian phase of sleep spindle frequency we calculated the individual mean of the phases of slow and fast sleep spindle frequency minima.

Fig. 1. Critical steps in determining the individual slow and fast sleep spindle frequencies. Amplitude spectra, as well as their frontal and centro-parietal averages (continuous blue and red lines, respectively) are calculated by means of mixed-radix Fast Fourier Transformation (4 sec, Hanning window, 2 sec overlap, see upper panel). The definition of frontal and centro-parietal regions is provided in the lower insert (circle representing the head model). Amplitude spectra are submitted to a numerical derivation procedure in two steps, resulting in second derivatives ('', lower panel). The latter are averaged (continuous green line) followed by the detection of their zero crosses (interpolated on a scale of 1/16 Hz, see dotted, light blue arrows). Last, the individual spectral peak frequencies are determined by combining the frontal and parietal spectral peaks with the zero crosses determined above. Note that the procedure is the first step of the Individual Adjustment Method (IAM) of sleep spindle analysis.

Statistical analyses

In coherence with our earlier report (Bódizs et al. 2017), subjects of the night sleep study were classified according to the following age ranges: children (age < 10 years; N = 31, 15 females), teenagers (10 years ≤ age < 20 years; N = 36, 18 females), young adults (20 years ≤ age < 40 years; N = 150, 75 females) and middle-aged adults (age ≥ 40 years; N = 34, 14 females). The sleep cycle effect in oscillatory spindle frequency was tested with the control of the between-subject factors sex and age group, the within-subject factor spindle type (slow/fast), as well as all possible interactions of the above mentioned predictors, by using the general linear model approach. The comparability of nap and night sleep records (N = 249, 98 females, mean age: 23.79) was ensured by matching a subset of the night sleep records (the young adult group, N = 141, 69 females) to the nap sleep database (N = 108, 29 females) in terms of age and there were no significant difference between the two groups (t(247) = -0.318, p = .75). Besides our main focus (the nap vs night between subject factor), sex (between-subject) and spindle type (slow/fast, within-subject) were included in the general linear model. Given the homogenous age of the subjects of the nap vs night sleep study, the age factor was not included in the statistical model.

Table 1. Details of the recording procedures of different subsamples

Nap/ night	Subsample	Original setting/aim	Nr. of subjects (females)	Age range (ys)	Available EEG derivations (10-20 system)	Recording apparatus	Precision (bit)	Hardware prefiltering (Hz)	Sampling rate (Hz/channel)	Recording software
Night sleep	MPIP[1] – I	Lab/sleep & IQ	95 (43)	18–69	Fp1, Fp2, Fpz, AF1, AF2, F3, F4, Fz, F7, F8, C3, C4, Cz, P3, P4, Pz, T3, T4, T5, T6, O1, O2	Comlab 32 Digiltal SleepLab	8	0.53–70	250	Brainlab 3.3
	MPIP[1] – II		20 (12)		Fp1, Fp2, F3, F4, C3, C4, P3, P4, O1, O2					
	PPCU[2] – I	Home/Williams syndrome study (controls included here)	20 (14)	6–28	Fp1, Fp2, Fpz, F3, F4, F7, F8, Fz, C3, C4, Cz, T3, T4, T5, T6, P3, P4, Pz, O1, O2, Oz	SD-LTM 32BS (Micromed Ltd, Italy)	22	0.15-250 (plus <463.3 Hz digital antialising filtering before downsampling from 4096 to 1024 Hz)	1024	BRAIN QUICK System PLUS (Micromed)
	PPCU[2] – II	Home/Adolescent sleep	23 (12)	15–22						
	SU[3] – I	Lab/sleep & IQ, sleep spindle methodology, wake-sleep transition analysis	49 (19)	17–55	Fp1, Fp2, F3, F4, F7, F8, Fz, C3, C4, Cz, T3, T4, T5, T6, P3, P4, O1, O2	Flat Style SLEEP La Mont Headbox, HBX32-SLP preamplifier (La Mont Medical)	12	0.5–70	249	Datalab (Medcare)
	SU[3] – II	Lab/nightmare study (controls included here)	16 (7)	19–21	Fp1, Fp2, F3, F4, Fz, F7, F8, C3, C4, Cz, P3, P4, Pz, T3, T4, T5, T6, O1, O2	Brain-Quick BQ132S (Micromed)	12	0.33–1500 (plus <450 Hz anti-aliasing digital filtering before downsampling from 4096 to 1024 Hz)	1024	System 98 (Micromed)
	SU[3] – III	Lab/home/ children's dreaming	29 (15)	3.84–8.42		Brain-Quick BQ132S/ SD LTM 32BS (Micromed)	12/22	0.33–1500/0.15-250 (plus <450/<463.3 Hz anti-aliasing digital filtering before downsampling from 4096 to 1024 Hz)		System 98/System Plus Evolution (Micromed)
Nap sleep	MPIP – III	Lab/Nap sleep & memory consolidation	55 (35)	18–30	C3, C4	Comlab 32 Digiltal SleepLab	8	0.53–70	250	Brainlab 3.3
	MPIP - IV	Lab/Nap sleep & IQ	79 (0)	18–30	F3, F4, C3, C4, O1, O2	Comlab 32 Digiltal SleepLab	8	0.53–70	250	Brainlab 3.3

[1]Max Planck Institute of Psychiatry, Munich, Germany; [2]Pázmány Péter Catholic University, Budapest Hungary; [3]Semmelweis University Budapest, Hungary

Results

Overnight dynamics in sleep spindle frequency

Number of complete sleep (NREM-REM) cycles in night sleep records varied between 2 and 5 (mean:

4.24). In order to keep the number of subjects high, first we will focus on the NREM periods of the

first four sleep cycles (~6.8 hours after sleep onset). Mean duration of sleep cycles was 102.22,

106.33, 103.97, 95.68, and 81.59 minutes for cycle 1, 2, 3, 4, and 5 respectively. No significant age,

sex, cycle main effects or interactions were revealed for sleep cycle durations (GLM for cycles 1–4,

repeated measures with the categorical predictors age group and sex).

Significant sleep cycle effects ($F = 18.57$; d.f. = 3, 693; $p < .001$) revealed a U-shaped dynamic in

spindle frequencies, with highest values in the first and the fourth cycles and lowest values in the

second and third cycles. This pattern was evident for both slow and fast sleep spindle types,

characterized by overall frequency drops of 0.09 and 0.13 Hz, respectively (Fig. 2; post-hoc Fisher LSD

tests indicate decelerated sleep spindles in cycles 2 and 3, as compared to cycles 1 and 4 in terms of

both slow and fast spindles, whereas cycles 1 and 4 do not differ significantly). The sleep cycle effect

interacted with the between-subject factor age group ($F = 2.07$; d.f. = 9, 693; $p = .03$), indicating an

attenuated spindle deceleration in middle aged adults (flattening of the U-shaped dynamic), but also

in the lack of this pattern in the time course of slow sleep spindles in girls below 10 years of age. The

age-related flattening of overnight dynamic is supported by the negative correlation between age

and a measure of deceleration of spindles (mean spindle frequency in cycles 1 and 4 minus mean

spindle frequency in cycles 2–3): $r = -.12$ ($p = .049$) for slow and $r = -.21$ ($p = .001$) for fast types.

Nominal sleep spindle deceleration reached 0.18 and 0.20 in case of slow and fast sleep spindles of

teenagers, respectively, whereas it was only 0.03 and 0.04 in middle aged adults.

As expected, sleep spindle frequency varied between age groups ($F = 66.46$; d.f. = 3, 693; $p < .001$)

and as a function of sex ($F = 8.94$; d.f. = 1, 693; $p = .003$), indicating increasing oscillatory frequencies

with increasing ages, and higher frequencies in females as compared to males, respectively.

A subset of records contained a complete 5[th] sleep cycle (N = 75, that is 29 children, 23 teenagers, 22

young adults, and 1 middle aged adult). The above statistical analyses were extended over the 5[th]

sleep cycle by relying on this subset of records, with the evident exclusion of the single middle aged

subject (N = 74). Results still reflected a significant sleep cycle effect on spindle frequency ($F = 17.08$;

d.f. = 4, 272; $p < .001$), but not interaction between cycle effect and age group (because the middle

aged group of subjects with flattened U-shaped dynamic was not included in this latter analysis: $F =$

0.50; d.f. = 8, 272; $p = .85$). The reason of implementing the analysis on this subset of records was the

Fig. 2. Sleep spindle oscillations are decelerated in the middle of the habitual sleep period. The highest oscillatory frequencies are seen in the first and in the fourth sleep cycles, whereas cycles 2 and 3 are characterized by lower values. A. Slow and fast sleep spindle frequencies as functions of sleep cycle, age group and sex. Note the U-shaped distribution in the overnight dynamics, as well as its attenuation in middle aged subjects. In addition, a lack of this effect is also evident for slow sleep spindles in girls below 10 years of age. B. Example for the deceleration of sleep spindles in the middle of the habitual sleep period in a 20 years old male. Slow sleep spindles measured at recording location Fp2 are averaged to the maxima of the spindle-filtered signal. Averaged sleep spindles of successive sleep cycles are aligned according to their maxima. Note the progressive misalignment of averaged sleep spindle waves of sleep cycles 2 and 3, indicating their decelerated frequency.

focus on cycle 5. sleep spindle frequencies, which proved to exceed all other sleep spindle

frequencies measured in cycles 1–4, according to post-hoc Fisher LSD tests.

The phase of the nadir in sleep spindle frequency (NSSF)

The sample mean of the phase of the NSSF was 2.63 (SD = 2.16) and 2.49 (SD = 2.01) hours for slow

and fast sleep spindles, respectively. That is most of the participants reach their lowest sleep spindle

frequencies roughly two and a half hours after midnight (the sample mean of the individually

averaged phases of slow and fast NSSF was 2.56 hours). A general linear model targeting the effects

of age group, sex, spindle type and their interaction on NSSF resulted in a significant main effect of

age group (F = 5.30; d.f. = 3, 240; p = .001). Based on the means and standard errors depicted in

figure 3, it is evident, that the hypothesized age effects are best approximated by slow sleep spindle

frequencies: phase of the NSSF is seen circa one hour after midnight in children, whereas it reaches 3

hours in teenagers, followed by a slow decrease to 2.5 hours in upcoming age groups. Post-hoc

Fig. 3. Phase of the nadir in sleep spindle frequency (NSSF) as expressed in fraction of hours relative to
midnight. Vertical bars denote 95% confidence intervals.

Fisher LSD tests revealed a significantly earlier phase of the NSSF (slow spindles) in children as
compared to all other age groups. Nominal differences among teenagers and adults (young and
middle aged) did not result in significant post-hoc effects.

Nap vs night sleep spindle frequency

Nap sleep spindles were faster than night sleep spindles (F = 72.2; d.f. = 1, 249; p < .001). Although

this effect is evident for both spindle types (Fig. 4), day vs night slow spindle frequency differences

are higher than corresponding fast spindle differences (0.57 vs 0.39 Hz; interaction of nap vs night

and spindle type: F = 4.6; d.f. = 1, 249; p = .033). Oscillatory frequency of sleep spindles was higher in

females as compared to males, but this sex-difference did not interact with the nap vs night effect,

which is the focus of the current study.

Fig. 4. Nap sleep spindles are higher than night sleep spindles. Individual-specific slow- and fast sleep
spindles are compared in two groups of age- and sex-matched subjects sleeping during the afternoon
and during the night, respectively.

Discussion

Our findings indicate that sleep spindles decelerate in the middle of the habitual sleeping period, by

roughly 0.1 Hz. This effect is evident for both slow and fast sleep spindles, thus it is not caused by the

redistribution of slow and fast sleep spindle occurrences during the night. Younger subjects are

characterized by more accentuated deceleration of sleep spindling in the middle of their sleeping

period. This U-shaped dynamics, characterized by an equality of the first and the fourth sleep cycle,

and the decrease in the second and the third cycle cannot be explained by homeostatic sleep

regulatory mechanisms, which latter would imply a strictly decreasing or increasing function.

Findings reported by the relevant forced desynchrony (Wei et al. 1999), constant routine (Knoblauch

et al. 2005) and sleep displacement (Aeschbach et al. 1997) studies suggest the involvement of

circadian regulation in modulating sleep spindle frequency in humans. Former studies analysing the

overnight dynamics of sleep spindle frequency in habitually-timed polysomnography records, did not

deliberately discern slow and fast sleep spindles (Himanen et al. 2002), thus could at least partially

reflect the changing dominance of slow over fast sleep spindles. The latter was reported in our

former study, suggesting that both the incidence and the amplitude of slow spindles decreased over

successive sleep cycles, whereas an opposite trend was observable for fast sleep spindles (Bódizs et

al. 2009). Overnight changes in slow and fast sleep spindle incidence might explain the roughly 2 Hz

acceleration of sleep spindles reported by Himanen et al. (2002).

The deceleration of slow and fast sleep spindles we report in our current study could be a result of

the increased melatonin levels and the associated decrease in core body temperature in the middle

of the habitual sleep period. The acrophase of the human plasma melatonin rhythm is known to

emerge between 2:00 and 5:00 AM (Voultsios, Kennaway, and Dawson 1997). In addition, melatonin

receptors are expressed in the reticular thalamic nucleus (Ng et al. 2017), that is a critical

neuroanatomical structure in the process of sleep spindle generation (Fan, Liao, and Wang 2017).

Thus, both the timing and one of the target organs of melatonin release are ideally suited to support

the involvement of the pineal hormone in shaping the evolution of sleep spindle frequency during

the night. Last, but not least, sleep spindle frequency varies as a function of locally manipulated brain

temperature (Csernai et al. 2019), whereas melatonin is known for its hypothermic effect (Marrin et

al. 2013). As a consequence, the hypothermia induced by increased melatonin levels in the middle of

the sleeping period is potentially involved in the deceleration process of sleep spindling revealed in

the current report. This assumption is further supported by differences in sleep spindle frequency

activity during and outside melatonin secretory phase (Knoblauch et al. 2005), as well as by the

findings indicating that spindle frequency reaches its nadir at the trough of the body temperature

cycle (Wei et al. 1999).

The age-dependent attenuation of sleep spindle deceleration in the middle of the habitual sleep

period reported herein is reminiscent of the age-related decline in the amplitude of the circadian

rhythm (Hood and Amir 2017), the flattening of the body temperature rhythm in the aged (Weitzman

et al. 1982), as well as the associated decline in melatonin release (Waldhauser et al. 1988).

Decreased circadian modulation and melatonin production might fail to induce an efficient reduction

in core body temperature and consequently a suboptimal modulation of sleep spindle frequency.

Our findings cohere with the forced desynchrony study reporting that older subjects express notably

smaller circadian variation of sleep spindle frequency than young subjects (Wei et al. 1999).

However, the reason for a lack of the above detailed U-shaped distribution in the frequency of slow

sleep spindles in young girls do not cohere with the above reasoning.

Sleep spindles speed up in the fourth sleep cycle and continue to accelerate on the rising limb of the

circadian cycle up to the 5th sleep cycle in the subset of subjects having five complete NREM-REM

periods in our dataset. Thus, we assume that sleep spindles are even faster during daytime as

compared to night time sleep. This coheres with our findings reported in the nap study. That is, in

addition to the reported deceleration of spindle oscillations in the middle of the habitual night sleep

period, we report accelerated spindle frequency during afternoon naps as compared to nocturnal

averages (cycles 1–4). The finding that the oscillatory frequency of nap sleep spindles exceeds that of

night spindles further strengthens our assumption that the above discussed sleep cycle effects might

reflect circadian modulation and/or melatonin release. Core body temperature is known to be higher

during afternoon hours as compared to late night periods (Baehr, Revelle, and Eastman 2000),

whereas melatonin release is evidently at its lowest level during these daytime hours (Aulinas 2000).

Thus, the acceleration of sleep spindles in afternoon naps coheres with our assumption regarding the

circadian regulation of the duration of thalamocortical hyperpolarization-rebound sequences. To the

best of our knowledge this is the first study explicitly reporting a nap vs night sleep difference in

spindle frequencies.

We hypothesized that the phase of the middle night drop in sleep spindle frequency could serve as

an EEG-marker of the circadian phase in humans. Available data in the current set is only suitable for

an indirect test of this assumption. That is, we tested the known age-effects in circadian phase by

comparing the different age groups. Our findings partially support the hypothesis: children were

characterized by earlier phases than teenagers or adults. Although, this latter finding coheres with

known age-differences in circadian phase, additional age effects, namely the phase advancement in

middle aged adults was not unequivocally supported by our findings. This could reflect the lack of

aged subjects in our sample or the insufficient precision of measuring the phase of the nadir of sleep

spindle frequency (in the middle of the sleep cycle characterized by lowest sleep spindle frequency).

Later studies could invoke a more instantaneous phase measure, providing investigators with

increased temporal resolution. Given the partial statistical support, we consider our measure as a

potentially suitable EEG-index of circadian phase, which might bridge the methodological and

conceptual gap between chronobiology and somnology in future investigations.

Besides of the U-shaped overnight dynamics and afternoon nap effect, formerly reported sex-differences and age-effects in sleep spindle frequencies were also supported by our current analyses. That is, sleep spindles were of higher frequency in females as compared to males, an effect which was already reported by different research groups (Ujma et al., 2014; Markovic, Kaess, and Tarokh 2020). Moreover, sleep spindle frequencies were significantly lower in prepubertal ages (below 10 years), as compared to teenagers and adults. This effect was also reported by our own former study (Ujma et al., 2016), as well as by other teams analysing sleep spindles in children (Campbell and Feinberg 2016). The convergence of these findings with the available reports in the literature strengthens the validity of our current approach.

The strength of our study is the high number of subjects and wide age range, whereas limitations are the lack of repeated day vs night sleep measurements in the same groups of subjects, as well as the lack of valid circadian measures (body temperature, melatonin release) against which we could perform tests of convergent validity. We consider our findings as a first step in defining an EEG-index of the circadian modulation of sleep, which could efficiently complete the already available and widely used measure of sleep homeostasis. We hypothesize that the reliable measurement of the overnight dynamics of slow and fast sleep spindle frequencies might convey information about the amplitude and perhaps the phase of the circadian rhythm in future translational and clinical studies, including investigations on patients suffering from circadian rhythm sleep disorders.

Acknowledgements

Research supported by the Hungarian National Research, Development and Innovation Office (K-128117; https://nkfih.gov.hu/about-the-office), the Higher Education Institutional Excellence Program of the Ministry of Human Capacities in Hungary, within the framework of the Neurology thematic program of the Semmelweis University (http://semmelweis.hu/english/), the Netherlands Organization for Scientific Research (NWO; https://www.nwo.nl/en), the European Cooperation in Science and Technology (COST Action CA18106; https://www.cost.eu/), as well as the general budgets of the Institute of Behavioural Sciences, Semmelweis University (http://semmelweis.hu/magtud/en/) and the Max Planck Institute of Psychiatry (https://www.psych.mpg.de/en). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. Authors wish to thank the assistance of Dorina Ali and Dávid Rottmayer for their assistance in data analysis.

References

Achermann, Peter. 2004. "The Two-Process Model of Sleep Regulation Revisited." in *Aviation Space*
and Environmental Medicine. Vol. 75.

Achermann, Peter. 2009. "EEG Analysis Applied to Sleep." *Epileptologie* 26:28–33.

Achermann, Peter, and Alexander A. Borbély. 2003. "Mathematical Models of Sleep Regulation."
Frontiers in Bioscience 8(SUPPL.):683–93.

Aeschbach, Daniel, and Alexander A. Borbély. 1993. "All-night Dynamics of the Human Sleep EEG."
Journal of Sleep Research 2(2):70–81. doi: 10.1111/j.1365-2869.1993.tb00065.x.

Aeschbach, Daniel, Derk Jan Dijk, and Alexander A. Borbély. 1997. "Dynamics of EEG Spindle
Frequency Activity during Extended Sleep in Humans: Relationship to Slow-Wave Activity and
Time of Day." *Brain Research* 748(1–2):131–36. doi: 10.1016/S0006-8993(96)01275-9.

Agnew, H. W., Wilse B. Webb, Robert L. Williams, and J. Hillis Miller. 1966. "The First Night Effect: An
EEG Study of Sleep." *Psychophysiology* 2(3):263–66.

Aulinas, Anna. 2000. *Physiology of the Pineal Gland and Melatonin (Updated 2019)*. edited by K.
Feingold, B. Anawalt, A. Boyce, G. Chrousos, W. de Herder, K. Dhatariya, K. Dungan, A.
Grossman, J. Hershman, J. Hofland, S. Kalra, G. Kaltsas, C. Koch, P. Kopp, M. Korbonits, C.
Kovacs, W. Kouhung, B. Laferrére, E. McGee, R. McLachlan, J. Morley, M. New, J. Purnell, R.
Sahay, F. Singer, C. Stratakis, D. Trence, and D. Wilson. South Darthmouth (MA).

Baehr, Erin K., William Revelle, and Charmane I. Eastman. 2000. "Individual Differences in the Phase
and Amplitude of the Human Circadian Temperature Rhythm: With an Emphasis on
Morningness-Eveningness." *Journal of Sleep Research* 9(2):117–27. doi: 10.1046/j.1365-
2869.2000.00196.x.

Berry, RB, CL Albertario, SM Harding, and et al. 2018. *For the American Academy Os Sleep Medicine.*
The AASM Manual for the Scoring of Sleep and Associated Events: Rules, Terminology and
Technical Specifications. Version 2. Darien, IL: American Academy of Sleep Medicine.

Bódizs, Róbert, Ferenc Gombos, Péter P. Ujma, Sára Szakadát, Piroska Sándor, Péter Simor, Adrián
Pótári, Boris️ Nikolai Konrad, Lisa Genzel, Axel Steiger, Martin Dresler, and Ilona Kovács. 2017.
"The Hemispheric Lateralization of Sleep Spindles in Humans." *Sleep Spindles & Cortical Up*
States 1(1):42–54. doi: 10.1556/2053.01.2017.002.

Bódizs, Róbert, János Körmendi, Péter Rigó, and Alpár Sándor Lázár. 2009. "The Individual

Adjustment Method of Sleep Spindle Analysis: Methodological Improvements and Roots in the Fingerprint Paradigm." *Journal of Neuroscience Methods* 178(1):205–13. doi: 10.1016/j.jneumeth.2008.11.006.

Bódizs, Róbert, Orsolya Szalárdy, Csenge Horváth, Péter P. Ujma, Ferenc Gombos, Péter Simor, Adrián Pótári, Marcel Zeising, Axel Steiger, and Martin Dresler. 2021. "A Set of Composite, Non-Redundant EEG Measures of NREM Sleep Based on the Power Law Scaling of the Fourier Spectrum." *Scientific Reports* 11(1):2041. doi: 10.1038/s41598-021-81230-7.

Borbély, A. A. 1982. "A Two Process Model of Sleep Regulation." *Human Neurobiology* 1(3):195–204.

Borbély, Alexander A., Peter Achermann, Lorenz Trachsel, and Irene Tobler. 1989. "Sleep Initiation and Initial Sleep Intensity: Interactions of Homeostatic and Circadian Mechanisms." *Journal of Biological Rhythms* 4(2). doi: 10.1177/074873048900400205.

Borbély, Alexander A., Serge Daan, Anna Wirz-Justice, and Tom Deboer. 2016. "The Two-Process Model of Sleep Regulation: A Reappraisal." *Journal of Sleep Research* 25(2):131–43. doi: 10.1111/jsr.12371.

Cajochen, Christian, K. Kräuchi, and A. Wirz-Justice. 2003. "Role of Melatonin in the Regulation of Human Circadian Rhythms and Sleep." *Journal of Neuroendocrinology* 15(4).

Campbell, Ian G., and Irwin Feinberg. 2016. "Maturational Patterns of Sigma Frequency Power across Childhood and Adolescence: A Longitudinal Study." *Sleep* 39(1):193–201. doi: 10.5665/sleep.5346.

Cox, Roy, Anna C. Schapiro, Dara S. Manoach, and Robert Stickgold. 2017. "Individual Differences in Frequency and Topography of Slow and Fast Sleep Spindles." *Frontiers in Human Neuroscience* 11. doi: 10.3389/fnhum.2017.00433.

Csernai, Márton, Sándor Borbély, Kinga Kocsis, Dávid Burka, Zoltán Fekete, Veronika Balogh, Szabolcs Káli, Zsuzsa Emri, and Péter Barthó. 2019. "Dynamics of Sleep Oscillations Is Coupled to Brain Temperature on Multiple Scales." *Journal of Physiology* 597(15):4069–86. doi: 10.1113/JP277664.

Dijk, Derk Jan, and Charles A. Czeisler. 1995. "Contribution of the Circadian Pacemaker and the Sleep Homeostat to Sleep Propensity, Sleep Structure, Electroencephalographic Slow Waves, and Sleep Spindle Activity in Humans." *Journal of Neuroscience* 15(5 I). doi: 10.1523/jneurosci.15-05-03526.1995.

Driver, Helen S., Derk Jan Dijk, Esther Werth, Kurt Biedermann, and Alexander A. Borbély. 1996.

"Sleep and the Sleep Electroencephalogram across the Menstrual Cycle in Young Healthy

Women." *Journal of Clinical Endocrinology and Metabolism* 81(2). doi: 10.1210/jc.81.2.728.

Fan, Denggui, Fucheng Liao, and Qingyun Wang. 2017. "The Pacemaker Role of Thalamic Reticular

Nucleus in Controlling Spike-Wave Discharges and Spindles." *Chaos* 27(7). doi:

10.1063/1.4991869.

Fernandez, Laura M. J., and Anita Lüthi. 2020. "Sleep Spindles: Mechanisms and Functions."

Physiological Reviews 100(2):805–68. doi: 10.1152/physrev.00042.2018.

Genzel, Lisa, Anna Bäurle, Alina Potyka, Renate Wehrle, Marek Adamczyk, Elisabeth Friess, Axel

Steiger, and Martin Dresler. 2014. "Diminished Nap Effects on Memory Consolidation Are Seen

under Oral Contraceptive Use." *Neuropsychobiology* 70(4):253–61. doi: 10.1159/000369022.

Genzel, Lisa, Teresa Kiefer, Lisa Renner, Renate Wehrle, Michael Kluge, Michael Grözinger, Axel

Steiger, and Martin Dresler. 2012. "Sex and Modulatory Menstrual Cycle Effects on Sleep

Related Memory Consolidation." *Psychoneuroendocrinology* 37(7):987–98. doi:

10.1016/j.psyneuen.2011.11.006.

Gibbs, F. A., and E. L. Gibbs. 1951. *Atlas of Electroencephalography*. Vol. 1. Cambridge: Addison-

Wesley.

Himanen, Sari Leena, Jussi Virkkala, Heini Huhtala, and Joel Hasan. 2002. "Spindle Frequencies in

Sleep EEG Show U-Shape within First Four NREM Sleep Episodes." *Journal of Sleep Research*

11(1):35–42. doi: 10.1046/j.1365-2869.2002.00273.x.

Hood, Suzanne, and Shimon Amir. 2017. "The Aging Clock: Circadian Rhythms and Later Life." *Journal

of Clinical Investigation* 127(2):437–46.

Iguchi, Haruo, Ken Ichi Kato, and Hiroshi Ibayashi. 1982. "Age-Dependent Reduction in Serum

Melatonin Concentrations in Healthy Human Subjects." *Journal of Clinical Endocrinology and

Metabolism* 55(1). doi: 10.1210/jcem-55-1-27.

Ishizuka, Yoshikazu, Charles P. Pollak, Shuichiro Shirakawa, Tatsuyuki Kakuma, Kazuo Azumi, Akira

Usui, Kouichi Shiraishi, Hitoshi Fukuzawa, and Tetsuhiko Kariya. 1994. "Sleep Spindle Frequency

Changes during the Menstrual Cycle." *Journal of Sleep Research* 3(1):26–29. doi:

10.1111/j.1365-2869.1994.tb00100.x.

Jobert, Marc, Eric Poiseau, Peter Jähnig, Hartmut Schulz, and Stanislaw Kubicki. 1992. "Topographical

Analysis of Sleep Spindle Activity." *Neuropsychobiology* 26(4). doi: 10.1159/000118923.

Knoblauch, Vera, Mirjam Münch, Katharina Blatter, Wim L. J. Martens, Carmen Schröder, Corina
Schnitzler, Anna Wirz-Justice, and Christian Cajochen. 2005. "Age-Related Changes in the
Circadian Modulation of Sleep-Spindle Frequency during Nap Sleep." *Sleep* 28(9):1093–1101.
doi: 10.1093/sleep/28.9.1093.

Loomis, Alfred L., E. Newton Harvey, and Garret Hobart. 1935. "Further Observations on the
Potential Rhythms of the Cerebral Cortex during Sleep." *Science* 82(2122).

Markovic, Andjela, Michael Kaess, and Leila Tarokh. 2020. "Gender Differences in Adolescent Sleep
Neurophysiology: A High-Density Sleep EEG Study." *Scientific Reports* 10(1):undefined. doi:
10.1038/s41598-020-72802-0.

Marrin, K., B. Drust, W. Gregson, and G. Atkinson. 2013. "A Meta-Analytic Approach to Quantify the
Dose-Response Relationship between Melatonin and Core Temperature." *European Journal of
Applied Physiology* 113(9):2323–29. doi: 10.1007/s00421-013-2668-x.

Ng, Khuen Yen, Mun Kit Leong, Huazheng Liang, and George Paxinos. 2017. "Melatonin Receptors:
Distribution in Mammalian Brain and Their Respective Putative Functions." *Brain Structure and
Function* 222(7):2921–39.

Oster, Henrik, Etienne Challet, Volker Ott, Emanuela Arvat, E. Ronald de Kloet, Derk Jan Dijk, Stafford
Lightman, Alexandros Vgontzas, and Eve Van Cauter. 2017. "The Functional and Clinical
Significance of the 24-Hour Rhythm of Circulating Glucocorticoids." *Endocrine Reviews* 38(1):3–
45.

Purcell, S. M., D. S. Manoach, C. Demanuele, B. E. Cade, S. Mariani, R. Cox, G. Panagiotaropoulou, R.
Saxena, J. Q. Pan, J. W. Smoller, S. Redline, and R. Stickgold. 2017. "Characterizing Sleep
Spindles in 11,630 Individuals from the National Sleep Research Resource." *Nature
Communications* 8. doi: 10.1038/ncomms15930.

Reid, Kathryn J. 2019. "Assessment of Circadian Rhythms." *Neurologic Clinics* 37(3):505–26.

Roenneberg, Till. 2012. "What Is Chronotype?" *Sleep and Biological Rhythms* 10(2):75–76.

Roenneberg, Till, Tim Kuehnle, Peter P. Pramstaller, Jan Ricken, Miriam Havel, Angelika Guth, and
Martha Merrow. 2004. "A Marker for the End of Adolescence." *Current Biology* 14(24).

Rosinvil, T., M. Lafortune, Z. Sekerovic, M. Bouchard, J. Dubé, A. Latulipe-Loiselle, N. Martin, J. M.
Lina, and J. Carrier. 2015. "Age-Related Changes in Sleep Spindles Characteristics during
Daytime Recovery Following a 25-Hour Sleep Deprivation." *Frontiers in Human Neuroscience*
9(JUNE). doi: 10.3389/fnhum.2015.00323.

Ujma, Péter P., Róbert Bódizs, Ferenc Gombos, Johannes Stintzing, Boris N. Konrad, Lisa Genzel, Axel
Steiger, and Martin Dresler. 2015. "Nap Sleep Spindle Correlates of Intelligence." *Scientific*
Reports 5. doi: 10.1038/srep17159.

Ujma, Péter P., Ferenc Gombos, Lisa Genzel, Boris Nikolai Konrad, Péter Simor, Axel Steiger, Martin
Dresler, and Róbert Bódizs. 2015. "A Comparison of Two Sleep Spindle Detection Methods
Based on All Night Averages: Individually Adjusted vs. Fixed Frequencies." *Frontiers in Human*
Neuroscience 9(FEB). doi: 10.3389/fnhum.2015.00052.

Ujma, Péter P., Boris Nikolai Konrad, Lisa Genzel, Annabell Bleifuss, Péter Simor, Adrián Pótári, János
Körmendi, Ferenc Gombos, Axel Steiger, Róbert Bódizs, and Martin Dresler. 2014. "Sleep
Spindles and Intelligence: Evidence for a Sexual Dimorphism." *Journal of Neuroscience*
34(49):16358–68. doi: 10.1523/JNEUROSCI.1857-14.2014.

Ujma, Péter P., Piroska Sándor, Sára Szakadát, Ferenc Gombos, and Róbert Bódizs. 2016. "Sleep
Spindles and Intelligence in Early Childhood-Developmental and Trait-Dependent Aspects."
Developmental Psychology 52(12):2118–29. doi: 10.1037/dev0000233.

Voultsios, Athena, David J. Kennaway, and Drew Dawson. 1997. "Salivary Melatonin as a Circadian
Phase Marker: Validation and Comparison to Plasma Melatonin." *Journal of Biological Rhythms*
12(5):457–66. doi: 10.1177/074873049701200507.

Waldhauser, F., G. Weiszenbacher, E. Tatzer, B. Gisinger, M. Waldhauser, M. Schemper, and H.
Frisch. 1988. "Alterations in Nocturnal Serum Melatonin Levels in Humans with Growth and
Aging." *Journal of Clinical Endocrinology and Metabolism* 66(3):648–52. doi: 10.1210/jcem-66-
3-648.

Wei, Henry G., Eymard Riel, Charles A. Czeisler, and Derk Jan Dijk. 1999. "Attenuated Amplitude of
Circadian and Sleep-Dependent Modulation of Electroencephalographic Sleep Spindle
Characteristics in Elderly Human Subjects." *Neuroscience Letters* 260(1):29--32. doi:
10.1016/S0304-3940(98)00851-9.

Weitzman, Elliot D., Margaret L. Moline, Charles A. Czeisler, and Janet C. Zimmerman. 1982.
"Chronobiology of Aging: Temperature, Sleep-Wake Rhythms and Entrainment." *Neurobiology*
of Aging 3(4):299–309. doi: 10.1016/0197-4580(82)90018-5.

Wirz-Justice, Anna. 2007. "How to Measure Circadian Rhythms in Humans." *Medicographia* 29(1):84–
90.