A centronuclear myopathy-causing mutation in dynamin-2 perturbs the actin-dependent structure of dendritic spines leading to excitatory synaptic defects in a murine model of the disease.

Running title: Actin-dependent synaptic defects in dynamin-2-linked CNM

Jorge Arriagada-Diaz[1], Bárbara Gómez[2], Lorena Prado-Vega[1], Michelle Mattar-Araos[3], Marjorie Labraña-Allende[3], Fernando Hinostroza[4], Ivana Gajardo[5], María José Guerra-Fernández[6], Jorge A. Bevilacqua[7], Ana M. Cárdenas[6], Marc Bitoun[8], Alvaro O. Ardiles[6,9,10]*, Arlek M. Gonzalez-Jamett[6,11]*.

[1]Programa de Magister en Ciencias, Mención Neurociencia, Universidad de Valparaíso, Valparaíso, Chile.
[2] Programa de Magister en Ciencias Médicas, Mención Biología Celular y Molecular, Universidad de Valparaíso, Valparaíso, Chile.
[3] Escuela de Tecnología Médica, Facultad de Medicina, Universidad de Valparaíso, Valparaíso, Chile.
[4] Centro de Investigación de Estudios Avanzados del Maule, CIEAM, Vicerrectoría de Investigación y Postgrado, Universidad Católica del Maule, Talca, Chile
[5] Departamento de Biología Celular y Molecular, Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile, Santiago, Chile.
[6]Centro Interdisciplinario de Neurociencia de Valparaíso, Universidad de Valparaíso, Valparaíso, Chile.
[7] Departamento de Neurología y Neurocirugía, Hospital Clínico Universidad de Chile and Departamento de Anatomía y Medicina Legal, Facultad de Medicina, Universidad de Chile, Santiago, Chile.
[8] Sorbonne Université, Inserm, Institut de Myologie, Centre de Recherche en Myologie, F-75013 Paris, France.
[9]Centro de Neurología Traslacional, Facultad de Medicina, Universidad de Valparaíso, Valparaíso, Chile.
[10]Centro Interdisciplinario de Estudios en Salud, Facultad de Medicina, Universidad de Valparaíso, Viña del Mar, Chile
[11] Escuela de Química y Farmacia, Facultad de Química y Farmacia, Universidad de Valparaíso, Valparaíso, Chile.

*correspondence should be addressed to Dr. Arlek M. Gonzalez-Jamett +56-32-2508203 arlek.gonzalez@uv.cl or Dr. Alvaro O. Ardiles +56-32-2507354 alvaro.ardiles@uv.cl

Abstract

Dynamin-2 is a large GTP-ase, member of the dynamin superfamily, that regulates membrane remodeling and cytoskeleton dynamics. In the mammalian nervous system dynamin-2 modulates synaptic vesicle (SV)-recycling at the nerve terminals and receptor-trafficking to and from postsynaptic densities (PSDs). Mutations in dynamin-2 cause autosomal dominant centronuclear myopathy (CNM), a congenital neuromuscular disorder characterized by progressive weakness and atrophy of distal skeletal muscles. Cognitive defects have also been reported in dynamin-2-linked CNM patients suggesting a concomitant impairment of the central nervous system. Here we addressed the mechanisms that lead to cognitive defects in dynamin-2-linked CNM using a knock-in mouse model that harbors the p.R465W mutation in dynamin-2, the most common causing CNM. Our results show that these mice exhibit reduced capability to learn and acquire spatial and recognition memory, impaired long-term potentiation of the excitatory synaptic strength and perturbed dendritic spine morphology, which seem to be associated with actin defects. Together, these data reveal for the first time that structural and functional synaptic defects underlie cognitive defects in the CNM context. In addition our results contribute to the still scarce knowledge about the importance of dynamin-2 at central synapses.

Introduction

Dynamin super-family (DSF) is a group of large GTP-ases that act as mechano-enzymes promoting membrane remodeling in several processes including exocytosis, endocytosis, intracellular trafficking and mitochondrial dynamics among others (Ferguson, et al., 2012; González-Jamett, et al., 2013; Lomash, et al., 2015; Arriagada-Diaz, et al., 2020). Classical dynamins are the best understood members of the DSF. In mammals, these are encoded by three different genes: *DNM1*, *DNM2* and *DNM3* located on chromosomes 9, 19 and 1, respectively (Newman-Smith, et al., 1997; Züchner, et al., 2005; Noakes, et al., 1999). Classical dynamins are composed by five highly conserved domains: an amino-terminal GTP-ase domain that binds and hydrolyze GTP, a middle structural domain, a PH-domain involved in lipid membrane interaction, a GTP-ase effector domain (GED) and an arginine- and proline-rich domain (PRD) that allows dynamin association with SH3-containing proteins (Praefcke, et al., 2004; Ferguson, et al., 2012; Antonny, et al., 2016; Singh, et al., 2017; Arriagada-Diaz, et al., 2020). These domains organize in three regions: a "bundle signaling element" (BSE) composed by helices from the GTP-ase domain and GED, a "stalk" composed by helices from the middle domain and GED, and a membrane-inserting "foot" formed by the PH domain (Chappie, et al., 2009; Faelber, et al., 2011; Ford, et al., 2011; Kong, et al., 2018). In response to the binding of GTP, the BSE convey conformational changes from the GTP-ase domain to the "stalk" promoting dynamin oligomerization in helical structures that enhance its GTP-ase activity and favor membrane scission (Hinshaw, et al., 1995; Warnock, et al., 1997; Antonny, et al., 2016). Dynamin assembly/disassembly cycles and GTP-ase activity also regulate the bundling of the actin cytoskeleton (Gu, et al., 2010; Zhang, et al., 2020; Lin, et al., 2020; Schiffer, et al., 2015a).

The three classical dynamins are expressed at the central nervous system (CNS) and play functions at the pre- and post-synapses (Arriagada-Diaz, et al., 2020). At the pre-synapses they support the endocytic recycling of synaptic vesicles (SVs), exerting differential roles depending on the pattern of neuronal activity (Tanifuji, et al., 2013). In this regard, dynamin-1 mediates SV recycling upon high-frequency stimulation but responding in a slow time window after the arrival of action potentials (APs); dynamin-3 works in a faster time window independent of the frequency of APs and dynamin-2 mediates SV resupply upon high-frequency stimulation with a fast kinetics (Tanifuji, et al., 2013).

At the postsynaptic level dynamin-2 is an important regulator of surface-membrane availability of
neurotransmitter-receptors (Carroll, et al., 1999; Kabbani, et al., 2004; Bhatnagar, et al., 2000; Wang, et
al., 2017). The latter appears to be especially relevant for glutamatergic excitatory synapses, as
dynamin-2 modulate insertion (Jaskolski, et al., 2009), removal (Carroll, et al., 1999; Chowdhury, et al.,
2006), and recycling (Lu, et al., 2007; Zheng, et al., 2015) of the α-amino-3-hydroxy-5-methyl-4-
isoxazolepropionic acid (AMPA) receptor to and from PSDs.
Although the isoforms -1 and -3 are expressed at a more greater level than dynamin-2 at the CNS
(Okamoto, et al., 2001), dynamin-2 seems to play a more critical role for synapse development
(Ferguson, et al., 2009). In fact, while embryos of dynamin-1 and/or dynamin-3 knockout (KO) mice
survive for weeks after born (Ferguson, et al., 2007; Raimondi, et al., 2011), the KO of dynamin-2 results
in early embryonic lethality (Ferguson, et al., 2009).
Mutations in dynamin-2 cause autosomal dominant human diseases (Züchner, et al., 2005; Bitoun, et al.,
2005; Tanabe, et al., 2009; Liu, et al., 2011; Koutsopoulos, et al., 2011; Böhm, et al., 2012; Sambuughin,
et al., 2015; Ali, et al., 2019). Among them, mutations in the middle and PH domain of dynamin-2 cause
the autosomal dominant form of centronuclear myopathy (CNM) a rare neuromuscular disorder
clinically manifested by myalgia, fatigability, weakness, and progressive atrophy of distal skeletal
muscles (Jeannet, et al., 2004; Fischer, et al., 2006; Böhm, et al., 2012). Cognitive deficiencies,
manifested as learning disabilities and limited intelligent quotient, have also been reported in CNM
patients (Jeannet, et al., 2004; Fischer, et al., 2006; Echaniz-Laguna, et al., 2007; Böhm, et al., 2012)
although the mechanisms involved have never been described. Here we show that a knock-in (KI) mouse
model bearing the CNM-causing p.R465W mutation in dynamin-2 (Durieux, et al., 2010; Durieux, et al.,
2012) exhibit a deficient capability to learn and acquire spatial and recognition memory. These cognitive
defects correlate with impaired hippocampal excitatory synaptic transmission and plasticity.
Furthermore, reduced dendritic spine density and changes in spine morphology are also evident in the
brain of these mice. This latter seem to rely on defects in the actin cytoskeleton organization. As the
catalytic activity of dynamin-2 is an important regulator of the actin dynamics (Mooren, et al., 2009; Gu,
et al., 2010; Yamada, et al., 2013; Yamada, et al., 2016; Lin, et al., 2020) and this is a mechanism affected
in dynamin-2 linked CNM (González-Jamett, et al., 2017) these data strongly suggest that CNM-causing
mutations perturb the synaptic role of dynamin-2, impacting on the actin-dependent dendritic-spine
structure and consequently affecting synaptic transmission and cognitive functions.

Results

Impaired learning and memory in adult CNM-mice

As early learning disabilities and cognitive defects have been reported in dynamin-2-related CNM
patients (Jeannet, et al., 2004; Fischer, et al., 2006; Echaniz-Laguna, et al., 2007; Böhm, et al., 2012) we
first evaluated learning and memory in a murine model of CNM. This is a knock-in mouse harboring the
p.R465W mutation located in the middle domain of dynamin-2, the most common mutation causing
CNM (Durieux, et al., 2010; Durieux, et al., 2012). Heterozygous mice (HTZ) recapitulate most of the
signs of the myopathy, starting at 1 month old (m.o) and progressing as the animal ages (Durieux, et al.,
2010). In order to assay learning and memory capabilities, adult HTZ mice over 6 m.o and their age-
matched wild type (WT) littermates were subjected to two different behavioral paradigms: the novel
object recognition test (NOR) and the Barnes maze (BM). Whilst NOR is useful to evaluate recognition

memory (Ennaceur, et al., 1988), BM is used to measure the ability of mice to learn and remember a
spatial location (Rizzo, et al., 2017; Negrón-Oyarzo, et al., 2015; Pitts, et al., 2018).

First, we performed the NOR test. A schematic representation of the test is shown in Figure 1a. As
expected WT mice spent significantly more time exploring a novel object (N) than a familial object (F)
along sessions (Figure 1b). However HTZ mice did not modify their preference between N and F (Figure
1b) suggesting defects in the recognition memory. Moreover, HTZ mice exhibited a significantly reduced
recognition index (D1) compared to their WT littermates (Figure 1c) strongly suggesting a decline in the
recognition memory in the CNM context.

Then we applied the BM test. A schematic representation of the test is shown in Figure 1d. WT and HTZ
mice were trained during 4 days (training) to find and enter into a hidden scape box using spatial clues.
On the fifth day a reversal phase started in which the position of the scape-box was changed but the
visual clues were kept to evaluate memory flexibility. As shown in Figure 1e the latency to find and enter
into the scape box was decreasing throughout the training sessions for both WT and HTZ mice.
However, HTZ mice took a longer time than WT to access the scape box throughout the reversal phase
(Figure 1f). In fact, on the last day of the reversal phase WT mice took on average 57.3 ± 18.2 s whereas,
HTZ mice took 100.6 ± 20.5 s to end the test (Figure 1f). Remarkably the percentage of HTZ mice that
completed the test at the last day of the reversal phase was significantly lower than WT mice (Figure 1g)
suggesting defects in spatial memory flexibility. Importantly, these observations did not appear to be
due to locomotion defects in HTZ mice as walking time percentage (Figure supplement S1a) and total
distance traveled in an open-field arena (Figure supplement S1b) were not different between groups.

Together these data show that the DNM2-linked CNM mouse model shares cognitive impairment with
CNM patients and represents a pertinent model to identify the underlying molecular mechanisms

Excitatory synaptic transmission and plasticity are defective in adult hippocampal slices from CNM-
mice

Changes in neuronal activity induce modifications in the efficacy of the synaptic transmission; this is
known as synaptic plasticity and the mechanisms regulating it are the basis of learning and memory
(Shors, et al., 1997). As synaptic plasticity in the hippocampus is particularly important for memory
formation (Lee, et al., 2011) and the most studied forms of synaptic plasticity are long-term potentiation
(LTP) and long-term depression (LTD) of the synaptic strength at the CA3-to-CA1 synapses (Malenka, et
al., 2004) we evaluated excitatory synaptic transmission and plasticity in hippocampal slices isolated
from HTZ mice over 6 m.o and their age-matched WT littermates.

The strength of the basal excitatory synaptic transmission was estimated as the I/O relationship. Figure
2a schematizes the experimental configuration for electrophysiological field recordings in hippocampal
slices. Figure 2b show representative I/O traces for WT and HTZ hippocampal slices. As shown in Figure
2c the I/O relationship was significantly impaired in HTZ compared to WT slices. HTZ slices exhibited
significantly lower fEPSP slopes (Figure 2d) but significantly higher FV-amplitudes (Figure 2e) compared
to the WT condition. To evaluate whether plasticity of the synaptic strength is also affected in the CNM-
context we induced NMDAR-dependent LTP in WT and HTZ hippocampal slices by using a standard theta

burst stimulation (TBS) protocol that drives a compound-LTP, involving pre- and post-synaptic
mechanisms (Bayazitov, et al., 2007). Whilst LTP induction relies on the activation of NMDAR- that
requires both presynaptic glutamate release and post-synaptic depolarization (Citri, et al., 2008), LTP
maintenance mainly relies on AMPAR trafficking at the PSDs (Anggono, et al., 2012a). As shown in Figure
3 the magnitude of the TBS-induced LTP was significantly reduced after 50 min of record in HTZ
compared to WT hippocampal slices (Figures 3a and 3c) suggesting a role of dynamin-2 in the
maintenance of the tetanus-induced LTP. Furthermore, the early potentiation of the synaptic response
measured during the first 5 minutes after the application of TBS was also reduced in HTZ compared to
WT slices (Figure 3b) suggesting that LTP-induction is also dependent on dynamin-2. To evaluate the
presynaptic contribution to the defects observed in HTZ synapses we estimated a paired-pulse
facilitation (PPF) index (Figure 3d-e). PPF is a form of short-term synaptic plasticity that occurs when two
stimulation-pulses of similar intensities are given in rapid succession, causing the second synaptic
response to be bigger than the first one (Zucker, et al., 2002). Facilitation appears to be due to an
increased presynaptic Ca^{2+} concentration that leads to enhanced neurotransmitter-release probability
(Zucker, et al., 2002). The PPF index was unchanged between HTZ and WT hippocampal slices (Figure 3d-
e) suggesting that, although presynaptic mechanisms could operate in the CNM-context, those appear
to be independent of the release probability. Hence, the p.R465W mutation in dynamin-2 seems to
mainly affect post-synaptic mechanisms, impairing synaptic potentiation upon sustained neuronal
activity. The main post-synaptic impact was confirmed by a chemically induced LTP using the NMDAR-
agonist Glycine (chLTP) which activates NMDAR without mediating a presynaptic glutamate release (Lu,
et al., 2001; Zhang, et al., 2014). Upon chLTP a significant reduction in the magnitude of synaptic
potentiation was observed in HTZ compared to WT slices (Figure supplement S2). Interestingly, these
postsynaptic mechanisms appear to predominantly affect excitatory synaptic transmission, as we
observed similar differences between HTZ and WT slices when the TBS protocol was elicited in the
presence of *picrotoxin* (*PTX*), a blocker of the GABAR-dependent inhibitory transmission (Figure
supplement S3).

Like LTP, LTD is a relevant long-term synaptic plasticity phenomena also implicated in learning and
memory (Malenka, et al., 2004; Citri, et al., 2008). We evaluated weather LTD is also affected in the
CNM-context and observed a significant reduction in the LTD-magnitude in HTZ compared to WT slices
(Figure supplement S4).

Altogether these data demonstrate that the excitatory synaptic transmission and plasticity are DNM2-
dependent processes impaired in the CNM context.

Dendritic-spine morphology is perturbed in neurons from adult CNM-mice

Besides a functional excitatory synaptic plasticity that mainly relies on changes in AMPAR activity at the
PSDs (Anggono, et al., 2012b) there is a structural plasticity involving morphological changes at dendritic
spines. Dendritic spines are actin-enriched protrusions where occur most of the excitatory synapses
(Bosch, et al., 2012). Their density, size and shape correlate with the degree of synapse maturity and
functionality and undergo modifications upon synaptic plasticity. In this regard it has been
demonstrated that LTP-induction increases dendritic spine density and promotes the transit towards a

mature-spine shape with higher head diameter leading to a higher head/spine length ratio (Matsuzaki,
et al., 2001; Okamoto, et al., 2009). On the contrary, LTD-induction reduces dendritic spine density and
leads to the shrinkage of pre-existing spines (Okamoto, et al., 2004). In order to evaluate whether the
defects in excitatory synaptic transmission and plasticity observed in HTZ slices were related to
structural defects in dendritic spines, we marked brains from 6 m.o WT and HTZ mice with a Golgi- -
staining and quantified dendritic spine density and morphology in hippocampal and cortical neurons.
Figures 4 and 5 summarize observations made on hippocampal and cortical pyramidal neurons,
respectively. The comparison of the mean morphological parameters revealed a significant reduction in
spine density (number of spines per 1 μm of dendritic shaft) in HTZ compared to WT hippocampal
neurons (Figure 4b). Although no significant differences were observed in the spine shape on average
(Figure 4c) HTZ hippocampal neurons tended to exhibit a lower percentage of filopodial-protrusions and
a higher percentage of stubby-like spines. To further evaluate dendritic spine structure, we quantified
the cumulative frequency of spine lengths, head diameters and head/length ratios. Remarkably,
although head diameters were not different between WT and HTZ hippocampal neurons (Figure 4d)
these latter exhibited a significantly higher frequency of shorter spines, producing a leftward shift in the
cumulative distribution curve compared to WT spines (Figure 4e). As a consequence of having shorter
dendritic spines, HTZ hippocampal neurons also exhibited a significantly higher head diameter/spine
length ratio compared to WT neurons (Figure 4f).
Similar to that observed in hippocampal neurons, HTZ cortical neurons exhibited a significantly lower
mean spine density (Figure 5b) and tended to show a lower proportion of filopodial-protrusions
compared to the WT condition (Figure 5c). Remarkably, cortical HTZ neurons exhibited a higher
frequency of spines with smaller heads (Figure 5d) and shorter lengths (Figure 5e) although the
head/length ratio was unchanged (Figure 5f).
Together, these data suggest that pyramidal neurons bearing the dynamin-2 p.R465W mutation may
undergo modifications in density, head width and length of dendritic spines.

Impaired F-actin organization in CNM-neurons

As dendritic spine morphology critically depends on the underlying actin network (Okamoto, et al.,
2009; Hotulainen, et al., 2010; Matus, et al., 2000) and actin dynamics is significantly affected in the
CNM context (González-Jamett, et al., 2017) we next evaluated actin organization in brains of WT and
HTZ mice. Relative amounts of filamentous (F-) and monomeric (G-) actin were quantified in the
hippocampus and cortex. Figure 6 shows representative blots from hippocampal (Figure 6a) and cortical
tissues (Figure 6b) and the quantification of the respective F/G ratios. Our results indicated that the
balance between F- and G-actin is perturbed in brains of HTZ compared to WT mice. Specifically the F/G
ratio increased from 1.9 ± 0.5 to 3.9 ± 1.3 in HTZ compared to WT hippocampal tissue (Figure 6a). In the
cortex the F/G ratio increased from 0.6 ± 0.01 to 2.1 ±0.8 in HTZ compared to WT brains (Figure 6b).
These data suggest that the modifications in dendritic spine morphology experienced by HTZ neurons
(Figures 4-5) could rely on actin cytoskeleton disturbances in adult mice. To further explore this
possibility early in the disease development, we evaluated dendritic spine morphology in hippocampal
neurons from WT and HTZ neonates (P0). 7 days *in vitro* (7*DIV*) neuron primary cultures were fixed,
stained with the F-actin-binding toxin phalloidin-rhodamine-B and visualized by confocal microscopy.
Unlike what we observed in neurons from adult mice (Figures 4-5), the mean density of spines stained
with phalloidin was not different between neonate-derived WT and HTZ neurons (Figure 7b). However
HTZ neurons tended to exhibit a higher percentage of filopodial protrusions and a lower proportion of
stubby-like spines compared to WT neurons (Figure 7c) suggesting that the actin-dependent spine
development is delayed in HTZ neurons. It is noteworthy that we were not able to observe branched
mature spines in cultured neurons. We also quantified the cumulative frequency of spine head

diameter, spine length and head/length ratio. Although head diameters of phalloidin stained-spines
were not different between WT and HTZ neurons (Figure 7d), HTZ neurons exhibited a significantly
higher frequency of longer spines (Figure 7e) and lower head/spine ratios compared to WT neurons
(Figure 7f). This latter supports the idea that actin-dependent dendritic spine features could be early
impaired in the CNM context. In this regard, the phalloidin-fluorescence intensity was significantly lower
in HTZ compared to WT spines (Figure supplement S5) suggesting that the F-actin network underlying
HTZ spines is impaired.
All together the results presented here demonstrate that dynamin-2 is an important regulator of the
excitatory synaptic transmission in the mammalian brain, by modulating actin-dependent dendritic
spine structure. This dynamin-2 function appears to be perturbed by the p.R465W mutation, impairing
synaptic structure and function and consequently leading to cognitive defects in the CNM context.
259
Discussion

Dynamin-2 is a large GTP-ase, member of the dynamin superfamily, that regulates membrane
remodeling and actin dynamics, orchestrating actin and membrane-based processes such as exocytosis
(González-Jamett, et al., 2013; Shin, et al., 2018), endocytosis (Van der Bliek, et al., 1993; Damke, et al.,
1994; Loerke, et al., 2009) and endosomal recycling (Nicoziani, et al., 2000) among others.
In the nervous system, dynamin-2 is expressed at the pre-and post-synaptic levels (Okamoto, et al.,
2001). At the pre-synapses dynamin-2 participates in the endocytic recycling of SVs, allowing the
resupply of the ready-releasable pool in response to high frequency neuronal activity (Tanifuji, et al.,
2013). At the post-synapses dynamin-2 regulates the availability of neurotransmitter receptors at the
surface membranes (Carroll, et al., 1999; Kabbani, et al., 2004; Bhatnagar, et al., 2001; Wang, et al.,
2017). Of particular relevance for excitatory synaptic transmission is the role played by dynamin-2 in
AMPAR trafficking. The GTP-ase activity of dynamin-2 is required for the insertion of AMPAR into spines,
by mediating their lateral diffusion from dendritic shafts to PSDs (Jaskolski, et al., 2009). In addition,
dynamin-2 GTP-ase activity has been implicated in the removal (Carroll, et al., 1999; Chowdhury, et al.,
2006) and recycling (Lu, et al., 2007; Zheng, et al., 2015) of AMPARs from and to PSDs. Hence, as AMPAR
mediates the majority of fast excitatory synaptic transmission in the mammalian brain (Anggono, et al.,
2012a), dynamin-2 could play a key regulatory function, supporting synaptic transmission and plasticity.
In agreement with this idea, pharmacological inhibition of dynamin´s GTP-ase activity with dynole 34-2
(Arriagada-Diaz, et al., 2020) or dynasore (Fa, et al., 2014) significantly reduces LTP in hippocampal
slices. Here, we demonstrate that the CNM-causing p.R465W mutation negatively impacts on the
synaptic function of dynamin-2, leading to defects in excitatory synaptic transmission and plasticity in
the brain of HTZ mice. As dynamin-2 is mostly expressed at the post-synaptic level (Okamoto, et al.,
2001) it is likely that these HTZ defects in synaptic transmission rely on post-synaptic mechanisms such
as AMPAR trafficking. Indeed, we observed a reduction in the magnitude of LTP in HTZ slices when we
induced potentiation both electrically as well as when we did it chemically with glycine. As the chLTP
protocol elicits potentiation of the synaptic strength by directly activating NMDARs without mediating
the presynaptic release of glutamate (Lu, et al., 2001; Zhang, et al., 2014), these results are consistent
with major post-synaptic mechanisms underlying HTZ synaptic defects. In fact, the PPF index was
unchanged between WT and HTZ hippocampal slices (Figure 3d). As PPF is an estimation of the
neurotransmitter release probability (Zucker, et al., 2002) it is unlikely that the changes that we
observed in excitatory synaptic transmission in HTZ hippocampal slices are related to changes in the
glutamate release probability. This is agreement with a recent report of Moro and collaborators who
found that dynamins are dispensable for synaptic vesicle exocytosis (Moro, et al., 2021).However, we
cannot rule out presynaptic defects in the CNM context. The fact that the amplitude of the fiber volley
was significantly higher in HTZ compared to WT slices (Figure 2) could mean that more presynaptic

terminals are recruited in order to compensate a reduction in the SV releasable pool. As dynamins´s
GTP-ase activity of dynamin-2 regulates SV recycling (Tanifuji, et al., 2013; Watanabe, et al., 2013;
Cheung, et al., 2019) it is possible to visualize a scenario in which the p.R465W mutation in dynamin-2
affects the size of the SV releasable pool, impacting neurotransmission. More experiments are needed
to address this possibility.
Regardless of whether defects were pre or post-synaptic, there was a significant impairment in synaptic
transmission and plasticity in HTZ compared to WT brains. Congruently, HTZ mice exhibit a cognitive
decline, manifested as defects in recognition and spatial memory which can be explained by
perturbations in the synaptic function. In this regard, even though CNM primarily affects skeletal
muscles (Romero, et al., 2011; González-Jamett, et al., 2017) learning disabilities and limited intelligent
quotient have been reported in dynamin-2-linked CNM patients (Jeannet, et al., 2004; Fischer, et al.,
2006; Echaniz-Laguna, et al., 2007; Böhm, et al., 2012). As we observed here in HTZ mice, cognitive
defects manifested by CNM patients could be due to disturbances in the synaptic function. Remarkably,
in addition to changes in the synaptic strength, we also observed structural modifications in dendritic
spines of the HTZ compared to WT neurons. In this regard, dendritic spine density was significantly
lower in pyramidal neurons from adult HTZ mice (Figures 4-5). As the majority of the synaptic contacts
between excitatory neurons are made on dendritic spines (Harris, et al., 1994) a net reduction in spine
density can lead to significant defects in synaptic transmission, such as those observed in HTZ
hippocampal slices. Moreover, dendritic spine morphology, which is closely related to the maturity and
functionality of synapses (Holtmaat, et al., 2009; Gipson, et al., 2017), also underwent modifications in
HTZ compared to WT neurons (Figures 4, 5 and 7). Although on average we did not observe significant
differences in spine morphology between genotypes, the cumulative distribution of the spine length,
spine head diameter and head/length ratio yielded important variations. Whilst mature HTZ
hippocampal neurons exhibited a higher frequency of shorter spines, neonatal hippocampal neurons
cultured from HTZ pups showed a higher frequency of immature larger spines with a significant
reduction in the head diameter/spine length ratio. Moreover, cortical neurons from adult HTZ brains
exhibited higher frequency of shorter spines, with smaller heads compared to WT neurons. As spines
with large head and large head/spine ratio have been found to correlate with an increased PSD area and
AMPAR density (Matsuzaki, et al., 2001) a reduction in spine head diameter or the head/length ratio
could explain defects in synaptic transmission and plasticity.
A major factor regulating spine morphology is the actin cytoskeleton (Okamoto, et al., 2004; Bosch, et al., 2014). A
dynamic F-actin network exists in dendritic spines which act primarily as a cytoskeleton component defining spine
architecture, but also as a scaffold for the recruitment of post-synaptic proteins including AMPAR (Okamoto, et al.,
2009; Cingolani, et al., 2008; Stefen, et al., 2016).Therefore, remodeling of spine-actin is critical for structural and
functional synaptic plasticity (Kim, et al., 1999; Okamoto, et al., 2004) and it is not surprising that perturbations in
actin dynamics lead to different synaptopathies (Penzes, et al., 2012; Pelucchi, et al., 2020). Thus, the changes that
we observed in the density and shape of dendritic spines in HTZ neurons are compatible with defects in actin
dynamics. Indeed, we previously demonstrated that actin organization and remodeling are disrupted in muscle
cells of HTZ mice (González-Jamett, et al., 2017) and our results here suggest an imbalance between F- and G-actin
in the hippocampus and cortex of adult HTZ mice. The fact that the observed differences did not reach statistical
significance when compared total brain homogenates (Figure 6) could be due to the F/G imbalance is more
important in neurons than in other cell types. If that is true such differences may be attenuated in total
homogenates in which more of the cells are glia. An increase in the F/G ratio as we observed in HTZ compared to
WT brains (Figure 6) could be related to a less dynamic neuronal actin network, with a reduced ability to remodel
itself to support synaptic transmission. In the case of neonatal cultured neurons, the fluorescence intensity of the
F-actin-binding toxin phalloidin at the dendritic spines was significantly lower in the HTZ compared to WT condition
(Figure supplement S5). This latter strongly suggests that the F-actin network that supports dendritic spine
structure is perturbed in HTZ neurons. But, how the p.R465W mutation could impact on dynamin-2 function
leading to defects in actin and actin-based mechanisms? The residue 465 localizes near to a putative direct actin-
binding motif in dynamin-2 (Gu, et al., 2010, red region in Figure 8a). As these authors proposed, unassembled

dynamins could bind to short actin filaments via their middle domain and in turn oligomerize in rings promoting
actin polymerization (Gu, et al., 2010; Gu, et al., 2014; Shiffer, et al., 2015). If that is true, the substitution of a
positive-charged arginine by a non-polar tryptophan in the oligomerization-middle domain of dynamin-2 could
certainly influence its actin-binding properties affecting actin dynamics. In this regard, by using full-atom and
coarse-grained molecular simulations, we previously demonstrated that the p.R465W mutation affects the
interaction between the residue 465 and surrounding amino acids, modifying the conformation of the BSE and
consequently impacting on the GTPase-domain position and dynamin-2 self-assembly (Hinostroza, et al., 2020).
Recent evidences suggest that helix of dynamin-2 interact with F-actin via their outer-rim (Zhang, et al., 2020). This
region includes the C-terminus PRD that is adjacent to the GTPase domain and to the BSE (Figure 8b). It is
noteworthy that the R/W 465 residue interacts with the BSE in a region that is close to the starting point of the
PRD (Figures 8b-c). As the p.R465W mutation in dynamin-2 extends the angle between the G domain and the BSE
(Hinostroza, et al., 2020) in the tetramer and helical configuration (Figure 8d) we speculate that the
arginine/tryptophan substitution in the residue 465 not only modifies the BSE conformation but also changes the
position of the PRD, being able to affect the PRD-actin interaction. This assumption is in agreement with a model in
which the actin-remodeling activity of dynamin-2 is disrupted by CNM-causing mutations. In accordance with this
idea, Lin and collaborators recently demonstrated that dynamin-2 regulates the bundling of actin at the
postsynaptic membranes of neuromuscular junctions (NMJs). Remarkably, CNM-linked mutations seem to disrupt
this dynamin-2 function impairing actin remodeling and perturbing postsynaptic structures at the NMJs (Lin, et al.,
2020). A similar mechanism could be operating at central excitatory synapses in the brain of HTZ mice, explaining
the defects that we observed in dendritic spine morphology, synaptic transmission and cognitive functions.

In our knowledge, this is the first report addressing the mechanisms that lead to cognitive defects in dynamin-2-
linked CNM. Our data support a model in which the actin cytoskeleton organization is impaired, leading to
structural and functional synaptic defects. In addition, our results contribute to the better understanding of the
still scarce knowledge about the function of dynamin-2 at central synapses.

Methods

Animals
HTZ mice harboring the p.R465W mutation in dynamin-2 is a mammalian model of CNM. These mice
recapitulate most of the CNM-signs, which start at 1 mo and progress as animals age (Durieux et al.,
2010). HTZ and WT littermates (C57BL/6 strain) were housed at room temperature with *ad libitum*
access to food and water. Mice were maintained on light-darkness cycles of 12–12 h, according to
standard protocols. Genotyping was performed by PCR as previously described (Durieux et al., 2010)
using DNA extracted from ears. Primers used were: 3'-CTGCGAGAGGAGACCGAGC-5' (forward) and 3'-
GCTGAGCACTGGAGAGTGTATGG-5 (reverse). PCR products were electrophorated in agarose gels and
bands of 445 bp and 533 bp, representing the WT and p.R465W mutated allele respectively, were
detected. All animal protocols described here were conducted in accordance with the approved
protocols of the Institutional committee for the care of laboratory animals of Universidad de Valparaíso
(BEA131-18).

383
384
385
Behavioral tests
Recognition and spatial learning and memory were assessed in 6 m.o WT and HTZ mice by application of
the Novel Object Recognition (NOR) test and the Barnes maze test (BM), respectively. Locomotion
capabilities in mice were evaluated by application of a standard Open Field (OF) test. All behavioral

tests were carried out in a room conditioned for this purpose, with controlled temperature (21 ° ± 2 ° C),
white noise (50 dB) and regulated luminosity (200 lux).
392
NOR. This test is based on the spontaneous tendency of rodents to spend more time exploring a novel
object than a familial one. Maze consisted of a 50 cm x 40 cm x 63 cm square white acrylic box in which
mice explored freely during 5 min throughout three phases: (i) sample, where mice explored a pair of
identical objects; (ii) retention, where mice were removed for cleaning and changing objects; and (iii)
choice, where mice explored a pair of different objects: a familiar object (F) and a new object (N). Each
session was repeated during 3 days. The time that mice spent exploring F or N was quantified. A
recognition index (D1) was calculated as [N exploration time-F exploration time.
BM. This task is dependent on the intrinsic inclination of rodents to escape from an aversive
environment (Barnes 1979). It consists in a 70 cm diameter white circular platform elevated 90 cm from
the floor with 20 equally spaced holes along the perimeter (each 7 cm in diameter) located 2 cm from
the edge of the platform. The animals are oriented by means of spatial clues located on the walls of the
room, in order to find and remember the position of a black plexiglass escape box (17 × 13 × 7 cm) that
is hidden under one of the holes. The maze was illuminated with 4 incandescent lights to produce a light
level of around 600 lux fall on the circular platform. BM test consisted of 9 session days, 4 trials per
session along which three phases were carried out: (i) pre-training phase, where mice were trained to
find the escape platform (ii) training phase, where mice remember the position of the escape box
guided by the spatial clues arranged in the room, (iii) reversion phase, where the location of the escape
box was changed but maintaining the position of the spatial clues to evaluate memory flexibility. In all
phases, if mice did not find the target hole and did not enter into the escape box during 3 minutes, the
test was terminated. Subsequently, the experimenter guided the animal to the escape box and allowed
it to remain in the box for 1 minute before returning to their holding box. Latency or time that took to
animals to find the target hole and enter into the scape box was quantified.
OF. Mice were placed in the center of a 50 cm x 40 cm x 63 cm white acrylic box and left to explore
freely during 5 minutes, along 3 sessions (1 session per day). Total distance traveled and the percentage
of walking time was quantified.
418
Field electrophysiological recordings
To evaluate basal strength of synaptic transmission and synaptic plasticity we evoked field excitatory
postsynaptic potentials (fEPSPs) in WT and HTZ hippocampal slices by stimulating the Schaffer
Collaterals (CA3) and recording the synaptic responses in *stratum radiatum* of CA1. Hippocampal slices
were prepared as previously described (Ardiles et al. 2014). Animals were euthanized under deep
anesthesia using isoflurane at saturation and their brains were quickly removed after craniotomy. Brains
were sectioned in two hemispheres, removing hippocampus to be sectioned in 300 μm thick slices and
immersed in dissection buffer (in mM: 212.7, sucrose 26 $NaHCO_3$, 1.23 NaH_2PO_4, 10 D-glucose, 5 KCl, 2
$CaCl_2$, 1 MgCl, 3 pyruvate) using vibratome (Vibratome 1000 plus, Ted Pella Inc., CA, USA). Slices were
transferred and kept for 1 hour at room temperature in artificial cerebrospinal fluid (ACSF, in mM: 124
NaCl, 26 $NaHCO_3$, 1.23 NaH_2PO_4, 10 D-glucose, 5 KCl, 2 $CaCl_2$, 1 MgCl, 3 pyruvate). All recordings were
obtained in an immersion chamber perfused with ACSF (30 ± 0.5°C; 2 ml / min). Stimulation was
performed with pulses of 0.2 ms duration, administered through concentric bipolar stimulation
electrodes. Basal responses were recorded using mean stimulation intensity compared to maximum.
Basal synaptic transmission was analyzed by determining input-output relationship (I /O) of the fEPSP

generated by gradually increasing the stimulation intensity. Input corresponded to the amplitude of the
fiber volley and the output to the initial slope of the fEPSP. To evaluate long-term potentiation (LTP),
slices were subjected to a high-frequency electrical stimulation protocol or a chemical stimulation
protocol. LTP using high-frequency electrical stimulation was performed using a standard theta burst
(TBS) protocol, which consists of 4 trains at 100 Hz for 1 min (Flores-Muñoz et al. 2020; Gajardo et al.
2018). On the other hand, chemical LTP (chLTP) was induced by perfusing a glycine solution (glycine, Gly
600 μM + *picrotoxin*, PTX 50 μM + strychnine Stric 3 μM) (X. Y. Zhang et al. 2014), directly on the
hippocampal slices during 5 minutes. Gly is co-agonist of NMDAR, PTX is an inhibitor of the inhibitory
GABAR-dependent transmission and Stric is an inhibitor of Gly-receptors. Once the application time of
solution had elapsed, it was removed and the registration buffer was replaced with a new one. In both
cases, fEPSPs were recorded for 20 minutes prior to stimulation and for 50 minutes after the application
of the electrical stimulus (TBS) or chLTP.
To assess long-term depression (LTD), a standard low-frequency electrical protocol was induced which
consists of a 1 Hz stimulation for 15 minutes (Gajardo et al. 2018). fEPSPs were recorded for 20 minutes
prior to stimulation and for 60 minutes after the application of LFS.
Paired pulse facilitation (PPF) was obtained by stimulating at different intervals in a range between 25-
300 ms and recording excitatory post-synaptic potential (EPSP). A PPF index was estimated by dividing
the amplitude of the second response over the first one (R2/R1).
452
Spine density and morphology
HTZ mice over 6 mo and age-matched WT littermates were sacrificed, brains were quickly removed and
processed using the FD Rapid Golgi-Stain-TM kit (FD Neuro Technologies) according to the manufacturer
instructions. Golgi impregnated cortical and hippocampal neurons were imaged using a Leica DM500
microscope equipped with a camera system, a 63X /1.40 oil (HCPL Apo, Leica) objective and the Leica
acquisition software. Pyramidal neurons were analyzed and processed using the ImageJ software (NIH,
USA). To evaluate the spine density and morphology, measurements were performed in dendritic
segments of 10 to 20 μm, visualized from four apical dendrites per neuron. Dendritic- protrusions were
categorized following the parameters revised by Risher and collaborators (Risher et al., 2014). Long, thin
without-head protrusions of more than 2 um long were classified as immature filopodia; wide-head
(>0.6 um width) and short protrusions (<1 um) were classified as mature mushroom-spines; wide-
head/without neck protrusions (length: width ratio <1 um) were classified as stubby-spines; thin and
short (<2 um) headed protrusions were classified as thin-spines. Cup-shaped protrussions were
classified as branched spines (Figure Supplement S6). Measurements were made by two different
experimenters, blind to the genotype of the samples.
468
469
470
Neuron primary culture
Hippocampal neurons were primary cultured from post-natal mice (P0) following the protocol described
by Beduoain and collaborators (Beduoain et al., 2012). Briefly, hippocampi from individual pups were
dissected, trypsinized for neurons dissociation and plated in poly-L-lysine treated coverslips and
maintained in medium containing B27-supplement (GIBCO), glutamax (GIBCO) and
Penicilin/Streptomycin mixture of antibiotics (GIBCO). Genotype of each single culture was confirmed a
posteriori from tail-segments of pups. 2 days after culture neurons were treated with 5μM cytosisne-
arabinoside (Ara-C) to inhibit the proliferation of non-neuronal cells. At 7 *DIV,* neurons were fixed with

4%PFA + 4% sucrose during 10 min at 37°C, permeabilized with 0.1% triton-X100 in saline phosphate
buffer (PBS, mM: 137 NaCl, 2.7 KCl, 10 Na$_2$HPO$_4$, 2 KH$_2$PO$_4$, pH 7.4) during 10 min at room temperature
and stained during 1 h with 1μM phalloidin-Rhodamin-B (Sigma) before confocal visualization. Images
were acquired in a confocal microscope (upright Eclipse Nikon 80i) using a 60x magnification objective.
Single confocal images were acquired with the EZ-C1 (Nikon) software using similar acquisition settings
between comparable samples. Analyses of dendritic spines stained with phalloidin were made with the
ImageJ software (NIH, USA) by two different experimenters, blind to the genotype of the samples. The
phalloidin fluorescence intensity was measured inside spines, without including the neighboring
dendritic shaft.
488
F/G actin assay
Hippocampal and cortical tissue from WT and HTZ mice were lysed and homogenized in conditions that
stabilize filamentous (F-) and monomeric (G-) actin using an F/G actin commercial assay (Cytoskeleton
Inc.). Lysed extracts were ultracentrifuged at 100,000 g for 1 h at 37 °C in order to separate pellets (F-
actin) and soluble (G-actin) fractions. The F-actin pellet was resuspended in a depolymerizing buffer
(Cytoskeleton Inc) and then F-and G-actin fractions were diluted in loading buffer (50 mM Tris–HCl, 2%
SDS, 10% glycerol, 1% beta-mercaptoethanol and bromophenol blue). Samples were subjected to
electrophoresis in 12%-SDS-PAGE, transferred to a PVDF membrane, blocked for 1 h with 5% non-fat
milk in TBS-T (mM:150 NaCl, 50 Tris/HCl, pH 7.4, 0.05% Tween-20), incubated with a polyclonal antibody
against actin (1:500; Cytoskeleton Inc.) and developed by chemiluminescence. Densitometric analysis of
the F- and G-actin bands was performed with the ImageJ software.
500
Statistical analyses
GraphPad Prism 9 software was used to perform the statistical analyses. Paired or unpaired non-
parametric Mann-Whitney test were used to compare WT and HTZ conditions. Two-way ANOVA
followed by Bonferroni multiple comparison post-test was also used in electrophysiological experiments.
Data in graphs represent the mean ± SEM, with a p-value <0.05 being significant. If not specified,
differences were not statistically significant.
507
508 **Declaration of Conflicting Interests**
509

510 The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or
511 publication of this article.
512
513 **Acknowledgements**
514 We thank Dr. Isaac García-Carrillo for facilitating facilities for western blot development. This work was
515 supported by the grants Fondecyt 11180731 (to AMG-J), Fondecyt 1201342 (to AOA), and ICM-ANID
516 ICN09-022 CINV (to AMG-J, AOA, and AMCD) and by ANID scholarship 22200154 (to JA-D).

Figure 1: HTZ mice exhibit reduced performance in recognition and spatial memory tasks. WT (black circles) and HTZ (red circles) mice over 6 m.o were evaluated in the novel object recognition (NOR) and the Barnes maze tests to estimate recognition and spatial memory, respectively. **(a)** Representative scheme of the NOR test. Mice are faced with two identical objects (F1 and F2) in the sample phase and then in the choice phase, the objects are replaced by an extra copy of the familiar object (F) and a novel object (N). **(b)** Mean exploration time for familial (F) and novel (N) objects. Note that HTZ mice spend significantly less time exploring new objects compared to WT mice. Wilcoxon test, *p =0.0312 compared to the WT condition. **(c)** Mean recognition index (D1). Positive values indicate recognition of the novel over the familial object. Mann-Whitney test, *p =0.0221 compared to the WT condition **(d)** Schematic representation of the BM test. Mice were trained for 4 days to find and escape to a hidden dark box using visual cues (Training phase). On the fifth day, the escape box is located in the opposing quadrant keeping the visual cues to evaluate spatial memory flexibility (Reversal phase). **(e)** Quantification of the latency to entry into the escape box throughout the training and reversal sessions.

(f) Mean latency to entry into the escape box at day 8 of the test (last day of the reversal phase). **(6)** Quantification of the percentage of animals that completed the task at the end of the reversal phase. Note that the the time to enter into the escape box was not statistically significant, but the percentage of HTZ mice that completed the task on the last day of the reversal phase was significantly lower than WT. Mann-Whitney test, *p =0.0286 compared to the WT condition All data are represented as mean ± SEM. In parentheses is the number of WT and HTZ mice.

Figure 2: Impaired basal excitatory synaptic transmission in HTZ hippocampal slices **(a)** Scheme of the
experimental configuration for field recordings in Schaffer Collaterals (Sc) to CA1 hippocampal synapses.
The stimulation electrode and the recording electrode (R) were positioned in *stratum radiatum* of CA1.
(b) Representative fEPSP traces of both experimental groups HTZ (red) and WT (black**).** The numbers in
parentheses are slices, animals recorded. **c)** Relationship between the slope values of fEPSPs and the
amplitude of FVs obtained between each experimental groups. The R^2 values for the respective linear
regressions are shown. **(d)** Mean fEPSP slopes at different intensity values are plotted per each
experimental group. Note the significant difference in fEPSPs as the intensity of stimulation increased
between WT and HTZ slices. **(e)** Mean FV amplitudes for WT and HTZ slices. Note the significant
difference in FV amplitudes between WT and HTZ slices as the stimulus intensity increased. (WT: n = 18
slices, 5 mice; HTZ: n = 9 slices, 3 mice). All data are represented as mean ± SEM. Statistical differences
were calculated using a t-test, * $p < 0.0001$ compared to the WT condition.

598
599
Figure 3: TBS-induced LTP impairment in excitatory synapses of HTZ hippocampal slices. **(a-c)**
Hippocampal slices of 6 m.o of WT (black circles) and HTZ (red circles) mice were stimulated with a
standard TBS protocol to induce LTP and excitatory postsynaptic field potentials (fEPSP) were recorded
during 50 min after the application of the TBS protocol. 2-Way ANOVA, $F_{(1,2257)}$ = 1414, *p <0.0001
followed by Bonferroni post hoc test (p < 0.05) compared to the WT condition. Representative traces
before (gray) and after (black and red) the application of TBS are shown as inserts. Note that at the end
of the recording (c), as well as after 5 min of LTP-induction (b) HTZ slices exhibited a significantly
reduced potentiation of the synaptic strength compared to WT slices (WT: n = 17 slices, 5 mice; HTZ: n =
22 slices, 7 mice). Mann-Whitney test, *p <0.0001 compared to the WT condition in both 5 or 50 min
after TBS. The numbers in parentheses are slices, animals recorded **(d)** Representative traces of HTZ
(red) and WT (black) slices stimulated with two stimuli of the same intensity separated by a time of 50
ms for PPF index estimation. Note that the amplitude of the second response was greater than the first
one in both cases. **(e)** Paired pulse facilitation by different stimulation intervals. Note that no significant
differences were observed between the experimental groups (WT: n = 26, slices 6 mice; HTZ: n = 30
slices, 7 mice). All data are represented as mean ± SEM.
615

Figure 4: Reduced spine density and spine length in hippocampal neurons from adult HTZ brains. Brains
from WT (black bars) and HTZ (red bars) mice over 6 m.o were isolated and Golgi-stained for
visualization of pyramidal hippocampal neurons **(a)** Representative neurons and dendritic shafts per
genotype. Black and red arrows point spines in WT and HTZ dendrites respectively. Black arrowhead
highlights a branched spine. Scale bar= 2.5 μm **(b)** Mean spine density per neuron (number of spines per
1 μm of dendritic segment) **(c)** Mean percentage of filopodium, thin, stubby, mushroom and branched
spines in WT and HTZ hippocampal neurons. 4 dendritic shafts per neuron were analyzed. N=8 WT
neurons and 19 HTZ neurons from at least 2 different animals per genotype. Bars are mean ± SEM **(d-f)**
Average (i) and cumulative distribution (ii) of spine head diameters (d), spine lengths (e) and
head/length ratios (f) in WT and HTZ Golgi-stained neurons. N=217 WT spines, 332 HTZ spines. *p<0.005
versus WT condition, Mann-Whitney 2-tail test for non-parametric data.

Figure 5: Reduced spine density and spine head diameter in cortical neurons from adult HTZ brains.

Brains from WT (black bars) and HTZ (red bars) mice over 6 m.o were isolated and Golgi- stained for

visualization of pyramidal cortical neurons **(a)** Representative neurons and dendritic shafts per

genotype. Black and red arrows point spines in WT and HTZ dendrites respectively. Scale bar= 2.5 µm **(b)**

Mean spine density per neuron **(c)** Mean percentage of filopodium, thin, stubby, mushroom or

branched spines in WT and HTZ cortical neurons. 4 dendritic shafts per neuron were analyzed N= 8 WT

neurons and 19 HTZ neurons from at least two different animals per genotype. Bars are mean ± SEM **(d-**

f) Average (i) and cumulative distribution (ii) of spine head diameters (d), spine lengths (e) and

head/length ratios (f) in WT and HTZ Golgi-stained neurons. N=155 WT spines, 328 HTZ spines. *p<0.005

versus WT condition, Mann-Whitney 2-tail test for non-parametric data.

Figure 6: Imbalance in the F/G actin ratio in brains of HTZ mice. Representative blots of the relative amounts of F- and G-actin in the hippocampus (a) and cortex (b) of WT and HTZ mice are shown at the top. Graphs below summarize the quantification of the densitometry analysis of the F/G actin ratio in hippocampus (a) and cortex (b) isolated from WT and HTZ mouse brains. Bars are mean ± SEM of tissues from 3 WT and 3 HTZ mice. Statistical comparison was done using Mann-Whitney test for non-parametric data. p=0.200 and 0.400 for hippocampus and cortex, respectively. Although not significant HTZ brain tissue tended to exhibit an increased F/G ratio compared to the WT condition.

Figure 7: Reduced spine head/length ratio in cultured HTZ neurons. 7 *DIV* hippocampal neurons, primary
cultured from WT and HTZ pups (P0), were fixed, stained with the F-actin binding toxin phalloidin-
rhodamin-B and visualized by confocal microscopy **(a)** Representative phalloidin-stained neurons and
dendritic shafts per genotype. Yellow arrows point typical dendritic spines. Scale bar= 1.5 µm **(b)** Mean
spine density per neuron **(c)** Mean percentage of filopodium, thin, stubby or mushroom spines in WT
and HTZ phalloidin-stained neurons. Branched spines were not observed neither in WT nor in HTZ
neurons. 3 dendritic shafts per neuron N= 3 WT neurons and 18 HTZ neurons from 2 different cultures.
Bars are mean ± SEM **(d-f)** Average (i) and cumulative distribution (ii) of spine head diameters (d), spine
lengths (e) and head/length ratios (f) in WT and HTZ phalloidin-stained neurons. N=61 WT spines, 166
HTZ spines. *$p<0.005$ versus WT condition, Mann-Whitney 2-tail test for non-parametric data.

707
708
Figure 8: The CNM-causing p.R465W mutation localizes near to a putative direct actin-binding motif in
dynamin-2 **(a)** Corse-grained model of a dynamin-2 helix surrounding a lipid nanotube. Dynamin-2 is
represented in silver and gray; Putative actin-binding region proposed by Gu et al., (2010) (residues 399
to 444 in the middle domain) are shown in red; the residue R/W465 is represented in green and the lipid
nanotube in yellow. **(b)** Tetramer configuration in the dynamin-2 helix. G-domains are colored in blue,
BSEs are represented in yellow, stalks are shown in red and PH domains in lime. The position of PRD is
represented only in two dynamins-2 as a silver circles. Green dot represent R/W465. **(c)** Amino acids of
the BSE that are close to the R/W465. This interaction occurs in a zone close to the PRD. **(d)** G-domains
and BSE of WT and mutant dynamin-2. The extension of BSE in the mutant dynamin-2 changes the
position of the G-domain.

719
720

721
722 **Figure S1**: Locomotion activity is not different between WT and HTZ mice. WT (black circles) and HTZ
723 (red circles) mice over 6 m.o were left to explore freely in an open field arena during 5 min. (a) Mean
724 percentage of the walking time and (b) total distance traveled are plotted per each experimental group.
725 All data are represented as mean ± SEM. N= 6 WT and 7 HTZ mice. Statistical differences were
726 calculated using a non-parametric Mann-Whitney test.

746
Figure S2: Impaired glycine-induced LTP in HTZ hippocampal slices. **(a-b)** Hippocampal slices from 6 m.o
WT (black circles) and HTZ (red circles) mice were stimulated with the application of 600 μM glycine + 50
μM PTX + 3 μM strychnine in order to chemically induce LTP and fEPSPs were recorded during 50 min. 2-
Way ANOVA, $F_{(1,480)}$ = 208.2, *p <0.0001 followed by Bonferroni post hoc test (p < 0.05) compared to the
WT condition. Representative traces before (gray) and after (black and red) the application of glycine
solution are shown as inserts. Note that HTZ hippocampal slices exhibit a significantly reduced
potentiation at the end of the recording (b) compared to the WT slices (N= 5 slices, 2 WT mice; 5 slices, 2
HTZ mice). Mann-Whitney test * p <0.005 compared to the WT condition. All data are represented as
mean ± SEM.

Figure S3. LTP impairment in HTZ hippocampal slices is independent of inhibitory synaptic transmission.

(a-b) Hippocampal slices from 6 m.o WT (black circles) and HTZ (red circles) mice were stimulated with a

standard TBS protocol in the presence of the GABAR-inhibitor Picrotoxin (PTX). fEPSPs were recorded

during 50 min after TBS-application. 2-Way ANOVA, $F_{(1,671)}$ = 245.8, *p <0.0001 followed by Bonferroni

post hoc test (p < 0.05) compared to the WT condition. Representative traces before (gray) and after

TBS are shown as inserts. Note that even in the presence of PTX, HTZ hippocampal slices exhibit a

significantly reduced potentiation compared to WT slices, suggesting that inhibitory transmission is not

involved in the synaptic plasticity defects in HTZ slices. N= 6 slices, 2 WT mice/ 7 slices, 2 HTZ mice).

Mann-Whitney test * p <0.005 compared to the WT condition. All data are represented as mean ± SEM.

780
Figure S4. Impaired LFS-induced LTD in HTZ hippocampal slices. **(a-b)** Hippocampal slices from 6 m.o WT
(black circles) and HTZ (red circles) mice were stimulated with a standard low-frequency electrical
stimulation (LFS) protocol and fEPSPs were recorded during 60 min to estimate depression of the
synaptic response (LTD). 2-Way ANOVA, $F_{(1,2793)}$ = 1039, *p <0.0001 followed by Bonferroni post hoc test
(p < 0.05) compared to the WT condition. Representative traces before (gray) and after LFS are shown as
inserts. Note that HTZ hippocampal slices exhibit a significantly reduced capability to depress the
synaptic response at the end of the recording compared to WT slices (WT: n = 23 slices, 7 mice; HTZ: n =
28 slices, 8 mice). Mann-Whitney test * p <0.005 compared to the WT condition. All data are
represented as mean ± SEM.
.
791
792
793
794

795
796
Figure S5. Reduced phalloidin-reactivity in dendritic spines of HTZ neurons. 7 *DIV* hippocampal neurons,
cultured from WT and HTZ pups (P0), were fixed, stained with the F-actin binding toxin phalloidin-
rhodamin-B and visualized by confocal microscopy using the same acquisition parameters **(a)**
Representative phalloidin-stained WT and HTZ dendritic spines are shown above. Dotted yellow lines
draw the spine perimeters. Phalloidin fluorescence intensity was measured inside spines. Scale bar= 1.5
µm **(b)** Mean phalloidin-fluorescence intensity in WT (black bar) and HTZ (red bar) dendritic spines.
N=61 WT spines and 166 HTZ from 3 WT and 18 HTZ neurons. Bars are mean ± SEM *p<0.0002
compared to WT condition, Mann-Whitney 2-tail test for non-parametric data.

815
816

Figure S6. Representative dendritic spine-morphologies. Filopodia are long protrusions (> 2 um) without head. Thin-spines are short (<2 um) protrusions with small heads (<0.5 um). Stubby-spines are wide-head protrusions without neck. Mushroom spines are short protrusions (<1 um) with wide-heads (> 0.6 um width). Branched spines are short cup-shaped protrussions. Yellow arrows point dendritic spines of each morphology. Scale bar=2 um

References

Ali, Tayyibah, Joanna Bednarska, Stéphane Vassilopoulos, Martin Tran, Ivan A. Diakonov, Azza Ziyadeh-Isleem, Pascale Guicheney, et al. 2019. "Correlative SICM-FCM Reveals Changes in Morphology and Kinetics of Endocytic Pits Induced by Disease-Associated Mutations in Dynamin." *FASEB Journal* 33 (7): 8504–18. https://doi.org/10.1096/fj.201802635R.

Anggono, Victor, and Richard L. Huganir. 2012a. "Regulation of AMPA Receptor Trafficking and Synaptic Plasticity." *Current Opinion in Neurobiology*. Curr Opin Neurobiol. https://doi.org/10.1016/j.conb.2011.12.006.

2012b. "Regulation of AMPA Receptor Trafficking and Synaptic Plasticity." *Current Opinion in Neurobiology*. Curr Opin Neurobiol. https://doi.org/10.1016/j.conb.2011.12.006.

Antonny, Bruno, Christopher Burd, Pietro De Camilli, Elizabeth Chen, Oliver Daumke, Katja Faelber, Marijn Ford, et al. 2016. "Membrane Fission by Dynamin: What We Know and What We Need to Know." *The EMBO Journal*. https://doi.org/10.15252/embj.201694613.

Ardiles, Alvaro O., Carolina Flores-Muñoz, Gabriela Toro-Ayala, Ana M. Cárdenas, Adrian G. Palacios, Pablo Muñoz, Marco Fuenzalida, Juan C. Sáez, and Agustín D. Martínez. 2014. "Pannexin 1 Regulates Bidirectional Hippocampal Synaptic Plasticity in Adult Mice." *Frontiers in Cellular Neuroscience* 8 (OCT). https://doi.org/10.3389/fncel.2014.00326.

Arriagada-Diaz, Jorge, Lorena Prado-Vega, Ana M. Cárdenas Díaz, Alvaro O. Ardiles, and Arlek M. Gonzalez-Jamett. 2020. "Dynamin Superfamily at Pre- and Postsynapses: Master Regulators of Synaptic Transmission and Plasticity in Health and Disease." *Neuroscientist*. SAGE Publications Inc. https://doi.org/10.1177/1073858420974313.

Barnes, C. A. 1979. "Memory Deficits Associated with Senescence: A Neurophysiological and Behavioral Study in the Rat." *Journal of Comparative and Physiological Psychology*. https://doi.org/10.1037/h0077579.

Bayazitov, Ildar T., Robert J. Richardson, Robert G. Fricke, and Stanislav S. Zakharenko. 2007. "Slow Presynaptic and Fast Postsynaptic Components of Compound Long-Term Potentiation." *Journal of Neuroscience* 27 (43): 11510–21. https://doi.org/10.1523/JNEUROSCI.3077-07.2007.

Bhatnagar, Anushree, David L. Willins, John A. Gray, Jason Woods, Jeffrey L. Benovic, and Bryan L. Roth. 2001. "The Dynamin-Dependent, Arrestin-Independent Internalization of 5-Hydroxytryptamine 2A (5-HT2A) Serotonin Receptors Reveals Differential Sorting of Arrestins and 5-HT2A Receptors during Endocytosis." *Journal of Biological Chemistry*. https://doi.org/10.1074/jbc.M006968200.

Bhatnagar, Anushree, David L Willins, John A Gray, Jason Woods, Jeffrey L Benovic, and Bryan L Roth. 2000. "The Dynamin-Dependent, Arrestin-Independent Internalization of 5-Hydroxytryptamine 2A (5-HT 2A) Serotonin Receptors Reveals Differential Sorting of Arrestins and 5-HT 2A Receptors during Endocytosis* □ S." *THE JOURNAL OF BIOLOGICAL CHEMISTRY* 276 (11): 8269–77. https://doi.org/10.1074/jbc.M006968200.

Bitoun, Marc, Svetlana Maugenre, Pierre Yves Jeannet, Emmanuelle Lacène, Xavier Ferrer, Pascal Laforêt, Jean Jacques Martin, et al. 2005. "Mutations in Dynamin 2 Cause Dominant Centronuclear Myopathy." *Nature Genetics*. https://doi.org/10.1038/ng1657.

Bliek, A. M. Van der, T. E. Redelmeier, H. Damke, E. J. Tisdale, E. M. Meyerowitz, and S. L. Schmid. 1993. "Mutations in Human Dynamin Block an Intermediate Stage in Coated Vesicle Formation." *Journal of Cell Biology*. https://doi.org/10.1083/jcb.122.3.553.

Böhm, Johann, Valerie Biancalana, Elizabeth T. DeChene, Marc Bitoun, Christopher R. Pierson, Elise Schaefer, Hatice Karasoy, et al. 2012. "Mutation Spectrum in the Large Gtpase Dynamin 2, and Genotype-Phenotype Correlation in Autosomal Dominant Centronuclear Myopathy." *Human Mutation*. https://doi.org/10.1002/humu.22067.

Bosch, Miquel, Jorge Castro, Takeo Saneyoshi, Hitomi Matsuno, Mriganka Sur, and Yasunori Hayashi. 2014. "Structural and Molecular Remodeling of Dendritic Spine Substructures during Long-Term Potentiation." *Neuron* 82 (2): 444–59. https://doi.org/10.1016/j.neuron.2014.03.021.

Bosch, Miquel, and Yasunori Hayashi. 2012. "Structural Plasticity of Dendritic Spines." *Current Opinion in Neurobiology*. Elsevier Current Trends. https://doi.org/10.1016/j.conb.2011.09.002.

Carroll, Reed C., Eric C. Beattie, Houhui Xia, Christian Lüscher, Yoram Altschuler, Roger A. Nicoll, Robert C. Malenka, and Mark Von Zastrow. 1999. "Dynamin-Dependent Endocytosis of Ionotropic Glutamate Receptors." *Proceedings of the National Academy of Sciences of the United States of America*. https://doi.org/10.1073/pnas.96.24.14112.

Chappie, Joshua S., Sharmistha Acharya, Ya Wen Liu, Marilyn Leonard, Thomas J. Pucadyil, and Sandra L. Schmid. 2009. "An Intramolecular Signaling Element That Modulates Dynamin Function in Vitro and in Vivo." *Molecular Biology of the Cell*. https://doi.org/10.1091/mbc.E09-04-0318.

Cheung, Giselle, and Michael A. Cousin. 2019. "Synaptic Vesicle Generation from Activity-Dependent Bulk Endosomes Requires a Dephosphorylation-Dependent Dynamin–Syndapin Interaction." *Journal of Neurochemistry* 151 (5): 570–83. https://doi.org/10.1111/jnc.14862.

Chowdhury, Shoaib, Jason D. Shepherd, Hiroyuki Okuno, Gregory Lyford, Ronald S. Petralia, Niels Plath, Dietmar Kuhl, Richard L. Huganir, and Paul F. Worley. 2006. "Arc/Arg3.1 Interacts with the Endocytic Machinery to Regulate AMPA Receptor Trafficking." *Neuron* 52 (3): 445–59. https://doi.org/10.1016/j.neuron.2006.08.033.

Cingolani, Lorenzo A., Agnes Thalhammer, Lily M.Y. Yu, Myriam Catalano, Timothy Ramos, Michael A. Colicos, and Yukiko Goda. 2008. "Activity-Dependent Regulation of Synaptic AMPA Receptor Composition and Abundance by B3 Integrins." *Neuron* 58 (5): 749–62. https://doi.org/10.1016/j.neuron.2008.04.011.

Citri, Ami, and Robert C. Malenka. 2008. "Synaptic Plasticity: Multiple Forms, Functions, and Mechanisms." *Neuropsychopharmacology*. Neuropsychopharmacology. https://doi.org/10.1038/sj.npp.1301559.

Damke, Hanna, Takeshi Baba, Dale E. Warnock, and Sandra L. Schmid. 1994. "Induction of Mutant Dynamin Specifically Blocks Endocytic Coated Vesicle Formation." *Journal of Cell Biology*. https://doi.org/10.1083/jcb.127.4.915.

Durieux, Anne Cécile, Bernard Prudhon, Pascale Guicheney, and Marc Bitoun. 2010. "Dynamin 2 and Human Diseases." *Journal of Molecular Medicine*. https://doi.org/10.1007/s00109-009-0587-4.

Durieux, Anne Cécile, Stéphane Vassilopoulos, Jeanne Lainé, Bodvaël Fraysse, Laura Briñas, Bernard

Prudhon, Josiane Castells, et al. 2012. "A Centronuclear Myopathy - Dynamin 2 Mutation Impairs
Autophagy in Mice." *Traffic* 13 (6): 869–79. https://doi.org/10.1111/j.1600-0854.2012.01348.x.

Durieux, Anne Cécile, Alban Vignaud, Bernard Prudhon, Mai Thao Viou, Maud Beuvin, Stéphane
Vassilopoulos, Bodvaël Fraysse, et al. 2010. "A Centronuclear Myopathy-Dynamin 2 Mutation
Impairs Skeletal Muscle Structure and Function in Mice." *Human Molecular Genetics*.
https://doi.org/10.1093/hmg/ddq413.

Echaniz-Laguna, Andoni, Anne Sophie Nicot, Sophie Carré, Jérôme Franques, Christine Tranchant,
Nicolas Dondaine, Valérie Biancalana, Jean Louis Mandel, and Jocelyn Laporte. 2007. "Subtle
Central and Peripheral Nervous System Abnormalities in a Family with Centronuclear Myopathy
and a Novel Dynamin 2 Gene Mutation." *Neuromuscular Disorders*.
https://doi.org/10.1016/j.nmd.2007.06.467.

Ennaceur, A., and J. Delacour. 1988. "A New One-Trial Test for Neurobiological Studies of Memory in
Rats. 1: Behavioral Data." *Behavioural Brain Research* 31 (1): 47–59. https://doi.org/10.1016/0166-
4328(88)90157-X.

Fa, Mauro, Agnieszka Staniszewski, Faisal Saeed, Yitshak I. Francis, and Ottavio Arancio. 2014. "Dynamin
1 Is Required for Memory Formation." *PLoS ONE* 9 (3).
https://doi.org/10.1371/journal.pone.0091954.

Faelber, Katja, York Posor, Song Gao, Martin Held, Yvette Roske, Dennis Schulze, Volker Haucke, Frank
Noé, and Oliver Daumke. 2011. "Crystal Structure of Nucleotide-Free Dynamin." *Nature*.
https://doi.org/10.1038/nature10369.

Ferguson, Shawn M., Gabor Brasnjo, Mitsuko Hayashi, Markus Wölfel, Chiara Collesi, Silvia Giovedi,
Andrea Raimondi, et al. 2007. "A Selective Activity-Dependent Requirement for Dynamin 1 in
Synaptic Vesicle Endocytosis." *Science*. https://doi.org/10.1126/science.1140621.

Ferguson, Shawn M., and Pietro De Camilli. 2012. "Dynamin, a Membrane-Remodelling GTPase." *Nature
Reviews Molecular Cell Biology*. https://doi.org/10.1038/nrm3266.

Ferguson, Shawn, Andrea Raimondi, Summer Paradise, Hongying Shen, Kumi Mesaki, Agnes Ferguson,
Olivier Destaing, et al. 2009. "Coordinated Actions of Actin and BAR Proteins Upstream of Dynamin
at Endocytic Clathrin-Coated Pits." *Developmental Cell*.
https://doi.org/10.1016/j.devcel.2009.11.005.

Fischer, Dirk, Muriel Herasse, Marc Bitoun, Héctor M. Barragán-Campos, Jacques Chiras, Pascal Laforêt,
Michel Fardeau, Bruno Eymard, Pascale Guicheney, and Norma B. Romero. 2006. "Characterization
of the Muscle Involvement in Dynamin 2-Related Centronuclear Myopathy." *Brain*.
https://doi.org/10.1093/brain/awl071.

Flores-Muñoz, Carolina, Bárbara Gómez, Elena Mery, Paula Mujica, Ivana Gajardo, Claudio Córdova,
Daniela Lopez-Espíndola, et al. 2020. "Acute Pannexin 1 Blockade Mitigates Early Synaptic Plasticity
Defects in a Mouse Model of Alzheimer's Disease." *Frontiers in Cellular Neuroscience* 14 (March).
https://doi.org/10.3389/fncel.2020.00046.

Ford, Marijn G.J., Simon Jenni, and Jodi Nunnari. 2011. "The Crystal Structure of Dynamin." *Nature*.
https://doi.org/10.1038/nature10441.

Gajardo, Ivana, Claudia S. Salazar, Daniela Lopez-Espíndola, Carolina Estay, Carolina Flores-Muñoz,
Claudio Elgueta, Arlek M. Gonzalez-Jamett, Agustín D. Martínez, Pablo Muñoz, and Álvaro O.
Ardiles. 2018. "Lack of Pannexin 1 Alters Synaptic GluN2 Subunit Composition and Spatial Reversal
Learning in Mice." *Frontiers in Molecular Neuroscience* 11 (April).
https://doi.org/10.3389/fnmol.2018.00114.

Gipson, C. D., and M. F. Olive. 2017. "Structural and Functional Plasticity of Dendritic Spines – Root or
Result of Behavior?" *Genes, Brain and Behavior*. Blackwell Publishing Ltd.
https://doi.org/10.1111/gbb.12324.

González-Jamett, Arlek M., Ximena Baez-Matus, María José Olivares, Fernando Hinostroza, Maria José
Guerra-Fernández, Jacqueline Vasquez-Navarrete, Mai Thao Bui, et al. 2017. "Dynamin-2
Mutations Linked to Centronuclear Myopathy Impair Actin-Dependent Trafficking in Muscle Cells."
Scientific Reports. https://doi.org/10.1038/s41598-017-04418-w.

González-Jamett, Arlek M., Fanny Momboisse, María José Guerra, Stéphane Ory, Ximena Báez-Matus,
Natalia Barraza, Valerie Calco, et al. 2013. "Dynamin-2 Regulates Fusion Pore Expansion and
Quantal Release through a Mechanism That Involves Actin Dynamics in Neuroendocrine
Chromaffin Cells." *PLoS ONE*. https://doi.org/10.1371/journal.pone.0070638.

González-Jamett, Arlek M., Fanny Momboisse, Valentina Haro-Acuña, Jorge A. Bevilacqua, Pablo
Caviedes, and Ana María Cárdenas. 2013. "Dynamin-2 Function and Dysfunction along the
Secretory Pathway." *Frontiers in Endocrinology*. https://doi.org/10.3389/fendo.2013.00126.

Gu, Changkyu, Suma Yaddanapudi, Astrid Weins, Teresia Osborn, Jochen Reiser, Martin Pollak, John
Hartwig, and Sanja Sever. 2010. "Direct Dynaming-Actin Interactions Regulate the Actin
Cytoskeleton." *EMBO Journal*. https://doi.org/10.1038/emboj.2010.249.

Harris, Kristen M., and S. B. Kater. 1994. "Dendritic Spines: Cellular Specializations Imparting Both
Stability and Flexibility to Synaptic Function." *Annual Review of Neuroscience*. Annual Reviews Inc.
https://doi.org/10.1146/annurev.ne.17.030194.002013.

Hinshaw, Jenny E., and Sandra L. Schmid. 1995. "Dynamin Self-Assembles into Rings Suggesting a
Mechanism for Coated Vesicle Budding." *Nature* 374 (6518): 190–92.
https://doi.org/10.1038/374190a0.

Holtmaat, Anthony, and Karel Svoboda. 2009. "Experience-Dependent Structural Synaptic Plasticity in
the Mammalian Brain." *Nature Reviews Neuroscience*. Nat Rev Neurosci.
https://doi.org/10.1038/nrn2699.

Hotulainen, Pirta, and Casper C. Hoogenraad. 2010. "Actin in Dendritic Spines: Connecting Dynamics to
Function." *Journal of Cell Biology* 189 (4): 619–29. https://doi.org/10.1083/jcb.201003008.

Jaskolski, Frédéric, Belen Mayo-Martin, David Jane, and Jeremy M. Henley. 2009. "Dynamin-Dependent
Membrane Drift Recruits AMPA Receptors to Dendritic Spines." *Journal of Biological Chemistry*.
https://doi.org/10.1074/jbc.M808401200.

Jeannet, P. Y., G. Bassez, B. Eymard, P. Laforêt, J. A. Urtizberea, A. Rouche, P. Guicheney, M. Fardeau,
and Norma Beatriz Romero. 2004. "Clinical and Histologic Findings in Autosomal Centronuclear
Myopathy." *Neurology*. https://doi.org/10.1212/01.WNL.0000124388.67003.56.

Kabbani, Nadine, Andreas Jeromin, and Robert Levenson. 2004. "Dynamin-2 Associates with the
Dopamine Receptor Signalplex and Regulates Internalization of Activated D2 Receptors." *Cellular*
Signalling. https://doi.org/10.1016/j.cellsig.2003.09.011.

Kim, Chong Hyun, and John E. Lisman. 1999. "A Role of Actin Filament in Synaptic Transmission and
Long-Term Potentiation." *Journal of Neuroscience* 19 (11): 4314–24.
https://doi.org/10.1523/jneurosci.19-11-04314.1999.

Kong, Leopold, Kem A. Sochacki, Huaibin Wang, Shunming Fang, Bertram Canagarajah, Andrew D. Kehr,
William J. Rice, Marie Paule Strub, Justin W. Taraska, and Jenny E. Hinshaw. 2018. "Cryo-EM of the
Dynamin Polymer Assembled on Lipid Membrane." *Nature* 560 (7717): 258–62.
https://doi.org/10.1038/s41586-018-0378-6.

Koutsopoulos, Olga S., Catherine Koch, Valerie Tosch, Johann Böhm, Kathryn N. North, and Jocelyn
Laporte. 2011. "Mild Functional Differences of Dynamin 2 Mutations Associated to Centronuclear
Myopathy and Charcot-Marie-Tooth Peripheral Neuropathy." Edited by Gisela Nogales-Gadea.
PLoS ONE 6 (11): e27498. https://doi.org/10.1371/journal.pone.0027498.

Lee, Hey Kyoung, and Alfredo Kirkwood. 2011. "AMPA Receptor Regulation during Synaptic Plasticity in
Hippocampus and Neocortex." *Seminars in Cell and Developmental Biology*. Elsevier Ltd.
https://doi.org/10.1016/j.semcdb.2011.06.007.

Lin, Shan Shan, Tsung Lin Hsieh, Gunn Guang Liou, Tsai Ning Li, Hsin Chieh Lin, Chiung Wen Chang,
Hsiang Yi Wu, Chi Kuang Yao, and Ya Wen Liu. 2020. "Dynamin-2 Regulates Postsynaptic
Cytoskeleton Organization and Neuromuscular Junction Development." *Cell Reports* 33 (4).
https://doi.org/10.1016/j.celrep.2020.108310.

Liu, Ya Wen, Vasyl Lukiyanchuk, and Sandra L. Schmid. 2011. "Common Membrane Trafficking Defects of
Disease-Associated Dynamin 2 Mutations." *Traffic*. https://doi.org/10.1111/j.1600-
0854.2011.01250.x.

Loerke, Dinah, Marcel Mettlen, Defne Yarar, Khuloud Jaqaman, Henry Jaqaman, Gaudenz Danuser, and
Sandra L. Schmid. 2009. "Cargo and Dynamin Regulate Clathrin-Coated Pit Maturation." *PLoS*
Biology. https://doi.org/10. 1371/journal.pbio.1000057.

Lomash, Richa Madan, Xinglong Gu, Richard J. Youle, Wei Lu, and Katherine W. Roche. 2015.
"Neurolastin, a Dynamin Family GTPase, Regulates Excitatory Synapses and Spine Density." *Cell*
Reports 12 (5): 743–51. https://doi.org/10.1016/j.celrep.2015.06.064.

Lu, Jiuyi, Thomas D. Helton, Thomas A. Blanpied, Bence Rácz, Thomas M. Newpher, Richard J. Weinberg,
and Michael D. Ehlers. 2007. "Postsynaptic Positioning of Endocytic Zones and AMPA Receptor
Cycling by Physical Coupling of Dynamin-3 to Homer." *Neuron* 55 (6): 874–89.
https://doi.org/10.1016/j.neuron.2007.06.041.

Lu, Wei Yang, Heng Ye Man, William Ju, William S. Trimble, John F. MacDonald, and Yu Tian Wang. 2001.
"Activation of Synaptic NMDA Receptors Induces Membrane Insertion of New AMPA Receptors
and LTP in Cultured Hippocampal Neurons." *Neuron* 29 (1): 243–54.
https://doi.org/10.1016/S0896-6273(01)00194-5.

Malenka, Robert C., and Mark F. Bear. 2004. "LTP and LTD: An Embarrassment of Riches." *Neuron*.
Neuron. https://doi.org/10.1016/j.neuron.2004.09.012.

Matsuzaki, Masanori, Graham C.R. Ellis-Davies, Tomomi Nemoto, Yasushi Miyashita, Masamitsu Iino,
and Haruo Kasai. 2001. "Dendritic Spine Geometry Is Critical for AMPA Receptor Expression in
Hippocampal CA1 Pyramidal Neurons." *Nature Neuroscience* 4 (11): 1086–92.
https://doi.org/10.1038/nn736.

Matus, A. 2000. "Actin-Based Plasticity in Dendritic Spines." *Science*. Science.
https://doi.org/10.1126/science.290.5492.754.

Mooren, Olivia L., Tatyana I. Kotova, Andrew J. Moore, and Dorothy A. Schafer. 2009. "Dynamin2 GTPase
and Cortactin Remodel Actin Filaments." *Journal of Biological Chemistry*.
https://doi.org/10.1074/jbc.M109.024398.

Moro, Alessandro., van Nifterick Ann., Toonen Ruud., Verhage Matthijs. 2021. "Dynamin controls
neuropeptides secretion by organizing dense-core vesicle fusion sites"Science Advances. https://DOI:
10.1126/sciadv.abf0659

Negrón-Oyarzo, Ignacio, David Neira, Nelson Espinosa, Pablo Fuentealba, and Francisco Aboitiz. 2015.
"Prenatal Stress Produces Persistence of Remote Memory and Disrupts Functional Connectivity in
the Hippocampal-Prefrontal Cortex Axis." *Cerebral Cortex*. https://doi.org/10.1093/cercor/bhu108.

Newman-Smith, Erin D., Dixie Lee Shurland, and Alexander M. Van Der Bliek. 1997. "Assignment of the
Dynamin-1 Gene (DNM1) to Human Chromosome 9q34 by Fluorescence in Situ Hybridization and
Somatic Cell Hybrid Analysis." *Genomics*. https://doi.org/10.1006/geno.1996.4596.

Nicoziani, Paolo, Frederik Vilhardt, Alicia Llorente, Leila Hilout, Pierre J. Courtoy, Kirsten Sandvig, and Bo
Van Deurs. 2000. "Role for Dynamin in Late Endosome Dynamics and Trafficking of the Cation-
Independent Mannose 6-Phosphate Receptor." *Molecular Biology of the Cell*.
https://doi.org/10.1091/mbc.11.2.481.

Noakes, P. G., D. Chin, S. S. Kim, S. Liang, and W. D. Phillips. 1999. "Expression and Localisation of
Dynamin and Syntaxin during Neural Development and Neuromuscular Synapse Formation."
Journal of Comparative Neurology. https://doi.org/10.1002/(SICI)1096-
9861(19990809)410:4<531::AID-CNE2>3.0.CO;2-C.

Okamoto, Ken Ichi, Takeharu Nagai, Atsushi Miyawaki, and Yasunori Hayashi. 2004. "Rapid and
Persistent Modulation of Actin Dynamics Regulates Postsynaptic Reorganization Underlying
Bidirectional Plasticity." *Nature Neuroscience* 7 (10): 1104–12. https://doi.org/10.1038/nn1311.

Okamoto, Kenichi, Miquel Bosch, and Yasunori Hayashi. 2009. "The Roles of CaMKII and F-Actin in the
Structural Plasticity of Dendritic Spines: A Potential Molecular Identity of a Synaptic Tag?"
Physiology 24 (6): 357–66. https://doi.org/10.1152/physiol.00029.2009.

Okamoto, Patricia M., Chantal Gamby, David Wells, Justin Fallon, and Richard B. Vallee. 2001. "Dynamin
Isoform-Specific Interaction with the Shank/ProSAP Scaffolding Proteins of the Postsynaptic
Density and Actin Cytoskeleton." *Journal of Biological Chemistry*.
https://doi.org/10.1074/jbc.M104927200.

Pelucchi, Silvia, Ramona Stringhi, and Elena Marcello. 2020. "Dendritic Spines in Alzheimer's Disease:
How the Actin Cytoskeleton Contributes to Synaptic Failure." *International Journal of Molecular
Sciences*. MDPI AG. https://doi.org/10.3390/ijms21030908.

Penzes, Peter, and Igor Rafalovich. 2012. "Regulation of the Actin Cytoskeleton in Dendritic Spines."
Advances in Experimental Medicine and Biology 970: 81–95. https://doi.org/10.1007/978-3-7091-
0932-8_4.

Pitts, Matthew. 2018. "Barnes Maze Procedure for Spatial Learning and Memory in Mice." *BIO-
PROTOCOL*. https://doi.org/10.21769/bioprotoc.2744.

Praefcke, Gerrit J.K., and Harvey T. McMahon. 2004. "The Dynamin Superfamily: Universal Membrane
Tubulation and Fission Molecules?" *Nature Reviews Molecular Cell Biology*.
https://doi.org/10.1038/nrm1313.

Raimondi, Andrea, Shawn M. Ferguson, Xuelin Lou, Moritz Armbruster, Summer Paradise, Silvia Giovedi,
Mirko Messa, et al. 2011. "Overlapping Role of Dynamin Isoforms in Synaptic Vesicle Endocytosis."
Neuron. https://doi.org/10.1016/j.neuron.2011.04.031.

Romero, Norma B., and Marc Bitoun. 2011. "Centronuclear Myopathies." *Seminars in Pediatric
Neurology* 18 (4): 250–56. https://doi.org/10.1016/j.spen.2011.10.006.

Schiffer, Mario, Beina Teng, Changkyu Gu, Valentina A. Shchedrina, Marina Kasaikina, Vincent A. Pham,
Nils Hanke, et al. 2015. "Pharmacological Targeting of Actin-Dependent Dynamin Oligomerization
Ameliorates Chronic Kidney Disease in Diverse Animal Models." *Nature Medicine* 21 (6): 601–9.
https://doi.org/10.1038/nm.3843.

Shin, Wonchul, Lihao Ge, Gianvito Arpino, Seth A. Villarreal, Edaeni Hamid, Huisheng Liu, Wei Dong
Zhao, Peter J. Wen, Hsueh Cheng Chiang, and Ling Gang Wu. 2018. "Visualization of Membrane
Pore in Live Cells Reveals a Dynamic-Pore Theory Governing Fusion and Endocytosis." *Cell* 173 (4):
934-945.e12. https://doi.org/10.1016/j.cell.2018.02.062.

Shors, Tracey J., and Louis D. Matzel. 1997. "Long-Term Potentiation: What's Learning Got to Do with
It?" *Behavioral and Brain Sciences*. Behav Brain Sci. https://doi.org/10.1017/S0140525X97001593.

Singh, Mahaveer, Hemant R. Jadhav, and Tanya Bhatt. 2017. "Dynamin Functions and Ligands: Classical
Mechanisms Behind." *Molecular Pharmacology*. https://doi.org/10.1124/mol.116.105064.

Stefen, Holly, Chanchanok Chaichim, John Power, and Thomas Fath. 2016. "Regulation of the
Postsynaptic Compartment of Excitatory Synapses by the Actin Cytoskeleton in Health and Its
Disruption in Disease." *Neural Plasticity* 2016. https://doi.org/10.1155/2016/2371970.

Sukoff Rizzo, Stacey J., and Jacqueline N. Crawley. 2017. "Behavioral Phenotyping Assays for Genetic
Mouse Models of Neurodevelopmental, Neurodegenerative, and Psychiatric Disorders." *Annual
Review of Animal Biosciences*. https://doi.org/10.1146/annurev-animal-022516-022754.

Tanabe, Kenji, and Kohji Takei. 2009. "Dynamic Instability of Microtubules Requires Dynamin 2 and Is
Impaired in a Charcot-Marie-Tooth Mutant." *Journal of Cell Biology*.
https://doi.org/10.1083/jcb.200803153.

Tanifuji, Shota, Megumi Funakoshi-Tago, Fumihito Uedas, Tadashi Kasaharas, and Sumiko Mochida.
2013. "Dynamin Isoforms Decode Action Potential Firing for Synaptic Vesicle Recycling." *Journal of
Biological Chemistry*. https://doi.org/10.1074/jbc.M112.445874.

Wang, Weisheng, Yun Yue Ju, Qi Xin Zhou, Jian Xin Tang, Meng Li, Lei Zhang, Shuo Kang, et al. 2017. "The

Small GTPase Rac1 Contributes to Extinction of Aversive Memories of Drug Withdrawal by
Facilitating GABAa Receptor Endocytosis in the VmPFC." *Journal of Neuroscience*.
https://doi.org/10.1523/JNEUROSCI.3859-16.2017.

Warnock, Dale E., Takeshi Baba, and Sandra L. Schmid. 1997. "Ubiquitously Expressed Dynamin-II Has a
Higher Intrinsic GTPase Activity and a Greater Propensity for Self-Assembly than Neuronal
Dynamin-I." *Molecular Biology of the Cell*. https://doi.org/10.1091/mbc.8.12.2553.

Watanabe, Shigeki, Benjamin R. Rost, Marcial Camacho-Pérez, M. Wayne Davis, Berit Söhl-Kielczynski,
Christian Rosenmund, and Erik M. Jorgensen. 2013. "Ultrafast Endocytosis at Mouse Hippocampal
Synapses." *Nature* 504 (7479): 242–47. https://doi.org/10.1038/nature12809.

Yamada, Hiroshi, Tadashi Abe, Ayano Satoh, Nana Okazaki, Shota Tago, Kinue Kobayashi, Yumi Yoshida,
et al. 2013. "Stabilization of Actin Bundles by a Dynamin 1/Cortactin Ring Complex Is Necessary for
Growth Cone Filopodia." *Journal of Neuroscience* 33 (10): 4514–26.
https://doi.org/10.1523/JNEUROSCI.2762-12.2013.

Zhang, Ruihui, Donghoon M. Lee, John R. Jimah, Nathalie Gerassimov, Changsong Yang, Sangjoon Kim,
Delgermaa Luvsanjav, et al. 2020. "Dynamin Regulates the Dynamics and Mechanical Strength of
the Actin Cytoskeleton as a Multifilament Actin-Bundling Protein." *Nature Cell Biology* 22 (6): 674–
88. https://doi.org/10.1038/s41556-020-0519-7.

Zhang, Xiao Yan, Fang Ji, Ning Wang, Lin Lin Chen, Tian Tian, and Wei Lu. 2014. "Glycine Induces
Bidirectional Modifications in N-Methyl-D-Aspartate Receptor-Mediated Synaptic Responses in
Hippocampal CA1 Neurons." *Journal of Biological Chemistry* 289 (45): 31200–211.
https://doi.org/10.1074/jbc.M114.570630.

Zheng, Ning, Okunola Jeyifous, Charlotte Munro, Johanna M. Montgomery, and William N. Green. 2015.
"Synaptic Activity Regulates AMPA Receptor Trafficking through Different Recycling Pathways."
ELife 4 (MAY). https://doi.org/10.7554/eLife.06878.

Züchner, Stephan, Maher Noureddine, Marina Kennerson, Kristien Verhoeven, Kristl Claeys, Peter De
Jonghe, John Merory, et al. 2005. "Mutations in the Pleckstrin Homology Domain of Dynamin 2
Cause Dominant Intermediate Charcot-Marie-Tooth Disease." *Nature Genetics*.
https://doi.org/10.1038/ng1514.

Zucker, Robert S., and Wade G. Regehr. 2002. "Short-Term Synaptic Plasticity." *Annual Review of
Physiology*. Annu Rev Physiol. https://doi.org/10.1146/annurev.physiol.64.092501.114547.

1141

1142

1143

1144

1145

1146

A

B