THERAPEUTIC EFFECTS OF HYPOXIC AND PRO-INFLAMMATORY PRIMING OF

MESENCHYMAL STEM CELL-DERIVED EXTRACELLULAR VESICLES IN

INFLAMMATORY ARTHRITIS

Alasdair G. Kay[1], Kane Treadwell[2], Paul Roach[3], Rebecca Morgan[2], Rhys Lodge[4], Mairead

Hyland[2], Anna M.Piccinini[5], Nicholas Forsyth[6], *Oksana Kehoe[2]

[1]Department of Biology, University of York, York, UK; [2]School of Medicine, Keele University

at the RJAH Orthopaedic Hospital, Oswestry, UK; [3]Chemistry Department, Loughborough

University, Loughborough, UK; [4]School of Chemistry, University of Nottingham, Nottingham,

UK; [5]School of Pharmacy, University of Nottingham, Nottingham, UK; [6]School of Pharmacy

and Bioengineering, Keele University, The Guy Hilton Research Laboratories, Hartshill, Stoke

on Trent, UK

*Correspondence and requests for materials should be addressed to O.K. (email:

o.kehoe@keele.ac.uk)

Abstract

Novel biological therapies have revolutionised the management of Rheumatoid Arthritis (RA)

but no cure currently exists. Mesenchymal stem cells (MSCs) immunomodulate inflammatory

responses through paracrine signalling, including via secretion of extracellular vesicles (EVs)

in the cell secretome. We evaluated the therapeutic potential of MSCs-derived small EVs in

an antigen-induced model of arthritis (AIA).

EVs isolated from MSCs cultured normoxically (21% O_2, 5% CO_2), hypoxically (2% O_2, 5% CO_2)

or with a pro-inflammatory cytokine cocktail were applied into the AIA model. Disease

pathology was assessed post-arthritis induction through swelling and histopathological

analysis of synovial joint structure. Activated CD4+ T cells from healthy mice were cultured

with EVs or MSCs to assess deactivation capabilities prior to application of standard EVs in

vivo to assess T cell polarisation within the immune response to AIA.

All EVs treatments reduced knee-joint swelling whilst only normoxic and pro-inflammatory

primed EVs improved histopathological outcomes. In vitro culture with EVs did not achieve T

cell deactivation. Polarisation towards CD4+ helper cells expressing IL17a (Th17) was reduced

when normoxic and hypoxic EV treatments were applied in vitro. Normoxic EVs applied into

the AIA model reduced Th17 polarisation and improved Th17:Treg homeostatic balance.

Priming of MSCs in EV production can be applied to alter the therapeutic efficacy however

normoxic EVs present the optimal strategy for broad therapeutic benefit. The varied

outcomes observed in MSCs priming may promote EVs optimised for therapies targeted for

specific therapeutic priorities. EVs present an effective novel technology with potential for

cell-free therapeutic translation.

Keywords

Rheumatoid Arthritis, inflammation, immunomodulation, extracellular vesicles, mesenchymal stem cells

Introduction

Mesenchymal stem cells (MSCs) are a promising therapeutic option owing potential for tissue repair through trilineage differentiation capacity, immunomodulatory properties disrupting T cell proliferation, B cell function and dendritic cell (DCs) maturation; and promoting anti-inflammatory responses mediated through macrophage interactions[1]. Widespread introduction of stem cell therapies has been hindered by inconsistent outcomes at clinical trial and donor variability. Our group has demonstrated the immunomodulatory capacity of both MSCs and their conditioned medium (CM-MSC) to reduce inflammation in a murine antigen-induced arthritis (AIA) model through enhanced Treg function and restored Treg:Th17 ratio[2,3]. MSCs convey their immunomodulatory properties through cell-to-cell contact, autocrine responses and paracrine signalling[1], including through secretion of . extracellular vesicles (EVs)[4]. EVs are membrane bound particles that carry a cargo of microRNA (miRNA), mRNA, lipid, carbohydrate and protein signals to facilitate intercellular communication[5-9]. Notably, disease severity in seropositive RA has been linked to EVs signalling[10-15]. MSC-derived EVs have been shown to be beneficial in autoimmune disorders in functioning to modulate autoimmune responses, particularly related to graft rejection and hypertension[16-18] but also in inflammatory arthritis and rheumatic diseases[19-21]. T cells have been shown to recruit EVs released by DCs, suggesting an integral action in orchestrating *in*

vivo immune responses[22,23]. Furthermore, tolerance in DCs is influenced by CD4+ T cells

through EVs-mediated intercellular signalling mediated by transfer of EVs-packaged miRNA[24].

We hypothesised that EVs recapitulate the anti-inflammatory and immunomodulatory effects

of MSCs in the arthritis model and that priming MSCs during the generation of EVs through

hypoxic culture or pro-inflammatory cytokine cocktails will enhance therapeutic outcomes.

Identification of the most effective cell source and maintenance conditions for generation of

optimally efficacious EVs will accelerate development of EV therapeutics for RA and other

regenerative medicine targets. To date, no research has directly examined and contrasted the

influence of cell isolation and priming on EVs generation using both hypoxia and pro-

inflammatory pre-conditioning of MSCs during production.

Here we present novel data utilising priming strategies for MSC-derived EVs applied to the

AIA model of inflammatory arthritis. We investigate amelioration of symptoms through

reduced swelling and histopathological improvement, and EVs influence on T cell polarisation

in vitro and in vivo. We hypothesise that EVs represent a potential therapeutic approach for

the treatment of inflammatory arthritis that may encounter less obstacles than cell therapy

to widespread application in the clinic.

Results

EVs isolated through differential ultracentrifugation characterise as EVs meeting ISEV criteria

All EVs isolated in this study were enriched using 100,000 x g ultracentrifugation with EV

pellets resuspended based on counts of source cells taken at the point of collection, with

resuspension at vesicles from 1.0×10^6 MSCs per 30µl.

A combination of quantitative and qualitative analyses was performed to confirm isolation of

EVs. EV preparations from MSCs cultured under normoxic conditions were assessed by flow

cytometry analysis using the MACSPlex exosome kit (Miltenyi Biotec) to identify 37 exosomal

surface epitopes, including characteristic exosome markers CD9, CD63 and CD81. EV

isolations showed elevated enrichment in mean expression values for CD9 (81.25±5.03%),

CD63 (94.59±2.23%) and CD81 (79.41±9.07%) (n=11) (representative example Figure S1A)

confirming the identity as EVs. Western blot analysis of MSC lysate and EVs (EV-NormO2) for

detection of Alix, a protein involved in MVB biogenesis, and cytochrome C, an ubiquitous

mitochondrial protein acting as negative control, showed that isolated MSC-EVs are highly

enriched in Alix, which is expressed by their cells of origin, but do not contain cytochrome C

as expected (Figure S1B). TEM imaging demonstrated the presence of spherical vesicles in

isolated preparations in accordance with ISEV criteria for single EVs characterisation (Figure

S1C). Finally, EVs were characterised for their size distribution and concentration using the

Nanopore technology (Izon Science). EVs preparations showed a distribution of EV sizes with

most prevalent diameter of ~ 200 nm with maximal diameter ~500 nm (Figure S1D-E).

Together, these results demonstrate that our differential ultracentrifugation methodology

successfully isolates EVs meeting ISEV criteria. We observed that pro-inflammatory priming

of MSCs using the methodology adopted here was possible for a period of 48 hours for

collection of CM-MSC, but extended cell culture in the presence of pro-inflammatory cocktail

to 72 or 96 hours resulted in consistent reductions in cell numbers below the originally seeded

cell numbers, suggestive of induced crisis in the cultured cells (data not shown).

Priming of MSCs does not affect total protein content in EV cargo

We set out to determine whether the total protein content in the cargo of EVs was impacted

by cell priming. A 4 parameter polynomial nonlinear regression of log-transformed BCA assay

data revealed no significant increase in total protein between conditions EV-2%O2

preparations (81.29±34.82 pg/1.0 x 10^6 cells; n=8) and EV-Pro-Inflam (69.09±37.38 pg/1.0 x

10^6 cells; n=9) compared to EV-NormO2 (40.43±14.73 pg/1.0 x 10^6 cells; n=11) (p>0.05, data

not shown).

Application of MSC-derived EVs ameliorates histopathology and joint swelling in AIA

To assess the therapeutic potential of MSC-derived EVs we introduced each of our primed

MSCs EVs treatments into a murine model of inflammatory arthritis and measured effects of

treatments on histopathological outcomes. AIA is an acute model of inflammatory arthritis

that typically exhibits peak joint swelling at 24 hours post induction with clinical symptoms

and histopathological signs that resemble rheumatoid arthritis. We have previously

demonstrated the efficacy of CM-MSC in ameliorating swelling in AIA[3]. In this study, we

compared three EV treatments to PBS only vehicle controls, measuring the reduction of joint

diameter from peak swelling (day 1). Local administration of EVs into joints significantly

reduced joint diameter post-arthritis induction, a quantitative measure proportional to joint

swelling. Specifically, EV-NormO2 (day 2 = 4.4±0.6mm, day 3 = 8.0±0.6mm, p<0.01), EV-2%O2

(day 2 = 7.7±0.8mm, day 3 = 11.0±0.9mm, p<0.001) and EV-Pro-Inflam (day 2 = 6.7±1.0mm,

day 3 = 11.3±0.9mm, p<0.001) all very significantly reduce joint swelling in comparison to PBS

vehicle control treatments (day 2 = 0.2±0.8mm, day 3 = 2.8±1.0mm) (Figure 1A). Reductions

in joint diameter from peak swelling were proportional for all treatment conditions.

Histopathological symptoms include immune cell infiltration into the synovium; hyperplasia

of the synovial membrane; and extravasation of leukocytes into the synovial joint cavity. We

have previously shown that the intra-articular injection of MSC conditioned medium in mice

with AIA ameliorates damage as observed through histopathological analysis[3]. Whilst EV-

2%O2 overall showed a tendency to reduce total arthritis index (AI) scores (4.92±0.84)

compared to controls, this was not statistically significant. For EV-NormO2, this reduced AI

score reflected decreases from control scores in hyperplasia of the synovial membrane

(1.35±0.22 vs 2.30±0.17) and joint exudate (0.50±0.21 vs 1.68±0.30), whilst EV-Pro-Inflam

demonstrated reduced synovial infiltrate (2.0±0.39 vs 3.49±0.24) (Figure 1B). Overall AI

showed significant reductions in mice treated with EV-NormO2 (4.20±0.79) and EV-Pro-Inflam

(4.08±1.04) compared to EV-depleted medium control (7.46±0.59) (Figure 1C). Taken

together, these results demonstrate EV-NormO2 treatment produced a broad reduction in

swelling and improvement in histological outcomes, whilst primed MSCs were effective in

reducing swelling, but evidence suggested this was not exclusively through reductions in

synovial infiltrate or reduced synovial damage.

Figure 1 – MSC-derived EVs treatment of mice with AIA. (A) Alleviation of joint swelling as a measure of therapeutic efficacy following EV treatments shows a significant effect of EVs compared to vehicle controls at day 2 and day 3 after arthritis induction; normalised to 0 at peak swelling (day 1) (2 Way ANOVA with Repeated Measures and Bonferroni multiple comparisons test post hoc) (B) Examination of histological signs of arthritis pathogenesis following EV treatment shows significant therapeutic effects of EVs sourced from MSCs cultured under normoxia and MSCs cultured in the presence of pro-inflammatory cytokines 3 days post-induction. (C) Combined score arthritis index shows significant reductions in arthritis index 3 days after induction when treated with EVs (1 way ANOVA with Repeated Measures and Bonferroni Multiple Comparisons test post hoc); (D) Representative images for synovial infiltrate, hyperplasia of the synovial lining and synovial exudate into joint cavity for vehicle control and EVs treatments (Control n=21, EV-NormO2 n=8, EV-2%O2 n=6, EV-Pro-Inflam n=6. *p<0.05; **p<0.01; ***p<0.001)

EV-Pro-Inflam treatments showed reductions in synovial infiltrate sufficient to affect a significantly improved overall arthritis index, however this treatment methodology was not as effective at reducing synovial exudate or hyperplasia of the synovial membrane (Figure 1D). The reduction in synovial infiltrate may therefore contribute towards reduced swelling

however the results seen in EV-2%O2 suggests this is not a single factor underpinning the

mechanism of action.

EV treatments do not affect expression of pro- (TNF-α) and anti-inflammatory (IL-10) cytokines

detectable in serum of AIA mice

The pro-inflammatory cytokine TNF-α is a key driver of disease pathogenesis in RA and a

therapeutic target in biological treatments[25,26]. Conversely, IL-10 is a master regulator of anti-

inflammatory immune responses[6]. To investigate the effects on circulating cytokines, TNF-α

and IL-10 were measured at day 3 in the serum of mice following EV treatments. The

circulating concentration of TNF-α in serum of treated mice did not vary significantly between

controls (7.02±0.47pg/mL, n=15) and EV treated conditions or between treatments EV-

NormO2 (7.38±1.04pg/mL), EV-2%O2 (6.93±0.65pg/mL) or EV-Pro-Inflam (5.73±0.47pg/mL)

(p>0.05, n=5 per treatment). ELISA on mouse serum could not detect IL-10 in treatment or

control conditions (data not shown). Owing to the small volume of synovial fluid in the murine

articular cavity, it was not possible to isolate sufficient synovial fluid to accurately detect

cytokine expression within the localised joint cavity. The consistent concentrations of

circulating TNF-α in serum of mice undergoing treatment and in untreated control mice

suggests that immunomodulation of TNF-α regulated cytokine/chemokine induction is not a

primary mechanism underpinning efficacy of EVs treatments in reducing swelling and/or

histological improvement. This is in contrast to results previously observed in MSC

treatments[2].

EVs with hypoxic priming or unprimed decrease Th17 polarization but only MSCs suppress

proliferation when co-cultured with activated CD4+ T cells in vitro

Prior to introduction in vivo, we investigated the in vitro effect of MSC-derived EVs on T cell

polarisation and proliferation to establish an alternative mechanistic rationale for efficacies

observed in swelling and histological scores for EVs treatments. MSCs have been shown to

deactivate T cells in co-culture (2,3). MSCs and EVs produced following conditions EV-

NormO2, EV-2%O2 and EV-Pro-Inflam were therefore co-cultured for 5 days with activated T

cells isolated from healthy mice and polarisation and proliferation of activated cells assessed

as a measure of immune response and T cell deactivation.

T cell polarisation was assessed by flow cytometry analysis of surface CD4 and intracellular

markers characteristic of Th1 (IFN-γ), Th2 (IL-4) and Th17 (IL-17a). The ability of EV treatments

to affect deactivation of T cells, and therefore influence T cell proliferation and extent of

immune response, was examined to determine experimental conditions for assessment of EV

treatment interactions with T cells in vivo.

Replicating previous studies[2,3], MSC/CD4+ T cell co-cultures prompted significantly increased

numbers of CD4+ T cells compared to activated T cells cultured alone. MSCs also elicited

increased CD4+ T cells in comparison to all EV treatment conditions (n=3, p<0.05) (Figure 2A).

Further analysis of IL17a expressing cells within the CD4+ population demonstrated that MSC

co-cultures increased cell proportions in comparison to PBS controls and all EV conditions,

whereas all EV treatments showed a decrease in Th17 cell polarisation (p>0.05, n=3) (Figure

2B). Whilst no differences were seen for IFN-γ expressing cells (Th1) (Figure 2C), IL-4

expressing cells (Th2) were elevated in MSC co-cultures (6.00±0.24%) compared to EV-Pro-

Inflam (2.74±0.52%) (p<0.01) but not in comparison PBS control (4.79±0.99%); EV-NormO2

(2.86±0.30%) or EV-2%O2 (3.22±0.03%) (p>0.05, n=3) (Figure 2D). Statistical analysis was

performed using log transformed data.

Additionally, the expression of IL17a per cell, represented by mean fluorescence intensity

(MFI) of signal, was significantly reduced in EV-NormO2 (3311.74±33.71) and EV-2%O2/CD4+

T cell co-cultures (3367.19±39.92) compared to CD4+ T cells cultured alone (3728.69±50.28)

(p<0.01, p<0.05, n=3). The impact of this change would need to be assessed in vivo as this had

not translated to reduced Th17 numbers in vitro.

These results demonstrate that *in vitro* MSC-derived EVs do not prompt increases in CD4+ T

cell numbers or Th17 effector cell polarisation as seen with MSC treatments. Interpretation

of this data is limited by the lack of Treg data preventing the examination of the Treg:Th17

ratio which is offset in RA.

We have previously shown the ability of MSCs to suppress T cell proliferation in co-culture,

with MSCs being more effective than CM-MSC[2,3]. MSC/CD4+ T cell co-culture reduced the

proportion of proliferating T cells, measured via a proliferative index[27] and the number of

proliferative cycles[28] undergone by T cells in co-culture (5.31±0.38, 4.63±0.17 respectively) in

comparison to T cells cultured alone (6.16±0.10, 5.44±0.25) (n=8, p<0.05). In contrast, CD4+

T cells from healthy (not AIA induced) mice showed no significant inhibition of proliferative

cycles when cultured with EV-NormO2 (5.85±0.08, 5.01±0.17), EV-2%O2 (6.01±0.18,

4.97±0.14) or EV-Pro-Inflam (6.28±0.12, 4.95±0.09) (n=8, p>0.05) (Figure 2E).

Figure 2 – Outcomes of EV treatments co-cultured with T cells isolated from healthy murine spleens **(A)** Increased CD4+ T cells in MSC co-cultures compared to EV-2%O2; EV-Pro-Inflam; and T cells alone control **(B)** Increased pro-inflammatory Th17 cells (IL17a+) in MSC co-cultures over to EV-NormO2; EV-2%O2; EV-Pro-Inflam; and T cells alone control with EV-NormO2 significantly reduced on PBS control and EV-Pro-Inflamm also **(C)** EV or MSC treatments did not alter Th1 polarisation **(D)** EV treatments all reduced Th2 polarisation in comparison to MSC treatment, with EV-NormO2 and EV-Pro-Inflam also reduced in comparison to PBS controls (Index of proliferation only) (n=3; *p<0.05; **p<0.01; ***p<0.001; 1-Way ANOVA with Bonferroni post-hoc of log transformed data. **(E)** MSC co-cultures significantly inhibited T cell proliferation compared with T cells cultured alone (both measures) and EV-Pro-Inflam co-culture (Index of proliferation only) (n=3; *p<0.05; **p<0.01; ***p<0.001; 1-Way ANOVA with Bonferroni post-hoc of log transformed data.

EV treatments showed greater impact on proliferative cycles than on the proliferative index,

though treatments did not significantly vary from either MSC co-cultures or T cells cultured

alone, with the exception of pro-inflammatory primed EVs which allowed a higher

proliferative index in comparison to MSC co-culture (n=8, p<0.05) (Figure 2E). Previous studies

have shown similar results through the application of whole secretome[3]. The results found

here suggest that direct deactivation of T cells and immunomodulated T helper polarisation

does not provide a primary mechanism for therapeutic action of EVs in vivo. Whilst MSCs do

display direct immunomodulation of T cells, it may be that MSC-secreted EVs require an

intermediate such as DCs to affect a similar influence on T cell responses.

MSC-derived EVs reduce Th17 polarisation and restore the Th17:Treg balance ex vivo

Following on from our in vitro study of T cell deactivation using CD4+ T cells of healthy mice,

we hypothesised that direct T cell interactions were not likely to be the mechanism through

which therapeutic effects are observed following EVs infusion. EV-Pro-Inflam treatments did

not influence T cell polarisation however hypoxic priming of MSCs generated EVs capable of

influencing Th2 polarisation and Th17 expression (observed through increased mean

fluorescence intensity, i.e. per cell expression). At this stage, data suggested MSC priming

(either 2%O2 or pro-inflammatory cocktail) conveys no beneficial advantage over

normoxically produced MSC-EVs in modulating T cell responses or in reducing swelling and

affecting improved histological outcomes. Our previous study demonstrated the ability of

whole secretome conditioned medium from normoxically grown MSCs to influence T cell

polarisation, restoring Treg:Th17 homeostasis, and to affect reductions in swelling and tissue

damage in vivo, similarly not via T cell deactivation when investigated in vitro[3]. We again

hypothesis the need for a mediating cell to affect T cell deactivation. To investigate the

fraction of the secretome responsible for in vivo immunomodulatory outcomes observed with

CM-MSC, and to reinforce the principles of 3R's, we determined to apply the AIA model to

examine T cell interactions in only EV-NormO2 vesicles comparative to EV depleted

secretome. Here we aim to elucidate whether EVs or the non-vesicular secretome affects T

helper cells deactivation. Spleens and lymph nodes (inguinal and popliteal) of EV-NormO2 and

EV-depleted CM-MSC treated AIA mice and PBS controls were dissociated, and CD4+ T cells

isolated. These cells represent *in vivo* primed T cells within the inflammatory arthritis

environment prior to EVs treatment, isolation and subsequent *in vitro* activation.

EV-NormO2 did not increase the proportion of CD4+ T cells in spleen (13.28±0.51%) or lymph

nodes (15.04±1.04%) over PBS controls in spleen (14.21±1.07%) or lymph nodes

(15.52±1.15%) (p>0.05, n=4). However, EV-depleted CM-MSC treatment resulted in a small

but statistically significant increase in the proportion of CD4+ T cells in spleens (16.94±0.81%)

(p<0.05, n=4) compared to EV-NormO2 treatment, but not in lymph nodes (18.83±1.13%)

(p>0.05, n=4) (Figure S2A, S2B shown with log transformed data).

Examination of *in vivo* primed CD4+ T cells activated and cultured *in vitro* for 4 hours in the

presence of a membrane transport blocker facilitated evaluation of T cell polarisation. These

results demonstrated MSC polarisation of T cells favoured Th17 outcomes as shown

previously[2,3]. RA (and other autoimmune disease) sufferers experience an imbalanced

Th17:Treg ratio leading to inappropriate immune responses and tissue damage[29-32] so for the

in vivo analysis we additionally examined regulatory T cells (Tregs, CD25+FOXP3+) polarisation

to allow calculation of ratios of Th17:Treg and Th1:Th2 to highlight differential changes in

polarisation of subsets of helper T cell that could affect the overall homeostatic balance.

When compared to PBS control (3.23±0.81%), spleen T cell polarisation towards Th17 effector

cells following both EV-NormO2 (0.83± 0.10%) and EV-depleted CM-MSC (0.90±0.19%)

treatments showed significant decreases in the proportion of pro-inflammatory Th17 cells

induced in AIA mice (p<0.05, n=4), with no significant change in FOXP3 (Treg) proportions

(Figure 3A). This translated to a trend for improved Treg:Th17 ratios in spleens of EV-NormO2

(12.06±2.12:1) and EV-depleted CM-MSC (12.70±2.95:1) treated mice compared to PBS

controls (5.03±1.40:1) (p=0.12, p=0.09 respectively, n=4) (Figure 3B). This result indicates that

EVs can prompt an advantageous shift in the Treg:Th17 balance towards a healthy state,

demonstrating a similar efficacy of applying EVs alone to that previously seen when applying

CM-MSC[3]. A similar trend was seen in T cells from lymph nodes with a significant reduction in

Th17 polarisation in EV (0.90±0.09%) and EV-depleted CM-MSC (0.93±0.25%) compared with

PBS controls (4.26±1.12%, p<0.05, n=4) (Figure 3C) leading to improved Treg:Th17 ratio

(Figure 3D). Taken together, these data show that EV treatments and EV-depleted medium

were both capable of affecting alleviation of symptoms of inflammatory arthritis through

reduced Th17 polarisation and shift in the Treg:Th17 balance. We also observed an increase

in pro-inflammatory Th1 cells in spleens of EV-NormO2 treated mice (6.38±0.40%) compared

to PBS controls (4.00±0.35%, p<0.05, n=4) (Figure S2C).

The outcomes of treatment using EVs *in vivo* demonstrate a reduction in polarisation towards

IL17a secreting T helper cells for both spleen and lymph nodes of AIA treated mice that is not

seen *in vitro* using EV-treated activated T cells from healthy mice that have not experienced

AIA induction. We therefore hypothesise that the therapeutic effect seen in vitro may require

the presence of a mediator and we propose the candidate for this would be DCs. Regulatory

T cells influence DCs through EVs mediated intercellular communication to provide

immunomodulation, including regulation of tolerance in autoimmune conditions[24].

Potentially the variation observed through *in vivo* treatments compared with *in vitro*

responses reflect the absence of mediating cell types such as DCs.

Figure 3 – Outcomes of intracellular staining for T cell polarisation analysis. Intracellular staining
for FACS analysis of IFN-γ, IL4 and IL17a in CD4+ T cells from EV-NormO2, EV-depleted CM-MSC or
PBS control. Significant reductions in (A/C) IL17a expression suggestive of reduced Th17
polarisation in Spleen and Lymph Node with no change in regulatory T cell polarisation resulted in
(B/D) trend for restoration of the Treg:Th17 balance in EV-NormO2 treated mice which was found
significant in lymph node cells only (n=4, *p<0.05; **p<0.01; ***p<0.001)(1 Way ANOVA with
Bonferroni Multiple Comparisons Test post hoc using log transformed data.

Discussion

This study builds on the growing knowledge for the therapeutic efficacy of EVs as a treatment

for inflammatory autoimmune disorders. We aim to inform researchers and clinicians on the

efficacy of priming strategies to prepare EVs as an enhanced therapy. In our study, EV

treatments prompted amelioration of clinical symptoms of AIA with reduction of joint

swelling both with and without priming. Examination of joint sections showed improved

histological features following standard and pro-inflammatory primed MSC-derived EV

treatments compared to controls but not when using hypoxically derived EVs.

Mechanistically, EVs acted remarkably different from their cells of origin (MSCs) when co-

cultured with CD4+ T cells from healthy murine spleens. Standard and hypoxically-derived EVs

reduced Th17 polarization without affecting T cell proliferation, while MSCs increased Th17

polarization, and lessened T cell proliferation. Our previous study indicated that MSCs do

indeed increase Th17 polarisation but this is offset by increases in Treg cells to restore a more

homeostatic balance between the two. Here we examined the Treg:Th17 ratio following AIA

initiation and treatment with EV-NormO2. With lack of significant improvements in hypoxic

priming outcomes, we opted to test only EV-NormO2 and EV-depleted CM to compare MSC-

CM fractions for influence on T cell polarisation following treatment application in the pro-

inflammatory environment. *Ex vivo*, CD4+ T cells isolated from spleens of arthritic mice

treated with standard EV-NormO2 again showed significantly reduced Th17 polarization that

rebalanced the Treg:Th17 ratio. Together, this suggests the reduction in Th17 cells led to

restoration of the Treg:Th17 ratio, which is typically unbalanced in inflammatory arthritis, and

reduced immune cell recruitment.

To further dissect the therapeutic mechanism of action of MSC-derived EVs, circulating levels

of TNF-α, which is a key driver of pathogenesis in RA and therapeutic target in biological

treatments, and IL-10, which is a master regulator of anti-inflammatory immune responses,

were measured in mice with AIA at day 3. While IL-10 was not detected in our assay, similarly

low levels of TNF-α were detected in serum of untreated and EV-treated mice. However,

whilst this suggests that TNF-α blockade and IL-10 modulation are not the mechanisms by

which MSC-derived EVs improve AIA, it is noted that circulating serum levels of TNF-α rise

rapidly in 24 hours post-induction then fall rapidly in the AIA model, so detectable serum TNF-

α may not be significantly different in AIA and control mice at 3 days post-induction[33,34]. This

mechanism may be more suited to assessment following prophylactic treatment using EVs

administered at the point of induction rather than after onset of symptoms.

Our previous study demonstrated reductions in Th17 cells following CM-MSC treatment

compared to control and MSC treatments. Whilst EVs have been shown to increase Il-10 in

macrophages[35], this study indicates that the presence of MSCs, and not just their secretome,

may be driving increased IL-10 expression influencing T cell polarisation in vivo, and were

responsible for the reduction in Th1 and increase in Th2 cells seen in our previous study[3]. EVs

are capable of significantly reducing Th17 cell numbers however and this represents a

significant finding in the search for immunomodulatory therapeutics for treating autoimmune

disorders where an imbalance in T cell polarisation (Th1:Th2 or Treg:Th17) is integral in

disease pathology. Moreover, EV treatment in this study resulted in an improved (2.40-fold

over controls) Treg:Th17 ratio compared to our previous results using whole CM-MSC

treatment (2.13-fold over control) or MSC treatment (1.47-fold over control)[3].

The proportions of Treg and Th17 cells in RA sufferers has been directly linked to the severity

of disease, and restoring the Treg:Th17 balance has the potential to promote homeostasis

and positive clinical outcomes. Here, we demonstrate that untreated (PBS control) spleens of

mice with AIA display a Treg:Th17 ratio of 5:1 and that this is improved to 12:1 upon EV-

NormO2 and control EV-depleted CM-MSC treatments. This result reinforces the use of CM-

MSC or vesicular components as a therapeutic option for RA treatment.

We observed effects on T cell polarisation that were consistent between in vitro and ex vivo

outcomes, with the exception of reduced Th2 cells in vitro unmatched ex vivo.

It has been suggested that the immunomodulatory properties of EVs are contingent on pro-

inflammatory priming[19,36]. However, the present work demonstrates that whilst priming

strategies may enhance some therapeutic outcomes, the mechanisms are unclear and the

overall immunomodulatory property may be inherent to MSC-EVs regardless of priming

methodology. EV-Pro-Inflam were the least effective treatment methodology for T cell

deactivation however did reduce swelling and immune cell recruitment to the synovium. In

contrast, hypoxically primed MSCs produced EVs that reduced swelling but failed to have any

impact upon the histological outcomes measured in AIA joints.

Our results suggest that the *in vivo* response to pro-inflammatory primed EV treatment in the

AIA model of inflammatory arthritis is not primarily through inhibition of T cell activation, but

through suppression of CD4+ Th17 effector polarisation combined with reduction in synovial

infiltration through reduced recruitment of immune cells to the affected joint. Although this

effect is more pronounced following pro-inflammatory priming of MSCs in comparison to

standard EV production, the reduction in exudate and hyperplasia seen with EV-NormO2

treatments is likely to convey a greater therapeutic response. Pannus tissue is thought to

develop from cells of the synovial membrane and cells infiltrating the joint cavity in response

to pro-inflammatory signalling[37,38]. The reduction of synovial infiltrate, hyperplasia and

synovial exudate observed in EV-NormO2 treatments in vivo may therefore translate to

reduced pannus formation. These benefits, in combination with Th17 modulation and

improved Th17:Treg balance would convey advantageous outcomes in comparison to either

priming methodology.

We hypothesise that whilst the overall effectiveness of these treatment methodologies is

similar in potency, the outcomes differ due to variations in underlying mechanisms of action.

Our results demonstrate hypoxic preconditioning reduces swelling in vivo and affects Th17

polarisation of T cells in vitro, however insufficient evidence was found to propose a clear

mechanism for therapeutic capacity. MSCs under hypoxic culture have been shown to

promote anti-inflammatory M2 macrophage polarization[39], reduce reactive oxygen species

(ROS) and upregulate TGF-β, IL-8, IL-10 and PGE2, which are also implicated in macrophage

polarisation and MSC immunomodulatory capacity[39-41]. Consequently, we hypothesise that

effects seen in vivo following hypoxic preconditioning may be related to reduction in

macrophage mediated immune responses and more appropriate for tissue regenerative

applications. Furthermore, hypoxic culture of MSCs has been shown to promote

vascularisation[42,43] which, whilst beneficial for tissue repair, may actually contribute to

pannus formation in the RA joint[38]. Given these considerations, the evidence here suggests

that normal culture conditions are sufficient to elicit an optimal response in MSCs for EVs

production ahead of therapeutic application into the inflammatory environment.

Previously, we determined that joint swelling in induced arthritis as a clinical indication of

joint inflammation is reduced in the presence of administered MSCs and these cells migrate

into the inflamed synovium[2]. Joint swelling is common with different types of arthritis and is

caused by oedema due to the endothelial cells of blood vessels becoming leaky in the

inflamed synovium. We hypothesised that soluble factors produced by MSCs are responsible

for permeability changes in the synovial endothelial cells as these cells did not co-localise,

although further studies are required in this regard. The therapeutic potential of MSC-derived

EVs has been documented in animal model of radiation-induced lung injury where EVs

attenuated radiation-induced lung vascular damage, inflammation, and fibrosis via miRNA-

214-3p[44] and in a rodent model of haemorrhagic shock-induced lung injury model where EVs

attenuated pulmonary vascular permeability and lung injury[45].

Given the observed outcomes, we hypothesise that the similarities in immunomodulation

between EV-2%O2 and EV-Pro-Inflam treatment may lie in their influence on STAT3

activation. Increased HIF-1α activates STAT3[46]. STAT3 induces RORγt expression and is

activated by TNF-α, IL-6, IL-21 and IL-23, which are key cytokines in promoting Th17

differentiation, and RORγt is a master regulator of Th17 differentiation[47,48]. Moreover, STAT3

activation has been shown to be necessary for the development of Th17 cells and Th17

related autoimmunity, such as seen in RA[48] so EVs inhibiting STAT3 activation could lead to

the drop in Th17 polarisation observed. EVs from other mesenchymal cell types are implicated

in influencing STAT3 signalling to induce mesenchymal transition[49] and EVs from

macrophages are able to trigger nuclear translocation of STAT-3[35], demonstrating STAT3

activation as a target of EVs influence. STAT3 has been suggested as a therapeutic target in

the treatment of autoimmune inflammatory disorders such as RA[50] and inhibition of STAT3

by EVs would have direct therapeutic potential. Further work can elucidate this mechanism of

action.

In this study, our primary aim was to evaluate the efficacy of EV treatments *in vivo* during AIA.

We show that MSC priming leads to the release of EVs that if administered to mice with acute

inflammatory arthritis significantly ameliorate disease pathogenesis, mainly by inhibiting

Th17 polarization. Future studies will define the composition and sub-vesicular localisation of

proteins in EV cargos.

Conclusions

This study provides new evidence that supports the use of EVs in clinical therapies for RA and

similar autoimmune disorders. The possibility to manipulate the protein and nucleic acid

cargo of EVs through control of parental MSC cultures offers a novel opportunity for targeted

therapies that can be tailored to individual pathological features of RA, advancing

personalised medicine. Further work on the control of EV cargo will elucidate the molecular

mechanism of action underlying varied responses between priming strategies and assist in

the efficacy of cell-based therapies in the clinic. This study aims to support the growing body

of evidence for the introduction of EVs into the therapeutic milieu.

Methods

Cells and EVs

Primary human MSCs were isolated from commercially available bone marrow aspirate

(Lonza, USA) using an adherence technique[40] and each donor material cultured in both

normoxic ($21\%O_2$) or hypoxic ($2\%O_2$) conditions from isolation of MSCs in Dulbecco's

Modified Eagles Medium (DMEM) with 10% foetal bovine serum (FBS) and 1% penicillin-

streptomycin (n=3). Hypoxic cell culture was achieved using 2% O_2 in hypoxic workstation

(InvivO$_2$ Physiological Cell Culture Workstation, Baker Ruskinn) to ensure cells were not

exposed to environmental oxygen levels at any stage of isolation or culture. Cells (P3-P5) were

characterised as MSC through immunophenotyping of surface markers with flow cytometry

and tri-lineage differentiation[2]. EVs were therefore isolated over a 48 hour culture period

from MSCs in normoxic 21% oxygen culture (EV-NormO2, n=11); MSCs in hypoxic 2% oxygen

culture (EV-2%O2, n=4) and from MSCs previously cultured for 48 hours with pro-

inflammatory cytokine cocktail comprising interferon-gamma (IFN-γ, 10ng/ml), tumour

necrosis factor-alpha (TNFα, 10ng/ml) and interleukin 1 beta (IL-1β, 5ng/ml) prior to transfer

to serum-free medium for 48 hours for EV collection (EV-Pro-Inflam, n=4).

Differential ultracentrifugation for isolation of EVs

MSCs from the same batch used for cell treatment (P3-P5) were cultured to 80-90% confluence in T75 flasks, washed with PBS three times and then serum-free DMEM. Flasks were incubated for 48 hours with 12ml serum-free DMEM at 37°C, 5% CO_2. After 48 hours, conditioned medium (CM-MSC) was removed and a cell count performed to determine the number of cells used to produce EVs. EVs were isolated from CM-MSC by differential ultracentrifugation. In brief, CM-MSC was centrifuged for 10 minutes at 300 x g to remove cell debris then again for 10 minutes at 2000 x g to remove apoptotic bodies and residual dead cells. Supernatant was again taken and centrifuged at 10,000 x g for 45 minutes in an ultracentrifuge (Beckman Coulter) to remove larger/denser EVs. Supernatant was again retrieved and passed through a 0.22µM syringe filter. Filtered supernatant was then centrifuged using 70.1Ti rotor (Beckman Coulter) in a Beckman L8-55M ultracentrifuge at 100,000 x g for 90 minutes to isolate a pellet comprising small EVs. EVs depleted medium supernatant was removed and stored at -20°C (EVs-depleted CM-MSC), whilst the visible EVs pellet was resuspended in 5ml of PBS to wash EVs and then spun again at 100,000 x g for 60 minutes, the supernatant discarded and the residual EV pellet resuspended in PBS at 30µL per 1.0×10^6 cells used in EV generation and stored at 4°C to be used within 24 hours or at -20°C for later use.

Characterisation of EVs

Successful isolation of EVs was confirmed through multiple methods in accordance with ISEV guidelines[51]. Samples of vesicle preparations underwent bicinchoninic acid assay (BCA Assay,

Pierce Biotechnology) for total protein concentration in EV preparations following

manufacturer's instructions with EV disruption in a sonicating water bath (30 seconds

sonication at 1 minute intervals for 3 cycles) to release intracellular protein and assess cargo

proteins as well as surface proteins. Data passed a Kolmogorov-Smirnov test for normality

and was analysed with a repeated measures 1 way ANOVA with Tukey's post hoc multiple

comparison test.

Vesicles were examined using flow cytometry for expression of characteristic markers CD9,

CD63 and CD81 (Miltenyi MACSPlex Exosome Identification Kit, human) following

manufacturer's instructions for analysis. Particle by particle analysis of vesicle size was

assessed using Nanopore technology (Izon Science Ltd) tuned in the region ~80-300nm, to

confirm the size fractions being applied experimentally.

Images of EVs were obtained using Transmission Electron Microscopy (TEM). EVs to be

imaged by TEM were isolated as detailed above, however residual EV pellet after PBS wash

was resuspended in 30µl Milli-Q water per 3.0×10^6 cells. The resulting EV suspension was

spotted onto a glow discharged holey carbon mesh copper grids (Quantifoil, R2/1) and

incubated at room temperature for 4 mins 45 seconds before 5µL of 0.22µm filtered 2% (v/v)

uranyl acetate was added. This was left to incubate at room temperature for a further 90

seconds before the excess liquid was removed using blotting paper. Grids were then imaged

using a FEI Tecnai G2 12 Biotwin (LaB_6, accelerating voltage 100kV).

EV protein content was examined by Western Blotting to identify the presence of Alix and

absence of mitochondrial Cytochrome C (CYC1) in isolated particles, as per the minimal

criteria for identification of EVs[52]. Briefly, ten micrograms of EVs and cells were lysed with 2x

Laemlli buffer with β-mercaptoethanol and denatured by heating at 70ºC for 5min. Samples

were electrophoresed on a 4-12% TGX stain free gel (Bio-Rad, CA, USA) at 200V for 30-40min.

Samples were then blotted onto a nitrocellulose membrane using the wet transfer method

overnight at 100V. After blotting, membranes were blocked for 2 hours in 5% semi-skimmed

milk in Tris-buffered saline with 0.05% Tween (TBST) followed by an incubation in primary

antibodies Alix (1:200), CYC1 (1:200) (Santa Cruz Biotechnology, TX, US) for 2 hours at RT.

After three five-minute washes, the membrane was probed with a goat anti-mouse IgG-HRP

conjugated secondary antibody (1:1,000 in TBST, Life Technologies Limited, Paisley, UK). HRP-

conjugated secondary antibody was added for 1h and the wash step repeated. SuperSignal

West Femto Chemiluminescent Substrate (Thermo Fisher Scientific, Waltham, MA, USA) was

added to the membrane and imaged with ChemiDoc™ Touch Imaging System (Bio-Rad, CA,

USA) using Image Lab v.6.0.1 software (Bio-Rad, CA, USA).

Co-culture of healthy T cells with EVs

Proliferation of activated T cells was assessed as a measure of T cell deactivation. Initially, for

positive controls $5x10^4$ MSCs were cultured in 96-well plates for 24 hours at 37°C. CD4+ T cells

were purified from spleens and lymph nodes (popliteal/inguinal) of healthy C57Bl/6 mice

using the CD4+ T Cell Isolation Kit (Miltenyi) following manufacturer's instructions. T cells

were seeded at a density of $5.0x10^5$/well in 250µl RPMI medium with 10% FBS and cultured

for 5 days with MSCs (ratio 10:1) or with EV-NormO2, EV-2%O2 or EV-Pro-Inflam (ratio

equivalent to secretions from 10:1 cells) (n=8). T cells alone served as control. Cells were

activated using anti-Biotin MACSiBead Particles (Miltenyi) (ratio 2:1). EV treatments were

refreshed after 2 days of culture. Polarisation was assessed as above and proliferation

measured through reduction in signal intensity using VPD450 Violet proliferation dye (BD

Biosciences)[3]. Data was analysed using a 2 way repeated measures ANOVA with Bonferroni

multiple comparison test post hoc.

Antigen-induced arthritis (AIA) model of inflammatory arthritis

Animal procedures were undertaken in accordance with Home Office project licence

PPL40/3594. AIA was induced in male C57Bl/6 mice (7-8 weeks) as previously described[2,3,53].

Swelling was assessed by measuring the difference in diameter between the arthritic (right)

and non-arthritic (left) knee joints (in mm) using a digital micrometer (Kroeplin GmbH) before

and at set time points after treatment. Independent experiments were performed to assess

swelling and histopathological effects of EV-NormO2 (n=10), EV-2%O2 (n=6) and EV-Pro-

Inflam (n=6) compared with PBS controls (n=21) with all EVs sourced from matching MSCs

and statistical analysis performed using 2 Way ANOVA for Repeated Measures with

Bonferroni Multiple Comparisons Test post hoc; or to for in vivo primed T cell collections using

EV-NormO2 compared with EV-depleted CM-MSC (matched origin) with PBS control (n=4)

and log transformed data analysed with 1-Way ANOVA with Bonferroni Multiple Comparisons

Test post hoc.

Peak swelling was observed at 24 hours post-induction and results at subsequent timepoints

are expressed in millimetres as reduction from peak swelling.

Intra-articular injection of EVs

Treatments comprised 15µl of EVs suspension corresponding to EV secretions from ~5.0 x 10^5

cells, or EV-depleted CM-MSC (basal medium using serum free DMEM). Treatments or PBS

vehicle control were injected intra-articularly 1 day post arthritis induction with 0.5ml

monoject (29G) insulin syringes (BD Micro-Fine, Franklyn Lakes, USA) through the patellar

ligament into the right knee joint. Joint diameters were measured at 1, 2 and 3 days post

injection. Blood, joints, spleen, inguinal and popliteal lymph nodes were collected

immediately post-mortem. Four independent experiments were performed to assess primed

EVs impact on joint swelling and histopathology (Control n=21, EV-NormO2 n=10, EV-2%O2

and EV-Pro-Inflam n=6). A further independent experiment was conducted to address the

mechanism behind conditioned medium efficacy in immunomodulation of T cell polarisation

in AIA by specifically identifying the contribution of EVs to secretomic effects (EV-NormO2

versus EV-depleted conditioned medium with PBS controls, n=4 per condition). All measures

were taken to reduce animal numbers wherever possible.

Arthritis Index

Animals were sacrificed for histological analysis at day 3 post arthritis induction. Joints were

fixed in 10% neutral buffered formal saline and decalcified in formic acid for 4 days at 4°C

before paraffin embedding. Sections (5µm) were stained with haematoxylin and eosin (H&E)

and mounted in Hydromount (National Diagnostics) as described previously[2,3]. H&E sections

were scored for hyperplasia of the synovial intima (0=normal to 3=severe), cellular exudate

(0=normal to 3=severe) and synovial infiltrate (0=normal to 5=severe) by two independent

observers blinded to experimental groups[54]. Scores were summated, producing a mean

arthritis index. Data was analysed using a 1 way repeated measures analysis of variance test

with Bonferroni multiple comparison test post hoc.

Cytokine Quantification.

IL-10 and TNFα in serum and CM-MSC were quantified using mouse Quantikine ELISA IL-10

immunoassay (R&D Systems) and TNFα ELISA high sensitivity (eBioscience) respectively,

following manufacturer's instructions.

T cell polarisation

Spleens and lymph nodes (popliteal/ inguinal) were collected from mice 3 days post-arthritis

induction and dissociated as described previously[3]. Splenocytes and pooled lymph node cells

were seeded separately at 1.0×10^6 cells/well in 96-well plates (Sarstedt) in RPMI-1640 with

10% FBS, 0.05μg/mL IL-2 and activated with cell stimulation cocktail (eBioscience) for 1 hour

prior to adding 10μg/ml brefeldin A (Sigma) and culturing for a further 4 hours. Unstimulated

T cells and cells without brefeldin A served as negative controls. Following activation, cells

were resuspended in 2mM EDTA in PBS and Tregs (CD4+CD25+FOXP3+) were isolated using

the CD4+CD25+ Regulatory T Cell Isolation Kit (Miltenyi) following manufacturer's

instructions; or stained for T cell subset identification. For this, cells were permeabilised using

permeabilisation buffer kit (eBioscience) and intracellularly stained with anti-mouse IFN-γ

(Th1), IL-4 (Th2) or IL17a (Th17) (eBioscience). Cells were analysed on a BD FACS Canto II flow

cytometer and comparisons drawn for percentage CD4+ cells and signal intensity (XGeoMean)

for each antibody.

Statistical analysis

Data were tested for equal variance and normality using D'Agostino & Pearson omnibus

normality test. Differences between groups were compared using 1-way ANOVA with

Bonferroni post hoc for parametric data or Kruskal-Wallis ANOVA with Dunn's post hoc test

for non-parametric, or 2-way ANOVA with Bonferroni correction, as stated. Repeated

measures tests using log transformed data was applied to Protein concentration was analysed

with log transformed data using a 4 parameter polynomial nonlinear regression to interpolate

results from a standard curve. All statistical analysis was carried out using Prism 5 (GraphPad

software) or IBM SPSS Statistics 24.0, with $P < 0.05$ deemed statistically significant. Results are

expressed as mean ± standard error of the mean.

List of abbreviations

AIA – Antigen Induced Arthritis model

BCA – Bicinchoninic Acid Protein Assay

CM-MSC – MSCs derived conditioned medium

CTSD – Cathepsin D

CYC1 – Cytochrome C

DMEM – Dulbecco's Modified Eagle's Medium

EV-2%O2 – EVs derived from cells cultured in 2% oxygen

EV-NormO2 – EVs derived from cells cultured in normal environmental oxygen ~21%

EV-Pro-Inflam – EVs derived from MSCs cultured with pro-inflammatory cytokine stimulation

EVs – Extracellular vesicles

FBS – Foetal bovine serum

FIH – Factor inhibiting Hif-1 protein

H&E – Haematoxylin and eosin stain

HIF-1α – Hypoxia inducible factor 1 alpha

ICAM-1 – Intercellular adhesion molecule 1

IDO – Indoleamine 2,3-dioxygenase

IFN-γ – Interferon gamma

IL-10 – Interleukin 10

IL-1β – Interleukin 1 beta

ISEV – International Society of Extracellular Vesicles

LFA-1 – Leukocyte function associated antigen 1

MFI – Mean fluorescence intensity

miRNA – micro ribonucleic acid

MMPs – Matrix metalloproteins

MSCs – Mesenchymal stem/stromal cells

MVBs/MVEs – Multivesicular bodies/endosomes

NO – Nitric oxide

PBS – Phosphate buffered saline

RA – Rheumatoid Arthritis

RF – Rheumatoid factor

RISC – RNA-induced silencing complexes

RORγt – Retinoic acid receptor-related orphan receptor-γt

ROS – Reactive oxygen species

TIMPs – Tissue inhibitor of metalloproteinases

TNF-α – Tissue necrosis factor alpha

Treg – Regulatory T cell

VEGF – Vascular endothelial growth factor

VHL – von Hippel-Lindau ubiquitin ligase

Acknowledgements

The authors would like to thank the Nanoscale and Microscale Research Centre, University of

Nottingham, for access to TEM facilities. We would like to thank Dr. Mark Platt (Loughborough

University, UK) for his assistance in Nanopore technology.

Declaration of Interest Statement

The authors declare that they have no competing interests

Funding

This work was supported by the RJAH Orthopaedic Hospital Charity under Grant [G08028];

the UK EPSRC/MRC CDT in Regenerative Medicine (Keele University, Loughborough

University, the University of Nottingham) under Grant [EP/F500491/1]; Orthopaedic

Institute, Ltd under Grant [RPG 171].

Author's contributions

A.K. performed experimental work, wrote the manuscript and prepared all figures. K.T.

provided support and technical skills for T cell investigations. P.R. gave technical expertise

on Nanopore technology. R.M. provided support for histological investigations, TEM

experiments and discussed and commented on the manuscript. R.L. performed TEM

experiments. M.H. performed ELISAs and provided support for *in vivo* work and histological

experiments. A.M.P. gave technical advice on data interpretations and presentation and

reviewed the manuscript. N.R.F. contributed to experimental design. O.K. conceived and

designed the research study, performed *in vivo* experiments, analysed data, wrote and

supervised the manuscript. All authors reviewed the manuscript before submission.

References

1   Prockop, D. J. & Oh, J. Y. Mesenchymal stem/stromal cells (MSCs): role as guardians of
inflammation. *Mol Ther* **20**, 14-20, doi:10.1038/mt.2011.211 (2012).
2   Kehoe, O., Cartwright, A., Askari, A., El Haj, A. J. & Middleton, J. Intra-articular injection of
mesenchymal stem cells leads to reduced inflammation and cartilage damage in murine
antigen-induced arthritis. *J Transl Med* **12**, 157, doi:10.1186/1479-5876-12-157 (2014).
3   Kay, A. G. *et al.* Mesenchymal Stem Cell-Conditioned Medium Reduces Disease Severity and
Immune Responses in Inflammatory Arthritis. *Sci Rep* **7**, 18019, doi:10.1038/s41598-017-
18144-w (2017).
4   Stanton, H. *et al.* ADAMTS5 is the major aggrecanase in mouse cartilage in vivo and in vitro.
Nature **434**, 648-652 (2005).
5   Beer, K. B. *et al.* Extracellular vesicle budding is inhibited by redundant regulators of TAT-5
flippase localization and phospholipid asymmetry. *Proc Natl Acad Sci U S A* **115**, E1127-
E1136, doi:10.1073/pnas.1714085115 (2018).
6   Couper, K. N., Blount, D. G. & Riley, E. M. IL-10: the master regulator of immunity to
infection. *J Immunol* **180**, 5771-5777, doi:10.4049/jimmunol.180.9.5771 (2008).
7   Kusuma, G. D. *et al.* To Protect and to Preserve: Novel Preservation Strategies for
Extracellular Vesicles. *Front Pharmacol* **9**, 1199, doi:10.3389/fphar.2018.01199 (2018).
8   Muralidharan-Chari, V., Clancy, J. W., Sedgwick, A. & D'Souza-Schorey, C. Microvesicles:
mediators of extracellular communication during cancer progression. *J Cell Sci* **123**, 1603-
1611, doi:10.1242/jcs.064386 (2010).
9   Raposo, G. & Stoorvogel, W. Extracellular vesicles: exosomes, microvesicles, and friends. *J
Cell Biol* **200**, 373-383, doi:10.1083/jcb.201211138 (2013).
10   Burbano, C. *et al.* Extracellular vesicles are associated with the systemic inflammation of
patients with seropositive rheumatoid arthritis. *Sci Rep* **8**, 17917, doi:10.1038/s41598-018-
36335-x (2018).

11 Arntz, O. J. *et al.* Rheumatoid Arthritis Patients With Circulating Extracellular Vesicles
Positive for IgM Rheumatoid Factor Have Higher Disease Activity. *Front Immunol* **9**, 2388,
doi:10.3389/fimmu.2018.02388 (2018).
12 Messer, L. *et al.* Microparticle-induced release of B-lymphocyte regulators by rheumatoid
synoviocytes. *Arthritis Res Ther* **11**, R40, doi:10.1186/ar2648 (2009).
13 Lo Cicero, A., Majkowska, I., Nagase, H., Di Liegro, I. & Troeberg, L. Microvesicles shed by
oligodendroglioma cells and rheumatoid synovial fibroblasts contain aggrecanase activity.
Matrix Biol **31**, 229-233, doi:10.1016/j.matbio.2012.02.005 (2012).
14 Zhang, H. G. *et al.* A membrane form of TNF-alpha presented by exosomes delays T cell
activation-induced cell death. *J Immunol* **176**, 7385-7393,
doi:10.4049/jimmunol.176.12.7385 (2006).
15 Kim, S. H. *et al.* Exosomes derived from IL-10-treated dendritic cells can suppress
inflammation and collagen-induced arthritis. *J Immunol* **174**, 6440-6448,
doi:10.4049/jimmunol.174.10.6440 (2005).
16 Lee, C. *et al.* Exosomes mediate the cytoprotective action of mesenchymal stromal cells on
hypoxia-induced pulmonary hypertension. *Circulation* **126**, 2601-2611,
doi:10.1161/CIRCULATIONAHA.112.114173 (2012).
17 Mokarizadeh, A. *et al.* Microvesicles derived from mesenchymal stem cells: potent
organelles for induction of tolerogenic signaling. *Immunol Lett* **147**, 47-54,
doi:10.1016/j.imlet.2012.06.001 (2012).
18 Zhang, B. *et al.* Mesenchymal stem cells secrete immunologically active exosomes. *Stem*
Cells Dev **23**, 1233-1244, doi:10.1089/scd.2013.0479 (2014).
19 Cosenza, S. *et al.* Mesenchymal stem cells-derived exosomes are more immunosuppressive
than microparticles in inflammatory arthritis. *Theranostics* **8**, 1399-1410,
doi:10.7150/thno.21072 (2018).
20 Alcaraz, M. J., Compan, A. & Guillen, M. I. Extracellular Vesicles from Mesenchymal Stem
Cells as Novel Treatments for Musculoskeletal Diseases. *Cells* **9**, doi:10.3390/cells9010098
(2019).
21 Ma, D. *et al.* Immunomodulatory effect of human umbilical cord mesenchymal stem cells on
T lymphocytes in rheumatoid arthritis. *Int Immunopharmacol* **74**, 105687,
doi:10.1016/j.intimp.2019.105687 (2019).
22 Thery, C., Ostrowski, M. & Segura, E. Membrane vesicles as conveyors of immune responses.
Nat Rev Immunol **9**, 581-593, doi:10.1038/nri2567 (2009).
23 Nolte-'t Hoen, E. N., Buschow, S. I., Anderton, S. M., Stoorvogel, W. & Wauben, M. H.
Activated T cells recruit exosomes secreted by dendritic cells via LFA-1. *Blood* **113**, 1977-
1981, doi:10.1182/blood-2008-08-174094 (2009).
24 Tung, S. L. *et al.* Regulatory T cell-derived extracellular vesicles modify dendritic cell function.
Sci Rep **8**, 6065, doi:10.1038/s41598-018-24531-8 (2018).
25 Djouad, F. *et al.* Reversal of the immunosuppressive properties of mesenchymal stem cells
by tumor necrosis factor alpha in collagen-induced arthritis. *Arthritis Rheum* **52**, 1595-1603,
doi:10.1002/art.21012 (2005).
26 Firestein, G. S. Rheumatoid arthritis in a mouse? *Nat Clin Pract Rheumatol* **5**, 1,
doi:10.1038/ncprheum0973 (2009).
27 Angulo, R. & Fulcher, D. A. Measurement of Candida-specific blastogenesis: comparison of
carboxyfluorescein succinimidyl ester labelling of T cells, thymidine incorporation, and CD69
expression. *Cytometry* **34**, 143-151 (1998).
28 Que, J., Lian, Q., El Oakley, R. M., Lim, B. & Lim, S.-K. PI3 K/Akt/mTOR-mediated translational
control regulates proliferation and differentiation of lineage-restricted RoSH stem cell lines.
Journal of Molecular Signaling **2**, doi:10.1186/1750-2187-2-9 (2007).

29   Al-Zifzaf, D. S. *et al.* FoxP3+T regulatory cells in Rheumatoid arthritis and the imbalance of
748   the Treg/TH17 cytokine axis. *The Egyptian Rheumatologist* **37**, 7-15,
749   doi:10.1016/j.ejr.2014.06.004 (2015).

750   30   Arroyo-Villa, I. *et al.* Frequency of Th17 CD4+ T cells in early rheumatoid arthritis: a marker
751   of anti-CCP seropositivity. *PLoS One* **7**, e42189, doi:10.1371/journal.pone.0042189 (2012).

752   31   Diller, M. L., Kudchadkar, R. R., Delman, K. A., Lawson, D. H. & Ford, M. L. Balancing
753   Inflammation: The Link between Th17 and Regulatory T Cells. *Mediators of Inflammation*
754   **2016**, 1-8, doi:10.1155/2016/6309219 (2016).

755   32   Noack, M. & Miossec, P. Th17 and regulatory T cell balance in autoimmune and
756   inflammatory diseases. *Autoimmun Rev* **13**, 668-677, doi:10.1016/j.autrev.2013.12.004
757   (2014).

758   33   Lewthwaite, J. *et al.* Role of TNF alpha in the induction of antigen induced arthritis in the
759   rabbit and the anti-arthritic effect of species specific TNF alpha neutralising monoclonal
760   antibodies. *Ann Rheum Dis* **54**, 366-374, doi:10.1136/ard.54.5.366 (1995).

761   34   Kobayashi, H. *et al.* Antigen induced arthritis (AIA) can be transferred by bone marrow
762   transplantation: Evidence that interleukin 6 is essential for induction of AIA. *Journal of*
763   *Rheumatology* **29**, 1176-1182 (2002).

764   35   Bellmunt À, M., López-Puerto, L., Lorente, J. & Closa, D. Involvement of extracellular vesicles
765   in the macrophage-tumor cell communication in head and neck squamous cell carcinoma.
766   *PLoS One* **14**, e0224710, doi:10.1371/journal.pone.0224710 (2019).

767   36   Maumus, M. *et al.* Utility of a Mouse Model of Osteoarthritis to Demonstrate Cartilage
768   Protection by IFNgamma-Primed Equine Mesenchymal Stem Cells. *Front Immunol* **7**, 392,
769   doi:10.3389/fimmu.2016.00392 (2016).

770   37   Fassbender, H. G. & Simmling-Annefeld, M. The potential aggressiveness of synovial tissue in
771   rheumatoid arthritis. *The Journal of Pathology* **139**, 399-406,
772   doi:https://doi.org/10.1002/path.1711390314 (1983).

773   38   Shiozawa, S., Shiozawa, K. & Fujita, T. Morphologic observations in the early phase of the
774   cartilage-pannus junction. Light and electron microscopic studies of active cellular pannus.
775   *Arthritis Rheum* **26**, 472-478, doi:10.1002/art.1780260404 (1983).

776   39   Takizawa, N. *et al.* Bone marrow-derived mesenchymal stem cells propagate
777   immunosuppressive/anti-inflammatory macrophages in cell-to-cell contact-independent and
778   -dependent manners under hypoxic culture. *Exp Cell Res* **358**, 411-420,
779   doi:10.1016/j.yexcr.2017.07.014 (2017).

780   40   Kay, A. G. *et al.* BMP2 repression and optimized culture conditions promote human bone
781   marrow-derived mesenchymal stem cell isolation. *Regen Med* **10**, 109-125,
782   doi:10.2217/rme.14.67 (2015).

783   41   Li, B. *et al.* Hypoxia-Induced Mesenchymal Stromal Cells Exhibit an Enhanced Therapeutic
784   Effect on Radiation-Induced Lung Injury in Mice due to an Increased Proliferation Potential
785   and Enhanced Antioxidant Ability. *Cell Physiol Biochem* **44**, 1295-1310,
786   doi:10.1159/000485490 (2017).

787   42   Bader, A. M. *et al.* Hypoxic Preconditioning Increases Survival and Pro-Angiogenic Capacity of
788   Human Cord Blood Mesenchymal Stromal Cells In Vitro. *PLoS One* **10**, e0138477,
789   doi:10.1371/journal.pone.0138477 (2015).

790   43   Leroux, L. *et al.* Hypoxia preconditioned mesenchymal stem cells improve vascular and
791   skeletal muscle fiber regeneration after ischemia through a Wnt4-dependent pathway. *Mol*
792   *Ther* **18**, 1545-1552, doi:10.1038/mt.2010.108 (2010).

793   44   Lei, X. *et al.* Mesenchymal Stem Cell-Derived Extracellular Vesicles Attenuate Radiation-
794   Induced Lung Injury via miRNA-214-3p. *Antioxid Redox Signal*, doi:10.1089/ars.2019.7965
795   (2020).

45   Potter, D. R. *et al.* Mesenchymal stem cell-derived extracellular vesicles attenuate
pulmonary vascular permeability and lung injury induced by hemorrhagic shock and trauma.
J Trauma Acute Care Surg **84**, 245-256, doi:10.1097/ta.0000000000001744 (2018).
46   Almiron Bonnin, D. A. *et al.* Secretion-mediated STAT3 activation promotes self-renewal of
glioma stem-like cells during hypoxia. *Oncogene* **37**, 1107-1118, doi:10.1038/onc.2017.404
(2018).
47   Zhou, L. *et al.* TGF-beta-induced Foxp3 inhibits T(H)17 cell differentiation by antagonizing
RORgammat function. *Nature* **453**, 236-240, doi:10.1038/nature06878 (2008).
48   Harris, T. J. *et al.* Cutting edge: An in vivo requirement for STAT3 signaling in TH17
development and TH17-dependent autoimmunity. *J Immunol* **179**, 4313-4317,
doi:10.4049/jimmunol.179.7.4313 (2007).
49   Schweiger, M. W. *et al.* Extracellular Vesicles Induce Mesenchymal Transition and
Therapeutic Resistance in Glioblastomas through NF-kappaB/STAT3 Signaling. *Adv Biosyst* **4**,
e1900312, doi:10.1002/adbi.201900312 (2020).
50   Gaffen, S. L., Jain, R., Garg, A. V. & Cua, D. J. The IL-23-IL-17 immune axis: from mechanisms
to therapeutic testing. *Nat Rev Immunol* **14**, 585-600, doi:10.1038/nri3707 (2014).
51   Thery, C. *et al.* Minimal information for studies of extracellular vesicles 2018 (MISEV2018): a
position statement of the International Society for Extracellular Vesicles and update of the
MISEV2014 guidelines. *J Extracell Vesicles* **7**, 1535750, doi:10.1080/20013078.2018.1535750
(2018).
52   Lotvall, J. *et al.* Minimal experimental requirements for definition of extracellular vesicles
and their functions: a position statement from the International Society for Extracellular
Vesicles. *J Extracell Vesicles* **3**, 26913, doi:10.3402/jev.v3.26913 (2014).
53   Glant, T. T. *et al.* Characterization and Localization of Citrullinated Proteoglycan Aggrecan in
Human Articular Cartilage. *PLoS One* **11**, e0150784, doi:10.1371/journal.pone.0150784
(2016).
54   Misjak, P. *et al.* The role of citrullination of an immunodominant proteoglycan (PG) aggrecan
T cell epitope in BALB/c mice with PG-induced arthritis. *Immunol Lett* **152**, 25-31,
doi:10.1016/j.imlet.2013.03.005 (2013).