Delayed effects of climate on vital rates lead to demographic divergence in Amazonian forest fragments

Eric R. Scott[1,✉], María Uriarte[2], and Emilio M. Bruna[1,3,4]

08 September, 2021

[1] Department of Wildlife Ecology and Conservation, University of Florida, Gainesville, Florida 32611-0430

[2] Department of Ecology, Evolution and Environmental Biology, Columbia University 1200 Amsterdam Avenue, New York, New York 10027 USA

[3] Center for Latin American Studies, University of Florida, Gainesville, Florida 32611-5530

[4] Biological Dynamics of Forest Fragments Project, INPA-PDBFF, CP 478, Manaus, Amazonas 69011-970 Brazil

✉ Correspondence: Eric R. Scott <scottericr@gmail.com>

Keywords: Amazon; delayed effects; demography; distributed lag non-linear models; drought; ENSO; habitat fragmentation; *Heliconia acuminata*; Heliconiaceae

Running title: Delayed effects of climate on vital rates

Abstract

Deforestation often results in landscapes where remaining forest habitat is highly

fragmented, with remnants of different sizes embedded in an often highly contrasting

matrix. Local extinction of species from individual fragments is common, but the

demographic mechanisms underlying these extinctions are poorly understood. It is often

hypothesized that altered environmental conditions in fragments drive declines in

reproduction, recruitment, or survivorship. The Amazon basin, in addition to experiencing

continuing fragmentation, is experiencing climate change related increases in the

frequency and intensity of droughts and unusually wet periods. Whether plant populations

in tropical forest fragments are particularly susceptible to extremes in precipitation

remains unclear. Most studies of plants in fragments are relatively short (1–6 years), focus

on a single life-history stage, and often do not compare to populations in continuous forest.

Even fewer studies consider delayed effects of climate on demographic vital rates despite

the importance of delayed effects in studies that consider them. Using a decade of

demographic and climate data from an experimentally fragmented landscape in the Central

Amazon, we assess the effects of climate on populations of an understory herb (*Heliconia*

acuminata, Heliconiaceae). We used distributed lag non-linear models to understand the

delayed effects of climate (measured as standardized precipitation evapotranspiration

index, SPEI) on survival, growth, and flowering. We detected delayed effects of climate up

to 36 months. Extremes in SPEI in the previous year reduced survival, drought in the wet

season 8–11 months prior to the February census increased growth, and drought two dry

seasons prior increased flowering probability. Effects of extremes in precipitation on

survival and growth were more pronounced in forest fragments compared to continuous

40   forest. The complex delayed effects of climate and habitat fragmentation in our study point

41   to the importance of long-term demography experiments in understanding the effects of

42   anthropogenic change on plant populations.

Introduction

The expansion of agriculture and other human activities is a primary driver of deforestation in the tropics (Alroy, 2017; Haddad et al., 2015). It also results in landscapes where the remaining forest can be highly fragmented, with patches of different sizes embedded in a matrix of often contrasting habitat (Bianchi & Haig, 2013; Taubert et al., 2018). This fragmentation is associated with myriad ecological changes, including the local and regional extinction of plant species (da Silva & Tabarelli, 2000; Laurance et al., 2006). It is often hypothesized that the dramatically altered environmental conditions in tropical forest fragments (Arroyo-Rodríguez et al., 2017; Didham & Lawton, 1999; Ewers & Banks-Leite, 2013) drive declines in plant reproduction, recruitment, or survivorship (Bruna, 1999; Laurance et al., 1998; Zartman et al., 2015), although the demographic mechanisms responsible for these extinctions are poorly understood (Bruna et al., 2009). Despite the prevalence of this hypothesis (Betts et al., 2019; Didham & Lawton, 1999; Laurance et al., 2001), efforts to link population-level demographic responses with altered environmental conditions in fragments remains scarce.

Studies in temperate systems have shown that the demography of species can also be altered by climate change (Doak & Morris, 2010; Selwood et al., 2015; Sletvold, 2005; Williams et al., 2015). While the demographic consequences of climate change for tropical species are expected to be similarly severe (Brodie et al., 2012; Scheffers et al., 2017), surprisingly little is known about the responses of these species to climatic variability (Paniw et al., 2021). Climate models predict increases in the areas of the Amazon effected by severe drought and unusual wetness by the year 2100 (Duffy et al., 2015). Tropical plants may be particularly sensitive to climate change—they typically have narrow ranges

of climatic tolerance (Feeley et al., 2012), and recent results suggest increases in the

frequency and severity of extreme precipitation events (both drought and extreme wet)

reduce survival and reproduction (Esteban et al., 2021; Gaoue et al., 2019). This sensitivity

to climatic fluctuations, coupled with evidence that plant growth and survivorship are

lower in fragments (Bruna et al., 2002; Laurance et al., 1998; Zartman et al., 2015), has led

to speculation that plants in forest fragments will be especially susceptible to climate

change (Laurance et al., 2001; Opdam & Wascher, 2004; Selwood et al., 2015).

The simultaneous pressures of climate change and habitat fragmentation may also result in

worse than additive impacts on demography (Holyoak & Heath, 2016; Oliver et al., 2015).

For example, if fragments have a reduced capacity to buffer changes in microclimate

(Didham & Lawton, 1999; Ewers & Banks-Leite, 2013) or if fragment connectivity is climate

dependent (Honnay et al., 2002), populations experiencing habitat fragmentation will fare

worse under climate change.

Whether the demography of plant populations in tropical forest fragments is more

susceptible to climatic extremes remains unclear for three primary reasons. First, most

studies of plants in fragments have focused on a single life-history stage or process (Bruna

et al., 2009; Ehrlen et al., 2016), making it challenging to draw broader demographic

conclusions. Second, there is a growing literature on how tropical plants respond to

droughts (Esquivel-Muelbert et al., 2019; González-M et al., 2020; Uriarte et al., 2016), but

few studies have compared the responses of plants in continuous forest with those of

plants in forest fragments (Laurance et al., 2001). Finally, the multi-year data needed to

test population-level hypotheses about climate change and fragmentation are scant,

especially for tropical systems (Crone et al., 2011; Salguero-Gómez et al., 2015). These data

are critical not simply because they allow for capturing variation in climatic conditions and

the resulting demographic responses (Morris & Doak, 2002; Teller et al., 2016). They are

also essential because while some demographic effects of fragmentation or drought can be

detected immediately, others may take years to manifest (*e.g.,* Gagnon et al., 2011). Indeed,

lagged responses of demographic vital rates to climate may be the rule rather than the

exception (Anderegg et al., 2015; Evers et al., 2021; Kannenberg et al., 2020; Schwalm et al.,

2017).

Herbaceous plants represent up to 25% of plant diversity in tropical forests (Gentry &

Dodson, 1987), are critical food and habitat for myriad species (Snow, 1981), and are

economically and culturally vital (Nakazono et al., 2004; Ticktin, 2003). Nevertheless, the

impacts of global change phenomena on their demography remain conspicuously

understudied (Bruna et al., 2009). We used a decade of demographic and climatic data from

an experimentally fragmented landscape in the Central Amazon to assess the effects of

climate on populations of a tropical understory herb (*Heliconia acuminata*, Heliconiaceae).

This time series, which included the severe droughts of 1997 (McPhaden, 1999) and 2005

(Marengo et al., 2008; Zeng et al., 2008), allowed us to address the following questions: (1)

Does drought increase or decrease the growth, survival, and fertility of plant populations in

continuous forest? (2) Are there delayed effects of drought on demographic vital rates, and

if so what lag times are most critical? (3) Are the effects of drought on the vital rates of

populations in fragments similar in direction and magnitude to those in continuous forest?

Methods

Study site

The Biological Dynamics of Forest Fragments Project (BDFFP) is located ~70 km north of Manaus, Brazil (2º30' S, 60ºW). In addition to large areas of continuous forest, the BDFFP has forest fragment reserves isolated from 1980–1984 by felling the trees surrounding the area chosen for isolation and, in most cases, burning the downed trees once they dried (Bierregaard et al., 1992). In subsequent decades the vegetation regenerating around fragments has been periodically cleared to ensure fragment isolation (Bierregaard et al., 2001).

The BDFFP reserves are located in nonflooded (i.e., *terra firme*) tropical lowland forest with a 30–37m tall canopy (Rankin-de-Mérona et al., 1992) and an understory dominated by stemless palms (Scariot, 1999). The soils in the reserves are nutrient-poor xanthic ferrosols; their water retention capacity is poor despite having a high clay content. Mean annual temperature in the region is 26º C (range=19–39º C), and annual rainfall ranges from 1900–2300 mm. There is a pronounced dry season from June to October (Figure S1).

Focal species

Heliconia acuminata (LC Rich.) (Heliconiaceae) is a perennial monocot distributed throughout Central Amazonia (Kress, 1990) and is the most abundant understory herb at the BDFFP (Ribeiro et al., 2010). While many *Heliconia* species grow in large patches in treefall gaps and other disturbed areas, others, such as *H. acuminata,* are found at lower densities in the darker and cooler forest understory (Rundel et al., 2020). These species produce fewer inflorescences and are pollinated by traplining rather than territorial

hummingbirds (Bruna et al., 2004; Stouffer & Bierregaard, 1996). In our sites *H. acuminata*

is pollinated by *Phaeothornis superciliosus* and *P. bourcieri* (Bruna et al., 2004). Plants begin

flowering at the start of the rainy season; reproductive plants have $\bar{x} = 1.1$ flowering

shoots (range = 1–7), each of which has an inflorescence with 20–25 flowers (Bruna &

Kress, 2002). Fruits mature April-May, have 1–3 seeds per fruit ($\bar{x} = 2$), and are eaten by a

thrush and several species of manakin (Uriarte et al., 2011). Dispersed seeds germinate

approximately 6 months after dispersal at the onset of the subsequent rainy season, with

rates of germination and seedling establishment higher in continuous forest than forest

fragments (Bruna, 1999; Bruna & Kress, 2002).

Demographic data collection

This study uses data collected in four 1-ha fragment reserves and six continuous forest

sites. In 1997–1998 we established a 5000 m² plots (50 × 100m) in each of these sites in

which we marked and measured all *Heliconia acuminata*; plots in 1-ha fragments were on

one randomly selected half of the fragment, while plots in continuous forest were located

500–4000 m from the borders of secondary and mature forest. The distance between plots

ranged from 500 m–41 km. Our dataset comprised 4,886 plants in continuous forest and

1,375 plants in forest fragments. Plots in CF had on average more than twice as many

plants than plots in 1-ha fragments (CF median = 788, range = [201, 1549]; 1-ha median =

339, range = [297, 400]).

Each plot was subdivided into 50 quadrats (10 × 10m) to simplify annual surveys, during

which we recorded the number of vegetative shoots each plant had, the height of each plant

(i.e. distance from the ground to the tallest leaf tip), and whether each plant was flowering

(height and shoot number are correlated with leaf area, the probability of flowering, and

rates of survivorship (Bruna, 2002; Bruna & Kress, 2002). In this study, we used the

product of shoot number and plant height as our measure of plant size. Preliminary

analysis showed that the product of shoot number and height was a better predictor of

total leaf area (which in turn is assumed to be a strong predictor of aboveground biomass)

than either shoot number or height alone (Table S2). After the initial census year, between

80% and 97% of marked plants were found at each survey. Of plants that had missing

values for some years, but were found again in a subsequent year, 95% had 2 or fewer

years of missing values. Therefore, plants that were not found for three consecutive

surveys, and no subsequent survey, were considered to have died in the transition year

after their last observation. We also surveyed plots regularly during the rainy season to

identify any that flowered after the survey. For additional details on the location of plots,

survey methods, and *H. acuminata* population structure see Bruna & Kress (2002).

Climate data

Data on precipitation and potential evapotranspiration in our sites were obtained from a

published gridded dataset (0.25º × 0.25º resolution) built using data from 3,625 ground-

based weather stations across Brazil (Xavier et al., 2016). We used these data to calculate

the standardized precipitation evapotranspiration index (SPEI) (Vicente-Serrano et al.,

2010). SPEI is a proxy for meteorological drought that integrates precipitation and

evapotranspiration anomalies over a specified time scale. Positive SPEI values for a given

month indicate conditions wetter than the historical average for that month, while negative

values of SPEI indicate droughts with intensity categorized as mild (0 to -1), moderate (-1

to -1.5), severe (-1.5 to -2), or extreme (< -2) (McKee et al., 1993). SPEI can be calculated to

represent different temporal scales of drought; we used 3-month SPEI because—given its

shallow roots and rhizome—*H. acuminata* relies primarily on soil moisture rather than

deeper water sources that can persist for longer timescales (McKee et al., 1993; Vicente-

Serrano et al., 2010). Note that 3-month SPEI is still monthly data—each month's SPEI

value simply takes into account precipitation and evapotranspiration of the previous three

months. SPEI calculations were made using the SPEI package in R (Beguería & Vicente-

Serrano, 2017). The timing of drought events based on these SPEI calculations is consistent

with that resulting from SPEI calculated with other data sources, though the magnitude of

drought sometimes differed (Figure S2; Table S1).

Statistical Modeling of Vital Rates

To assess the effects of drought history on plant vital rates we used Distributed Lag Non-

linear Models (DLNMs, Gasparrini et al., 2017). DLNMs capture how potentially delayed

effects of predictor variables (e.g. SPEI) affect an outcome (e.g. growth) well beyond the

event period. DLNMs avoid the problems of including all possible climate windows in a

single model—namely collinearity of predictors and the arbitrary choice of window length.

They do so by fitting a bi-dimensional predictor-lag-response association spline, referred to

as a crossbasis function. This models a non-linear relationship between predictor and

response (e.g. between SPEI and vital rates) and allows the shape of that relationship to

vary smoothly over lag time. The shape of the function can be penalized so that it is only as

complex as the data supports, thus reducing effective degrees of freedom. We used a

crossbasis function of SPEI and lag time with possible lags from 0–36 months. We chose 36

months as a maximum lag because prior transplant experiments with *H. acuminata* showed

they typically recover from transplant shock in less than 36 months (Bruna et al., 2002) so

this is a reasonable upper bound for lagged effects of drought. The general form of the vital

rate (y) models was as follows:

202
$$g[E(y_{ij})] = \alpha_j + s_1(z_i) + s_2(d_{i,t}, \ldots, d_{i,t-36}) + \beta x_i$$
$$\alpha_j \sim N\left(\mu_{\alpha_j}, \sigma^2\right), \text{ for plot j}$$
(1)

where y_{ij} is the ith plant observation in the jth plot $s_1(z_i)$ is a smooth function of plant size

(natural log of height × shoot number), fit using a penalized cubic regression spline, $s_2(\cdot)$ is

the crossbasis function in which $d_{i,t}$ is the SPEI value during the census month of an

observation (February) and $d_{i,t-l}$ is the SPEI l months prior (see Gasparrini et al. 2017 for

details). To determine if plot characteristics influenced average vital rates we included a

random effect of plot ID on the intercept; this was represented by a_j in eq. 1. We modeled a

potential cost of reproduction by including flowering in the previous year as covariate, x_i,

in eq. 1. Individual-level random effects were not included due to computational limitations

in fitting random intercepts for all of the nearly 5,000 plants in continuous forest plots.

Fitting models with a data subset with and without individual-level random effects showed

that excluding individual-level random effects reduced model R^2, but did not affect the

shape or statistical significance of the smooths.

The crossbasis function, $s_2(\cdot)$ can also be written:

216
$$s_2(d_{i,t}, \ldots, d_{i,t-24}) = \sum_{l=l_0}^{L} f \cdot w(d_{i,t-l}, l)$$
(2)

where the crossbasis function, $f \cdot w(d, l)$, is composed of two marginal basis functions: the

standard predictor-response function $f(d)$, and the additional lag-response function $w(l)$.

These marginal functions are combined as a tensor product smooth such that the shape of

one marginal function varies smoothly along the other dimension (see chapter 5 of Wood

(2017) and Gasparrini et al. (2017) for more detail). Penalized cubic regression splines

were used for both marginal bases of the crossbasis function. We increased k, the

maximum basis dimension, as necessary to be sure it was adequate (Appendix A). Because

of penalization, the maximum basis dimension is generally not important as long it is large

enough to allow the smooth to represent the 'true' relationship (Wood, 2017). Estimated

degrees of freedom (edf) represent the 'true' complexity of the smooth after penalization

with edf = 1 being equivalent to a straight line and larger numbers representing more

complex curves.

The crossbasis function was fit to the data in the context of a generalized additive model

(GAM) with restricted maximum likelihood using the `mgcv` R package (Wood, 2017). The

crossbasis smooth was coded as:

232
```
s(spei_history, L, bs = "cb", xt = list(bs = "cr"))
```

Where `spei_history` is a $n \times m$ matrix with each row (n) containing the climate history of

a single plant in a year and $m = 36$ columns representing the past weather conditions

starting at the month of the census and going backwards in time 36 months. `L` is an $n \times m$

matrix with columns describing the lag structure, i.e. the integers 0 through 36. The spline

basis, `"cb"` is provided by the `dlnm` package (Gasparrini, 2011), and we specified the

marginal bases as cubic regression splines with the `xt` argument. For the full code used to

fit the models, see the Data Availability Statement.

We determined the effects of SPEI on plant growth using plant size in year t+1 as a

response variable. This was modeled with a scaled t family error distribution because

residuals were leptokurtic when a Gaussian family was used. Because number of

inflorescences was highly zero-inflated, we converted this to a binary response to model

reproduction (i.e., 1 for ≥1 inflorescence, 0 for no inflorescences). We modeled both

reproduction and survival (i.e., from year t to year t+1) using a binomial family error

distribution with a logit link function.

In the process of fitting the models, the penalty on the crossbasis smooth (and other

smoothed terms) is optimized such that more linear shapes are favored unless the data

supports non-linearity (Wood, 2017). We applied an additional penalty to shrink linear

portions toward zero with the `select=TRUE` option to the `gam()` function, and inferred

statistical significance of model terms with p-values from the `summary.gam()` function as

recommended in Marra & Wood (2011).

The `dlnm` package does not currently allow the modeling of interaction terms, which means

we could not asses the interaction of habitat type and lagged effects. We therefore fit

separate models for plants in fragments and in continuous forest to allow the shape of the

crossbasis function to differ between habitats. Significant main effects of habitat type were

assessed by looking for overlap in the 84% confidence intervals of model intercepts; the

84% CIs of two samples drawn from the same population overlap about 95% of the time

(Payton et al., 2003).

To visualize results, we plotted partial effects plots for the one-dimensional smooth

function of log size in the previous year and the two-dimensional smooth function of lagged

SPEI (the crossbasis function) for each model. To make these plots more interpretable, we

added the model intercept and (when applicable) back-transformed the y-axis to be on the

scale of the response (e.g. a probability for survival and flowering).

All analyses were conducted in R version 4.0.2 (2020-06-22) (R Core Team, 2020) using

the `targets` package for workflow management (Landau, 2021). Figures were created with

the aid of the `gratia`, `ggplot2`, and `patchwork` packages (Pedersen, 2020; Simpson, 2021;

Wickham, 2016).

Results

The meteorological droughts in our focal region indicated by SPEI are generally consistent

with those reported in the literature (Table S1). For example, the drought associated with

the 1997 El Niño Southern Oscillation (ENSO) event was one of the most severe on record

for the Amazon (Williamson et al., 2000); correspondingly, 1997 has the lowest SPEI values

in our timeseries (Figure 1d). The 2005 dry season (June–October) was also reported as an

exceptionally dry year, although this drought mostly affected the southwestern Amazon

(Marengo et al., 2008; Zeng et al., 2008). Our SPEI data show the 2005 dry season to be a

moderate drought ($-1 >$ SPEI > -1.5).

Survival, size, and flowering in continuous forest vs. fragments

Survival: Across all plots, the proportion of plants surviving was lowest in the 2003–2004

transition year (93% survival in both continuous forest and forest fragments). This

coincided with droughts in both the 2003 and 2004 rainy seasons (Figure 1b,d) and was

preceded by a drop in average plant size in the 2002–2003 transition year (i.e. negative

growth in Figure 1a). There was no significant difference in survival between continuous

forest and 1 ha fragments (Table 1). Although differences in survival probability are small

and not statistically significant, these slight differences can be seen to compound over time

in the survivorship of plants labeled in the first year of the surveys (Figure 2). Survival in

both habitats was significantly affected by size in the previous year (Table 2) with larger

plants having much greater survival probabilities (Figure 3b). The survival probability of

large plants approached 1 in both habitat types, but the smallest plants had higher survival

in 1 ha fragments (although 95% confidence intervals overlap at all sizes).

Size: Plants in continuous forest had an average of 2.9 shoots (± 1.8 SD) and were on

average 39.8 cm tall (± 26.2 SD). Plants in 1-ha fragments had on average 14% fewer

shoots (2.5 ± 1.4 SD) and were 10.5% shorter (35.6 cm ± 23.8 SD). Because our proxy for

plant size was the product of these two metrics, plants in continuous forest were on

average 32% larger than those in forest fragments (144 ± 176 SD vs. 109 ± 139 SD,

respectively), with fragments having proportionately fewer large plants (Figure 3d). Plants

were significantly larger in continuous forest on average (Table 1), and the disparity in

plant size—which was most pronounced in the initial years of our surveys—diminished

over time.

Mean plant size dropped dramatically in 2003 in both habitat types (negative fold-change

in Figure 1a), corresponding with a severe drought during the February census (SPEI = -

1.39) (Figure 1d). As with survival, size in year t was a significant predictor of size in year

t+1 (Table 2). Although plants were significantly larger in continuous forest compared to

fragments (Table 1), the effect of size in year t on size in year t+1 is nearly identical in the

two habitats (Figure 3a).

Flowering: The overall proportion of plants that flowered was very low, but the proportion

of plants flowering was 31% higher in continuous forest than 1-ha fragments (4.6%

vs. 3.5%, respectively). The intercepts for the continuous forest and 1 ha fragment models

were significantly different, indicating a main effect of habitat on flowering probability

(Table 1). The observed disparity in proportion of flowering plants was largely due to the

fact that flowering is also significantly size-dependent (Table 2), with the probability of

flowering increasing dramatically once plants reached the threshold size of about 148 (i.e.,

log(size) > 5 in Figure 3c). Despite the flowering probability of the largest plants being

greater in fragments than continuous forest, populations in fragments had proportionately

fewer plants above the reproductive size threshold (Figure 3d). The most striking

differences between habitat types coincided with relatively extreme weather conditions. In

2003, during the onset of a severe drought, the proportion of potentially reproductive

plants (i.e. plants above the size-threshold for reproduction) that actually flowered was

28% in continuous forest vs. 17% in 1-ha fragments. In 2006, following a wet spring and

winter (SPEI > 1), the trend was reversed: only 14% of these plants flowered in continuous

forest vs. 26% in 1-ha fragments (Figure 1c). Note that plots in 1-ha fragments generally

have fewer and smaller plants than those in continuous forest, so total numbers of

flowering plants was always lower in fragments.

Delayed effects of drought on demographic vital rates

Drought history had a significant effect on the survival, growth, and flowering of plants in

both habitats (Table 2). Comparing the respective crossbasis surfaces, however, reveals

that the specific climatic drivers, their timing, and their impact on individual vital rates all

differed among habitats.

Survival: For 1-ha fragments, there was a significant effect on survival of SPEI in the

preceding 0–16 months. The highest survival was near SPEI of 0, with mortality increasing

as conditions became either drier or wetter (i.e., as SPEI values became increasingly

negative or positive, respectively; Figure 4b). Additionally, a positive, lagged effect of high

SPEI at 32–36 months lag reached statistical significance. There was less of a delayed effect

of SPEI on survival in continuous forest with only the preceding 0–4 months and 32–36

months reaching statistical significance (Figure 4a). The short-term effects of SPEI on

survival in continuous forest were also unidirectional—the probability of survival was

highest in wet conditions and declined, albeit only slightly, with increasingly negative

values of SPEI (i.e., as droughts became more severe; Figure 4a). Plants in both habitat

types showed an increase in survival probability with very high SPEI values (i.e., extremely

high precipitation) at a lag time of 32–36 months. It should be noted, however, that only the

first transition year of census data (1998–1999) met these conditions. We compared the

effects of SPEI history in continuous forest and fragments by subtracting the fitted values in

Figure 4b from Figure 4a to produce Figure 4c. This shows that in average conditions (SPEI

= 0), there is little difference in survival probability between continuous forest and forest

fragments (Figure 4c). However, under extreme conditions, survival probability is up to

0.042 higher in continuous forest than fragments.

Size: The effects of drought history on trends in plant size were generally similar for

continuous forest and fragments. Under all conditions, plant size is greater in continuous

forest (Figure 5c). Decreasing SPEI at lags of 8–11 months (i.e., the end of the preceding

year's wet season) led to increased growth in both habitats. In continuous forest, but not 1-

ha fragments, SPEI at lags of 22–25 months had a significant effect on plant size in the

opposite direction, with wet conditions at that lag time resulting in larger predicted sizes.

At longer lags of 26–36 months positive effects of drought on plant size are predicted for

both continuous forest and 1-ha fragments by our models, although the area of SPEI–lag

space that reaches statistical significance is larger for continuous forest.

Flowering: Overall, the probability of flowering was significantly higher in continuous

forest than in 1-ha fragments (Figure 6, Table 1). Recent and past SPEI had less of an effect

on flowering probability in 1-ha fragments as indicated by the narrower range of the

evaluated smooth (Figure 6b). This led to some important inter-habitat differences in plant

responses to prior droughts. In continuous forests, recent drought (i.e., at lag = 0–2 with

SPEI < -1) and droughts two dry seasons prior (lags 15–20) increased the probability of

flowering. The shape of the crossbasis smooth for 1-ha fragments suggested that moderate

drought (-1.5 <SPEI< -1) in the previous 0–18 months and at lags of 28–36 months

significantly increased flowering probability slightly (the highest probability estimated is

0.004; Figure 6b). The effects of drought on flowering probability appeared stronger in

continuous forest compared to 1-ha fragments (Figure 6c). We found no evidence for a cost

of reproduction: in both forest and fragments, plants that had flowered in the previous year

were significantly more likely to be larger and flower again. Finally, with the exception of

the model for survival in 1-ha fragments, the random effect of plot was significant,

indicating that vital rates varied among plots (Table 2).

Discussion

Understanding how landscape structure and abiotic conditions act to influence population

dynamics is central to many conceptual frameworks for studying and conserving

fragmented landscapes (Didham et al., 2012; Driscoll et al., 2013). Our results support the

emerging consensus that the effects of climatic extremes on demographic vital rates can be

delayed for months or even years (Evers et al., 2021; Teller et al., 2016; Tenhumberg et al.,

2018). We also found different magnitudes, directions, and lag times of climate effects in

fragments and continuous forests. This suggests that the hypothesized synergies between

climate and fragmentation on population dynamics (Laurance & Williamson, 2001; Opdam

& Wascher, 2004; Selwood et al., 2015) may be important in this system in a way far more

complex than previously thought.

Temporal variation in demographic responses to forest fragmentation

Many studies investigating the biological consequences of habitat fragmentation on plant

growth, survival, and reproduction comprise short-term ($\leq$3 year) experiments and

observations. Our results underscore the difficulty in extrapolating long-term trends from

such short-term studies, particularly when studying long-lived organisms or when the

responses of interest can vary with size or age. For instance, one would have reached a very

different conclusion regarding the effect of fragmentation on annual survival if the study

windows were 1999–2002 (i.e., higher survival in continuous forest), 2002–2005 (i.e.,

higher survival in fragments), or 2004–2007 (i.e, no clear effect of fragmentation) (Figure

1b). It is only when evaluating over longer time windows that it becomes apparent

mortality is elevated in fragments relative to continuous forest (Figure 2), and that the

observed interannual variation is largely driven by dynamic patterns of recruitment

(Bruna, 2002) coupled with low mortality for plants beyond the smallest size classes

(Bruna, 2003).

Similarly, conclusions regarding the effects of fragmentation on flowering—which is also

both rare and size-dependent (Brooks et al., 2019)—would also differ based on the year in

which they were investigated. This could lead to erroneous extrapolations regarding the

effects of fragmentation on reproductive mutualists or population genetic structure (Côrtes

et al., 2013; Uriarte et al., 2010; Uriarte et al., 2011). Conclusions based on short-term

observations of temporally variable vital rates could lead to conservation and management

practices that are ineffective or even counterproductive, especially when when failing to

consider how the consequences of this variation might be modulated by organismal life

history (Morris et al., 2008).

It is important to emphasize, however, that the range of the estimated response to SPEI for

survival and growth was greater in fragments compared with continuous forest (Figures 4,

5). This suggests that extremes in SPEI may be more detrimental in forest fragments

compared with continuous forest. While intact forest and its canopy buffer populations

from climatic extremes, populations in fragments—especially near edges with high

contrast matrix—likely lack this protection (Didham & Lawton, 1999; Ewers & Banks-Leite,

2013). We suggest it is these climate extremes, rather than trends in average temperature,

precipitation, or SPEI (Laurance et al., 2014), that that are the causal mechanism

underlying the observed reduced plant survival and size in forest fragments.

Delayed effects of climate on demographic vital rates

Climate anomalies are known to have immediate effects on the growth, survival, or

reproduction of plants (Esteban et al., 2021; Wright & Calderon, 2006), including *Heliconia*

(Stiles, 1975; Westerband et al., 2017) and other tropical herbs (Wright, 1992). These

effects can be complex or even contradictory—mild droughts can increase the growth rates

of tropical trees and seedling survival, perhaps due to reductions in cloud cover and

concomitant increases in solar radiation (Alfaro-Sánchez et al., 2017; Condit et al., 2004;

Huete et al., 2006; Jones et al., 2014; Uriarte et al., 2018), but in severe drought years

growth can be extremely low and mortality can be sharply elevated (Connell & Green,

2000; Edwards & Krockenberger, 2006; Engelbrecht et al., 2002). There is also evidence

that the effects can persist for multiple years (Phillips et al., 2010), such as a boom in

drought-year fruit production followed by severe post-drought "famine" (Pau et al., 2013;

Wright et al., 1999).

Despite these insights, models of plant population dynamics rarely include the effects of

environmental drivers [but see Williams et al. (2015); Tenhumberg et al. (2018); Molowny-

Horas et al. (2017)). This has largely been due to the challenge (both ecologically and

statistically) of detecting any demographic responses to climatic extremes that are delayed

for multiple growing seasons. To address this, researchers have begun to use a number of

statistical methods that test for time lags in demographic responses without *a priori*

assumptions about the influence of any particular climate window (Evers et al., 2021; Ogle

et al., 2015; Teller et al., 2016; Tenhumberg et al., 2018). Our results are consistent with

this emerging literature—that the effects of precipitation extremes on the demography of

Heliconia acuminata could be delayed for up to 3 growing seasons. Additionally, our

method allowed us to capture significant non-linear responses to SPEI at different lag

times, which appear to differ in shape between habitats.

While it appears that delayed effects of climate on demographic vital rates may be

ubiquitous (Evers et al., 2021), the extent to which they vary spatially or with habitat

remains an open question. Our results suggest that they may, with habitat-specific

differences in how environmental conditions influenced future vital rates. For example,

extreme values of SPEI—both positive (unusually high precipitation) and negative

(drought conditions)—led to declines in the probability of individual survival in forest

fragments while extreme wet conditions significantly increased survival in continuous

forest. Similarly, the marginal effects of SPEI on plant size were greater in fragments,

suggesting a more pronounced effect of extreme climates in fragments. In contrast,

variation in SPEI corresponded to a greater range of flowering probabilities in continuous

forest than fragments. These results should be interpreted with some caution, however, as

the relatively low number of plants in fragments that are above the threshold-size for

flowering could limit the power to detect delayed effects.

Interestingly, we found significant detrimental effects of unusually wet conditions at some

lag times for all vital rates except survival in continuous forest and flowering probability in

1-ha fragments. While most studies investigating how precipitation extremes influence

tropical plants have focused on droughts (Lewis et al., 2011; Phillips et al., 2009;

Williamson et al., 2000), more recent work has shown that unusually wet conditions can

also be detrimental (Esteban et al., 2021). For instance, the increased cloud cover

associated with elevated precipitation could lead to reduced photosynthetic activity and

growth; plants in the resulting saturated soils could also have lower growth and elevated

mortality (Parent et al., 2008). The wind storms accompanying extreme precipitation

events (Espírito-Santo et al., 2010; Negrón-Juárez et al., 2018) could increase the likelihood

of tree-, branch-, and litter-fall, all of which are sources of mortality for understory plants

(Scariot, 2000; Ssali et al., 2019). Finally, cool temperatures associated with unusually wet

conditions may decrease flowering (Pau et al., 2013).

There are several, non-mutually-exclusive explanations for delayed effects of SPEI on

demography. The first is that the physiological processes underlying vital rates might be

initiated long before they are demographically apparent (Evers et al., 2021), and hence be

shaped by climatic events at any point in that physiological window. For example, the

flowering shoots of *Heliconia chartacea* begin to develop 6–10 months prior to the

appearance of inflorescences (Criley & Lekawatana, 1994). Adverse conditions during the 6

months following initiation, rather than the months when inflorescences are starting to

expand, leads to the aborted production of flowering shoots. Our results showed delayed

increases in flowering probability after drought. Drought conditions could be favorable for

H. accuminata flowering due to increased temperatures or decreased cloudiness associated

with droughts (Pau et al., 2013), and the effects could be delayed due to the development

time of inflorescences.

Demographic responses will also be delayed if abiotic stress causes plants to invest in

belowground rhizomes (*sensu* Pumisutapon et al., 2012). The carbohydrates stored in

rhizomes allow *Heliconia* to regenerate aboveground biomass following damage (Rundel et

al., 1998) and protect the buds that give rise to new shoots from stressful conditions

(Klimešová et al., 2018). This may be why drought led to delayed increases in growth—by

shedding shoots and leaves and investing in rhizomes, plants could be generating

proportionately more buds with which to regenerate when conditions improve. This is

consistent with the results of prior experiments in which *H. acuminata* plants transplanted

into hotter, drier fragments lost and then recovered far more leaf area than control plants

(Bruna et al., 2002).

Third, it may be that the delayed demographic effects we observed are indirectly mediated

by the effect of SPEI on other species rather than the direct effects on individual physiology

(Evers et al., 2021). For example, topical trees may not die until three or more years after a

drought (Criley & Lekawatana, 1994). When they finally do, the resulting leaf drop (Janssen

et al., 2021) and treefalls allow for light penetration to the forest understory (Canham et al.,

1990; Leitold et al., 2018), triggering a boom in the growth and flowering of understory

plants (Bruna & Oli, 2005). Similar delayed changes in the local environment could also

influence the foraging behavior of a plant's pollinators (Bruna et al., 2004; Stouffer &

Bierregaard, 1996), seed dispersers (Uriarte et al., 2011), or herbivores (Scott et al., 2021).

While more work is needed to explain why the (delayed) effects of SPEI on *H. acuminata*

survival and growth are greater in fragments than forest interiors, one hypothesis,

motivated by recent intriguing results from other systems (Sapsford et al., 2017), is that the

greater litterfall on edges (Vasconcelos & Luizão, 2004) may be altering the abundance of

pathogens or mycorrhizae.

Finally it is important to clarify the distinction between "physiological delays" and

"observational delays." For example, a Fall drought could reduce cold tolerance and

therefore overwinter survival. Alternatively the drought could kill plants immediately. If

mortality is recorded in Spring, then mortality would appear as a delayed effect of the

drought in both hypothetical cases, but only the first case is a truly delayed physiological

response to drought. The apparent delayed response in the second case is due to the timing

of the response in relation to the census date. In our analysis, this potential explanation of

"observational delays" applies only to plant size and survival, as the flowering season

coincided with the yearly census. It also only applies to lags of up to 12 months since

mortality, size, and flowering are recorded yearly. This possibility is not unique to our

study, rather it is a consequence of conducting demographic censuses on an annual scale

while the climate is quantified monthly or seasonally. For some demographic models and

follow-up experiments, knowing the precise timing of mortality might be critical; for others

less so. To disentangle possible mechanisms for observed delayed effects, it may be

necessary to conduct demographic surveys at the same temporal scale at which climate is

aggregated.

Conclusions & Future Directions

Over 24 million ha of the Brazilian Amazon have been cleared in the last two decades (Silva

Junior et al., 2021), resulting in their extensive fragmentation (Broadbent et al., 2008).

Climate models predict a future of extremes for these forests—increases in the frequency

and geographic extent of droughts, but also increases in the frequency and area affected by

periods of unusual wetness (Duffy et al., 2015). Our results support the hypothesis that

populations in Amazonian forest fragments could be more susceptible to the effects of

changing climate than those in continuous forest (Laurance et al., 2014). However, they

also indicate that the demographic responses to climate change of populations in

fragmented landscapes may be far more complex than previously appreciated. Multi-

factorial, multi-season experiments (*sensu* Aguirre et al., 2021; Bruna & Ribeiro, 2005;

Markewitz et al., 2010; Westerband et al., 2017), ideally manipulating multiple

combinations of climatic variables (Mundim & Bruna, 2016), are needed to determine how

and why habitat-specific differences in environmental conditions interact to delay the

demographic responses of plants to climatic variability. Also needed are statistical tools

that can test for synergistic effects of fragmentation and climate in vital rates, as those

currently available do not allow for including interaction terms. This also limits the ability

to include size by climate interactions in a DLNM; although plant responses to both

fragmentation and climatic extremes can be size-specific (Bruna & Oli, 2005; Schwartz et

al., 2019). The ability to identify size-specific lagged responses may be especially

complicated given size is rarely measured at the same time scale (e.g. monthly) as climate

drivers.

Recent research, including our own presented here, shows that lagged effects of climate

drivers on demography may be the norm (Evers et al., 2021). As such, we suggest that all

demographers investigate the possibility of lagged environmental effects, ideally using

methods that do not require an *a priori* choice of lag times such as DLNMs, automated

algorithms for choosing critical weather windows (van de Pol et al., 2016), or Bayesian

methods that incorporate "ecological memory" (Ogle et al., 2015).

Finally, any analytical approach for assessing lagged effects on demography requires long-

term data and/or data collected from many sites with independent weather (but similar

climate and site conditions) (Evers et al., 2021; Tenhumberg et al., 2018). Teller et al.

(2016) used a simulation study to show that detecting lagged effects required 20–25 years

of data, although this depended on effect size and the range of climate extremes in the data.

Work in progress from Compagnoni et al. (2021) showed that spatial replication may

alleviate this requirement, and in fact in our study we were able to detect significant lagged

effects with only 10 years of data and multiple plots experiencing slightly different

weather. Unfortunately, long-term data monitoring the entire life-cycle of tropical taxa are

rare, and those doing so in fragmented landscapes are virtually nonexistent (Bruna &

Ribeiro, 2005). Without investing in collecting such data, generalizations regarding the

demographic consequences of climate change in these species rich and increasingly

fragmented habitats will continue to prove elusive. More generally, however, researchers

need to consider how delayed responses to climate could influence the interpretation of

data in studies where the organisms lifespan exceeds the study's duration.

Acknowledgments

We thank Collin Edwards, Andrew Mercadante, Ellie McDaniel and 3 anonymous reviewers

for helpful discussions and comments on the manuscript. We also thank the technicians

and students who helped conduct the *Heliconia* censuses and the BDFFP and INPA for

logistical support. Financial support was provided by the U.S. National Science Foundation

(awards DEB-0614339, DEB-0614149, INT 98-06351, and DEB-1948607). This is

publication No. ___ in the BDFFP Technical Series. The authors delcare no conflicts of

interest.

Data Availability Statement

Data used in this study are available at the Dryad Digital Repository [links included upon

acceptance]; R code for analyses and visualizations are archived with Zenodo at

https://doi.org/10.5281/zenodo.5364591.

Tables

Table 1: Model intercepts and their 84% confidence intervals. Non-overlapping confidence

intervals are bolded and can be interpreted as a significant difference in the vital rate

between continuous forest and 1 ha fragments.

vital rate	habitat	intercept
survival	CF	0.97 [0.964, 0.975]
	1 ha	0.962 [0.957, 0.966]
size	CF	**4.36 [4.313, 4.408]**
	1 ha	**4.18 [4.062, 4.297]**
flowering	CF	**0.006 [0.005, 0.009]**
	1 ha	**0.002 [0.001, 0.004]**

Table 2: Marginal hypothesis tests for models of survival probability, size, and flowering

probability of *H. acuminata* in continuous forest and 1 ha fragment plots. The adjusted R^2 is

reported as a measure of model fit. The model terms included the parametric fixed effect

factor of whether plants flowered the previous year ("flowering$_t$"), the smoothed fixed

effect of plant size ("s(log(size$_t$))"), the crossbasis smooth of lagged SPEI ("s(SPEI, lag)"),

and a random effect of plot ID, ("s(plot)"). Degrees of freedom are reported for the

parametric term, and estimated degrees of freedom (edf) are reproted for smooths. Larger

values for edf indicate more complex smooths and when edf is zero the term is effectively

dropped from the model. The test statistic reported is χ^2 for survival and flowering and *F*

for log(size$_{t+1}$).

vital rate	habitat	R^2	term	(e)df	test statistic	p-value
survival	CF	0.11	flowering$_t$	1	0.14	0.709
			s(log(size$_t$))	3.46	1,957.82	<0.001

				edf	F	p
			s(SPEI, lag)	12.8	186.4	<0.001
			s(plot)	4.34	37.83	<0.001
	1 ha	0.06	flowering$_t$	1	1.64	0.200
			s(log(size$_t$))	2.84	360.41	<0.001
			s(SPEI, lag)	10.78	73.09	<0.001
			s(plot)	0	0	0.481
log(size$_{t+1}$)	CF	0.78	flowering$_t$	1	3.74	0.053
			s(log(size$_t$))	9.39	6,717.1	<0.001
			s(SPEI, lag)	15.14	95.58	<0.001
			s(plot)	4.08	7.04	<0.001
	1 ha	0.7	flowering$_t$	1	5.71	0.017
			s(log(size$_t$))	4.19	3,006.2	<0.001
			s(SPEI, lag)	18.18	27.04	<0.001
			s(plot)	2.83	23.03	<0.001
flowering$_{t+1}$	CF	0.19	flowering$_t$	1	131.59	<0.001
			s(log(size$_t$))	5.8	1,748.08	<0.001
			s(SPEI, lag)	11.45	415.26	<0.001
			s(plot)	3.95	47.54	<0.001
	1 ha	0.24	flowering$_t$	1	38.76	<0.001
			s(log(size$_t$))	3.4	321.86	<0.001
			s(SPEI, lag)	14.13	120.82	<0.001
			s(plot)	2.51	16.44	<0.001

Figures

Figure 1: (a-c) Time series of *H. acuminata* vital rates in 1-ha fragments (solid orange lines)

and continuous forest (dashed blue lines) and (d) drought occurrence in the study region.

(a) Mean fold-change in plant plant size ($\log2(\text{size}_{t+1} / \text{size}_t)$) varies by year and habitat. On

average, plants grew in most years with the notable exception in 2003, in which on average

plants regressed in size in both habitats (i.e., fold-change < 0). Error bars represent the

standard deviation. (b) The proportion of plants surviving from one transition year to the

next varied from 0.98 (CF in 1998-1999) to 0.93 (CF in 2003-2004). (c) The proportion of

H. acuminata above the size threshold for reproduction that flowered each year is on

average low but variable. The size threshold is determined by the upper 90th percentile

size of flowering plants across all years. (d) Monthly 3-month SPEI for our study region.

Gray lines represent values from different grid cells encompassing BDFFP; the dark line

represents the site mean. Colored stripes represent drought intensity: yellow = mild,

orange = moderate, dark orange = severe, red = extreme.

Figure 2: Survivorship curve for plants marked in the 1998 survey year; these plants

comprise 49% of those in the complete demographic dataset. The percentage of these

plants that were still alive ten years later was 79.7% (1629/2055) in continuous forest

vs. 72.4% (393/543) in 1-ha fragments.

Figure 3: Marginal effect of plant size in the previous census on (a) survival, (b) log(size),

and (c) flowering probability. Plots a–c were created by evaluating the smooth functions of

$log(size_t)$ at observed values, adding the model intercepts, and back-transforming (when

appropriate) to the response scale (e.g. probability for a and c). The bands depicting the

95% confidence interval include uncertainty in the intercept and uncertainty due to

smoothness selection; the smooths for 1-ha fragments and continuous forest are fit in

separate models. (d) Plant size distribution by habitat type (solid line = 1-ha fragments,

dashed line = Continuous Forest). The curves in a–c are shown with raw data

superimposed in Figure S3.

Figure 4: Marginal effect of lagged SPEI on *H. acuminata* survival in (a) continuous forest,

(b) 1-ha fragments, (c) and the difference between the two. Outlines show combinations of

SPEI and lag time that have a significant effect on survival, defined as areas where the 95%

confidence interval around the response does not overlap the intercept. The bar on the

bottom of each panel indicates wet seasons (black, November–May) and dry seasons

(white, June–October). For a and b, the model intercepts were added to the evaluated

crossbasis smooths and values were back-transformed to the response scale

(i.e. probabilities). Areas of the fitted smooth far from observed values (i.e. combinations of

lag time and SPEI) are shown in grey.

Figure 5: Marginal effect of lagged SPEI on *H. acuminata* size in (a) continuous forest, (b) 1-

ha fragments, (c) and the difference between the two. Outlines show combinations of SPEI

and lag time that have a significant effect on plant size, defined as areas where the 95%

confidence interval around the response does not overlap the intercept. The bar on the

bottom of each panel indicates wet seasons (black, November–May) and dry seasons

634   (white, June–October). For a and b, the model intercepts were added to the evaluated

635   crossbasis smooths. Areas of the fitted smooth far from observed values (i.e. combinations

636   of lag time and SPEI) are shown in grey.

638   Figure 6: Marginal effect of lagged SPEI on *H. acuminata* flowering probability in (a)

639   continuous forest, (b) 1-ha fragments, (c) and the difference between the two. Outlines

show combinations of SPEI and lag time that have a significant effect on probability of

flowering, defined as areas where the 95% confidence interval around the response does

not overlap the intercept. The bar on the bottom of each panel indicates wet seasons (black,

November–May) and dry seasons (white, June–October). For a and b, the model intercepts

were added to the evaluated crossbasis smooths and values were back-transformed to the

response scale (i.e. probabilities). Areas of the fitted smooth far from observed values

(i.e. combinations of lag time and SPEI) are shown in grey.

References

Aguirre, B. A., Hsieh, B., Watson, S. J., & Wright, A. J. (2021). The experimental manipulation of atmospheric drought: Teasing out the role of microclimate in biodiversity experiments. *Journal of Ecology*, *109*(5), 1986–1999. https://doi.org/https://doi.org/10.1111/1365-2745.13595

Alfaro-Sánchez, R., Muller-Landau, H. C., Wright, S. J., & Camarero, J. J. (2017). Growth and reproduction respond differently to climate in three Neotropical tree species. *Oecologia*, *184*(2), 531–541. https://doi.org/10.1007/s00442-017-3879-3

Alroy, J. (2017). Effects of habitat disturbance on tropical forest biodiversity. *Proceedings of the National Academy of Sciences*, *114*(23), 6056–6061. https://doi.org/10.1073/pnas.1611855114

Anderegg, W. R. L., Schwalm, C., Biondi, F., Camarero, J. J., Koch, G., Litvak, M., Ogle, K., Shaw, J. D., Shevliakova, E., Williams, A. P., Wolf, A., Ziaco, E., & Pacala, S. (2015). Pervasive drought legacies in forest ecosystems and their implications for carbon cycle models. *Science*, *349*(6247), 528–532. https://doi.org/10.1126/science.aab1833

Arroyo-Rodríguez, V., Saldaña-Vázquez, R. A., Fahrig, L., & Santos, B. A. (2017). Does forest fragmentation cause an increase in forest temperature? *Ecological Research*, *32*(1), 81–88. https://doi.org/10.1007/s11284-016-1411-6

Beguería, S., & Vicente-Serrano, S. M. (2017). *SPEI: Calculation of the standardised precipitation-evapotranspiration index*. https://CRAN.R-project.org/package=SPEI

Betts, M. G., Wolf, C., Pfeifer, M., Banks-Leite, C., Arroyo-Rodríguez, V., Ribeiro, D. B., Barlow, J., Eigenbrod, F., Faria, D., Fletcher, R. J., Hadley, A. S., Hawes, J. E., Holt, R. D., Klingbeil, B., Kormann, U., Lens, L., Levi, T., Medina-Rangel, G. F., Melles, S. L., … Ewers, R. M. (2019). Extinction filters mediate the global effects of habitat fragmentation on animals. *Science*, *366*(6470), 1236–1239. https://doi.org/10.1126/science.aax9387

Bianchi, C. A., & Haig, S. M. (2013). Deforestation trends of tropical dry forests in Central Brazil. *Biotropica*, *45*(3), 395–400. https://doi.org/https://doi.org/10.1111/btp.12010

Bierregaard, R. O., Gascon, C., Lovejoy, T. E., & Mesquita, R. (Eds.). (2001). *Lessons from Amazonia: The ecology and conservation of a fragmented forest*. Yale University Press.

Bierregaard, R. O., Lovejoy, T. E., Kapos, V., & Hutchings, R. W. (1992). The biological dynamics of tropical rainforest fragments. *BioScience*, *42*(11), 859–866. https://doi.org/10.2307/1312085

Broadbent, E. N., Asner, G. P., Keller, M., Knapp, D. E., Oliveira, P. J. C., & Silva, J. N. (2008). Forest fragmentation and edge effects from deforestation and selective logging in the Brazilian Amazon. *Biological Conservation*, *141*, 1745–1757. https://doi.org/10.1016/j.biocon.2008.04.024

Brodie, J., Post, E., & Laurance, W. F. (2012). Climate change and tropical biodiversity: A new focus. *Trends in Ecology & Evolution*, *27*(3), 145–150. https://doi.org/10.1016/j.tree.2011.09.008

Brooks, M. E., Kristensen, K., Darrigo, M. R., Rubim, P., Uriarte, M., Bruna, E. M., & Bolker, B. M. (2019). Statistical modeling of patterns in annual reproductive rates. *Ecology*, *100*(7). https://doi.org/10.1002/ecy.2706

Bruna, E. M. (1999). Seed germination in rainforest fragments. *Nature*, *402*(6758), 139. https://doi.org/10.1038/45963

Bruna, E. M. (2002). Effects of forest fragmentation on *Heliconia acuminata* seedling recruitment in central Amazonia. *Oecologia*, *132*(2), 235–243. https://doi.org/10.1007/s00442-002-0956-y

Bruna, E. M. (2003). Are plant populations in fragmented habitats recruitment limited? Tests with an Amazonian herb. *Ecology*, *84*(4), 932–947. https://doi.org/10.1890/0012-9658(2003)084[0932:APPIFH]2.0.CO;2

Bruna, E. M., Fiske, I. J., & Trager, M. D. (2009). Habitat fragmentation and plant populations: Is what we know demographically irrelevant? *Journal of Vegetation Science*, *20*(3), 569–576. https://doi.org/10.1111/j.1654-1103.2009.01060.x

Bruna, E. M., & Kress, W. J. (2002). Habitat fragmentation and the demographic structure of an Amazonian understory herb (*Heliconia acuminata*). *Conservation Biology*, *16*(5), 1256–1266. https://doi.org/10.1046/j.1523-1739.2002.99494.x

Bruna, E. M., Kress, W. J., Marques, F., & Silva, O. F. da. (2004). *Heliconia acuminata* reproductive success is independent of local floral density. *Acta Amazonica*, *34*(3), 467–471. https://doi.org/10.1590/S0044-59672004000300012

Bruna, E. M., Nardy, O., Strauss, S. Y., & Harrison, S. (2002). Experimental assessment of *Heliconia acuminata* growth in a fragmented Amazonian landscape. *Journal of Ecology*, *90*(4), 639–649. https://doi.org/10.1046/j.1365-2745.2002.00707.x

Bruna, E. M., & Oli, M. K. (2005). Demographic effects of habitat fragmentation on a tropical
herb: Life-table response experiments. *Ecology*, *86*(7), 1816–1824.
https://doi.org/10.1890/04-1716

Bruna, E. M., & Ribeiro, M. B. N. (2005). The compensatory responses of an understory herb
to experimental damage are habitat-dependent. *American Journal of Botany*, *92*(12), 2101–
2210. https://doi.org/10.3732/ajb.92.12.2101

Canham, C. D., Denslow, J. S., Platt, W. J., Runkle, J. R., Spies, T. A., & White, P. S. (1990). Light
regimes beneath closed canopies and tree-fall gaps in temperate and tropical forests.
Canadian Journal of Forest Research, *20*(5), 620–631. https://doi.org/10.1139/x90-084

Compagnoni, A., Evers, S., & Knight, T. M. (2021). Spatial replication provides substantial
opportunities to estimate plant population responses to temporal climatic variation.
Ecological Society of America Annual Meeting.

Condit, R., Aguilar, S., Hernandez, A., Perez, R., Lao, S., Angehr, G., Hubbell, S. P., & Foster, R.
B. (2004). Tropical forest dynamics across a rainfall gradient and the impact of an El Niño
dry season. *Journal of Tropical Ecology*, *20*(01), 51–72.
https://doi.org/10.1017/S0266467403001081

Connell, J. H., & Green, P. T. (2000). Seedling dynamics over thirty-two years in a tropical
rain forest tree. *Ecology*, *81*(2), 568–584.

Côrtes, M. C., Uriarte, M., Lemes, M. R., Gribel, R., Kress, W. J., Smouse, P. E., & Bruna, E. M.
(2013). Low plant density enhances gene dispersal in the Amazonian understory herb
Heliconia Acuminata. *Molecular Ecology*, *22*(22), 5716–5729.
https://doi.org/10.1111/mec.12495

Criley, R., & Lekawatana, S. (1994). Year around production with high yields may be a
possibility for *Heliconia chartacea*. *Acta Horticulturae*, *397*, 95–102.
https://doi.org/10.17660/ActaHortic.1995.397.7

Crone, E. E., Menges, E. S., Ellis, M. M., Bell, T., Bierzychudek, P., Ehrlen, J., Kaye, T. N., Knight,
T. M., Lesica, P., Morris, W. F., Oostermeijer, G., Quintana-Ascencio, P. F., Stanley, A., Ticktin,
T., Valverde, T., & Williams, J. L. (2011). How do plant ecologists use matrix population
models? *Ecology Letters*, *14*(1), 1–8.

da Silva, J. M. C., & Tabarelli, M. (2000). Tree species impoverishment and the future flora of
the Atlantic forest of northeast Brazil. *Nature*, *404*(6773), 72–74.
https://doi.org/10.1038/35003563

Didham, R. K., Kapos, V., & Ewers, R. M. (2012). Rethinking the conceptual foundations of
habitat fragmentation research. *Oikos*, *121*(2), 161–170. https://doi.org/10.1111/j.1600-
0706.2011.20273.x

Didham, R. K., & Lawton, J. H. (1999). Edge structure determines the magnitude of changes
in microclimate and vegetation structure in tropical forest fragments. *Biotropica*, *31*(1), 17–
30. https://doi.org/10.1111/j.1744-7429.1999.tb00113.x

Doak, D. F., & Morris, W. F. (2010). Demographic compensation and tipping points in
climate-induced range shifts. *Nature*, *467*(7318), 959–962.

Driscoll, D. A., Banks, S. C., Barton, P. S., Lindenmayer, D. B., & Smith, A. L. (2013).
Conceptual domain of the matrix in fragmented landscapes. *Trends in Ecology & Evolution*,
28(10), 605–613. https://doi.org/10.1016/j.tree.2013.06.010

Duffy, P. B., Brando, P., Asner, G. P., & Field, C. B. (2015). Projections of future
meteorological drought and wet periods in the Amazon. *Proceedings of the National
Academy of Sciences*, *112*(43), 13172–13177. https://doi.org/10.1073/pnas.1421010112

Edwards, W., & Krockenberger, A. (2006). Seedling mortality due to drought and fire
associated with the 2002 El Niño event in a tropical rain forest in north-east Queensland,
Australia. *Biotropica*, *38*(1), 16–26. https://doi.org/10.1111/j.1744-7429.2006.00098.x

Ehrlen, J., Morris, W. F., von Euler, T., & Dahlgren, J. P. (2016). Advancing environmentally
explicit structured population models of plants. *Journal of Ecology*, *104*(2), 292–305.
https://doi.org/10.1111/1365-2745.12523

Engelbrecht, B. M. J., Wright, S. J., & De Steven, D. (2002). Survival and ecophysiology of tree
seedlings during El Niño drought in a tropical moist forest in Panama. *Journal of Tropical
Ecology*, *18*, 569–579.

Espírito-Santo, F. D. B., Keller, M., Braswell, B., Nelson, B. W., Frolking, S., & Vicente, G.
(2010). Storm intensity and old-growth forest disturbances in the Amazon region.
Geophysical Research Letters, *37*(11). https://doi.org/10.1029/2010GL043146

Esquivel-Muelbert, A., Baker, T. R., Dexter, K. G., Lewis, S. L., Brienen, R. J. W., Feldpausch, T.
R., Lloyd, J., Monteagudo-Mendoza, A., Arroyo, L., Álvarez-Dávila, E., Higuchi, N., Marimon, B.
S., Marimon-Junior, B. H., Silveira, M., Vilanova, E., Gloor, E., Malhi, Y., Chave, J., Barlow, J., …
Phillips, O. L. (2019). Compositional response of Amazon forests to climate change. *Global
Change Biology*, *25*(1), 39–56. https://doi.org/10.1111/gcb.14413

Esteban, E. J. L., Castilho, C. V., Melgaço, K. L., & Costa, F. R. C. (2021). The other side of
droughts: Wet extremes and topography as buffers of negative drought effects in an
Amazonian forest. *New Phytologist*, *229*(4), 1995–2006.
https://doi.org/10.1111/nph.17005

Evers, S. M., Knight, T. M., Inouye, D. W., Miller, T. E. X., Salguero-Gómez, R., Iler, A. M., &
Compagnoni, A. (2021). Lagged and dormant season climate better predict plant vital rates
than climate during the growing season. *Global Change Biology*, *27*(9), 1927–1941.
https://doi.org/10.1111/gcb.15519

Ewers, R. M., & Banks-Leite, C. (2013). Fragmentation impairs the microclimate buffering
effect of tropical forests. *PLoS ONE*, *8*(3), e58093.
https://doi.org/10.1371/journal.pone.0058093

Feeley, K. J., Malhi, Y., Zelazowski, P., & Silman, M. R. (2012). The relative importance of
deforestation, precipitation change, and temperature sensitivity in determining the future

distributions and diversity of Amazonian plant species. *Global Change Biology*, *18*(8), 2636–2647. https://doi.org/10.1111/j.1365-2486.2012.02719.x

Gagnon, P. R., Bruna, E. M., Rubim, P., Darrigo, M. R., Littell, R. C., Uriarte, M., & Kress, W. J. (2011). Growth of an understory herb is chronically reduced in Amazonian forest fragments. *Biological Conservation*, *144*(2), 830–835. https://doi.org/10.1016/j.biocon.2010.11.015

Gaoue, O. G., Horvitz, C. C., Steiner, U. K., & Tuljapurkar, S. (2019). Climate, rather than human disturbance, is the main driver of age-specific mortality trajectories in a tropical tree. *Ecological Modelling*, *400*, 34–40. https://doi.org/10.1016/j.ecolmodel.2019.03.007

Gasparrini, A. (2011). Distributed lag linear and non-linear models in R: The package dlnm. *Journal of Statistical Software*, *43*(8), 1–20. https://doi.org/10.18637/jss.v043.i08

Gasparrini, A., Scheipl, F., Armstrong, B., & Kenward, M. G. (2017). A penalized framework for distributed lag non-linear models: Penalized DLNMs. *Biometrics*, *73*(3), 938–948. https://doi.org/10.1111/biom.12645

Gentry, A. H., & Dodson, C. (1987). Contribution of nontrees to species richness of a tropical rain forest. *Biotropica*, *19*(2), 149–156. https://doi.org/10.2307/2388737

González-M, R., Posada, J. M., Carmona, C. P., Garzón, F., Salinas, V., Idárraga-Piedrahita, Á., Pizano, C., Avella, A., López-Camacho, R., Norden, N., Nieto, J., Medina, S. P., Rodríguez-M, G. M., Franke-Ante, R., Torres, A. M., Jurado, R., Cuadros, H., Castaño-Naranjo, A., García, H., & Salgado-Negret, B. (2020). Diverging functional strategies but high sensitivity to an extreme drought in tropical dry forests. *Ecology Letters*, *n/a*(n/a). https://doi.org/10.1111/ele.13659

Haddad, N. M., Brudvig, L. A., Clobert, J., Davies, K. F., Gonzalez, A., Holt, R. D., Lovejoy, T. E., Sexton, J. O., Austin, M. P., Collins, C. D., Cook, W. M., Damschen, E. I., Ewers, R. M., Foster, B. L., Jenkins, C. N., King, A. J., Laurance, W. F., Levey, D. J., Margules, C. R., … Townshend, J. R. (2015). Habitat fragmentation and its lasting impact on Earth's ecosystems. *Science Advances*, *1*(2), e1500052. https://doi.org/10.1126/sciadv.1500052

Holyoak, M., & Heath, S. K. (2016). The integration of climate change, spatial dynamics, and habitat fragmentation: A conceptual overview. *Integrative Zoology*, *11*(1), 40–59. https://doi.org/https://doi.org/10.1111/1749-4877.12167

Honnay, O., Verheyen, K., Butaye, J., Jacquemyn, H., Bossuyt, B., & Hermy, M. (2002). Possible effects of habitat fragmentation and climate change on the range of forest plant species. *Ecology Letters*, *5*(4), 525–530. https://doi.org/10.1046/j.1461-0248.2002.00346.x

Huete, A. R., Didan, K., Shimabukuro, Y. E., Ratana, P., Saleska, S. R., Hutyra, L. R., Yang, W., Nemani, R. R., & Myneni, R. (2006). Amazon rainforests green-up with sunlight in dry season. *Geophysical Research Letters*, *33*(6). https://doi.org/10.1029/2005GL025583

Janssen, T., van der Velde, Y., Hofhansl, F., Luyssaert, S., Naudts, K., Driessen, B., Fleischer,
K., & Dolman, H. (2021). Drought effects on leaf fall, leaf flushing and stem growth in
Neotropical forest; reconciling remote sensing data and field observations. *Biogeosciences*
Discussions, 1–41. https://doi.org/10.5194/bg-2021-30

Jones, M. O., Kimball, J. S., & Nemani, R. R. (2014). Asynchronous Amazon forest canopy
phenology indicates adaptation to both water and light availability. *Environmental Research*
Letters, *9*(12), 124021. https://doi.org/10.1088/1748-9326/9/12/124021

Kannenberg, S. A., Schwalm, C. R., & Anderegg, W. R. L. (2020). Ghosts of the past: How
drought legacy effects shape forest functioning and carbon cycling. *Ecology Letters*, *23*(5),
891–901. https://doi.org/10.1111/ele.13485

Klimešová, J., Martínková, J., & Ottaviani, G. (2018). Belowground plant functional ecology:
Towards an integrated perspective. *Functional Ecology*, *32*(9), 2115–2126.
https://doi.org/10.1111/1365-2435.13145

Kress, W. J. (1990). The diversity and distribution of *Heliconia* (Heliconiaceae) in Brazil.
Acta Botanica Brasileira, *4*(1), 159–167.

Landau, W. M. (2021). The targets R package: A dynamic Make-like function-oriented
pipeline toolkit for reproducibility and high-performance computing. *Journal of Open*
Source Software, *6*(57), 2959. https://doi.org/ghwrqs

Laurance, W. F., Andrade, A. S., Magrach, A., Camargo, J. L. C., Campbell, M., Fearnside, P. M.,
Edwards, W., Valsko, J. J., Lovejoy, T. E., & Laurance, S. G. (2014). Apparent environmental
synergism drives the dynamics of Amazonian forest fragments. *Ecology*, *95*(11), 3018–
3026. https://doi.org/10.1890/14-0330.1

Laurance, W. F., Ferreira, L. V., Rankin de Merona, J. M., & Laurance, S. G. (1998). Rain forest
fragmentation and the dynamics of Amazonian tree communities. *Ecology*, *79*(6), 2032–
2040.

Laurance, W. F., Nascimento, H. E. M., Laurance, S. G., Andrade, A., Ribeiro, J. E. L. S., Giraldo,
J. P., Lovejoy, T. E., Condit, R., Chave, J., Harms, K. E., & D'Angelo, S. (2006). Rapid decay of
tree-community composition in Amazonian forest fragments. *Proceedings of the National*
Academy of Sciences, *103*(50), 19010–19014. https://doi.org/10.1073/pnas.0609048103

Laurance, W. F., & Williamson, G. B. (2001). Positive feedbacks among forest fragmentation,
drought, and climate change in the Amazon. *Conservation Biology*, *15*(6), 1529–1535.
https://doi.org/10.1046/j.1523-1739.2001.01093.x

Laurance, W. F., Williamson, G. B., Delamônica, P., Oliveira, A., Lovejoy, T. E., Gascon, C., &
Pohl, L. (2001). Effects of a strong drought on Amazonian forest fragments and edges.
Journal of Tropical Ecology, *17*(6), 771–785.

Leitold, V., Morton, D. C., Longo, M., dos-Santos, M. N., Keller, M., & Scaranello, M. (2018). El
Niño drought increased canopy turnover in Amazon forests. *New Phytologist*, *219*(3), 959–
971. https://doi.org/10.1111/nph.15110

Lewis, S. L., Brando, P. M., Phillips, O. L., Heijden, G. M. F. van der, & Nepstad, D. (2011). The
2010 Amazon Drought. *Science*, *331*(6017), 554–554.
https://doi.org/10.1126/science.1200807

Marengo, J. A., Nobre, C. A., Tomasella, J., Oyama, M. D., Sampaio de Oliveira, G., de Oliveira,
R., Camargo, H., Alves, L. M., & Brown, I. F. (2008). The drought of Amazonia in 2005.
Journal of Climate, *21*(3), 495–516. https://doi.org/10.1175/2007JCLI1600.1

Markewitz, D., Devine, S., Davidson, E. A., Brando, P., & Nepstad, D. C. (2010). Soil moisture
depletion under simulated drought in the Amazon: impacts on deep root uptake. *New
Phytologist*, *187*(3), 592–607. https://doi.org/10.1111/j.1469-8137.2010.03391.x

Marra, G., & Wood, S. N. (2011). Practical variable selection for generalized additive models.
Computational Statistics & Data Analysis, *55*(7), 2372–2387.
https://doi.org/10.1016/j.csda.2011.02.004

McKee, T. B., Doesken, N. J., Kleist, J., & others. (1993). The relationship of drought
frequency and duration to time scales. *Proceedings of the 8th Conference on Applied
Climatology*, *17*, 179–183.

McPhaden, M. J. (1999). The child prodigy of 1997-98. *Nature*, *398*(6728), 559–561.
https://doi.org/10.1038/19193

Molowny-Horas, R., Suarez, M. L., & Lloret, F. (2017). Changes in the natural dynamics of
Nothofagus Dombeyi forests: Population modeling with increasing drought frequencies.
Ecosphere, *8*(3), e01708. https://doi.org/10.1002/ecs2.1708

Morris, W. F., & Doak, D. F. (2002). *Quantitative conservation biology: Theory and practice of
population viability analysis.* Sinauer.

Morris, W. F., Pfister, C. A., Tuljapurkar, S., Haridas, C. V., Boggs, C. L., Boyce, M. S., Bruna, E.
M., Church, D. R., Coulson, T., Doak, D. F., Forsyth, S., Gaillard, J.-M., Horvitz, C. C., Kalisz, S.,
Kendall, B. E., Knight, T. M., Lee, C. T., & Menges, E. S. (2008). Longevity can buffer plant and
animal populations against changing climatic variability. *Ecology*, *89*(1), 19–25.
https://doi.org/10.1890/07-0774.1

Mundim, F. M., & Bruna, E. M. (2016). Is there a temperate bias in our understanding of how
climate change will alter plant–herbivore interactions? A meta- analysis of experimental
studies. *American Naturalist*, *188*, S74–S89.

Nakazono, E. M., Bruna, E. M., & Mesquita, R. C. G. (2004). Experiemental harvesting of the
non-timber forest product *Ischnosiphon polyphyllus* in central Amazonia. *Forest Ecology and
Management*, *190*(2-3), 219–225. https://doi.org/10.1016/j.foreco.2003.10.013

Negrón-Juárez, R. I., Holm, J. A., Marra, D. M., Rifai, S. W., Riley, W. J., Chambers, J. Q., Koven,
C. D., Knox, R. G., McGroddy, M. E., Vittorio, A. V. D., Urquiza-Muñoz, J., Tello-Espinoza, R.,
Muñoz, W. A., Ribeiro, G. H. P. M., & Higuchi, N. (2018). Vulnerability of Amazon forests to
storm-driven tree mortality. *Environmental Research Letters*, *13*(5), 054021.
https://doi.org/10.1088/1748-9326/aabe9f

Ogle, K., Barber, J. J., Barron-Gafford, G. A., Bentley, L. P., Young, J. M., Huxman, T. E., Loik, M.
E., & Tissue, D. T. (2015). Quantifying ecological memory in plant and ecosystem processes.
Ecology Letters, *18*(3), 221–235. https://doi.org/10.1111/ele.12399

Oliver, T. H., Marshall, H. H., Morecroft, M. D., Brereton, T., Prudhomme, C., & Huntingford, C.
(2015). Interacting effects of climate change and habitat fragmentation on drought-
sensitive butterflies. *Nature Climate Change*, *5*(10), 941–945.
https://doi.org/10.1038/nclimate2746

Opdam, P., & Wascher, D. (2004). Climate change meets habitat fragmentation: Linking
landscape and biogeographical scale levels in research and conservation. *Biological*
Conservation, *117*(3), 285–297.

Paniw, M., James, T. D., Ruth Archer, C., Römer, G., Levin, S., Compagnoni, A., Che-Castaldo, J.,
Bennett, J. M., Mooney, A., Childs, D. Z., Ozgul, A., Jones, O. R., Burns, J. H., Beckerman, A. P.,
Patwary, A., Sanchez-Gassen, N., Knight, T. M., & Salguero-Gómez, R. (2021). The myriad of
complex demographic responses of terrestrial mammals to climate change and gaps of
knowledge: A global analysis. *Journal of Animal Ecology*. https://doi.org/10.1111/1365-
2656.13467

Parent, C., Nicolas, C., Audrey, B., Crevècoeur, M., & Dat, J. (2008). An overview of plant
responses to soil waterlogging. *Plant Stress*, *2*(1), 20–27.

Pau, S., Wolkovich, E. M., Cook, B. I., Nytch, C. J., Regetz, J., Zimmerman, J. K., & Joseph
Wright, S. (2013). Clouds and temperature drive dynamic changes in tropical flower
production. *Nature Climate Change*, *3*(9), 838–842. https://doi.org/10.1038/nclimate1934

Payton, M. E., Greenstone, M. H., & Schenker, N. (2003). Overlapping confidence intervals or
standard error intervals: What do they mean in terms of statistical significance? *Journal of*
Insect Science, *3*(34), 1–6. https://doi.org/10.1673/031.003.3401

Pedersen, T. L. (2020). *Patchwork: The composer of plots* [Manual].

Phillips, O. L., Aragao, L. E. O. C., Lewis, S. L., Fisher, J. B., Lloyd, J., Lopez-Gonzalez, G., Malhi,
Y., Monteagudo, A., Peacock, J., Quesada, C. A., van der Heijden, G., Almeida, S., Amaral, I.,
Arroyo, L., Aymard, G., Baker, T. R., Banki, O., Blanc, L., Bonal, D., … Torres-Lezama, A.
(2009). Drought sensitivity of the Amazon rainforest. *Science*, *323*(5919), 1344–1347.
https://doi.org/10.1126/science.1164033

Phillips, O. L., van der Heijden, G., Lewis, S. L., López-González, G., Aragão, L. E. O. C., Lloyd, J.,
Malhi, Y., Monteagudo, A., Almeida, S., Dávila, E. A., Amaral, I., Andelman, S., Andrade, A.,
Arroyo, L., Aymard, G., Baker, T. R., Blanc, L., Bonal, D., de Oliveira, Á. C. A., … Vilanova, E.
(2010). Drought–mortality relationships for tropical forests. *New Phytologist*, *187*(3), 631–
646. https://doi.org/10.1111/j.1469-8137.2010.03359.x

Pumisutapon, P., Visser, R. G. F., & de Klerk, G.-J. (2012). Moderate abiotic stresses increase
rhizome growth and outgrowth of axillary buds in *Alstroemeria* cultured in vitro. *Plant Cell,*

Tissue and Organ Culture (PCTOC), *110*(3), 395–400. https://doi.org/10.1007/s11240-012-0160-7

R Core Team. (2020). *R: A language and environment for statistical computing*.

Rankin-de-Mérona, J. M., Prance, G. T., Hutchings, R. W., da Silva, M. F., Rodrigues, W. A., & Uehling, M. E. (1992). Preliminary results of a large-scale tree inventory of upland rain forest in the central Amazon. *Acta Amazonica*, *22*(4), 493–534. https://doi.org/10.1590/1809-43921992224534

Ribeiro, M. B. N., Bruna, E. M., & Mantovani, W. (2010). Influence of post-clearing treatment on the recovery of herbaceous plant communities in Amazonian secondary forests. *Restoration Ecology*, *18*, 50–58.

Rundel, P. W., Cooley, A. M., Gerst, K. L., Riordan, E. C., Sharifi, M. R., Sun, J. W., & Tower, J. A. (2020). Functional traits of broad-leaved monocot herbs in the understory and forest edges of a Costa Rican rainforest. *PeerJ*, *8*, e9958. https://doi.org/10.7717/peerj.9958

Rundel, P. W., Sharifi, M. R., Gibson, A. C., & Esler, K. J. (1998). Structural and physiological adaptation to light environments in Neotropical *Heliconia* (Heliconiaceae). *Journal of Tropical Ecology*, *14*(6), 789–801. https://doi.org/10.1017/S0266467498000571

Salguero-Gómez, R., Jones, O. R., Archer, C. R., Buckley, Y. M., Che-Castaldo, J., Caswell, H., Hodgson, D., Scheuerlein, A., Conde, D. A., Brinks, E., Buhr, H., Farack, C., Gottschalk, F., Hartmann, A., Henning, A., Hoppe, G., Römer, G., Runge, J., Ruoff, T., … Vaupel, J. W. (2015). The COMPADRE Plant Matrix Database: An open online repository for plant demography. *Journal of Ecology*, *103*(1), 202–218. https://doi.org/10.1111/1365-2745.12334

Sapsford, S. J., Paap, T., Hardy, G. E. St. J., & Burgess, T. I. (2017). The 'chicken or the egg': Which comes first, forest tree decline or loss of mycorrhizae? *Plant Ecology*, *218*(9), 1093–1106. https://doi.org/10.1007/s11258-017-0754-6

Scariot, A. (1999). Forest fragmentation effects on palm diversity in central Amazonia. *Journal of Ecology*, *87*(1), 66–76. https://doi.org/10.1046/j.1365-2745.1999.00332.x

Scariot, A. (2000). Seedling Mortality by Litterfall in Amazonian Forest Fragments. *Biotropica*, *32*, 662–669. https://doi.org/10.1111/j.1744-7429.2000.tb00513.x

Scheffers, B. R., Edwards, D. P., Macdonald, S. L., Senior, R. A., Andriamahohatra, L. R., Roslan, N., Rogers, A. M., Haugaasen, T., Wright, P., & Williams, S. E. (2017). Extreme thermal heterogeneity in structurally complex tropical rain forests. *Biotropica*, *49*(1), 35–44. https://doi.org/10.1111/btp.12355

Schwalm, C. R., Anderegg, W. R. L., Michalak, A. M., Fisher, J. B., Biondi, F., Koch, G., Litvak, M., Ogle, K., Shaw, J. D., Wolf, A., Huntzinger, D. N., Schaefer, K., Cook, R., Wei, Y., Fang, Y., Hayes, D., Huang, M., Jain, A., & Tian, H. (2017). Global patterns of drought recovery. *Nature*, *548*(7666), 202–205. https://doi.org/10.1038/nature23021

Schwartz, N. B., Budsock, A. M., & Uriarte, M. (2019). Fragmentation, forest structure, and topography modulate impacts of drought in a tropical forest landscape. *Ecology*, *100*(6), e02677. https://doi.org/10.1002/ecy.2677

Scott, E. R., Wei, J.-P., Li, X., Han, W.-Y., & Orians, C. M. (2021). Differing non-linear, lagged effects of temperature and precipitation on an insect herbivore and its host plant. *Ecological Entomology*, *early view*. https://doi.org/10.1111/een.13023

Selwood, K. E., McGeoch, M. A., & Nally, R. M. (2015). The effects of climate change and land-use change on demographic rates and population viability. *Biological Reviews*, *90*(3), 837–853. https://doi.org/10.1111/brv.12136

Silva Junior, C. H. L., Pessôa, A. C. M., Carvalho, N. S., Reis, J. B. C., Anderson, L. O., & Aragão, L. E. O. C. (2021). The Brazilian Amazon deforestation rate in 2020 is the greatest of the decade. *Nature Ecology & Evolution*, *5*(2), 144–145. https://doi.org/10.1038/s41559-020-01368-x

Simpson, G. L. (2021). *Gratia: Graceful 'ggplot'-Based graphics and other functions for GAMs fitted using 'mgcv'* [Manual].

Sletvold, N. (2005). Density-dependent growth and survival in a natural population of the facultative biennial *Digitalis purpurea*. *Journal of Ecology*, *93*(4), 727–736. https://doi.org/10.1111/j.1365-2745.2005.01008.x

Snow, D. W. (1981). Tropical frugivorous birds and their food plants - a world survey. *Biotropica*, *13*(1), 1–14. https://doi.org/10.2307/2387865

Ssali, F., Moe, S. R., & Sheil, D. (2019). Damage to artificial seedlings across a disturbed Afromontane forest landscape. *Biotropica*, *51*(5), 652–663. https://doi.org/https://doi.org/10.1111/btp.12703

Stiles, F. G. (1975). Ecology, flowering phenology, and hummingbird pollination of some Costa Rican *Heliconia* species. *Ecology*, *56*(2), 285–301. https://doi.org/10.2307/1934961

Stouffer, P. C., & Bierregaard, R. O. (1996). Forest fragmentation and seasonal patterns of hummingbird abundance in Amazonian Brazil. *Ararajuba*, *4*(1), 9–14.

Taubert, F., Fischer, R., Groeneveld, J., Lehmann, S., Müller, M. S., Rödig, E., Wiegand, T., & Huth, A. (2018). Global patterns of tropical forest fragmentation. *Nature*, *554*(7693), 519–522. https://doi.org/10.1038/nature25508

Teller, B. J., Adler, P. B., Edwards, C. B., Hooker, G., & Ellner, S. P. (2016). Linking demography with drivers: Climate and competition. *Methods in Ecology and Evolution*, *7*(2), 171–183. https://doi.org/10.1111/2041-210X.12486

Tenhumberg, B., Crone, E. E., Ramula, S., & Tyre, A. J. (2018). Time-lagged effects of weather on plant demography: Drought and *Astragalus scaphoides*. *Ecology*, *99*(4), 915–925. https://doi.org/10.1002/ecy.2163

Ticktin, T. (2003). Relationships between El Niño Southern Oscillation and demographic
patterns in a substitute food for collared peccaries in Panama. *Biotropica*, *35*(2), 189–197.
https://doi.org/10.1111/j.1744-7429.2003.tb00278.x

Uriarte, M., Anciães, M., da Silva, M. T. B., Rubim, R., Johnson, E., & Bruna, E. M. (2011).
Disentangling the drivers of reduced long-distance seed dispersal by birds in an
experimentally fragmented landscape. *Ecology*, *92*(4), 924–937.
https://doi.org/10.1890/10-0709.1

Uriarte, M., Bruna, E. M., Rubim, P., Anciães, M., & Jonckheere, I. (2010). Effects of forest
fragmentation on the seedling recruitment of a tropical herb: assessing seed vs. safe-site
limitation. *Ecology*, *91*(5), 1317–1328. https://doi.org/10.1890/09-0785.1

Uriarte, M., Muscarella, R., & Zimmerman, J. K. (2018). Environmental heterogeneity and
biotic interactions mediate climate impacts on tropical forest regeneration. *Global Change*
Biology, *24*(2), e692–e704. https://doi.org/10.1111/gcb.14000

Uriarte, M., Schwartz, N., Powers, J. S., Marín-Spiotta, E., Liao, W., & Werden, L. K. (2016).
Impacts of climate variability on tree demography in second growth tropical forests: The
importance of regional context for predicting successional trajectories. *Biotropica*, *48*(6),
780–797. https://doi.org/10.1111/btp.12380

van de Pol, M., Bailey, L. D., McLean, N., Rijsdijk, L., Lawson, C. R., & Brouwer, L. (2016).
Identifying the best climatic predictors in ecology and evolution. *Methods in Ecology and*
Evolution, *7*(10), 1246–1257. https://doi.org/10.1111/2041-210X.12590

Vasconcelos, H. L., & Luizão, F. J. (2004). Litter production and litter nutrient
concentrations in a fragmented Amazonian landscape. *Ecological Applications*, *14*(3), 884–
892. https://doi.org/10.1890/03-5093

Vicente-Serrano, S. M., Beguería, S., & López-Moreno, J. I. (2010). A multiscalar drought
index sensitive to global warming: The standardized precipitation evapotranspiration
index. *Journal of Climate*, *23*(7), 1696–1718. https://doi.org/10.1175/2009JCLI2909.1

Westerband, A. C., Horvitz, C. C., & Gilliam, F. (2017). Early life conditions and precipitation
influence the performance of widespread understorey herbs in variable light
environments. *Journal of Ecology*, *105*(5), 1298–1308. https://doi.org/10.1111/1365-
2745.12757

Wickham, H. (2016). *Ggplot2: Elegant graphics for data analysis*. Springer-Verlag New York.

Williams, J. L., Jacquemyn, H., Ochocki, B. M., Brys, R., Miller, T. E. X., & Shefferson, R. (2015).
Life history evolution under climate change and its influence on the population dynamics of
a long-lived plant. *Journal of Ecology*, *103*(4), 798–808. https://doi.org/10.1111/1365-
2745.12369

Williamson, G. B., Laurance, W. F., Oliveira, A. A., Delamônica, P., Gascon, C., Lovejoy, T. E., &
Pohl, L. (2000). Amazonian tree mortality during the 1997 El Niño drought. *Conservation*
Biology, *14*(5), 1538–1542. https://doi.org/10.1046/j.1523-1739.2000.99298.x

Wood, S. N. (2017). *Generalized additive models: An introduction with R* (Second edition).
CRC Press/Taylor & Francis Group.

Wright, S. J. (1992). Seasonal drought, soil fertility and the species density of tropical forest
plant communities. *Trends in Ecology & Evolution*, *7*(8), 260–263.
https://doi.org/10.1016/0169-5347(92)90171-7

Wright, S. J., & Calderon, O. (2006). Seasonal, El Niño and longer term changes in flower and
seed production in a moist tropical forest. *Ecology Letters*, *9*(1), 35–44.
https://doi.org/10.1111/j.1461-0248.2005.00851.x

Wright, S. J., Carrasco, C., Calderon, O., & Paton, S. (1999). The El Niño Southern Oscillation
variable fruit production, and famine in a tropical forest. *Ecology*, *80*(5), 1632–1647.

Xavier, A. C., King, C. W., & Scanlon, B. R. (2016). Daily gridded meteorological variables in
Brazil (1980). *International Journal of Climatology*, *36*(6), 2644–2659.
https://doi.org/10.1002/joc.4518

Zartman, C. E., Amaral, J. A., Figueiredo, J. N., & Dambros, C. D. (2015). Drought impacts
survivorship and reproductive strategies of an epiphyllous leafy liverwort in central
Amazonia. *Biotropica*, *47*(2), 172–178.

Zeng, N., Yoon, J. H., Marengo, J. A., Subramaniam, A., Nobre, C. A., Mariotti, A., & Neelin, J. D.
(2008). Causes and impacts of the 2005 Amazon drought. *Environmental Research Letters*,
3(1), 1–9. https://doi.org/10.1088/1748-9326/3/1/014002