Delayed effects of climate on vital rates lead to demographic divergence in Amazonian forest fragments

Eric R. Scott[1,✉], María Uriarte[2], and Emilio M. Bruna[1,3,4]

28 June, 2021

[1] Department of Wildlife Ecology and Conservation, University of Florida, Gainesville,

Florida 32611-0430

[2] Department of Ecology, Evolution and Environmental Biology, Columbia University 1200

Amsterdam Avenue, New York, New York 10027 USA

[3] Center for Latin American Studies, University of Florida, Gainesville, Florida 32611-5530

[4] Biological Dynamics of Forest Fragments Project, INPA-PDBFF, CP 478, Manaus,

Amazonas 69011-970 Brazil

✉ Correspondence: Eric R. Scott <scottericr@gmail.com>

Keywords: Amazon; delayed effects; demography; distributed lag non-linear models;

drought; ENSO; habitat fragmentation; *Heliconia acuminata*; Heliconiaceae

Running title: Delayed effects of climate on vital rates

Abstract

Deforestation often results in landscapes where remaining forest habitat is highly

fragmented, with remnants of different sizes embedded in an often highly contrasting

matrix. Local extinction of species from individual fragments is common, but the

demographic mechanisms underlying these extinctions are poorly understood. It is often

hypothesized that altered environmental conditions in fragments drive declines in

reproduction, recruitment, or survivorship. The Amazon basin, in addition to experiencing

continuing fragmentation, is warming and experiencing changes in precipitation leading to

altered frequency and intensity of droughts and unusually wet periods. Whether plant

populations in tropical forest fragments are particularly susceptible to extremes in

precipitation remains unclear. Most studies of plants in fragments are relatively short (1–6

years), focus on a single life-history stage, and often do not compare to populations in

continuous forest. Even fewer studies consider delayed effects of climate on demographic

vital rates despite the importance of delayed effects in studies that consider them. Using a

decade of demographic and climate data from an experimentally fragmented landscape in

the Central Amazon, we assess the effects of climate on populations of an understory herb

(*Heliconia acuminata*, Heliconiaceae). We used distributed lag non-linear models to

understand the delayed effects of temperature and precipitation on survival, growth, and

flowering. We detected delayed effects of climate up to 36 months. Drought two dry

seasons prior to the February census decreased survival and increased flowering

probability while drought in the wet season a year prior to the census decreased flowering

probability and increased growth. The effects of extremes in precipitation on survival and

growth were more pronounced in forest fragments compared to continuous forest. The

40   complex delayed effects of climate and habitat fragmentation in our study point to the

41   importance of long-term demography experiments in understanding the effects of

42   anthropogenic change on plant populations.

Introduction

The expansion of agriculture and other human activities is a primary driver of
deforestation in the tropics (Alroy, 2017; Haddad et al., 2015). It also results in landscapes
where the remaining forest can be highly fragmented, with patches of different sizes
embedded in a matrix of often contrasting habitat (Bianchi & Haig, 2013; Taubert et al.,
2018). This fragmentation is associated with myriad ecological changes, including the local
and regional extinction of plant species (da Silva & Tabarelli, 2000; Laurance et al., 2006).
Although the demographic mechanisms responsible these extinctions are poorly
understood (Bruna et al., 2009), it is often hypothesized that the dramatically altered
environmental conditions in tropical forest fragments (Arroyo-Rodríguez et al., 2017;
Didham & Lawton, 1999; Ewers & Banks-Leite, 2013) drive declines in plant reproduction,
recruitment, or survivorship (Bruna, 1999; Laurance et al., 1998; Zartman et al., 2015).
Despite the prevalence of this hypothesis (Betts et al., 2019; Didham & Lawton, 1999;
Laurance et al., 2001), efforts to link population-level demographic responses with altered
environmental conditions in fragments remains scarce.

Studies in temperate systems have shown that the demography of species can also be
altered by climate change (Doak & Morris, 2010; Selwood et al., 2015; Sletvold, 2005;
Williams et al., 2015), and that the effects of climate change could be more pronounced
when habitat is fragmented (Holyoak & Heath, 2016; Oliver et al., 2015). While the
demographic consequences of climate change for tropical species are expected to be
similarly severe (Brodie et al., 2012; Scheffers et al., 2017), surprisingly little is known
about the responses of these species to climatic variability (Paniw et al., 2021). Tropical
plants may be particularly sensitive to climate change—they typically have narrow ranges

of climatic tolerance (Feeley et al., 2012), and recent results suggest increases in the

frequency and severity of extreme precipitation events reduce survival and reproduction

(Esteban et al., 2021; Gaoue et al., 2019). This sensitivity to climatic fluctuations, coupled

with evidence that plant growth and survivorship are lower in fragments (Bruna et al.,

2002; Laurance et al., 1998; Zartman et al., 2015), has led to speculation that plants in

forest fragments will be especially susceptible to climate change (Laurance et al., 2001;

Opdam & Wascher, 2004; Selwood et al., 2015).

Whether the demography of plant populations in tropical forest fragments is more

susceptible to climatic extremes remains unclear for three primary reasons. First, most

studies of plants in fragments have focused on a single life-history stage or process (Bruna

et al., 2009; Ehrlen et al., 2016), making it challenging to draw broader demographic

conclusions. Second, there is a growing literature on how tropical plants respond to

droughts (Esquivel-Muelbert et al., 2019; González-M et al., 2020; Uriarte et al., 2016), but

few studies have compared the responses of plants in continuous forest with those of

plants in forest fragments (Laurance et al., 2001). Finally, the multi-year data needed to

test population-level hypotheses about climate change and fragmentation are scant,

especially for tropical systems (Crone et al., 2011; Salguero-Gómez et al., 2015). These data

are critical not simply because they allow for capturing variation in climatic conditions and

the resulting demographic responses (Morris & Doak, 2002; Teller et al., 2016). They are

also essential because while some demographic effects of fragmentation or drought can be

detected immediately, others may take years to manifest (*e.g.,* Gagnon et al., 2011). Indeed,

lagged responses of demographic vital rates to climate may be the rule rather than the

exception (Anderegg et al., 2015; Evers et al., 2021; Kannenberg et al., 2020; Schwalm et al.,

2017).

Herbaceous plants represent up to 25% of plant diversity in tropical forests (Gentry &

Dodson, 1987), are critical food and habitat for myriad species (Snow, 1981), and are

economically and culturally vital (Nakazono et al., 2004; Ticktin, 2003). Nevertheless, the

impacts of global change phenomena on their demography remain conspicuously

understudied (Bruna et al., 2009). We used a decade of demographic and climatic data from

an experimentally fragmented landscape in the Central Amazon to assess the effects of

climate on populations of a tropical understory herb (*Heliconia acuminata*, Heliconiaceae).

This time series, which included the severe droughts of 1997 (McPhaden, 1999) and 2005

(Marengo et al., 2008; Zeng et al., 2008), allowed us to address the following questions: (1)

Does drought increase or decrease the growth, survival, and fertility of plant populations in

continuous forest? (2) Are there delayed effects of drought on demographic vital rates, and

if so what lag times are most critical? (3) Are the effects of drought on the vital rates of

populations in fragments similar in direction and magnitude to those in continuous forest?

Methods

Study site

The Biological Dynamics of Forest Fragments Project (BDFFP) is located ~70 km north of

Manaus, Brazil (2º30' S, 60ºW). In addition to large areas of continuous forest, the BDFFP

has forest fragment reserves isolated from 1980–1984 by felling the trees surrounding the

area chosen for isolation and, in most cases, burning the downed trees once they dried

(Bierregaard et al., 1992). In subsequent decades the vegetation regenerating around

fragments has been periodically cleared to ensure fragment isolation (Bierregaard et al.,

2001).

The BDFFP reserves are located in nonflooded (i.e., *terra firme*) tropical lowland forest

with a 30–37m tall canopy (Rankin-de-Mérona et al., 1992) and an understory dominated

by stemless palms (Scariot, 1999). The soils in the reserves are nutrient-poor xanthic

ferrosols; their water retention capacity is poor despite having a high clay content. Mean

annual temperature in the region is 26º C (range=19–39º C), and annual rainfall ranges

from 1900–2300 mm. There is a pronounced dry season from June to October (Figure S1).

Focal species

Heliconia acuminata (LC Rich.) (Heliconiaceae) is a perennial monocot distributed

throughout Central Amazonia (Kress, 1990) and is the most abundant understory herb at

the BDFFP (Ribeiro et al., 2010). While many *Heliconia* species grow in large patches in

treefall gaps and other disturbed areas, others, such as *H. acuminata,* are found at lower

densities in the darker and cooler forest understory (Rundel et al., 2020). These species

produce fewer inflorescences and are pollinated by traplining rather than territorial

hummingbirds (Bruna et al., 2004; Stouffer & Bierregaard, 1996). In our sites *H. acuminata*

is pollinated by *Phaeothornis superciliosus* and *P. bourcieri* (Bruna et al., 2004). Plants begin

flowering at the start of the rainy season; reproductive plants have $\bar{x} = 1.1$ flowering

shoots (range = 1–7), each of which has an inflorescence with 20–25 flowers (Bruna &

Kress, 2002). Fruits mature April-May, have 1–3 seeds per fruit ($\bar{x} = 2$), and are eaten by a

thrush and several species of manakin (Uriarte et al., 2011). Dispersed seeds germinate

approximately 6 months after dispersal at the onset of the subsequent rainy season, with

rates of germination and seedling establishment higher in continuous forest than forest

fragments (Bruna, 1999; Bruna & Kress, 2002).

Demographic data collection

This study uses data collected in four 1-ha fragment reserves and six continuous forest

sites. In 1997–1998 we established a 5000 m^2 plots (50 × 100m) in each of these sites in

which we marked and measured all *Heliconia acuminata*; plots in 1-ha fragments were on

one randomly selected half of the fragment, while plots in continuous forest were located

500–4000 m from the borders of secondary and mature forest. The distance between plots

ranged from 500 m–41 km. Our dataset comprised 4,083 plants in continuous forest and

1,010 plants in forest fragments. Plots in CF had on average 2.7-fold more plants than plots

in 1-ha fragments (CF = 681 ± 493 SD; 1-ha = 253 ± 30 SD).

Each plot was subdivided into 50 quadrats (10 × 10m) to simplify annual surveys, during

which we recorded the number of vegetative shoots each plant had, the height of each plant

to the tallest leaf, and whether each plant was flowering (height and shoot number are

correlated with leaf area, the probability of flowering, and rates of survivorship (Bruna,

2002; Bruna & Kress, 2002). In this study, we used the product of shoot number and plant

height as our measure of plant size. Preliminary analysis showed that the product of shoot

number and height was a better predictor of total leaf area (which in turn is assumed to be

a strong predictor of aboveground biomass) than either shoot number or height alone

(Table S2). Plants that were not found for three consecutive surveys were considered

dead. We also surveyed plots regularly during the rainy season to identify any that

flowered after the survey. For additional details on the location of plots, survey methods,

and *H. acuminata* population structure see Bruna & Kress (2002).

Climate data

Data on precipitation and potential evapotranspiration in our sites were obtained from a

published gridded dataset ($0.25º \times 0.25º$ resolution) built using data from 3,625 ground-

based weather stations across Brazil (Xavier et al., 2016). We used these data to calculate

the standardized precipitation evapotranspiration index (SPEI) (Vicente-Serrano et al.,

2010). SPEI is a proxy for meteorological drought that integrates precipitation and

evapotranspiration anomalies over a specified time scale. Positive SPEI values for a given

month indicate conditions wetter than the historical average for that month, while negative

values of SPEI indicate droughts with intensity categorized as mild (0 to -1), moderate (-1

to -1.5), severe (-1.5 to -2), or extreme (< -2) (McKee et al., 1993). SPEI can be calculated to

represent different temporal scales of drought; we used 3-month SPEI because—given its

shallow roots and rhizome—*H. acuminata* relies primarily on soil moisture rather than

deeper water sources that can persist for longer timescales (Vicente-Serrano et al., 2010).

Note that 3-month SPEI is still monthly data—each month's SPEI value simply takes into

account precipitation and evapotranspiration of the previous three months. SPEI

calculations were made using the SPEI package (Beguería & Vicente-Serrano, 2017). The

timing of drought events based on these SPEI calculations is consistent with that resulting

from SPEI calculated with other data sources, though the magnitude of drought sometimes

differed (Figure S2; Table S1).

Statistical Modeling of Vital Rates

To assess the effects of drought history on plant vital rates we used Distributed Lag Non-

linear Models (DLNMs, Gasparrini et al., 2017). DLNMs capture how potentially delayed

effects of predictor variables (e.g. SPEI) affect an outcome (e.g. growth) well beyond the

event period. They do so by fitting a bi-dimensional predictor-lag-response association

spline, referred to as a crossbasis function. This models a non-linear relationship between

predictor and response (e.g. between SPEI and vital rates) and allows the shape of that

relationship to vary smoothly over lag time. Using the `dlnm` package (Gasparrini, 2011; R

Core Team, 2020), we created crossbasis functions with possible lags from 0–36 months.

We chose 36 months as a maximum lag because prior transplant experiments with *H.*

acuminata showed they typically recover from transplant shock in less than 36 months

(Bruna et al., 2002) so this is a reasonable upper bound for lagged effects of drought.

The crossbasis function was fit to the data in the context of a generalized additive model

(GAM) with restricted maximum likelihood using the `mgcv` package (Wood, 2017). The

general form of the vital rate (y) models was as follows:

190
$$g[\mathrm{E}(y_i)] = \alpha_j + s_1(z_i) + s_2(d_{i,t}, \ldots, d_{i,t-36}) + \beta x_i$$
$$\alpha_j \sim N\left(\mu_{\alpha_j}, \sigma^2_{\alpha_j}\right), \text{ for plot j} \tag{1}$$

191   where $s_1(z_i)$ is a smooth function of plant size (natural log of height × shoot number), fit

192   using a penalized cubic regression spline, $s_2(\cdot)$ is the crossbasis function in which $d_{i,t}$ is the

193   SPEI value during the census month of an observation (February) and $d_{i,t-l}$ is the SPEI l

194   months prior (see Gasparrini et al. 2017 for details). The crossbasis function, $s_2(\cdot)$ can also

195   be written:

$$196 \qquad s_2\big(d_{i,t}, \ldots, d_{i,t-24}\big) = \sum_{l=l_0}^{L} f \cdot w\big(d_{i,t-l}, l\big) \qquad (2)$$

where the crossbasis function, $f \cdot w(d,l)$, is composed of two marginal basis functions: the

standard predictor-response function $f(d)$, and the additional lag-response function $w(l)$.

These marginal functions are combined as a tensor product smooth such that the shape of

one marginal function varies smoothly along the other dimension (see chapter 5 of Wood

(2017) and Gasparrini et al. (2017) for more detail). Penalized cubic regression splines

were used for both marginal bases of the crossbasis function, with 35 knots for the lag

dimension (i.e. number of lagged SPEI values for each observation with 36 months as a

maximum lag) and 3 knots for the drought response dimension to restrict the shape of the

fitted response to drought to bimodal when most complex. Because of penalization, the

number of knots is generally not important as long it is large enough to allow the smooth to

represent the 'true' relationship (Wood, 2017). Estimated degrees of freedom (edf)

represent the 'true' complexity of the smooth after penalization with edf = 1 being

equivalent to a straight line and larger numbers representing more complex curves.

To determine if plot characteristics influenced average vital rates we included a random

effect of plot ID on the intercept; this was represented by a_j in eq. 1. We determined the

effects of SPEI on plant growth using plant size in year t+1 as a response variable. This was

modeled with a scaled t family error distribution because residuals were leptokurtic with a

Gaussian error structure. Because number of inflorescences was highly zero-inflated, we

converted this to a binary response to model reproduction (i.e., 1 for ≥1 inflorescence, 0 for

no inflorescences). We modeled both reproduction and survival (i.e., from year t to year

t+1) using a binomial family error distribution with a logit link function. We modeled a

potential cost of reproduction by including flowering in the previous year as covariate, x_i,

in eq. 1. Additionally, in models for flowering probability and size, we included plant ID as a

random effect to account for variation among individuals. Preliminary analyses showed

that this random effect was not significant in the growth models (edf ~ 0, p > 0.2) and as

such it was dropped to improve computational efficiency. A random effect of plant ID was

not included in survival models since each plant only dies once.

In the process of fitting the models, the penalty on the crossbasis smooth (and other

smoothed terms) is optimized such that more linear shapes are favored unless the data

supports non-linearity (Wood, 2017). We applied an additional penalty to shrink linear

portions toward zero with the `select=TRUE` option to the `gam()` function, and inferred

statistical significance of model terms with p-values from the `summary.gam()` function as

recommended in Marra & Wood (2011).

The `dlnm` package does not currently allow the modeling of interaction terms, which means

we could not asses the interaction of habitat type and lagged effects. We therefore fit

separate models for plants in fragments and in continuous forest to allow the shape of the

crossbasis function to differ between habitats. Significant main effects of habitat type were

assessed by looking for overlap in the 84% confidence intervals of model intercepts; the

84% CIs of two samples drawn from the same population overlap about 95% of the time

(Payton et al., 2003).

All analyses were conducted in R version 4.0.2 (2020-06-22) (R Core Team, 2020) using

the `targets` package for workflow management (Landau, 2021). Figures were created with

the aid of the `gratia`, `ggplot2`, and `patchwork` packages (Pedersen, 2020; Simpson, 2021;

Wickham, 2016).

Results

The meteorological droughts in our focal region indicated by SPEI are generally consistent

with those reported in the literature (Table S1). For example, the drought associated with

the 1997 El Niño Southern Oscillation (ENSO) event was one of the most severe on record

for the Amazon (Williamson et al., 2000); correspondingly, 1997 has the lowest SPEI values

in our timeseries (Figure 1d). The 2005 dry season (June–October) was also reported as an

exceptionally dry year, although this drought mostly affected the southwestern Amazon

(Marengo et al., 2008; Zeng et al., 2008). Our SPEI data show the 2005 dry season to be a

moderate drought (-1 > SPEI > -1.5).

Survival, growth, and flowering in continuous forest vs. fragments

Survival: Across all plots, the proportion of plants surviving was lowest in the 2003–2004

transition year ($P_{surv} = 0.92$). This coincided with droughts in both the 2003 and 2004

rainy seasons (Figure 1b,d) and was preceded by a drop in average plant size in the 2002–

2003 transition year (Figure 1a). The lowest survival for 1-ha fragment plots ($P_{surv} = 0.93$)

was for the 2005–2006 transition year, which encompassed a moderate drought in October

2005 and and wetter than average conditions (SPEI > 0.5) in December 2005 and January

2006 (Figure 1b,d). The lowest survival for continuous forest was in 2004 ($P_{surv} = 0.91$).

When summarizing across years, plots, and plant sizes, the survival probability of *Heliconia*

acuminata was similarly high in both continuous forest and fragments ($P_{surv} = 0.95$; Figure

1b; the overlapping 85% CI of model intercepts indicate no significant difference).

However, survival in both habitats was size dependent ($p < 0.001$ for the effect of log-

transformed plant size in year t on survival in year $t+1$ in both habitats). The survival

probability of large plants approached 1 in both habitat types (Figure 3b), but the smallest

plants had higher survival in 1-ha fragments.

Growth: Plants in continuous forest had an average of 2.9 shoots (± 1.8 SD) and were on

average 40.6 cm tall (± 26.5 SD). Plants in 1-ha fragments had on average 13.8% fewer

shoots (2.5 ± 1.5 SD) and were 10.8% shorter (36.3 cm ± 24.1 SD). Because our proxy for

plant size was the product of these two metrics, plants in continuous forest were on

average 34% larger than those in forest fragments (150 ± 175 SD vs. 112 ± 141 SD,

respectively), with fragments having proportionately fewer large plants (Figure 3d). This

difference was not significant, however (overlap in 84% CI of model intercepts), and the

disparity in plant size—which was most pronounced in the initial years of our surveys—

diminished over time (Figure 1a).

Mean plant size dropped dramatically in 2003 in both habitat types, corresponding with a

severe drought during the February census (SPEI = -1.39) (Figure 1d). As with survival,

size in year t was a significant predictor of size in year t+1 ($p < 0.001$ in both habitats).

While the effect was generally similar across size classes and habitat types, the impact of

plant size on growth was greatest for mid-sized plants in continuous forest (Figure 3a).

Flowering: The overall proportion of plants that flowered was very low. While it was 27%

higher in continuous forest than 1-ha fragments (0.048 ± 0.213 vs. 0.038 ± 0.19,

respectively), this difference was not statistically significant (84% CIs of model intercepts

overlapped). The observed disparity was largely due to the fact that flowering is also size-

dependent ($p < 0.001$ in both habitats), with the probability of flowering increasing

dramatically once plants reached the threshold size of about 148 (i.e., log(size) > 5 in

Figure 3c). Despite the flowering probability of large plants being greater in fragments than

continuous forest, populations in fragments had proportionately fewer plants above the

reproductive size threshold (Figure 3d). The most striking difference between habitat

types coincided with a severe drought in 2003, when the percentage of flowering

reproductive-sized plants was 28% in continuous forest vs. only 13.6% in 1-ha fragments

(Figure 1c).

Delayed effects of drought on demographic vital rates

Drought history had a significant ($p < 0.001$) effect on the survival, growth, and flowering

of plants in both habitats. Comparing the respective crossbasis surfaces, however, reveals

that the specific climatic drivers, their timing, and their impact on individual vital rates all

differed among habitats.

Survival: For 1-ha fragments, there was a significant effect on survival of SPEI in the

preceding 13 months. The highest survival was near SPEI of 0, with mortality increasing as

conditions became either drier or wetter (i.e., as SPEI values became increasingly negative

or positive, respectively; Figure 4b). Wet conditions in the preceding 23 months (i.e., SPEI >

1) also had a significant negative effect on survival in fragments (Figure 4b). In contrast,

the effect of recent SPEI in continuous forest was weaker, with only the 6 months preceding

a census having a significant effect on survival (Figure 4a). These short-term effects of SPEI

on survival were also unidirectional—the probability of survival declined, albeit only

slightly, with increasingly negative values of SPEI (i.e., as droughts became more severe;

Figure 4a). In contrast, the most pronounced negative effects of SPEI on the survival of

plants in continuous forest were at lag times of 15–20 months and 32–36 months. Drought

15–20 months prior to a census (i.e. two dry seasons prior to a census) was significantly

associated with reduced survival, while high precipitation (i.e., SPEI > 1) was significantly

associated with higher survival. Finally, plants in both habitat types showed an increase in

survival probability with very high SPEI values (i.e., extremely high precipitation) at a lag

time of 32–36 months. It should be noted, however, that only the first transition year of

census data (1998–1999) met these conditions. We compared the effects of SPEI history in

continuous forest and fragments by subtracting the fitted values in Figure 4b from Figure

4a to produce Figure 4c. This shows that in average conditions (SPEI = 0), there is little

difference in survival probability between continuous forest and forest fragments (Figure

4c). However, under extreme conditions, survival probability is higher in continuous forest

by up to 0.025.

Growth: The effects of drought history on trends in plant size were generally similar for

continuous forest and fragments. However, the crossbasis function for 1-ha fragments

indicated more complex responses in some situations (edf = 17.8 for 1-ha fragments; edf =

13.0 for continuous forest; see also Figure 5). For example, under average conditions (i.e.,

SPEI = 0), growth is similar or slightly higher in continuous forest over all lag periods

(Figure 5c). However, the growth of plants in fragments is reduced when the current wet

season is unusually wet (i.e., SPEI>2), and as a result the plants in continuous forests will

be larger by up to log(size) = 0.57. In contrast, drought at lags of 8–11 months (i.e., the end

of the preceding year's wet season) led to increased growth in both habitats, with a more

pronounced response in 1-ha fragments.

Flowering: Overall, the the probability of flowering was higher in continuous forest than in

1-ha fragments for all values of SPEI (Figure 6), although this difference was not significant

(84% CIs of intercepts overlap). The responses in 1-ha fragments were also more muted as

indicated by the shape of the crossbasis function (1-ha edf = 8.3, continuous forest edf =

10.6) (Figure 6). This led to some important inter-habitat differences in plant responses to

prior droughts. In continuous forests, recent drought (i.e., at lag = 0–2 with SPEI < -1),

drought two dry seasons prior (lags 15–20) and in the wet season 34–36 months prior all

increased the probability of flowering. However, drought at the end of the rainy season one

year prior (lags 7–13) significantly reduced flowering probability (Figure 6a). In fragments,

recent drought had no significant effect on flowering probability and only drought at two

dry seasons prior (lags 16–20) and in the wet season 33–36 months prior increased

flowering probability (Figure 6b). The effects of drought on flowering probability were

strong in continuous forest compared to 1 ha fragments (Figure 6c). We found no evidence

for a cost of reproduction: in both forest and fragments, plants that had flowered in the

previous year were on average more likely to be larger (CF: $p = 0.048$; 1-ha: $p = 0.030$)

and flower again (CF: $p < 0.001$; 1-ha: $p = 0.004$). The random effect of plant ID on

flowering probability was also significant (CF: $p < 0.001$; 1-ha: $p = 0.003$), indicating

significant individual-level variation in flowering probability.

Finally, with the exception of survival in 1-ha fragments ($p = 0.253$), the delayed effects of

SPEI on all three vital rates varied significantly among plots ($p < 0.01$ for the random effect

of plot).

Discussion

Understanding how landscape structure and abiotic conditions act to influence population dynamics is central to many conceptual frameworks for studying and conserving fragmented landscapes (Didham et al., 2012; Driscoll et al., 2013). Our results support the emerging consensus that the effects of climatic extremes on demographic vital rates can be delayed for months or even years (Evers et al., 2021; Teller et al., 2016; Tenhumberg et al., 2018). We also found that the delayed responses of populations in fragments can differ significantly in magnitude, direction, and lag time from those of populations in continuous forest. This suggests that the hypothesized synergies between climate and fragmentation on population dynamics (Laurance & Williamson, 2001; Opdam & Wascher, 2004; Selwood et al., 2015) are likely to be pervasive, but also far more complex than previously thought.

Temporal variation in demographic responses to forest fragmentation

Many studies investigating the biological consequences of habitat fragmentation on plant growth, survival, and reproduction comprise short-term (≤3 year) experiments and observations. Our results underscore the difficulty in extrapolating long-term trends from such short-term studies, particularly when studying long-lived organisms or when the responses of interest can vary with size or age. For instance, one would have reached a very different conclusion regarding the effect of fragmentation on annual survival if the study windows were 1999–2002 (i.e., higher survival in continuous forest), 2002–2005 (i.e., higher survival in fragments), or 2004–2007 (i.e, no clear effect of fragmentation) (Figure 1b). It is only when evaluating over longer time windows that it becomes apparent mortality is elevated in fragments relative to continuous forest (Figure 2), and that the observed interannual variation is largely driven by dynamic patterns of recruitment

(Bruna, 2002) coupled with low mortality for plants beyond the smallest size classes

(Bruna, 2003).

Similarly, conclusions regarding the effects of fragmentation on flowering—which is also

both rare and size-dependent (Brooks et al., 2019)—would also differ based on the year in

which they were investigated. This could lead to erroneous extrapolations regarding the

effects of fragmentation on reproductive mutualists or population genetic structure (Côrtes

et al., 2013; Uriarte et al., 2010; Uriarte et al., 2011). Conclusions based on short-term

observations of temporally variable vital rates could lead to conservation and management

practices that are ineffective or even counterproductive, especially when when failing to

consider how the consequences of this variation might be modulated by organismal life

history (Morris et al., 2008).

It is important to emphasize, however, that the overall effects of SPEI on survival and

growth are more severe in fragments than continuous forest (Figures 4, 5). Furthermore,

the magnitude of plant responses to climatic extremes is also greater in habitat

fragments—extreme drought in dry seasons and extreme precipitation in during rainy

seasons are most detrimental to growth and survival in fragments. While intact forest and

its canopy buffer populations from climatic extremes, populations in fragments—especially

near edges with high contrast matrix—likely lack this protection (Didham & Lawton, 1999;

Ewers & Banks-Leite, 2013). We suggest it is these climate extremes, rather than trends in

average temperature, precipitation, or SPEI (Laurance et al., 2014), that that are the causal

mechanism underlying reduced plant growth and survival in forest fragments.

Delayed effects of climate on demographic vital rates

Climate anomalies are known to have immediate effects on the growth, survival, or reproduction of plants (Esteban et al., 2021; Wright & Calderon, 2006), including *Heliconia* (Stiles, 1975; Westerband et al., 2017) and other tropical herbs (Wright, 1992). These effects can be complex or even contradictory—mild droughts can increase the growth rates of tropical trees and seedling survival, perhaps due to reductions in cloud cover and concomitant increases in solar radiation (Alfaro-Sánchez et al., 2017; Condit et al., 2004; Huete et al., 2006; Jones et al., 2014; Uriarte et al., 2018), but in severe drought years growth can be extremely low and mortality can be sharply elevated (Connell & Green, 2000; Edwards & Krockenberger, 2006; Engelbrecht et al., 2002). There is also evidence that the effects can persist for multiple years (Phillips et al., 2010), such as a boom in drought-year fruit production followed by severe post-drought "famine" (Pau et al., 2013; Wright et al., 1999).

Despite these insights, models of plant population dynamics rarely include the effects of environmental drivers [but see Williams et al. (2015); Tenhumberg et al. (2018); Molowny-Horas et al. (2017)). This has largely been due to the challenge (both ecologically and statistically) of detecting any demographic responses to climatic extremes that are delayed for multiple growing seasons. To address this, researchers have begun to use a number of statistical methods that test for time lags in demographic responses without *a priori* assumptions about the influence of any particular climate window (Evers et al., 2021; Teller et al., 2016; Tenhumberg et al., 2018). Our expansion of this approach, which offers an unbiased way of identifying these delayed effects without overfitting (but see Pierre et al. (2020) and Ogle et al. (2015) for alternative methods) yielded results consistent with

this emerging literature—that the effects of precipitation extremes on the demography of

Heliconia acuminata could be delayed for up to 3 growing seasons.

While it appears that delayed effects of climate on demographic vital rates may be

ubiquitous (Evers et al., 2021), the extent to which they vary spatially or with habitat

remains an open question. Our results clearly indicate that they can, with habitat-specific

differences in how environmental conditions influenced future vital rates. For example,

extreme values of SPEI—both positive (unusually high precipitation) and negative

(drought conditions)—led to declines in the probability of individual survival in both

habitat types. However, the magnitude of these declines was far greater in forest

fragments. Similarly, the detrimental effects of extremes in SPEI on growth rates were also

more pronounced in fragments. In contrast, variation in SPEI had far stronger effects on the

probability of flowering in continuous forest than fragments. These results should be

interpreted with some caution, however, as the relatively low number of plants in

fragments that are above the threshold-size for flowering could limit the power to detect

delayed effects.

There are several, non-mutually-exclusive explanations for delayed effects of SPEI on

demography. The first is that the physiological processes underlying vital rates might be

initiated long before they are demographically apparent (Evers et al., 2021), and hence be

shaped by climatic events at any point in that physiological window. For example, the

flowering shoots of *Heliconia chartacea* begin to develop 6–10 months prior to the

appearance of inflorescences (Criley & Lekawatana, 1994). Adverse conditions during the 6

months following initiation, rather than the months when inflorescences are starting

expand, leads to the aborted production of flowering shoots. Interestingly, we observed the

opposite effect—drought conditions increased the probability of flowering two years later.

While this could reflect bet-hedging in response to stress (Nihad et al., 2018), this does not

appear to be the case, as growth or survival do not appear to decrease following

reproduction (see also (Horvitz & Schemske, 1988). In fact, flowering in one year is

associated with increased reproduction and growth in the next .

Demographic responses will also be delayed if abiotic stress causes plants to invest in

belowground rhizomes (*sensu* Pumisutapon et al., 2012). The carbohydrates stored in

rhizomes allow *Heliconia* to regenerate aboveground biomass following damage (Rundel et

al., 1998) and protect the buds that give rise to new shoots from stressful conditions

(Klimešová et al., 2018). This may be why drought led to delayed increases in growth—by

shedding shoots and leaves (Bruna et al., 2002) and investing in rhizomes, plants are

generating proportionately more buds with which to regenerate when conditions improve.

This would also be consistent with the results of prior experiments, in which the growth

rates of *H. acuminata* 8 months after they were mechanically damaged far exceeded those

of control plants (Bruna & Ribeiro, 2005).

Third, it may be that the delayed demographic effects we observed are indirectly mediated

by the effect of SPEI on other species rather than the direct effects on individual physiology

(Evers et al., 2021). For example, topical trees may not die until three or more years after a

drought (Criley & Lekawatana, 1994). When they finally do, the resulting leaf drop (Janssen

et al., 2021) and treefalls allow for light penetration to the forest understory (Canham et al.,

1990; Leitold et al., 2018), triggering a boom in the growth and flowering of understory

plants (Bruna & Oli, 2005). Similar delayed changes in the local environment could also

influence the foraging behavior of a plant's pollinators (Bruna et al., 2004; Stouffer &

Bierregaard, 1996), seed dispersers (Uriarte et al., 2011), or herbivores (Scott et al., 2021).

While more work is needed to explain why the (delayed) effects of SPEI on *H. acuminata*

survival and growth are greater in fragments than forest interiors, one hypothesis,

motivated by recent intriguing results from other systems (Sapsford et al., 2017), is that the

greater litterfall on edges (Vasconcelos & Luizão, 2004) may be altering the abundance of

pathogens or mycorrhizae.

Finally, demographic delays could be an artifact of the timing of responses in relation to the

census date. If extreme drought in the dry season before the census increased plant

mortality during that season, for example, this would nevertheless appear in models as a

delayed effect (e.g. in Figure 4b). In our case, this potential explanation for delayed effects

applies only to plant size and survival, as plots were surveyed regularly throughout the

reproductive season to identify flowering plants. This possibility is not unique to our study,

rather it is a consequence of conducting demographic censuses on an annual scale while

the climate is quantified monthly or seasonally. While the very slow growth and extremely

low mortality rates of *H. acuminata* mean this effect is unlikely to be acting in our system, it

may be that for some species it will be important to conduct demographic surveys at the

same temporal scale at which climate is aggregated.

Conclusions & Future Directions

Over 24 million ha of the Brazilian Amazon have been cleared in the last two decades (Silva

Junior et al., 2021), resulting in their extensive fragmentation (Broadbent et al., 2008).

Climate models predict a future of extremes for these forests—increases in the frequency

and geographic extent of droughts, but also increases in the frequency and area affected by

periods of unusual wetness (Duffy et al., 2015). Our results support the hypothesis that

populations in Amazonian forest fragments could be more susceptible to the effects of

changing climate than those in continuous forest (Laurance et al., 2014). However, they

also indicate that the demographic responses to climate change of populations in

fragmented landscapes may be far more complex than previously appreciated. Multi-

factorial, multi-season experiments (*sensu* Aguirre et al., 2021; Bruna & Ribeiro, 2005;

Markewitz et al., 2010; Westerband et al., 2017), ideally manipulating multiple

combinations of climatic variables (Mundim & Bruna, 2016), are needed to determine how

and why habitat-specific differences in environmental conditions interact to delay the

demographic responses of plants to climatic variability. Also needed are statistical tools

that can test for synergistic effects of fragmentation and climate in vital rates, as those

currently available do not allow for including interaction terms. This also limits the ability

to include size by climate interactions in a DLNM; although plant responses to both

fragmentation and climatic extremes can be size-specific (Bruna & Oli, 2005; Schwartz et

al., 2019). The ability to identify size-specific lagged responses may be especially

complicated given size and growth are rarely measured at the same time scale as SPEI and

other putative climatic drivers.

Finally, no analytical approach assessing the potential for demographic lags can

compensate for a lack of long-term data (Evers et al., 2021; Tenhumberg et al., 2018).

Unfortunately, long-term data monitoring the entire life-cycle of tropical taxa are rare, and

those doing so in fragmented landscapes are virtually nonexistent (Bruna & Ribeiro, 2005).

Without investing in collecting such data, generalizations regarding the demographic

consequences of climate change in these species rich and increasingly fragmented habitats

will continue to prove elusive. More generally, however, researchers need to consider how

delayed responses to climate could influence the interpretation of data in studies where the

organisms lifespan exceeds the study's duration.

Acknowledgments

We thank Collin Edwards, ___, ___, and ___ anonymous reviewers for helpful discussions and

comments on the manuscript. We also thank the technicians and students who helped

conduct the *Heliconia* censuses and the BDFFP and INPA for logistical support. Financial

support was provided by the U.S. National Science Foundation (awards DEB-0614339,

DEB-0614149, INT 98-06351, and DEB-1948607). This is publication No. ___ in the BDFFP

Technical Series. The authors delcare no conflicts of interest.

Data Availability Statement

518   Data used in this study are available at the Dryad Digital Repository; R code for analyses

519   and visualizations are archived at Zenodo [links included upon acceptance].

Figures

Figure 1: (a-c) Time series of *H. acuminata* vital rates in 1-ha fragments (solid orange lines) and continuous

forest (dashed blue lines) and (d) drought occurrence in the study region. (a) Mean fold-change in plant plant

size ($\log2(\text{size}_{t+1} / \text{size}_t)$) varies by year and habitat. On average, plants grew in most years with the notable

exception in 2003, in which on average plants regressed in size in both habitats (i.e., fold-change < 0). Error

bars represent the standard deviation. (b) The proportion of plants surviving from one transition year to the

next varied from 0.98 (CF in 1998-1999) to 0.91 (CF in 2003-2004). (c) The proportion of *H. acuminata* above

the size threshold for reproduction that flowered each year is on average low but variable. The size threshold

is determined by the upper 90th percentile size of flowering plants across all years. (d) Monthly 3-month

SPEI for our study region. Gray lines represent values from different grid cells encompassing BDFFP; the dark

line represents the site mean. Colored stripes represent drought intensity: yellow = mild, orange = moderate,

dark orange = severe, red = extreme.

Figure 2: Survivorship curve for plants marked in the 1998 survey year; these plants comprise 49% of those

in the complete demographic dataset. The percentage of these plants that were still alive ten years later was

79.7% (1629/2055) in continuous forest vs. 72.4% (393/543) in 1-ha fragments.

Figure 3: Smooth effect from models of plant size in the previous census on (a) survival, (b) log(size), and (c)

flowering probability; these values correspond to the additive term $s_1(z_i)$ in eq. 1. The bands depicting the

95% confidence interval include uncertainty in the intercept and uncertainty due to smoothness selection;

the smooths for 1-ha fragments and continuous forest are fit in separate models. (d) Plant size distribution by

habitat type (solid line = 1-ha fragments, dashed line = Continuous Forest).

Figure 4: Smooth effect of lagged SPEI on *H. acuminata* survival in (a) continuous forest, (b) 1-ha fragments,

(c) and the difference between the two. Outlines show regions where the effect of SPEI is significant, defined

as those where the 95% confidence interval around the response does not overlap the intercept. The bar on

the bottom of each panel indicates the wet seasons (black, November–May) and dry seasons (white, June–

October). Surface is modeled as a crossbasis function with cubic regression splines for each marginal basis.

549   Model intercepts were added to fitted values of the crossbasis function and back-transformed to the response

550   scale.

552   Figure 5: Smooth effect of lagged SPEI on *H. acuminata* growth in (a) continuous forest, (b) 1-ha fragments,

553   (c) and the difference between the two. Outlines show regions where the effect of SPEI is significant, defined

554   as those where the 95% confidence interval around the response does not overlap the intercept. The bar on

a)

b)

c)

Figure 6: Smooth effect of lagged SPEI on *H. acuminata* flowering probability in (a) continuous forest, (b) 1-ha

fragments, (c) and the difference between the two. Outlines show regions where the effect of SPEI is

significant, defined as those where the 95% confidence interval around the response does not overlap the

intercept. The bar on the bottom of each panel indicates the wet seasons (black, November–May) and dry

seasons (white, June–October). Surface is modeled as a crossbasis function with cubic regression splines for

each marginal basis. Model intercepts were added to fitted values of the crossbasis function and back-

transformed to the response scale.

References

Aguirre, B. A., Hsieh, B., Watson, S. J., & Wright, A. J. (2021). The experimental manipulation
of atmospheric drought: Teasing out the role of microclimate in biodiversity experiments.
Journal of Ecology, *109*(5), 1986–1999. https://doi.org/https://doi.org/10.1111/1365-
2745.13595

Alfaro-Sánchez, R., Muller-Landau, H. C., Wright, S. J., & Camarero, J. J. (2017). Growth and
reproduction respond differently to climate in three Neotropical tree species. *Oecologia*,
184(2), 531–541. https://doi.org/10.1007/s00442-017-3879-3

Alroy, J. (2017). Effects of habitat disturbance on tropical forest biodiversity. *Proceedings of
the National Academy of Sciences*, *114*(23), 6056–6061.
https://doi.org/10.1073/pnas.1611855114

Anderegg, W. R. L., Schwalm, C., Biondi, F., Camarero, J. J., Koch, G., Litvak, M., Ogle, K., Shaw,
J. D., Shevliakova, E., Williams, A. P., Wolf, A., Ziaco, E., & Pacala, S. (2015). Pervasive
drought legacies in forest ecosystems and their implications for carbon cycle models.
Science, *349*(6247), 528–532. https://doi.org/10.1126/science.aab1833

Arroyo-Rodríguez, V., Saldaña-Vázquez, R. A., Fahrig, L., & Santos, B. A. (2017). Does forest
fragmentation cause an increase in forest temperature? *Ecological Research*, *32*(1), 81–88.
https://doi.org/10.1007/s11284-016-1411-6

Beguería, S., & Vicente-Serrano, S. M. (2017). *SPEI: Calculation of the standardised
precipitation-evapotranspiration index*. https://CRAN.R-project.org/package=SPEI

Betts, M. G., Wolf, C., Pfeifer, M., Banks-Leite, C., Arroyo-Rodríguez, V., Ribeiro, D. B., Barlow,
J., Eigenbrod, F., Faria, D., Fletcher, R. J., Hadley, A. S., Hawes, J. E., Holt, R. D., Klingbeil, B.,
Kormann, U., Lens, L., Levi, T., Medina-Rangel, G. F., Melles, S. L., … Ewers, R. M. (2019).
Extinction filters mediate the global effects of habitat fragmentation on animals. *Science*,
366(6470), 1236–1239. https://doi.org/10.1126/science.aax9387

Bianchi, C. A., & Haig, S. M. (2013). Deforestation trends of tropical dry forests in Central
Brazil. *Biotropica*, *45*(3), 395–400. https://doi.org/https://doi.org/10.1111/btp.12010

Bierregaard, R. O., Gascon, C., Lovejoy, T. E., & Mesquita, R. (Eds.). (2001). *Lessons from
Amazonia: The ecology and conservation of a fragmented forest*. Yale University Press.

Bierregaard, R. O., Lovejoy, T. E., Kapos, V., & Hutchings, R. W. (1992). The biological
dynamics of tropical rainforest fragments. *BioScience*, *42*(11), 859–866.
https://doi.org/10.2307/1312085

Broadbent, E. N., Asner, G. P., Keller, M., Knapp, D. E., Oliveira, P. J. C., & Silva, J. N. (2008).
Forest fragmentation and edge effects from deforestation and selective logging in the
Brazilian Amazon. *Biological Conservation*, *141*, 1745–1757.
https://doi.org/10.1016/j.biocon.2008.04.024

Brodie, J., Post, E., & Laurance, W. F. (2012). Climate change and tropical biodiversity: A
new focus. *Trends in Ecology & Evolution*, *27*(3), 145–150.
https://doi.org/10.1016/j.tree.2011.09.008

Brooks, M. E., Kristensen, K., Darrigo, M. R., Rubim, P., Uriarte, M., Bruna, E. M., & Bolker, B.
M. (2019). Statistical modeling of patterns in annual reproductive rates. *Ecology*, *100*(7).
https://doi.org/10.1002/ecy.2706

Bruna, E. M. (1999). Seed germination in rainforest fragments. *Nature*, *402*(6758), 139.
https://doi.org/10.1038/45963

Bruna, E. M. (2002). Effects of forest fragmentation on *Heliconia acuminata* seedling
recruitment in central Amazonia. *Oecologia*, *132*(2), 235–243.
https://doi.org/10.1007/s00442-002-0956-y

Bruna, E. M. (2003). Are plant populations in fragmented habitats recruitment limited?
Tests with an Amazonian herb. *Ecology*, *84*(4), 932–947. https://doi.org/10.1890/0012-
9658(2003)084[0932:APPIFH]2.0.CO;2

Bruna, E. M., Fiske, I. J., & Trager, M. D. (2009). Habitat fragmentation and plant
populations: Is what we know demographically irrelevant? *Journal of Vegetation Science*,
20(3), 569–576. https://doi.org/10.1111/j.1654-1103.2009.01060.x

Bruna, E. M., & Kress, W. J. (2002). Habitat fragmentation and the demographic structure of
an Amazonian understory herb (*Heliconia acuminata*). *Conservation Biology*, *16*(5), 1256–
1266. https://doi.org/10.1046/j.1523-1739.2002.99494.x

Bruna, E. M., Kress, W. J., Marques, F., & Silva, O. F. da. (2004). *Heliconia acuminata*
reproductive success is independent of local floral density. *Acta Amazonica*, *34*(3), 467–
471. https://doi.org/10.1590/S0044-59672004000300012

Bruna, E. M., Nardy, O., Strauss, S. Y., & Harrison, S. (2002). Experimental assessment of
Heliconia acuminata growth in a fragmented Amazonian landscape. *Journal of Ecology*,
90(4), 639–649. https://doi.org/10.1046/j.1365-2745.2002.00707.x

Bruna, E. M., & Oli, M. K. (2005). Demographic effects of habitat fragmentation on a tropical
herb: Life-table response experiments. *Ecology*, *86*(7), 1816–1824.
https://doi.org/10.1890/04-1716

Bruna, E. M., & Ribeiro, M. B. N. (2005). The compensatory responses of an understory herb
to experimental damage are habitat-dependent. *American Journal of Botany*, *92*(12), 2101–
2210. https://doi.org/10.3732/ajb.92.12.2101

Canham, C. D., Denslow, J. S., Platt, W. J., Runkle, J. R., Spies, T. A., & White, P. S. (1990). Light regimes beneath closed canopies and tree-fall gaps in temperate and tropical forests. *Canadian Journal of Forest Research*, *20*(5), 620–631. https://doi.org/10.1139/x90-084

Condit, R., Aguilar, S., Hernandez, A., Perez, R., Lao, S., Angehr, G., Hubbell, S. P., & Foster, R. B. (2004). Tropical forest dynamics across a rainfall gradient and the impact of an El Niño dry season. *Journal of Tropical Ecology*, *20*(01), 51–72. https://doi.org/10.1017/S0266467403001081

Connell, J. H., & Green, P. T. (2000). Seedling dynamics over thirty-two years in a tropical rain forest tree. *Ecology*, *81*(2), 568–584.

Côrtes, M. C., Uriarte, M., Lemes, M. R., Gribel, R., Kress, W. J., Smouse, P. E., & Bruna, E. M. (2013). Low plant density enhances gene dispersal in the Amazonian understory herb *Heliconia Acuminata*. *Molecular Ecology*, *22*(22), 5716–5729. https://doi.org/10.1111/mec.12495

Criley, R., & Lekawatana, S. (1994). Year around production with high yields may be a possibility for *Heliconia chartacea*. *Acta Horticulturae*, *397*, 95–102. https://doi.org/10.17660/ActaHortic.1995.397.7

Crone, E. E., Menges, E. S., Ellis, M. M., Bell, T., Bierzychudek, P., Ehrlen, J., Kaye, T. N., Knight, T. M., Lesica, P., Morris, W. F., Oostermeijer, G., Quintana-Ascencio, P. F., Stanley, A., Ticktin, T., Valverde, T., & Williams, J. L. (2011). How do plant ecologists use matrix population models? *Ecology Letters*, *14*(1), 1–8.

da Silva, J. M. C., & Tabarelli, M. (2000). Tree species impoverishment and the future flora of the Atlantic forest of northeast Brazil. *Nature*, *404*(6773), 72–74. https://doi.org/10.1038/35003563

Didham, R. K., Kapos, V., & Ewers, R. M. (2012). Rethinking the conceptual foundations of habitat fragmentation research. *Oikos*, *121*(2), 161–170. https://doi.org/10.1111/j.1600-0706.2011.20273.x

Didham, R. K., & Lawton, J. H. (1999). Edge structure determines the magnitude of changes in microclimate and vegetation structure in tropical forest fragments. *Biotropica*, *31*(1), 17–30. https://doi.org/10.1111/j.1744-7429.1999.tb00113.x

Doak, D. F., & Morris, W. F. (2010). Demographic compensation and tipping points in climate-induced range shifts. *Nature*, *467*(7318), 959–962.

Driscoll, D. A., Banks, S. C., Barton, P. S., Lindenmayer, D. B., & Smith, A. L. (2013). Conceptual domain of the matrix in fragmented landscapes. *Trends in Ecology & Evolution*, *28*(10), 605–613. https://doi.org/10.1016/j.tree.2013.06.010

Duffy, P. B., Brando, P., Asner, G. P., & Field, C. B. (2015). Projections of future meteorological drought and wet periods in the Amazon. *Proceedings of the National Academy of Sciences*, *112*(43), 13172–13177. https://doi.org/10.1073/pnas.1421010112

Edwards, W., & Krockenberger, A. (2006). Seedling mortality due to drought and fire
associated with the 2002 El Niño event in a tropical rain forest in north-east Queensland,
Australia. *Biotropica*, *38*(1), 16–26. https://doi.org/10.1111/j.1744-7429.2006.00098.x

Ehrlen, J., Morris, W. F., von Euler, T., & Dahlgren, J. P. (2016). Advancing environmentally
explicit structured population models of plants. *Journal of Ecology*, *104*(2), 292–305.
https://doi.org/10.1111/1365-2745.12523

Engelbrecht, B. M. J., Wright, S. J., & De Steven, D. (2002). Survival and ecophysiology of tree
seedlings during El Niño drought in a tropical moist forest in Panama. *Journal of Tropical
Ecology*, *18*, 569–579.

Esquivel-Muelbert, A., Baker, T. R., Dexter, K. G., Lewis, S. L., Brienen, R. J. W., Feldpausch, T.
R., Lloyd, J., Monteagudo-Mendoza, A., Arroyo, L., Álvarez-Dávila, E., Higuchi, N., Marimon, B.
S., Marimon-Junior, B. H., Silveira, M., Vilanova, E., Gloor, E., Malhi, Y., Chave, J., Barlow, J., …
Phillips, O. L. (2019). Compositional response of Amazon forests to climate change. *Global
Change Biology*, *25*(1), 39–56. https://doi.org/10.1111/gcb.14413

Esteban, E. J. L., Castilho, C. V., Melgaço, K. L., & Costa, F. R. C. (2021). The other side of
droughts: Wet extremes and topography as buffers of negative drought effects in an
Amazonian forest. *New Phytologist*, *229*(4), 1995–2006.
https://doi.org/10.1111/nph.17005

Evers, S. M., Knight, T. M., Inouye, D. W., Miller, T. E. X., Salguero-Gómez, R., Iler, A. M., &
Compagnoni, A. (2021). Lagged and dormant season climate better predict plant vital rates
than climate during the growing season. *Global Change Biology*, *27*(9), 1927–1941.
https://doi.org/10.1111/gcb.15519

Ewers, R. M., & Banks-Leite, C. (2013). Fragmentation impairs the microclimate buffering
effect of tropical forests. *PLoS ONE*, *8*(3), e58093.
https://doi.org/10.1371/journal.pone.0058093

Feeley, K. J., Malhi, Y., Zelazowski, P., & Silman, M. R. (2012). The relative importance of
deforestation, precipitation change, and temperature sensitivity in determining the future
distributions and diversity of Amazonian plant species. *Global Change Biology*, *18*(8), 2636–
2647. https://doi.org/10.1111/j.1365-2486.2012.02719.x

Gagnon, P. R., Bruna, E. M., Rubim, P., Darrigo, M. R., Littell, R. C., Uriarte, M., & Kress, W. J.
(2011). Growth of an understory herb is chronically reduced in Amazonian forest
fragments. *Biological Conservation*, *144*(2), 830–835.
https://doi.org/10.1016/j.biocon.2010.11.015

Gaoue, O. G., Horvitz, C. C., Steiner, U. K., & Tuljapurkar, S. (2019). Climate, rather than
human disturbance, is the main driver of age-specific mortality trajectories in a tropical
tree. *Ecological Modelling*, *400*, 34–40. https://doi.org/10.1016/j.ecolmodel.2019.03.007

Gasparrini, A. (2011). Distributed lag linear and non-linear models in R: The package dlnm.
Journal of Statistical Software, *43*(8), 1–20. https://doi.org/10.18637/jss.v043.i08

Gasparrini, A., Scheipl, F., Armstrong, B., & Kenward, M. G. (2017). A penalized framework
for distributed lag non-linear models: Penalized DLNMs. *Biometrics*, *73*(3), 938–948.
https://doi.org/10.1111/biom.12645

Gentry, A. H., & Dodson, C. (1987). Contribution of nontrees to species richness of a tropical
rain forest. *Biotropica*, *19*(2), 149–156. https://doi.org/10.2307/2388737

González-M, R., Posada, J. M., Carmona, C. P., Garzón, F., Salinas, V., Idárraga-Piedrahita, Á.,
Pizano, C., Avella, A., López-Camacho, R., Norden, N., Nieto, J., Medina, S. P., Rodríguez-M, G.
M., Franke-Ante, R., Torres, A. M., Jurado, R., Cuadros, H., Castaño-Naranjo, A., García, H., &
Salgado-Negret, B. (2020). Diverging functional strategies but high sensitivity to an
extreme drought in tropical dry forests. *Ecology Letters*, *n/a*(n/a).
https://doi.org/10.1111/ele.13659

Haddad, N. M., Brudvig, L. A., Clobert, J., Davies, K. F., Gonzalez, A., Holt, R. D., Lovejoy, T. E.,
Sexton, J. O., Austin, M. P., Collins, C. D., Cook, W. M., Damschen, E. I., Ewers, R. M., Foster, B.
L., Jenkins, C. N., King, A. J., Laurance, W. F., Levey, D. J., Margules, C. R., … Townshend, J. R.
(2015). Habitat fragmentation and its lasting impact on Earth's ecosystems. *Science*
Advances, *1*(2), e1500052. https://doi.org/10.1126/sciadv.1500052

Holyoak, M., & Heath, S. K. (2016). The integration of climate change, spatial dynamics, and
habitat fragmentation: A conceptual overview. *Integrative Zoology*, *11*(1), 40–59.
https://doi.org/https://doi.org/10.1111/1749-4877.12167

Horvitz, C. C., & Schemske, D. W. (1988). Demographic cost of reproduction in a Neotropical
herb: An experimental field study. *Ecology*, *69*(6), 1741–1745.
https://doi.org/10.2307/1941152

Huete, A. R., Didan, K., Shimabukuro, Y. E., Ratana, P., Saleska, S. R., Hutyra, L. R., Yang, W.,
Nemani, R. R., & Myneni, R. (2006). Amazon rainforests green-up with sunlight in dry
season. *Geophysical Research Letters*, *33*(6). https://doi.org/10.1029/2005GL025583

Janssen, T., van der Velde, Y., Hofhansl, F., Luyssaert, S., Naudts, K., Driessen, B., Fleischer,
K., & Dolman, H. (2021). Drought effects on leaf fall, leaf flushing and stem growth in
Neotropical forest; reconciling remote sensing data and field observations. *Biogeosciences*
Discussions, 1–41. https://doi.org/10.5194/bg-2021-30

Jones, M. O., Kimball, J. S., & Nemani, R. R. (2014). Asynchronous Amazon forest canopy
phenology indicates adaptation to both water and light availability. *Environmental Research*
Letters, *9*(12), 124021. https://doi.org/10.1088/1748-9326/9/12/124021

Kannenberg, S. A., Schwalm, C. R., & Anderegg, W. R. L. (2020). Ghosts of the past: How
drought legacy effects shape forest functioning and carbon cycling. *Ecology Letters*, *23*(5),
891–901. https://doi.org/10.1111/ele.13485

Klimešová, J., Martínková, J., & Ottaviani, G. (2018). Belowground plant functional ecology:
Towards an integrated perspective. *Functional Ecology*, *32*(9), 2115–2126.
https://doi.org/10.1111/1365-2435.13145

Kress, W. J. (1990). The diversity and distribution of *Heliconia* (Heliconiaceae) in Brazil. *Acta Botanica Brasileira*, *4*(1), 159–167.

Landau, W. M. (2021). The targets R package: A dynamic Make-like function-oriented pipeline toolkit for reproducibility and high-performance computing. *Journal of Open Source Software*, *6*(57), 2959. https://doi.org/ghwrqs

Laurance, W. F., Andrade, A. S., Magrach, A., Camargo, J. L. C., Campbell, M., Fearnside, P. M., Edwards, W., Valsko, J. J., Lovejoy, T. E., & Laurance, S. G. (2014). Apparent environmental synergism drives the dynamics of Amazonian forest fragments. *Ecology*, *95*(11), 3018–3026. https://doi.org/10.1890/14-0330.1

Laurance, W. F., Ferreira, L. V., Rankin de Merona, J. M., & Laurance, S. G. (1998). Rain forest fragmentation and the dynamics of Amazonian tree communities. *Ecology*, *79*(6), 2032–2040.

Laurance, W. F., Nascimento, H. E. M., Laurance, S. G., Andrade, A., Ribeiro, J. E. L. S., Giraldo, J. P., Lovejoy, T. E., Condit, R., Chave, J., Harms, K. E., & D'Angelo, S. (2006). Rapid decay of tree-community composition in Amazonian forest fragments. *Proceedings of the National Academy of Sciences*, *103*(50), 19010–19014. https://doi.org/10.1073/pnas.0609048103

Laurance, W. F., & Williamson, G. B. (2001). Positive feedbacks among forest fragmentation, drought, and climate change in the Amazon. *Conservation Biology*, *15*(6), 1529–1535. https://doi.org/10.1046/j.1523-1739.2001.01093.x

Laurance, W. F., Williamson, G. B., Delamônica, P., Oliveira, A., Lovejoy, T. E., Gascon, C., & Pohl, L. (2001). Effects of a strong drought on Amazonian forest fragments and edges. *Journal of Tropical Ecology*, *17*(6), 771–785.

Leitold, V., Morton, D. C., Longo, M., dos-Santos, M. N., Keller, M., & Scaranello, M. (2018). El Niño drought increased canopy turnover in Amazon forests. *New Phytologist*, *219*(3), 959–971. https://doi.org/10.1111/nph.15110

Marengo, J. A., Nobre, C. A., Tomasella, J., Oyama, M. D., Sampaio de Oliveira, G., de Oliveira, R., Camargo, H., Alves, L. M., & Brown, I. F. (2008). The drought of Amazonia in 2005. *Journal of Climate*, *21*(3), 495–516. https://doi.org/10.1175/2007JCLI1600.1

Markewitz, D., Devine, S., Davidson, E. A., Brando, P., & Nepstad, D. C. (2010). Soil moisture depletion under simulated drought in the Amazon: impacts on deep root uptake. *New Phytologist*, *187*(3), 592–607. https://doi.org/10.1111/j.1469-8137.2010.03391.x

Marra, G., & Wood, S. N. (2011). Practical variable selection for generalized additive models. *Computational Statistics & Data Analysis*, *55*(7), 2372–2387. https://doi.org/10.1016/j.csda.2011.02.004

McKee, T. B., Doesken, N. J., Kleist, J., & others. (1993). The relationship of drought frequency and duration to time scales. *Proceedings of the 8th Conference on Applied Climatology*, *17*, 179–183.

McPhaden, M. J. (1999). The child prodigy of 1997-98. *Nature*, *398*(6728), 559–561.
https://doi.org/10.1038/19193

Molowny-Horas, R., Suarez, M. L., & Lloret, F. (2017). Changes in the natural dynamics of
Nothofagus Dombeyi forests: Population modeling with increasing drought frequencies.
Ecosphere, *8*(3), e01708. https://doi.org/10.1002/ecs2.1708

Morris, W. F., & Doak, D. F. (2002). *Quantitative conservation biology: Theory and practice of*
population viability analysis. Sinauer.

Morris, W. F., Pfister, C. A., Tuljapurkar, S., Haridas, C. V., Boggs, C. L., Boyce, M. S., Bruna, E.
M., Church, D. R., Coulson, T., Doak, D. F., Forsyth, S., Gaillard, J.-M., Horvitz, C. C., Kalisz, S.,
Kendall, B. E., Knight, T. M., Lee, C. T., & Menges, E. S. (2008). Longevity can buffer plant and
animal populations against changing climatic variability. *Ecology*, *89*(1), 19–25.
https://doi.org/10.1890/07-0774.1

Mundim, F. M., & Bruna, E. M. (2016). Is there a temperate bias in our understanding of how
climate change will alter plant–herbivore interactions? A meta- analysis of experimental
studies. *American Naturalist*, *188*, S74–S89.

Nakazono, E. M., Bruna, E. M., & Mesquita, R. C. G. (2004). Experiemental harvesting of the
non-timber forest product *Ischnosiphon polyphyllus* in central Amazonia. *Forest Ecology and*
Management, *190*(2-3), 219–225. https://doi.org/10.1016/j.foreco.2003.10.013

Nihad, K., Krishnakumar, V., & Sheela, V. L. (2018). Relationship between stress and
flowering in *Heliconia* (*Heliconia stricta*). *International Journal of Agriculture Sciences(IJAS)*,
10(3), 5137. https://doi.org/10.9735/0975-3710.10.3.5137-5139

Ogle, K., Barber, J. J., Barron-Gafford, G. A., Bentley, L. P., Young, J. M., Huxman, T. E., Loik, M.
E., & Tissue, D. T. (2015). Quantifying ecological memory in plant and ecosystem processes.
Ecology Letters, *18*(3), 221–235. https://doi.org/10.1111/ele.12399

Oliver, T. H., Marshall, H. H., Morecroft, M. D., Brereton, T., Prudhomme, C., & Huntingford, C.
(2015). Interacting effects of climate change and habitat fragmentation on drought-
sensitive butterflies. *Nature Climate Change*, *5*(10), 941–945.
https://doi.org/10.1038/nclimate2746

Opdam, P., & Wascher, D. (2004). Climate change meets habitat fragmentation: Linking
landscape and biogeographical scale levels in research and conservation. *Biological*
Conservation, *117*(3), 285–297.

Paniw, M., James, T. D., Ruth Archer, C., Römer, G., Levin, S., Compagnoni, A., Che-Castaldo, J.,
Bennett, J. M., Mooney, A., Childs, D. Z., Ozgul, A., Jones, O. R., Burns, J. H., Beckerman, A. P.,
Patwary, A., Sanchez-Gassen, N., Knight, T. M., & Salguero-Gómez, R. (2021). The myriad of
complex demographic responses of terrestrial mammals to climate change and gaps of
knowledge: A global analysis. *Journal of Animal Ecology*. https://doi.org/10.1111/1365-
2656.13467

Pau, S., Wolkovich, E. M., Cook, B. I., Nytch, C. J., Regetz, J., Zimmerman, J. K., & Wright, S. J.
(2013). Clouds and temperature drive dynamic changes in tropical flower production.
Nature Climate Change, *3*(9), 838–842.

Payton, M. E., Greenstone, M. H., & Schenker, N. (2003). Overlapping confidence intervals or
standard error intervals: What do they mean in terms of statistical significance? *Journal of*
Insect Science, *3*(34), 1–6. https://doi.org/10.1673/031.003.3401

Pedersen, T. L. (2020). *Patchwork: The composer of plots* [Manual].

Phillips, O. L., van der Heijden, G., Lewis, S. L., López-González, G., Aragão, L. E. O. C., Lloyd, J.,
Malhi, Y., Monteagudo, A., Almeida, S., Dávila, E. A., Amaral, I., Andelman, S., Andrade, A.,
Arroyo, L., Aymard, G., Baker, T. R., Blanc, L., Bonal, D., de Oliveira, Á. C. A., … Vilanova, E.
(2010). Drought–mortality relationships for tropical forests. *New Phytologist*, *187*(3), 631–
646. https://doi.org/10.1111/j.1469-8137.2010.03359.x

Pierre, J.-S., Hullé, M., Gauthier, J.-P., & Rispe, C. (2020). Critical windows: A method for
detecting lagged variables in ecological time series. *Ecological Informatics*, *61*, 101178.
https://doi.org/10.1016/j.ecoinf.2020.101178

Pumisutapon, P., Visser, R. G. F., & de Klerk, G.-J. (2012). Moderate abiotic stresses increase
rhizome growth and outgrowth of axillary buds in *Alstroemeria* cultured in vitro. *Plant Cell,*
Tissue and Organ Culture (PCTOC), *110*(3), 395–400. https://doi.org/10.1007/s11240-012-
0160-7

R Core Team. (2020). *R: A language and environment for statistical computing*.

Rankin-de-Mérona, J. M., Prance, G. T., Hutchings, R. W., da Silva, M. F., Rodrigues, W. A., &
Uehling, M. E. (1992). Preliminary results of a large-scale tree inventory of upland rain
forest in the central Amazon. *Acta Amazonica*, *22*(4), 493–534.
https://doi.org/10.1590/1809-43921992224534

Ribeiro, M. B. N., Bruna, E. M., & Mantovani, W. (2010). Influence of post-clearing treatment
on the recovery of herbaceous plant communities in Amazonian secondary forests.
Restoration Ecology, *18*, 50–58.

Rundel, P. W., Cooley, A. M., Gerst, K. L., Riordan, E. C., Sharifi, M. R., Sun, J. W., & Tower, J. A.
(2020). Functional traits of broad-leaved monocot herbs in the understory and forest edges
of a Costa Rican rainforest. *PeerJ*, *8*, e9958. https://doi.org/10.7717/peerj.9958

Rundel, P. W., Sharifi, M. R., Gibson, A. C., & Esler, K. J. (1998). Structural and physiological
adaptation to light environments in Neotropical *Heliconia* (Heliconiaceae). *Journal of*
Tropical Ecology, *14*(6), 789–801. https://doi.org/10.1017/S0266467498000571

Salguero-Gómez, R., Jones, O. R., Archer, C. R., Buckley, Y. M., Che-Castaldo, J., Caswell, H.,
Hodgson, D., Scheuerlein, A., Conde, D. A., Brinks, E., Buhr, H., Farack, C., Gottschalk, F.,
Hartmann, A., Henning, A., Hoppe, G., Römer, G., Runge, J., Ruoff, T., … Vaupel, J. W. (2015).
The COMPADRE Plant Matrix Database: An open online repository for plant demography.
Journal of Ecology, *103*(1), 202–218. https://doi.org/10.1111/1365-2745.12334

Sapsford, S. J., Paap, T., Hardy, G. E. St. J., & Burgess, T. I. (2017). The 'chicken or the egg': Which comes first, forest tree decline or loss of mycorrhizae? *Plant Ecology*, *218*(9), 1093–1106. https://doi.org/10.1007/s11258-017-0754-6

Scariot, A. (1999). Forest fragmentation effects on palm diversity in central Amazonia. *Journal of Ecology*, *87*(1), 66–76. https://doi.org/10.1046/j.1365-2745.1999.00332.x

Scheffers, B. R., Edwards, D. P., Macdonald, S. L., Senior, R. A., Andriamahohatra, L. R., Roslan, N., Rogers, A. M., Haugaasen, T., Wright, P., & Williams, S. E. (2017). Extreme thermal heterogeneity in structurally complex tropical rain forests. *Biotropica*, *49*(1), 35–44. https://doi.org/10.1111/btp.12355

Schwalm, C. R., Anderegg, W. R. L., Michalak, A. M., Fisher, J. B., Biondi, F., Koch, G., Litvak, M., Ogle, K., Shaw, J. D., Wolf, A., Huntzinger, D. N., Schaefer, K., Cook, R., Wei, Y., Fang, Y., Hayes, D., Huang, M., Jain, A., & Tian, H. (2017). Global patterns of drought recovery. *Nature*, *548*(7666), 202–205. https://doi.org/10.1038/nature23021

Schwartz, N. B., Budsock, A. M., & Uriarte, M. (2019). Fragmentation, forest structure, and topography modulate impacts of drought in a tropical forest landscape. *Ecology*, *100*(6), e02677. https://doi.org/10.1002/ecy.2677

Scott, E. R., Wei, J.-P., Li, X., Han, W.-Y., & Orians, C. M. (2021). Differing non-linear, lagged effects of temperature and precipitation on an insect herbivore and its host plant. *Ecological Entomology*, *early view*. https://doi.org/10.1111/een.13023

Selwood, K. E., McGeoch, M. A., & Nally, R. M. (2015). The effects of climate change and land-use change on demographic rates and population viability. *Biological Reviews*, *90*(3), 837–853. https://doi.org/10.1111/brv.12136

Silva Junior, C. H. L., Pessôa, A. C. M., Carvalho, N. S., Reis, J. B. C., Anderson, L. O., & Aragão, L. E. O. C. (2021). The Brazilian Amazon deforestation rate in 2020 is the greatest of the decade. *Nature Ecology & Evolution*, *5*(2), 144–145. https://doi.org/10.1038/s41559-020-01368-x

Simpson, G. L. (2021). *Gratia: Graceful 'ggplot'-Based graphics and other functions for GAMs fitted using 'mgcv'* [Manual].

Sletvold, N. (2005). Density-dependent growth and survival in a natural population of the facultative biennial *Digitalis purpurea*. *Journal of Ecology*, *93*(4), 727–736. https://doi.org/10.1111/j.1365-2745.2005.01008.x

Snow, D. W. (1981). Tropical frugivorous birds and their food plants - a world survey. *Biotropica*, *13*(1), 1–14. https://doi.org/10.2307/2387865

Stiles, F. G. (1975). Ecology, flowering phenology, and hummingbird pollination of some Costa Rican *Heliconia* species. *Ecology*, *56*(2), 285–301. https://doi.org/10.2307/1934961

Stouffer, P. C., & Bierregaard, R. O. (1996). Forest fragmentation and seasonal patterns of hummingbird abundance in Amazonian Brazil. *Ararajuba*, *4*(1), 9–14.

Taubert, F., Fischer, R., Groeneveld, J., Lehmann, S., Müller, M. S., Rödig, E., Wiegand, T., &
Huth, A. (2018). Global patterns of tropical forest fragmentation. *Nature*, *554*(7693), 519–
522. https://doi.org/10.1038/nature25508

Teller, B. J., Adler, P. B., Edwards, C. B., Hooker, G., & Ellner, S. P. (2016). Linking
demography with drivers: Climate and competition. *Methods in Ecology and Evolution*, *7*(2),
171–183. https://doi.org/10.1111/2041-210X.12486

Tenhumberg, B., Crone, E. E., Ramula, S., & Tyre, A. J. (2018). Time-lagged effects of weather
on plant demography: Drought and *Astragalus scaphoides*. *Ecology*, *99*(4), 915–925.
https://doi.org/10.1002/ecy.2163

Ticktin, T. (2003). Relationships between El Niño Southern Oscillation and demographic
patterns in a substitute food for collared peccaries in Panama. *Biotropica*, *35*(2), 189–197.
https://doi.org/10.1111/j.1744-7429.2003.tb00278.x

Uriarte, M., Anciães, M., da Silva, M. T. B., Rubim, R., Johnson, E., & Bruna, E. M. (2011).
Disentangling the drivers of reduced long-distance seed dispersal by birds in an
experimentally fragmented landscape. *Ecology*, *92*(4), 924–937.
https://doi.org/10.1890/10-0709.1

Uriarte, M., Bruna, E. M., Rubim, P., Anciães, M., & Jonckheere, I. (2010). Effects of forest
fragmentation on the seedling recruitment of a tropical herb: assessing seed vs. safe-site
limitation. *Ecology*, *91*(5), 1317–1328. https://doi.org/10.1890/09-0785.1

Uriarte, M., Muscarella, R., & Zimmerman, J. K. (2018). Environmental heterogeneity and
biotic interactions mediate climate impacts on tropical forest regeneration. *Global Change
Biology*, *24*(2), e692–e704. https://doi.org/10.1111/gcb.14000

Uriarte, M., Schwartz, N., Powers, J. S., Marín-Spiotta, E., Liao, W., & Werden, L. K. (2016).
Impacts of climate variability on tree demography in second growth tropical forests: The
importance of regional context for predicting successional trajectories. *Biotropica*, *48*(6),
780–797. https://doi.org/10.1111/btp.12380

Vasconcelos, H. L., & Luizão, F. J. (2004). Litter production and litter nutrient
concentrations in a fragmented Amazonian landscape. *Ecological Applications*, *14*(3), 884–
892. https://doi.org/10.1890/03-5093

Vicente-Serrano, S. M., Beguería, S., & López-Moreno, J. I. (2010). A multiscalar drought
index sensitive to global warming: The standardized precipitation evapotranspiration
index. *Journal of Climate*, *23*(7), 1696–1718. https://doi.org/10.1175/2009JCLI2909.1

Westerband, A. C., Horvitz, C. C., & Gilliam, F. (2017). Early life conditions and precipitation
influence the performance of widespread understorey herbs in variable light
environments. *Journal of Ecology*, *105*(5), 1298–1308. https://doi.org/10.1111/1365-
2745.12757

Wickham, H. (2016). *Ggplot2: Elegant graphics for data analysis*. Springer-Verlag New York.

Williams, J. L., Jacquemyn, H., Ochocki, B. M., Brys, R., Miller, T. E. X., & Shefferson, R. (2015).
Life history evolution under climate change and its influence on the population dynamics of
a long-lived plant. *Journal of Ecology*, *103*(4), 798–808. https://doi.org/10.1111/1365-
2745.12369

Williamson, G. B., Laurance, W. F., Oliveira, A. A., Delamônica, P., Gascon, C., Lovejoy, T. E., &
Pohl, L. (2000). Amazonian tree mortality during the 1997 El Niño drought. *Conservation*
Biology, *14*(5), 1538–1542. https://doi.org/10.1046/j.1523-1739.2000.99298.x

Wood, S. N. (2017). *Generalized additive models: An introduction with R* (Second edition).
CRC Press/Taylor & Francis Group.

Wright, S. J. (1992). Seasonal drought, soil fertility and the species density of tropical forest
plant communities. *Trends in Ecology & Evolution*, *7*(8), 260–263.
https://doi.org/10.1016/0169-5347(92)90171-7

Wright, S. J., & Calderon, O. (2006). Seasonal, El Niño and longer term changes in flower and
seed production in a moist tropical forest. *Ecology Letters*, *9*(1), 35–44.
https://doi.org/10.1111/j.1461-0248.2005.00851.x

Wright, S. J., Carrasco, C., Calderon, O., & Paton, S. (1999). The El Niño Southern Oscillation
variable fruit production, and famine in a tropical forest. *Ecology*, *80*(5), 1632–1647.

Xavier, A. C., King, C. W., & Scanlon, B. R. (2016). Daily gridded meteorological variables in
Brazil (1980). *International Journal of Climatology*, *36*(6), 2644–2659.
https://doi.org/10.1002/joc.4518

Zartman, C. E., Amaral, J. A., Figueiredo, J. N., & Dambros, C. D. (2015). Drought impacts
survivorship and reproductive strategies of an epiphyllous leafy liverwort in central
Amazonia. *Biotropica*, *47*(2), 172–178.

Zeng, N., Yoon, J. H., Marengo, J. A., Subramaniam, A., Nobre, C. A., Mariotti, A., & Neelin, J. D.
(2008). Causes and impacts of the 2005 Amazon drought. *Environmental Research Letters*,
3(1), 1–9. https://doi.org/10.1088/1748-9326/3/1/014002