The myopathic transcription factor DUX4 induces the production of truncated RNA-

binding proteins in human muscle cells

Amy E. Campbell[1*], Michael C. Dyle[1*], Lorenzo Calviello[2*], Tyler Matheny[3], Michael A. Cortazar[1],

Thomas Forman[4,5], Rui Fu[3], Austin E. Gillen[3], Stephen N. Floor[2,6], and Sujatha Jagannathan[1,3†]

[1]Department of Biochemistry and Molecular Genetics, University of Colorado Anschutz Medical

Campus, Aurora, CO 80045, USA.

[2]Department of Cell and Tissue Biology, University of California, San Francisco, San Francisco,

CA 94143, USA.

[3]RNA Bioscience Initiative, University of Colorado Anschutz Medical Campus, Aurora, CO

80045, USA.

[4]Department of Craniofacial Biology, University of Colorado Anschutz Medical Campus, Aurora,

CO 80045, USA.

[5]Medical Scientist Training Program, University of Colorado Anschutz Medical Campus, Aurora,

CO 80045, USA.

[6]Helen Diller Family Comprehensive Cancer Center, University of California, San Francisco,

San Francisco, CA 94143, USA.

*These authors contributed equally

†Correspondence: sujatha.jagannathan@cuanschutz.edu

Running title: DUX4 promotes truncated protein production

Keywords: DUX4, FSHD, NMD, translation, splicing

ABSTRACT

DUX4 is an embryonic transcription factor whose misexpression in skeletal muscle causes facioscapulohumeral muscular dystrophy (FSHD). DUX4 induces the transcription of thousands of RNAs and dysregulates multiple pathways that could contribute to FSHD pathophysiology. However, lack of temporal data and the knowledge of which RNAs are actively translated following DUX4 expression has hindered our understanding of the cascade of events that lead to muscle cell death. Here, we interrogate the DUX4 transcriptome and translatome over time and find dysregulation of most key pathways as early as 4 hours after DUX4 induction, demonstrating the potent effect of DUX4 in disrupting muscle biology. We also observe extensive transcript downregulation as well as induction, and a high concordance between mRNA abundance and translation status. Significantly, DUX4 triggers widespread production of truncated protein products derived from aberrant RNAs that are degraded in normal muscle cells. One such protein, truncated serine/arginine-rich splicing factor 3 (SRSF3-TR), is present in FSHD muscle cells and disrupts splicing autoregulation when ectopically expressed in myoblasts. Taken together, the temporal dynamics of DUX4 induction show how the pathologic presence of an embryonic transcription factor in muscle cells alters gene expression to ultimately perturb RNA homeostasis.

INTRODUCTION

Facioscapulohumeral muscular dystrophy (FSHD) is a prevalent progressive myopathy caused by misexpression of an early embryonic transcription factor, DUX4, in skeletal muscle (Hamel & Tawil, 2018; Tawil, van der Maarel, & Tapscott, 2014). Sustained DUX4 expression is toxic to somatic cells and induces apoptotic death, leading to skeletal muscle atrophy in individuals with FSHD (Bosnakovski et al., 2008; Kowaljow et al., 2007; Rickard, Petek, & Miller, 2015; Wallace et al., 2011). In the decade since the elucidation of a unifying genetic model for FSHD (Lemmers et al., 2010), much work investigating the molecular consequences of DUX4 expression has led to the discovery of myriad altered genes and pathways that are associated with FSHD pathophysiology (Campbell, Belleville, Resnick, Shadle, & Tapscott, 2018; Lim, Nguyen, & Yokota, 2020). Efforts to determine which genes or pathways downstream of DUX4 misexpression cause FSHD are just beginning. Such understanding is critical for the development of effective therapeutics for FSHD.

Previous genome-wide studies of DUX4-mediated gene expression have been performed as endpoint assays in cell populations that have seen long periods of DUX4 misexpression and are likely already undergoing apoptosis (Bosnakovski et al., 2019; Geng et al., 2012; Jagannathan, Ogata, Gafken, Tapscott, & Bradley, 2019; Jagannathan et al., 2016; Lek et al., 2020; Resnick et al., 2019; Shadle et al., 2019; Shadle et al., 2017; Sharma, Harafuji, Belayew, & Chen, 2013; Whiddon, Langford, Wong, Zhong, & Tapscott, 2017). Although mechanisms underlying DUX4-induced cytotoxicity have been uncovered this way, it has also been shown that important pathologic processes occur in distinct temporal order. For example, inhibition of the RNA quality control pathway nonsense-mediated decay (NMD) by DUX4 was shown to occur early in the course of DUX4 expression while proteotoxic stress was a later event (Feng et al., 2015; Jagannathan et al., 2019). Therefore, genome-wide time course studies are needed to identify early changes in gene expression or cellular pathways that lead to skeletal muscle cell death.

While DUX4 induces widespread transcriptomic changes, proteomics shows that many genes display discordant transcript and protein levels (Jagannathan et al., 2019). Given the sparse nature of proteomics data, sequencing-based measurement of active translation is necessary to determine if the observed discordance between RNA and protein levels in DUX4-expressing cells is the result of altered translation regulation or protein stability. Such a measure would also reveal whether the aberrant RNAs stabilized by DUX4-mediated NMD inhibition (Feng et al., 2015) produce truncated proteins.

To identify early transcript- and translation-level changes induced by DUX4, we performed

paired RNA-sequencing (RNA-seq) and ribosome profiling (Ribo-seq) at 0, 4, 8, and 14 h

following the expression of DUX4 in MB135-iDUX4 human skeletal muscle myoblasts. MB135-

iDUX4 cells are a well-characterized model of FSHD that robustly recapitulates the

consequences of endogenous DUX4 expression in FSHD muscle (Jagannathan et al., 2016;

Yao et al., 2014). While RNA-seq measures transcript abundance, Ribo-seq measures

ribosome-protected RNA fragments, allowing quantification of ribosome density along an mRNA

that serves as a proxy for active translation (Ingolia, 2014). Ribo-seq also enables precise

delineation of translation start and end sites to characterize the protein products made from

aberrant RNAs.

We found that ~1600 genes show a significant change at the transcript level after 4 h of DUX4

induction, including many that are repressed; most pathways known to be misregulated by

DUX4 are altered at this early time point. We also found a high concordance of changes in

mRNA abundance and translation status, suggesting that post-transcriptional modulation by

DUX4 (Jagannathan et al., 2019) occurs primarily at the level of protein stability. Notably, the

hundreds of aberrant RNAs stabilized by DUX4-mediated inhibition of NMD (Feng et al., 2015)

are actively translated to produce truncated proteins, including truncated RNA-binding proteins

(RBPs) and splicing factors. We show that one such truncated splicing factor, truncated

serine/arginine-rich splicing factor 3 (SRSF3-TR), is expressed in FSHD muscle cell cultures

and perturbs splicing homeostasis when ectopically expressed in myoblasts. Together, our

results illustrate the importance of misregulated RNA quality control in DUX4-induced pathology.

 RESULTS

Identification of time points for paired RNA-seq and Ribo-seq in DUX4-expressing cells

Misexpression of DUX4 in skeletal muscle cells is cytotoxic (Geng et al., 2012; Jagannathan et

al., 2016; Rickard et al., 2015). To identify time points at which to measure transcript- and

translation-level changes induced by DUX4 before the onset of overt cytotoxicity, we utilized a

well-characterized doxycycline-inducible DUX4 human myoblast line, MB135-iDUX4

(Jagannathan et al., 2016), harboring a DUX4-responsive fluorescent reporter (**Figure 1A**). We

live imaged these cells every 15 min for 28 h following doxycycline treatment to induce DUX4

(**Figure 1B**, **Video 1**). In the absence of doxycycline, a very low level of 'leaky' DUX4

expression was observed, which is consistent with studies in other doxycycline-driven DUX4

induction systems (Bosnakovski, Chan, et al., 2017; Dandapat et al., 2014). In the presence of

doxycycline, expression of the DUX4-responsive fluorescent reporter was rapid and nearly

synchronous, with fluorescence detectable after 2 h. Most cells fluoresced after 4 h, with

fluorescence intensity increasing until 6 h at which point it plateaued. Cytotoxicity was first

observed 9 h following DUX4 induction (**Figure 1B**). Myoblasts continued to round up and

detach, resulting in a culture where most cells were dead or dying by 18 h. Given these data, we

limited our study to time points ≤ 14 h to identify early and direct gene expression changes

induced by DUX4.

To study the temporal trajectory of known DUX4-induced gene expression changes in important

cellular pathways, we carried out quantitative reverse transcription PCR (RT-qPCR) using RNA

extracted from the parental MB135-iDUX4 myoblast line. At 4 h we observed robust induction of

the DUX4 target gene ZSCAN4 (**Figure 1C**, top left) and repression of the myogenic program

(**Figure 1C**, top right), which is exquisitely sensitive to DUX4 misexpression (Bosnakovski et al.,

2018; Bosnakovski et al., 2008). In contrast, other consequences of DUX4 expression, such as

upregulation of pericentric human satellite II repeats (Shadle et al., 2019), activation of the

unfolded protein response (Jagannathan et al., 2019), perturbed Wnt signaling (C. R. Banerji et

al., 2015), and downregulation of oxidative stress response genes (Bosnakovski et al., 2008)

are more prominent at later time points (**Figure 1C**, remaining panels)**.** Based on our measures

of DUX4 activity, perturbation of downstream pathways, and myoblast cell death, 4, 8, and 14 h

were chosen as informative early, mid, and late time points to measure transcript- and

translation-level changes induced by misexpression of DUX4, with the 0 h time point serving as

a control. Therefore, we performed standard RNA-seq paired with Ribo-seq in parental MB135-

iDUX4 myoblasts induced to express DUX4 for 0, 4, 8, or 14 h (**Figure 1D**).

Figure 1. Synchronous expression of DUX4 in MB135-iDUX4 myoblasts enables time course analyses of downstream gene expression changes. (**A**) Schematic representation of the inducible DUX4 expression systems used in this study and the experimental outline to identify informative time points prior to the onset of cell death. (**B**) Still images from live cell fluorescence microscopy of MB135-iDUX4/ZSCAN4-mCherry myoblasts every 2 h following treatment with doxycycline to induce DUX4. Arrowheads indicate overtly dying cells. Scale bar, 150 μm. (**C**) Relative RNA levels of ZSCAN4, MYOD1, HSATII, HSPA5, TCF3, and HAGHL determined by RT-qPCR following treatment with doxycycline to induce DUX4 in MB135-iDUX4 myoblasts. Error bars denote the standard deviation from the mean of three biological replicates, which are shown as individual data points. (**D**) A schematic representation of the paired RNA-seq and Ribo-seq experiment. RNA was harvested from MB135-iDUX4 myoblasts treated with doxycycline to induce DUX4 for 0, 4, 8, or 14 h and either directly sequenced (RNA-seq) or digested with nuclease to degrade non-ribosome-protected RNA and the protected fragments sequenced (Ribo-seq).

DUX4 triggers early-onset disruption of key pathways involved in FSHD

We examined DUX4-induced transcriptome changes revealed by our RNA-seq dataset

(**Supplementary Table 1**). As expected, the housekeeping gene RPL27 had constant, robust

RNA expression throughout the time course **(Figure 2A**, top**).** Transcripts of a DUX4 target

gene, ZSCAN4, were absent in uninduced cells but highly expressed at 4 h and increased with

time, while transcripts from a myogenic gene, MYOD1, displayed the opposite trend (**Figure 2A**,

middle and bottom). Genome-wide, DUX4 altered the expression of thousands of transcripts,

with known DUX4 targets (Yao et al., 2014) showing increasing upregulation throughout the

time course (**Figure 2B**). Notably, there were 1674 genes whose expression significantly

changed after only 4 h of DUX4 induction with similar numbers activated and repressed. Genes

underlying pathways known to be involved in FSHD pathology are already significantly altered

at the 4 h time point (**Figure 2 – figure supplement 1A**), suggesting that DUX4 sets up

cascades of misregulation very early after its expression in muscle cells.

To identify novel pathways altered by DUX4, we used k-means clustering to group genes

significantly altered (defined as absolute log2 fold change > 1 and adjusted p-value < 0.01) at

any point during the time course into five clusters (**Figure 2 – figure supplement 1B**) and

carried out gene ontology (GO) analysis on each cluster (**Figure 2C-E**, **Supplementary Table**

2). We observed that Cluster 1, which is comprised of 181 genes rapidly induced upon DUX4

expression, returns GO categories that could underlie DUX4's normal role in establishing an

early embryonic program – negative regulation of cell differentiation, positive regulation of cell

proliferation, and DNA-templated transcription. The 692 genes that are rapidly silenced (Cluster

5) are enriched for GO categories related to myogenesis, positive regulation of cell

differentiation, and cytoskeleton organization. Together, this is suggestive of a general switch

away from a differentiated muscle program and towards a proliferative phenotype. Cluster 2

(583 genes) that is less robustly induced and Cluster 4 (2462 genes) that is less robustly

repressed are enriched for broad GO categories illustrative of the many fundamental cellular

processes that are altered upon DUX4 misexpression. Cluster 3 (2243 genes), which is induced

only at the late 14 h time point is enriched for GO categories mRNA splicing, ribonucleoprotein

transport, and ubiquitin-dependent process, which have previously been reported as some of

the major signatures of DUX4-induced gene expression (Geng et al., 2012; Jagannathan et al.,

2016; Rickard et al., 2015). Together, these time course RNA-seq data show that each pathway

misregulated by DUX4 has a unique temporal signature following DUX4 induction and suggest

that a combination of perturbed cellular systems underlies DUX4 toxicity.

Figure 2. Time course RNA-seq in MB135-iDUX4 myoblasts reveals early transcript-level changes in pathways underlying FSHD pathology. (**A**) RNA-seq read coverage over a time course of DUX4 expression for housekeeping gene RPL27 (top), DUX4 target gene ZSCAN4 (middle), and myogenic factor MYOD1 (bottom). (**B**) M-A plots for RNA-seq data after 4, 8, and 14 h of DUX4 induction compared to the 0 h control. DUX4 target status defined as in (Yao et al., 2014). (**C**) Heatmap depicting the clustering of all genes significantly differentially expressed upon DUX4 induction. The numbers within the heatmap represent the mean log2 fold change for the genes within that cluster. C, cluster. (**D**) Log2 fold change in RNA expression from the 0 h time point is shown for each gene in each cluster defined in (**C**). C, cluster. The thick black line represents the cluster mean. (**E**) GO analysis results of selected gene sets (biological process complete) that are significantly enriched in each cluster defined in (**C**).

Most DUX4-induced coding transcripts are robustly translated

A previous study that used stable isotope labeling by amino acids in cell culture (SILAC) mass spectrometry to measure the DUX4 proteome revealed extensive post-transcriptional gene regulation including discordant changes at the RNA versus protein level, particularly for genes

involved in RNA quality control (Jagannathan et al., 2019). While powerful, this study could not

provide a complete index of all expressed proteins due to inherent limitations in mass

spectrometry technology. Ribo-seq, by measuring which mRNAs are actively translated, is

capable of identifying the proteins being produced with greater depth than proteomics.

Additionally, DUX4 induces eIF2α phosphorylation at later time points in MB135 myoblasts

(Jagannathan et al., 2019; Shadle et al., 2017), which is thought to generally inhibit cap-

dependent translation (Proud, 2005), but the effect of this on protein expression in DUX4-

expressing cells is unknown. Therefore, we asked whether transcript level changes driven by

DUX4 misexpression were echoed at the level of translation by comparing the RNA-seq and

Ribo-seq datasets. To verify the quality of our Ribo-seq data, we confirmed that reads exhibited

the characteristic 3 nucleotide periodicity indicative of ribosome-protected RNA fragments

(**Figure 3 – figure supplement 1**). Representative Ribo-seq read coverage plots of the

housekeeping gene RPL27 showed constant, robust translation throughout the time course

(**Figure 3A**, top). In contrast, a DUX4 target gene, ZSCAN4, showed no coverage in uninduced

cells, low ribosome density beginning at 4 h, and active translation at 8 and 14 h (**Figure 3A**,

middle). MYOD1 was robustly translated at 0 h but rapidly downregulated (**Figure 3A**, bottom).

The changes in translation status at these specific genes mirrored the differences seen in their

mRNA levels (**Figure 2A**).

On a genome scale, DUX4 altered the translation status for thousands of transcripts, with later

time points showing larger differences and known DUX4 targets being translated at increasing

levels throughout the time course (**Figure 3B**, **Supplementary Table 3**). Strikingly, most genes

were concordantly up or downregulated at the level of transcript and translation status (**Figure**

3C, Materials and Methods) at all three time points with only a small number of genes showing

some discordance. GO analysis of the discordantly regulated genes returned significant results

only for the gene set (n = 137) that showed a mild translation downregulation at the 14 h time

point with pathways such as protein targeting to ER and viral transcription being enriched

(**Supplementary Table 3**). This likely represents a small subset of genes that are affected by

the inhibition of cap-dependent translation due to eIF2α phosphorylation. In contrast, a pairwise

correlation analysis of the RNA-seq, Ribo-seq and our previous SILAC proteomics

(Jagannathan et al., 2019) data demonstrated that both RNA and translation status are

discordant with protein levels (**Figure 3D**). Overall, these data show a high concordance of

transcript level and ribosome occupancy in DUX4-expressing cells, demonstrating that post-

transcriptional modulation by DUX4 likely occurs at the level of protein stability rather than

protein synthesis.

Figure 3. Ribo-seq shows high concordance between transcript levels and translation status. (**A**) Ribo-seq read coverage over a time course of DUX4 expression for housekeeping gene RPL27 (top), DUX4 target gene ZSCAN4 (middle), and myogenic factor MYOD1 (bottom). Green triangle, translation start. Red triangle, translation stop. (**B**) M-A plots for Ribo-seq data after 4, 8, and 14 h of DUX4 induction compared to the 0 h control. (**C**) Scatter plot of RNA-seq versus Ribo-seq log2 fold change after 4, 8, and 14 h of DUX4 expression. Significance defined as adjusted p-value < 0.01 for Ribo-seq fold change. (**D**) Pearson's correlation coefficient for the log2 fold change values computed for genes with quantification in all datasets (RNA-seq, Ribo-seq, and SILAC proteomics; n = 4854 genes). (**E**) Heatmap showing RNA-seq, Ribo-seq, and proteomics log2 fold change values of NMD factors.

We have previously shown that DUX4 induction leads to detectably lower protein abundance of

key NMD factors such as UPF1, SMG6, and XRN1, and that UPF1 is proteolytically degraded

(Feng et al., 2015; Jagannathan et al., 2019). We sought to determine if other NMD factors

follow a mechanism similar to UPF1 or if their downregulation could be due to lowered

translation of these proteins. Comparing RNA abundance from our RNA-seq data, translation

status from our Ribo-seq data, and protein level from our previous proteomics data

(Jagannathan et al., 2019) for all NMD factors showed that transcript level and translation is

highly concordant, while protein level is not (**Figure 3E**). This confirms that multiple key NMD

factors are indeed downregulated at the protein level without a corresponding change in their

RNA level or translation status, pointing to protein stability as the mode of regulation by DUX4.

DUX4 causes widespread truncated protein production

Having established that most transcripts induced by DUX4 are robustly translated, we wanted to

determine if truncated proteins are produced from RNAs containing premature translation

termination codons (PTCs) stabilized as a result of DUX4-mediated NMD inhibition (Feng et al.,

2015). Western blot analysis showed that levels of the key NMD factor UPF1 were reduced by

65% after only 4 h of DUX4 induction (**Figure 4A**). UPF1 levels fall further to 13% and 1% of

baseline at 8 and 14 h, respectively. This confirmed our previous observation that NMD

inhibition is an early event in the course of DUX4 expression (Feng et al., 2015), and suggested

that proteins resulting from the translation of NMD-targeted, stabilized aberrant RNAs should

appear in our dataset.

To ask if and when aberrant RNAs are translated, we used ORFquant (Calviello, Hirsekorn, &

Ohler, 2020), a new pipeline that identifies isoform-specific translation events from Ribo-seq

data. We then used DEXSeq (Anders, Reyes, & Huber, 2012) to conduct exon-level differential

analysis on the set of ORFquant-derived open reading frames, using Ribo-seq data. This

analysis identifies changes in relative exon usage to measure differences in the expression of

individual exons that are not simply the consequence of changes in overall transcript level. After

4 h of DUX4 induction 397 genes showed differential expression of specific exons, of which only

24 are predicted NMD targets (**Figure 4B**), whereas later time points showed a greater fraction

of exons that are unique to NMD targets as differentially expressed (**Supplementary Table 4**).

We grouped exons based on their NMD target status and calculated their fold change in

ribosome footprints at 4, 8, and 14 h of DUX4 expression compared to the 0 h time point

(**Figure 4C**). We observed a progressive and significant increase in the translation status of

NMD-targeted exons at 8 and 14 h, confirming the translation of stabilized aberrant RNAs in

DUX4-expressing cells.

Translation of an NMD target typically generates a prematurely truncated protein. To ask how

the specific truncated proteins being produced in DUX4-expressing myoblasts might functionally

impact cell homeostasis, we conducted GO analysis of the 74 truncated proteins being actively

translated at 14 h of DUX4 induction (**Figure 4D**, **Supplementary Table 4**). Strikingly, the

truncated proteins are enriched for genes encoding RBPs involved in mRNA metabolism and

specifically, splicing (**Supplementary Table 4**). Examples include genes such as IVNS1ABP,

SRSF3, SRSF6, and SRSF7 (**Figure 4E**). Thus, not only are NMD targets stabilized by DUX4

expression but they also produce truncated versions of many RBPs, which could have

significant downstream consequences to mRNA processing in DUX4-expressing cells.

Figure 4. Exon-level analysis of Ribo-seq data demonstrates robust translation of NMD-targeted aberrant RNAs. (**A**) Western blot analysis for UPF1, DUX4, and Histone H3 (loading control) over a time course of DUX4 expression following doxycycline induction in MB135-iDUX4 myoblasts. (**B**) M-A plot of exon-level analysis of Ribo-seq data from 4, 8, and 14 h of DUX4 induction. The x-axis represents mean expression calculated at the level of each exon within a gene. (**C**) Exon-level log2 fold change Ribo-seq values at 4, 8, or 14 h of DUX4 expression for canonical and NMD exons. Statistical testing was performed using a two-sided Wilcoxon test. (**D**) GO analysis results of selected gene sets (biological process complete) for all NMD targets that are translated at 14 h. (**E**) RNA-seq and Ribo-seq coverage over splicing-related genes IVNS1ABP, SRSF3, SRSF6, and SRSF7. The orange triangles denote PTCs.

Truncated SRSF3 is present in FSHD myotubes and perturbs RNA homeostasis

To explore the role truncated proteins play in DUX4-induced cellular phenotypes, we chose

SRSF3 for further characterization. SRSF3 is an SR family protein that possesses an N-terminal

RNA-binding RNA recognition motif (RRM) and a C-terminal arginine/serine (RS)-rich domain

responsible for protein-protein and protein-RNA interactions. SRSF3 is a multifunctional splicing

factor involved in transcriptional, co-transcriptional, and post-transcriptional regulation and has

been implicated in a variety of human pathologies including heart disease, Alzheimer's, and

cancer (More & Kumar, 2020; Zhou et al., 2020). Examination of our paired RNA-seq and Ribo-

seq data revealed robust expression of SRSF3 NMD-targeted exon 4 after 8 h of DUX4

induction, followed by robust translation of this exon that ends at the site of the PTC (**Figure**

5A).

To confirm translation of the aberrant SRSF3 RNA stabilized by DUX4, we carried out polysome

profiling using sucrose density gradient separation. As expected, the polysome profile after 14 h

of DUX4 expression showed a higher fraction of 80S monosomes compared to polysomes

(**Figure 5B**, top left), consistent with our observation of eIF2α phosphorylation and general

downregulation of translation at this and later time points of DUX4 expression (Jagannathan et

al., 2019; Shadle et al., 2017). We extracted RNA from various fractions and found that RPL27

mRNA was ribosome-bound in both control and DUX4-expressing cells, as evidenced by its

association with monosomes and polysomes and as expected for a housekeeping gene (**Figure**

5B, top right). In contrast, DUX4 mRNA and mRNA of the DUX4 target gene ZSCAN4 are highly

translated only in DUX4-expressing myoblasts, as indicated by their enrichment in heavy

polysomes (**Figure 5B**, middle). The normal SRSF3 isoform (SRSF3-Excl) is enriched in heavy

polysomes in both control and DUX4-expressing myoblasts but is less abundant in the latter

(**Figure 5B**, bottom left). In contrast, the NMD-targeted isoform of SRSF3 (SRSF3-Incl), also

enriched in heavy polysomes in both conditions, was significantly more abundant in DUX4-

expressing cells (**Figure 5B**, bottom right). These data confirm that aberrant SRSF3 mRNA is

being actively translated into truncated protein in DUX4-expressing myoblasts.

To determine if we could detect truncated SRSF3 protein, we generated a custom antibody

recognizing a 10 amino acid C-terminal neo-peptide unique to SRSF3-TR. This custom SRSF3-

TR antibody was able to recognize FLAG-tagged SRSF3-TR exogenously expressed in 293T

cells and endogenous SRSF3-TR immunoprecipitated from DUX4-expressing MB135-iDUX4

myoblasts (**Figure 5C**). We also used a commercial SRSF3 antibody that recognizes an N-

Figure 5. Truncated SRSF3 protein could disrupt RNA processing in FSHD myotubes. (**A**) RNA-seq and Ribo-seq coverage over SRSF3. The PTC-containing exon 4 is highlighted. The red hexagon indicates the normal stop codon while the orange hexagon denotes the PTC. (**B**) Absorbance at 254 nm across a sucrose density gradient of lysates from control MB135-iDUX4 myoblasts and MB135-iDUX4 myoblasts expressing DUX4 for 14 h (top left). RT-qPCR measurement of RPL27, DUX4, ZSCAN4, SRSF3-Excl, and SRSF3-Incl mRNA levels in DUX4-expressing myoblasts relative to control myoblasts from collected fractions (remaining panels). (**C**) Detection of SRSF3 and SRSF3-TR in whole cell extracts from 293T cells exogenously expressing FLAG-tagged full-length (FL) or truncated (TR) SRSF3 as compared to protein lysates from MB135-iDUX4 myoblasts treated with (+) or without (-) doxycycline (DOX) to induce DUX4 and immunoprecipitated with a custom anti-SRSF3-TR antibody, no antibody (Ab), or a commercial SRSF3 antibody. IP, immunoprecipitation. Asterisks denote proteins of interest. (**D**) Immunofluorescence staining in MB135 control and MB200 FSHD myotubes differentiated for 72 h and stained with DAPI (blue) and rabbit anti-DUX4 (green) or custom rabbit anti-SRSF3-TR (pink) antibody. (**E**) SRSF3-Excl and -Incl mRNA isoform levels as determined by RT-qPCR following induction of full-length (FL) or truncated (TR) SRSF3 in MB135-iFLAG-SRSF3-FL or -TR myoblasts by treatment with doxycycline for 72 h.

terminal epitope common to both the full-length and truncated SRSF3. This antibody detected

both exogenously expressed full-length and truncated FLAG-SRSF3, and endogenous full-

length SRSF3, but was insufficient to visualize endogenous SRSF3-TR (**Figure 5C**), possibly

due to lower affinity for this protein isoform in an immunoprecipitation assay. To determine if

SRSF3-TR was present in FSHD myotubes expressing endogenous levels of DUX4, we carried

out immunofluorescence for SRSF3-TR or DUX4 in differentiated FSHD and control muscle

cells. While there was no detectable SRSF3-TR staining in control cells, in DUX4-expressing

FSHD cultures SRSF3-TR appeared in cytoplasmic puncta (**Figure 5D**). Together, these data

show that endogenous SRSF3-TR is present at detectable levels in DUX4-expressing MB135-

iDUX4 and FSHD cells.

To explore the functional consequences of SRSF3-TR expression, we exogenously expressed

FLAG-tagged full-length or truncated SRSF3 in healthy muscle cells. As previously described,

full-length SRSF3 decreased the level of the normal SRSF3 mRNA isoform and increased the

aberrant isoform ((Jumaa & Nielsen, 1997); **Figure 5E**). Strikingly, SRSF3-TR led to a 5-fold

upregulation of the endogenous SRSF3 aberrant mRNA isoform (**Figure 5F**). These data

suggest that DUX4-induced translation of SRSF3-TR may feedback to create more aberrant

RNA and therefore more truncated protein, highlighting the importance of misregulated post-

transcriptional processes in DUX4-induced pathology.

DISCUSSION

We paired RNA-seq and Ribo-seq across a time course of DUX4 expression in MB135-iDUX4 human skeletal muscle myoblasts, providing an integrative temporal view of the transcriptome and translatome in a well-accepted cellular model of FSHD. Our choice to examine early time points, before overt DUX4-induced cytotoxicity, provides a view into early DUX4-driven regulation that builds on prior work at later time points (Bosnakovski et al., 2019; Chew et al., 2019; DeSimone, Leszyk, Wagner, & Emerson, 2019; Geng et al., 2012; Jagannathan et al., 2019; Jagannathan et al., 2016; Lek et al., 2020; Resnick et al., 2019; Shadle et al., 2019; Shadle et al., 2017; Sharma et al., 2013; Whiddon et al., 2017). Taken together, our results demonstrate that critical pathways underlying FSHD pathophysiology are perturbed earlier than previously understood and provide datasets that should serve as an informative resource for the FSHD community as it works to develop effective FSHD therapeutics.

Our time course RNA-seq data revealed a large number of both up and downregulated genes after only 4 h of DUX4 expression. While the former result is expected given DUX4's role as a transcriptional activator (Geng et al., 2012), this degree of early gene silencing is remarkable and would require rapid cessation of transcription. There is no evidence that DUX4 acts as a direct repressor of transcription (Bosnakovski et al., 2018); therefore, our results suggest that there exists one or more unstudied DUX4-induced transcriptional repressors responsible for early gene downregulation. Indeed, of the 883 genes significantly upregulated by DUX4 at 4 h, 28 are DNA-binding transcription factors known to effect gene repression and at least 5 more are DNA-binding proteins with an as-yet-undetermined role in gene regulation. This novel hypothesis provides a mechanism by which DUX4 inhibits gene expression that is unique from previously suggested models invoking competition of DUX4 with the homeodomain transcription factors PAX3 and PAX7 (C. R. S. Banerji et al., 2017; Bosnakovski, Daughters, Xu, Slack, & Kyba, 2009; Bosnakovski, Toso, et al., 2017; Bosnakovski et al., 2008) or transcriptional interference from DUX4-induced non-coding transcripts (Bosnakovski et al., 2018). Future work could determine which, if any, of the DNA-binding repressors induced by DUX4 cause early DUX4-mediated gene downregulation.

The profound consequences of DUX4 misexpression on cell identity were also revealed in our time course RNA-seq results. GO analyses showed that DUX4-expressing myoblasts begin to turn on genes that promote cell proliferation and inhibit cell differentiation, and silence genes that promote myogenesis, early after DUX4 induction. Thus, skeletal muscle – a post-mitotic,

differentiated cell type – is pushed towards a proliferative, naïve embryonic state. This conflicted
cell identity may contribute to apoptosis, as has been proposed before (Ashoti, Alemany, Sage,
& Geijsen, 2021; Geng et al., 2012). Interestingly, a few hours after DUX4 begins forcing
expression of an embryonic gene expression program, and just as overt apoptosis appears,
cells respond by altering core regulatory processes such as RNA splicing and localization, and
protein ubiquitination. Whether these changes underlie cellular "coping" mechanisms that delay
death or underpin cytotoxicity remains to be determined.
346
We recently described the DUX4-induced proteome using MB135-iDUX4 cells (Jagannathan et
al., 2019). Although this catalog of expressed proteins remains incomplete due to partial
coverage and restricted dynamic range inherent to mass spectrometry, our proteomics data
demonstrated that hundreds of genes are post-transcriptionally modulated by DUX4. But, these
experiments could not determine if this regulation was mediated at the level of protein synthesis
or degradation. The time course Ribo-seq presented here complements and extends this earlier
work, and suggests that most DUX4 post-transcriptional regulation, including of multiple key
NMD factors, occurs via proteolysis. We did uncover a small number of genes regulated at the
level of protein synthesis at the 14 h time point of DUX4 expression, likely representing the
subset of transcripts impacted by eIF2α phosphorylation induced by DUX4, as previously
reported (Jagannathan et al., 2019; Shadle et al., 2017).
358
Loss of NMD leads to the stabilization of aberrant RNAs (Kurosaki & Maquat, 2016). However,
few studies have looked at whether aberrant RNAs are translated and what proteins they might
produce. Here, we demonstrate that DUX4-induced NMD inhibition causes truncated protein
production in muscle cells. The confirmed existence of truncated proteins in DUX4-expressing
cells has implications for how we understand cytotoxicity. Protein truncation could result in a
dominant negative function that inhibits the activity of the remaining, cell critical full-length
protein. Truncated proteins might misfold and facilitate the formation of protein aggregates, such
as those observed in FSHD myotubes (Homma, Beermann, Boyce, & Miller, 2015; Homma,
Beermann, Yu, Boyce, & Miller, 2016; Shadle et al., 2017). Additionally, some DUX4-induced
truncated proteins contain unique C-terminal extensions, or neo-peptides, that could serve as
novel antigens and might induce inflammation in FSHD muscle. Importantly, we were able to
detect truncated proteins in FSHD muscle cell cultures, thereby validating our findings from the
inducible DUX4 system in a more physiological setting and raising the possibility that these
molecules could be used clinically as functionally relevant and FSHD-specific biomarkers.

Future work is required to establish whether truncated proteins, either individually or in
combination, contribute to cell death.
375
Many of the identified DUX4-induced truncated proteins are RBPs and splicing factors. It is well-
established that DUX4 alters RNA splicing (Geng et al., 2012; Jagannathan et al., 2019;
Jagannathan et al., 2016; Rickard et al., 2015) and therefore interesting to speculate that
truncated RBPs and splicing proteins might be responsible for inducing global RNA processing
defects. Such misprocessing would generate aberrant RNAs that could act to further overwhelm
the already inhibited NMD pathway. As an example of this phenomenon, we observed that
exogenous expression of the truncated splicing factor SRSF3-TR causes a significant increase
in the level of the corresponding aberrant SRSF3 RNA. Splicing factors are known to regulate
and coordinate their expression via various forms of autoregulation (Konigs et al., 2020; Leclair
et al., 2020; Muller-McNicoll, Rossbach, Hui, & Medenbach, 2019). Overexpression of full-length
SRSF3 leads to a reduction in the level of normal SRSF3 transcript while at the same time
activating the production of aberrant SRSF3 RNA that gets degraded by NMD in a balancing act
mechanism that prevents SRSF3 accumulation (Jumaa & Nielsen, 1997). Our results suggest
that SRSF3-TR facilitates its own production and disrupts normal autoregulatory processes,
leading to a new model in which DUX4-expressing cells hijack the splicing protein
autoregulatory network and flip it – instead of acting to limit the production of full-length protein
the creation of truncated protein is amplified.
393
In summary, our study provides a critical missing piece towards a comprehensive
characterization of the major steps in DUX4-induced gene expression. We demonstrate that
DUX4 induces cascading changes at all levels of gene regulation that are invisible at steady
state, offering a glimpse into how developmentally regulated transcription factors expressed at
the wrong time in the wrong context can wreak havoc leading to human disease.

MATERIALS AND METHODS
400
401 **RESOURCE AVAILABILITY**
402
403 **Lead contact**
404 Further information and requests for resources and reagents should be directed to and will be
405 fulfilled by the lead contact, Sujatha Jagannathan (sujatha.jagannathan@cuanschutz.edu).
406
407 **Materials availability**
408 The cell lines and antibody generated in this study are available upon request. Plasmids
409 generated in this study have been deposited to Addgene (plasmid #171951, #171952, #172345,
410 and #172346).
411
412 **Data and code availability**
413 The RNA-seq and Ribo-seq data generated during this study are available at GEO (accession
414 number GSE178761). The code generated during this study are available at GitHub
415 (https://github.com/sjaganna/2021-campbell_dyle_calviello_et_al).
416
417 **EXPERIMENTAL MODEL AND SUBJECT DETAILS**
418
Cell lines and culture conditions
293T cells were obtained from ATCC (CRL-3216). MB135, MB135-iDUX4, MB135-
iDUX4/ZSCAN4-mCherry, and MB200 immortalized human myoblasts were a gift from Dr.
Stephen Tapscott and originated from the Fields Center for FSHD and Neuromuscular
Research at the University of Rochester Medical Center. MB135-iDUX4 cells have been
described previously (Jagannathan et al., 2016). MB135-iFLAG-SRSF3-FL, and MB135-iFLAG-
SRSF3-TR immortalized human myoblasts were generated in this study. All parental cell lines
were authenticated by karyotype analysis and determined to be free of mycoplasma by PCR
screening. 293T cells were maintained in Dulbecco's Modified Eagle Medium (DMEM) (Thermo
Fisher Scientific) supplemented with 10% EqualFETAL (Atlas Biologicals). Myoblasts were
maintained in Ham's F-10 Nutrient Mix (Thermo Fisher Scientific) supplemented with 20% Fetal
Bovine Serum (Thermo Fisher Scientific), 10 ng/mL recombinant human basic fibroblast growth
factor (Promega), and 1 µM dexamethasone (Sigma-Aldrich). MB135-iDUX4/ZSCAN4-mCherry
and MB135-iDUX4 myoblasts were additionally maintained in 2 µg/mL puromycin

dihydrochloride (VWR). MB135-iFLAG-SRSF3-FL and -TR myoblasts were additionally

maintained in 10 µg/mL blasticidin S HCl (Thermo Fisher Scientific). Induction of DUX4 and

SRSF3 transgenes was achieved by culturing cells in 1 µg/mL doxycycline hyclate (Sigma-

Aldrich). Differentiation of myoblasts into myotubes was achieved by switching the fully

confluent myoblast monolayer into DMEM containing 1% horse serum (Thermo Fisher

Scientific) and Insulin-Transferrin-Selenium (Thermo Fisher Scientific). All cells were incubated

at 37 °C with 5% CO_2.

METHOD DETAILS

Cloning

pTwist-FLAG-SRSF3_Full.Length_Codon.Optimized and pTwist-FLAG-

SRSF3_Truncated_Codon.Optimized plasmids were synthesized by Twist Bioscience. To

construct pCW57.1-FLAG-SRSF3_Full.Length_Codon.Optimized-Blast and pCW57.1-FLAG-

SRSF3_Truncated_Codon.Optimized-Blast plasmids, the SRSF3 open reading frames were

subcloned into pCW57-MCS1-P2A-MCS2 (Blast) (a gift from Adam Karpf, Addgene plasmid

#80921) (Barger, Branick, Chee, & Karpf, 2019) by restriction enzyme digest using EcoRI and

BamHI (New England Biolabs).

Antibody generation

Purified SRSF3-TR peptide (Cys-PRRRVTIMSLLTTL) was used as an immunogen and

polyclonal rabbit anti-SRSF3-TR antibody production was done in collaboration with Pacific

Immunology (Ramona, CA). The antisera from all animals were screened for reactivity by ELISA

against the immunogen and with western blots and immunofluorescence against transfected

SRSF3-TR.

Transgenic cell line generation

Lentiviral particles expressing doxycycline-inducible FLAG-SRSF3-FL or -TR transgenes were

generated by co-transfecting 293T cells with the appropriate lentivector, pMD2.G (a gift from

Didier Trono, Addgene plasmid #12259), and psPAX2 (a gift from Didier Trono, Addgene

plasmid #12260) using Lipofectamine 2000 Transfection Reagent (Thermo Fisher Scientific). To

generate polyclonal SRSF3 transgenic cell lines, MB135 myoblasts were transduced with

lentivirus in the presence of 8 µg/mL polybrene (Sigma-Aldrich) and selected using 10 µg/mL

blasticidin S HCl.

Plasmid transfections

293T cells were transfected with pTwist-FLAG-SRSF3_Full.Length_Codon.Optimized and

pTwist-FLAG-SRSF3_Truncated_Codon.Optimized plasmids using Lipofectamine 2000

Transfection Reagent following the manufacturer's instructions.

Live cell imaging

MB135-iDUX4/ZSCAN4-mCherry myoblasts were induced with doxycycline hyclate to turn on

DUX4 expression and subjected to time lapse imaging using the IncuCyte S3 incubator

microscope system (Sartorius). Images were collected every 15 min from the time of

doxycycline addition (t = 0 h) to 28 h.

RNA extraction and RT-qPCR

Total RNA was extracted from whole cells using TRIzol Reagent (Thermo Fisher Scientific)

following the manufacturer's instructions. Isolated RNA was treated with DNase I (Thermo

Fisher Scientific) and reverse transcribed to cDNA using SuperScript III reverse transcriptase

(Thermo Fisher Scientific) and random hexamers (Thermo Fisher Scientific) according to the

manufacturer's protocol. Quantitative PCR was carried out on a CFX384 Touch Real-Time PCR

Detection System (Bio-Rad) using primers specific to each gene of interest and iTaq Universal

SYBR Green Supermix (Bio-Rad). The expression levels of target genes were normalized to

that of the reference gene *RPL27* using the delta-delta-Ct method (Livak & Schmittgen, 2001).

The primers used in this study are listed in the Key Resources Table.

RNA-seq library preparation and sequencing

Total RNA was extracted from whole cells using TRIzol Reagent following the manufacturer's

instructions. Isolated RNA was subjected to ribosomal RNA depletion using the Ribo-Zero rRNA

Removal Kit (Illumina). RNA-seq libraries were prepared using the NEXTflex Rapid Directional

qRNA-Seq Kit (Bioo Scientific) following the manufacturer's instructions and sequenced using

75 bp single-end sequencing on the Illumina NextSeq 500 platform by the BioFrontiers Institute

Next-Gen Sequencing Core Facility.

Ribosome footprinting

Ribo-seq was performed as described previously (Calviello et al., 2020) using six 70% confluent

10 cm dishes of MB135-iDUX4 cells per condition. Briefly, cells were washed with ice-cold

phosphate-buffered saline (PBS) supplemented with 100 µg/mL cycloheximide (Sigma-Aldrich),

flash frozen on liquid nitrogen, and lysed in Lysis Buffer (PBS containing 1% (v/v) Triton X-100

and 25 U/mL TurboDNase (Ambion)). Cells were harvested by scraping and further lysed by

trituration ten times through a 26-gauge needle. The lysate was clarified by centrifugation at

20,000 g for 10 min at 4 °C. The supernatants were flash frozen in liquid nitrogen and stored at -

80 °C. Thawed lysates were treated with RNase I (Ambion) at 2.5 U/μL for 45 min at room

temperature with gentle mixing. Further RNase activity was stopped by addition of SUPERaseIn

RNase Inhibitor (Thermo Fisher Scientific). Next, ribosome complexes were enriched using

MicroSpin S-400 HR Columns (GE Healthcare) and RNA extracted using the Direct-zol RNA

Miniprep Kit (Zymo Research). Ribo-Zero rRNA Removal Kit was used to deplete rRNAs and

the ribosome-protected fragments were recovered by running them in a 17% Urea gel, staining

with SYBR Gold (Invitrogen), and extracting nucleic acids that are 27 to 30 nucleotides long

from gel slices by constant agitation in 0.3 M NaCl at 4 °C overnight. The recovered nucleic

acids were precipitated with isopropanol using GlycoBlue Coprecipitant (Ambion) as carrier and

treated with T4 polynucleotide kinase (Thermo Fisher Scientific). Libraries were prepared using

the NEXTflex Small RNA-Seq Kit v3 (Bioo Scientific) following the manufacturer's instructions

and sequenced using 75 bp single-end reads on an Illumina NextSeq 500 by the BioFrontiers

Institute Next-Gen Sequencing Core Facility.

RNA-seq and Ribo-seq data analysis

Fastq files were stripped of the adapter sequences using cutadapt. UMI sequences were

removed, and reads were collapsed to fasta format. Reads were first aligned against rRNA

(accession number U13369.1), and to a collection of snoRNAs, tRNAs, and miRNA (retrieved

using the UCSC table browser) using bowtie2 (Langmead & Salzberg, 2012). Remaining reads

were mapped to the hg38 version of the genome (without scaffolds) using STAR 2.6.0a (Dobin

et al., 2013) supplied with the GENCODE 25 .gtf file. A maximum of two mismatches and

mapping to a minimum of 50 positions was allowed. De-novo splice junction discovery was

disabled for all datasets. Only the best alignment per each read was retained. Quality control

and read counting of the Ribo-seq data was performed with Ribo-seQC (Calviello, Sydow,

Harnett, & Ohler, 2019).

Differential gene expression analysis of the RNA-seq data was conducted using DESeq2 (Love,

Huber, & Anders, 2014). Briefly, featureCounts from the subread R package (Liao, Smyth, &

Shi, 2014) was used to assign aligned reads (in BAM format) to genomic features supplied with

the GENCODE 25. gtf file. The featureCounts output was then supplied to DESeq2 and

differential expression analysis was conducted with the 0 h time point serving as the reference
sample. Genes with very low read count were filtered out by requiring at least a total of 10 reads
across the 12 samples (3 replicates each of the 0, 4, 8, and 14 h samples). Log2 fold change
shrinkage was done using the apeglm function (Zhu, Ibrahim, & Love, 2019).

Differential analysis of the RNA-seq and Ribo-seq data was performed using DESeq2, as
previously described (Calviello et al., 2021; Chothani et al., 2019), using an interaction model
between the tested condition and RNA-seq – Ribo-seq counts. Only reads mapping uniquely to
coding sequence regions were used. In addition, ORFquant (Calviello et al., 2020) was used to
derive de-novo isoform-specific translation events, by pooling the Ribo-seQC output from all
Ribo-seq samples, using uniquely mapping reads. DEXSeq (Anders et al., 2012) was used to
perform differential exon usage along the DUX4 time course data, using Ribo-seq counts on
exonic bins and junctions belonging to different ORFquant-derived translated regions. NMD
candidates were defined by ORFquant as open reading frames ending with a stop codon
upstream of an exon-exon junction.

GO category analysis
Gene Ontology (GO) analysis was conducted using the web tool http://geneontology.org,
powered by pantherdb.org. Briefly, statistical overrepresentation test using the complete GO
biological process annotation dataset was conducted and p-values were calculated using the
Fisher's exact test and False Discovery Rate was calculated by the Benjamini-Hochberg
procedure.

Polysome profiling
Polysome profiling was performed as previously described (Merrick & Hensold, 2001; Miura,
Andrews, Holcik, & Jasmin, 2008) with the following modifications. Four 70% confluent 15 cm
dishes of MB135-iDUX4 cells per condition were treated with 100 μg/mL cycloheximide for 10
min, transferred to wet ice, washed with ice-cold PBS containing 100 μg/mL cycloheximide, and
then lysed in 400 μL Lysis Buffer (20 mM HEPES pH 7.4, 15 mM MgCl$_2$, 200 mM NaCl, 1%
Triton X-100, 100 μg/mL cycloheximide, 2 mM DTT, and 100 U/mL SUPERaseIn RNase
Inhibitor) per 15 cm dish. The cells and buffer were scraped off the dish and centrifuged at
13,000 rpm for 10 min at 4 °C. Lysates were fractionated on a 10% to 60% sucrose gradient
using the SW 41 Ti Swinging-Bucket Rotor (Beckman Coulter) at 36,000 rpm for 3 h and 10
min. Twenty-four fractions were collected using a Gradient Station ip (BioComp) and an FC

203B Fraction Collector (Gilson) with continuous monitoring of absorbance at 254 nm. RNA

from each fraction was extracted using TRIzol LS Reagent (Thermo Fisher Scientific) following

the manufacturer's instructions. RT-qPCR was carried out as described above.

Protein extraction

Total protein was extracted from whole cells using TRIzol Reagent following the manufacturer's

instructions, excepting that protein pellets were dissolved in Protein Resuspension Buffer (0.5 M

Tris base, 5% SDS). Isolated protein was quantified using the Pierce BCA Protein Assay Kit

(Thermo Fisher Scientific) according to the manufacturer's protocol. Protein was mixed with 4X

NuPAGE LDS Sample Buffer (Thermo Fisher Scientific) containing 50 mM DTT and heated to

70 °C before immunoblotting.

Immunoprecipitation

MB135-iDUX4 myoblasts were treated with or without doxycycline for 14 h and then trypsinized

prior to lysis on ice in 1 mL of Lysis Buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 1% NP-40)

containing protease inhibitors (Sigma Aldrich). Lysates were precleared using Protein G

Sepharose (Thermo Fisher Scientific) for 1 h prior to an overnight incubation at 4 °C with either

anti-SRSF3 or anti-SRSF3-TR antibody. Protein G Sepharose was added the following morning

for 5 h to bind the antibody, and beads were subsequently washed 5 times with 1 mL cold Lysis

Buffer. After the final wash, 4X NuPAGE LDS Sample Buffer containing 50 mM DTT was added

directly to the beads and samples heated to 70 °C for protein elution before immunoblotting.

Immunoblotting

Protein was run on NuPAGE Bis-Tris precast polyacrylamide gels (Thermo Fisher Scientific)

alongside PageRuler Plus Prestained Protein Ladder (Thermo Fisher Scientific) and transferred

to Odyssey nitrocellulose membrane (LI-COR Biosciences). Membranes were blocked in

Intercept (PBS) Blocking Buffer (LI-COR Biosciences) before overnight incubation at 4 °C with

primary antibodies diluted in Blocking Buffer containing 0.2% Tween 20. Membranes were

incubated with IRDye-conjugated secondary antibodies (LI-COR Biosciences) for 1 h and

fluorescent signal visualized using a Sapphire Biomolecular Imager (Azure Biosystems) and

Sapphire Capture software (Azure Biosystems). When appropriate, membranes were stripped

with Restore Western Blot Stripping Buffer (Thermo Fisher Scientific) before being re-probed.

Band intensities were quantified by densitometry using ImageJ (Schneider, Rasband, & Eliceiri,

2012).

Immunofluorescence

Cells were fixed in 10% Neutral Buffered Formalin (Research Products International) for 30 min and permeabilized for 10 min in PBS with 0.1% Triton X-100. Samples were then incubated overnight at 4 °C with primary antibodies, followed by incubation with 488- or 594-conjugated secondary antibodies for 1 h prior to counterstaining and mounting with Prolong Diamond Antifade Mountant with DAPI (Thermo Fisher Scientific). Slides were imaged with a DeltaVision Elite deconvolution microscope, CoolSNAP HQ^2 high-resolution CCD camera, and Resolve3D softWoRx-Acquire v7.0 software. Image J software (Schneider et al., 2012) was used for image analysis.

Antibodies

The antibodies used in this study are anti-DUX4 (Abcam 124699), anti-Histone H3 (Abcam 1791), anti-SRSF3 (Thermo Fisher Scientific 33-4200), anti-SRSF3-TR (this paper), anti-RENT1/hUPF1 (Abcam ab109363), Drop-n-Stain CF 488A Donkey Anti-Rabbit IgG (Biotium 20950), Drop-n-Stain CF 594 Donkey Anti-Rabbit IgG (Biotium 20951), IRDye 650 Goat anti-Mouse IgG Secondary Antibody (LI-COR Biosciences 926-65010), and IRDye 800CW Goat anti-Rabbit IgG Secondary Antibody (LI-COR Biosciences 926-32211).

QUANTIFICATION AND STATISTICAL ANALYSIS

Data analysis, statistical tests, and visualization

All data analysis and statistical tests were performed in the R programming environment and relied on Bioconductor (Huber et al., 2015) and ggplot2 (Wickham, 2016). Plots were generated using R plotting functions and/or the ggplot2 package. Bar and line graphs were generated using GraphPad Prism software version 9.0. Biological replicates were defined as experiments performed separately on distinct samples (i.e. cells cultured in different wells) representing identical conditions and/or time points. No outliers were eliminated in this study. All statistical tests were performed using R functions.

ACKNOWLEDGEMENTS

We thank Stephen Tapscott for the MB135-iDUX4/ZSCAN4-mCherry cell line. We thank Jeffrey Kieft for his guidance in carrying out polysome profiling. We thank Neelanjan Mukherjee, Olivia Rissland, Srinivas Ramachandran, and all members of the Jagannathan laboratory for insightful manuscript feedback. We thank the BioFrontiers Institute Next-Gen Sequencing Core Facility, which performed the Illumina sequencing and library construction. This work was supported by the RNA Bioscience Initiative, University of Colorado Anschutz Medical Campus (S.J.), Friends of FSH Research and The Chris Carrino Foundation for FSHD AWD-194864 (S.J.), the FSHD Society FSHS-82018-01 (A.E.C. and M.D.), the California Tobacco-Related Disease Research Grants Program 27KT-0003 (S.N.F.), and the National Institutes of Health DP2GM132932 (S.N.F.).

AUTHOR CONTRIBUTIONS

A.E.C., M.C.D., and S.J. conceived and designed the study. A.E.C, M.C.D., M.A.C., and T.F. performed experiments. A.E.C., L.C., T.M., R.F., A.E.G., S.N.F., and S.J. analyzed data. A.E.C. and S.J. wrote the paper with input from all authors.

DECLARATION OF INTERESTS

The authors declare no competing interests.

Figure 2 – figure supplement 1. Gene-level analysis of early changes in pathways impacted by DUX4. (**A**) Heat maps of transcript-level changes in genes that belong to various biological pathways known to be misregulated by DUX4, as defined by the Gene Ontology Biological Process classification system accessed through GSEA/MSigDB database. The subset of genes that showed significant change in expression at the 4 h time point of DUX4 induction are plotted. (**B**) Elbow method analysis was used to determine appropriate cluster number for Figure 2C-E.

Figure 3 – figure supplement 1. Ribo-seq quality control. Aggregate profile of P-sites coverage (as calculated by Ribo-seQC) depicting single nucleotide resolution of Ribo-seq data along the time course. Each frame is shown with a different color. Error bars represent the standard deviation from three biological replicates.

Video 1. Time course imaging following DUX4 expression.

Live cell fluorescence microscopy recording of MB135-iDUX4/ZSCAN4-mCherry myoblasts

treated with doxycycline to induce DUX4 expression.

Supplementary Table 1.

DESeq2 differential gene expression analysis results for DUX4 time course RNA-seq data at 4,

8, or 14 h post-induction compared to 0 h (control).

Supplementary Table 2.

Cluster analysis of RNA-seq log2 fold change at 4, 8, or 14 h of DUX4 induction compared to 0

h (control); and Gene Ontology analysis of genes within each cluster.

Supplementary Table 3.

ORFquant analysis results for DUX4 time course RNA-seq and Ribo-seq data at 4, 8, or 14 h

post-induction compared to 0 h (control); and Gene Ontology analysis of genes with

downregulated ribosome density at 14 h.

Supplementary Table 4.

DEXSeq analysis results for DUX4 time course Ribo-seq data at 4, 8, or 14 h post-induction

compared to 0 h (control); and Gene Ontology analysis of NMD targets upregulated at 14 h.

REFERENCES

Anders, S., Reyes, A., & Huber, W. (2012). Detecting differential usage of exons from RNA-seq

data. *Genome Res, 22*(10), 2008-2017. doi:10.1101/gr.133744.111

Ashoti, A., Alemany, A., Sage, F., & Geijsen, N. (2021). DUX4 induces a homogeneous

sequence of molecular changes, culminating in the activation of a stem-cell-like transcriptional

network and induction of apoptosis in somatic cells. *bioRxiv*. doi:10.1101/2021.05.04.442407

Banerji, C. R., Knopp, P., Moyle, L. A., Severini, S., Orrell, R. W., Teschendorff, A. E., &

Zammit, P. S. (2015). beta-Catenin is central to DUX4-driven network rewiring in

facioscapulohumeral muscular dystrophy. *J R Soc Interface, 12*(102), 20140797.

doi:10.1098/rsif.2014.0797

Banerji, C. R. S., Panamarova, M., Hebaishi, H., White, R. B., Relaix, F., Severini, S., & Zammit,

P. S. (2017). PAX7 target genes are globally repressed in facioscapulohumeral muscular

dystrophy skeletal muscle. *Nat Commun, 8*(1), 2152. doi:10.1038/s41467-017-01200-4

Barger, C. J., Branick, C., Chee, L., & Karpf, A. R. (2019). Pan-Cancer Analyses Reveal

Genomic Features of FOXM1 Overexpression in Cancer. *Cancers (Basel), 11*(2).

doi:10.3390/cancers11020251

Bosnakovski, D., Chan, S. S. K., Recht, O. O., Hartweck, L. M., Gustafson, C. J., Athman, L. L.,

. . . Kyba, M. (2017). Muscle pathology from stochastic low level DUX4 expression in an FSHD

mouse model. *Nat Commun, 8*(1), 550. doi:10.1038/s41467-017-00730-1

Bosnakovski, D., da Silva, M. T., Sunny, S. T., Ener, E. T., Toso, E. A., Yuan, C., . . . Kyba, M.

(2019). A novel P300 inhibitor reverses DUX4-mediated global histone H3 hyperacetylation,

target gene expression, and cell death. *Sci Adv, 5*(9), eaaw7781. doi:10.1126/sciadv.aaw7781

Bosnakovski, D., Daughters, R. S., Xu, Z., Slack, J. M., & Kyba, M. (2009). Biphasic myopathic

phenotype of mouse DUX, an ORF within conserved FSHD-related repeats. *PLoS One, 4*(9),

e7003. doi:10.1371/journal.pone.0007003

Bosnakovski, D., Gearhart, M. D., Toso, E. A., Ener, E. T., Choi, S. H., & Kyba, M. (2018). Low

level DUX4 expression disrupts myogenesis through deregulation of myogenic gene expression.

Sci Rep, 8(1), 16957. doi:10.1038/s41598-018-35150-8

Bosnakovski, D., Toso, E. A., Hartweck, L. M., Magli, A., Lee, H. A., Thompson, E. R., . . . Kyba,

M. (2017). The DUX4 homeodomains mediate inhibition of myogenesis and are functionally

exchangeable with the Pax7 homeodomain. *J Cell Sci, 130*(21), 3685-3697.

doi:10.1242/jcs.205427

Bosnakovski, D., Xu, Z., Gang, E. J., Galindo, C. L., Liu, M., Simsek, T., . . . Kyba, M. (2008).

An isogenetic myoblast expression screen identifies DUX4-mediated FSHD-associated

molecular pathologies. *EMBO J, 27*(20), 2766-2779. doi:10.1038/emboj.2008.201

Calviello, L., Hirsekorn, A., & Ohler, U. (2020). Quantification of translation uncovers the

functions of the alternative transcriptome. *Nat Struct Mol Biol, 27*(8), 717-725.

doi:10.1038/s41594-020-0450-4

Calviello, L., Sydow, D., Harnett, D., & Ohler, U. (2019). Ribo-seQC: comprehensive analysis of

cytoplasmic and organellar ribosome profiling data. *bioRxiv*. doi:10.1101/601468

Calviello, L., Venkataramanan, S., Rogowski, K. J., Wyler, E., Wilkins, K., Tejura, M., . . . Floor,

S. N. (2021). DDX3 depletion represses translation of mRNAs with complex 5' UTRs. *Nucleic*

Acids Res, 49(9), 5336-5350. doi:10.1093/nar/gkab287

Campbell, A. E., Belleville, A. E., Resnick, R., Shadle, S. C., & Tapscott, S. J. (2018).

Facioscapulohumeral dystrophy: activating an early embryonic transcriptional program in human

skeletal muscle. *Hum Mol Genet, 27*(R2), R153-R162. doi:10.1093/hmg/ddy162

Chew, G. L., Campbell, A. E., De Neef, E., Sutliff, N. A., Shadle, S. C., Tapscott, S. J., &

Bradley, R. K. (2019). DUX4 Suppresses MHC Class I to Promote Cancer Immune Evasion and

Resistance to Checkpoint Blockade. *Dev Cell, 50*(5), 658-671 e657.

doi:10.1016/j.devcel.2019.06.011

Chothani, S., Adami, E., Ouyang, J. F., Viswanathan, S., Hubner, N., Cook, S. A., . . . Rackham,
O. J. L. (2019). deltaTE: Detection of Translationally Regulated Genes by Integrative Analysis of
Ribo-seq and RNA-seq Data. *Curr Protoc Mol Biol, 129*(1), e108. doi:10.1002/cpmb.108

Dandapat, A., Bosnakovski, D., Hartweck, L. M., Arpke, R. W., Baltgalvis, K. A., Vang, D., . . .
Kyba, M. (2014). Dominant lethal pathologies in male mice engineered to contain an X-linked
DUX4 transgene. *Cell Rep, 8*(5), 1484-1496. doi:10.1016/j.celrep.2014.07.056

DeSimone, A. M., Leszyk, J., Wagner, K., & Emerson, C. P., Jr. (2019). Identification of the
hyaluronic acid pathway as a therapeutic target for facioscapulohumeral muscular dystrophy.
Sci Adv, 5(12), eaaw7099. doi:10.1126/sciadv.aaw7099

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., . . . Gingeras, T. R.
(2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics, 29*(1), 15-21.
doi:10.1093/bioinformatics/bts635

Feng, Q., Snider, L., Jagannathan, S., Tawil, R., van der Maarel, S. M., Tapscott, S. J., &
Bradley, R. K. (2015). A feedback loop between nonsense-mediated decay and the retrogene
DUX4 in facioscapulohumeral muscular dystrophy. *Elife, 4*. doi:10.7554/eLife.04996

Geng, L. N., Yao, Z., Snider, L., Fong, A. P., Cech, J. N., Young, J. M., . . . Tapscott, S. J.
(2012). DUX4 activates germline genes, retroelements, and immune mediators: implications for
facioscapulohumeral dystrophy. *Dev Cell, 22*(1), 38-51. doi:10.1016/j.devcel.2011.11.013

Hamel, J., & Tawil, R. (2018). Facioscapulohumeral Muscular Dystrophy: Update on
Pathogenesis and Future Treatments. *Neurotherapeutics, 15*(4), 863-871. doi:10.1007/s13311-
018-00675-3

Homma, S., Beermann, M. L., Boyce, F. M., & Miller, J. B. (2015). Expression of FSHD-related
DUX4-FL alters proteostasis and induces TDP-43 aggregation. *Ann Clin Transl Neurol, 2*(2),
151-166. doi:10.1002/acn3.158

Homma, S., Beermann, M. L., Yu, B., Boyce, F. M., & Miller, J. B. (2016). Nuclear bodies
reorganize during myogenesis in vitro and are differentially disrupted by expression of FSHD-
associated DUX4. *Skelet Muscle, 6*(1), 42. doi:10.1186/s13395-016-0113-7

Huber, W., Carey, V. J., Gentleman, R., Anders, S., Carlson, M., Carvalho, B. S., . . . Morgan,
M. (2015). Orchestrating high-throughput genomic analysis with Bioconductor. *Nat Methods,*
12(2), 115-121. doi:10.1038/nmeth.3252

Ingolia, N. T. (2014). Ribosome profiling: new views of translation, from single codons to
genome scale. *Nat Rev Genet, 15*(3), 205-213. doi:10.1038/nrg3645

Jagannathan, S., Ogata, Y., Gafken, P. R., Tapscott, S. J., & Bradley, R. K. (2019). Quantitative
proteomics reveals key roles for post-transcriptional gene regulation in the molecular pathology
of facioscapulohumeral muscular dystrophy. *Elife, 8*. doi:10.7554/eLife.41740

Jagannathan, S., Shadle, S. C., Resnick, R., Snider, L., Tawil, R. N., van der Maarel, S. M., . . .
Tapscott, S. J. (2016). Model systems of DUX4 expression recapitulate the transcriptional
profile of FSHD cells. *Hum Mol Genet, 25*(20), 4419-4431. doi:10.1093/hmg/ddw271

Jumaa, H., & Nielsen, P. J. (1997). The splicing factor SRp20 modifies splicing of its own mRNA
and ASF/SF2 antagonizes this regulation. *EMBO J, 16*(16), 5077-5085.
doi:10.1093/emboj/16.16.5077

Konigs, V., de Oliveira Freitas Machado, C., Arnold, B., Blumel, N., Solovyeva, A., Lobbert, S., .
. . Muller-McNicoll, M. (2020). SRSF7 maintains its homeostasis through the expression of Split-
ORFs and nuclear body assembly. *Nat Struct Mol Biol, 27*(3), 260-273. doi:10.1038/s41594-
020-0385-9

Kowaljow, V., Marcowycz, A., Ansseau, E., Conde, C. B., Sauvage, S., Matteotti, C., . . . Rosa,
A. L. (2007). The DUX4 gene at the FSHD1A locus encodes a pro-apoptotic protein.
Neuromuscul Disord, 17(8), 611-623. doi:10.1016/j.nmd.2007.04.002

Kurosaki, T., & Maquat, L. E. (2016). Nonsense-mediated mRNA decay in humans at a glance.
J Cell Sci, 129(3), 461-467. doi:10.1242/jcs.181008

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat*

Methods, 9(4), 357-359. doi:10.1038/nmeth.1923

Leclair, N. K., Brugiolo, M., Urbanski, L., Lawson, S. C., Thakar, K., Yurieva, M., . . . Anczukow,

O. (2020). Poison Exon Splicing Regulates a Coordinated Network of SR Protein Expression

during Differentiation and Tumorigenesis. *Mol Cell, 80*(4), 648-665 e649.

doi:10.1016/j.molcel.2020.10.019

Lek, A., Zhang, Y., Woodman, K. G., Huang, S., DeSimone, A. M., Cohen, J., . . . Kunkel, L. M.

(2020). Applying genome-wide CRISPR-Cas9 screens for therapeutic discovery in

facioscapulohumeral muscular dystrophy. *Sci Transl Med, 12*(536).

doi:10.1126/scitranslmed.aay0271

Lemmers, R. J., van der Vliet, P. J., Klooster, R., Sacconi, S., Camano, P., Dauwerse, J. G., . . .

van der Maarel, S. M. (2010). A unifying genetic model for facioscapulohumeral muscular

dystrophy. *Science, 329*(5999), 1650-1653. doi:10.1126/science.1189044

Liao, Y., Smyth, G. K., & Shi, W. (2014). featureCounts: an efficient general purpose program

for assigning sequence reads to genomic features. *Bioinformatics, 30*(7), 923-930.

doi:10.1093/bioinformatics/btt656

Lim, K. R. Q., Nguyen, Q., & Yokota, T. (2020). DUX4 Signalling in the Pathogenesis of

Facioscapulohumeral Muscular Dystrophy. *Int J Mol Sci, 21*(3). doi:10.3390/ijms21030729

Lin, J. H., Li, H., Yasumura, D., Cohen, H. R., Zhang, C., Panning, B., . . . Walter, P. (2007).

IRE1 signaling affects cell fate during the unfolded protein response. *Science, 318*(5852), 944-

949. doi:10.1126/science.1146361

Livak, K. J., & Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-

time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods, 25*(4), 402-408.

doi:10.1006/meth.2001.1262

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol, 15*(12), 550. doi:10.1186/s13059-014-
0550-8

Merrick, W. C., & Hensold, J. O. (2001). Analysis of eukaryotic translation in purified and
semipurified systems. *Curr Protoc Cell Biol, Chapter 11*, Unit 11 19.
doi:10.1002/0471143030.cb1109s08

Miura, P., Andrews, M., Holcik, M., & Jasmin, B. J. (2008). IRES-mediated translation of
utrophin A is enhanced by glucocorticoid treatment in skeletal muscle cells. *PLoS One, 3*(6),
e2309. doi:10.1371/journal.pone.0002309

More, D. A., & Kumar, A. (2020). SRSF3: Newly discovered functions and roles in human health
and diseases. *Eur J Cell Biol, 99*(6), 151099. doi:10.1016/j.ejcb.2020.151099

Muller-McNicoll, M., Rossbach, O., Hui, J., & Medenbach, J. (2019). Auto-regulatory feedback
by RNA-binding proteins. *J Mol Cell Biol, 11*(10), 930-939. doi:10.1093/jmcb/mjz043

Proud, C. G. (2005). eIF2 and the control of cell physiology. *Semin Cell Dev Biol, 16*(1), 3-12.
doi:10.1016/j.semcdb.2004.11.004

Resnick, R., Wong, C. J., Hamm, D. C., Bennett, S. R., Skene, P. J., Hake, S. B., . . . Tapscott,
S. J. (2019). DUX4-Induced Histone Variants H3.X and H3.Y Mark DUX4 Target Genes for
Expression. *Cell Rep, 29*(7), 1812-1820 e1815. doi:10.1016/j.celrep.2019.10.025

Rickard, A. M., Petek, L. M., & Miller, D. G. (2015). Endogenous DUX4 expression in FSHD
myotubes is sufficient to cause cell death and disrupts RNA splicing and cell migration
pathways. *Hum Mol Genet, 24*(20), 5901-5914. doi:10.1093/hmg/ddv315

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of
image analysis. *Nat Methods, 9*(7), 671-675. doi:10.1038/nmeth.2089

Shadle, S. C., Bennett, S. R., Wong, C. J., Karreman, N. A., Campbell, A. E., van der Maarel, S.
M., . . . Tapscott, S. J. (2019). DUX4-induced bidirectional HSATII satellite repeat transcripts

form intranuclear double-stranded RNA foci in human cell models of FSHD. *Hum Mol Genet,*
28(23), 3997-4011. doi:10.1093/hmg/ddz242

Shadle, S. C., Zhong, J. W., Campbell, A. E., Conerly, M. L., Jagannathan, S., Wong, C. J., . . .
Tapscott, S. J. (2017). DUX4-induced dsRNA and MYC mRNA stabilization activate apoptotic
pathways in human cell models of facioscapulohumeral dystrophy. *PLoS Genet, 13*(3),
e1006658. doi:10.1371/journal.pgen.1006658

Sharma, V., Harafuji, N., Belayew, A., & Chen, Y. W. (2013). DUX4 differentially regulates
transcriptomes of human rhabdomyosarcoma and mouse C2C12 cells. *PLoS One, 8*(5),
e64691. doi:10.1371/journal.pone.0064691

Tawil, R., van der Maarel, S. M., & Tapscott, S. J. (2014). Facioscapulohumeral dystrophy: the
path to consensus on pathophysiology. *Skelet Muscle, 4*, 12. doi:10.1186/2044-5040-4-12

Wallace, L. M., Garwick, S. E., Mei, W., Belayew, A., Coppee, F., Ladner, K. J., . . . Harper, S.
Q. (2011). DUX4, a candidate gene for facioscapulohumeral muscular dystrophy, causes p53-
dependent myopathy in vivo. *Ann Neurol, 69*(3), 540-552. doi:10.1002/ana.22275

Whiddon, J. L., Langford, A. T., Wong, C. J., Zhong, J. W., & Tapscott, S. J. (2017).
Conservation and innovation in the DUX4-family gene network. *Nat Genet, 49*(6), 935-940.
doi:10.1038/ng.3846

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. New York, NY: Springer-
Verlag.

Yao, Z., Snider, L., Balog, J., Lemmers, R. J., Van Der Maarel, S. M., Tawil, R., & Tapscott, S.
J. (2014). DUX4-induced gene expression is the major molecular signature in FSHD skeletal
muscle. *Hum Mol Genet, 23*(20), 5342-5352. doi:10.1093/hmg/ddu251

Zeisel, A., Yitzhaky, A., Bossel Ben-Moshe, N., & Domany, E. (2013). An accessible database
for mouse and human whole transcriptome qPCR primers. *Bioinformatics, 29*(10), 1355-1356.
doi:10.1093/bioinformatics/btt145

Zhou, Z., Gong, Q., Lin, Z., Wang, Y., Li, M., Wang, L., . . . Li, P. (2020). Emerging Roles of

SRSF3 as a Therapeutic Target for Cancer. *Front Oncol, 10*, 577636.

doi:10.3389/fonc.2020.577636

Zhu, A., Ibrahim, J. G., & Love, M. I. (2019). Heavy-tailed prior distributions for sequence count

data: removing the noise and preserving large differences. *Bioinformatics, 35*(12), 2084-2092.

doi:10.1093/bioinformatics/bty895

APPENDIX

KEY RESOURCES TABLE

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
cell line (*Homo sapiens*)	293T	ATCC	Cat# CRL-3216, RRID:CVCL_0063	
cell line (*H. sapiens*)	MB135	Stephen Tapscott		Female control myoblast line
cell line (*H. sapiens*)	MB135-iDUX4	(Jagannathan et al., 2016)		Harbors doxycycline-inducible DUX4 transgene
cell line (*H. sapiens*)	MB135-iDUX4/ZSCAN4-mCherry	Stephen Tapscott		Harbors doxycycline-inducible DUX4 transgene and DUX4-responsive fluorescent reporter
cell line (*H. sapiens*)	MB135-iFLAG-SRSF3-FL	This paper		Harbors doxycycline-inducible, FLAG-tagged, codon-optimized, full-length SRSF3
cell line (*H. sapiens*)	MB135-iFLAG-SRSF3-TR	This paper		Harbors doxycycline-inducible, FLAG-tagged, codon-optimized, truncated SRSF3
cell line (*H. sapiens*)	MB200	Stephen Tapscott		Male FSHD myoblast line
antibody	anti-DUX4 (Rabbit monoclonal)	Abcam	Cat#ab124699, RRID:AB_10973363	IF(1:200), WB(1:1000)
antibody	anti-Histone H3 (Rabbit polyclonal)	Abcam	Cat# ab1791, RRID:AB_302613	WB(1:5000)

antibody	anti-SRSF3 (Mouse monoclonal)	Thermo Fisher Scientific	Cat# 33-4200, RRID:AB_2533119	WB(1:250)
antibody	anti-SRSF3-TR (Rabbit polyclonal)	This paper		IF(1:200), WB(1:500)
antibody	anti-RENT1/hUPF1 (Rabbit monoclonal)	Abcam	Cat# ab109363, RRID:AB_10861979	WB(1:1000)
recombinant DNA reagent	pCW57.1-FLAG-SRSF3_Full.Length_Codon.Optimized-Blast (plasmid)	This paper		All-in-one lentiviral vector for doxycycline-inducible expression of FLAG-tagged, codon-optimized, full-length SRSF3
recombinant DNA reagent	pCW57.1-FLAG-SRSF3_Truncated_Codon.Optimized-Blast (plasmid)	This paper		All-in-one lentiviral vector for doxycycline-inducible expression of FLAG-tagged, codon-optimized, truncated SRSF3
recombinant DNA reagent	pCW57-MCS1-P2A-MCS2 (Blast) (plasmid)	Addgene	Plasmid #80921, RRID:Addgene_80921	
recombinant DNA reagent	pMD2.G (plasmid)	Addgene	Plasmid #12259; RRID:Addgene_12259	
recombinant DNA reagent	psPAX2 (plasmid)	Addgene	Plasmid #12260; RRID:Addgene_12260	
recombinant DNA reagent	pTwist-FLAG-SRSF3_Full.Length_Codon.Optimized (plasmid)	This paper		For expression of FLAG-tagged, codon-optimized full-length SRSF3 in mammalian cells; synthesized by Twist Bioscience
recombinant DNA reagent	pTwist-FLAG-SRSF3_Truncated_Codon.Optimized (plasmid)	This paper		For expression of FLAG-tagged, codon-optimized truncated SRSF3 in mammalian cells; synthesized by Twist Bioscience
sequence-based reagent	DUX4 transgene F	(Shadle et al., 2019)	qPCR primers	TAGGGGAAGAGGTAGACGGC

sequence-based reagent	DUX4 transgene R	(Shadle et al., 2019)	qPCR primers	CGGTTCCGGGATTC CGATAG
sequence-based reagent	HAGHL F	This paper	qPCR primers	AGTTTGCCCAGAAAG TGGAG
sequence-based reagent	HAGHL R	This paper	qPCR primers	ATCCTCATCCCTCTT CTTAGCC
sequence-based reagent	HSATII F	(Shadle et al., 2019)	qPCR primers	TGAATGGAATCGTCA TCGAA
sequence-based reagent	HSATII R	(Shadle et al., 2019)	qPCR primers	CCATTCGATAATTCC GCTTG
sequence-based reagent	HSPA5 F	(Lin et al., 2007)	qPCR primers	CGGGCAAAGATGTC AGGAAAG
sequence-based reagent	HSPA5 R	(Lin et al., 2007)	qPCR primers	TTCTGGACGGGCTT CATAGTAGAC
sequence-based reagent	MYOD1 F	Stephen Tapscott	qPCR primers	AGCACTACAGCGGC GACT
sequence-based reagent	MYOD1 R	Stephen Tapscott	qPCR primers	GCGACTCAGAAGGC ACGTC
sequence-based reagent	SRSF3 F	(Feng et al., 2015)	qPCR primers	TGGAACTGTCGAATG GTGAA
sequence-based reagent	SRSF3-Excl R	(Feng et al., 2015)	qPCR primers	CTTGGAGATCTGCG ACGAG
sequence-based reagent	SRSF3-Incl R	(Feng et al., 2015)	qPCR primers	GGGTGGTGAGAAGA GACATGA
sequence-ce-	TCF3 F	(Zeisel, Yitzhaky,	qPCR primers	CCCCAGGAGAATGA ACCAG

based reagent		Bossel Ben-Moshe, & Domany, 2013)		
sequence-based reagent	TCF3 R	(Zeisel et al., 2013)	qPCR primers	CCTTCCCGTTGGTGACAG
sequence-based reagent	RPL27 F	(Jagannathan et al., 2016)	qPCR primers	GCAAGAAGAAGATCGCCAAG
sequence-based reagent	RPL27 R	(Jagannathan et al., 2016)	qPCR primers	TCCAAGGGGATATCCACAGA
sequence-based reagent	ZSCAN4 F	(Geng et al., 2012)	qPCR primers	TGGAAATCAAGTGGCAAAAA
sequence-based reagent	ZSCAN4 R	(Geng et al., 2012)	qPCR primers	CTGCATGTGGACGTGGAC
peptide, recombinant protein	Insulin-Transferrin-Selenium	Thermo Fisher Scientific	41400045	
peptide, recombinant protein	Protein G Sepharose	Thermo Fisher Scientific	101241	
peptide, recombinant protein	Recombinant human basic fibroblast growth factor	Promega	G5071	
chemical compound, drug	Blasticidin S HCl	Thermo Fisher Scientific	R21001	
chemical compound, drug	Cycloheximide	Sigma-Aldrich	239765	

chemical compound, drug	Doxycycline hyclate	Sigma-Aldrich	D9891	
chemical compound, drug	Lipofectamine 2000 Transfection Reagent	Thermo Fisher Scientific	11668-030	
chemical compound, drug	Puromycin dihydrochloride	VWR	97064-280	
chemical compound, drug	TRIzol Reagent	Thermo Fisher Scientific	15596018	
commerical assay or kit	BCA Protein Assay Kit	Pierce	23225	
commerical assay or kit	Direct-zol RNA Miniprep Kit	Zymo Research	R2051	
commerical assay or kit	NEXTflex Rapid Directional qRNA-Seq Kit	Bioo Scientific	NOVA-5130-02D	
commerical assay or kit	NEXTflex Small RNA-Seq Kit v3	Bioo Scientific	NOVA-5132-06	
commerical assay or kit	Ribo-Zero rRNA Removal Kit	Illumina	MRZH11124, discontinued	
commerical assay or kit	SuperScript III First-Strand Synthesis System	Thermo Fisher Scientific	18080051	
software, algorithm	Code used for RNA-seq and Ribo-seq figure generation	This paper		The code used for the RNA-seq and Ribo-seq figure generation can be accessed via github at https://github.com/sjag

				anna/2021-campbell_dyle_calviello_et_al
software, algorithm	GraphPad Prism	GraphPad Prism (https://graphpad.com)	RRID:SCR_002798	Version 9
software, algorithm	ImageJ	ImageJ (https://imagej.nih.gov)	RRID:SCR_003070	