Title: Pleiotropy is associated with fitness consequences of new mutations in mutation accumulation lines

Author list: Sydney Kreutzmann[1], Elizabeth Pompa[2], Nhan D. Nguyen[2], Liya Tilahun[3], Matthew T. Rutter[4], Mao-Lun Weng[5], Charles B. Fenster[6], *Carrie F. Olson-Manning[1]

Department of Biology, Augustana University, Sioux Falls, SD, 57105, USA

Department of Chemistry and Biochemistry, Augustana University, Sioux Falls, SD, 57105, USA

Make School, Dominican University of California, San Francisco, CA, 94102, USA

Department of Biology, College of Charleston, Charleston, SC, 29424, USA

Department of Biology, Westfield State University, Westfield, MA 01086, USA

Director, Oak Lake Field Station, South Dakota State University, Brookings, SD 57007,USA

* Author or Correspondence: Carrie F. Olson-Manning Department of Biology, Augustana University, Sioux Falls, SD, 57105, USA, colsonmanning@augie.edu

ORC IDs

M.T.R. 0000-0002-0389-7293

M.L. 0000-0001-7299-8736

C.B.F. 0000-0002-1655-4409

C.F.O.-M. 0000-0001-8054-3889

Keywords: mutation accumulation, spontaneous mutations, transcriptional regulatory activity, metabolomics

Significance Statement:

As the source of all genetic variation, new mutations are crucial for understanding how organisms adapt to their environment. However, the ways in which new mutations affect the range of metabolic reactions that occur in the cell is unknown. With a combination of gene functional analyses and measurement of the small molecules that drive cellular function and physiology we found that mutations associated with high

fitness are enriched in proteins that regulate the specific timing and location of gene expression and are less disruptive of cellular physiology.

ABSTRACT

Understanding the mechanisms by which mutations affect fitness and the distribution of mutational effects are central goals in evolutionary biology. Mutation accumulation (MA) lines have long been an important tool for understanding the effect of new mutations on fitness, phenotypic variation, and mutational parameters. However, there is a clear gap in predicting the effect of new mutations to their effects on fitness. Here we complete gene ontology analysis and metabolomics experiments on *Arabidopsis thaliana* MA lines to determine how spontaneous mutations affect global metabolic output in lines that have measured fitness consequences. For these analyses, we compared three lines with relative fitness consistently higher than the unmutated progenitor and three lines with lower relative fitness. In a gene ontology analysis, we find that the high fitness lines were significantly enriched in mutations in or near genes with transcription regulator activity. We also find that although they do not have an average difference in the number of mutations, low fitness lines have significantly more metabolic subpathways disrupted than high fitness lines. Taken together, these results suggest that the effect of a new mutation on fitness depends less on the specific metabolic pathways disrupted and more on the pleiotropic effects of those mutations and that organisms can explore a considerable amount of physiological space with only a few mutations.

Introduction

Adaptation requires mutations that both affect a certain trait under selection and, importantly, do not have deleterious effects on other traits. Proteins that regulate the timing and location of gene expression may satisfy both of these conditions. Transcription regulatory proteins interact with the regulatory DNA to direct the spatial and temporal gene expression in eukaryotes. These proteins drive gene expression in a highly specific manner by interacting with the DNA regulatory elements of genes in a combinatorial fashion. A growing number of studies have found that mutations in transcription factors make important contributions to adaptive evolution (Des Marais and Rausher 2010; Streisfeld and Rausher 2009, 2011; Durbin 2003; Whittall et al. 2006; Hopkins and Rausher 2011; Quattrocchio et al. 1999; Hoballah et al. 2007). While some transcription regulatory proteins are widely expressed, a majority are expressed in only a few tissues or under specific environmental conditions (Streisfeld, Liu, and Rausher 2011). Due to this tissue-specific functioning, mutations that affect the regulation of genes are less likely to have widespread pleiotropic consequences and are thus less likely to be deleterious (consistent with theory (Fisher 1930)). Given the demonstrated importance of regulatory evolution for adaptation, we would like to understand the prevalence of transcriptional regulatory mutations in the mutational spectrum (i.e. the range of spontaneous mutations that occur) and to predict the effect of mutations on organismal physiology and fitness.

An important tool for examining the types and effects of new spontaneous mutations are mutation accumulation (MA) lines (Halligan and Keightley 2009). MA lines are created when sublines of an original progenitor are allowed to accumulate spontaneous mutations in the absence of selection through many generations. These lines have shed light on the distribution of phenotypic effects (Chang and Shaw 2003; Shaw, Geyer, and Shaw 2002; Weng et al. 2021), the mutational spectrum (Weng et al. 2019; Schrider et al. 2013; Denver et al. 2006; Grey Monroe et al. 2020; Nguyen et al. 2020), and the average fitness effect of new mutations (Rutter, Roles, and Fenster 2018; Rutter et al. 2012; Roles et al. 2016; Rutter, Shaw, and Fenster 2010) (reviewed in (Katju and Bergthorsson 2019; Eyre-Walker and Keightley 2007)).

The *Arabidopsis thaliana* MA lines (Chang and Shaw 2003; Shaw, Geyer, and Shaw 2002) offer an opportunity to connect genotype to fitness through physiology. The *A. thaliana* MA lines have been fully sequenced (Weng et al. 2019) and the fitness measured in field trials (Rutter, Roles, and Fenster 2018; Rutter et al. 2012). However, there has been little use of this resource to examine the mutational effect of fitness through physiology. In accordance with theory (Fisher 1930) and supported by empirical studies (e.g. (Cooper, Nickerson, and Eichler 2007)), we predict that lines that have higher fitness than the progenitor should also have proportionally more mutations that display low levels of pleiotropy, and these mutations may be enriched in genes responsible for regulation. With these high and low fitness lines and their associated mutations, we ask 1) are regulatory proteins enriched in MA lines that have high relative fitness compared to the unmutated progenitor? and 2) Are there more overall changes in metabolic output in the low fitness lines compared to the high fitness lines?

In an analysis of the gene ontology of the mutations in or near the genes in the high and low fitness lines, we find that mutations in MA lines with high relative fitness are enriched in proteins with transcription regulator activity. We also find an intriguing trend where high fitness lines have fewer changes to their metabolic pathways than the low fitness lines, suggesting that the mutations associated with the high fitness lines are less pleiotropic than mutations in the low fitness lines. Taken together, these results suggest that mutations that affect transcription regulator activity are common in the mutational spectrum and that lines that harbor these mutations have less disrupted metabolic expression.

Results and Discussion

The ways in which mutations affect organismal fitness is of great interest in evolutionary biology and molecular evolution. Although other evolutionary forces determine the fate of the mutations after they arise, bias in the mutational spectrum can affect the composition of the genome (Nguyen et al. 2020)

how the composition of the genome affects mutation rate (Grey Monroe et al. 2020), and polymorphism in populations (Weng et al. 2019). In our analyses, we seek to connect the composition of spontaneous mutations in the *A. thaliana* MA lines with high and low relative fitness (measured under field conditions) and the effect of those mutations on organismal metabolism.

High fitness lines are enriched in transcription regulator activity

In a Gene Ontology (GO) term analysis for molecular function (Mi et al. 2021) we found that the genes mutated in the high fitness lines submitted for metabolomics in this study have a 7.7-fold enrichment (Fisher's Exact Test with Bonferroni correction, $p<0.01$) in transcription regulatory activity (8 of the 16 mapped gene IDs and represent between 25-50% of the mutations in each line). We repeated this GO term analysis for all 18 high fitness MA lines (Supplementary Table 4) and found that together they also have a greater than 3.1-fold enrichment for transcription regulator activity ($p<0.001$, 21 of 114 mapped gene IDs have transcription regulator activity). The same analyses on all 11 low fitness lines revealed no enrichment in any activity and only one out of 32 genes with mutations in the low fitness lines has transcriptional regulatory activity.

These results suggest that mutations in or near genes encoding transcriptional regulatory proteins make up a substantial portion of the mutational spectrum in MA lines with high relative fitness and are less prevalent in lines with low fitness. Proteins with transcriptional regulatory activity assemble into varied complexes consisting of other proteins and cis-regulatory sequences in a combinatorial fashion to tightly control the timing and location of gene expression (Brkljacic and Grotewold 2017). In this way, a small handful of transcription factors can control the expression of a wide variety of genes. The combinatorial nature of these regulatory networks have led many to hypothesize that mutations in transcription factor proteins could play an outsized role in adaptive evolution because they are less pleiotropic (Jeong et al. 2008; Wray et al. 2003). The evolutionary genetics literature has many examples implicating mutations in proteins with transcriptional regulatory activity in adaptations, including flower color evolution (Streisfeld, Liu, and Rausher 2011; Wessinger and Rausher 2013; Lin and Rausher 2021; Quattrocchio et al. 1999; Whittall et al. 2006), adaptation to high temperature (Koini et al. 2009), cold

acclimation (Van Buskirk and Thomashow 2006), drought (Haake et al. 2002; Leng and Zhao 2020; Jan et al. 2019), Zinc deficiency (Inaba et al. 2015), and the interface between stress and growth response (Danisman 2016). Given the purported role of proteins with transcriptional regulatory activity (Wagner and Lynch 2008), and their prevalence in the mutational spectrum in lines with high relative fitness, it is suggested that the types of low pleiotropy mutations are readily available for adaptation.

Metabolic disruption in high and low fitness lines

We find a consistent trend of more metabolic subpathways disruption in the low fitness lines than in the high fitness lines. The low fitness lines have an average of 25 disrupted subpathways compared to 15.3 disrupted subpathways in the high fitness lines when we consider metabolites that approach significance (two-sided unpaired t-test, p-value = 0.02) (Table 1, Supplementary Table 5). This pattern of a higher average number of disrupted metabolites in low fitness lines is consistent at the super pathway, subpathway, and individual metabolite levels, regardless of whether we include only statistically significant metabolites or metabolites that approached significance (Table 1). Overall, these results suggest that mutations in the low fitness lines affect more parts of metabolism than mutations in high fitness lines.

Taken together, the lower levels of pleiotropy and enrichment in mutations in or near genes with transcription factor activity in the high fitness lines is consistent with studies that find transcription

Table 1: The number of enriched metabolic subpathways and individual metabolites calculated for metabolites that met the 0.05 significance threshold and those that met the 0.1 threshold. Significant difference between the number of enriched subpathways that reach the 0.1 threshold denoted with * (p<0.05 unpaired t-test).

Line number	Relative Fitness	Enriched Subpathways p<0.05	Enriched Subpathways p<0.1 *	Total metabolites p<0.05	Total metabolites p<0.1
44	Low	10	21	13	37
73		20	29	32	60
75		12	25	13	39
53	High	14	17	16	25
61		10	14	18	35
119		7	15	9	25

factors are over-represented in adaptive evolution due to their non-pleiotropic nature (Smith 2016;

Wagner and Lynch 2008; Cheatle Jarvela and Hinman 2015). Many transcription factors are expressed in a tissue-specific manner and the level of constraint in different transcription factors is reliably predicted by level of integration into gene regulatory networks (Streisfeld, Liu, and Rausher 2011). Our results suggest that mutations in genes coding or transcription factors are not only common in the mutational spectrum but they may also have predictably less pleiotropic effects.

Pleiotropic effects of mutations on metabolic pathways

While the high and low fitness lines have no average difference in the number of unique mutations (average of 9.0 for high fitness and 10.7 for low fitness lines, Supplementary Table 1), there is considerable variance in the number of mutations among the lines. There is no relationship between the number of mutations and the number significantly disrupted metabolic subpathways (F-statistic 0.017, p = 0.9026). On average, each mutation affects multiple metabolic pathways. Mutations in four of the six lines affect 1.5-3.5 metabolic subpathways per mutation (Supplementary Table 1). High fitness line 119 and low fitness line 44 both affect fewer than one metabolic subpathways per mutation. With this small sample size and the confounding effects of individual mutations in each line, we cannot know with certainty how many pathways each mutation affects. However, based on these data, it appears that *A. thaliana* can explore a considerable amount of physiological space with only a few mutations.

Spontaneous mutations cause excess down-regulation of metabolic pathways

We found that overall, mutations caused significantly more down-regulation to metabolic pathways than up-regulation in both the high and low fitness lines (Figure 1, blue indicates down-regulation, red is up-regulation, Chi-square p<0.007). Down-regulated metabolites are 4-times more common than up-regulated metabolites and there was no significant difference in the number of up- and down-regulation in a comparison of the high and low fitness lines. It is possible that mutations are more likely to lead to down-regulation because mutations tend to disrupt steps in metabolic pathways.

Figure 1: Log fold change of all significant and near significant metabolites. The columns from left to right are the low fitness lines (44, 73, 75) and high fitness lines (53, 61, 119). The rows are ordered based on superpathway (full data in Supplemental Table 3).

While it might be reasonable to assume that general degradation of metabolic function is deleterious, there are many instances of loss of function alleles contributing to adaptation (Stower 2013; Xu et al. 2019; Grey Monroe et al. 2020; Monroe et al. 2018). In *A. thaliana*, loss-of-function alleles are common in coding genes, and an estimated 1% of loss-of-function alleles are under positive selection (Xu et al. 2019). Therefore, without further measure of allele frequency changes in the MA lines following selection, it is not possible to speculate on the evolutionary importance in an overall reduction in metabolic activity.

Spontaneous mutations affect metabolism in varied ways

The effect of new spontaneous mutations on metabolism is varied (Fig. 1 and 2) and suggests that the mutations accumulated in these lines are idiosyncratic with respect to effect on metabolic pathways. In a comparison of the six lines with Euclidean distance (Supplementary Figure 1) the high and low fitness

lines are intermixed suggesting that there are not consistent changes to the types of metabolic disruption that led to high or low fitness. This suggests that the spontaneous mutations were indeed random with respect to their effect on metabolism.

Figure 2: Up- (reds) and down-regulation (blues) metabolites in their super pathways context with

subpathways in individual boxes. Lighter colors are metabolites that approach significance. We excluded super pathways from this figure that did not show any significant or near-significant differences (hormone metabolism and partially characterized molecules metabolism)

The potential and limitations of metabolomics studies in understanding the effects of spontaneous mutations

Understanding and predicting the effect of mutations is a primary goal across many fields of biology ((Eyre-Walker and Keightley 2007)). In this study, we used metabolomics to try to close the gap between a mutation and its effect on fitness. Even with hundreds of metabolites measured in this study in well-known pathways and the well-annotated *A. thaliana* genome (Lamesch et al. 2012) we failed to infer any direct connections between mutations in or near enzymes with an effect on metabolism. Among the reasons for this failed connection could be that most of the mutations in the MA lines in this study are intergenic and have direct or indirect effects on the expression genes far away, that the small number of mutations failed to disrupt any known pathways, or that our annotations of gene function are woefully incomplete.

This study only measures the metabolic effects in leaf tissue at one time point on only six of the 107 *A. thaliana* MA lines, so the conclusions relating to the pleiotropic nature of the mutations should be viewed in that light. We also chose lines that showed consistent relative fitness in field trials (Rutter, Roles, and Fenster 2018); however, most of the lines had strong genotype-by-environment effects. Additionally, studies on the effect of new, spontaneous mutations on transcriptional expression have shed light on some of the biases of the mutational spectrum (Zalts and Yanai 2017; Huang et al. 2016) but lack fitness measures to connect genotypic changes to fitness effects. Future work that includes transcriptional studies combining metabolic effects with sampling of additional tissues and an analysis of more lines will further elucidate the effect of spontaneous mutations on gene expression, fitness, and physiology.

Materials and Methods

Choice in plants to grow and analyze

Of the 107 total *A. thaliana* mutation accumulation (MA) lines (Chang and Shaw 2003; Shaw, Geyer, and Shaw 2002), we chose lines that showed relative fitness that was higher and lower in most

environments compared to the unmutated Col-0 progenitor (Rutter et al. 2018) for metabolic analysis (Supplementary Table 1 and 2). The focal MA lines in this study (line numbers 44, 53, 61, 73, 75, and 119) are numbered based on the original propagation line numbers (Chang and Shaw 2003; Shaw, Geyer, and Shaw 2002). Briefly, in a previous study, Rutter et al. (2018), all MA lines were grown in four temporal environments at a single field site and plant fitness of the MA lines was compared to the unmutated progenitor. While most lines displayed genotype-by-environment dependent fitness, for this study we chose lines that consistently ranked among the highest or lowest in relative fitness compared to the progenitor (Supplementary Table 2). As defined in this manuscript, the "high fitness lines" (numbers 53, 61, and 119) had higher relative fitness than the progenitor in at least three of four environments and the "low fitness lines" (lines 44, 73, and 75) had lower relative fitness than the progenitor in at least three of four environments (Rutter et al. 2018). These lines have accumulated an average of 17 mutations over 30 generations of mutation accumulation (Weng et al. 2019).

GO Term Enrichment comparison in high and low fitness lines

To identify gene ontology terms that might be over- or under-represented in the high or low fitness lines, we entered the *A. thaliana* gene model names for mutations (or the gene model names surrounding intergenic mutations) into the Protein Analysis Through Evolutionary Relationships (PANTHER) resource (Mi et al. 2021) (Supplementary Table 3). We compared the set of mutations in the high and low fitness for biological processes, molecular function complete, and cellular component complete with the Fisher's exact test with Bonferroni correction for multiple testing. The analyses were performed separately on the set of three high and three low fitness lines of focus in this study. We also analyzed the additional 15 lines that had average relative fitness that was higher and 8 with relative fitness lower than the progenitor (Rutter, Roles, and Fenster 2018).

Plant growth and tissue harvest

Individual seeds from the 25th generation of mutation accumulation were sown on moistened soil (Promix BX with Osmocote fertilizer added per manufacturer's instructions), randomized in a 24-cell pallet and allowed to imbibe. The seeds were placed in the dark for 3 days at 4 C to stratify and overcome

dormancy. Plants were grown for 3 weeks under long day conditions (16h light) at 18 C. The plant rosettes at time of harvest were vigorous and did not display any obvious signs of stress. To control for the developmental stage, all above-ground tissue was harvested before the plant began bolting and was immediately frozen on liquid nitrogen. Each of the high and low fitness lines had four biological replicates and the progenitor had six. The tissue was stored at -80 C until further processing. Frozen leaves were pulverized on liquid nitrogen and added to a frozen collection tube and stored at -80 C until shipment. Samples were shipped on dry ice to Metabolon Inc. (North Carolina). Frozen samples were lyophilized and each sample was standardized with dry weight.

Metabolon sample extraction and metabolite identification

Briefly, Metabolon Inc. extracted the samples with the MicroLab STAR System (Hamilton Company) with the internal standards. Each extract was divided into four fractions: two were analyzed using acidic positive ion conditions (one optimized for hydrophilic compounds, one for hydrophobic) and two used basic negative ion conditions (one optimized for hydrophilic compounds, one for hydrophobic). Metabolomic analyses were performed using Ultra High Performance Liquid Chromatography-Tandem Mass Spectroscopy (UPLC MS/MS). These methods utilized a Waters Aquity UPLC and Q-Exactive high resolution mass spectrometer (Thermo Scientific) with a heated electrospray ionization (HESI-II) source. The MS analysis used dynamic exclusion, alternating between MS and data-dependent MS^n, and varied between 70-1000 m/z. The MS/MS scores were derived from a comparison of the ions in the experimental samples and the ions in a known library spectrum (within the Metabolon LIMS system. The MS/MS scores (forward and reverse), retention time/index, and molecular mass (m/z) were all processed using Metabolon's software and compared to purified standards within the internal library to identify each metabolite. A number of data curation and data normalization procedures were utilized to ensure high quality data and to decrease minor day-to-day variations from instrumentation tuning. These steps provided the high quality data needed to perform accurate statistical analyses and data interpretation.

Identification of disrupted metabolites and pathways

A total of 386 known compounds were identified in this dataset. Missing data points were given the minimum observed value for each compound and all data were log transformed. Each metabolite was categorized into one of nine super pathways (e.g. Amino Acid, Carbohydrate, Peptide, Secondary Metabolite) and into one of 58 subpathways (e.g. Serine family amino acid, TCA cycle, Dipeptide, Benzenoids) (Supplementary Table 4) (Kanehisa and Goto 2000). Biological replicate measures of each metabolite were compared to the value in the progenitor with Welch's two-sample t-test. A metabolite was judged to be disrupted in a certain line if its value was significantly different from the value in the progenitor based on the t-test based on an alpha value of 0.05.

We took two approaches to identify enrichment in metabolic subpathway disruption. With a pathway enrichment metric (Xia and Wishart 2011), the number of significant ($p \leq 0.05$) compounds were calculated for individual metabolic subpathways. The Enrichment score (Xia and Wishart 2011) compares the ratio of significant metabolites in a pathway (k) to total detected metabolites in that pathway (m), and standardized it by considering the total number of significant metabolites in the entire dataset (n) and the total metabolites detected in the dataset (N) (Eq 1) (Table 1).

Eq 1   Enrichment score = $(k/m) / ((n-k)/(N-m))$

The traditional enrichment score is a conservative method of identifying pathways with more differences than expected as it ignores individual pathways with many metabolites that approach significance in the Welch's two-sample t-test ($p < 0.1$). As a less conservative test, we also calculated an expanded enrichment score that includes metabolites that approach significance. This test allowed us to identify metabolic subpathways that have sets of metabolites that are disrupted and share common biological function. The results we get from the typical enrichment analysis and this extended analysis are consistent.

Comparison of up and down regulation of metabolic pathways

We compared the number of metabolites that were significantly up- or down-regulated as compared to the Col-0 progenitor with a Chi-square statistic. The null expectation is that spontaneous mutations will cause an equal number of up- and down-regulation in metabolites.

Data Availability Statement: All data used in this study is represented in the supplementary tables. The datasets will also be uploaded to the Dryad repository upon acceptance.

References

Brkljacic, Jelena, and Erich Grotewold. 2017. "Combinatorial Control of Plant Gene Expression." *Biochimica et Biophysica Acta, Gene Regulatory Mechanisms* 1860 (1): 31–40.

Chang, Shu-Mei, and Ruth G. Shaw. 2003. "The Contribution of Spontaneous Mutation to Variation in Environmental Response in Arabidopsis Thaliana: Responses to Nutrients." *Evolution; International Journal of Organic Evolution* 57 (5): 984–94.

Cheatle Jarvela, Alys M., and Veronica F. Hinman. 2015. "Evolution of Transcription Factor Function as a Mechanism for Changing Metazoan Developmental Gene Regulatory Networks." *EvoDevo* 6 (1): 3.

Cooper, Gregory M., Deborah A. Nickerson, and Evan E. Eichler. 2007. "Mutational and Selective Effects on Copy-Number Variants in the Human Genome." *Nature Genetics* 39 (7 Suppl): S22–29.

Danisman, Selahattin. 2016. "TCP Transcription Factors at the Interface between Environmental Challenges and the Plant's Growth Responses." *Frontiers in Plant Science*. https://doi.org/10.3389/fpls.2016.01930.

Denver, Dee R., Seth Feinberg, Catherine Steding, Matthew D. Durbin, and Michael Lynch. 2006. "The Relative Roles of Three DNA Repair Pathways in Preventing Caenorhabditis Elegans Mutation Accumulation." *Genetics* 174 (1): 57–65.

Des Marais, David L., and Mark D. Rausher. 2010. "Parallel Evolution at Multiple Levels in the Origin of Hummingbird Pollinated Flowers in Ipomoea." *Evolution; International Journal of Organic Evolution* 64 (7): 2044–54.

Durbin, M. 2003. "Genes That Determine Flower Color: The Role of Regulatory Changes in the Evolution of Phenotypic Adaptations." *Molecular Phylogenetics and Evolution*. https://doi.org/10.1016/s1055-7903(03)00196-9.

Eyre-Walker, Adam, and Peter D. Keightley. 2007. "The Distribution of Fitness Effects of New Mutations." *Nature Reviews. Genetics* 8 (8): 610–18.

Fisher, R. A. 1930. "The Genetical Theory of Natural Selection Oxford University Press." *London, UK*.

Grey Monroe, J., Thanvi Srikant, Pablo Carbonell-Bejerano, Moises Exposito-Alonso, Mao-Lun Weng, Matthew T. Rutter, Charles B. Fenster, and Detlef Weigel. 2020. "Mutation Bias Shapes Gene Evolution in Arabidopsis Thaliana." *Cold Spring Harbor Laboratory*. https://doi.org/10.1101/2020.06.17.156752.

Haake, Volker, Daniel Cook, José Luis Riechmann, Omaira Pineda, Michael F. Thomashow, and James Z. Zhang. 2002. "Transcription Factor CBF4 Is a Regulator of Drought Adaptation in Arabidopsis." *Plant Physiology* 130 (2): 639–48.

Halligan, Daniel L., and Peter D. Keightley. 2009. "Spontaneous Mutation Accumulation Studies in Evolutionary Genetics." *Annual Review of Ecology, Evolution, and Systematics* 40 (1): 151–72.

Hoballah, Maria Elena, Thomas Gübitz, Jeroen Stuurman, Larissa Broger, Mario Barone, Therese Mandel, Alexandre Dell'Olivo, Maeva Arnold, and Cris Kuhlemeier. 2007. "Single Gene-Mediated Shift in Pollinator Attraction in Petunia." *The Plant Cell* 19 (3): 779–90.

Hopkins, Robin, and Mark D. Rausher. 2011. "Identification of Two Genes Causing Reinforcement in the Texas Wildflower Phlox Drummondii." *Nature* 469 (7330): 411–14.

Huang, Wen, Richard F. Lyman, Rachel A. Lyman, Mary Anna Carbone, Susan T. Harbison, Michael M. Magwire, and Trudy Fc Mackay. 2016. "Correction: Spontaneous Mutations and the Origin and Maintenance of Quantitative Genetic Variation." *eLife* 5 (October). https://doi.org/10.7554/eLife.22300.

Inaba, Shoko, Rie Kurata, Mami Kobayashi, Yoko Yamagishi, Izumi Mori, Yoshiyuki Ogata, and Yoichiro Fukao. 2015. "Identification of Putative Target Genes of bZIP19, a Transcription Factor Essential for Arabidopsis Adaptation to Zn Deficiency in Roots." *The Plant Journal: For Cell and Molecular Biology* 84 (2): 323–34.

Jan, Sumira, Nazia Abbas, Muhammad Ashraf, and Parvaiz Ahmad. 2019. "Roles of Potential Plant

Hormones and Transcription Factors in Controlling Leaf Senescence and Drought Tolerance." *Protoplasma* 256 (2): 313–29.

Jeong, Sangyun, Mark Rebeiz, Peter Andolfatto, Thomas Werner, John True, and Sean B. Carroll. 2008. "The Evolution of Gene Regulation Underlies a Morphological Difference between Two Drosophila Sister Species." *Cell* 132 (5): 783–93.

Kanehisa, M., and S. Goto. 2000. "KEGG: Kyoto Encyclopedia of Genes and Genomes." *Nucleic Acids Research* 28 (1): 27–30.

Katju, Vaishali, and Ulfar Bergthorsson. 2019. "Old Trade, New Tricks: Insights into the Spontaneous Mutation Process from the Partnering of Classical Mutation Accumulation Experiments with High-Throughput Genomic Approaches." *Genome Biology and Evolution*. https://doi.org/10.1093/gbe/evy252.

Koini, Maria A., Liz Alvey, Trudie Allen, Ceinwen A. Tilley, Nicholas P. Harberd, Garry C. Whitelam, and Keara A. Franklin. 2009. "High Temperature-Mediated Adaptations in Plant Architecture Require the bHLH Transcription Factor PIF4." *Current Biology: CB* 19 (5): 408–13.

Lamesch, Philippe, Tanya Z. Berardini, Donghui Li, David Swarbreck, Christopher Wilks, Rajkumar Sasidharan, Robert Muller, et al. 2012. "The Arabidopsis Information Resource (TAIR): Improved Gene Annotation and New Tools." *Nucleic Acids Research* 40 (Database issue): D1202–10.

Leng, Pengfei, and Jun Zhao. 2020. "Transcription Factors as Molecular Switches to Regulate Drought Adaptation in Maize." *TAG. Theoretical and Applied Genetics. Theoretische Und Angewandte Genetik* 133 (5): 1455–65.

Lin, Rong-Chien, and Mark D. Rausher. 2021. "R2R3-MYB Genes Control Petal Pigmentation Patterning in Clarkia Gracilis Ssp. Sonomensis (Onagraceae)." *The New Phytologist* 229 (2): 1147–62.

Mi, Huaiyu, Dustin Ebert, Anushya Muruganujan, Caitlin Mills, Laurent-Philippe Albou, Tremayne Mushayamaha, and Paul D. Thomas. 2021. "PANTHER Version 16: A Revised Family Classification, Tree-Based Classification Tool, Enhancer Regions and Extensive API." *Nucleic Acids Research* 49 (D1): D394–403.

Monroe, J. Grey, Tyler Powell, Nicholas Price, Jack L. Mullen, Anne Howard, Kyle Evans, John T. Lovell, and John K. McKay. 2018. "Drought Adaptation in Arabidopsis Thaliana by Extensive Genetic Loss-of-Function." *eLife* 7 (December). https://doi.org/10.7554/eLife.41038.

Nguyen, Duong T., Baojun Wu, Shujie Xiao, and Weilong Hao. 2020. "Evolution of a Record-Setting AT-Rich Genome: Indel Mutation, Recombination, and Substitution Bias." *Genome Biology and Evolution*. https://doi.org/10.1093/gbe/evaa202.

Quattrocchio, Francesca, John Wing, Karel van der Woude, Erik Souer, Nick de Vetten, Joseph Mol, and Ronald Koes. 1999. "Molecular Analysis of the anthocyanin2 Gene of Petunia and Its Role in the Evolution of Flower Color." *The Plant Cell*. https://doi.org/10.2307/3870973.

Roles, Angela J., Matthew T. Rutter, Ian Dworkin, Charles B. Fenster, and Jeffrey K. Conner. 2016. "Field Measurements of Genotype by Environment Interaction for Fitness Caused by Spontaneous Mutations in Arabidopsis Thaliana." *Evolution; International Journal of Organic Evolution* 70 (5): 1039–50.

Rutter, Matthew T., Angela Roles, Jeffrey K. Conner, Ruth G. Shaw, Frank H. Shaw, Korbinian Schneeberger, Stephan Ossowski, Detlef Weigel, and Charles B. Fenster. 2012. "Fitness of Arabidopsis Thaliana Mutation Accumulation Lines Whose Spontaneous Mutations Are Known." *Evolution; International Journal of Organic Evolution* 66 (7): 2335–39.

Rutter, Matthew T., Angela J. Roles, and Charles B. Fenster. 2018. "Quantifying Natural Seasonal Variation in Mutation Parameters with Mutation Accumulation Lines." *Ecology and Evolution* 8 (11): 5575–85.

Rutter, Matthew T., Frank H. Shaw, and Charles B. Fenster. 2010. "Spontaneous Mutation Parameters for Arabidopsis Thaliana Measured in the Wild." *Evolution; International Journal of Organic Evolution* 64 (6): 1825–35.

Schrider, Daniel R., David Houle, Michael Lynch, and Matthew W. Hahn. 2013. "Rates and Genomic Consequences of Spontaneous Mutational Events in Drosophila Melanogaster." *Genetics* 194 (4):

937–54.

Shaw, Frank H., Charles J. Geyer, and Ruth G. Shaw. 2002. "A Comprehensive Model of Mutations Affecting Fitness and Inferences for Arabidopsis Thaliana." *Evolution; International Journal of Organic Evolution* 56 (3): 453–63.

Smith, Stacey D. 2016. "Pleiotropy and the Evolution of Floral Integration." *The New Phytologist* 209 (1): 80–85.

Stower, Hannah. 2013. "Adaptation by Loss of Function." *Nature Reviews Genetics*. https://doi.org/10.1038/nrg3557.

Streisfeld, Matthew A., Danya Liu, and Mark D. Rausher. 2011. "Predictable Patterns of Constraint among Anthocyanin-Regulating Transcription Factors in Ipomoea." *The New Phytologist* 191 (1): 264–74.

Streisfeld, Matthew A., and Mark D. Rausher. 2009. "Altered Trans-Regulatory Control of Gene Expression in Multiple Anthocyanin Genes Contributes to Adaptive Flower Color Evolution in Mimulus Aurantiacus." *Molecular Biology and Evolution*. https://doi.org/10.1093/molbev/msn268.

———. 2011. "Population Genetics, Pleiotropy, and the Preferential Fixation of Mutations during Adaptive Evolution." *Evolution; International Journal of Organic Evolution* 65 (3): 629–42.

Van Buskirk, Heather A., and Michael F. Thomashow. 2006. "Arabidopsis Transcription Factors Regulating Cold Acclimation." *Physiologia Plantarum* 126 (1): 72–80.

Wagner, Günter P., and Vincent J. Lynch. 2008. "The Gene Regulatory Logic of Transcription Factor Evolution." *Trends in Ecology & Evolution* 23 (7): 377–85.

Weng, Mao-Lun, Jon Ågren, Eric Imbert, Henning Nottebrock, Matthew T. Rutter, and Charles B. Fenster. 2021. "Fitness Effects of Mutation in Natural Populations of Arabidopsis Thaliana Reveal a Complex Influence of Local Adaptation." *Evolution; International Journal of Organic Evolution* 75 (2): 330–48.

Weng, Mao-Lun, Claude Becker, Julia Hildebrandt, Manuela Neumann, Matthew T. Rutter, Ruth G. Shaw, Detlef Weigel, and Charles B. Fenster. 2019. "Fine-Grained Analysis of Spontaneous Mutation Spectrum and Frequency in Arabidopsis Thaliana." *Genetics* 211 (2): 703–14.

Wessinger, Carolyn A., and Mark D. Rausher. 2013. "PREDICTABILITY AND IRREVERSIBILITY OF GENETIC CHANGES ASSOCIATED WITH FLOWER COLOR EVOLUTION IN PENSTEMON BARBATUS." *Evolution; International Journal of Organic Evolution*, December. http://doi.wiley.com/10.1111/evo.12340.

Whittall, Justen B., Claudia Voelckel, Daniel J. Kliebenstein, and Scott A. Hodges. 2006. "Convergence, Constraint and the Role of Gene Expression during Adaptive Radiation: Floral Anthocyanins in Aquilegia." *Molecular Ecology* 15 (14): 4645–57.

Wray, Gregory A., Matthew W. Hahn, Ehab Abouheif, James P. Balhoff, Margaret Pizer, Matthew V. Rockman, and Laura A. Romano. 2003. "The Evolution of Transcriptional Regulation in Eukaryotes." *Molecular Biology and Evolution* 20 (9): 1377–1419.

Xia, Jianguo, and David S. Wishart. 2011. "Metabolomic Data Processing, Analysis, and Interpretation Using MetaboAnalyst." *Current Protocols in Bioinformatics / Editoral Board, Andreas D. Baxevanis ... [et Al.]* Chapter 14 (June): Unit 14.10.

Xu, Yong-Chao, Xiao-Min Niu, Xin-Xin Li, Wenrong He, Jia-Fu Chen, Yu-Pan Zou, Qiong Wu, Yong E. Zhang, Wolfgang Busch, and Ya-Long Guo. 2019. "Adaptation and Phenotypic Diversification in Arabidopsis through Loss-of-Function Mutations in Protein-Coding Genes." *The Plant Cell* 31 (5): 1012–25.

Zalts, Harel, and Itai Yanai. 2017. "Developmental Constraints Shape the Evolution of the Nematode Mid-Developmental Transition." *Nature Ecology & Evolution* 1 (5): 113.

Acknowledgments

We thank Detlef Weigel for thoughtful comments and for the generation of the sequencing data. This work was supported by the National Science Foundation (DEB 2017485 to C.F.O.-M, OIA EPSCoR 1920954, DEB 1257902 to C.B.F., DEB 0844820 to C.B.F., DEB 1258053 to M.T.R., and DEB 0845413 to M.T.R.); and the National Institute of Health (INBRE 2P20GM103443-19).