DNA METHYLTRANSFERASE 3 (MET3) is regulated by Polycomb Group

complex during Arabidopsis endosperm development

Louis Tirot[1] and Pauline E. Jullien[1†]

[1] Institute of Plant Sciences, University of Bern, Bern, Switzerland.

[†]*Corresponding author:* Pauline E. Jullien (pauline.jullien@ips.unibe.ch)

Key message

The DNA METHYLTRANSFERASE MET3 is controlled by Polycomb group complex during

endosperm development.

Abstract

Complex epigenetic changes occur during plant reproduction. These regulations ensure the proper

transmission of epigenetic information as well as allowing for zygotic totipotency. In *Arabidopsis*, the

main DNA methyltransferase is called MET1 and is responsible for methylating cytosine in the CG

context. The *Arabidopsis* genome encodes for three additional reproduction-specific homologs of

MET1, namely *MET2a*, *MET2b* and *MET3.* In this paper, we show that the DNA methyltransferase

MET3 is expressed in the seed endosperm and its expression is further restricted to the chalazal

endosperm. *MET3* is biallelically expressed in the endosperm but displays a paternal expression bias.

We found that *MET3* expression is regulated by the Polycomb complex proteins FIE and MSI1. Seed

development is not impaired in *met3* mutant, and we could not observe significant transcriptional

changes in *met3* mutant. Interestingly, we found that *MET3* regulates gene expression in a Polycomb

mutant background suggesting a further complexification of the interplay between H3K27me3 and

DNA methylation in the seed endosperm.

Keywords

DNA methylation, *Arabidopsis*, endosperm, Polycomb, *MET3*

Introduction

Sexual reproduction in Angiosperm is initiated by a double fertilization event. Fertilization of the haploid egg cell by one of the sperm cells gives rise to the diploid embryo whereas fertilization of the homodiploid central cell gives rise to the triploid endosperm (Berger 2003; Costa et al. 2004). The endosperm represents a nourishing tissue supporting embryo growth and is therefore key for proper seed development. Cell divisions in the endosperm are initiated very rapidly following fertilization. These divisions are initially occurring without cellularization and form a syncytium that will later cellularize (Brown et al. 1999, 2003; Boisnard-Lorig et al. 2001). An additional peculiarity of the endosperm, beyond its triploid syncytial nature, is being the seat of interesting epigenetic phenomena and complex epigenetic regulation.

Endosperm development and cellularization are indeed regulated by the FIS Polycomb group complex known to mediate Histone H3 Lysine 27 tri-methylation, a key silencing epigenetic mark. Some of the key members of the FIS Polycomb group complex (FIS-PcG) are MEA, FIS2, FIE, and MSI1 (Chaudhury et al. 1997; Luo et al. 1999; Kiyosue et al. 1999; Yadegari et al. 2000; Köhler et al. 2003; Guitton et al. 2004). In mutants affecting those genes, the endosperm fails to cellularize, resulting in an arrest of embryo development and eventually seed abortion. Additionally, several genes were found to be imprinted, *i.e.* only one parental allele is expressed whereas the other is epigenetically silent, in the endosperm. This is the case, for example, of genes such as *FIS2*, *FWA*, *MEA* or *PHE1* (Kinoshita et al. 1999, 2004; Luo et al. 2000; Köhler et al. 2005; Jullien et al. 2006). The silencing of those endosperm imprinted genes relies principally on two epigenetic mechanisms: either regulation by the FIS-PcG itself like for *MEA* or *PHE1* or silencing by DNA methylation like for *FWA* and *FIS2* (Jullien and Berger 2009; Gehring 2013; Batista and Köhler 2020). Another epigenetic singularity of the endosperm is of being relatively hypomethylated compared to other plant tissues (Gehring et al. 2009; Hsieh et al. 2009). This hypomethylation is due in part to the activity of a DNA glycosylase called DEMETER (Choi et al. 2002; Gehring et al. 2006; Hsieh et al. 2009) and likely also to the low expression of canonical actors of the DNA methylation pathway (Jullien et al. 2012).

DNA methylation is a key epigenetic mark regulating gene expression and protecting genome integrity by repressing transposons. In plant genomes, DNA methylation is found in three cytosine contexts: CG, CHG and CHH (where H is any base except C). Methylation on these different contexts relies on specific DNA methyltransferases. DNA methylation on CG sites relies on maintenance DNA METHYLTRANSFERASE (MET) where the main ubiquitous enzyme is MET1. DNA methylation on CHG sites relies on CHROMOMETHYLASE3 (CMT3) and an interplay with histone methylation (Lindroth et al. 2001). DNA methylation on CHH site, due to its non-symmetrical nature, relies on the constant *de novo* methylation pathway involving small RNA molecules as well as DOMAIN REARANGED METHYLTRANSFERASE2 (DRM2) (Cao and Jacobsen 2002). In centromeric sequences, CHH methylation also relies on CHROMOMETHYLASE2 (CMT2) (Stroud et al. 2013).

Although we know a lot about the main actors of this pathway, the *Arabidopsis*'s genome encodes multiple copies of DNA methyltransferase genes (4 *METs*, 3 *CMTs* and 3 *DRMs*) some of which might have a more complex or similar function in discreet cell types. For example, *CMT1* (the third CHROMOMETHYLASE encoded by the *Arabidopsis* genome) is principally detected in reproductive tissue (Henikoff and Comai 1998; Klepikova et al. 2016) and the reconstitution of a full-length transcript relies on the splicing out of a transposable element situated in its 13[th] exon (Yadav et al. 2018). The *DOMAIN REARANGED METHYLTRANSFERASE1* (*DRM1*) seems to also solely play a role in reproductive tissue, where a redundancy between *DRM1* and *DRM2* was observed in the early embryo (Jullien et al. 2012). Similarly, data concerning the potential function of non-canonical *METs* are scarce. *MET2a* and *MET2b* are detected in the central cell, but their function is unknown (Jullien et al. 2012). Nonetheless, correlative evidence suggest *MET2a* might be important to regulate transposon reactivation in wild *Arabidopsis* accessions (Quadrana et al. 2016) and involved in fungal response (Salvador-Guirao et al. 2018).

As mentioned, little is known about the DNA methyltransferase *MET3*. *MET3* is also named *MATERNAL EFFECT EMBRYO ARREST 57* (*MEE57*) as a transposon insertion associated with the *MET3*

locus led to an arrest in endosperm development (Pagnussat et al. 2005). *MET3* is also reported to be

the sole *MET* expressed in the endosperm (Jullien et al. 2012). Here, we show that *MET3* is specifically

expressed in the endosperm in a biallelic fashion with a paternal bias. *MET3* expression is controlled

by the FIS-PcG complex. Despite the initial report of a seed arrest phenotype in the *mee57* line, we

did not see any seed developmental phenotype in two independent *met3* mutant alleles. Additionally,

we could not see major changes in the seed transcriptome of *met3* mutant. Nevertheless, we could

see an effect on the seed transcriptome in a *fie* mutant background suggesting that *MET3* might

interplay with PcG gene regulation in the developing endosperm.

Material and Methods

Plant Materials, Growth Conditions and Genotyping.

The wild-types Col-0 and Gr-1, the *MET3* mutant lines *met3-3* (GABI404F04), *met3-4* (GABI659H03)

and PRC2 mutant lines *fie-362* (GABI_362D08) (Bouyer et al. 2011) and *msi1* (SAIL_429_B08) were

provided by the Nottingham Arabidopsis Stock Center (NASC). *pMET3:H2B-RFP* line was previously

described (Jullien et al. 2012). After sowing, plants were stratified in the dark at 4°C between 2 and 4

days. Plants were germinated and grown in growing chambers under long-day conditions (16h light

22°C / 8h dark 18°C). For the transmission analysis (Fig. S5b) , plants were grown on Murashige and

Skoog (MS1/2) media agarose plate in long-day conditions for 12 days before genotyping. Primers for

genotyping are listed in Table S1.

Microscopy and phenotype observation

DIC seed phenotype and GUS observations were done using a Leica DM2000 as previously described

(Jullien et al. 2006). For seed development observation and counting (Fig. 4a) , plants were

synchronized by removing all open flowers from the inflorescences. After 6 days, the two first siliques

situated above the previously removed flowers were picked for each inflorescence. We refer to this

stage as 6 days after synchronization (6 DAS). Siliques were dissected, cleared using chloride hydrate

solution and mounted on a slide for observation. *pMET3:H2B-tdTomato* and *pMET3:H2B-RFP* reporter

lines were imaged using a laser scanning confocal microscope (Leica SP5). When necessary, brightness

and contrast were uniformly modified by using ImageJ.

Cloning and transformation

pMET3:H2B-tdTomato and *pMET3:H2B-GUS* were generated using the Gateway Cloning System

(Invitrogen). All PCR fragments were amplified by PCR using the Phusion High-Fidelity DNA Polymerase

(Thermo). Primer sequences used for cloning can be found in Table S1. All plasmids were transformed

into wild-type Columbia-0 plants the by floral dipping method (Clough and Bent 1998). At least ten

transgenic lines were analyzed per construct, which showed a consistent fluorescence expression

pattern. An Illustration of the constructs can be found in Fig. S2a.

RNA extraction, qPCR & RT-PCR

Total RNAs were extracted using RNeasy Plant Minikit (Qiagen). All samples were treated with DNase

I (ThermoScientific) at 37°C for 30 minutes. DNAse I was subsequently inactivated by the addition of

EDTA and heat treatment (65°C for 10 minutes). First-strand cDNAs were synthesized using between

500 and 1000 ng of DNase treated total RNAs as a template. The RT reaction was performed using

either Maxima First Strand cDNA Synthesis Kit (Fig. 1a, Fig. 2a) (ThermoScientific) , containing both

oligo-dT and random hexamer primers, or RevertAid RT Reverse Transcription Kit (Fig. 3e, Fig. S4b)

(ThermoScientific) , containing only oligo-dT primers. The qPCR reactions were performed with a

QuantStudio 5 thermocycler (ThermoScientific) using SYBR green (KAPA SYBR FAST qPCR Master Mix

or ORA qPCR Green ROX H Mix). The qPCR mix was prepared according to the manufacturer's protocol.

An RNA equivalent of 25 ng of cDNA templates was used for each reaction. The qPCR program was as

follow: 95 °C for 3 minutes followed by 45 cycles of 95 °C for 5 seconds and 60 °C for 30 seconds.

ACTIN2 (*AT3G18780*) expression was used to normalize the transcript level in each sample. For each

condition, RNA abundance of target genes was calculated from the average of three independent

biological replicates with three qPCR technical replicates. Real-time PCR primers used in this study are

listed in Table S1. For the allele-specific RT-PCR (Fig.2a) , cDNAs were amplified for 22 cycles for *ACT2*

primers and 35 cycles for *MET3* specific primers. Half of the *MET3* PCR product was digested for 1h30

at 37°C with XbaI restriction enzyme (Sigma-Aldrich). *ACT2* amplification was used as a control.

RNA sequencing and Bioinformatics

Total RNAs were extracted and DNAseI treated as previously mentioned. mRNA libraries were

prepared and sequenced by Novogene (https://en.novogene.com/). Bioinformatic analyses were

performed on the Galaxy web platform (https://usegalaxy.org) (Afgan et al. 2018). Our Galaxy

workflow including the exact parameters and tool versions used can be downloaded on

https://usegalaxy.org/u/pej/w/pejrnaseq and be freely reused. Briefly, Paired-end raw mRNA

sequencing reads were controlled using FastQC (Galaxy version 0.72) and trimmed using Trimmomatic

(Galaxy version 0.36.6) (Andrews 2010; Bolger et al. 2014). Clean reads were aligned to the *Arabidopsis*

thaliana TAIR 10 genome assembly using HISAT2 (Galaxy version 2.1.0+galaxy4) (Kim et al. 2015).

Aligned sequencing reads were assigned to genomic features using featureCounts (Galaxy version

1.6.2) (Liao et al. 2014). Differential expression was analyzed using DESeq2 default parameters (Galaxy

Version 2.11.40.6+galaxy1) (Love et al. 2014). Differentially expressed genes (DEGs) were defined by

an absolute logFC > 2 and an FDR < 0.05. Gene ontology (GO) enrichment analysis has been performed

on PANTHER (Mi et al. 2019) and visualized using REVIGO (Supek et al. 2011). All plots have been

generated using R-studio (www.rstudio.com). Raw data are deposited on GEO under reference XXXX.

Results

MET3 is expressed biallelically with a paternal bias in the endosperm

Our previous analysis has shown that *MET3* is expressed in the endosperm of developing

Arabidopsis seeds (Jullien et al. 2012). However, the detail and exclusivity of its expression pattern

remain to be investigated. To get a better characterization of *MET3* expression pattern, we performed

a qPCR of *MET3* transcript in different tissue types of wild-type Col-0 (Fig. 1a). Our result shows that

MET3 is principally expressed in siliques and its expression peaks at 5 Days After Pollination (DAP).

These results could be confirmed using publicly available transcriptome datasets (Fig. S1a) (Klepikova

et al. 2016). To gain a better spatial and temporal characterization of *MET3* expression, we generated

two new transcriptional *MET3* reporter constructs, encompassing 2kb of the *MET3* promoter driving

either H2B-tdTomato or H2B-GUS (Fig. S2a). The analysis of the *pMET3:H2B-GUS* in different plant

tissues confirmed the specificity of *MET3* expression to the seed endosperm (Fig. S1b-d). To

characterize in detail the temporal expression of *MET3* in the endosperm, we performed confocal

microscopy on the *pMET3:H2B-tdTomato* lines at different stages of seed development. We could

detect *pMET3:H2B-tdTomato* expression from as early as the four nuclei stage endosperm (Fig. 1b).

pMET3:H2B-tdTomato remains express throughout the endosperm (Fig. 1c) until the globular stage of

embryo development where its expression starts to be higher in the chalazal endosperm and chalazal

cyst (Fig. 1d). At later stages, *pMET3:H2B-tdTomato* expression is limited to the chalazal endosperm

and chalazal cyst (Fig. 1e). From 7 DAP, *pMET3:H2B-tdTomato* expression can no longer be detected.

A similar expression pattern was observed with *pMET3:H2B-GUS* (Fig. S2b) and *pMET3:H2B-RFP* (Fig.

S2c) as well as online transcriptomic data (Fig. S2d) (Belmonte et al. 2013). MET3 protein expression

and localization could not be determined as we, so far, failed in expressing a fluorescently tagged

MET3 protein in *Arabidopsis* (LT personal communication).

Such endosperm expression pattern is common in imprinted genes, like *FWA*, *FIS2*, *MEA* or *PHE1*

(Kinoshita et al. 1999, 2004; Köhler et al. 2005; Jullien et al. 2006). In order to examine if *MET3* is

biallelically or mono-allelically expressed, we performed allele-specific RT-PCR. We are making use of

a Short Nucleotide Polymorphism (SNP) consisting of a substitution from a C to a T within *MET3* 9th

exon in the Gr-1 ecotype which is abolishing a XbaI restriction site present in Col-0. We did reciprocal

crosses using Col-0 and Gr-1 ecotypes and analyzed *MET3* parental expression at 5 DAP following XbaI

digestion (Fig. 2a). We could observe, for both reciprocal crosses, bands corresponding to *MET3*

transcript from Col-0 (505 and 329 bp) and from Gr-1 (834 bp) with a bias toward the paternal allele.

This result shows that *MET3* is expressed from both maternal and paternal allele but displays a

paternal bias of expression. *MET3* paternally biased expression was also observed using the

pMET3:H2B-RFP transgene at 1 DAP (Fig.2b). Taking together our results shows that *MET3* expression

is biallelic with a paternal bias and confined to the endosperm. *MET3* expression, initially throughout

the endosperm, becomes restricted to the chalazal pole at later stages.

MET3 expression is regulated by Polycomb group proteins

Beyond imprinted genes, *MET3* expression pattern is reminiscent of the expression of genes

controlled by the endosperm FIS Polycomb group complex (PcG) composed of FIE, MSI1, FIS2 and MEA

(Guitton et al. 2004). To investigate if *MET3* could be regulated by PcG, we introgressed a *MET3*

transcriptional reporter into *fie* and *msi1* mutant background (Fig. 3a-d). We could observe increased

pMET3:H2B-RFP reporter expression in 49% of the seeds (n=249) in *fie/+* mutant and 44% (n=240) in

msi1/+ mutant background characteristic of the maternal gametophytic effect of those mutations (Fig.

3d). In *msi1* and *fie* mutants, *pMET3:H2B-RFP* expression is higher and observed throughout the

endosperm (Fig. 3b-c) whereas at the same developmental stage the expression of *pMET3:H2B-RFP* is

already restricted to the chalazal pole in wild-type seeds (Fig. 3a). In order to confirm that the

regulation of *MET3* by *MSI1* and *FIE* was not only restricted to our transgene, we performed a RT-qPCR

of *MET3* expression in wild-type and mutant selected seeds at 10 DAP (Fig. 3e). We could observe a

clear upregulation of *MET3* expression in both *fie* and *msi1* seeds compare to wild-type seeds (t-test

p-value of 0.0543 and 0.0114 respectively). Using publicly available data, we could see that the

upregulation of *MET3* is also observed in other PcG mutants (Fig. S3a-b). *MET3* upregulation was

observed in silique samples of *clf* mutant (Fig. S3a) (Liu et al. 2016) and in *fis2* seeds (Fig. S3b)

(Weinhofer et al. 2010).

Subsequently, we wanted to know if the effect of the PcG complex was direct or indirect. PcG

complexes are known to repress gene expression by tri-methylating the Lysine 27 of the Histone H3

tail (H3K27me3) inducing a closed chromatin state at the targeted loci. We, therefore, analyzed

available H3K27me3 genome-wide Chromatin Immunoprecipitation (ChIP) data. We could see that

the *MET3* locus is covered by H3K27me3 in *Arabidopsis* seedling samples (Fig. 3f) (Zhang et al. 2007).

Additionally, using the RepMap2020 tool (Chèneby et al. 2020) , we could observe H3K27me3 on the

MET3 locus in several independent ChIP experiments including some performed on endosperm tissue

(Fig. S3c). We conclude that *MET3* expression is directly repressed by PcG complex induced H3K27me3

in the endosperm.

MET3 does not affect seed development

Considering that *MET3* is specifically expressed in the developing endosperm during seed

development, we then ask if *MET3* function influences seed and/or endosperm development. We

characterized two mutant alleles from the GABI collection: *met3-3* (GABI_404F04) and *met3-4*

(GABI_659H03) (Fig. S4a). The mutations are located on the 10th and the 2nd exon respectively and are

expected to abolish *MET3* function. To confirm the downregulation of *MET3* in the mutants, we

performed RT-qPCR. We could observe that *MET3* is downregulated in both mutant alleles (Fig. S4b).

To investigate if the *met3* mutation could result in seed lethality we first investigated the presence of

aborted seeds at the green seed stage (~12DAP). We could not see any significant seed abortion in

both *met3-3* and *met3-4* alleles (Fig. S5a). To further confirm the absence of defects, we analyzed the

transmission rate of the *met3* mutations in *met3-3/+* and *met3-4/+* selfed progeny. We could not

observe any segregation distortion from the Mendelian ratio (Fig. S5b). Mutations affecting the main

Arabidopsis DNA methyltransferase *MET1* display variation in seed size. We, therefore, investigated if

met3 mutants would display a seed size phenotype. We could not see any significant variation in seed

size using both *met3* alleles (Fig. S5c-d). We conclude that *MET3* mutation does not severely impair

seed development.

As shown above, *MET3* expression is regulated by PcG complex in the endosperm, we then

analyzed if *MET3* mutation could influence the PcG *fie* mutant phenotype. To answer this question,

we generated *fie/+;met3-3* and *fie/+;met3-4* double mutants and analyzed their seed development

phenotype using DIC (Fig. 4a-f). In order to minimize the stress to the plant due to handling during

emasculation and crossing, we used "synchronized seeds". In practice, we remove the open flowers

of the day, and we wait an x number of days before collecting two siliques above our cutting. This is

allowing us to have age synchronized siliques without the physical disturbance of

emasculation/pollination and is, therefore, closer to normal growth and fertilization. We are using the

term Day After Synchronization (DAS). To our experience, 6 DAS is corresponding to around 4-5DAP.

At 6 DAS, we could not see any delay in embryo development and endosperm cellularization

comparing Col-0 to *met3-3* and *met3-4* mutants (Fig. 4a-c, 4g-h). In *fie*/+, we could clearly see the

delayed endosperm cellularization characteristic of FIS-PcG mutants (Fig. 4d-f and h) (56% n=407)

(Ohad et al. 1999; Sørensen et al. 2001). In the double mutants, we could not see variations in the

quantification of the *fie*/+ phenotype for both embryo development (Fig. 4g) and endosperm

cellularization (Fig. 4h). We conclude that *MET3* mutations do not influence *fie* mutant seed

phenotype.

Effect of MET3 mutation on the seed transcriptome

In order to investigate the potential effect of *met3* mutation on the seed transcriptome, we

performed a mRNA deep sequencing experiment of 3 DAP (Fig. S6) and 10 DAS (Fig. 5a) seeds (*i.e.*

seeds attached to the septum). We then compared the seed transcriptome of *met3* seeds to wild-type

seeds. At 3 DAP, we could only detect one differentially expressed gene: *ESM1* (*AT3G14210*). *ESM1*

was up-regulated in both *met3-3* and *met3-4* mutant seeds (Fig. S6a-b). At 10 DAS, we could also only

see very minor changes to the seed transcriptome (Fig. 5a). We could identify only 16 differentially

expressed genes with an absolute logFC>2 and FDR<0.05. These results show that reminiscent of the

absence of seed phenotype in *met3* mutants, the seed transcriptome is also mostly unaffected by the

met3 mutation. As *MET3* expression is regulated by the FIS-PcG complex in the endosperm, we then

ask if *MET3* mutation could influence the *fie* transcriptome. We, therefore, sequenced the

transcriptome of 10 DAS seeds where we selected for *fie* mutant seeds under the dissecting

microscope (white seeds). We compared the transcriptome of *fie* seeds to the transcriptome of

fie;met3-3 double mutant seeds. We could observe 87 differentially expressed genes (DEG) with an

absolute logFC>2 and FDR<0.05 (Fig 5b). A Goterm enrichment analysis revealed that these genes are

enriched for genes involved in pectin metabolism (GO:0045490, FDR=4.46E-10; GO:0045488,

FDR=6.59E-09) and cell-wall related processes (GO:0042545, FDR= 8.97E-09; GO:0071555, FDR=4.87E-

10, GO:0071554, FDR=5.67E-08) (Fig. 5c and Table S2). We then analyzed if among these 87 DEGs

some are also modified in *fie* mutant compared to wild-type. We could find that a large proportion of

either *met3* DEGs (Col vs *met3*, 11/16) and *met3;fie* DEGs (*fie* versus *met3fie*, 44/87) are miss-

regulated in *fie* mutant seeds (Fig. 5d). We conclude that the *met3* mutation does not drastically

change the seed transcriptome, but a set of genes are miss-regulated by both *met3 and fie mutants*.

Discussion

Our study highlights an additional connection between DNA Methylation pathways and Polycomb

group H3K27 tri-methylation in the seed endosperm via the regulation of *MET3* by the FIS-PcG

complex. *MET3* is specifically expressed in the endosperm, and its expression becomes restricted to

the chalazal pole at later stages of endosperm development. In our study, using two independent

insertion lines, *met3-3* and *met3-4*, we could not observe any major seed developmental phenotype.

Both mutants show a significant decrease in *MET3* expression by qPCR suggesting they both represent

knockout mutants. It was previously documented that a mutation affecting *MET3* named *mee57*

displayed a strong seed developmental phenotype (Pagnussat et al. 2005). The *mee57* mutation shows

an early maternal embryo and endosperm arrest. The discrepancy between our lines and the

previously published *mee57* could have several causes: the mutagenesis method used (T-DNA versus

transposition) , the presence of additional genetic modifications, or a difference between the two

ecotypes used, Columbia-0 in our case and Landsberg for *mee57*. If the latter is true, *MET3* function

could vary between different *Arabidopsis* accessions.

In this work, we show that repression of *MET3* expression at later stages of endosperm

development is linked to the direct action of the FIS-PcG complex on the *MET3* locus. *MET3* is over-

expressed in PcG mutant seeds, such as *fie* and *msi1* mutant seeds. Interestingly, it was previously

shown that *fie* mutant endosperm display higher CG methylation compared to wild-type endosperm

and lower CHG and CHH (Ibarra et al. 2012). This increased CG methylation in *fie* mutant seems to be

restricted to the endosperm as it is not observed in *fie* seedling or *fie* embryo methylome (Bouyer et

al. 2017). We propose that higher expression of *MET3* in the *fie* endosperm could be the cause for the

increase in CG methylation specifically in the endosperm. Indeed, in our study, we could not observe

any change in *MET3* tissue expression pattern in PcG mutants but the increased expression is still

restricted to the endosperm. In further studies, the Investigation of the endosperm methylome in

met3 and *met3;fie* mutants will allow to test this hypothesis. Additionally, further experiments are

required to determine if MET3 is a functional DNA methyltransferase.

As we previously mentioned, *MET3* function could vary in between wild *Arabidopsis* accessions.

Similarly, to CMT1, several SNPs can be found at the *MET3* locus in-between different ecotypes

suggesting that *MET3* might not be fully functional in all of them. Additionally, looking closer at the

MET3 locus, we could detect the presence of a Class 2 DNA transposon (*AT4TE34810*) situated in the

third intron of the *MET3* gene. In addition to PcG regulation, *MET3* expression and possibly function

could be influenced by the presence or regulation of *AT4TE34810* like it is the case for *CMT1* transcript

(Yadav et al. 2018). The study of *MET3* function and imprint in different accession could reveal more

about its function in natural growth conditions. One hint of *MET3* potential function in Columbia-0

came from the study of methylome stability across several generations (Becker et al. 2011). In this

study, methylome stability was investigated in several Col-0 selfed lineages (30 generations). The line

accumulating the most methylation polymorphisms had concomitantly acquired a SNP in the *MET3*

gene. It was therefore suggested that *MET3* could be the cause of such methylome instability (Becker

et al. 2011; Schmitz and Ecker 2012). Analyzing the methylome of selfed *met3* mutants after 30

generations would allow to test this hypothesis. If true and taking into account the endosperm

specificity of *MET3* expression, it would indicate that the endosperm methylome influences the

embryonic methylome, a hypothesis that has stimulated a lot of interest over the last 10 years but

remains to be demonstrated.

Acknowledgments

304   We would like to thank the following people for their help: Jasmine Sekulovski for support

305   concerning plant growth, Diane Bonnet for technical support. We thank Mathieu Ingouff and Diane

306   Bonnet for critical reading of the manuscript.

Funding

308   PEJ and LT are supported by an SNF professorship grant (no.163946) attributed to PEJ.

Author contributions

310   PEJ conceived the study. LT performed the experiments. LT and PEJ analyzed the data. PEJ wrote

311   the manuscript with the help of LT.

Conflicts of interest

313   Authors state that there is no conflict of interest.

References

Afgan E, Baker D, Batut B, et al (2018) The Galaxy platform for accessible, reproducible and

collaborative biomedical analyses: 2018 update. Nucleic Acids Res 46:W537–W544.

https://doi.org/10.1093/nar/gky379

Andrews S (2010) FastQC: a quality control tool for high throughput sequence data.

Batista RA, Köhler C (2020) Genomic imprinting in plants-revisiting existing models. Genes Dev

34:24–36. https://doi.org/10.1101/gad.332924.119

Becker C, Hagmann J, Müller J, et al (2011) Spontaneous epigenetic variation in the Arabidopsis

thaliana methylome. Nature 480:245–9. https://doi.org/10.1038/nature10555

Belmonte MF, Kirkbride RC, Stone SL, et al (2013) Comprehensive developmental profiles of gene

activity in regions and subregions of the Arabidopsis seed. Proc Natl Acad Sci 110:E435–E444.

https://doi.org/10.1073/pnas.1222061110

Berger F (2003) Endosperm: The crossroad of seed development. Curr Opin Plant Biol 6:42–50.

https://doi.org/10.1016/S1369526602000043

Boisnard-Lorig C, Colon-Carmona A, Bauch M, et al (2001) Dynamic analyses of the expression of the

HISTONE::YFP fusion protein in arabidopsis show that syncytial endosperm is divided in mitotic

domains. Plant Cell 13:495–509

Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: A flexible trimmer for Illumina sequence data.

Bioinformatics 30:2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Bouyer D, Kramdi A, Kassam M, et al (2017) DNA methylation dynamics during early plant life.

Genome Biol 18:179. https://doi.org/10.1186/s13059-017-1313-0

Bouyer D, Roudier F, Heese M, et al (2011) Polycomb repressive complex 2 controls the embryo-to-

seedling phase transition. PLoS Genet 7:. https://doi.org/10.1371/journal.pgen.1002014

Brown RC, Lemmon BE, Nguyen H (2003) Events during the first four rounds of mitosis establish

three developmental domains in the syncytial endosperm of Arabidopsis thaliana. Protoplasma

222:167–74. https://doi.org/10.1007/s00709-003-0010-x

Brown RC, Lemmon BE, Nguyen H, Olsen OA (1999) Development of endosperm in Arabidopsis

thaliana. Sex Plant Reprod 12:32–42. https://doi.org/10.1007/s004970050169

Cao X, Jacobsen SE (2002) Locus-specific control of asymmetric and CpNpG methylation by the DRM

and CMT3 methyltransferase genes. Proc Natl Acad Sci U S A 99:16491–16498.

https://doi.org/10.1073/pnas.162371599

Chaudhury a M, Ming L, Miller C, et al (1997) Fertilization-independent seed development in

Arabidopsis thaliana. Proc Natl Acad Sci U S A 94:4223–8

Chèneby J, Ménétrier Z, Mestdagh M, et al (2020) ReMap 2020: A database of regulatory regions

from an integrative analysis of Human and Arabidopsis DNA-binding sequencing experiments.

Nucleic Acids Res 48:D180–D188. https://doi.org/10.1093/nar/gkz945

Choi Y, Gehring M, Johnson L, et al (2002) DEMETER, a DNA glycosylase domain protein, is required

for endosperm gene imprinting and seed viability in arabidopsis. Cell 110:33–42

Clough SJ, Bent AF (1998) Floral dip: A simplified method for Agrobacterium-mediated

transformation of *Arabidopsis thaliana*. Plant J 16:735–743. https://doi.org/10.1046/j.1365-

313X.1998.00343.x

Costa LM, Gutièrrez-Marcos JF, Dickinson HG (2004) More than a yolk: The short life and complex

times of the plant endosperm. Trends Plant Sci 9:507–514.

https://doi.org/10.1016/j.tplants.2004.08.007

Gehring M (2013) Genomic imprinting: insights from plants. Annu Rev Genet 47:187–208.

https://doi.org/10.1146/annurev-genet-110711-155527

Gehring M, Bubb KL, Henikoff S (2009) Extensive demethylation of repetitive elements during seed

development underlies gene imprinting. Science (80-) 324:1447–51.

https://doi.org/10.1126/science.1171609

Gehring M, Huh JH, Hsieh TF, et al (2006) DEMETER DNA Glycosylase Establishes MEDEA Polycomb

Gene Self-Imprinting by Allele-Specific Demethylation. Cell 124:495–506.

https://doi.org/10.1016/j.cell.2005.12.034

Guitton A, Page DR, Chambrier P, et al (2004) Identification of new members of Fertilisation

Independent Seed Polycomb Group pathway involved in the control of seed development in

Arabidopsis thaliana. Development 2971–2981. https://doi.org/10.1242/dev.01168

Henikoff S, Comai L (1998) A DNA Methyltransferase Homolog With a Chromodomain Exists in

Multiple Polymorphic Forms in Arabidopsis. Genetics 149:307–318

Hsieh TT-F, Ibarra CA, Silva P, et al (2009) Genome-wide demethylation of Arabidopsis endosperm.

Science (80-) 324:1451–1454. https://doi.org/10.1126/science.1172417

Ibarra CA, Feng X, Schoft VK, et al (2012) Active DNA Demethylation in Plant Companion Cells

Reinforces Transposon Methylation in Gametes. Science (80-) 337:1360–1364.

https://doi.org/10.1126/science.1224839

Jullien PE, Berger F (2009) Gamete-specific epigenetic mechanisms shape genomic imprinting. Curr

Opin Plant Biol 12:637–642. https://doi.org/10.1016/j.pbi.2009.07.004

Jullien PE, Kinoshita T, Ohad N, Berger F (2006) Maintenance of DNA Methylation during the

Arabidopsis Life Cycle Is Essential for Parental Imprinting. Plant Cell 18:1360–1372.

381 https://doi.org/10.1105/tpc.106.041178.1

382 Jullien PE, Susaki D, Yelagandula R, et al (2012) DNA methylation dynamics during sexual

383 reproduction in Arabidopsis thaliana. Curr Biol 22:1825–1830.

384 https://doi.org/10.1016/j.cub.2012.07.061

385 Kim D, Langmead B, Salzberg SL (2015) HISAT: A fast spliced aligner with low memory requirements.

386 Nat Methods 12:357–360. https://doi.org/10.1038/nmeth.3317

387 Kinoshita T, Miura A, Choi Y, et al (2004) One-Way Control of FWA Imprinting in Arabidopsis

388 Endosperm by DNA Methylation. Science (80-) 303:521–523.

389 https://doi.org/10.1126/science.1089835

390 Kinoshita T, Yadegari R, Harada JJ, et al (1999) Imprinting of the MEDEA polycomb gene in the

391 Arabidopsis endosperm. Plant Cell 11:1945–52

392 Kiyosue T, Ohad N, Yadegari R, et al (1999) Control of fertilization-independent endosperm

393 development by the MEDEA polycomb gene in Arabidopsis. Proc Natl Acad Sci 96:4186–4191.

394 https://doi.org/10.1073/pnas.96.7.4186

395 Klepikova A V., Kasianov AS, Gerasimov ES, et al (2016) A high resolution map of the Arabidopsis

396 thaliana developmental transcriptome based on RNA-seq profiling. Plant J 88:1058–1070.

397 https://doi.org/10.1111/tpj.13312

398 Köhler C, Hennig L, Bouveret R, et al (2003) Arabidopsis MSI1 is a component of the MEA/FIE

399 Polycomb group complex and required for seed development. EMBO J 22:4804–14.

400 https://doi.org/10.1093/emboj/cdg444

401 Köhler C, Page DR, Gagliardini V, Grossniklaus U (2005) The Arabidopsis thaliana MEDEA Polycomb

402 group protein controls expression of PHERES1 by parental imprinting. Nat Genet 37:28–30.

https://doi.org/10.1038/ng1495

Liao Y, Smyth GK, Shi W (2014) FeatureCounts: An efficient general purpose program for assigning

sequence reads to genomic features. Bioinformatics 30:923–930.

https://doi.org/10.1093/bioinformatics/btt656

Lindroth a M, Cao X, Jackson JP, et al (2001) Requirement of CHROMOMETHYLASE3 for maintenance

of CpXpG methylation. Science 292:2077–80. https://doi.org/10.1126/science.1059745

Liu J, Deng S, Wang H, et al (2016) CURLY LEAF regulates gene sets coordinating seed size and lipid

biosynthesis. Plant Physiol 171:424–436. https://doi.org/10.1104/pp.15.01335

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-Seq

data with DESeq2. bioRxiv 1–21. https://doi.org/10.1101/002832

Luo M, Bilodeau P, Dennis ES, et al (2000) Expression and parent-of-origin effects for FIS2, MEA, and

FIE in the endosperm and embryo of developing Arabidopsisseeds. PNAS.

https://doi.org/10.1073/pnas.170292997

Luo M, Bilodeau P, Koltunow a, et al (1999) Genes controlling fertilization-independent seed

development in Arabidopsis thaliana. Proc Natl Acad Sci U S A 96:296–301

Mi H, Muruganujan A, Huang X, et al (2019) Protocol Update for large-scale genome and gene

function analysis with the PANTHER classification system (v.14.0). Nat Protoc 14:703–721.

https://doi.org/10.1038/s41596-019-0128-8

Ohad N, Yadegari R, Margossian L, et al (1999) Mutations in FIE, a WD polycomb group gene, allow

endosperm development without fertilization. Plant Cell 11:407–415.

https://doi.org/10.1105/tpc.11.3.407

Pagnussat GC, Yu HJ, Ngo QA, et al (2005) Genetic and molecular identification of genes required for

425   female gametophyte development and function in Arabidopsis. Development 132:603–614.

426   https://doi.org/10.1242/dev.01595

427 Quadrana L, Silveira AB, Mayhew GF, et al (2016) The Arabidopsis thaliana mobilome and its impact

428   at the species level. Elife 5:1–25. https://doi.org/10.7554/eLife.15716

429 Salvador-Guirao R, Baldrich P, Weigel D, et al (2018) The MicroRNA miR773 Is Involved in the

430   *Arabidopsis* Immune Response to Fungal Pathogens. Mol Plant-Microbe Interact 31:249–259.

431   https://doi.org/10.1094/MPMI-05-17-0108-R

432 Schmitz RJ, Ecker JR (2012) Epigenetic and epigenomic variation in Arabidopsis thaliana. Trends Plant

433   Sci 17:149–154. https://doi.org/10.1016/j.tplants.2012.01.001

434 Sørensen MB, Chaudhury AM, Robert H, et al (2001) Polycomb group genes control pattern

435   formation in plant seed. Curr Biol 11:277–281

436 Stroud H, Do T, Du J, et al (2013) Non-CG methylation patterns shape the epigenetic landscape in

437   Arabidopsis. Nat Struct Mol Biol 21:64–72. https://doi.org/10.1038/nsmb.2735

438 Supek F, Bošnjak M, Škunca N, Šmuc T (2011) Revigo summarizes and visualizes long lists of gene

439   ontology terms. PLoS One 6:. https://doi.org/10.1371/journal.pone.0021800

440 Weinhofer I, Hehenberger E, Roszak P, et al (2010) H3K27me3 Profiling of the Endosperm Implies

441   Exclusion of Polycomb Group Protein Targeting by DNA Methylation. PLoS Genet 6:e1001152.

442   https://doi.org/10.1371/journal.pgen.1001152

443 Winter D, Vinegar B, Nahal H, et al (2007) An "electronic fluorescent pictograph" Browser for

444   exploring and analyzing large-scale biological data sets. PLoS One 2:1–12.

445   https://doi.org/10.1371/journal.pone.0000718

446 Yadav NS, Khadka J, Domb K, et al (2018) CMT3 and SUVH4/KYP silence the exonic Evelknievel

retroelement to allow for reconstitution of CMT1 mRNA 06 Biological Sciences 0604 Genetics.

Epigenetics and Chromatin 11:1–12. https://doi.org/10.1186/s13072-018-0240-y

Yadegari R, Kinoshita T, Lotan O, et al (2000) Mutations in the FIE and MEA genes that encode

interacting polycomb proteins cause parent-of-origin effects on seed development by distinct

mechanisms. Plant Cell 12:2367–2382

Zhang X, Clarenz O, Cokus S, et al (2007) Whole-genome analysis of histone H3 lysine 27

trimethylation in Arabidopsis. PLoS Biol 5:e129. https://doi.org/10.1371/journal.pbio.0050129

Figure legends

Figure 1. MET3 expression pattern

(a) *MET3* expression measured by RT-qPCR. DAP = Day After Pollination. The histogram displays the

mean, and each dot represents a biological replicate. *ACT2* is used as a normalizer. RQ = Relative

Quantification (b-e) Single-plan Confocal images representing the expression of *pMET3:H2B-*

tdTomato construct in *Arabidopsis* 1 Day After Synchronization (DAS) seed (b), 2 DAS seed (c), 3 DAS

(d) and 5 DAS (e). cze = chalazal endosperm, endo = endosperm nuclei. Scale bars represent 25μm.

Figure 2. MET3 is biallelically expressed with a paternal bias

(a) Allele specific RT-PCR of *MET3* parental expression in 5DAP silique samples. The *XbaI* restriction

enzyme digests the Col-0 *MET3* transcript but not the Gr-1 transcript. *ACT2* is used as loading control.

(b) Single-plan confocal images of *pMET3:H2B-RFP* parental expression in the endosperm of 1 DAP

seeds. endo = endosperm nuclei. Scale bars represent 25μm. ♂ symbol indicates the genotype of the

male parent while ♀ indicates the genotype of the female parent.

Figure 3. MET3 expression is controlled by FIE and MSI1 Polycomb proteins

(a-c) Confocal images representing the expression of *pMET3:H2B-RFP* construct in *Arabidopsis*

wildtype (a), *fie* (b) and *msi1* (c) selfed seeds at 5 days after pollination (DAP). Scale bars represent 50

μm. (d) Proportion of seeds with high RFP signal in the endosperm at 5DAP in wild-type, *fie* and *msi1*

mutants. Grey bars represent the seeds with a restricted RFP expression to the chalazal endosperm

(as illustrated in (a)). Black bars represent the seeds with ectopic expression of *pMET3:H2B-RFP*

throughout the endosperm (as illustrated in (b-c)). (e) *MET3* expression measured by RT-qPCR in 10

DAP selected seeds of wild-type, *fie* and *msi1*. The histogram displays the mean, and each dot

represents a biological replicate. *ACT2* is used as a normalizer. (f) Snapshot showing that the *MET3*

locus contains H3K27me3 and LHP1 but no H3K4me1, H3K4me2, H3K4me3 and H3K9me2 in seedlings.

Data from http://epigenomics.mcdb.ucla.edu/H3K27m3/ (Zhang et al. 2007).

Figure 4. MET3 does not influence fie phenotype

(a-f) Seed developmental phenotype observed after clearing by Difference Interference Contrast (DIC)

of wild-type (a), *met3-3* (b), *met3-4* (c), *fie/+* (d), *met3-3;fie/+* (e) and *met3-4;fie/+* (f) at 6 days after

synchronization (DAS). Scale bars represent 50 μm. (g-h) Quantification of the embryo developmental

stages (g) and endosperm cellularization (h) in wild-type, *met3-3*, *met3-4*, *fie/+*, *met3-3;fie/+* and

met3-4;fie/+ at 6 DAS. mce = micropylar endosperm.

Figure 5. Transcriptome of met3 and met3;fie mutant seeds

(a-b) Volcano plot depicting differentially expressed genes (DEGs) comparing Col-0 to *met3-3* (a) and

fie to *met3-3;fie* in 10 DAS seeds (b). Red dots represent up-regulated DEGs and blue dots represent

down-regulated DEGs. The top 10 DEGs are annotated. We use a threshold of [absolute logFC > 2,

FDR < 0.05]. (c) GO term analysis for the 87 DEGs obtained with the *fie vs met3-3;fie* contrast. (d)

Venn diagram showing the overlapping DEGs between the different contrasts.

Figure S1. MET3 is only expressed in seeds

(a) Snapshot of the *MET3* expression pattern obtained from ebar using the data from Klepikova *et al.*

(Winter et al. 2007; Klepikova et al. 2016)(b-d) The absence of *pMET3:H2B-GUS* expression observed

in *Arabidopsis* seedling (b), stamen (c) and ovule (d).

Figure S2. MET3 reporters and expression

(a) Representation of *pMET3:H2B-GUS*, *pMET3:H2B-tdTomato* and *pMET3:H2B-RFP*. (b) *pMET3:H2B-*

GUS expression observed in 1 DAP and 3 DAP seeds. (c) *pMET3:H2B-RFP* expression observed in 1 DAP

and 3 DAP seeds. Scale bars represent 50 μm. (d) Snapshot of expression data from LCM dissected

seeds from http://seedgenenetwork.net/plotprobe?name=254720_at (Belmonte et al. 2013).

Figure S3. MET3 regulation by Polycomb group

(a) *MET3* expression from an RNA-seq experiment in inflorescence, root, shoot and siliques from wild-

type plants and *clf28* mutant plants, data from Liu *et al* 2016 (Liu et al. 2016). (b) *MET3* expression

from transcriptomic data obtained from 3 DAP seeds using MicroArray comparing wild-type to *fis2-1*

endosperm, data from Weinhofer *et al* 2010 (Weinhofer et al. 2010). (c) Snapshot of H3K27me3 ChIP

data from several tissues including endosperm (colored in blue), flowers (colored in yellow) and

various sporophytic (colored in green). Green line represent the presence of ChIP peaks of H3K27me3.

Data from ReMap (Chèneby et al. 2020).

Figure S4. Characterization of MET3 mutants

(a) Representation of the *MET3* locus. Blue triangles are representing the T-DNA insertion site of both

met3-3 (GABI_404F04) and *met3-4* (GABI_659H03) (b) *MET3* expression measured by RT-qPCR in Col-

0 and *met3* mutants seeds at 3DAP. The histogram displays the mean, and each dot represents a

biological replicate. *ACT2* is used as a normalizer. 2 stars indicate a p value <0.01.

Figure S5. MET3 mutants do not show developmental defect

(a) Percentage of seed abortion in Col-0 and *met3* mutants (*met3-3* and *met3-4*) siliques. (b) met3

mutants transmission in self progeny. +/+ represent wild-type, +/- heterozygotes and -/- homozygotes

plants for the mutation. (c) Seed size measurement for Col-0 and *met3* mutants dry seeds. No

significant differences observed. (d) Pictures of Col-0 and *met3* mutants dry seeds. The scale bars

represent 1 mm.

Figure S6. Transcriptome of met3 mutant seeds at 3DAP

(a-b) Volcano plot depicting one differentially expressed gene comparing Col-0 to *met3-3* (a) and

met3-4 (b) at 3DAP.

Figure 1. *MET3* expression pattern

Figure 2. *MET3* is biallelically expressed with a paternal bias

Figure 3. *MET3* expression is controlled by *FIE* and *MSI1* Polycomb proteins

Figure 4. *MET3* does not influence *fie* phenotype

Figure 5. Transcriptome of *met3* and *met3fie* mutant seeds

a

Klepikova eFP (RNA-Seq data): AT4G13610 / MEE57

Figure S1. *MET3* is only expressed in seeds

Figure S2. *MET3* reporters and expression

Figure S3. *MET3* regulation by Polycomb group

a

"*met3-4*" (GABI659H03)   "*met3-3*" (GABI404F04)

MET3 locus

1000   2000   3000   4000   5000   6000

b

Figure S4. Characterization of *MET3* mutants

Figure S5. *MET3* mutant do not show developmental defect

Figure S6. Transcriptome of *met3* mutant seeds at 3DAP