Robust chronic convulsive seizures, high frequency oscillations, and human seizure onset patterns in an intrahippocampal kainic acid model in mice

Christos Panagiotis Lisgaras[1, 2] and Helen E. Scharfman*[1, 2]

[1]Departments of Child & Adolescent Psychiatry, Neuroscience & Physiology, and Psychiatry, and the Neuroscience Institute
New York University Langone Health
550 First Ave.
New York, NY 10016

[2]Center for Dementia Research
The Nathan Kline Institute for Psychiatric Research
New York State Office of Mental Health
140 Old Orangeburg Road, Bldg. 35
Orangeburg, NY 10962

Conflicts of interest: None

*Corresponding author:
Helen E. Scharfman
E-Mail: hscharfman@nki.rfmh.org
The Nathan Kline Institute
Center for Dementia Research
140 Old Orangeburg Rd. Bldg. 35
Orangeburg, NY 10962
Phone: 845-398-5427
Fax: 845-398-5422

Abbreviations:
IHKA = Intrahippocampal kainic acid
KA = Kainic acid
TLE = Temporal lobe epilepsy
LVF = Low-voltage fast
HYP = Hypersynchronous
HFOs = High frequency oscillations
SE = Status Epilepticus
vEEG = video electroencephalography

47 **HIGHLIGHTS**
48

49 • **Our implementation of the IHKA model led to robust chronic spontaneous**
50 **convulsive seizures in mice**
51

52 • **Convulsive seizures were synchronized in both hippocampi and two**
53 **cortical sites**
54

55 • **Seizure frequency increased from 2-4 wks to 10-12 wks in 50% of mice**
56 **and declined in others**
57

58 • **Convulsive seizures fit LVF and HYP types found in human temporal lobe**
59 **epilepsy**
60

61 • **HFOs (>250 Hz) were common, at >1 location, and were both ictal and**
62 **interictal**
63
64
65

ABSTRACT

Intrahippocampal kainic acid (IHKA) has been widely implemented to simulate temporal lobe epilepsy (TLE), but evidence of robust seizures is usually limited. To resolve this ambiguity, we slightly modified previous methods and employed continuous wideband video-EEG monitoring from 4 recording sites to best detect and characterize chronic epilepsy outcomes in both male and female mice. We found many more convulsive seizures than most studies have reported. Mortality was low. Analysis of convulsive seizures at 2-4 and 10-12 wks post-IHKA showed a robust frequency (2-4 per day on average) and duration (typically 20-30 sec) at each time. Comparison of the two timepoints showed that seizure burden became more severe in approximately 50% of the animals. We show that almost all convulsive seizures could be characterized as either low-voltage fast or hypersynchronous onset seizures, which has not been reported in a mouse model of epilepsy and is important because these seizure types are found in humans. In addition, we report that high frequency oscillations (>250 Hz) occur, resembling findings from IHKA in rats and TLE patients. Pathology in the hippocampus at the site of IHKA injection was similar to mesial temporal lobe sclerosis and reduced contralaterally. In summary, our methods produce a model of TLE in mice with robust convulsive seizures, and there is variable progression. HFOs are robust also, and seizures have onset patterns and pathology like human TLE.

SIGNIFICANCE STATEMENT

Although the IHKA model has been widely used in mice for epilepsy research, there is variation in outcomes, with many studies showing few robust seizures long-term, especially convulsive seizures. We present an implementation of the IHKA model with frequent convulsive seizures that are robust, meaning they are >10 sec and associated with complex high frequency rhythmic activity recorded from 2 hippocampal and 2 cortical sites. Seizure onset patterns usually matched the low-voltage fast and hypersynchronous seizures in TLE. Importantly, there is low mortality, and both sexes can be used. We believe our results will advance the ability to use the IHKA model of TLE in mice. The results also have important implications for our understanding of HFOs, progression, and other topics of broad interest to the epilepsy research community. Finally, the results have implications for preclinical drug screening because seizure frequency increased in approximately half of the mice after a 6 wk interval, suggesting that the typical 2 wk period for monitoring seizure frequency is insufficient.

INTRODUCTION

Temporal lobe epilepsy (TLE) is the most common type of focal epilepsy in adults (Thijs et al., 2019), with up to 30% of the patients achieving poor seizure control by currently available anti-seizure drugs (ASDs; Kwan et al., 2011; Kapur et al., 2019). Moreover, several comorbidities challenge the quality of life of patients and their families (Keezer et al., 2016; Ravizza et al., 2017). As a result, it is important to facilitate research to identify new therapeutic targets, and to that end, animal models of epilepsy have played a key role (Holmes, 2007; Löscher, 2017; Pitkänen, 2017). Kainic acid (KA) has been used for decades in rats (Ben-Ari and Lagowska, 1978; Nadler et al., 1978; Schwarcz et al., 1978; Nadler, 1981; Ben-Ari and Cossart, 2000; Lévesque and Avoli, 2013), where it can induce status epilepticus (SE), a pattern of neuronal loss similar to human TLE (mesial temporal sclerosis; MTS; (Houser, 1999; Scharfman, 2007; Blümcke et al., 2012; Thom, 2014), and spontaneous recurrent seizures (epilepsy), most of which are convulsive (Williams et al., 2007; Lévesque and Avoli, 2013).

Although useful as a rat model of TLE, there often was mortality during or shortly after SE. Mortality was decreased by the use of anticonvulsants 1-2 hrs after SE (Scharfman et al., 2000; Scharfman et al., 2002) or by the use of repetitive low dosing (Meier and Dudek, 1996). However, there has been a major shift away from use of KA in rats to the use of KA in mice, a result of the powerful methods that were tailored for mice and can be used in epilepsy research.

Initial studies of mice often injected KA systemically. Unfortunately, initial studies in mice showed high mortality after systemic injection (Schauwecker and Steward, 1997). Use of anticonvulsants before KA led to reduced mortality (Iyengar et al., 2015) but the degree of hippocampal neuronal loss was sometimes weak even if electrographic SE lasted for several hours (Iyengar et al., 2015). In some studies, neuronal loss was greater (VonDran et al., 2014) but studies of long-term consequences showed that there were few spontaneous seizures (McKhann et al., 2003). Investigators tried to inject KA into the hippocampus instead and found pathology like MTS (Bouilleret et al., 1999). In addition, there was robust GC dispersion (GCD) which is also found in human TLE

(Houser, 1992). The pathology following IHKA was useful because it simulated the
cases of TLE where there is unilateral hippocampal sclerosis. In contrast, prior studies
of systemic KA in rats produced severe pathology in both hippocampi. Another benefit
of IHKA was low mortality (Bouilleret et al., 1999; Riban et al., 2002).
However, chronic convulsive seizures in the wks and months after IHKA in mice
were not discussed in detail (Bouilleret et al., 1999; Riban et al., 2002) so the frequency
and severity were not clear. EEG recordings from the site of IHKA injection showed that
most frequent epileptiform abnormalities in mice were short-lasting non-convulsive
episodes (Bouilleret et al., 1999; Maroso et al., 2011). Brief (3-7 sec) abnormalities were
reported by others where they could be as frequent as 100 per hr (Kim et al., 2018;
Sandau et al., 2019; Lai et al., 2020). Sometimes the EEG examples that were
published suggested trains of spikes or epileptiform activity occurred in the wks after
IHKA, but seizures were either rare, not shown, or the evidence that they occurred was
not strong (e.g., Kiasalari et al., 2016; Zhu et al., 2016; Runtz et al., 2018; Bielefeld et
al., 2019; Li et al., 2020). One study that described methods for IHKA in mice concluded
that IHKA did not lead to epilepsy (Bielefeld et al., 2017).
In addition to these problems, there have been other questions about the IHKA
model in mice. For example, one lead is often used for the EEG and it is placed where
KA was injected, without recording from other brain areas. This may lead to difficulty
assessing how much the seizures spread beyond the hippocampus and mistaken
interpretation that seizures are focal. How many seizures were convulsive and how
many were non-convulsive is not always clear. In addition, many studies occurred
before the mandate at the National Institutes of Health (NIH) to study both sexes
(Clayton and Collins, 2014), and this mandate is not in place outside the U.S. Therefore,
most published data have used males.
To address these issues, we modified the IHKA procedure to produce frequent
convulsive seizures while maintaining low mortality. We report the characteristics of
these seizures that make them robust, such as multiple seizures per day or wk, severity
in that most fit Racine scale 4-5, long duration (up to 100 sec), and postictal depression,
a hallmark of convulsive seizures in TLE (So and Blume, 2010). We also report data for
non-convulsive seizures and both sexes.
We also addressed other questions about IHKA in mice. For example, despite the
report that post-IHKA epileptiform activity may change with time (Riban et al., 2002;
Häussler et al., 2016), data about chronic seizures over time is limited, especially
beyond 2 months post-IHKA (Henshall, 2017). Two studies mentioned in the text that
convulsive seizures occurred many months after IHKA but data were not shown
(Bouilleret et al., 1999; Bui et al., 2018). In contrast, quantified data from rats after
systemic KA showed that spontaneous seizures increased over the first 6 months after
IHKA injection from ~9 per wk up to 50 per wk (Rattka et al., 2013). Therefore, we
quantified seizures occurring 2-4 and 10-12 wks following IHKA injection. Remarkably,
we show that approximately half of the mice exhibited a rise in seizure frequency
whereas the other half declined.
Another question we addressed is the degree seizure types found in human TLE
are also observed in the IHKA model. The seizures in humans have been characterized
according to the pattern of the EEG during the initial phase of the seizure, called the
seizure onset pattern. From a translational perspective, the identification of different

seizure onset patterns is important because the type of seizure onset appears to correlate with the size of the seizure onset zone (Avoli et al., 2016) and may predict post-surgical outcomes (Zaher et al., 2020). The different seizure onset patterns are still being debated (Velasco et al., 2000; Perucca et al., 2014; Gnatkovsky et al., 2019; Saggio et al., 2020), but there appears to be a consensus that at least two types exist: low-voltage fast (LVF) or hypersychronous (HYP) seizures (Lévesque et al., 2012; Gnatkovsky et al., 2019). Although LVF and HYP seizures have been identified in IHKA-treated rats (Bragin et al., 2005), pilocarpine-treated rats (Behr et al., 2017), 4-aminopyridine or picrotoxin-treated rats (Salami et al., 2015) and slices of mice (Shiri et al., 2016), their presence in IHKA-treated mice has been unclear. This is an important gap because presently IHKA in mice is very common in studies of TLE. We show that almost all seizures in our IHKA-treated mice fit either the LVF or HYP classification.

In humans with TLE, high frequency oscillations (HFOs) are of great interest because they appear at seizure onset and may mark the best area for removal in surgical resection for intractable TLE (Zijlmans et al., 2012). Although observed in IHKA-treated rats (Bragin et al., 1999a), HFOs have not been described in epileptic mice to our knowledge. Furthermore, an important question is whether HFOs are confined to the focus, which is assumed to be the KA injection site in the IHKA model (Bouilleret et al., 1999; Riban et al., 2002). In addition, prior studies using one recording electrode at the injection site limited the assessment of seizure focus. We examined when HFOs occur, and where they occur.

In summary, we employed a slightly different approach to induce SE using IHKA and used continuous (24 hrs per day, 7 days per wk) wideband video-EEG (vEEG) monitoring from 4 recording electrodes at 2 different timepoints after IHKA. We asked several questions about our IHKA treated mice: 1) Were there frequent convulsive seizures? Did they generalize? Were non-convulsive seizures present? 2) Were seizures characterized by an onset pattern similar to LVF or HYP seizures? 3) Did seizures progress or remit? 4) Were there HFOs, and if so, where and when? Did HFOs mark a focus, and where was the focus? In addition, we examined hippocampal pathology to determine if it was similar to past reports of the IHKA model, i.e., MTS (Bouilleret et al., 1999). Finally, we made observations in both sexes to determine if there were sex differences.

MATERIALS AND METHODS

I. Animals, breeding, genotyping and animal care

The number of animals used for the present study is summarized in **Table 1**. All experimental procedures were performed in accordance with the NIH guidelines and approved by the Institutional Animal Care and Use Committee at the Nathan Kline Institute.

Amigo2-Cre transgenic mice were kindly provided by Dr. Steven Siegelbaum at Columbia University (Hitti and Siegelbaum, 2014). Hemizygous *Amigo2*-Cre males were bred to C57BL/6 females (Stock number 027, Charles River Laboratories). *Amigo2*-Cre+/- and *Amigo2*-Cre-/- adult males and females were used for experiments in anticipation of future closed-loop seizure intervention studies using this transgenic

mouse line. Note that there was no effect of genotype on acute and chronic outcomes
as described in more detail in **Table 2**.
Prior to IHKA injection, mice were housed 2-4 per cage in standard laboratory
cages and after IHKA injection they were housed 1 per cage. Animals were handled
daily by the experimenter when they were single housed to reduce behavioral stress
related to the lack of social housing (Bernard, 2019; Manouze et al., 2019). Cages were
filled with corn cob bedding and there was a 12 hr light:dark cycle (7:00 a.m. lights on,
7:00 p.m. lights off). Red plastic mini igloos (W.F. Fisher) were placed at the base of the
cage to provide a location that was partially hidden for the mice. This was done to
reduce behavioral stress. Food (Purina 5001, W.F. Fisher) and water was provided ad
libitum.
All breeding pairs were fed Purina 5008 rodent chow (W.F. Fisher) and provided
with one 2''x 2'' nestlet (W.F. Fisher). Mice were weaned on postnatal day 23-30, and
after that time the chow was Purina 5001 (W.F. Fisher).
Genotyping was done using tail samples collected at approximately 23-30 days of
age. Genotyping was performed by the Mouse Genotyping Core Laboratory at New
York University Langone Medical Center.
Although this study used *Amigo2*-Cre mice we do not think the presence of Cre
recombinase in cells expressing *Amigo2* influenced the results. One reason is that our
results from *Amigo2*-Cre+/- and *Amigo2*-Cre-/- mice were similar in several acute
(during SE) and chronic (during chronic seizures) outcome measures as described in
more detail in **Table 2**. In addition, *Amigo2* has been localized only to CA2 pyramidal
cells and hilar neurons (Hitti and Siegelbaum, 2014), which is a small fraction of
neurons in the brain. However, it is notable that in studies of others, *DLX*-Cre+/-mice did
appear to have more seizures than *DLX*-Cre-/- mice despite the absence of viral
injection to experimentally manipulate cells expressing *DLX* (Kim et al., 2013). Other
laboratories have used Cre+/- lines without viral expression in their research, however,
and it has not been shown that hemizygous Cre on its own has major effects on
endpoints we investigated, e.g., seizures after SE.
278
II. Kainic acid (KA) injection
A. KA preparation
KA monohydrate (#0222, Tocris Bioscience) was dissolved in 0.9% sterile saline
and pH was adjusted to 7.4 with 20-30 µl 1 N NaOH (#SS266-1, Fisher Scientific)
according to Tocris Bioscience's guidelines. The final concentration of the stock solution
was 20 mM and it is similar to the concertation used by other investigators (Bui et al.,
2018; Zeidler et al., 2018; Li et al., 2020). The stock solution was then sonicated for 1 hr
to ensure good solubility as indicated in the manufacturer's recommendations. After
sonication, the solution was aliquoted in 0.5 ml portions and kept at -80°C for a
maximum of 1 month. On the day of IHKA injection, an aliquot was allowed to come to
room temperature, and the solution was sonicated for 30 min before the start of the
surgical procedure. The aliquot was discarded after use and a new aliquot was used on
the day of every experiment. Using this approach, we did not notice any precipitation at
any step of KA preparation.
293
294

B. Stereotaxic injection of KA

At approximately 8 wks of age, mice were injected with KA and then implanted with electrodes 10 days or 9 wks later (see methods for KA injection in the next paragraph and implantation in section IV), which was 4-7 days before starting vEEG monitoring. Although implantation at 10 days after IHKA may have altered epileptogenesis, animals exhibited spontaneous seizures in their home cage by 10 days after IHKA, so epilepsy had already developed. Furthermore, in our experience, implantation reduces seizures rather than increasing them (Jain et al., 2019), so it is likely that our results underestimate (rather than overestimate) seizures. This is important because our goal was to show robust seizures rather than few, equivocal seizures. Note that implantation prior to IHKA injection would have potentially allowed detailed recordings of SE, but that was not the major focus of this study. Moreover, prior implantation can reduce seizures as mentioned above, so it was avoided.

To begin the procedure, the 8 wk-old mice were brought to the laboratory for acclimation to the location where KA would be injected. Acclimation typically included two 5 min-long sessions per day for the 2 days before IHKA injection. In each session, the investigator who would be injecting KA conducted the acclimation. For acclimation, the mouse was removed from the cage using gloves and the mouse was allowed to walk on the part of the lab coat covering the lower forearm. One M&M (chocolate-coated peanut) was used as a food reward.

KA was injected between 8 a.m.-1 p.m. Mice were anesthetized with 3% isoflurane (Aerrane, Henry Schein) for 2 min in a rectangular transparent plexiglass box (induction chamber) and then transferred to a stereotaxic apparatus (Model #502063, World Precision Instruments). Anesthesia was then lowered to 1-2%. Mice were frequently monitored to confirm there was adequate respiration and there was no reflex in response to a toe pinch. Mice were placed on top of a homeothermic blanket (Model #50-7220F, Harvard Apparatus) and body temperature was maintained at 37°C by feedback from a lubricated probe inserted into the rectum. Eye ointment was applied to prevent dehydration (Artificial tears, Pivetal). The scalp was shaved and swabbed with Betadine (Purdue Products) using sterile cotton-tipped applicators (Puritan) followed by 70% ethanol. A midline incision exposing the skull surface was made with a sterile scalpel and the skull was cleaned with sterile saline.

One burr hole was drilled using a drill bit (Model #514552, 60 mm, Stoelting) mounted to a surgical drill (Model C300, Grobert) above the left hippocampus. Stereotaxic coordinates for the burr hole were (-2 mm anterior-posterior (A-P) to Bregma, - 1.25 mm medio-lateral (M-L)). Care was taken to leave the dura mater intact by regularly monitoring it with a stereoscope during drilling (Stemi SV6, Zeiss). The drilling area was regularly hydrated with 0.9% sterile saline solution and drilling was done in steps to avoid depressing the underlying tissue during drilling. This approach was followed because we measured the dorso-ventral (D-V) zero point from dural surface, so we wanted the measurement of the dural surface to be as accurate as possible.

Next, a 0.5 ml Hamilton syringe (Model 7001, Hamilton) was lowered from brain surface 1.6 mm into the left dorsal hippocampus (-1.6 mm D-V) and 70-100 nL of 20 mM KA dissolved in 0.9% sterile saline was manually injected over approximately 5 min. A range of IHKA volumes were used instead of a fixed volume because in pilot studies

all doses were successful in triggering robust convulsive SE with minimal
(approximately 20%) acute (during SE) mortality (see the legend for **Table 2** for a
detailed explanation). The speed was controlled by depressing the syringe 1 unit (10
nL) every 15 sec. The needle remained in place for an additional 3 min and was then
slowly removed to prevent backflow of the injected solution. The incision was quickly
closed using tissue adhesive (Vetbond, 3M). Mice were transferred to a clean cage on
an autoclaved paper towel (without bedding) and placed over a 37°C heating pad. After
approximately 3-4 hrs of monitoring the animal, bedding was used instead of the towel.

III. Behavioral monitoring of IHKA-induced status epilepticus (SE)
        Behavior during IHKA-induced SE was visually monitored and seizure severity
during SE was scored using stages 1-5 of the Racine scale (Racine, 1972) and stages
6-7 using the Pinel and Ronver scale (Pinel and Rovner, 1978) because all stages were
observed in our experiments. Using the scale with stages 1-7, stages 1-2 were
considered non-convulsive and stages 3 and higher were considered convulsive. Stage
1 was accompanied by intense mastication and facial movements such as blinking,
repetitive ear movements, a sudden change in the behavior to a frozen stance. Stage 2
was head nodding. Stage 3 was unilateral forelimb clonus, stage 4 was bilateral
forelimb clonus with rearing, and stage 5 was a stage 4 seizure with loss of postural
tone. Stage 6 seizures were accompanied by jumping or repetitive falling and stage 7
included vigorous jumping and running around the cage. Notably, all IHKA-injected
animals experienced vigorous convulsive SE meaning that there were several stage 5
seizures. After the first stage 5 seizure, the cage was removed from the heating pad to
prevent hyperthermia. The body temperature was regularly (approximately every 15
min) monitored with an infrared thermometer (Model #800048, Sper Scientific) to ensure
that body temperature was within the physiological range (36-37°C).
        Mice were visually monitored until normal behavior (defined as exploration and/or
grooming) resumed. After 3-4 hrs of observation following IHKA injection, mice resumed
normal behavior, and they were returned to their home cages with normal bedding and
no paper towel. The cages were kept overnight in the same location. After 1 day, mice
were moved close to the EEG equipment so that they would acclimate to that area.
        Starting with the day after IHKA injection, animals were handled for approximately 5
min daily as described above and their body weight was measured for the next 7 days
once daily. In 3 IHKA-injected mice (1 male, 2 female), body weight declined transiently
by 10-18% so they were administered 3 ml of warm (36-37°C) sterile saline solution s.c.
which did not affect chronic outcome measures as described in **Table 2**. The solution
was administered once and within 24 hrs post-IHKA. These and other aspects of the
IHKA injection procedure and subsequent 7 days are listed in **Table 2.**

IV. EEG Recording
A. Surgical implantation of EEG electrodes
        Two wks after IHKA injection, animals were implanted with 6 subdural electrodes (4
recording, 1 reference, 1 ground; see below for details). As mentioned above, this time
was chosen because animals were already exhibiting convulsive seizures in their
homecage, so the period of epileptogenesis appeared to be over.

Mice were anesthetized with 3% isoflurane (Aerrane, Henry Schein) for 2 min and then transferred to the stereotaxic apparatus (Model #502063, World Precision Instruments). Anesthesia was then lowered to 1-2% and monitored the same way as described above for IHKA injection. Note that vendor information is already specified above for many of the items used below and where not the vendor information is provided. Mice were placed on top of a homeothermic blanket, eye ointment was applied, and the scalp was shaved and cleaned following the same procedures described above. A midline incision was made with a sterile scalpel and the skull was cleaned with sterile saline. Two burr holes were drilled over the left and right hippocampus (-2.5 mm A-P, ± 1.75 mm M-L), slightly posterior to the septotemporal level where KA was injected. Two additional burr holes were drilled anterior to the IHKA injection site (-0.5 mm A-P, ± 1.5 mm M-L) to serve as the cortical recording leads. Subdural screw electrodes (0.10'' length stainless steel) were placed in the burr holes and secured using dental cement (Jet Set-4 Denture Repair, Lang Dental). Two additional burr holes were drilled above the cerebellar region to serve as reference (-5.7 mm A-P, +1.25 mm M-L) and ground (-5.7 mm A-P, -1.25 mm M-L). The reference was placed contralateral to the IHKA injection site and the ground was ipsilateral. The subdural screw electrodes were attached to an 8-pin connector (#ED83100-ND, Digikey) which was secured to the skull with dental cement.

B. Continuous wide-band video-EEG monitoring

One-day after EEG surgery and until the vEEG recording started, mice were housed in the room where the vEEG equipment was located so that they could acclimate to the recording environment. For recording, mice were placed into a 21 cm x 19 cm square transparent plexiglass cage which had access to food and water and had no cage lid. Food was placed on the bottom of the cage and a water bottle was attached to one of the walls of the cage. A pre-amplifier (Pinnacle Technologies) was connected to the 8-pin connector and then to a commutator (Pinnacle Technologies) which allowed free movement of the mouse in the entire cage. EEG signals were amplified 10 times and recorded at 2 kHz sampling rate using a bandpass filter (0.5-500 Hz) and Sirenia Acquisition System (Pinnacle Technologies). High frame rate video (30 fs) was recorded simultaneously using an infrared LED camera (Model #AP-DCS100W, Apex CCTV).

C. Quantification of video-EEG
1. Seizure analyses

Seizures were detected from vEEG by a blinded investigator (CPL) using replay of the EEG and video records. The left hippocampal lead was chosen for most measurements because it was the site of the IHKA injection. However, seizures were typically present in all leads. Therefore, they demonstrated robust synchronization and generalization.

The electrographic component of a seizure, whether convulsive or non-convulsive, was defined by a sudden change in amplitude >2x of the standard deviation (SD) of the baseline mean. The threshold, 2x the SD of the baseline mean, was chosen because it was adequate to differentiate seizures from normal EEG. The baseline mean was calculated from the 30 sec prior to the event that was being considered to be a possible seizure. During this 30 sec baseline, care was taken to choose a time when the

behavioral state was not interrupted by large artifacts or abrupt behavioral state changes. An electrographic seizure was also defined by high frequency rhythmic activity (>5 Hz) which consisted of an abnormal pattern (large amplitude spikes and clusters of spikes lasting for at least 10 sec). Ten seconds was chosen because seizures in TLE typically last at least 10 sec and often are 20-60 sec (Balish et al., 1991). In addition, most seizures recorded in a standard Epilepsy Monitoring Unit typically last more than 10 sec (Jenssen et al., 2006). We also have found that the majority of seizures in epileptic mice after pilocarpine-induced SE in our laboratory were at least 10 sec and usually 20-60 sec (Iyengar et al., 2015; Botterill et al., 2019; Jain et al., 2019). **Seizure onset** was defined as the time when the baseline of the left hippocampal lead exceeded 2x the SD of the baseline mean. The **end of a seizure** (seizure termination) was defined as the time when high amplitude activity declined to <2x the SD of the baseline mean. **Seizure duration** was calculated by subtracting the time of seizure termination (the end of the seizure as defined above) from the time of seizure onset. **Seizure burden** was quantified as the number of days that animals sustained chronic seizures (either non-convulsive or convulsive) divided by the total number of days that continuous vEEG was performed at 2-4 and 10-12 wks (i.e., 14 days). For each seizure, the light cycle was noted as lights on (7:00 a.m.) or off (7:00 p.m.) and the time of the day with a.m. and p.m. The behavioral state when the seizure began was determined as either wakefulness or sleep based on video and the EEG for a period of 30 sec before seizure onset. The **awake** state included awake exploration and awake immobility. **Awake exploration** was defined by walking or other movement around the cage and the predominance of theta rhythm (4-12 Hz in the hippocampal leads) with eyes open (when the mouse was facing the camera). **Awake immobility** was not accompanied by exploration, but eyes were open. **Sleep** was defined either as non-rapid eye movement (NREM) or rapid eye movement (REM) sleep. **NREM** was dominated by slow wave activity in the delta (<5 Hz) frequency range and eyes were closed with no movement of the body. **REM** sleep was defined by theta rhythm (4-12 Hz) and eyes closed with no movement of the body. For defining sleep stages, we relied on leads contralateral to the IHKA injection site because of previous reports about loss of theta rhythm predominantly adjacent to the IHKA injection site (Riban et al., 2002; Arabadzisz et al., 2005).

    For quantification, the percent of seizures in each category (on or off, a.m. or p.m.; awake or asleep) was calculated by dividing the number of seizures in a category by the total number of seizures. The total was either the total number per animal or the total number of all seizures in all animals.

    Seizure onset patterns were defined LVF, HYP (as described above) or 'unclear' which was a seizure that had an onset pattern that was not possible to classify as LVF or HYP. LVF seizures started with a sentinel spike (amplitude >2x SD above the baseline mean) followed by brief suppression of the background EEG whereas HYP seizures began with high amplitude (>2x SD above the baseline mean) spiking at a frequency that was much greater than interictal spikes. These definitions are similar to those described in other animal models of epilepsy (Bragin et al., 2005; Behr et al., 2017) and human patients (Velasco et al., 2000).

2. High frequency oscillations (HFOs)

HFOs were defined as oscillations >250 Hz consistent with past studies in humans (Staba et al., 2004) and animals (Bragin et al., 1999a). To sample HFOs, NREM sleep was selected because prior studies suggested that HFOs occur mainly during slow waves and sharp waves occurring in NREM sleep (Staba et al., 2004; Bagshaw et al., 2009). For each animal, an interictal period was chosen, and 10 min was sampled at least 1 hr after the last seizure and at least 1 hr before the next seizure. All 4 channels of wideband interictal EEG were visually inspected for the presence of HFOs and a 10 min NREM sleep epoch was selected. This was possible because high frequency activity was visually discernable from the ongoing background EEG activity. We then applied an automated approach to detect HFOs using the RippleLab application written in MATLAB (Navarrete et al., 2016) and the algorithm developed by Staba and colleagues (Staba et al., 2002). Automated detection was followed by visual inspection of putative HFOs and inclusion/exclusion criteria were used to either include or exclude the putative events as HFOs. These criteria are described below. In brief, each channel was band-pass filtered between 250-500 Hz using a finite impulse response (FIR) digital filter and the root mean square (RMS) amplitude of the filtered signal was quantified. Successive RMS amplitudes >5x SDs above the mean amplitude of the RMS signal calculated for the entire 10 min epoch (during NREM) that lasted >6 msec in duration were defined as putative HFOs. Putative HFOs that did not have at least 4 peaks in the rectified band-passed signal >3x SDs greater than the mean of the baseline signal were excluded. We then reviewed each putative HFO to exclude artifactual events associated with movement or other sources of noise (Bénar et al., 2010; Menendez de la Prida et al., 2015; Amiri et al., 2016). To better appreciate HFO power in time, we applied time-frequency analyses to visualize HFO power. To that end, we used the time-frequency function which is part of RippleLab (Navarrete et al., 2016) and applied a 64-point window to analyze HFO power in the 250-500 Hz frequency domain.

Whether HFOs were interictal or ictal was visually determined by inspecting all 4 channels. **Interictal HFOs** were determined from the EEG at least 1 hr before or 1 hr after a seizure. **Ictal HFOs** were visually defined as HFOs that occurred at seizure onset and during seizures by inspecting all 4 channels for the presence of HFOs >250 Hz.

V. Anatomy
A. Perfusion-fixation and sectioning

Mice were deeply anesthetized with isoflurane and they were injected with an overdose of urethane (2.5 g/kg, i.p.; Sigma Aldrich). During surgical anesthesia, defined as the time when mice did not respond to a toe pinch with a withdrawal reflex, the abdominal cavity was opened with a scalpel and a 25 gauge butterfly-style needle (Model #J454D, Jorvet) was inserted into the heart. When the needle was clamped in place with a hemostat, mice were transcardially perfused with 10 ml of room temperature (25°C) saline, followed by 30 ml of cold 4% paraformaldehyde (#1920, Electron Microscopy Systems) in 0.1 M phosphate buffer (PB; pH=7.4) using a peristaltic pump at a rate of 10-12 ml/min (Minipuls2, Gilson). The brain was quickly removed and stored overnight in 4% paraformaldehyde in 0.1 M PB. An incision was made laterally on the right side of the brain to mark the side that was contralateral to the

IHKA injection. On the next day, the brains were sectioned (50 µm thickness) in the coronal plane using a vibratome (Vibratome 3000, Ted Pella) in Tris buffer. Sections were stored in 24 or 48 well tissue culture plates containing cryoprotectant solution (25% glycerol, 30% ethylene glycol, 45% 0.1 M PB, pH 6.7) at 4°C until use.

B. Nissl staining

For Nissl staining, sections were mounted on 3% gelatin-coated slides and left to dry overnight at room temperature. Then slides were dehydrated in increasing concentrations of ethanol (70%, 95%, 100%, 100%) for 2.5 min each, cleared in Xylene (Sigma Aldrich) for 10 min, and then dehydrated again (70%, 95%, 100%, 100%) followed by hydration in double-distilled (dd) H_2O for 30 sec. Then sections were stained with 0.25% cresyl violet (Sigma Aldrich) dissolved in ddH_2O for 1.5 min followed by 30 sec in 4% acetic acid solution dissolved in ddH_2O. Then sections were dehydrated (70%, 95%, 100%, 100%), cleared in Xylene for 4 min, and cover-slipped with Permount (Electron Microscopy Systems).

C. Quantification

To estimate neuronal loss and granule cell dispersion in dorsal hippocampus, one coronal section per animal was selected slightly anterior to both the IHKA or saline injection site. This site also allowed us to assess tissue integrity near the injection track and subdural electrode near the injection site. This dorsal section corresponded to approximately -1.94 to -2.06 mm posterior to Bregma in a common mouse stereotaxic atlas (Figures 47-48; Franklin and Paxinos, 1997). We chose this coronal level to conduct our anatomical procedures based on previous reports suggesting that neuronal loss is maximal adjacent to the IHKA injection site (Bouilleret et al., 1999; Riban et al., 2002). In both IHKA- and saline-injected mice, good tissue integrity was observed near the injection site, and the injection track was actually hard to see. This absence of damage by the injection needle rendered anatomical assessments possible at this tissue level. The absence of a visible injection track was probably achieved because we used a syringe and not a permanently implanted (guide) cannula to inject KA or saline. The absence of damage from the EEG recording electrode was because we used subdural screw electrodes instead of depth electrodes for our vEEG monitoring. All these approaches significantly minimized tissue damage near the injection site, providing an opportunity to assess the injection area with limited technical confounds.

Nissl-stained sections were imaged using a microscope (Model BX51, Olympus of America) and digital camera (Infinity3-6URC) at 2752x2192 pixel resolution. Photographs were taken with Infinity capture software (version 6.5.6). After images were imported to ImageJ software (NIH), we measured the pyramidal cell layer length (PCL length), granule cell layer (GCL) area, and GCL width or thickness. Details are shown in **Supplemental Figure 1**. For PCL length, a line was drawn using the freehand line tool from the border of the PCL in CA3c with the DG hilus to the CA1 border with the subiculum. The CA3c cell layer at the border with the hilus was defined as the point where pyramidal cells became sufficiently close so that there was <1 pyramidal cell body width apart from adjacent PCs. The CA1 cell layer at the border with the subiculum was defined in an analogous manner because the cell layer broadens significantly as the subiculum begins. For GCL area measurements, we traced the

borders of the GCL using the freehand tool. The GCL was distinguished from the hilus
as the point where GC somata had <1 cell width distance between them. For GCL
thickness, we used the straight-line tool to measure the thickest part of the layer. The
line to measure thickness was made perpendicular to the GCL, starting at the border
with the hilus and ending at the border with the inner molecular layer (IML). Two
measurements were made of the widest part of the upper and the lower blade, and then
we averaged the 2 measurements.
576
VI. Statistics
Data are presented as mean ± standard error of the mean (SEM). Statistical
significance was set at $p < 0.05$ and is denoted by asterisks on all graphs. All statistical
analyses were performed using Prism (version 8.4.2, Graphpad). Comparisons of
parametric data of two groups were done using unpaired or paired two-tailed Student's
t-test. To determine if data fit a normal distribution the D'Agostino-Pearson test was
used in Graphpad. To determine if the variance of groups was homogeneous, the
Brown-Forsythe test was used in Graphpad. When data did not fit a normal distribution
or transformation was unable to resolve differences in variance, nonparametric statistics
were used. The non-parametric tests were the Mann-Whitney U-test to compare two
groups and the Kruskal-Wallis ANOVA for multiple groups. For correlation analyses we
used linear regression and the Pearson correlation coefficient (r) in Graphpad.
589
VII. Experimental design and data collection
The present study used *Amigo2*-Cre+/- or *Amigo2*-Cre-/- mice for IHKA and saline
injections. Animals were first acclimated and then KA or saline was injected in the
hippocampus. A cohort of 5 IHKA-injected animals was recorded at 2-4 wks post-IHKA
and the same mice were recorded again at 10-12 wks with continuous wideband vEEG
monitoring. In addition, 3 IHKA-injected animals were recorded only at 10-12 wks.
For anatomy, animals were perfusion-fixed after vEEG, 12 wks post-IHKA (n=8).
Earlier times were also checked to confirm a similar degree of PCL damage (3 days,
n=1; 3 wks, n=1), consistent with the idea that most pyramidal cell loss occurs shortly
after SE (Sutula and Pitkänen, 2002). Saline-treated mice were examined also (3 days,
n=1; 2-4 wks, n=3) and did not show neuronal loss, as described in more detail in the
Results. A list of all animals included in the study is presented in **Table 1**. Only one
animal died in this study, and that was unrelated to the study.
603
VIII. Data collection
Although, EEG analyses and quantification of Nissl-stained sections were done by
an investigator (CPL) who was blinded to the experimental group and sex, data
collection was unblinded because animals were more active after IHKA injection
compared to saline-injected controls.
Data collection was done using Case Report Forms (CRFs), where each CRF was
specific for a procedure such as IHKA injection. Common Data Elements (CDEs) were
compiled as previously described (Harte-Hargrove et al., 2018; Scharfman et al., 2018).
We provide all CDEs we used in each procedure-specific CRF as a supplemental .xls
file.
614

RESULTS

I. IHKA leads to epilepsy with frequent spontaneous convulsive seizures

To determine whether convulsive seizures occurred in the mice that were injected with KA (IHKA-treated mice), mice were monitored by continuous vEEG from 2-4 and 10-12 wks after IHKA-induced convulsive SE (**Figure 1A**). At both the early (2-4 wks) and late (10-12 wks) times, convulsive seizures occurred spontaneously and were typically frequent (multiple seizures per day) and severe (stage 4-5). A representative example of a convulsive seizure is shown in **Figure 1B1** and is expanded in **Figure 1B2**.

Figure 1. Example of a spontaneous convulsive seizure recorded 2-4 wks after IHKA

(A) Experimental timeline of the study. Animals were injected with KA in the left dorsal hippocampus and 10 days later they were implanted with 4 subdural screw electrodes. They were then vEEG monitored 2-4 wks post-IHKA continuously (red arrows, with data from this time shown in B) and then recorded again at 10-12 wks post-IHKA.

(B) Representative example of a chronic convulsive seizure recorded during the 2-4 wk session after IHKA. Four leads were used to record right and left frontal cortices and left and right hippocampal regions. A 5 min-long EEG trace with seizure activity in all 4 leads is shown in B1. The start and the end of the convulsive seizure is indicated by green arrows (Conv SZ on/off). The same seizure is shown in a 2 min-long time window in B2 and further expanded (a, b) in 10 sec epochs to better show the EEG complexity. Inset a of the seizure in B2 shows complex and often rhythmic activity with fast and slow components. Inset b of the seizure in B2 shows prolonged suppression of the background EEG in all 4 leads after the termination of the electrographic manifestations of the seizure.

The EEG corresponding to the seizures reflected robust seizure activity in that there was abnormal high frequency activity lasting 20-60 sec (**Figure 1**). Specifically, there was a sudden increase in amplitude (>2x SD above the mean) of rhythmic (>5 Hz) activity that was followed by a suppression of the EEG (**Figure 1B1, 1B2**) lasting 21.5±2.3 sec. For this measurement, the mean was first calculated for 4 seizures at random for one animal. Then the mean was taken for 5 animals. Insets a (**Figure 1B2a**) and b (**Figure 1B2b**) show EEG traces where the complex rhythmic activity that is typical of a seizure is clear. The first inset shows the EEG during the seizure and the second inset shows the postictal suppression. Although there was some variability between leads, the EEG correlates of a severe seizure and postictal suppression were evident at each recording location. During the part of the EEG that appeared to be a seizure, convulsions typically developed in the middle or later parts of the EEG seizure activity. Thus, the EEG showed evidence of a robust seizure first, and then a convulsion developed before the EEG manifestations of the seizure ended. At the same time as loss of postural tone, or after the convulsive behavior ended, the EEG manifestations of the seizure ended.

Next, we quantified chronic spontaneous convulsive seizures during the 2-4wks post-IHKA (**Figure 2A**) and results are shown in **Figure 2B-H**. The total number of convulsive seizures during the 2 wks is shown in **Figure 2B,** highlighting that every animal sustained multiple convulsive seizures. A breakdown of the number of seizures per day showed some variability between animals, although all animals had multiple convulsive seizures per day for many days in a row (**Figure 2C**). Note that one of the females (76-1) had fewer seizures than other mice (**Figure 2B-D**), but that mouse did have 5 convulsive seizures in 2 wks which is still a clear demonstration of chronic epilepsy.

The mean convulsive seizure frequency was 2.1±0.5 seizures per day (range 0.3-3.7, n=5 mice; **Figure 2D**). Convulsive seizures lasted for 37.9±2.3 sec (range 29.9-43.0) when mean duration was calculated per animal, and when all seizures were pooled, the mean seizure duration was 36.2±0.8 sec (range 19.7-94.5; **Figure 2E**).

Analyses of simultaneous video records revealed that convulsive seizures fit the modified Racine scale with a mean severity score of 4.3±0.4 (range 3-6) when the mean was calculated per animal, and 4.1±1.1 (range 3-7) when all seizures were pooled (**Figure 2F**). Next, we quantified the number of days that animals sustained convulsive seizures in 2-4 wks because this metric gives us an estimate of seizure burden (**Figure 2G**). The mean percent of days spent with seizures (57.0±10.5%, range 27-87%) was not different from the days spent without seizures (43.0±10.5%, range 13-73%; Wilcoxon signed rank test, p=0.62). In order to account for any potential circadian effects on seizures, we calculated the percent of seizures occurring during the 12 hr-long period when lights were on and the 12 hr-long period when lights were off (**Figure 2H1a, 2H1b**), and no significant differences were found (paired t-test, t_{crit} = 1.472, p=0.21). We also did not find any differences when we examined the a.m. or p.m. (12:00 a.m.-12:00 p.m. vs. 12:00 p.m. to 12:00 a.m.; paired t-test, t_{crit} = 0.015, p=0.98; **Figure 2H2a, 2H2b)**. Finally, we examined the times when mice were awake and times they were asleep (awake and sleep are defined in the Methods) and no difference was found (paired t-test, t_{crit} = 2, p=0.11;

Figure 2H3a, 2H3b). Although no significant differences were detected for any of these analyses, some individual animals showed a striking difference in seizures experienced in the time period when lights were on or off, a.m. vs. p.m., or during awake vs. sleep states (**Figure 2H**).

Figure 2. Quantification of chronic spontaneous convulsive seizures 2-4 wks post-IHKA

(A) Experimental timeline of the study. The 2-4 wk data were used in this figure (red arrows). For this timepoint, there were 5 mice (males, blue; females, pink; total # seizures=415).

(B) The total number of chronic convulsive seizures recorded 2-4 wks post-IHKA is shown per animal. Note that animals showed frequent convulsive seizures although there was variability. One of the female mice, 76-1, had 5 seizures in 2 wks which is fewer than other animals, but nevertheless is a demonstration of chronic epilepsy.

(C) The number of convulsive seizures per day and per animal is shown for all recording days. The animal ID is shown in the inset.

(D) Convulsive seizure frequency was calculated as a mean number of convulsive seizures per day for each animal.

(E) Convulsive seizure duration was calculated as the mean per animal (left) or the mean of all convulsive seizure durations (right; n=143 seizures).

(F) Convulsive seizure severity was calculated as a mean per animal (left) or the mean of all convulsive seizures (right; n=143 seizures).

(G) Convulsive seizure burden was defined as the percent of days spent with (SZ) or without (No SZ) seizures.

(H) The percent of convulsive seizures is shown, either occurring during the light period or dark period of the light:dark cycle (Lights ON or OFF; H1), a.m. or p.m. (AM, PM; H2), and in awake (AW) or sleep (SL) state (H3). The percentages were calculated as the mean per animal (H1a, H2a, H3a) or the mean of all seizures (H1b, H2b, H3b). There were no significant differences (Wilcoxon signed rank tests; all p>0.05).

In summary, convulsive seizures were robust both qualitatively and quantitively in IHKA-treated mice. Seizures were typically severe and observed with all recording electrodes, indicating that they were generalized. They also showed additional

characterizing that have been discussed in human TLE, including HFOs, which are
discussed further below.

II. Chronic convulsive seizure frequency varies over time

Next, we continued to ask whether our IHKA model was robust. To that end, we
determined whether convulsive seizures persisted at 10-12 wks, 2.5-3.0 months after
IHKA.

Figure 3. Quantification of chronic spontaneous convulsive seizures 10-12 wks post-IHKA

(A) Experimental timeline of the study. The 10-12 wk data were used for this figure (green arrows; total # seizures=415). Animals were recorded 2-4 wks post-IHKA and then again at 10-12 wks post-IHKA (2 male, 2 female). Additionally, 3 animals were recorded at 10-12 wks only (3 females).

(B) The total number of chronic convulsive seizures is shown per animal. Note that all animals showed convulsive seizures at 10 wks suggesting epilepsy persisted.

(C) The number of convulsive seizures per day and per animal is shown for all recording days. The animal ID is shown in the inset.

(D) Convulsive seizure frequency was calculated as the mean number of convulsive seizures per day for each animal.

(E) Convulsive seizure duration was calculated as the mean per animal (left) or the mean for all convulsive seizures (right; n=415 seizures).

(F) Convulsive seizure severity was calculated as the mean per animal (left) or the mean of all convulsive seizures (right; n=415 seizures).

(G) Convulsive seizure burden was defined as the percent of days spent with (SZ) or without (No SZ) convulsive seizures.

(H) The percent of convulsive seizures is shown. either occurring during the light period or dark period of the light:dark cycle (Lights ON or OFF; H1), a.m. or p.m. (AM/PM; H2), and awake (AW) or sleep (SL) state (H3). The percentages were calculated as the mean per animal (H1a, H2a, H3a) or the mean of all seizures (H1b, H2b, H3b). There were no significant differences (Wilcoxon signed rank tests; all p>0.05).

Four out of 5 animals recorded at 2-4 wks were recorded again at 10-12 wks
(**Figure 3A**) along with 3 other IHKA-injected animals (78-3, 80-4, 81-1) that were
recorded at 10-12 wks only. A total of 415 seizures were analyzed and results are
presented in **Figure 3B-H**. When all seizures during the two 2 wk-long periods (2-4 wks
vs. 10-12 wks) were compared, 2 out of 4 animals that were recorded at both timepoints
showed more seizures at 10-12 wks, an increase or progression (defined here as
seizures that worsen with time), and the remaining 2 showed a decrease (see **Figure
3B vs. Figure 2B**). We use the term progression conservatively as we cannot exclude
the possibility that if a third timepoint was used the seizures that had increased from 2-4
to 10-12 wks might decrease eventually or vice-versa.
Similar to the 2-4 wk timepoint, convulsive seizures at the 10-12 wk timepoint were
evident for most recording days (**Figure 3C**). There was a mean frequency of 4.0±1.3
(range 0.2-8.4) seizures per day per animal (**Figure 3D**). Convulsive seizures lasted for
41.2±3.4 sec (range 33.2-56.0) when the mean duration was calculated per animal and
41.3±0.5 sec (range 22.5-74.1) when all seizures were pooled (**Figure 3E**). These
durations were not significantly different from those at 2-4 wks (**Supplemental Figure
2D**, paired t-test, t_{crit} = 2.41, p=0.09). Regarding severity, most seizures were severe
because seizures scores were 4.4±0.3 (range 3-6) when the mean was calculated per
animal and 4.6±0.04 (range 3-6) when all seizures were pooled (**Figure 3F**). These
seizures scores were not significantly different from those at 2-4 wks (Wilcoxon signed
rank test, p=0.62). The mean percent of days with seizures (72.4±13.7%, range 14-
100%) was not significantly different from the mean percent of days without seizures
(27.4±13.6%, range 0-85%; Wilcoxon signed rank test, p=0.10), similar to the 2-4 wk
timepoint.
For the "progressed" animals, seizure burden was 100% and for "non-progressed"
mice it was 46.5±6.5% (**Figure 3G**). When comparing the percent of seizures occurring
during the time that lights were on vs. the time when lights were off, there were no
significant differences (Wilcoxon signed rank test, p=0.46; **Figure 3H1a, 3H1b**). There
also were no significant differences between the percent of seizures during the a.m. vs.
p.m. (Wilcoxon signed rank test, p=0.46; **Figure 3H2a, 3H2b**). Finally, there was no
difference in the percent of seizures occurring in awake vs. sleep states (Wilcoxon
signed rank test, p=0.57; **Figure 3H3a, 3H3b**).
A comparison of the data from 2-4 and 10-12 wks showed no significant differences
in the total number (paired t-test, t_{crit} = 0.61, p=0.58), frequency (paired t-test, t_{crit} = 0.55,
p=0.61), severity (paired t-test, t_{crit} =0.37, p=0.73) or days spent with seizures (unpaired
t-test, t_{crit} = = 0.35, p=0.73). Also, no differences were found in the percent of seizures
occurring during lights on vs. lights off, a.m. vs. p.m. or awake vs. sleep between the 2
timepoints (**Supplemental Figure 3A**). The only difference was found in seizure
duration as seizures at 2-4 wks were shorter than those recorded at 10-12 wks post-
IHKA (paired t-test t_{crit} = = 3.36, p=0.043). Therefore, taking seizure frequency into
account, only 50% of mice "progressed", but if seizure duration is used as a
measurement of progression, the mice did exhibit worsening of their seizures with time.
795
796
797

III. Chronic non-convulsive seizures occur after IHKA and are less frequent than convulsive seizures

Next, we analyzed non-convulsive seizures to determine whether they also are frequent post-IHKA. To that end, the same animals presented before in **Figures 1** and **2** were analyzed as shown in **Figure 4A**. An example of a non-convulsive seizure is shown in **Figure 4B1** and an expanded version of the same seizure is shown in **Figure 4B2.**

Figure 4. A spontaneous non-convulsive seizure recorded 10-12 wks post-IHKA

(A) Experimental timeline shows when the non-convulsive seizure in this figure was recorded, 10-12 wks post-IHKA (red arrows). For non-convulsive seizures, the mice were the same ones used to analyze convulsive seizures.

(B) Representative example of a non-convulsive seizure recorded 10-12 wks post-IHKA. A 5 min-long EEG trace illustrating seizure activity in all 4 leads is shown in B1. The same seizure is depicted in a 2 min-long time window in B2 and further expanded in 10 sec-long epochs to better highlight electrographic activity over time. Inset a shows electrographic activity during the non-convulsive seizure. Inset b shows seizure termination. Note seizure termination was more distinct for some leads and the lead that was most distinct (green arrow in b, LHPC) varied from seizure to seizure.

The electrographic correlate of non-convulsive seizures consisted of a sudden increase in amplitude of rhythmic activity (>5 Hz) in all 4 leads. Insets in **Figure 4** show expanded EEG traces where the electrographic correlate of a non-convulsive seizure can be further appreciated. The non-convulsive seizure was characterized by trains of spikes of variable amplitude which appeared to occur synchronously in

all 4 leads. Interestingly, post-ictal depression was not as pronounced as after a
convulsive seizure (**see Figure 1B2b**).
We next quantified non-convulsive seizures at 2-4 wks (**Figure 5A**) and results are
shown in **Figure 5B-H**. We found that both the total number and frequency of non-
convulsive seizures were less frequent than convulsive seizures at 2-4 wks (paired t-
test, t_{crit}=3.29, p=0.03; paired t-test, t_{crit}=3.32, p=0.02, respectively) with no differences
in mean duration (paired t-test, t_{crit}=1.21, p=0.34); **Supplemental Figure 2D**). More
831

Figure 5. Quantification of chronic spontaneous non-convulsive seizures 2-4 wks post-IHKA
(A) Experimental timeline of the study. The seizures for this figure were recorded 2-4 wks after IHKA (red
arrows; total # seizures=8). The mice were the same as those used for convulsive seizure
measurements.
(B) The total number of chronic non-convulsive seizures during the 2 wk-long recording period is shown
per animal. Blue and pink shades represent males and females, respectively.
(C) The number of non-convulsive seizures per day and per animal is shown for all recording days. The
animal ID is shown in the inset.
(D) Non-convulsive seizure frequency was calculated as the mean number of non-convulsive seizures per
day for each animal.
(E) Non-convulsive seizure duration was calculated as the mean per animal (left) or the mean of all non-
convulsive seizures (right; n=8 seizures).
(F) Non-convulsive seizure severity was calculated as the mean per animal (left) or the mean of all non-
convulsive seizures (right; n=8 seizures).
(G) Non-convulsive seizure burden was defined as the percent of days spent with (SZ) or without (No SZ)
non-convulsive seizures in the 2-4 wk-long recording period.
(H) The percent of non-convulsive seizures is shown, either occurring during the light period or dark period
of the light:dark cycle (Lights ON or OFF; H1), a.m. or p.m. (AM/PM; H2), and awake (AW) or sleep
(SL) state. Percentages are shown for the mean per animal (H1a, H2a, H3a) or the mean of all seizures
(H1b, H2b, H3b). Note that in H2a there are 3 animals but the lines for 2 of the animals overlap since
all of their seizures occurred during AM. In H3a, all seizures in all 3 animals occurred during the awake
state (lines are overlapping). Statistical comparisons: H1a, not significant (Wilcoxon signed rank test;
p>0.05); H2a, not significant (Wilcoxon signed rank test; p>0.05); H3a, not significant (Wilcoxon signed
rank test; p>0.05).

detailed comparisons about lights on vs. off, a.m. vs. p.m. and awake vs. sleep are
shown in **Supplemental Figure 3C and 3D**.
Figure 5C shows the numbers of non-convulsive seizures for the different
recording days, and it is evident that there could be few non-convulsive seizures.
The mean frequency of non-convulsive seizures was 0.1±0.04 seizures per day
(range 0-0.2; n=5 mice; **Figure 5D**) and the duration was 28.8±5.2 sec (range 19.4-
37.5 sec) when the mean was calculated per animal and 28.7±4.0 sec (range 15.0-
44.1 sec) when all seizures were pooled together (**Figure 5E**). The mean severity
score was 1.1±0.1 (range 1-2) per animal and 1.1±0.1 (range 1-2) when all seizures
were pooled (**Figure 5F**). The mean percent of days spent with vs. without seizures
was 8.0±3.8% (range 0-20%) and 92.0±3.8% (range 80-100%) respectively (**Figure
5G**) which was statistically significant (paired t-test, t_{crit}=10.88, p<0.001). No
significant differences were found for lights on vs. off (Wilcoxon signed rank test,
p=0.50; **Figure 5H1a, 5H1b**), but all seizures occurred in the awake state (**Figure
5H3a, 5H3b**).
To determine whether non-convulsive seizures persisted with time, 4 out of 5
animals recorded at 2-4 wks were recorded again at 10-12 wks, as well as 3 more
animals (78-3, 80-4, 81-1) that were recorded at 10-12 wks only (**Figure 6A**).
Quantified non-convulsive seizures at 10-12 wks post-IHKA are shown in **Figure 6B-
H**. Non-convulsive seizures at 2-4 wks and 10-12 wks were similar in number (paired
t-test, t_{crit} = 1.88, p=0.15; **Figure 6B, Supplemental Figure 2B**) and frequency
(paired t-test, t_{crit} = 1.89, p=0.15; **Figure 6C**, **Supplemental Figure 2C**). Seizure
duration and severity are summarized in **Figures 6E** and **F** respectively and there
was no significant difference from 2-4 wks (**Supplemental Figure 2D**). The percent
of days spent with vs. without seizures did not change between timepoints (Wilcoxon
signed rank test, p=0.25; **Figure 6G vs. Figure 5G**). There were also no significant
differences in the proportion of seizures occurring during lights on vs. off (Wilcoxon
signed rank test, p>0.99), a.m. vs. p.m. (Wilcoxon signed rank test, p=0.18) and
awake vs. sleep states (Wilcoxon signed rank test, p=0.50; **Figure 6H**), which did
not differ with the 2-4 wk timepoint (**Supplemental Figure 3B**).
888
**IV. Different seizure onset patterns can be recorded and their prevalence changes
with time**
Recently several laboratories have suggested that seizures in rodent models of TLE
can simulate human seizures (Velasco et al., 2000), and discussed several types of
seizures based on their onset (Bragin et al., 1999b; Avoli et al., 2016). A common
nomenclature refers to two types primarily: one type with low-voltage fast activity at the
seizure onset (LVF seizure; Bragin et al., 1999b) and another type with a
hypersynchronous onset (HYP; Bragin et al., 1999b).
897

898
899
Figure 6. Quantification of chronic spontaneous non-convulsive seizures 10-12 wks post-IHKA

(A) Experimental timeline of the study. The seizures for this figure were recorded 10-12 wks after IHKA (green arrows; total # seizures=58). The mice were the same as those used for convulsive seizure measurements.

(B) The total number of chronic non-convulsive seizures during the 2 wk-long recording period is shown per animal. Blue and pink shades represent males and females, respectively.

(C) The number of non-convulsive seizures per day and per animal is shown for all recording days. The animal ID is shown in the inset.

(D) Non-convulsive seizure frequency was calculated as the mean number of non-convulsive seizures per day for each animal.

(E) Non-convulsive seizure duration was calculated as the mean per animal (left) or the mean of all non-convulsive seizures (right; n=58 seizures).

(F) Non-convulsive seizure severity was calculated as the mean per animal (left) or the mean of all non-convulsive seizures (right; n=58 seizures).

(G) Non-convulsive seizure burden was defined as the percent of days spent with (SZ) or without (No SZ) non-convulsive seizures in the 2 wks of continuous vEEG starting at 10 wks.

(H) The percent of non-convulsive seizures is shown, either occurring during the light period or dark period of the light:dark cycle (Lights ON or OFF; H1), a.m. or p.m. (AM/PM; H2), and awake (AW) or sleep (SL) state (H3). The percentages were calculated as the mean per animal (H1a, H2a, H3a) or the mean of all seizures (H1b, H2b, H3b) respectively. They were no significant differences (Wilcoxon signed rank tests; all p>0.05).

The seizures we recorded showed onset patterns that remarkably, almost always fit the description of LVF and HYP seizures. The LVF pattern was characterized by a "sentinel" spike and by brief suppression of background EEG followed by rhythmic activity (**Figure 7A**). The HYP pattern was characterized by a progressive increase in spike frequency (2-5 Hz) that eventually escalated to the point where a seizure was clear (**Figure 7B**). However, there were some seizures that could not fit into the two categories, which we classified as "unclear" onset.

931
932
Figure 7. Chronic IHKA seizures show different seizure onset patterns
(A) Representative example of a low-voltage fast (LVF) onset seizure recorded during the 10-12 wk recording session post-IHKA. A 30 sec-long EEG trace (Top) and an expanded 10 sec-long epoch (Bottom) taken from the seizure onset are shown. Note that the LVF pattern starts with a sentinel spike (asterisk and single red arrow in the expanded trace at the Bottom) followed by brief suppression of the background EEG (2 red arrows pointing at each side of the EEG suppression) and subsequent series of spikes.
(B) Representative example of a hypersynchronous (HYP) onset seizure recorded during the 2-4 wk recording period post-IHKA. A 2 min-long EEG trace (Top) and an expanded 30 sec-long epoch (Bottom) taken from the seizure onset are shown. Note that the HYP patterns start with a series of spikes (asterisks) followed by spikes at increased frequency.

Next, we quantified the different seizure onset patterns to understand if one of the two patterns was more prominent (**Figure 8**). For this analysis, we analyzed a total of 632 seizures (all data from each timepoint were pooled). Remarkably, almost all seizures either fit the LVF or HYP pattern (**Figure 8B**). There were no significant differences between LVF and HYP seizures in the total number (paired t-test, t_{crit}=2.02, p=0.08; **Figure 8B**) or frequency (paired t-test, t_{crit}=2.03, p=0.08; **Figure 8C**). When seizure durations were measured for each animal, the means were not different (Wilcoxon signed rank test, p=0.84; **Figure 8D1).** There also was no difference when all seizure durations were pooled (Wilcoxon signed rank test, p=0.20; **Figure 8D2**). The severity was not different when it was measured for each animal (Wilcoxon signed rank test, p=0.31; **Figure 8E1**), but HYP seizures were significantly more severe when all seizures were pooled (Wilcoxon signed rank test, p<0.0001; **Figure 8E2**), probably because the sample size for severity was animals and the sample size for all seizures was much larger

959
Figure 8. Quantification of low-voltage fast and hypersynchronous onset seizures
(A) Experimental timeline of the study. The data used for this figure were pooled from the 2-4 wk (red
arrows) and 10-12 wk timepoints (green arrows) and a total of 632 seizures were included in the
analyses. Seizures were distinguished as HYP, LVF, or seizure onset type was unclear. There were 8
mice (2 males, blue, recorded at both 2-4 and 10-12 wks; 6 females, pink, where 2 were recorded at
both times, and 4 were recorded only at one of the times).
(B) The total number of seizures is shown. There was no significant difference between HYP and LVF
seizures (paired t-test, t_{crit}=2.02, p=0.08).
(C) Same as B but the frequency of seizures is plotted. There was no significant difference between HYP
and LVF seizures (paired t-test, t_{crit}=2.03, p=0.08).
(D) Same as B but seizure duration is plotted. Seizure duration was calculated as an average per animal
(D1), or all seizures were pooled (D2). Neither was significant (Wilcoxon signed rank tests, p>0.05).
(E) Same as B but seizure severity is shown. Seizure severity was calculated as an average per animal
(E1) and for all seizures (E2). HYP seizures were significantly more severe than LVF seizures when all
seizures were included (E2; Wilcoxon signed rank test, p<0.0001). In other words, E1 was not
significant but E2 was, and that result is likely to be due to the large number of seizures in E2 compared
to animals in E1.
977

We next evaluated if there was a preference towards a particular seizure onset
pattern between the 2 different timepoints, 2-4 or 10-12 wks. Results are presented in
Figure 9. The total number of HYP and LVF seizures was similar between timepoints
(HYP: Wilcoxon signed rank test, p=0.12; LVF: Wilcoxon signed rank test, p=0.25;
Figure 9B). Interestingly, when the percent of HYP and LVF seizures were analyzed
instead of the total number, HYP and LVF seizures were significantly different between
timepoints with more HYP seizures dominating the earlier timepoint and more LVF
seizures dominating the later timepoint (Fisher's exact test, p<0.0001; **Figure 9C**).
Regarding the frequency of LVF or HYP seizures (**Figure 9D**), HYP seizures were
more frequent at 2-4 wks (paired t-test, t_{crit} = 3.50, p=0.03). When all seizures were
pooled, HYP seizures recorded 10-12 wks after IHKA lasted longer than their early
counterparts (Wilcoxon signed rank test, p=0.02; **Figure 9E2**) with no significant
differences in their severity (Wilcoxon signed rank test, p>0.99; **Figure 9F2**). No

991
992
Figure 9. Quantification of different seizure onset patterns between timepoints
(A) Experimental timeline of the study. The data used for this figure were both from the 2-4 wk (red arrows) and 10-12 wk timepoint (green arrows). Seizures were distinguished as HYP or LVF. Seizures that were unclear in onset are not included in this figure because they were relatively rare (see Figure 8).
(B) The total number of chronic HYP and LVF seizures recorded at the early (2-4 wks) and late (10-12 wks) timepoints are shown. No significant differences were found (Wilcoxon signed rank test, p>0.05).
(C) The percent of HYP vs. LVF seizures are shown for early and late timepoints. HYP seizures dominated early timepoints whereas LVF seizures dominated late timepoints (Fisher's exact test, p<0.0001).
(D) HYP and LVF seizure frequency was calculated as the mean number of seizures per day for each timepoint (early, late). HYP seizures were more frequent early vs. late (paired t-test, t_{crit} = 3.50, p=0.03), whereas LVF seizures did not differ between timepoints (Wilcoxon signed rank test, p>0.05).
(E) HYP and LVF seizure duration was calculated as the mean number of seizures per day for each animal at each timepoint (early, late; E1). No significant differences were found (Wilcoxon signed rank test, p>0.05). (E2) Same as in E1 but all HYP and LVF seizures were pooled and are presented according to timepoint and type (HYP, LVF). HYP seizures lasted longer in duration at the late timepoint (Wilcoxon signed rank test, p=0.02) and the late HYP seizures were longer than the late LVF seizures (Wilcoxon signed rank test, p=0.02).
(F) HYP and LVF seizure severity was calculated as the mean per animal at each timepoint (early, late; F1). (F2) Same as in F1 but all seizures were pooled and are presented according to timepoint and type (HYP, LVF). No significant differences were found (Wilcoxon signed rank test, p>0.05).

1013
differences between 2-4 and 10-12 wks were found for the total number of LVF seizures (Wilcoxon signed rank test, p=0.81; **Figure 9B**), mean frequency (Wilcoxon signed rank test, p=0.17; **Figure 9D**), duration (Wilcoxon signed rank test, p=0.50; **Figure 9E1**, Wilcoxon signed rank test, p=0.81; **Figure 9E2**) or severity (Wilcoxon signed rank test, p=0.75; **Figure 9F1**, Wilcoxon signed rank test, p=0.75; **Figure 9F2**). Similar results were obtained when convulsive seizures were analyzed only (**Supplemental Figure 4**). A more detailed comparison of convulsive seizure onset patterns among animals that showed an increase ('progressed') vs. a decrease in seizures ('non-progressed') between timepoints is shown in **Supplemental Figure 5**.

1023

V. HFOs (>250Hz) after IHKA are frequent at the site of IHKA injection, during seizures, and also occur interictally

Additional analysis of the EEG traces revealed the presence of HFOs in the frequency range of 250-500 Hz. HFOs were frequent during slow wave sleep, in line with previously published observations (Staba et al., 2004). Examples of interictal HFOs are shown in **Figure 10B** and their spectral properties are shown in **Figure 10D1, D2**.

Figure 10. HFOs (>250Hz) are frequent interictally, before and during seizures, and are recorded primarily from the IHKA injection site

(A) Experimental timeline of the study. The data used for this figure come from the 10-12 wk timepoint (red arrows).

(B) Representative example of interictal HFOs recorded from the left hemisphere. Note the presence of HFOs (red arrows) in the left hippocampal lead where they were in the trough of slow waves. Note that interictal HFOs were not always in the trough of slow waves, however.

(C) An example of an HFO recorded at the sentinel spike of an LVF seizure (black arrow) as well as during the seizure (red arrows). In this instance, HFOs were primarily recorded from the left hippocampal lead but this was not always the case.

(D) Two examples of HFOs (D1, D2). For both D1 and D2, the top is wideband recording (0.5-500 Hz) showing coupling of HFOs to the trough of slow waves. The centers are the filtered traces (250-500 Hz) and the bottom shows spectral properties of HFOs with frequencies >250 Hz in the 250-500 Hz time-frequency domain.

(E) Quantification of HFOs (events/min) based on either the left (hippocampus or cortex) or right (hippocampus or cortex) electrodes. There was a tendency towards more frequent HFOs in the left hemisphere compared to the right, but it did not reach statistical significance (paired t-test, p=0.057).

(F) The percent of seizures that showed HFOs at their onset (F1). The percentage of HYP vs. LVF seizures that showed HFOs at seizure onset was significantly different (Fisher's exact test, p<0.05; F2). Thus, a greater percent of HYP vs. LVF seizures seemed to be associated with HFOs at seizure onset (Fisher's exact test, p<0.05; F2). The percentage of HYP and LVF seizures for each category (HFOs, no HFOs) was calculated for the total number of seizures for each seizure type (HYP or LVF).

.

Although HFOs tended to be more frequent in the left compared to the right
hemisphere (**Figure 10E**) this was not confirmed statistically (paired t-test, t_{crit} =

p=0.05). Importantly, we found that interictal HFOs occurred in both sexes. The
mean number of HFOs recorded from the left hemisphere (hippocampus or cortex)
in the males was 20±11.5 and in the females was 130.3±58.3. However, sex
differences were not significant (Mann-Whitney U-test, U = 7, p=0.32).

HFOs also appeared frequently at seizure onset and during seizures. An
example of HFOs recorded at seizure onset and during an LVF seizure is shown in
Figure 10C. Forty-four percent of all recorded seizures showed HFOs at seizure
onset (**Figure 10F1**). When we looked at the percentage of HYP vs. LVF seizures
that showed and did not show HFOs at their onset we found a significant difference
(Fisher's exact test, p<0.05; **Figure 10F2**). Thus, a greater percent of HYP vs. LVF
seizures seemed to be associated with HFOs at seizure onset and conversely, a
greater percent of LVF vs. HYP seizures showed no HFOs at seizure onset
(Fisher's exact test, p<0.05; **Figure 10F2**). The percentage of HYP and LVF
seizures for each category (HFOs, no HFOs) was calculated for the total number of
seizures for each seizure type (HYP or LVF). Our data suggest that HFOs may be a
better biomarker for the onset of HYP seizures than LVF seizures, which is
consistent with prior studies showing that HFOs >250 Hz predominate HYP seizure
onsets primarily (Lévesque et al., 2012).

**VI. Hippocampal damage to the ipsilateral hippocampus is consistent with MTS
and correlates with SE severity but not chronic seizure burden**

To determine post-IHKA neuropathological outcomes, animals were sacrificed
at 12 wks post-IHKA (**Figure 11A**) and coronal brain sections located adjacent to
the IHKA injection site were stained with cresyl violet (**Figure 11**). Neuronal loss
was primarily in the ipsilateral hippocampus (**Figure 11B, C**) and included loss in
the hilus and pyramidal cell layers (PCL) of CA3 and CA1. Damage to the
hippocampal pyramidal cells near the IHKA injection site was estimated by
measuring the length of the PCL where neurons survived. For this purpose,
measurements were made from, the border of CA1 with the subiculum through the
entirety of CA1, CA3, and CA3, ending at the border of CA3 with the hilus
(**Supplemental Figure 1**). We found a significant reduction in PCL length after
IHKA compared to controls (Mann-Whitney U-test, U = 0, p=0.002, n=12; **Figure
11D1**).

To quantify granule cell dispersion, the granule cell layer (GCL) area was
measured, as well as the thickness (**Supplemental Figure 1**). There was a greater
GCL area in IHKA-treated animals compared to saline-treated controls (Mann-Whitney
U-test, U = 5, p=0.036; **Figure 11D2**), suggesting granule cell dispersion after IHKA. In
addition, there was increased GCL thickness in IHKA-treated animals vs. saline-injected
controls (Mann-Whitney U-test, U = 4, p=0.024; **Figure 11D3**).

1099

Figure 11. Quantification of hippocampal damage at 12 wks post-IHKA

(A) Experimental timeline of the study indicating the time animals were perfused at 12 wks post-IHKA (red arrows). Neuronal damage was quantified based on cresyl violet-stained coronal sections that were adjacent to the IHKA injection site (-2 mm A-P, ± 1.25 mm M-L, 1.6 mm D-V) corresponding approximately to plate 47 of a standard mouse stereotaxic atlas (Franklin and Paxinos, 1997).

(B) An example of mild hippocampal damage meaning that there was neuronal loss but less than the severe damage in C. The white arrow points to neuronal loss in the left CA1 pyramidal cell layer. Note that mild hippocampal damage was rare in the animals we studied. Calibration bar: 200 μm.

(C) An example of severe hippocampal damage. Note extensive neuronal loss in all pyramidal layers of the left hippocampus. Also, note extensive granule cell dispersion (white arrow). Calibration bar: 200 μm.

(D) Measurements of the IHKA-injected hippocampus are shown. Neuronal damage in the pyramidal cell layers was quantified by the length of the pyramidal cell layer (PCL) which did not exhibit neuronal loss. Note the reduced PCL length in IHKA vs. saline-injected animals (Mann-Whitney U-test, U = 0, p=0.002) D1). Granule cell layer (GCL) area was used as a reflection of granule cell dispersion with larger GCL area in IHKA vs. saline-injected animals (Mann-Whitney U-test, U = 5, p=0.036; D2) The width of the GCL was used as another measure of granule cell dispersion, confirming that the GCL is wider in IHKA vs. saline-injected animals (Mann-Whitney U-test, U = 4, p=0.024; D3).

(E) There was a significant correlation between the number of stage 5 seizures during IHKA-induced SE and PCL length (r=-0.68, p=0.04; E1), but not GCL area (r=0.35, p=0.34; E2) and not GCL thickness (r=0.30, p=0.42; E3). All IHKA-injected animals were included in the analyses.

(F) There was no significant correlation between the total number of chronic seizures (at both timepoints; 2-4 and 10-12 wks) and PCL length (r=0.13, p=0.83; F1), GCL area (r=-0.73, p=0.15; F2) or GCL thickness (r=-0.61, p=0.27; F3) at 12 wks post-IHKA. For the total number of chronic seizures, seizures were pooled for the two timepoints and only those animals with recordings at both timepoints are included.

To address the relationship between hippocampal damage and seizures during SE, PCL length, GCL area, and GCL thickness were plotted relative to the number of stage 5 seizures during SE (**Figure 11E, F**). PCL length was significantly correlated with the number of stage 5 seizures, such that PCL length was shorter when mice had more stage 5 seizures (r=-0.68, p=0.04; **Figure 11E1**), indicating the more severe seizures

during SE were associated with more neuronal loss in the PCL. The idea that more neuronal damage occurs when SE is more severe is consistent with prior studies of SE (Dingledine et al., 2014). GCL area was not correlated significantly with the number of stage 5 seizures during SE (r=0.35, p=0.34; **Figure 11E2**) and GCL thickness was not either (r=0.30, p=0.42; **Figure 11E3).** These data are consistent with the idea that GCL dispersion may not be a result of the severity of SE but additional factors (Haas et al., 2002; Kobow et al., 2009).

To address the relationship between hippocampal damage and the number of chronic seizures, the PCL length, GCL area, or GCL thickness were plotted relative to the total number of chronic seizures at both timepoints (2-4 and 10-12 wks) post-IHKA (**Figure 11F**). The correlations were not significant (PCL length: r=0.13, p=0.83, **Figure 11F1;** GCL area: r=-0.73, p=0.15, **Figure 11F2**; GCL thickness: r=-0.61, p=0.27, **Figure 11F3)**. The reasons that PCL length did not correlate with chronic seizure burden is consistent with the idea that PCL neuronal loss is primarily due to SE-associated excitotoxicity (Dingledine et al., 2014). However, one would expect more damage to lead to more chronic seizures. The reason that estimates of GCL dispersion were not correlated with chronic seizure burden may be due to the idea that it arises during epileptogenesis but not as a result of chronic seizures (Haas et al., 2002; Kobow et al., 2009).

IV. DISCUSSION

A. Summary of main findings

The results suggest that our use of IHKA in mice results in robust spontaneous convulsive seizures. The seizures are robust because of their severity and duration. Also, animals had frequent convulsive seizures that continued for over 2 wks. In addition, these seizures were robust because the electrographic correlates showed high amplitude, long-lasting rhythmic activity in each of our 4 leads. These seizures ended with postictal suppression of the EEG, which is another characteristic that supports the description of the seizures as robust. These seizures were evident 2-4 wks after IHKA and at 10-12 wks, but some animals showed an increase in frequency and others showed a decrease, suggesting variability but robust epilepsy, nevertheless. Notably, seizure duration consistently increased with time even if frequency did not, suggesting progressively worse epilepsy depending on the type of measurement of seizures. Non-convulsive seizures were also evident but were less frequent than convulsive seizures. Interestingly, seizures did not appear to be focal, meaning they were not confined to one electrode (such as the site where IHKA was infused).

Convulsive seizures showed seizure onset patterns other investigators have found in their rat models of epilepsy (Bragin et al., 1999b; Bragin et al., 2005; Lévesque et al., 2012), as well as humans with TLE (Velasco et al., 2000; Perucca et al., 2014; Gnatkovsky et al., 2019; Saggio et al., 2020). HFOs were frequent not only at the site of IHKA injection but the ipsilateral cortical site, suggesting a wider epileptogenic network beyond the IHKA injection site. Although HFOs were observed in the contralateral hemisphere, they were relatively rare. Nevertheless, diverse sites of HFOs suggest caution before assuming they are the site of the seizure focus, consistent with a previous study showing HFOs outside the IHKA focus in mice (Sheybani et al., 2018).

However, the rapid generalization of the seizures we recorded did not permit us to
ascertain where the focus was, and there could have been more than one focus.
Histopathological findings were consistent with prior reports about the IHKA model
(Bouilleret et al., 1999), including extensive hippocampal neuronal loss in the area of
IHKA injection, like MTS. Although other types of quantification are common, such as
cell counts, our conclusion that there was extensive pyramidal cell loss would have
been clear with almost any measurement, in our view. For this reason, and to avoid the
uncertainty of cell counts in the packed cell layers, we did not count cells. We mention
in the Results that we consider the measurement of PCL damage an estimate only.
Besides neuronal loss, granule cell dispersion was also evident in our animals, as
reported before (Bouilleret et al., 1999). Far less disruption of the contralateral
hippocampus is also consistent with prior reports (Bouilleret et al., 1999).
1189
B. IHKA produces robust chronic epilepsy
Several electrophysiologic elements of chronic convulsive seizures reported here
are distinct from the very frequent short-lasting epileptiform abnormalities reported
before (Kim et al., 2018; Sandau et al., 2019; Lai et al., 2020). One element is seizure
duration. We provide evidence of prolonged convulsive seizures similar to seizure
durations reported in human TLE (Balish et al., 1991) and data from an Epilepsy
Monitoring Unit (Jenssen et al., 2006). The longer-lasting seizures contrast with the 3-7
sec epileptiform abnormalities reported by some past studies of IHKA in mice (Kim et
al., 2018; Sandau et al., 2019; Lai et al., 2020). In addition, our results differ from what
appears to be continuous spiking at the site of IHKA injection indicated by some
investigators (Sandau et al., 2019).
Another aspect of the seizures we recorded was the presence of complex rhythmic
activity during a seizure. Thus, seizures we recorded were more complex
electrographically than some trains of spikes shown by others (Kim et al., 2018; Sandau
et al., 2019; Lai et al., 2020). For instance, our seizures showed fast and slow spikes as
well as repetitive spikes and waves at many different frequencies. These complex EEG
patterns were found for both convulsive and non-convulsive seizures. Finally, the
convulsive seizures we recorded were typically followed by prolonged post-ictal
suppression and this was not clear in studies by others (Kim et al., 2018; Sandau et al.,
2019; Lai et al., 2020).
Another aspect of the EEG that we found was a characteristic of seizures (both
convulsive and non-convulsive) was that they were recorded at 4 sites, both hippocampi
and two cortical locations. In the past, many studies of the IHKA model in mice did not
use as many recording sites so less information was available (Riban et al., 2002;
Zeidler et al., 2018). On the basis of their recording sites and the lack of convulsive
behavior, conclusions were sometimes made that the model showed focal seizures
primarily (Riban et al., 2002). An important point about our recordings is that we found
synchronized activity even from the start of the seizure. This was especially clear for
LVF seizures because at the onset there is a sentinel spike.
Regarding the site of the seizure focus, based on the findings from sentinel spikes,
one might argue that seizure onset is where the sentinel spike is largest, the
hippocampus. This interpretation seems logical since it was the site of IHKA injection.
However, the subdural screws for cortical recordings had less resolution than depth

electrodes in the hippocampus. In addition, it has been suggested that the origin of LVF
seizures is extrahippocampal (Velasco et al., 2000).
Chronic convulsive seizures were detected both at 2-4 and 10-12 wks, another
argument that chronic epilepsy was robust because convulsive seizures continued to
occur. The fact that seizures persisted is important to be noted as there is little
quantified longitudinal data of chronic seizure outcomes beyond 2 months post-IHKA in
mice (Henshall, 2017). However,1 or 2 motor seizures a wk and seizures at 8 months
were mentioned in the text of the study of Bouilleret et al. (Bouilleret et al., 1999).
We found that at 10-12 wks after IHKA, 50% of animals increased their seizure
frequency and the rest decreased their seizure frequency compared to the 2-4 wk
timepoint. None of the animals showed an absence of seizures at 10-12 wks. The
variability in seizure frequency is important because it is consistent with human TLE.
Thus, seizure diaries of patients with TLE suggest that some individuals may
experience an increase in seizures with time whereas others report a decrease (Bauer
and Burr, 2001). However, it is important to also note that patients were taking ASDs
and our animals were not. Also, seizure diaries are not as accurate as continuous
vEEG. The variable seizure frequencies we found have important implications for
preclinical drug testing, because testing is usually for only one time period, and that
lasts only 2 wks. Our data suggest that a 2 wk-long testing period may not predict
seizure frequency throughout the lifespan. Testing at different timepoints during chronic
epilepsy has been stressed by human studies already (Zimmermann and Trinka, 2020;
Thomson et al., 2021).
One study of IHKA is important to note because the investigators did study
progression of the epilepsy and no clear progression was found (Welzel et al., 2019).
Those data are consistent with our findings in that we found progression only in half of
the animals.
1249
C. Chronic IHKA seizures show different seizure onset patterns
The original studies of IHKA in mice have suggested no changes in seizure patterns
with time (Bouilleret et al., 1999). However, a limitation to assess this might be related
to the fact that convulsive seizures were rare, or discontinuous vEEG with minimal
bilateral recording was employed (Bouilleret et al., 1999; Riban et al., 2002). Different
chronic seizure onset patterns were evident in the same animal, and almost all seizures
could be categorized as LVF or HYP. Importantly, seizure onset patterns resembled
those in TLE (Velasco et al., 2000; Gnatkovsky et al., 2019) and following IHKA in rats
(Bragin et al., 1999b; Bragin et al., 2005; Lévesque et al., 2012). Our results suggest
that the prevalence of LVF or HYP seizures in an animal may change with time, a
finding that is consistent with earlier observations in the rat pilocarpine- and IHKA-
treated rats (Bragin et al., 1999b). This is important to be noted as it suggests changes
in the seizure-generating network as epilepsy continues. On the other hand,
characteristics of the LVF or HYP seizures did not significantly change. For example,
both seizure types continued to be accompanied by stage 4-5 convulsions and were
followed by prolonged postictal depression. On the other hand, HYP seizure duration
increased with time. Therefore, the data presented here suggest that once initiated,
HYP seizures may reverberate more with time, leading to a prolongation of the HYP
seizure.

D. High frequency oscillations (HFOs) occurred during seizures, at seizure onsets, as well as interictally

HFOs are considered hallmarks of epileptogenicity by numerous clinical (Jacobs et al., 2008; Weiss et al., 2016) and animal (Bragin et al., 1999a) studies and are currently used in the presurgical evaluation of patients with epilepsy (Zijlmans et al., 2019). Although definitions and terms vary, we defined HFOs in a way that distinguished them from most normal oscillations that have high frequency, such as those between 100 and 200 Hz. Our definition was >250 Hz. We also used spectrograms to ensure peak frequency was indeed above 250 Hz. Note that others have used the term "pathological HFO (pHFO)" to refer to HFOs in epilepsy, but we have avoided that term because some HFOs occur in normal tissue (Engel et al., 2009; Pearce et al., 2014).

HFOs in our study were present at seizure onsets and during seizures, which has been reported in acute and chronic IHKA-induced seizures in the rat (Bragin et al., 1999b; Lévesque et al., 2012; Li et al., 2018) and in TLE (Weiss et al., 2016). We found frequent HFOs during periods of slow wave activity similar to reports of HFOs during delta activity in humans (Staba et al., 2004) and rats (Bragin et al., 1999a; Bragin et al., 2016). Importantly the NREM stage of sleep where delta oscillations are prominent appears to be the most useful to identify the seizure onset zone in TLE (Klimes et al., 2019). We also report HFOs outside the IHKA injection site, which suggests a wider epileptogenic network involving adjacent cortical and remote hippocampal areas (Sheybani et al., 2018).

There is currently only one mouse IHKA study that reported HFOs, but it mostly focused on power (Häussler et al., 2012) rather than the identification of individual HFO events during sleep, which is standard for the detection and evaluation of HFOs in clinical practice (Frauscher et al., 2017). Importantly, we found that both male and female mice showed HFOs. This is important because both sexes have not been studied before. These data suggest that HFOs, an abnormality with an important translational value, are in both sexes. The findings support the increasing need to expand the bandwidth normally used for preclinical studies to capture high frequency abnormalities in the EEG of animals with epilepsy.

E. Hippocampal pathology at the IHKA injection site was consistent with Mesial Temporal Sclerosis (MTS)

Human tissue from many patients with intractable TLE shows extensive pyramidal cell loss. The pattern called MTS is characterized by neuronal death primarily in the hilus, CA3 and CA1 (Houser, 1999; Scharfman and Pedley, 2006; Blümcke et al., 2012; Thom, 2014). In addition, the dentate gyrus is characterized by granule cell dispersion (Houser, 1990; Suzuki et al., 1995; Bouilleret et al., 1999; Riban et al., 2002). The original studies of IHKA showed extensive pyramidal cell loss and granule cell dispersion near the site of IHKA injection, with little evidence of neuropathology in the contralateral hippocampus (Suzuki et al., 1995; Bouilleret et al., 1999). In the animals that we studied after IHKA, we also saw extensive pyramidal cell loss near the site of IHKA injection, although there was variability. Quantification showed a significant reduction in pyramidal cells near the IHKA injection site in both CA3 and CA1, like MTS. We also observed significant granule cell dispersion in most animals, and when

quantified, granule cell dispersion was reflected by an increase in granule cell area and
thickness (post-IHKA compared to saline).
The mechanisms by which KA induces cell death in pyramidal layers are thought to
be excitotoxicity mediated by KA receptors, and due to prolonged seizures (Ben-Ari et
al., 1980). To address a possible correlation between histopathology and seizures
during SE, we plotted the number of stage 5 seizures during IHKA-induced SE and
histopathological measurements (pyramidal cell length, GCL area, and GCL average
thickness). We also plotted the relation between chronic seizures and histopathological
measurements. The data showed a significant correlation between neuronal loss in the
pyramidal cell layers and the number of stage 5 seizures during SE. The results are
similar to those of the intraamygdala KA model in rats (Ben-Ari et al., 1980; Henshall et
al., 2000) and mice (Araki et al., 2002) although in prior studies the investigators used
other methods to quantify neuronal damage than the ones we implemented. Together
these data support the idea that severe SE is needed for the neuronal loss in the
pyramidal cell layers (Dingledine et al., 2014). This is notable because a long-standing
assumption is that chronic epilepsy is more likely when there is greater neuronal loss
(Dingledine et al., 2014).
It is important to note that we evaluated pyramidal cell loss at the end of our
experiments, typically 12 wks post-IHKA. Others typically examine neuronal loss in the
days following SE (Ben-Ari et al., 1980; Henshall et al., 2000; Araki et al., 2002).
However, we did assess some animals at early times and the neuronal loss in the
pyramidal cell layer was similar. Moreover, in SE models the vast majority of neuronal
loss occurs within 10 days of SE so we do not think that examining mice after 12 wks
was a major limitation.
In contrast to the length of the pyramidal cell layer, measurements of granule cell
dispersion were not correlated with acute or chronic seizures. This lack of correlation in
our data might be attributed to the fact that granule cell dispersion is an alteration which
is not dependent on SE severity or chronic seizures. Alternatively, the measures of SE
or chronic seizures may not have been those that are related to granule cell dispersion.
For example, we did not measure the power or duration of SE but the number of stage 5
seizures during SE because the animals were not vEEG monitored during SE. Power
and duration of SE may be more sensitive as a measure of SE severity than the number
of individual convulsive seizures. The same could be true of chronic seizures. On the
other hand, the mechanisms underlying granule cell dispersion may be initiated by the
way KA receptors modify the granule cells, independent of SE convulsive seizures or
chronic epilepsy. Other possibilities also exist, based on proposed mechanisms for
granule cell dispersion (Haas et al., 2002).
1351
F. Using IHKA to induce robust chronic epilepsy in mice
Our data suggest that the IHKA model in mice induces robust chronic epilepsy,
unlike several past reports where chronic convulsive seizures are not noted, or the EEG
evidence does not clearly reflect robust seizures (Kiasalari et al., 2016; Zhu et al., 2016;
Runtz et al., 2018; Bielefeld et al., 2019; Li et al., 2020). Notably, some investigators
have stated that they see chronic epilepsy after IHKA, but the evidence is not always
presented (Kim et al., 2018; Sandau et al., 2019; Lai et al., 2020).

Why would our methods lead to more convulsive seizures after IHKA-SE than other methods? One potential explanation is related to the fact that our animals sustained IHKA-SE with multiple stage 5 seizures. In some other studies of IHKA, non-convulsive SE was reported (Bouilleret et al., 1999; Riban et al., 2002; Arabadzisz et al., 2005; Maroso et al., 2011). If it is true that there is more neuronal loss when there are more convulsive seizures during SE, and furthermore if it is true that the greater neuronal loss is, the more robust the chronic seizures, then our initiation of IHKA-SE with many stage 5 seizures could have led to more robust epilepsy.

Another possibility is that we did not implant animals with electrodes or cannula prior to SE (or immediately after IHKA injection), which is notable because in the past we have found that implanted animals have a higher seizure threshold ((Jain et al., 2019) and HES, unpublished).

We also handled mice more than is typical. Thus, we handled mice once or twice per day for the 2 days before IHKA injection and a longer time period afterwards to account for the behavioral stress related to the lack of social housing (Bernard, 2019; Manouze et al., 2019). Animals also were housed with miniature enclosures so that animals could enter the enclosure and not be seen. This type of environment would be likely to lower behavioral stress even more. A study in the pilocarpine model suggested that single housing may increase seizures by a factor of 16 (Manouze et al., 2019). Single housing might thus explain why very frequent and brief (3-7 sec) epileptiform abnormalities have been reported in IHKA-treated mice in the past (Kim et al., 2018; Sandau et al., 2019; Lai et al., 2020; Rusina et al., 2021). It would be possible that our approach that included frequent handling combined with enriched housing reduced these very frequent abnormalities thereby allowing the expression of robust seizures. These ideas are consistent with past studies suggesting that stress plays a major role in seizure induction and in chronic epilepsy in both animals (MacKenzie and Maguire, 2015) and humans (Lang et al., 2018), although the relationship between stress and seizures/epilepsy is complex (Gunn and Baram, 2017).

Another factor is that we modified the pH of the solution of KA we injected so it was closer to a physiological pH. In the past, the pH may not have been monitored, and if not, a very acidic solution would have been injected because the pH of our solution, before adding a base to bring it closer to the physiological range, was approximately 4.0. In the past, we found that bringing the solution of pilocarpine to pH 7.4 before injection played a very large role in the consequences of pilocarpine with fewer seizures when pH was not controlled (Scharfman et al., unpublished). Finally, we found that seizures can be robust, but if a time when seizures are limited is the only time that is studied, one might get a false sense of few seizures when animals actually had robust epilepsy at other times.

G. Limitations of the study

One of the limitations of our study is that some mice were transgenic. However, the mice were designed to express Cre recombinase conditionally, and only in small groups of neurons. Since we did not conditionally activate Cre, we assume the mice that were hemizygous acted like the background strain, C57BL/6. In addition, wild type mice were also used. In fact, we found no evidence that the hemizygous and wild type mice

differed in acute effects of SE or chronic seizures. Therefore, we do not think the use of transgenic mice was a major limitation.

We also did not assess the EEG during SE, because all mice exhibited robust convulsive SE. Also, the presence of cannulas and electrodes could have influenced SE.

ACKNOWLEDGEMENTS

We thank Drs. Annamaria Vezzani, Esther Krook-Magnuson, Ivan Soltesz, and Peter West for sharing their methods for injecting KA in mice. We also thank members of the Scharfman lab for fruitful discussions and John LaFrancois for sharing his method for intrahippocampal injection in mice. This work was supported by the National Institutes of Health [grant number NIH R01 NS106983] and the New York State Office of Mental Health.

CONTRIBUTION OF AUTHORS

Christos P. Lisgaras and Helen E. Scharfman: Conceptualization, **Christos P. Lisgaras:** Data curation, **Christos P. Lisgaras and Helen E. Scharfman:** Formal analysis; **Helen E. Scharfman:** Funding acquisition, **Christos P. Lisgaras:** Investigation; **Christos P. Lisgaras:** Methodology, **Helen E. Scharfman:** Project administration, **Helen E. Scharfman:** Resources, **Christos P. Lisgaras:** Software, **Helen E. Scharfman:** Supervision, **Christos Panagiotis Lisgaras:** Validation, **Christos P. Lisgaras and Helen E. Scharfman:** Visualization, **Christos P. Lisgaras:** Roles/Writing - original draft, **Helen E. Scharfman and Christos P. Lisgaras:** Writing - review & editing.

Table 1. Animals included in the study

Animal number	Subject ID	Sex	Genotype	Experimental use	Treatment	Dose (nL)	Times of EEG recording	Times of anatomical assessment
1	78a-5	Male	Cre -/-	EEG, Anatomy	KA	80	2 wks, 10 wks	12 wks
2	76-1	Female	Cre +/-	EEG, Anatomy	KA	100	2 wks	10 wks
3	76-4	Female	Cre -/-	EEG, Anatomy	KA	100	2 wks, 10 wks	12 wks
4	75a-5	Male	Cre +/-	EEG, Anatomy	KA	100	2 wks, 10 wks	12 wks
5	77-2	Female	Cre +/-	EEG, Anatomy	KA	100	2 wks, 10 wks	12 wks
6	78-3	Female	Cre -/-	EEG, Anatomy	KA	70	10 wks	12 wks
7	78a-6	Male	Cre +/-	Anatomy	KA	75	NA	3 wks
8	79a-6	Male	Cre -/-	Anatomy	KA	70	NA	3 days
9	80-4	Female	Cre -/-	EEG, Anatomy	KA	70	10 wks	12 wks
10	81-1	Female	Cre -/-	EEG, Anatomy	KA	70	10 wks	12 wks
11	82-1	Female	Cre +/-	Anatomy	Saline	100	NA	2 wks
12	82-2	Female	Cre +/-	Anatomy	Saline	100	NA	2 wks
13	82a-3	Male	Cre -/-	Anatomy	Saline	80	NA	3 days
14	82a-4	Male	Cre -/-	Anatomy	Saline	100	NA	4 wks

Table 1 Legend. Animals included in the study. For each animal we note the identification number, sex, genotype, experimental use, treatment (KA, saline), dose (nL of KA or saline), times of EEG recording and anatomical assessment. The dosing varied because pilot studies showed that doses between 70 and 100 nL all were successful in producing severe SE and in addition no correlation was found between dosing and the total number of chronic seizures (r=0.19, p=0.64). Therefore, doses were randomly assigned for this study. Orange: IHKA animals; Blue: saline-injected controls. NA: Not applicable.

Table 2. Variables for induction of IHKA-SE and chronic epilepsy outcomes

Animal number	Subject ID	Sex	Genotype	Experimental use	Treat-ment	Dose (nL)	Total duration of anesthesia (min)	Latency to Stg 5 (min) from IHKA injection	Total number of Stg5 during SE	Total number of chronic seizures	Administra-tion of 0.9% saline s.c. post-IHKA
1	78a-5	Male	Cre -/-	EEG, Anatomy	KA	80	22	44	19	46	Yes
2	76-1	Female	Cre +/-	EEG, Anatomy	KA	100	17	28	8	8	No
3	76-4	Female	Cre -/-	EEG, Anatomy	KA	100	15	27	10	136	Yes
4	75a-5	Male	Cre +/-	EEG, Anatomy	KA	100	17	11	4	168	No
5	77-2	Female	Cre +/-	EEG, Anatomy	KA	100	15	42	6	59	Yes
6	78-3	Female	Cre -/-	EEG, Anatomy	KA	70	16	49	5	132	No
7	78a-6	Male	Cre +/-	Anatomy	KA	75	20	63	2	NA	No
8	79a-6	Male	Cre -/-	Anatomy	KA	70	17	39	8	NA	No
9	80-4	Female	Cre -/-	EEG, Anatomy	KA	70	19	66	4	8	No
10	81-1	Female	Cre -/-	EEG, Anatomy	KA	70	19	79	2	79	No
11	82-1	Female	Cre +/-	Anatomy	Saline	100	21	NA	NA	NA	NA
12	82-2	Female	Cre +/-	Anatomy	Saline	100	21	NA	NA	NA	NA
13	82a-3	Male	Cre -/-	Anatomy	Saline	80	21	NA	NA	NA	NA
14	82a-4	Male	Cre -/-	Anatomy	Saline	100	19	NA	NA	NA	NA

Table 2 Legend. For each animal we note the subject ID, sex, genotype, experimental use, treatment (KA, saline), dose (nL of KA or saline), total duration of anesthesia (defined as the time anesthesia started to the time anesthesia ended), latency to the first stage 5 seizure (defined as the time to a stage 5 convulsion after the end of anesthesia). We also note the total number of stage 5 seizures and chronic seizures (2-4 and 10-12 wk seizures were pooled). We also note whether animals were administered with 0.9% of saline s.c. after IHKA (typically the day after IHKA i.e., 24 hrs post-IHKA).

IHKA dose: There was no significant correlation between IHKA dose and the total number of stage 5 seizures during SE (r=0.18, p=0.6), suggesting the dose had little influence on the severity of SE.

Anesthesia: There were no significant correlations between the total duration of anesthesia and the latency to the first stage 5 seizure (r=0.40, p=0.31), the total number of stage 5 seizures during SE (r=0.27, p=0.43), or the total number of chronic seizures (r=-0.43, p=0.28), suggesting that the duration of anesthesia had little influence on outcome after IHKA.

Latency to the first stage 5 and number of stage 5 seizures during SE: Like the lack of correlation between anesthesia and latency to the first stage 5 seizure, there was no correlation between the latency to the first stage 5 and the total number of stage 5 seizures during SE (r=-0.34, p=0.33) or total number of chronic seizures (r=-0.43, p=0.28). These data suggest little effect of the latency of the first stage 5 seizures on the severity of SE or subsequent chronic seizures. The total number of stage 5 seizures during SE was not correlated with the total number of chronic seizures (r=-0.19, p=0.64) which is surprising because one might expect more severe SE to lead to more severe epilepsy.

Genotype: We did not find any statistically significant differences between animals that were Cre -/- (n=6) or Cre +/- (n=4) in terms of the latency to
the first stage 5 seizure (Mann-Whitney U-test, U = 7, p=0.35) or the total number of stage 5 seizures during SE (Mann-Whitney U-test, U = 8.5,
p=0.50). Also, the total number of chronic seizures was not different between Cre -/- (n=5) vs. Cre+/- (n=3) animals (Mann-Whitney U-test, U = 7.5,
p>0.99). These data suggest Cre-/- and Cre+/- mice were similar, regarding the measurements in this table.
Administration of saline: Animals that received saline s.c. the day after IHKA (n=3) because of transient body weight loss were similar to those
that did not (n=5) in terms of total number of chronic seizures (Mann-Whitney U-test, U = 7, p=0.92).
Orange: IHKA animals; Blue: saline-injected controls. NA: Not applicable.
1474

 SUPPLEMENTAL MATERIAL

1476

A. Pyramidal cell length and granule cell layer area

B. Upper blade and lower blade thickness

1477
1478
Supplemental Figure 1. Quantification of hippocampal pathology 12 wks post-IHKA
Methods used for cresyl violet-stained sections from dorsal hippocampus near the area of the IHKA
injection. Measurements were made using ImageJ as discussed in the Methods.
(A) A cresyl violet-stained section (A1) and the same image without the stained section (A2) shows the
lines used for measurements. For pyramidal cell length (PCL), a line was drawn using the freehand line
tool from CA3c to the end of CA1. The line included those areas where pyramidal cells were contiguous,
i.e., there was <1 soma width between cell bodies. The CA1 border with the subiculum and the CA3c
border with the hilus was the end of CA1 and CA3c respectively. These areas were defined in the
Methods. For the measurement of the granule cell layer (GCL) area, we traced the borders of the GCL
using the freehand tool and calculated the area within the borders. The border of the GCL and hilus
and border of the GCL and IML were defined by the point where adjacent cells became >1 GC soma
width apart.
(B) For GCL width, we used the straight-line tool to measure the thickest part of the upper and lower blade
(red arrows; B1) based on the traced borders of GCL. The white line with arrows (B1) was perpendicular
to the length of the GCL as shown in B2. We averaged 2 measurements for an estimate of upper blade
width of a section and then did the same procedure for the lower blade to yield the estimate for the
lower blade width of that section.

Supplemental Figure 2. Quantification of convulsive and non-convulsive seizures for 2-4 and 10-12 wks post-IHKA for animals that recorded at both timepoints

(A) Experimental timeline shows the times when measurements were made; 2-4 wks (red arrows) and 10-12 wks (green arrows) post-IHKA.

(B) The total number of chronic convulsive and non-convulsive seizures recorded during 2 wks of continuous vEEG is shown for the 2-4 wk timepoint (n=5 animals; 2 males, light and royal blue; 3 females, pink and red) and 10-12 wk timepoint (n=4 animals; 2 males, 2 females). These animals were the same except for one that was removed from the study and therefore could not be included in the 10-12 wk timepoint. The number of convulsive seizures was significantly greater than the number of non-convulsive seizures (paired t-test, t_{crit} 3.29, p=0.030), suggesting that the majority of seizures were convulsive.

(C) Same as B, but seizure frequency is shown. The frequency of convulsive seizures was significantly greater than non-convulsive seizures (paired t-test, t_{crit} 3.32, p=0.029), again suggesting that there were more convulsive seizures in the animals.

(D) Same as C, but seizure duration is shown. Seizure duration was calculated as an average per animal. The duration of convulsive or non-convulsive seizures did not differ between timepoints (convulsive: Wilcoxon signed rank test, p=0.25; non-convulsive: Wilcoxon signed rank test, p>0.99). Note that elsewhere we discuss that seizure duration increases with time so these data might seem not to support that idea. However, we note elsewhere that it is only when seizures are differentiated between HYP and LVF that there is an ability to see increased seizure duration with time.

1518
1519
Supplemental Figure 3. Percent of convulsive and non-convulsive seizures at the two time points
(2-4, 10-12 wks): effects of light, time of day and sleep.
(A) Percent of convulsive seizures that occurred when lights were on or off, a.m. or p.m., and awake or
sleep for the two time points (2-4 and 10-12 wks). Only animals recorded at both timepoints are
included.
(B) Same as in A but for non-convulsive seizures.
(C) Data were compared using just the 2-4 wk timepoint.
(D) Same as C, but for the 10-12 wk timepoint. There were no significant differences (paired t-tests, p>0.05)
most likely due to the variability from one animal to the next.

1535
Supplemental Figure 4. Seizure onset patterns at 2-4 and 10-12 wks for convulsive seizures only
(A) Experimental timeline of the study. Data from the early (red arrows) and late (green arrows) timepoint
are included in this figure. There were 5 mice at 2-4 wks (3 males, 2 females). The same mice were
recorded at 10-12 wks except for one of the males that could not be recorded at 10-12 wks.
(B) The total number of chronic convulsive HYP and LVF seizures recorded during 2-4 wks and 10-12
wks. No significant differences were found for LVF seizures between the 2 timepoints (Wilcoxon
signed rank test, p=0.5). Statistical comparisons for HYP seizures were not possible because of the
small sample size of HYP seizures at the late timepoint.
(C) Percent of HYP and LVF seizures at the 2-4 and 10-12 wk timepoints. The percent of HYP vs. LVF
seizures is different between timepoints with HYP seizures dominating the earlier timepoint and LVF
seizures the late timepoint (Fisher's exact test, p<0.0001).
(D) Same as B, but convulsive seizure frequency is shown. The frequency of LVF seizures did not
change between timepoints (Wilcoxon signed rank test, p=0.25).
(E) Same as B, but convulsive seizure duration is plotted. Seizure duration was calculated as the
average per animal in E1 and for all seizures in E2. No significant differences were found for LVF
seizures between the 2 timepoints (Wilcoxon signed rank test, p=0.50). Statistical comparisons for
HYP seizures were not possible because of the small sample size of HYP seizures at the late
timepoint.
(F) Same as B, but convulsive seizure severity is shown. Convulsive seizure severity was calculated as
the average per animal in F1 and for all seizures in F2. There were no significant differences
(Wilcoxon signed rank tests, p>0.05).
1557

1558
1559
Supplemental Figure 5. Comparison of different convulsive seizure onset patterns in mice that did
or did not show progression at 10-12 wks
(A) Experimental timeline of the study as shown before. Data from the early timepoint (red arrows) are
used for this figure. The comparisons are between mice that did not "progress" (seizures were most
frequent at 2-4 wks) and mice that "progressed" (seizures were most frequent at 10-12 wks). There
were 2 mice/group. P, progressed; NP, non-progressed.
(B) The total number of convulsive HYP and LVF seizures recorded during 2-4 wks is shown for two
subsets of mice, those with seizures that increased between 2-4 wks and 10-12 wks (progressed, P)
and those mice with seizures that decreased (non-progressed, NP). There were 4 mice (2 males,
blue; 2 females, pink).
(C) The percentages of HYP or LVF seizures at 2-4 wks were compared between P and NP mice. Mice
that showed progression (P) had a significantly different proportion of HYP seizures relative to LVF
seizures depending on their progression (P) or non-progression (NP; Fisher's exact test, p<0.0001).
The data support the view that HYP seizures dominated non-progressed mice but LVF were more
prevalent in mice that progressed.
(D) Same as B, but seizure frequency is shown.
(E) Same as B, but seizure duration is plotted. Seizure duration was calculated as an average per
animal.
(F) Same as B, but seizure severity is shown. Convulsive seizure severity was calculated as an average
per animal.

REFERENCES

Amiri, M., Lina, J. M., Pizzo, F., Gotman, J., 2016. High frequency oscillations and spikes: Separating real hfos from false oscillations. *Clin Neurophysiol*. 127, 187-196.

Arabadzisz, D., Antal, K., Parpan, F., Emri, Z., Fritschy, J. M., 2005. Epileptogenesis and chronic seizures in a mouse model of temporal lobe epilepsy are associated with distinct EEG patterns and selective neurochemical alterations in the contralateral hippocampus. *Exp Neurol*. 194, 76-90.

Araki, T., Simon, R. P., Taki, W., Lan, J. Q., Henshall, D. C., 2002. Characterization of neuronal death induced by focally evoked limbic seizures in the C57BL/6 mouse. *J Neurosci Res*. 69, 614-621.

Avoli, M., de Curtis, M., Gnatkovsky, V., Gotman, J., Köhling, R., Lévesque, M., Manseau, F., Shiri, Z., Williams, S., 2016. Specific imbalance of excitatory/inhibitory signaling establishes seizure onset pattern in temporal lobe epilepsy. *J Neurophysiol*. 115, 3229-3237.

Bagshaw, A. P., Jacobs, J., LeVan, P., Dubeau, F., Gotman, J., 2009. Effect of sleep stage on interictal high-frequency oscillations recorded from depth macroelectrodes in patients with focal epilepsy. *Epilepsia*. 50, 617-628.

Balish, M., Albert, P. S., Theodore, W. H., 1991. Seizure frequency in intractable partial epilepsy: A statistical analysis. *Epilepsia*. 32, 642-649.

Bauer, J., Burr, W., 2001. Course of chronic focal epilepsy resistant to anticonvulsant treatment. *Seizure*. 10, 239-246.

Behr, C., Lévesque, M., Stroh, T., Avoli, M., 2017. Time-dependent evolution of seizures in a model of mesial temporal lobe epilepsy. *Neurobiol Dis*. 106, 205-213.

Ben-Ari, Y., Cossart, R., 2000. Kainate, a double agent that generates seizures: Two decades of progress. *Trends Neurosci*. 23, 580-587.

Ben-Ari, Y., Lagowska, J., 1978. Epileptogenic action of intra-amygdaloid injection of kainic acid. *C R Acad Hebd Seances Acad Sci D*. 287, 813-816.

1632 Ben-Ari, Y., Tremblay, E., Ottersen, O. P., 1980. Injections of kainic acid into the
1633   amygdaloid complex of the rat: An electrographic, clinical and histological study
1634   in relation to the pathology of epilepsy. *Neuroscience*. 5, 515-528.
1635

1636 Bénar, C. G., Chauvière, L., Bartolomei, F., Wendling, F., 2010. Pitfalls of high-pass
1637   filtering for detecting epileptic oscillations: A technical note on "false" ripples. *Clin
1638   Neurophysiol*. 121, 301-310.
1639

1640 Bernard, C., 2019. On the interpretation of results obtained in singly housed animals.
1641   *Epilepsia*. 60, 2013-2015.
1642

1643 Bielefeld, P., Schouten, M., Meijer, G. M., Breuk, M. J., Geijtenbeek, K., Karayel, S.,
1644   Tiaglik, A., Vuuregge, A. H., Willems, R. A. L., Witkamp, D., Lucassen, P. J.,
1645   Encinas, J. M., Fitzsimons, C. P., 2019. Co-administration of anti microRNA-124
1646   and -137 oligonucleotides prevents hippocampal neural stem cell loss upon non-
1647   convulsive seizures. *Front Mol Neurosci*. 12, 31.
1648

1649 Bielefeld, P., Sierra, A., Encinas, J. M., Maletic-Savatic, M., Anderson, A., Fitzsimons,
1650   C. P., 2017. A standardized protocol for stereotaxic intrahippocampal
1651   administration of kainic acid combined with electroencephalographic seizure
1652   monitoring in mice. *Front Neurosci*. 11, 160.
1653

1654 Blümcke, I., Coras, R., Miyata, H., Ozkara, C., 2012. Defining clinico-neuropathological
1655   subtypes of mesial temporal lobe epilepsy with hippocampal sclerosis. *Brain
1656   Pathol*. 22, 402-411.
1657

1658 Botterill, J. J., Lu, Y. L., LaFrancois, J. J., Bernstein, H. L., Alcantara-Gonzalez, D., Jain,
1659   S., Leary, P., Scharfman, H. E., 2019. An excitatory and epileptogenic effect of
1660   dentate gyrus mossy cells in a mouse model of epilepsy. *Cell Rep*. 29, 2875-
1661   2889.e2876.
1662

1663 Bouilleret, V., Ridoux, V., Depaulis, A., Marescaux, C., Nehlig, A., Le Gal La Salle, G.,
1664   1999. Recurrent seizures and hippocampal sclerosis following intrahippocampal
1665   kainate injection in adult mice: Electroencephalography, histopathology and
1666   synaptic reorganization similar to mesial temporal lobe epilepsy. *Neuroscience*.
1667   89, 717-729.
1668

1669 Bragin, A., Azizyan, A., Almajano, J., Wilson, C. L., Engel, J., 2005. Analysis of chronic
1670   seizure onsets after intrahippocampal kainic acid injection in freely moving rats.
1671   *Epilepsia*. 46, 1592-1598.
1672

1673   Bragin, A., Engel, J., Jr., Wilson, C. L., Fried, I., Mathern, G. W., 1999a. Hippocampal
1674   and entorhinal cortex high-frequency oscillations (100-500 Hz) in human epileptic
1675   brain and in kainic acid-treated rats with chronic seizures. *Epilepsia*. 40, 127-137.
1676

1677   Bragin, A., Engel, J., Wilson, C. L., Vizentin, E., Mathern, G. W., 1999b.
1678   Electrophysiologic analysis of a chronic seizure model after unilateral
1679   hippocampal KA injection. *Epilepsia*. 40, 1210-1221.
1680

1681   Bragin, A., Li, L., Almajano, J., Alvarado-Rojas, C., Reid, A. Y., Staba, R. J., Engel, J.,
1682   Jr., 2016. Pathologic electrographic changes after experimental traumatic brain
1683   injury. *Epilepsia*. 57, 735-745.
1684

1685   Bui, A. D., Nguyen, T. M., Limouse, C., Kim, H. K., Szabo, G. G., Felong, S., Maroso,
1686   M., Soltesz, I., 2018. Dentate gyrus mossy cells control spontaneous convulsive
1687   seizures and spatial memory. *Science*. 359, 787-790.
1688

1689   Clayton, J. A., Collins, F. S., 2014. Policy: NIH to balance sex in cell and animal studies.
1690   *Nature*. 509, 282-283.
1691

1692   Dingledine, R., Varvel, N. H., Dudek, F. E., 2014. When and how do seizures kill
1693   neurons, and is cell death relevant to epileptogenesis? *Adv Exp Med Biol*. 813,
1694   109-122.
1695

1696   Engel, J., Jr., Bragin, A., Staba, R., Mody, I., 2009. High-frequency oscillations: What is
1697   normal and what is not? *Epilepsia*. 50, 598-604.
1698

1699   Franklin, K. B. J., Paxinos, G., 1997. The mouse brain in stereotaxic coordinates.
1700   *Academic Press*, San Diego.
1701

1702   Frauscher, B., Bartolomei, F., Kobayashi, K., Cimbalnik, J., van 't Klooster, M. A.,
1703   Rampp, S., Otsubo, H., Höller, Y., Wu, J. Y., Asano, E., Engel, J., Jr., Kahane,
1704   P., Jacobs, J., Gotman, J., 2017. High-frequency oscillations: The state of clinical
1705   research. *Epilepsia*. 58, 1316-1329.
1706

1707   Gnatkovsky, V., Pelliccia, V., de Curtis, M., Tassi, L., 2019. Two main focal seizure
1708   patterns revealed by intracerebral electroencephalographic biomarker analysis.
1709   *Epilepsia*. 60, 96-106.
1710

1711   Gunn, B. G., Baram, T. Z., 2017. Stress and seizures: Space, time and hippocampal
1712   circuits. *Trends Neurosci*. 40, 667-679.

1713 Haas, C. A., Dudeck, O., Kirsch, M., Huszka, C., Kann, G., Pollak, S., Zentner, J.,
1714   Frotscher, M., 2002. Role for reelin in the development of granule cell dispersion
1715   in temporal lobe epilepsy. *J Neurosci*. 22, 5797-5802.
1716

1717 Harte-Hargrove, L. C., Galanopoulou, A. S., French, J. A., Pitkänen, A., Whittemore, V.,
1718   Scharfman, H. E., 2018. Common data elements (CDEs) for preclinical epilepsy
1719   research: Introduction to CDEs and description of core CDEs. A TASK3 report of
1720   the ILAE/AES joint translational task force. *Epilepsia Open*. 3, 13-23.
1721

1722 Häussler, U., Bielefeld, L., Froriep, U. P., Wolfart, J., Haas, C. A., 2012. Septotemporal
1723   position in the hippocampal formation determines epileptic and neurogenic
1724   activity in temporal lobe epilepsy. *Cereb Cortex*. 22, 26-36.
1725

1726 Häussler, U., Rinas, K., Kilias, A., Egert, U., Haas, C. A., 2016. Mossy fiber sprouting
1727   and pyramidal cell dispersion in the hippocampal CA2 region in a mouse model
1728   of temporal lobe epilepsy. *Hippocampus*. 26, 577-588.
1729

1730 Henshall, D. C., Chapter 41 - Poststatus epilepticus models: Focal kainic acid. In: A.
1731   Pitkänen, P. S. Buckmaster, A. S. Galanopoulou, S. L. Moshé, (Eds.), Models of
1732   Seizures and Epilepsy (second edition). *Academic Press*, 2017, pp. 611-624.
1733

1734 Henshall, D. C., Sinclair, J., Simon, R. P., 2000. Spatio-temporal profile of DNA
1735   fragmentation and its relationship to patterns of epileptiform activity following
1736   focally evoked limbic seizures. *Brain Res*. 858, 290-302.
1737

1738 Hitti, F. L., Siegelbaum, S. A., 2014. The hippocampal CA2 region is essential for social
1739   memory. *Nature*. 508, 88-92.
1740

1741 Holmes, G. L., 2007. Animal model studies application to human patients. *Neurology*.
1742   69, S28-32.
1743

1744 Houser, C. R., 1990. Granule cell dispersion in the dentate gyrus of humans with
1745   temporal lobe epilepsy. *Brain Res*. 535, 195-204.
1746

1747 Houser, C. R., 1992. Morphological changes in the dentate gyrus in human temporal
1748   lobe epilepsy. *Epilepsy Res Suppl*. 7, 223-234.
1749

1750 Houser, C. R., 1999. Neuronal loss and synaptic reorganization in temporal lobe
1751   epilepsy. *Adv Neurol*. 79, 743-761.
1752

1753   Iyengar, S. S., LaFrancois, J. J., Friedman, D., Drew, L. J., Denny, C. A., Burghardt, N.
1754   S., Wu, M. V., Hsieh, J., Hen, R., Scharfman, H. E., 2015. Suppression of adult
1755   neurogenesis increases the acute effects of kainic acid. *Exp Neurol*. 264, 135-
1756   149.
1757

1758   Jacobs, J., LeVan, P., Chander, R., Hall, J., Dubeau, F., Gotman, J., 2008. Interictal
1759   high-frequency oscillations (80-500 Hz) are an indicator of seizure onset areas
1760   independent of spikes in the human epileptic brain. *Epilepsia*. 49, 1893-1907.
1761

1762   Jain, S., LaFrancois, J. J., Botterill, J. J., Alcantara-Gonzalez, D., Scharfman, H. E.,
1763   2019. Adult neurogenesis in the mouse dentate gyrus protects the hippocampus
1764   from neuronal injury following severe seizures. *Hippocampus*. 29, 683-709.
1765

1766   Jenssen, S., Gracely, E. J., Sperling, M. R., 2006. How long do most seizures last? A
1767   systematic comparison of seizures recorded in the epilepsy monitoring unit.
1768   *Epilepsia*. 47, 1499-1503.
1769

1770   Kapur, J., Elm, J., Chamberlain, J. M., Barsan, W., Cloyd, J., Lowenstein, D., Shinnar,
1771   S., Conwit, R., Meinzer, C., Cock, H., Fountain, N., Connor, J. T., Silbergleit, R.,
1772   2019. Randomized trial of three anticonvulsant medications for status epilepticus.
1773   *N Engl J Med*. 381, 2103-2113.
1774

1775   Keezer, M. R., Sisodiya, S. M., Sander, J. W., 2016. Comorbidities of epilepsy: Current
1776   concepts and future perspectives. *Lancet Neurol*. 15, 106-115.
1777

1778   Kiasalari, Z., Khalili, M., Shafiee, S., Roghani, M., 2016. The effect of vitamin E on
1779   learning and memory deficits in intrahippocampal kainate-induced temporal lobe
1780   epilepsy in rats. *Indian J Pharmacol*. 48, 11-14.
1781

1782   Kim, J. Y., Kim, J. H., Lee, H. J., Kim, S. H., Jung, Y. J., Lee, H. Y., Kim, H. J., Kim, S.
1783   Y., 2018. Antiepileptic and anti-neuroinflammatory effects of red ginseng in an
1784   intrahippocampal kainic acid model of temporal lobe epilepsy demonstrated by
1785   electroencephalography. *Yeungnam Univ J Med*. 35, 192-198.
1786

1787   Kim, T. H., Richards, K., Heng, J., Petrou, S., Reid, C. A., 2013. Two lines of transgenic
1788   mice expressing cre-recombinase exhibit increased seizure susceptibility.
1789   *Epilepsy Res*. 104, 11-16.
1790

1791   Klimes, P., Cimbalnik, J., Brazdil, M., Hall, J., Dubeau, F., Gotman, J., Frauscher, B.,
1792   2019. NREM sleep is the state of vigilance that best identifies the epileptogenic
1793   zone in the interictal electroencephalogram. *Epilepsia*. 60, 2404-2415.
1794

1795 Kobow, K., Jeske, I., Hildebrandt, M., Hauke, J., Hahnen, E., Buslei, R., Buchfelder, M.,
1796   Weigel, D., Stefan, H., Kasper, B., Pauli, E., Blümcke, I., 2009. Increased reelin
1797   promoter methylation is associated with granule cell dispersion in human
1798   temporal lobe epilepsy. *J Neuropathol Exp Neurol*. 68, 356-364.
1799

1800 Kwan, P., Schachter, S. C., Brodie, M. J., 2011. Drug-resistant epilepsy. *N Engl J Med*.
1801   365, 919-926.
1802

1803 Lai, M. C., Wu, S. N., Huang, C. W., 2020. The specific effects of OD-1, a peptide
1804   activator, on voltage-gated sodium current and seizure susceptibility. *Int J Mol
1805   Sci*. 21.
1806

1807 Lang, J. D., Taylor, D. C., Kasper, B. S., 2018. Stress, seizures, and epilepsy: Patient
1808   narratives. *Epilepsy Behav*. 80, 163-172.
1809

1810 Lévesque, M., Avoli, M., 2013. The kainic acid model of temporal lobe epilepsy.
1811   *Neurosci Biobehav Rev*. 37, 2887-2899.
1812

1813 Lévesque, M., Salami, P., Gotman, J., Avoli, M., 2012. Two seizure-onset types reveal
1814   specific patterns of high-frequency oscillations in a model of temporal lobe
1815   epilepsy. *J Neurosci*. 32, 13264-13272.
1816

1817 Li, J., Leverton, L. K., Naganatanahalli, L. M., Christian-Hinman, C. A., 2020. Seizure
1818   burden fluctuates with the female reproductive cycle in a mouse model of chronic
1819   temporal lobe epilepsy. *Exp Neurol*. 334, 113492.
1820

1821 Li, L., Kriukova, K., Engel, J., Bragin, A., 2018. Seizure development in the acute
1822   intrahippocampal epileptic focus. *Scientific Reports*. 8.
1823

1824 Löscher, W., 2017. Animal models of seizures and epilepsy: Past, present, and future
1825   role for the discovery of antiseizure drugs. *Neurochem Res*. 42, 1873-1888.
1826

1827 MacKenzie, G., Maguire, J., 2015. Chronic stress shifts the GABA reversal potential in
1828   the hippocampus and increases seizure susceptibility. *Epilepsy Res*. 109, 13-27.
1829

1830 Manouze, H., Ghestem, A., Poillerat, V., Bennis, M., Ba-M'hamed, S., Benoliel, J. J.,
1831   Becker, C., Bernard, C., 2019. Effects of single cage housing on stress,
1832   cognitive, and seizure parameters in the rat and mouse pilocarpine models of
1833   epilepsy. *eNeuro*. 6.
1834

1835 Maroso, M., Balosso, S., Ravizza, T., Iori, V., Wright, C. I., French, J., Vezzani, A.,
1836   2011. Interleukin-1β biosynthesis inhibition reduces acute seizures and drug
1837   resistant chronic epileptic activity in mice. *Neurotherapeutics*. 8, 304-315.
1838

1839 McKhann, G. M., 2nd, Wenzel, H. J., Robbins, C. A., Sosunov, A. A., Schwartzkroin, P.
1840   A., 2003. Mouse strain differences in kainic acid sensitivity, seizure behavior,
1841   mortality, and hippocampal pathology. *Neuroscience*. 122, 551-561.
1842

1843 Meier, C. L., Dudek, F. E., 1996. Spontaneous and stimulation-induced synchronized
1844   burst afterdischarges in the isolated CA1 of kainate-treated rats. *J Neurophysiol*.
1845   76, 2231-2239.
1846

1847 Menendez de la Prida, L., Staba, R. J., Dian, J. A., 2015. Conundrums of high-
1848   frequency oscillations (80-800 Hz) in the epileptic brain. *J Clin Neurophysiol*. 32,
1849   207-219.
1850

1851 Nadler, J. V., 1981. Minireview. Kainic acid as a tool for the study of temporal lobe
1852   epilepsy. *Life Sci*. 29, 2031-2042.
1853

1854 Nadler, J. V., Perry, B. W., Cotman, C. W., 1978. Intraventricular kainic acid
1855   preferentially destroys hippocampal pyramidal cells. *Nature*. 271, 676-677.
1856

1857 Navarrete, M., Alvarado-Rojas, C., Le Van Quyen, M., Valderrama, M., 2016.
1858   RIPPLELAB: A comprehensive application for the detection, analysis and
1859   classification of high frequency oscillations in electroencephalographic signals.
1860   *PLoS One*. 11, e0158276.
1861

1862 Pearce, P. S., Friedman, D., Lafrancois, J. J., Iyengar, S. S., Fenton, A. A., Maclusky,
1863   N. J., Scharfman, H. E., 2014. Spike-wave discharges in adult Sprague-Dawley
1864   rats and their implications for animal models of temporal lobe epilepsy. *Epilepsy
1865   Behav*. 32, 121-131.
1866

1867 Perucca, P., Dubeau, F., Gotman, J., 2014. Intracranial electroencephalographic
1868   seizure-onset patterns: Effect of underlying pathology. *Brain*. 137, 183-196.
1869

1870 Pinel, J. P., Rovner, L. I., 1978. Electrode placement and kindling-induced experimental
1871   epilepsy. *Exp Neurol*. 58, 335-346.
1872

1873 Pitkänen, A., Buckmaster, P. S., Galanopoulou, A. S., Moshé, S. L., 2017. Models of
1874   Seizures and Epilepsy (second edition). *Academic Press*.
1875

1876   Racine, R. J., 1972. Modification of seizure activity by electrical stimulation. I. After-
1877   discharge threshold. *Electroencephalogr Clin Neurophysiol*. 32, 269-279.
1878

1879   Rattka, M., Brandt, C., Löscher, W., 2013. The intrahippocampal kainate model of
1880   temporal lobe epilepsy revisited: Epileptogenesis, behavioral and cognitive
1881   alterations, pharmacological response, and hippoccampal damage in epileptic
1882   rats. *Epilepsy Res*. 103, 135-152.
1883

1884   Ravizza, T., Onat, F. Y., Brooks-Kayal, A. R., Depaulis, A., Galanopoulou, A. S.,
1885   Mazarati, A., Numis, A. L., Sankar, R., Friedman, A., 2017. WONOEP appraisal:
1886   Biomarkers of epilepsy-associated comorbidities. *Epilepsia*. 58, 331-342.
1887

1888   Riban, V., Bouilleret, V., Pham-Lê, B. T., Fritschy, J. M., Marescaux, C., Depaulis, A.,
1889   2002. Evolution of hippocampal epileptic activity during the development of
1890   hippocampal sclerosis in a mouse model of temporal lobe epilepsy.
1891   *Neuroscience*. 112, 101-111.
1892

1893   Runtz, L., Girard, B., Toussenot, M., Espallergues, J., Fayd'Herbe De Maudave, A.,
1894   Milman, A., deBock, F., Ghosh, C., Guerineau, N. C., Pascussi, J. M., Bertaso,
1895   F., Marchi, N., 2018. Hepatic and hippocampal cytochrome P450 enzyme
1896   overexpression during spontaneous recurrent seizures. *Epilepsia*. 59, 123-134.
1897

1898   Rusina, E., Bernard, C., Williamson, A., 2021. The kainic acid models of temporal lobe
1899   epilepsy. *eNeuro*. 8.
1900

1901   Saggio, M. L., Crisp, D., Scott, J. M., Karoly, P., Kuhlmann, L., Nakatani, M., Murai, T.,
1902   Dümpelmann, M., Schulze-Bonhage, A., Ikeda, A., Cook, M., Gliske, S. V., Lin,
1903   J., Bernard, C., Jirsa, V., Stacey, W. C., 2020. A taxonomy of seizure
1904   dynamotypes. *Elife*. 9.
1905

1906   Salami, P., Lévesque, M., Gotman, J., Avoli, M., 2015. Distinct EEG seizure patterns
1907   reflect different seizure generation mechanisms. *J Neurophysiol*. 113, 2840-
1908   2844.
1909

1910   Sandau, U. S., Yahya, M., Bigej, R., Friedman, J. L., Saleumvong, B., Boison, D., 2019.
1911   Transient use of a systemic adenosine kinase inhibitor attenuates epilepsy
1912   development in mice. *Epilepsia*. 60, 615-625.
1913

1914   Scharfman, H. E., 2007. The neurobiology of epilepsy. *Current Neurology and
1915   Neuroscience Reports*. 7, 348-354.
1916

1917   Scharfman, H. E., Galanopoulou, A. S., French, J. A., Pitkänen, A., Whittemore, V.,
1918   Harte-Hargrove, L. C., 2018. Preclinical common data elements (CDEs) for
1919   epilepsy: A joint ILAE/AES and NINDS translational initiative. *Epilepsia Open*. 3,
1920   9-12.
1921

1922   Scharfman, H. E., Goodman, J. H., Sollas, A. L., 2000. Granule-like neurons at the
1923   hilar/CA3 border after status epilepticus and their synchrony with area CA3
1924   pyramidal cells: Functional implications of seizure-induced neurogenesis. *J*
1925   *Neurosci*. 20, 6144-6158.
1926

1927   Scharfman, H. E., Pedley, T. A., Temporal lobe epilepsy. In: Gilman, A., (Ed.), In the
1928   Neurobiology of Disease. *Academic Press*, New York, 2006.
1929

1930   Scharfman, H. E., Sollas, A. L., Smith, K. L., Jackson, M. B., Goodman, J. H., 2002.
1931   Structural and functional asymmetry in the normal and epileptic rat dentate gyrus.
1932   *J Comp Neurol*. 454, 424-439.
1933

1934   Schauwecker, P. E., Steward, O., 1997. Genetic determinants of susceptibility to
1935   excitotoxic cell death: Implications for gene targeting approaches. *Proc Natl Acad*
1936   *Sci U S A*. 94, 4103-4108.
1937

1938   Schwarcz, R., Zaczek, R., Coyle, J. T., 1978. Microinjection of kainic acid into the rat
1939   hippocampus. *Eur J Pharmacol*. 50, 209-220.
1940

1941   Sheybani, L., Birot, G., Contestabile, A., Seeck, M., Kiss, J. Z., Schaller, K., Michel, C.
1942   M., Quairiaux, C., 2018. Electrophysiological evidence for the development of a
1943   self-sustained large-scale epileptic network in the kainate mouse model of
1944   temporal lobe epilepsy. *J Neurosci*. 38, 3776-3791.
1945

1946   Shiri, Z., Manseau, F., Lévesque, M., Williams, S., Avoli, M., 2016. Activation of specific
1947   neuronal networks leads to different seizure onset types. *Ann Neurol*. 79, 354-
1948   365.
1949

1950   So, N. K., Blume, W. T., 2010. The postictal EEG. *Epilepsy Behav*. 19, 121-126.
1951

1952   Staba, R. J., Wilson, C. L., Bragin, A., Fried, I., Engel, J., Jr., 2002. Quantitative
1953   analysis of high-frequency oscillations (80-500 Hz) recorded in human epileptic
1954   hippocampus and entorhinal cortex. *J Neurophysiol*. 88, 1743-1752.
1955

1956 Staba, R. J., Wilson, C. L., Bragin, A., Jhung, D., Fried, I., Engel, J., Jr., 2004. High-
1957   frequency oscillations recorded in human medial temporal lobe during sleep. *Ann*
1958   *Neurol*. 56, 108-115.
1959

1960 Sutula, T. P., Pitkänen, A., 2002. Do seizures damage the brain? *Elsevier*, New York.
1961

1962 Suzuki, F., Junier, M. P., Guilhem, D., Sorensen, J. C., Onteniente, B., 1995.
1963   Morphogenetic effect of kainate on adult hippocampal neurons associated with a
1964   prolonged expression of brain-derived neurotrophic factor. *Neuroscience*. 64,
1965   665-674.
1966

1967 Thijs, R. D., Surges, R., O'Brien, T. J., Sander, J. W., 2019. Epilepsy in adults. *Lancet*.
1968   393, 689-701.
1969

1970 Thom, M., 2014. Review: Hippocampal sclerosis in epilepsy: A neuropathology review.
1971   *Neuropathol Appl Neurobiol*. 40, 520-543.
1972

1973 Thomson, K. E., West, P. J., Metcalf, C. S., Wilcox, K. S., 2021. Response: Usefulness
1974   of the post-kainate spontaneous recurrent seizure model for screening for
1975   antiseizure and for neuroprotective effects. *Epilepsia*.
1976

1977 Velasco, A. L., Wilson, C. L., Babb, T. L., Engel, J., Jr., 2000. Functional and anatomic
1978   correlates of two frequently observed temporal lobe seizure-onset patterns.
1979   *Neural Plast*. 7, 49-63.
1980

1981 VonDran, M. W., LaFrancois, J., Padow, V. A., Friedman, W. J., Scharfman, H. E.,
1982   Milner, T. A., Hempstead, B. L., 2014. p75NTP, but not proNGF, is upregulated
1983   following status epilepticus in mice. *ASN Neuro*. 6.
1984

1985 Weiss, S. A., Alvarado-Rojas, C., Bragin, A., Behnke, E., Fields, T., Fried, I., Engel, J.,
1986   Jr., Staba, R., 2016. Ictal onset patterns of local field potentials, high frequency
1987   oscillations, and unit activity in human mesial temporal lobe epilepsy. *Epilepsia*.
1988   57, 111-121.
1989

1990 Welzel, L., Schidlitzki, A., Twele, F., Anjum, M., Löscher, W., 2019. A face-to-face
1991   comparison of the intra-amygdala and intrahippocampal kainate mouse models
1992   of mesial temporal lobe epilepsy and their utility for testing novel therapies.
1993   *Epilepsia*.
1994

1995 Williams, P. A., Hellier, J. L., White, A. M., Staley, K. J., Dudek, F. E., 2007.
1996   Development of spontaneous seizures after experimental status epilepticus:
1997   Implications for understanding epileptogenesis. *Epilepsia*. 48 Suppl 5, 157-163.
1998

1999 Zaher, N., Urban, A., Antony, A., Plummer, C., Bagić, A., Richardson, R. M., Kokkinos,
2000   V., 2020. Ictal onset signatures predict favorable outcomes of laser thermal
2001   ablation for mesial temporal lobe epilepsy. *Front Neurol*. 11, 595454.
2002

2003 Zeidler, Z., Brandt-Fontaine, M., Leintz, C., Krook-Magnuson, C., Netoff, T., Krook-
2004   Magnuson, E., 2018. Targeting the mouse ventral hippocampus in the
2005   intrahippocampal kainic acid model of temporal lobe epilepsy. *eneuro*. 5,
2006   ENEURO.0158-0118.
2007

2008 Zhu, H., Zhao, Y., Wu, H., Jiang, N., Wang, Z., Lin, W., Jin, J., Ji, Y., 2016. Remarkable
2009   alterations of Nav1.6 in reactive astrogliosis during epileptogenesis. *Sci Rep*. 6,
2010   38108.
2011

2012 Zijlmans, M., Jiruska, P., Zelmann, R., Leijten, F. S., Jefferys, J. G., Gotman, J., 2012.
2013   High-frequency oscillations as a new biomarker in epilepsy. *Ann Neurol*. 71, 169-
2014   178.
2015

2016 Zijlmans, M., Zweiphenning, W., van Klink, N., 2019. Changing concepts in presurgical
2017   assessment for epilepsy surgery. *Nat Rev Neurol*. 15, 594-606.
2018

2019 Zimmermann, G., Trinka, E., 2020. Accounting for individual variability in baseline
2020   seizure frequencies when planning randomized clinical trials remains
2021   challenging. *Epilepsia*.

2022