Are microbes growing on flowers evil? Effects of old flower microbes on fruit set in a

wild ginger with one-day flowers, *Alpinia japonica* (Zingiberaceae)

Nuria Jiménez Elvira[1], Masayuki Ushio[2,3], Shoko Sakai[3]

[1]Centre de Recerca Ecològica i Aplicacions Forestals, 08193 Cerdanyola del Vallès,

Barcelona, Spain

[2]Hakubi Center, Kyoto University, Kyoto University, Kyoto 606-8501, Japan

[3]Center for Ecological Research, Kyoto University, Hirano 2-509-3, Otsu 520-2113,

Japan

Corresponding Author: shokosakai@ecology.kyotyo-u.ac.jp

Running title: Effects of old flower microbes on fruit set in *Alpinia japonicus*

Author contributions. NJE, MU and SS designed the study. NJE conducted field

observation, sampling and amplicon sequencing with the assistance of SS and MU,

respectively. SS and NJE wrote the manuscript. All authors gave final approval for

publication.

Keywords: flower longevity, fruit set, anthosphere, pollination, microbiome

Data availability statement: Original datasets of amplicon sequencing were deposited

in DDBJ (Accession No. DRA012415).

Acknowledgements: We are grateful to Otsu city for permission to conduct field work

in Seta Park. This research was supported by JSPS KAKENHI Grant Numbers 19K22455

and 20H03324.

Abstract

Flowers are colonized and inhabited by diverse microbes. Plants rapidly replace flowers

with short lifespan, and old flowers senesce. This may contribute to avoiding adverse

effects of the microbes. In this study, we investigate if the flower microbial community

on old flowers impedes fruit and seed production in a wild ginger with one-day flowers.

We inoculated newly opened flowers with old flower microbes, and monitored the

effects on fruit and seed set. We also assessed prokaryotic communities on the flowers

using amplicon sequencing. We found six bacterial amplicon sequence variants (ASVs)

whose proportions were increased on the inoculated flowers. These ASVs were also

found on flower buds and flowers that were bagged by net or paper during anthesis,

suggesting that they had been present in small numbers prior to flowering. Fruit set was

negatively associated with the proportions of these ASVs, while seed set was not. The

results suggest that old flowers harbor microbial communities different from those at

anthesis, and that the microbes abundant on old flowers negatively affect plant

reproduction. Though the short lifespan of flowers has gotten little attention, it might be

an essential defense mechanism to cope with antagonistic microbes that rapidly

proliferate on the flowers.

1. INTRODUCTON

Flowers are an elaborate device to exchange pollen among conspecific individuals. Instead of moving around to find a potential mate, the plants disperse and receive pollen relying on different vectors such as insects and wind to achieve sexual reproduction. As a consequence, flowers are also exposed to microbial colonization under natural conditions (Burdon et al. 2018; Kaltz and Shykoff 2001; Ushio et al. 2015). Flowers have various habitats for microbes. Flowers attract pollinators by secreting nectar, which is rich in sugars and often contains other nutrients, such as amino acids and lipids (Roy et al. 2017). The stigma is a germination bed for pollen grains connected to a growth chamber for pollen tubes. It maintains humidity and nutrients necessary for pollen tube growth (Taylor and Hepler 1997), which would also be beneficial for microbes. Recent studies using high-throughput sequencing have revealed highly diverse microbial communities on flowers of many species in a wide range of habitats (Gaube et al. 2021; Massoni et al. 2020; Wei and Ashman 2018).

Today, flower microbes are increasingly recognized as essential components in the ecology and evolution of plant reproduction (reviewed in Rebolleda-Gómez et al. 2019; Vannette 2020). Not surprisingly, some flower microbes are known to have strong negative effects on the survival and reproduction of the host plant. Many pathogens use flowers, especially the stigma and nectary, as a point of entry. Their infection of flowers often results in fruit abortion and systemic infection (Bubán et al. 2003; Vanneste 2000). Other microbes impede various processes of pollination and fertilization. Nectar yeasts are reported to modify pollinator foraging patterns and reduce pollination success (Herrera et al. 2013). *Acinetobacter* bacteria that commonly inhabit floral nectar exploit pollen nutrition by inducing pollen germination and bursting (Christensen et al. 2021). Most studies of these phenomena examined the effects of an individual microbial species or of separate strains rather than that of the whole community.

In general, even newly opened flowers already have microbes at least on some flower tissues. Their abundance increases over time on individual flowers (Vannette 2020). Flowers often have characteristics that may suppress microbial growth, such as flower volatiles (Burdon et al. 2018; Huang et al. 2012), reactive oxygen (McInnis et al. 2006) and secondary compounds (Dobson and Bergstrom 2000; Palmer-Young et al.

2019). An extremely short lifespan (from one to several days) of flowers (Primack

1985), especially in hot habitats (Bawa 1990), is also thought to reduce the chance of

microbial infection and the proliferation of microbes (Kaltz and Shykoff 2001; Rogers

2006; Shykoff et al. 1996). Unlike the reproductive structures of mammals, flowers, the

reproductive structures of the plants, are only retained while they are needed, and are

developed *de novo* every season in perennial species (Rogers 2006). We do not know

yet, however, to what extent microbes that are suppressed by abscission of flowers are

potentially detrimental to plant fecundity.

In this study, we examine if the flower microbial community on old flowers

impedes fruit and seed production in a wild ginger, *Alpinia japonica* (Zingiberaceae),

that produces short-lifespan flowers. This perennial herb grows on the humid forest

floor of evergreen and deciduous forests in western Japan. It flowers in June, when the

climate is hot and humid. Its flowers open in the morning and wilt at sunset within the

same day. Therefore, its flowering coincides with the optimal season for microbial

growth, while it may avoid the negative effects of microbes by renewing flowers daily.

To imitate microbial communities that would appear on the flowers of *A. japonica* if

their longevity were longer, we inoculate microbes from old flowers onto newly opened

flowers in the morning. Firstly, we characterize the prokaryote communities on *A.*

japonica flowers. Secondly, we identify prokaryotes that significantly increased after

inoculation by comparing prokaryotic communities between the inoculated flowers and

open flower controls. In addition, we examine whether these prokaryotes colonize via

air or via flower visitors after flowering, or if they had already been present before

anthesis. We covered the flowers by net or paper bags to exclude microbial colonization

by flower visitors or by both flower visitors and air, respectively. If the old-flower

microbes colonize only after anthesis, bagging would reduce the occurrences of these

microbes. Lastly, we observe fruit and seed sets of the flowers under the treatments. We

expected that the flowers that had more old-flower microbes would show lower fruit

and seed set if microbe proliferation on flowers is generally to be avoided.

2. MATERIALS AND METHODS

2.1 Study species

Alpinia japonica (Zingiberaceae) is a wild ginger distributed in temperate and
subtropical regions of eastern Asia. It is a perennial herb 0.5–0.7 m in height occurring
mostly in broad-leaved evergreen forests. Inflorescences are compound racemose with
10–60 flowers (37 on average in this population). Flowers are zygomorphic with a
prominent labellum 1.1–1.2 cm in length, and white with red stripes at the margin of the
labellum (Fig. 1a). Flowering occurs almost synchronously within a population and lasts
for 10–14 days for each plant between late May and June. From one to five flowers
open per inflorescence. At anthesis around 0700, the dorsal and lateral petals open, and
the labellum inside expands (Fig. 1b). The flowers are known to have volatiles,
including 1, 8-cineole, with antimicrobial activities (Asakawa et al. 2017; van Vuuren
and Viljoen 2007). The corolla wilts at sunset within the day. The ovary matures into an
ellipsoid, three-loculate fleshy capsule ~2 months after flowering. The species is self-
compatible, and pollinated by bees, mainly carpenter bees (*Xylocopa appendiculata*)
and bumble bees (*Bombus ardens*) (J. Nuria and S. Sakai, unpublished data).
123
2.2 Study site
Studies were conducted in May to July of 2018, in Seta Park, Otsu, Shiga
Prefecture, Japan (34°50'N, 135°50'E). Seta Park has a young secondary deciduous
forest where streams are abundant, which provides an optimal habitat for *A. japonica*.
The annual mean temperature is 14.9°C, and annual total rainfall is ~1530 mm in Otsu
city (Japan Meteorological Agency, https://www.data.jma.go.jp/). Mean daily maximum
temperatures in May, June and July 2018 were 24.0°C, 27.3°C and 33.7°C, respectively.
131
2.3 Flower treatments
Alpinia japonica typically presents 2–12 inflorescences per plant at the study
site. We selected six plants with at least eight inflorescences with more than 10 flowers
on each. Unfortunately, one of them had been damaged, probably by park visitors,
before the experiments and sampling. Therefore, we used the remaining five plants
(Plant ID = N1, N2, N4, N5, N6). In each target plant, eight inflorescences were tagged
and covered with a fine net (Cloth Cabin, Suminoe Teijin Techno, Osaka, Japan) prior
to flowering.

We applied one of five treatments to part of the flowers that opened on the

inflorescences between 22nd May and 7th June. The flowers under these treatments

were used for either identification of microbes on the flowers or for observation of fruit

set (Fig. 1c). We could not use the same flower for both, because we should collect the

flower on the same day for the former, while we should leave the flower intact for the

latter. The five treatments were as follows: 1) open control (OT), in which the mesh bag

on the inflorescence was removed during 0700–1700, so that microbes were transmitted

to the flower by both air and pollinators; 2) insect exclusion by a net bag (NB), in which

the inflorescence was left covered with the net, which allowed air but not insects to get

through; 3) paper bagged (PB), in which the mesh bag covering the inflorescence was

replaced by a new paper bag (Grape Bag, DAIICHI VINYL, Fukui, Japan) during

0700–1700 to exclude air flow and insect visits; 4) open and microbial inoculation (MI)

and 5) open and water sprayed mock (WI) treatments. In MI, the mesh bag on the

inflorescence was removed as in OT, and the flower was sprayed with microbial

solution. During the spray inoculation, the focal flower was isolated from the rest of the

inflorescence by a funnel-shaped plastic sheet to avoid contamination. The microbial

suspension for the inoculation was prepared by soaking old flowers in distilled water.

Nine old flowers that had opened the previous day from 1–3 individuals were collected

from individuals that did not receive the treatments. Microbes were detached by

ultrasonic dispersion in 10 ml of distilled water and kept at 4°C until the inoculation

(maximum of 36 hours). Inoculation was conducted on eight different days, and the

same inoculant was used for all flowers under MI treatment on the same day. In WI, the

mesh bag covering the inflorescence was removed as in OT, and the flower was sprayed

with distilled water as a control for the physical effect of the MI treatment. We did not

conduct the treatments on cloudy or rainy days. Flowers that opened on the same day on

the same inflorescence were assigned to the same treatments in the case of NB and PB,

while OT, MI and WI were occasionally mixed on an inflorescence. This is because we

bagged the entire inflorescence rather than a single flower to avoid damaging the

flowers. The sample size of each treatment is summarized in Table S1.

2.4 Identification of microbial communities

To identify microbial communities, we sampled 4–5 flowers with each treatment

from each plant. The flowers were cropped at ~1700 of the day of flower opening and

separately put into a 5 ml plastic tube after removing the stigma (to use for another

study) and petals (Fig. 1b). In addition, five flower buds were collected from each

individual to identify microbes that were present inside prior to anthesis with the same

procedure. The tube was kept on ice until being brought to the lab within 4 hours after

sampling and kept at –20°C until further processing.

To detach microbes from the flower surface, we added 3 ml of phosphate-

buffered saline solution (PBS buffer, pH 7.4, Nippon Gene Co., Ltd.) to the tube and

sonicated it using an Ultrasonic Disruptor Handy Sonic UR-21P (Digital Biology) at

Level 5 power for 30 sec. Microbes in the PBS solution were then filtered using a

Sterivex Millipore filter (Sterivex-HV 0.22, Merck). Residual PBS on the Sterivex filter

was removed using a Vac-Man® Laboratory Vacuum Manifold (Promega).

Microbial DNA was extracted with DNeasy Blood & Tissue Kits (QIAGEN)

following the manufacturer's instructions. A portion of the 16S small-subunit ribosomal

gene was PCR-amplified using the primer pairs 515F (5'-GTG YCA GCM GCC GCG

GTA A-3') and 806R (5'-GGA CTA CNV GGG TWT CTA AT-3') (Caporaso et al.

2011) with Nextera DNA index adapters. First PCR was performed in 12 μl reactions,

each containing 1 μl of DNA template, 6 μl of KAPA HiFi HotMasterMix (Kapa

Biosystems) and 0.7 μl of each primer (5 μM). The first PCR was conducted with a

temperature profile of 95°C for 3 min, followed by 35 cycles at 98°C for 20 s, 60°C for

15 s, 72°C for 30 s, and a final extension at 72°C for 5 min, followed by storage at 4°C

until use. Triplicate PCR products were pooled and purified with Agencourt AMPure

XP (Beckman Coulter). P5/P7 Illumina adaptors were then added in a second PCR

using fusion primers with 8-mer index sequences for sample identification (Hamady et

al. 2008) (forward, 5'-AAT GAT ACG GCG ACC ACC GAG ATC TAC AC-[8-mer

tag]-TCG TCG GCA GCG TCA GAT GTG TAT AAG AGA CAG -3'; reverse, 5'-

CAA GCA GAA GAC GGC ATA CGA GAT-[8-mer tag]-GTC TCG TGG GCT CGG

AGA TGT GTA TAA GAG ACA G-3'). This preparation had a total volume of 24 μl

with double the amount of each reagent in the first PCR and 10-times diluted first PCR

products as DNA template. The temperature profile was 95°C for 3 min, followed by 12

cycles at 98°C for 20 s, 68°C for 15 s, 72°C for 15 s, a final extension at 72°C for 5

min, and 4°C until use. Second PCR products were pooled at equal volumes after a

purification/equalization process with the same AMPure XP protocol and E-Gel

SizeSelect II Agarose Gel kit (Invitrogen). DNA quantitation was performed using a

Qubit® dsDNA HS Assay Kit (Thermo Fisher Scientific Inc). The pooled library was

sequenced on an Illumina Miseq using Reagent Kit V2 (500 cycle Nano Kit).

We processed the sequence data following Ushio (Ushio 2019). The raw MiSeq

data were converted into FASTQ files using the bcl2fastq program provided by Illumina

(bcl2fastq v2.18). The FASTQ files were then demultiplexed using the command

implemented in Claident (http://www.claid ent.org; (Tanabe and Toju 2013)).

Demultiplexed FASTQ files were analysed using the ASV method implemented in the

package DADA2 (Callahan et al. 2016) of R (R Core Team 2016). At the quality

filtering process, forward and reverse sequences were trimmed at the length of 215 and

160, respectively, using DADA2::filterAndTrim() function.

Taxonomic identification was performed for ASVs (Amplicon Sequence

Variants; proxy for Operational Taxonomic Units) inferred using DADA2 based on the

query-centric auto-k-nearest-neighbour (QCauto) method (Tanabe and Toju 2013) and

subsequent taxonomic assignment with the lowest common ancestor algorithm (Huson

et al. 2007) using the "merge" database and clidentseq and classigntax commands

implemented in claident v0.2.2018.05.29. The reads identified as Bacteria or Archaea

were used for further analysis. Taxonomy, ASV table, and sample information were

combined into a single R object for downstream manipulation using the R package

phyloseq (McMurdie and Holmes 2013). We also removed ASVs that only occurred in

negative control samples. Furthermore, potential contaminant ASVs were identified

statistically based on prevalence in negative control samples using the R package

decontam (Davis et al. 2018) based on ASV prevalence and a threshold of 0.5. This

resulted in elimination of 87 of 2915 16S ASVs.

2.5 Fruit set and seed production

The treated flowers for the observation of fruit set under all the treatments were

hand-pollinated around 1700 using outcrossing pollen to ensure that the fruit set was not

limited by pollen. Pollen was taken with a wood toothpick from a cut flower from a

pollen donor plant, of which inflorescences had been covered with a mesh bag prior to

anthesis to prevent removal of pollen by flower visitors. In total, 251 pollinated flowers

were monitored for fruit and seed set (Table S1). Six weeks after the treatment, we

surveyed whether the fruits under the treatments were still retained on the inflorescence,

and collected the fruits (131 fruits) and counted developing seeds and aborted ovules

inside. At that time, the fruits were still green but fully plump.

2.6 Statistical analysis of microbial community data

To examine differences in the prokaryotic diversity among the treatments, we

evaluated the Shannon diversity index of each sample using the

phyloseq::estimate_richness() function. Significant differences in the index among

treatments were tested using Generalized Linear Mixed Model (GLMM). Prior to the

analysis, we confirmed that distributions of the index were not significantly deviated

from the normal distribution (Shapiro-Wilk test, $W = 0.983$, $p = 0.0759$). We

constructed a GLMM with the treatment as a fixed term and Plant ID as a random factor

using the R package lme4 (Bates et al. 2015). Significance of the fixed term was tested

by comparing the full model and the model without the fixed term using the anova()

function.

We examined variation in the structure of the prokaryotic communities using the

ASVs that were present in 20 or more flower samples (36 ASVs satisfied this criterion).

The frequency of each ASV was divided by the number of the total sequences of the 36

ASVs in each sample to convert it to a proportion.

To compare microbial communities among flowers under the different

treatments, we calculated the Bray–Curtis dissimilarity index using the package vegan

(Oksanen et al. 2010) in R. We performed a permutational multivariate analysis of

variance (PERMANOVA) using the vegan::adonis function to examine if flowers of the

same individuals nested under the same treatment have similar microbial communities.

Ordinations were plotted with nonmetric multidimensional scaling (NMDS) using the

vegan::metaMDS procedure. We then calculated the distances of each flower sample

from the centroid of the treatment. Variation in the distances among the treatments were

tested using vegan::anova for betadisper object.

To find ASVs that tend to co-occur in the samples, we conducted hierarchical

clustering of ASVs based on Bray–Curtis dissimilarity using Ward's minimum variance

algorithm using the hclust procedure. The algorithm merges clusters that minimize the

increase in the sum of squared distances from the cluster centroid. We chose the number

of clusters that maximize the silhouette index (Rousseeuw 1987). The index was

calculated using silhouette in the R package cluster (Maechler et al. 2001). The

proportions of the ASVs in the sample were visualized by a heatmap created using the R

package ComplexHeatmap (Gu et al. 2016). We identified the cluster that showed the

largest difference in the median proportion in the samples under the microbial inoculate

(MI) treatment compared with that under the water sprayed (WI) treatment. Differences

in the proportions of the cluster between the MI and WI treatments were also evaluated

by the statistic of the Wilcoxon rank sum test. The identification of the ASVs in the

cluster was searched against the Nucleotide database of the National Center for

Biotechnology Information (NCBI) with BLAST (Altschul et al. 1990). Top matches of

the sequences were referred to in order to assign taxonomic identities (Class, Family,

Genus) of the ASVs.

2.7 Statistical analysis of fruit set and seed set data

To assess the factors associated with fruit set and seed production, we used

generalized linear mixed models (GLMMs). To evaluate differences in fruit set among

the treatments after controlling for the difference among the individuals, we modeled

fruit set (retained/aborted) with the treatments as a fixed factor and the individuals as a

random factor using a binomial error distribution. Then we calculated and plotted

estimated marginal means of fruit set for the treatments using the ggpredict() function of

the R package ggeffects (Lüdecke 2018). Then, to test the association of fruit set with

microbial communities, we constructed the second model for fruit set. The model

included microbial compositions as a fixed factor and the individuals as a random factor

using a binomial error distribution. We identified Cluster 6 to include ASVs that

showed the largest increase after the microbial inoculation (see below). Therefore, we

used the average proportion of Cluster 6 of each treatment as the variable for the

microbial composition. We evaluated the significance of the fixed factor by the Wald

test.

We also modeled seed set (retained/aborted) and tested its association with

microbial composition. The model included the proportion of Cluster 6 as a fixed factor

and the individuals and fruit as nested random factors. We fitted the model with a

binominal error distribution. The GLMM analyses for fruit and seed sets were

performed using the R package lme4 (Bates et al. 2015).

3. RESULTS

3.1 Microbial communities

We obtained 839,113 reads of prokaryotic sequences, which were clustered into

2,828 ASVs. We have removed samples that had < 250 prokaryotic reads (12 samples).

This reduced the number of samples to 134. According to the rarefaction curves (Fig.

S1), the sequencing captured most prokaryotic diversity in these 134 samples. The

median number of reads in the remaining samples was 1354.5. The microbial

communities on *Alpinia japonica* flowers were by far dominated by the bacterial

phylum Proteobacteria, which accounted for 47.8 % in the samples on average (Fig. 2).

At the class level, Gammaproteobaceteria was the most dominant (16.5 %), followed by

Alphaproteobacteria (6.1 %) and Betaproteobacteria (5.8 %). The Shannon diversity

index of the samples varied among samples, and the majority fell between 1.5 and 3.5

(Fig. S2). The diversity index was significantly different among the treatments, and was

the lowest in the paper bag (PB) treatment (GLMM, $\chi^2 = 22.23$, $p = 0.0005$).

The 36 ASVs that appeared in 20 or more samples were identified as bacteria,

and accounted for 75.7 % of the prokaryotic sequences. NMDS plots of the microbial

communities revealed that the flowers under different treatments in each plant

individual had different microbial communities, while the same treatment on different

plant individuals did not always result in microbial communities with similar

compositions. For example, the OT flowers are plotted on the right of the flower bud

samples in the plant N1, while the OT flowers are on the left side of the buds in N4 and

N5 (Fig. S3a). Only MI treatment resulted in the conversion of the prokaryotic

communities. PERMANOVA indicated that the communities were significantly

different among individuals ($F = 4.89$, $R^2 = 0.100$, $p < 0.0001$) and among the

treatments within each individual ($F = 2.85$, $R^2 = 0.366$, $p < 0.0001$). Variation in the

microbial communities within a treatment was significantly different among the

treatments (ANOVA, df = 5, $F = 8.60$, $p < 0.0001$), and it was the lowest in the MI

treatment, as suggested from the NMDS plots (Fig. S3b).

The cluster analyses grouped the 36 ASVs into 11 clusters (Fig. 3, Table S2).

The first and second most dominant clusters of the 11 include 4 and 6 ASVs and

accounted for 33.0 % and 30.6 % of the sequence reads of the 36 ASVs, respectively

(Table S2). Cluster 6, which accounted for the second highest proportion, showed the

largest difference in the proportion between the MI and WI treatments, followed by

Cluster 2, the most dominant cluster (Table S2, Fig. S4). The Wilcoxon rank-sum test

showed that the difference between the treatments was the most significant for Cluster 6

(w = 444, p < 0.0001 for Cluster 6), while Clusters 2 and 5 also showed significant

differences (w = 78.0, p = 0.0001 for Cluster 2 and w = 127.0, p = 0.0071 for Cluster 5).

The Claident pipeline identified the most abundant 4 ASVs out of the six in

Cluster 6 as belonging to *Pseudomonas* (T00005, T00006 and T0007) and *Erwinia*

(T0004). The three *Pseudomonas* ASV sequences were different for five to eight bases

among 253 (2.0–3.0 %). Blast search further identified the other two ASVs (T00028

and T00031) in the dataset as *Luteibacter* (Rhodanobacteraceae, Gammaproteobacteria)

and *Massilia* (Oxalobacteraceae, Betaproteobacteria) (Table 1).

3.2 Fruit set and seed production

The probability of fruit set of OT predicted by the first GLMM with the

treatment as the fixed factor was 57.3 %, and that of MI was the lowest (34.7 %) (Fig.

3b). Those of the other treatments were between these two (Fig. 3b). On the other hand,

the proportion of Cluster 6 showed the opposite patterns; it was the lowest in OT and

highest in MI. The GLMM including the proportion of Cluster 6 as the fixed factor

indicated that Cluster 6 was significantly associated with low fruit set (GLMM, z = –

2.21, p = 0.0271).

Each fruit that remained on the plants under the treatments had 12–24 ovules,

55.6 ± 20 % of which had developed. In contrast to fruit set, seed set was not

significantly associated with the microbial composition (GLMM, z = 0.685, p > 0.05).

4 DISCUSSION

4.1 Prokaryotic community on *A. japonica* flowers

Alpinia flowers harbored diverse prokaryote communities comparable with those

on flowers of other plant species reported so far (Berg et al. 2017; Müller et al. 2016;

Qian et al. 2021; Wei and Ashman 2018). They are dominated by the phylum

Proteobacteria, most of which belonged to Gammaproteobactera and

Alphaproteobacteria. *Erwinia* and *Pseudomonas* of the Gammaproteobactera are among

the most common constituents of the anthosphere (Vannette 2020).

Though these major taxa were shared among the samples, prokaryotic

community compositions were significantly different among these plant individuals.

The differences already existed in the buds prior to anthesis. Since flower microbes are

often shared by leaves of the same plant (Junker et al. 2011; Massoni et al. 2020; Wei

and Ashman 2018), the differences might have originated from vegetative parts of the

same individuals. The effects of most treatments were not consistent across the

individuals (Fig. S3). Although the paper bag treatment decreased the Shannon diversity

index (Fig. S2), we did not find a drastic increase of particular ASVs by this treatment

(Fig. 3). Growth of a small number of bacteria species in the bag with high humidity

and low UV-irradiance might have reduced the diversity (Hayes et al. 2021).

Prokaryotic diversity on open and pollinator-excluded net-bagged flowers was not

drastically different from that on the flower buds (Fig. S2). It has been reported that

flower visitors disperse microbes among flowers (Ushio et al. 2015), and often increase

microbial diversity (Allard et al. 2018). Flower visitors might not have had large effects

in our system, however, partly due to the relatively low flower visitor activities during

the year of this study (N. E. Jimenez and S. Sakai, personal observation).

4.2 Identification of bacteria that were increased by inoculation

We identified a group of ASVs that showed substantial changes after inoculation

of old-flower microbes. The cluster analyses grouped the ASVs into 11 clusters based

on the co-occurrence among the samples. The microbial inoculation treatment

drastically increased Cluster 6 (Figs. 3 and 4), which consists of 6 ASVs. Among the

ASVs, four ASVs, one *Erwinia* (T00004) and three *Pseudomonas* (T00005, T00006 and

T00007) ASVs accounted for 90 % of the reads of the cluster on average (Table 1).

Microbial communities on old flowers might be dominated by these taxa. The bacteria

may have already colonized before anthesis, because these bacteria were also found in

flower buds at lower abundances as well as in the paper- and net- bagged flowers. The

proportions of Cluster 6 were also moderately higher in the bagged and water-sprayed

flowers than the proportion under the open control treatment (Fig. S3b). Higher

humidity due to these treatments might have enhanced growth of these bacteria. In

agriculture, it has often been reported that high humidity or precipitation enhances

activities of pathogenic *Erwinia* and *Pseudomonas*, and increases damage by diseases

(Billing 1980; Llontop et al. 2020; Pietrarelli et al. 2006).

Though the number of studies that investigate microbial communities on flowers

is rapidly increasing, no study has examined microbial communities on old flowers as

far as we know. Our results suggested that old flowers *A. japonicus* flowers in our field

site were dominated by bacteria of *Erwinia* and *Pseudomonas*. Many bacterial species

of these genera are frequently found both on flowers and inside the plant and/or on

leaves (Vannette 2020). Interestingly, we found all the proteobacterial ASVs in Cluster

6 from flowers of an unrelated dioecious tree, *Mallotus japonicus* (Euphorbiaceae)*.* In

particular, T0007 was the most abundant and frequent ASV on male flowers of the plant

and was present in 72.3 % of male flower samples (Marre et al. 2021 (T0007 is referred

to as ASV02P1in this article), and the dataset used for the study (DRA013108 in

DDBJ)). Therefore, we consider that the bacteria of Group 6 may be generalist

inhabitants on plants, which proliferate on senesced plant parts. It is not uncommon that

flowers of unrelated plant species in the same habitats share flower bacteria species

(Massoni et al. 2020; Qian et al. 2021). It should be noted, however, that ASVs in this

study are distinguished based on relatively short DNA sequences of 253 base pairs, as is

often the case in most anthosphere and phylosphere studies. Besides, we have analyzed

the anthosphere of only two plant species so far. To confirm the generalist habit of the

bacteria, more detailed examination such as sequencing of additional regions and genes,

more extensive sampling and culture isolation would be necessary.

4.3 Effects of old-flower microbes on fruit and seed production

Do the microbes that become dominant on old flowers negatively affect

reproductive success of the plant? We observed the lowest fruit set in the microbial

inoculated treatment, which recorded the highest abundance of Cluster 6. Besides, we

found significant negative associations between the proportion of the ASVs of Cluster 6

and fruit set based on the GLMM analysis, while we did not find such associations for

seed set. Since *Erwinia* and *Pseudomonas* include plant pathogens that trigger abortion

of flowers and fruits (Goumas et al. 1999; Llontop et al. 2020; Marre et al. 2022; Young

1988), some of these bacteria might have negatively affected fruit development by

infecting a reproductive organ. However, we cannot rule out other potential

explanations about the correlation. Fungi, rather than bacteria, that show similar

changes among the treatments with Cluster 6 might have been the cause of the low fruit

set. Alternatively, bacteria and/or fungi dominant on old flowers might have indirectly

decreased fruit set by changing the microenvironment of flowers. Some flower microbes

are known to inhibit pollen germination or pollen tube growth (Christensen et al. 2021;

Eisikowitch et al. 1990), while the effects of microbes on other post-pollination

processes are largely understudied (Cullen et al. 2021). On the other hand, it is unlikely

that pollinator behavioral response affected the fruit set as reported in previous studies

(Sobhy et al. 2018; Vannette et al. 2013), because we hand-pollinated the flowers under

the fruit set monitoring.

5. CONCLUSION

Compared with other floral traits, lifespan of flowers has received surprisingly

little attention. Some previous studies elegantly explained variation among species

based on the balance of the production and maintenance cost and benefit in terms of

reproductive outputs (Ashman 2004). However, it has rarely been argued why plants so

quickly replace cheap, fragile flowers during the flowering period rather than

maintaining expensive but durable flowers. Short lifespan of the flower may be a

defense measure of the plant to escape from negative effects of the microbes that

proliferate on flowers during anthesis, though these microbes may not cause apparent

symptoms. It is increasingly recognized that flowers have characteristic microbial

communities, but the effects and implications of these communities have just started to

be explored and our knowledge about them is still very fragmentary (Cullen et al. 2021;

Rebolleda-Gómez et al. 2019; Rowe et al. 2020; Vannette 2020). Future studies will

elucidate the impacts of microbes on plant reproductive ecology deeply embedded in the

evolution of angiosperms (Rebolleda-Gómez et al. 2019).

REFERENCES

Allard, S. M., Ottesen, A. R., Brown, E. W., & Micallef, S. A. (2018). Insect exclusion
limits variation in bacterial microbiomes of tomato flowers and fruit. *Journal of*
Applied Microbiology, *125*(6), 1749–1760. https://doi.org/10.1111/jam.14087

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local
alignment search tool. *Journal of molecular biology*, *215*(3), 403–410.

Asakawa, Y., Ludwiczuk, A., Sakurai, K., Tomiyama, K., Kawakami, Y., & Yaguchi,
Y. (2017). Comparative study on volatile compounds of *Alpinia japonica* and
Elettaria cardamomum. *Journal of Oleo Science*, *66*(8), 871–876.
https://doi.org/10.5650/jos.ess17048

Ashman, T.-L. (2004). 24 - Flower Longevity. In L. D. B. T.-P. C. D. P. Noodén (Ed.),
(pp. 349–362). San Diego: Academic Press.
https://doi.org/https://doi.org/10.1016/B978-012520915-1/50027-8

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects
models using lme4. *Journal of Statistical Software*, *67*(1), 1–48.
https://doi.org/10.18637/jss.v067.i01

Bawa, K. S. (1990). Plant-pollinator interactions in tropical rain forests. *Annual Review*
of Ecology and Systematics, *21*(1), 399–422.
https://doi.org/10.1146/annurev.es.21.110190.002151

Berg, G., Köberl, M., Rybakova, D., Müller, H., Grosch, R., & Smalla, K. (2017). Plant
microbial diversity is suggested as the key to future biocontrol and health trends.
FEMS Microbiology Ecology, *93*(5), 1–9. https://doi.org/10.1093/femsec/fix050

Billing, E. (1980). Fireblight *Erwinia amylovora* and weather: a comparison of warning
systems. *Annals of Applied Biology*, *95*(3), 365–377.
https://doi.org/10.1111/j.1744-7348.1980.tb04756.x

Bubán, T., Orosz-Kovács, Z., & Farkas, Á. (2003). The nectary as the primary site of
infection by Erwinia amylovora (Burr.) Winslow et al.: A mini review. *Plant*
Systematics and Evolution, *238*(1–4), 183–194. https://doi.org/10.1007/s00606-
002-0266-1

Burdon, R. C. F., Junker, R. R., Scofield, D. G., & Parachnowitsch, A. L. (2018).
Bacteria colonising *Penstemon digitalis* show volatile and tissue-specific responses
to a natural concentration range of the floral volatile linalool. *Chemoecology*,
28(1), 11–19. https://doi.org/10.1007/s00049-018-0252-x

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., &

Holmes, S. P. (2016). DADA2: High-resolution sample inference from Illumina

amplicon data. *Nature Methods*, *13*(7), 581–583.

https://doi.org/10.1038/nmeth.3869

Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Lozupone, C. A.,

Turnbaugh, P. J., et al. (2011). Global patterns of 16S rRNA diversity at a depth of

millions of sequences per sample. *Proceedings of the National Academy of*

Sciences of the United States of America, *108*(SUPPL. 1), 4516–4522.

https://doi.org/10.1073/pnas.1000080107

Christensen, S. M., Munkres, I., & Vannette, R. (2021). Nectar bacteria stimulate pollen

germination and bursting to enhance their fitness. *bioRxiv*.

https://doi.org/https://doi.org/10.1101/2021.01.07.425766

Cullen, N. P., Fetters, A. M., & Ashman, T. L. (2021). Integrating microbes into

pollination. *Current Opinion in Insect Science*, *44*, 48–54.

https://doi.org/10.1016/j.cois.2020.11.002

Davis, N. M., Proctor, Di. M., Holmes, S. P., Relman, D. A., & Callahan, B. J. (2018).

Simple statistical identification and removal of contaminant sequences in marker-

gene and metagenomics data. *Microbiome*, *6*(1), 1–14.

https://doi.org/10.1186/s40168-018-0605-2

Dobson, H. E. M., & Bergstrom, G. (2000). The ecology and evolution of pollen odors.

Plant Syst. Evol, *222*, 63–87.

Eisikowitch, D., Lachance, M. A., Kevan, P. G., Willis, S., & Collins-Thompson, D. L.

(1990). The effect of the natural assemblage of microorganisms and selected

strains of the yeast *Metschnikowia reukaufii* in controlling the germination of

pollen of the common milkweed *Asclepias syriaca*. *Canadian Journal of Botany*,

68(5), 1163–1165. https://doi.org/10.1139/b90-147

Gaube, P., Junker, R. R., & Keller, A. (2021). Changes amid constancy: Flower and leaf

microbiomes along land use gradients and between bioregions. *Basic and Applied*

Ecology, *50*, 1–15. https://doi.org/10.1016/j.baae.2020.10.003

Goumas, D. E., Malathrakis, N. E., & Chatzaki, A. K. (1999). Characterization of

Pseudomonas viridiflava associated with a new symptom on tomato fruit.

European Journal of Plant Pathology, *105*(9), 927–932.

https://doi.org/10.1023/A:1008725818334

Gu, Z., Eils, R., & Schlesner, M. (2016). Complex heatmaps reveal patterns and

correlations in multidimensional genomic data. *Bioinformatics*, *32*(18), 2847–

2849.

Hamady, M., Walker, J. J., Harris, J. K., Gold, N. J., & Knight, R. (2008). Error-

correcting barcoded primers for pyrosequencing hundreds of samples in multiplex.

Nature methods, *5*(3), 235–237.

Hayes, R. A., Rebolleda-Gómez, M., Butela, K., Cabo, L. F., Cullen, N., Kaufmann, N.,

et al. (2021). Spatially explicit depiction of a floral epiphytic bacterial community

reveals role for environmental filtering within petals. *MicrobiologyOpen*, *10*(1), 1–

19. https://doi.org/10.1002/mbo3.1158

Herrera, C. M., Pozo, M. I., & Medrano, M. (2013). Yeasts in nectar of an early-

blooming herb: Sought by bumble bees, detrimental to plant fecundity. *Ecology*,

94(2), 273–279. https://doi.org/10.1890/12-0595.1

Huang, M., Sanchez-Moreiras, A. M., Abel, C., Sohrabi, R., Lee, S., Gershenzon, J., &

Tholl, D. (2012). The major volatile organic compound emitted from *Arabidopsis*

thaliana flowers, the sesquiterpene (E)-β-caryophyllene, is a defense against a

bacterial pathogen. *New Phytologist*, *193*(4), 997–1008.

https://doi.org/10.1111/j.1469-8137.2011.04001.x

Huson, D. H., Auch, A. F., Qi, J., & Schuster, S. C. (2007). MEGAN analysis of

metagenomic data. *Genome Research*, *17*(3), 377–386.

https://doi.org/10.1101/gr.5969107

Junker, R., Loewel, C., Gross, R., Dötterl, S., Keller, A., & Blüthgen, N. (2011).

Composition of epiphytic bacterial communities differs on petals and leaves. *Plant*

Biology, *13*(6), 918–924. https://doi.org/10.1111/j.1438-8677.2011.00454.x

Kaltz, O., & Shykoff, J. A. (2001). Male and female *Silene latifolia* plants differ in per-

contact risk of infection by a sexually transmitted disease. *Journal of Ecology*,

89(1), 99–109. https://doi.org/10.1046/j.1365-2745.2001.00527.x

Llontop, M. E. M., Hurley, K., Tian, L., Bernal Galeano, V. A., Wildschutte, H. K.,

Marine, S. C., et al. (2020). Exploring rain as source of biological control agents

for fire blight on apple. *Frontiers in Microbiology*, *11*(February). https://doi.org/10.3389/fmicb.2020.00199

Lüdecke, D. (2018). ggeffects: Tidy data frames of marginal rffects from regression models. *Journal of Open Source Software*, *3*(26), 772. https://doi.org/10.21105/joss.00772

Maechler, M., Rousseeuw, P., Struyf, A., Hubert, M., & Hornik, K. (2001). Cluster: Cluster analysis basics and extensions. https://cran.r-project.org/package=cluster

Marre, M., Ushio, M., & Sakai, S. (2021). Contrasting microbial communities on male and female flowers of a dioecious plant, Mallotus japonicus (Euphorbiaceae). *Environmental DNA*, (November), 1–15. https://doi.org/10.1002/edn3.271

Marre, M., Ushio, M., & Shoko, S. (2022). The effects of the floral infection by a bacterial pathogen in a dioecious plant , Mallotus japonicus (Euphorbiaceae), 1–9. https://doi.org/10.1002/1438-390X.12110

Massoni, J., Bortfeld-Miller, M., Jardillier, L., Salazar, G., Sunagawa, S., & Vorholt, J. A. (2020). Consistent host and organ occupancy of phyllosphere bacteria in a community of wild herbaceous plant species. *ISME Journal*, *14*(1), 245–258. https://doi.org/10.1038/s41396-019-0531-8

McInnis, S. M., Emery, D. C., Porter, R., Desikan, R., Hancock, J. T., & Hiscock, S. J. (2006). The role of stigma peroxidases in flowering plants: Insights from further characterization of a stigma-specific peroxidase (SSP) from Senecio squalidus (Asteraceae). *Journal of Experimental Botany*, *57*(8), 1835–1846. https://doi.org/10.1093/jxb/erj182

McMurdie, P. J., & Holmes, S. (2013). Phyloseq: An R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS ONE*, *8*(4). https://doi.org/10.1371/journal.pone.0061217

Müller, D. B., Vogel, C., Bai, Y., & Vorholt, J. A. (2016). The plant microbiota: Systems-level insights and perspectives. *Annual Review of Genetics*, *50*(1), 211– 234. https://doi.org/10.1146/annurev-genet-120215-034952

Oksanen, J., Blanchet, F., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B., et al. (2010). Community ecology package: vegan. *R: A Language and Environment for Statistical Computing*. https://cran.r-project.org/web/packages/vegan/index.html

Palmer-Young, E. C., Farrell, I. W., Adler, L. S., Milano, N. J., Egan, P. A., Junker, R.
R., et al. (2019). Chemistry of floral rewards: intra- and interspecific variability of
nectar and pollen secondary metabolites across taxa. *Ecological Monographs*,
89(1), 1–23. https://doi.org/10.1002/ecm.1335

Pietrarelli, L., Balestra, G. M., & Varvaro, L. (2006). Effects of simulated rain on
Pseudomonas syringae pv. *tomato* populations on tomato plants. *Journal of Plant*
Pathology, *88*, 245–251.

Primack, R. B. (1985). Longevity of individual flowers. *Annual Review of Ecology and*
Systematics, *16*, 15–37.

Qian, X., Li, X., Li, H., & Zhang, D. (2021). Floral fungal-bacterial community
structure and co-occurrence patterns in four sympatric island plant species. *Fungal*
Biology, *125*(1), 49–61. https://doi.org/10.1016/j.funbio.2020.10.004

R Core Team. (2016). R: A language and environment for statistical computing. *R*
Foundation for Statistical Computing. https://doi.org/10.1007/978-3-540-74686-7

Rebolleda-Gómez, M., Forrester, N. J., Russell, A. L., Wei, N., Fetters, A. M.,
Stephens, J. D., & Ashman, T. (2019). Gazing into the anthosphere: Considering
how microbes influence floral evolution. *New Phytologist*.
https://doi.org/10.1111/nph.16137

Rogers, H. J. (2006). Programmed cell death in floral organs: How and why do flowers
die? *Annals of Botany*, *97*(3), 309–315. https://doi.org/10.1093/aob/mcj051

Rousseeuw, P. J. (1987). Silhouettes: A graphical aid to the interpretation and validation
of cluster analysis. *Journal of Computational and Applied Mathematics*, *20*(C),
53–65. https://doi.org/10.1016/0377-0427(87)90125-7

Rowe, M., Veerus, L., Trosvik, P., Buckling, A., & Pizzari, T. (2020). The reproductive
microbiome: An emerging driver of sexual selection, sexual conflict, mating
systems, and reproductive isolation. *Trends in Ecology and Evolution*, *35*(3), 220–
234. https://doi.org/10.1016/j.tree.2019.11.004

Roy, R., Schmitt, A. J., Thomas, J. B., & Carter, C. J. (2017). Review: Nectar biology:
From molecules to ecosystems. *Plant Science*, *262*(May), 148–164.
https://doi.org/10.1016/j.plantsci.2017.04.012

Shykoff, J. A., Bucheli, E., & Kaltz, O. (1996). Flower lifespan and disease risk.
Nature, *379*(6568), 779.

Sobhy, I. S., Baets, D., Goelen, T., Herrera-Malaver, B., Bosmans, L., Van den Ende,

W., et al. (2018). Sweet scents: Nectar specialist yeasts enhance nectar attraction of

a generalist aphid parasitoid without affecting survival. *Frontiers in Plant Science*,

9(June). https://doi.org/10.3389/fpls.2018.01009

Tanabe, A. S., & Toju, H. (2013). Two new computational methods for universal DNA

barcoding: A benchmark using barcode sequences of bacteria, archaea, animals,

fungi, and land plants. *PLoS ONE*, *8*(10).

https://doi.org/10.1371/journal.pone.0076910

Taylor, L. P., & Hepler, P. K. (1997). Pollen germination and tube growth. *Annual*

Review of Plant Biology, *48*(July), 461–491.

https://doi.org/10.1146/annurev.arplant.48.1.461

Ushio, M. (2019). Use of a filter cartridge combined with intra-cartridge bead-beating

improves detection of microbial DNA from water samples. *Methods in Ecology*

and Evolution, *10*(8), 1142–1156. https://doi.org/10.1111/2041-210X.13204

Ushio, M., Yamasaki, E., Takasu, H., Nagano, A. J., Fujinaga, S., Honjo, M. N., et al.

(2015). Microbial communities on flower surfaces act as signatures of pollinator

visitation. *Scientific Reports*, *5*, 1–7. https://doi.org/10.1038/srep08695

van Vuuren, S. F., & Viljoen, A. M. (2007). Antimicrobial activity of limonene

enantiomers and 1,8-cineole alone and in combination, *22*, 540–544.

https://doi.org/10.1002/ffj.1843

Vanneste, J. L. (2000). *Fire blight: the disease and its causative agent,* Erwinia

amylovora. CABI.

Vannette, R. L. (2020). The floral microbiome: Plant, pollinator, and microbial

perspectives. *Annual Review of Ecology, Evolution, and Systematics*, *51*, 363–386.

https://doi.org/10.1146/annurev-ecolsys-011720-013401

Vannette, R. L., Gauthier, M. P. L., & Fukami, T. (2013). Nectar bacteria, but not yeast,

weaken a plant– pollinator mutualism. *Proceedings of the Royal Society B:*

Biological Sciences, *280*(1752). https://doi.org/10.1098/rspb.2012.2601

Wei, N., & Ashman, T.-L. L. (2018). The effects of host species and sexual dimorphism

differ among root, leaf and flower microbiomes of wild strawberries in situ.

Scientific Reports, *8*(1), 1–12. https://doi.org/10.1038/s41598-018-23518-9

Young, J. M. (1988). Bacterial blight of kiwifruit in New Zealand. *EPPO Bulletin*,

18(1), 131–140. https://doi.org/10.1111/j.1365-2338.1988.tb00358.x

TABLE 1. The 6 ASVs in Cluster 6. Taxonomic information was based on Claident (shown by normal face) supplemented by BLAST

search (shown by bold face with percent of identical bases). The results of BLAST search of the ASVs in Cluster 6, and their percent of

matches.

ASV	Phylum	Class	Family	Genus	Percent identity[1]	Average proportion among samples[2]
T0004	Proteobacteria	Gammaproteobacteria	**Erwiniaceae**	*Erwinia*	100.0	0.086
T0005	Proteobacteria	Gammaproteobacteria	Pseudomonadaceae	*Pseudomonas*		0.062
T0006	Proteobacteria	Gammaproteobacteria	Pseudomonadaceae	*Pseudomonas*		0.040
T0007	Proteobacteria	Gammaproteobacteria	Pseudomonadaceae	*Pseudomonas*		0.086
T0028	Proteobacteria	**Betaproteobacteria**	**Oxalobacteraceae**	*Massilia*	99.2	0.017
T0031	Proteobacteria	Gammaproteobacteria	Rhodanobacteraceae	*Luteibacter*	99.2	0.015

Note: 1, The highest percentage of the identical bases that resulted from the BLAST search; it is shown only for the ASVs that were

subjected to the BLAST search due to incomplete taxonomic assignments by Claident. 2, Average proportions of the ASV of 36 prevalent

ASVs used for the heatmap in Figure 3.

FIGURE 1. Experimental design used in this study. a, An open flower and flower buds

on an inflorescence of *Alpinia japonica*. b, Floral parts of *Alpinia japonica* in open state

(left) and bud state (right). c, Experimental and sampling procedures.

FIGURE 2. Composition of prokaryotic communities on the flowers of *Alpinia*

japonica. Proportions in the samples of the same treatment in the same individuals were

averaged. See Table S1 for sample sizes.

FIGURE 3. Heatmap of the prokaryotic communities of flower buds and flowers under

different treatments. Samples were vertically arranged and grouped by the treatments as

shown on the right. ASVs were clustered hierarchically based on Bray–Curtis

dissimilarity using Ward's minimum variance algorithm. The class of each ASV is

shown by the color below the tree. The color intensity in each panel shows the

proportion in a sample, referring to the color key on the left.

FIGURE 4. Association between the proportion of Cluster 6 and predicted probability

of fruit set. (a) Variation in the total proportions of the bacteria among the treatments.

(b) The predicted probabilities of fruit set 6 weeks after flowering under the different

treatments by the GLMM with the treatment as the fixed factor. The thick lines and the

boxes indicate the predicted values and their 95 % confidential intervals. Distribution of

flowers that set fruits (top) or were aborted (bottom) are shown by dots. Colors indicate

the five individual plants).

684 TABLE S1. The number of the samples used to monitor fruit set for the five treatments

685 and to analyze microbial communities.

Treatments	Fruit set and seed production					Microbial community				
	N1	N2	N4	N5	N6	N1	N2	N4	N5	N6
Bud	–	–	–	–	–	5	4	4	5	4
Open	10	26	14	16	19	5	5	5	5	5
Net bag	9	17	15	17	10	5	4	4	4	4
Paper bag	7	9	5	7	9	5	5	2	5	5
Microbe	6	11	5	7	5	3	5	4	5	4
Water	4	3	6	7	7	5	4	4	5	5

TABLE S2. Characteristics of the 11 clusters. The number of ASVs included, the
median proportion in the samples of the microbial inoculated (MI) and water sprayed
(WI) treatments, and p values of Wilcoxon rank sum tests in the comparisons of the
proportions between MI and WI are shown. The asterisks indicate that the proportions
are significantly different between the treatments (*, p < 0.01;**, p < 0.001; ***, p <
0.0001). Cluster 6, which shows significantly higher proportions in MI than in WI, is
highlighted with boldface type.

Cluster ID	No. ASVs included	Median of proportions		Wilcoxon rank sum test (MI vs. WI)	
		Microbe (MI)	Water (WI)	w	p-values
Cluster 1	2	0.000	0.000	186.0	0.1469
Cluster 2	4	0.038	0.190	78.0	0.0001**
Cluster 3	2	0.000	0.000	267.5	0.4143
Cluster 4	2	0.000	0.000	225.5	0.6198
Cluster 5	4	0.004	0.011	127.0	0.0071*
Cluster 6	**6**	**0.845**	**0.213**	**444.0**	**< 0.0001***
Cluster 7	4	0.023	0.036	204.0	0.3826
Cluster 8	4	0.014	0.097	162.5	0.0635
Cluster 9	2	0.000	0.000	224.0	0.6204
Cluster 10	3	0.000	0.000	213.0	0.4531
Cluster 11	3	0.000	0.000	276.0	0.3623

FIGURE S1. Rarefaction curves of the samples. The vertical axis shows the number of observed prokaryotic ASVs. The number of

sequences per sample is shown on the horizontal axis. The three panels show the same graph with different ranges of the horizontal axis.

FIGURE S2. Variation in the prokaryotic diversity among the treatments and samples.

The diversity was significantly different among the treatments (GLMM, $\chi^2 = 22.234$, p

$= 0.0005$). Colors indicate the five individual plants.

FIGURE S3. Variation of the prokaryotic communities on the flowers under different

treatments. (a) NMDS plots of the microbial communities of flower buds and flowers

under different treatments. In the five panels, all microbial samples are plotted with grey

circles, and samples from each plant are overlayed individually. The five treatments are

indicated by different colors. (b) Distribution of the distances from the treatment

centroid. Variation in the microbial communities within a treatment was significantly

different among the treatments (ANOVA, df = 5, F = 8.60, p < 0.0001). Colors indicate

the five individual plants.

FIGURE S4. Variation in the total proportions of the 10 ASV clusters among the flower bud and flowers under the five treatments. Colors
indicate the five individual plants.