1   Why are flowers ephemeral? Effects of old flower microbes on fruit set in a wild ginger
2   with one-day flowers, *Alpinia japonica* (Zingiberaceae)
3

4   Nuria Jiménez Elvira[1], Masayuki Ushio[2,3], Shoko Sakai[3]
5
6   [1]Centre de Recerca Ecològica i Aplicacions Forestals, 08193 Cerdanyola del Vallès,
7   Barcelona, Spain
8   [2]Hakubi Center, Kyoto University, Kyoto University, Kyoto 606-8501, Japan
9   [3]Center for Ecological Research, Kyoto University, Hirano 2-509-3, Otsu 520-2113,
10   Japan

11
12

Abstract: Flowers are colonized by and inhabited by diverse microbes. Plants rapidly
replace flowers of short lifespan, and old flowers senesce. This may contribute to
avoiding adverse effects of the microbes. In this study, we investigate if the flower
microbial community on old flowers impedes fruit and seed production in a wild ginger
with one-day flowers. We inoculated newly opened flowers with old flower microbes,
and monitored the effects on fruit and seed set. We also assessed prokaryotic
communities on the flowers using amplicon sequencing. We found five bacterial
amplicon sequence variants (ASVs) whose proportions were increased on the inoculated
flowers. These ASVs were also found on flower buds and flowers that were bagged by
net or paper during anthesis. Fruit set was negatively associated with the proportions of
these ASVs, while seed set was not. The results suggest that old flowers harbor
microbial communities different from those at anthesis, and that the microbes abundant
on old flowers negatively affect plant reproduction. Though the short lifespan of flowers
has gotten little attention, it might be an essential defense mechanism to cope with
antagonistic microbes that rapidly proliferate on the flowers.

Keywords: flower longevity, anthosphere, pollination, microbiome

INTRODUCTON

Flowers are an elaborated device to exchange pollen among conspecific individuals. Instead of moving around to find a potential mate, the plants disperse and receive pollen relying on different vectors such as insects and wind to achieve sexual reproduction. As a consequence, flowers are also exposed to microbial colonization under natural conditions [1–3]. Flowers have various habitats for microbes. Flowers attract pollinators by secreting nectar, which is rich in sugars and often contains other nutrients, such as amino acids and lipids [4]. The stigma is a germination bed for pollen grains connected to a growth chamber for pollen tubes. It maintains humidity and nutrients necessary for pollen tube growth [5], which would also be beneficial for microbes. Recent studies using high-throughput sequencing have revealed highly diverse microbial communities on flowers of many species in a wide range of habitats [6–8].

Today, flower microbes are increasingly recognized as essential components in the ecology and evolution of plant reproduction (reviewed in [9,10]). Not surprisingly, some flower microbes are known to have negative effects on the survival and reproduction of the host plant. Many pathogens use flowers, especially the stigma and nectary, as a point of entry. Their infection of flowers often results in fruit abortion and systemic infection [11,12]. Other microbes impede various processes of pollination and fertilization. Nectar yeasts are reported to modify pollinator foraging patterns and reduce pollination success [13]. *Acinetobacter* bacteria that commonly inhabit floral nectar exploit pollen nutrition by inducing pollen germination and bursting [14]. Most of these studies examined the effects of an individual microbial species or strains separately rather than that of the whole community.

In general, even newly opened flowers already have microbes at least on some flower tissues. Their abundance increases over time on individual flowers [10]. Flowers have various mechanisms to suppress microbial growth, such as flower volatiles [3,15], reactive oxygen [16] and secondary compounds [17,18]. Besides, an extremely short lifespan of the flowers from one to several days [19], especially in hot habitats [20], is also understood as a compromise to efficiently achieve sexual reproduction while avoiding adverse effects of flower antagonists, including microbes [1,21,22]. Unlike in mammals, flowers, reproductive structures of the plants, are only retained while they are

needed, and are developed *de novo* every season in perennial species [22]. We do not

know yet, however, to what extent microbes that are suppressed by the short lifespan of

flowers are potentially detrimental to plant fecundity.

In this study, we examine if the flower microbial community on old flowers

impedes fruit and seed production in a wild ginger, *Alpinia japonica* (Zingiberaceae),

that produces short-lifespan flowers. This perennial herb grows on the humid forest

floor of evergreen and deciduous forests in western Japan. It flowers in June, when the

climate is hot and humid. Its flowers open in the morning and wilt at sunset within the

same day. Therefore, its flowering matches with the most ideal season for microbial

growth, while it may avoid the negative effects of microbes by renewing flowers every

day. To imitate microbial communities that would appear on the flowers of *A. japonica*

if their longevity were longer, we inoculate microbes from old flowers onto newly

opened flowers in the morning. Firstly, we characterize the prokaryote communities on

A. japonica flowers. Secondly, we identify prokaryotes that significantly increased after

inoculation by comparing prokaryotic communities between the inoculated flowers and

open flower controls. In addition, we examine if these prokaryotes colonize by air or by

flower visitors after flowering, or if they had already existed before anthesis. We

covered the flowers by net or paper bags to exclude microbial colonization by flower

visitors or by both flower visitors and air, respectively. If the old-flower microbes

colonize only after anthesis, bagging would reduce the occurrences of these microbes.

Lastly, we observe fruit and seed sets of the flowers under the treatments. We expected

that the flowers that had more old-flower microbes would show lower fruit and seed set

if antagonistic microbes are abundant on old flowers.

MATERIALS AND METHODS

Study species

Alpinia japonica (Zingiberaceae) is a wild ginger distributed in temperate and

subtropical regions of eastern Asia. It is a perennial herb 0.5–0.7 m in height occurring

mostly in broad-leaved evergreen forests. Inflorescences are compound racemose with

10–60 flowers (37 on average in this population). Flowers are zygomorphic with a

prominent labellum 1.1–1.2 cm in length, and white with red stripes at the margin of the

labellum (Fig. 1a). Flowering occurs almost synchronously within a population and lasts

for 10–14 days for each plant between late May and June. From one to five flowers

open per inflorescence. At anthesis around 0700, the dorsal and lateral petals open, and

the labellum inside expands (Fig. 1b). The flowers are known to have volatiles,

including 1, 8-cineole, with antimicrobial activities [23,24]. The corolla wilts at sunset

within the day. The ovary matures into an ellipsoid, three-loculate fleshy capsule ~2

months after flowering. The species is self-compatible, and pollinated by bees, mainly

carpenter bees (*Xylocopa appendiculata*) and bumble bees (*Bombus ardens*) (J. Nuria

and S. Sakai, unpublished data).

Study site

Studies were conducted in May to July of 2018, in Seta Park, Otsu, Shiga

Prefecture, Japan (34°50'N, 135°50'E). Seta Park has a young secondary deciduous

forest where streams are abundant, which provides an optimal habitat for *A. japonica*.

The annual mean temperature is 14.9°C, and annual total rainfall is ~1530 mm in Otsu

city (Japan Meteorological Agency, https://www.data.jma.go.jp/). Mean daily maximum

temperatures in May, June and July 2018 were 24.0°C, 27.3°C and 33.7°C, respectively.

Flower treatments

Alpinia japonica typically presents 2–12 inflorescences per plant at the study

site. We selected six plants with at least eight inflorescences with more than 10 flowers

on each. Unfortunately, one of them had been damaged, probably by park visitors,

before the experiments and sampling. Therefore, we used the remaining five plants

(Plant ID = N1, N2, N4, N5, N6). In each target plant, eight inflorescences were tagged

and covered with a fine net (Cloth Cabin, Suminoe Teijin Techno, Osaka, Japan) prior

to flowering.

We applied one of five treatments to part of the flowers that opened on the

inflorescences between 22nd May and 7th June. The flowers under these treatments

were used for either identification of microbes on their flowers or for observation of

fruit set (Fig. 1c). We could not use the same flower for both, because we should collect

the flower on the same day for the former, while we should leave the flower intact for

the latter. The five treatments were as follows: 1) open control (OT), in which the mesh

bag on the inflorescence was removed during 0700–1700, so that microbes were

transmitted to the flower by both air and pollinators; 2) insect exclusion by a net bag

(NB), in which the inflorescence was left covered with the net, which allowed air but

not insects to get through; 3) paper bagged (PB), in which the mesh bag covering the

inflorescence was replaced by a new paper bag (Grape Bag, DAIICHI VINYL, Fukui,

Japan) during 0700–1700 to exclude air flow and insect visits; 4) open and microbial

inoculation (MI) and 5) open and water sprayed mock (WI) treatments. In MI, the mesh

bag on the inflorescence was removed as in OT, and the flower was sprayed with

microbial solution. During the spray inoculation, the focal flower was isolated from the

rest of the inflorescence by a funnel-shaped plastic sheet to avoid contamination. The

microbial suspension for the inoculation was prepared by soaking old flowers in

distilled water. Nine old flowers that had opened the previous day from 1–3 individuals

were collected from individuals that did not receive the treatments. Microbes were

detached by ultrasonic dispersion in 10 ml of distilled water and kept at 4°C until the

inoculation (maximum of 36 hours). In WI, the mesh bag covering the inflorescence

was removed as in OT, and the flower was sprayed with distilled water as a control for

the physical effect of the MI treatment. We did not conduct the treatments on cloudy or

rainy days. Flowers that opened on the same day on the same inflorescence were

assigned to the same treatments in the case of NB and PB, while OT, MI and WI were

occasionally mixed on an inflorescence. This is because we bagged the entire

inflorescence rather than a single flower to avoid damaging the flowers. The sample size

of each treatment is summarized in Table S1.

Identification of microbial communities

To identify microbial communities, we sampled 4–5 flowers with each treatment

from each plant. The flowers were cropped at ~1700 of the day of flower opening and

separately put into a 5 ml plastic tube after removing the stigma (to use for another

study) and petals (Fig. 1b). In addition, five flower buds were collected from each

individual to identify microbes that were present inside prior to anthesis with the same

procedure. The tube was kept on ice until being brought to the lab within 4 hours after

sampling and kept at –20°C until further processing.

To detach microbes from the flower surface, we added 3 ml of Phosphate-

Buffered Saline solution (PBS buffer, pH 7.4, Nippon Gene Co., Ltd.) to the tube and

sonicated it using an Ultrasonic Disruptor Handy Sonic UR-21P (Digital Biology) at

Level 5 power for 30 sec. Microbes in the PBS solution were then filtered using a

Sterivex Millipore filter (Sterivex-HV 0.22, Merck). Residual PBS on the Sterivex filter

was removed using a Vac-Man® Laboratory Vacuum Manifold (Promega).

Microbial DNA was extracted with DNeasy Blood & Tissue Kits (QIAGEN)

following the manufacturer's instructions. A portion of the 16S small-subunit ribosomal

gene was PCR-amplified using the primer pairs, 515F (5'-GTG YCA GCM GCC GCG

GTA A-3') and 806R (5'-GGA CTA CNV GGG TWT CTA AT-3') [25] with

Nextera DNA index adapters. First PCR was performed in 12 μl reactions, each

containing 1 μl of DNA template, 6 μl of KAPA HiFi HotMasterMix (Kapa

Biosystems) and 0.7 μl of each primer (5 μM). The first PCR was conducted with a

temperature profile of 95°C for 3 min, followed by 35 cycles at 98°C for 20 s, 60°C for

15 s, 72°C for 30 s, and a final extension at 72°C for 5 min, followed by storage at 4°C

until use. Triplicate PCR products were pooled and purified with Agencourt AMPure

XP (Beckman Coulter). P5/P7 Illumina adaptors were then added in a second PCR

using fusion primers with 8-mer index sequences for sample identification [26]

(forward, 5'-AAT GAT ACG GCG ACC ACC GAG ATC TAC AC-[8-mer tag]-TCG

TCG GCA GCG TCA GAT GTG TAT AAG AGA CAG -3'; reverse, 5'-CAA GCA

GAA GAC GGC ATA CGA GAT-[8-mer tag]-GTC TCG TGG GCT CGG AGA TGT

GTA TAA GAG ACA G-3'). This preparation had a total volume of 24 μl with double

the amount of each reagent in the first PCR and 10-times diluted first PCR products as

DNA template. The temperature profile was 95°C for 3 min, followed by 12 cycles at

98°C for 20 s, 68°C for 15 s, 72°C for 15 s, a final extension at 72°C for 5 min, and 4°C

until use. Second PCR products were pooled at equal volumes after a

purification/equalization process with the same AMPure XP protocol and E-Gel

SizeSelect II Agarose Gel kit (Invitrogen). DNA quantitation was performed using a

Qubit® dsDNA HS Assay Kit (Thermo Fisher Scientific Inc). The pooled library was

sequenced on the Illumina Miseq using Reagent Kit V2 (500 cycle Nano Kit).

We processed the sequence data following Ushio [27]. The raw MiSeq data were

converted into FASTQ files using the bcl2fastq program provided by Illumina

(bcl2fastq v2.18). The FASTQ files were then demultiplexed using the command

implemented in Claident (http://www.claid ent.org; [28]). Demultiplexed FASTQ files

were analysed using the ASV method implemented in the package DADA2 [29] of R

[30]. At the quality filtering process, forward and reverse sequences were trimmed at

the length of 215 and 160, respectively, using DADA2::filterAndTrim() function.

Taxonomic identification was performed for ASVs (Amplicon Sequence

Variants; proxy for Operational Taxonomic Units) inferred using DADA2 based on the

query-centric auto-k-nearest-neighbour (QCauto) method [28] and subsequent

taxonomic assignment with the lowest common ancestor algorithm [31] using the

"semiall" database and clidentseq and classigntax commands implemented in claident

v0.2.2018.05.29. The reads identified as Bacteria or Archaea were used for further

analysis. Blast search was also conducted for the most abundant ASVs for which the

above procedure did not provide the identification of the order or lower taxonomic

levels. Top matches of the sequences were referred to in order to assign taxonomic

identities of the ASVs. Taxonomy, ASV table, and sample information were combined

into a single R object for downstream manipulation using the R package phyloseq [32].

We also removed ASVs that only occurred in negative control samples. Furthermore,

potential contaminant ASVs were identified statistically based on prevalence in negative

control samples using the R package decontam [33] based on ASV prevalence and a

threshold of 0.5. This resulted in elimination of 79 of 2679 16S ASVs.

Fruit set and seed production

The treated flowers for the observation of fruit set were hand-pollinated around

1700 using outcrossing pollen to ensure that the fruit set was not limited by pollen.

Pollen was taken with a wood toothpick from a cut flower from a pollen donor plant, of

which inflorescences had been covered with a mesh bag prior to anthesis to prevent

removal of pollen by flower visitors. In total, 251 pollinated flowers were monitored for

fruit and seed set (Table S1). Six weeks after the treatment, we surveyed whether the

fruits under the treatments were still retained on the inflorescence, and collected the

fruits (131 fruits) and counted developing seeds and aborted ovules inside. At that time,

the fruits were still green but fully plump.

Statistical analysis of microbial community data

To examine differences in the prokaryotic diversity among the treatments, we

evaluated the Shannon diversity index of each sample using the

phyloseq::estimate_richness() function. The index did not show a significant association

with the total numbers of the reads (the results not shown). The differences among the

treatments were tested using a nonparametric Kruskal-Wallis test.

We examined variation in the structure of the prokaryotic communities using the

ASVs that were present in 20 or more flower samples (34 ASVs satisfied this criterion).

The frequency of each ASV was divided by the number of the total sequences of the 34

ASVs in each sample to convert it to a proportion. The ASVs that accounted for more

than 5.0 % in the samples (5 ASVs) on average were further identified by matching

with the sequences of the type strains in the NCBI database

(https://blast.ncbi.nlm.nih.gov/). The genus of the strain(s) with the highest similarity

was assigned.

To compare microbial communities among flowers under the different

treatments, we calculated the Bray–Curtis dissimilarity index using the package vegan

[34] in R. We performed a permutational multivariate analysis of variance

(PERMANOVA) using the vegan::adonis function to examine if flowers of the same

individuals nested within under the same treatment have similar microbial communities.

Ordinations were plotted with nonmetric multidimensional scaling (NMDS) using the

vegan::metaMDS procedure. We then calculated the distances of each flower sample

from the centroid of the treatment. Variation in the distances among the treatments were

tested using vegan::anova for betadisper object.

To find ASVs that tend to co-occur in the samples, we conducted hierarchical

clustering of ASVs based on Bray–Curtis dissimilarity using Ward's minimum variance

algorithm using the hclust procedure. The algorithm merges clusters that minimize the

increase in the sum of squared distances from the cluster centroid. We chose the number

of clusters that maximize the silhouette index [35]. The index was calculated using

silhouette in the R package cluster [36]. The proportions of the ASVs in the sample

were visualized by a heatmap created using the R package ComplexHeatmap [37]. We

identified the cluster that showed the largest difference in the median proportion in the

samples under the microbial inoculate (MI) treatment compared with that under the

water sprayed (WI) treatment. Differences in the proportions of the cluster between the

MI and WI treatments were also evaluated by the statistic of the Wilcoxon rank sum

test.

Statistical analysis of fruit set and seed set data

To assess the factors associated with fruit set and seed production, we used

generalized linear mixed models (GLMMs). To evaluate differences in fruit set among

the treatments after controlling the difference among the individuals, we modeled fruit

set (retained/aborted) with the treatments as a fixed factor and the individuals as a

random factor using a binomial error distribution. Then we calculated and plotted

estimated marginal means of fruit set for the treatments using the ggpredict() function of

the R package ggeffects [38]. Then, to test the association of fruit set with microbial

communities, we constructed the second model for fruit set. The model included

microbial compositions as a fixed factor and the individuals as a random factor using a

binomial error distribution. We identified Cluster 5 to include ASVs that showed the

largest increase after the microbial inoculation (see below). Therefore, we used the

average proportion of Cluster 5 of each treatment as the variable for the microbial

composition. We evaluated the significance of the fixed factor by the Wald test.

We also modeled seed set (retained/aborted) and tested its association with

microbial composition. The model included the proportion of Cluster 5 as a fixed factor

and the individuals and fruit as nested random factors. We fitted the model with a

binominal error distribution. The GLMM analyses for fruit and seed sets were

performed using the R package lme4 [39].

RESULTS

Microbial communities

We obtained 804,963 reads of prokaryotic sequences, which were clustered into

2,600 ASVs. Samples without prokaryotic reads were removed from further analysis

(four samples). This reduced the number of samples to 142. According to the rarefaction

curves (Fig. S1), the sequencing captured most prokaryotic diversity. The median

number of reads in the remaining samples was 670. The Shannon diversity index of the

samples varied among samples, and the majorities fell between 2.0 and 3.5 (Fig. S2).

The diversity index was significantly different among the treatments, and was the

lowest in the paper bag (PB) treatment (Kruskal-Wallis χ^2 =18.2, df = 5, p = 0.0054).

The 34 ASVs that appeared in 20 or more samples were identified as bacteria,

and accounted for 75.7 % of the prokaryotic sequences. NMDS plots of the microbial

communities revealed that the flowers under different treatments in each plant

individual had different microbial communities, while the same treatment on different

plant individuals did not always converge the microbial communities into similar

compositions. For example, the OT flowers are plotted on the right of the flower bud

samples in the plant N1, while the OT flowers are on the left side of the buds in N5

(Fig. S3a). Only MI treatment resulted in the conversion of the prokaryotic

communities. PERMANOVA indicated that the communities were significantly

different among individuals (F = 5.52, R^2 = 0.106, p < 0.0001) and among the

treatments within each individual (F = 2.96, R^2 = 0.355, p < 0.0001). Variation in the

microbial communities within a treatment was significantly different among the

treatments (ANOVA, df = 5, F = 13.31, p < 0.0001), and it was the lowest in the MI

treatment, as suggested from the NMDS plots (Fig. S3b).

The cluster analyses grouped the 34 ASVs into 10 clusters (Fig. 2, Table S2).

Cluster 5, which accounts for the second highest proportion, showed the largest

difference in the proportion between the MI and WI treatments, followed by Cluster 1,

the most dominant cluster (Table S2, Fig. S4). The Wilcoxon rank-sum test showed that

the difference between the treatments was significant for both clusters (w = 90, p =

0.0001 for Cluster 1; and w = 501, p < 0.0001 for Cluster 5), while the difference was

not significant in the other clusters.

Blast search identified the two most abundant ASVs (T00020 and T00030) in

the dataset as *Cutibacterium* (Propionibacteriaceae) and *Ochrobactrum* (Brucellaceae)

in Cluster 1 (Table 1, Fig. 2). The third most-abundant ASVs (T00004) showed the

highest matches with sequences of *Erwinia* (Erwiniaceae) and the fourth and fifth

(T00005 and T0000) did with *Pseudomonas* (Pseudomonodaceae) in

Gammaproteobacteria (Table 1). The two *Pseudomonas* ASV sequences were different

for only five bases among 253 (2.0 %), so they may belong to the same species. These

three ASVs were included in Cluster 5 (Fig. 2).

Fruit set and seed production

The probability of fruit set of OT predicted by the first GLMM was 55.7 %, and

that of MI was the lowest (34.7 %) (Fig. 3b). Those of the other treatments were

between these two (Fig. 3b). On the other hand, the proportion of Cluster 5 showed the

opposite patterns; it is the lowest in OT and highest in MI. The GLMM including the

proportion of Cluster 5 as the fixed factor indicated that Cluster 5 was significantly

associated with low fruit set (GLMM, $z = -2.19$, $p = 0.0288$).

Each fruit that remained on the plants under the treatments had 12–24 ovules,

55.6 ± 20 % of which had developed. In contrast to fruit set, seed set was not

significantly associated with the microbial composition (GLMM, $z = 0.685$, $p > 0.05$).

DISCUSSION

Prokaryotic community on *A. japonica* flowers

Alpinia flowers harbored diverse prokaryote communities comparable with those

on flowers of other plant species reported so far [6,40]. ASVs that accounted for more

than 5% on average were *Cutibacterium* (T00020), *Ochrobactrum* (T00030), *Erwinia*

(T0004), and two *Pseudomonas* ASVs (T0005, T0007) (Table 1). The genus

Cutibacterium (Propionibacteriaceae) has rarely been reported as a dominant taxon in a

plant microbiome (but see [41]), and little is known about the role of this genus on

plants. On the other hand, *Ochrobactrum* is often found in the rhizosphere of various

plant species (e.g., [42]). Some strains of the genus have been reported to positively

affect plant survival and growth [43,44]. *Erwinia* and *Pseudomonas* are among the most

common constituents of the anthosphere [10]. They include many well-known plant

pathogens. *Erwinia amylovora* is one of the best-studied bacterial plant pathogens

causing fire blight in apple trees. The pathogen is transmitted among trees by flower

visitor insects as well as wind and rain [12]. *Psudomonas syringae* is also a well-studied

pathogen with many pathovars, which infect different host plants. *Pseudomonas*

syringae pv. *actinidiae*, the causal agent of kiwifruit bacterial canker, is vectored by

flower visitors, resulting in systemic invasion [45].

Though these major taxa were shared among the samples, prokaryotic

community compositions were significantly different among these closely related plant

individuals. The differences already existed in the buds prior to anthesis. Since flower

microbes are often shared by leaves of the same plant [6,7,46], the differences might

have originated from vegetative parts of the same individuals. The effects of most

treatments were not consistent across the individuals (Fig. S3). Although the paper bag

treatment decreased the Shannon diversity index (Fig. S2), we did not find a drastic

increase of particular ASVs by the treatment (Fig. 3). Growth of a small number of

bacteria species in the bag with high humidity and low UV-irradiance might have

reduced the diversity [47]. Prokaryotic diversity on open and pollinator-excluded net-

bagged flowers was more or less similar to that on the flower buds (Fig. S2). It has been

reported that flower visitors disperse microbes among flowers [2], and often increase

microbial diversity [48]. Flower visitors might not have large effects in our system,

however, partly due to the relatively low flower visitor activities during this year (N. E.

Jimenez and S. Sakai, personal observation).

Identification of bacteria that were increased by inoculation

363   We identified a group of ASVs that showed substantial changes after inoculation

of old-flower microbes. The cluster analyses grouped the ASVs into 10 clusters based

on the co-occurrence among the samples. The first and second most dominant clusters

of the 10 accounted for 41.2 % and 29.3 % of the sequence reads of the samples,

respectively, and were significantly different among the treatments. The microbial

inoculation treatment drastically increased Cluster 5 (Fig. 4), which contains three of

the five most dominant ASVs, *Erwinia* (T00004) and two *Pseudomonas* (T00005,

T00007) (Table 1). Microbial communities on old flowers might be dominated by these

taxa. The bacteria may have already colonized before anthesis, because these bacteria

were also found in flower buds at lower abundances. The proportions of Cluster 5 were

also moderately higher in the paper- and net-bagged and water-sprayed flowers than the

proportion under the open control treatment (Fig. S3b). Higher humidity due to these

treatments might have enhanced growth of these bacteria. In agriculture, it has often

been reported that high humidity or precipitation enhance activities of pathogenic

Erwinia and *Pseudomonas*, and increase damage by diseases [49–51].

378   Though the number of studies that investigate microbial communities on flowers

is rapidly increasing, no study has examined microbial communities on old flowers as

far as we know. The bacterial families Erwiniaceae and Pseudomonodaceae probably

dominating on old flowers of *A. japonicus* are among the most dominant taxa on

flowers [10]. Many bacterial species of these families are frequently found both on

flowers and inside the plant and/or on leaves [10]. Besides, flowers of unrelated plant

species in the same habitats often share common flower bacteria species [7,40].

Therefore, we consider that the bacteria of Group 5 may be generalist inhabitants on

living plants and senesced plant parts. However, identification of bacteria in most

microbial community studies of the anthosphere and phyllosphere are based on short

DNA sequences of hundreds of base pairs. To confirm habitat specificity within a plant

and adaptation to a specific host, cell lineage tracking using a bacterial strain with a

marker gene, or genomic comparisons among flower and leaf isolates from the same

and different plant hosts would be necessary [10].

Effects of old-flower microbes on fruit and seed production

Do the microbes that become dominant on old flowers negatively affect

reproductive success of the plant? We observed the lowest fruit set in the microbial

inoculated treatment, which recorded the highest abundance of Cluster 5. Besides, we

found significant negative associations between the proportion of the ASVs of Cluster 5

and fruit set based on the GLMM analysis, while we did not for seed set. Since *Erwinia*

and *Pseudomonas* include plant pathogens that trigger abortion of flowers and fruits

[51–53], some of these bacteria might have negatively affected fruit development by

infecting a reproductive organ. However, we cannot rule out other potential

explanations about the correlation. Fungi, rather than bacteria, that show similar

changes among the treatments with Cluster 5 might be the cause of the low fruit set.

Alternatively, bacteria and/or fungi dominant on old flowers might have indirectly

decreased fruit set by changing the microenvironment of flowers. Some flower microbes

are known to inhibit pollen germination or pollen tube growth [14,54], while the effects

of microbes on other post-pollination processes are largely understudied [55]. On the

other hand, it is unlikely that pollinator behavioral response affected the fruit set as

reported in previous studies [56,57], because we hand-pollinated the flowers under the

fruit set monitoring.

CONCLUSION

Compared with other floral traits, lifespan of flowers has received surprisingly little attention. Some previous studies elegantly explained variation among species based on the balance of the production and maintenance cost and benefit in terms of reproductive outputs [44–46]. However, it has rarely been argued why plants so quickly replace cheap, fragile flowers during the flowering period rather than maintaining expensive but durable flowers [19]. This study is the first to show the potential negative consequence of longer flower lifespan by non-pathogenic (microbes that do not cause apparent symptoms) microbes. It is increasingly recognized that flowers have characteristic microbial communities, but their effects and implications have just started to be explored and our knowledge is still very fragmental [9,10,55,58]. Future studies will elucidate the impacts of microbes on plant reproductive ecology deeply embedded in the evolution of angiosperms [9].

REFERENCES

1. Kaltz O, Shykoff JA. 2001 Male and female *Silene latifolia* plants differ in per-contact risk of infection by a sexually transmitted disease. *J. Ecol.* **89**, 99–109. (doi:10.1046/j.1365-2745.2001.00527.x)

2. Ushio M, Yamasaki E, Takasu H, Nagano AJ, Fujinaga S, Honjo MN, Ikemoto M, Sakai S, Kudoh H. 2015 Microbial communities on flower surfaces act as signatures of pollinator visitation. *Sci. Rep.* **5**, 1–7. (doi:10.1038/srep08695)

3. Burdon RCF, Junker RR, Scofield DG, Parachnowitsch AL. 2018 Bacteria colonising *Penstemon digitalis* show volatile and tissue-specific responses to a natural concentration range of the floral volatile linalool. *Chemoecology* **28**, 11–19. (doi:10.1007/s00049-018-0252-x)

4. Roy R, Schmitt AJ, Thomas JB, Carter CJ. 2017 Review: Nectar biology: From molecules to ecosystems. *Plant Sci.* **262**, 148–164. (doi:10.1016/j.plantsci.2017.04.012)

5. Taylor LP, Hepler PK. 1997 Pollen germination and tube growth. *Annu. Rev. Plant Biol.* **48**, 461–491. (doi:10.1146/annurev.arplant.48.1.461)

6. Wei N, Ashman TL. 2018 The effects of host species and sexual dimorphism differ among root, leaf and flower microbiomes of wild strawberries in situ. *Sci. Rep.* **8**, 1–12. (doi:10.1038/s41598-018-23518-9)

7.   Massoni J, Bortfeld-Miller M, Jardillier L, Salazar G, Sunagawa S, Vorholt JA.
2020 Consistent host and organ occupancy of phyllosphere bacteria in a
community of wild herbaceous plant species. *ISME J.* **14**, 245–258.
(doi:10.1038/s41396-019-0531-8)
8.   Gaube P, Junker RR, Keller A. 2021 Changes amid constancy: Flower and leaf
microbiomes along land use gradients and between bioregions. *Basic Appl. Ecol.*
50, 1–15. (doi:10.1016/j.baae.2020.10.003)
9.   Rebolleda-Gómez M, Forrester NJ, Russell AL, Wei N, Fetters AM, Stephens
JD, Ashman T. 2019 Gazing into the anthosphere: Considering how microbes
influence floral evolution. *New Phytol.* (doi:10.1111/nph.16137)
10.   Vannette RL. 2020 The floral microbiome: Plant, pollinator, and microbial
perspectives. *Annu. Rev. Ecol. Evol. Syst.* **51**, 363–386. (doi:10.1146/annurev-
ecolsys-011720-013401)
11.   Bubán T, Orosz-Kovács Z, Farkas Á. 2003 The nectary as the primary site of
infection by Erwinia amylovora (Burr.) Winslow et al.: A mini review. *Plant*
Syst. Evol. **238**, 183–194. (doi:10.1007/s00606-002-0266-1)
12.   Vanneste JL. 2000 *Fire blight: the disease and its causative agent,* Erwinia
amylovora. CABI.
13.   Herrera CM, Pozo MI, Medrano M. 2013 Yeasts in nectar of an early-blooming
herb: Sought by bumble bees, detrimental to plant fecundity. *Ecology* **94**, 273–
279. (doi:10.1890/12-0595.1)
14.   Christensen SM, Munkres I, Vannette R. 2021 Nectar bacteria stimulate pollen
germination and bursting to enhance their fitness. *bioRxiv*
(doi:https://doi.org/10.1101/2021.01.07.425766)
15.   Huang M, Sanchez-Moreiras AM, Abel C, Sohrabi R, Lee S, Gershenzon J, Tholl
D. 2012 The major volatile organic compound emitted from *Arabidopsis thaliana*
flowers, the sesquiterpene (E)-β-caryophyllene, is a defense against a bacterial
pathogen. *New Phytol.* **193**, 997–1008. (doi:10.1111/j.1469-8137.2011.04001.x)
16.   McInnis SM, Emery DC, Porter R, Desikan R, Hancock JT, Hiscock SJ. 2006
The role of stigma peroxidases in flowering plants: Insights from further
characterization of a stigma-specific peroxidase (SSP) from Senecio squalidus
(Asteraceae). *J. Exp. Bot.* **57**, 1835–1846. (doi:10.1093/jxb/erj182)

17.   Dobson HEM, Bergstrom G. 2000 The ecology and evolution of pollen odors.
Plant Syst. Evol **222**, 63–87.

18.   Palmer-Young EC, Farrell IW, Adler LS, Milano NJ, Egan PA, Junker RR, Irwin
RE, Stevenson PC. 2019 Chemistry of floral rewards: intra- and interspecific
variability of nectar and pollen secondary metabolites across taxa. *Ecol. Monogr.*
89, 1–23. (doi:10.1002/ecm.1335)

19.   Primack RB. 1985 Longevity of individual flowers. *Annu. Rev. Ecol. Syst.* **16**,
15–37.

20.   Bawa KS. 1990 Plant-pollinator interactions in tropical rain forests. *Annu. Rev.*
Ecol. Syst. **21**, 399–422. (doi:10.1146/annurev.es.21.110190.002151)

21.   Shykoff JA, Bucheli E, Kaltz O. 1996 Flower lifespan and disease risk. *Nature*
379, 779.

22.   Rogers HJ. 2006 Programmed cell death in floral organs: How and why do
flowers die? *Ann. Bot.* **97**, 309–315. (doi:10.1093/aob/mcj051)

23.   Asakawa Y, Ludwiczuk A, Sakurai K, Tomiyama K, Kawakami Y, Yaguchi Y.
2017 Comparative study on volatile compounds of *Alpinia japonica* and *Elettaria*
cardamomum. *J. Oleo Sci.* **66**, 871–876. (doi:10.5650/jos.ess17048)

24.   van Vuuren SF, Viljoen AM. 2007 Antimicrobial activity of limonene
enantiomers and 1,8-cineole alone and in combination. **22**, 540–544.
(doi:10.1002/ffj.1843)

25.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA,
Turnbaugh PJ, Fierer N, Knight R. 2011 Global patterns of 16S rRNA diversity
at a depth of millions of sequences per sample. *Proc. Natl. Acad. Sci. U. S. A.*
108, 4516–4522. (doi:10.1073/pnas.1000080107)

26.   Hamady M, Walker JJ, Harris JK, Gold NJ, Knight R. 2008 Error-correcting
barcoded primers for pyrosequencing hundreds of samples in multiplex. *Nat.*
Methods **5**, 235–237.

27.   Ushio M. 2019 Use of a filter cartridge combined with intra-cartridge bead-
beating improves detection of microbial DNA from water samples. *Methods*
Ecol. Evol. **10**, 1142–1156. (doi:10.1111/2041-210X.13204)

28.   Tanabe AS, Toju H. 2013 Two new computational methods for universal DNA
barcoding: A benchmark using barcode sequences of bacteria, archaea, animals,
fungi, and land plants. *PLoS One* **8**. (doi:10.1371/journal.pone.0076910)

29.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP.
2016 DADA2: High-resolution sample inference from Illumina amplicon data.
Nat. Methods **13**, 581–583. (doi:10.1038/nmeth.3869)

30.   R Core Team. 2016 R: A language and environment for statistical computing. *R*
Found. Stat. Comput. **1**, 409. (doi:10.1007/978-3-540-74686-7)

31.   Huson DH, Auch AF, Qi J, Schuster SC. 2007 MEGAN analysis of metagenomic
data. *Genome Res.* **17**, 377–386. (doi:10.1101/gr.5969107)

32.   McMurdie PJ, Holmes S. 2013 Phyloseq: An R package for reproducible
interactive analysis and graphics of microbiome census data. *PLoS One* **8**.
(doi:10.1371/journal.pone.0061217)

33.   Davis NM, Proctor DiM, Holmes SP, Relman DA, Callahan BJ. 2018 Simple
statistical identification and removal of contaminant sequences in marker-gene
and metagenomics data. *Microbiome* **6**, 1–14. (doi:10.1186/s40168-018-0605-2)

34.   Oksanen J *et al.* 2010 Community ecology package: vegan. *R A Lang. Environ.*
Stat. Comput. , http://vegan.r-forge.r-project.org/. See https://cran.r-
project.org/web/packages/vegan/index.html.

35.   Rousseeuw PJ. 1987 Silhouettes: A graphical aid to the interpretation and
validation of cluster analysis. *J. Comput. Appl. Math.* **20**, 53–65.
(doi:10.1016/0377-0427(87)90125-7)

36.   Maechler M, Rousseeuw P, Struyf A, Hubert M, Hornik K. 2001 Cluster: Cluster
analysis basics and extensions. See https://cran.r-project.org/package=cluster.

37.   Gu Z, Eils R, Schlesner M. 2016 Complex heatmaps reveal patterns and
correlations in multidimensional genomic data. *Bioinformatics* **32**, 2847–2849.

38.   Lüdecke D. 2018 ggeffects: Tidy data frames of marginal rffects from regression
models. *J. Open Source Softw.* **3**, 772. (doi:10.21105/joss.00772)

39.   Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting linear mixed-effects
models using lme4. *J. Stat. Softw.* **67**, 1–48. (doi:10.18637/jss.v067.i01)

40. Qian X, Li X, Li H, Zhang D. 2021 Floral fungal-bacterial community structure
and co-occurrence patterns in four sympatric island plant species. *Fungal Biol.*
125, 49–61. (doi:10.1016/j.funbio.2020.10.004)

41. Anguita-Maeso M, Trapero JL, Olivares-García C, Ruano-Rosa D, Palomo-Ríos
E, Jiménez-Díaz RM, Navas-Cortés JA, Landa BB. 2020 Verticillium dahliae
inoculation and in vitro propagation modify the xylem microbiome and disease
reaction to verticillium wilt in a wild olive genotype. *bioRxiv* **12**, 1–15.
(doi:10.1101/2020.11.23.392977)

42. Mina D, Pereira JA, Lino-Neto T, Baptista P. 2020 Epiphytic and endophytic
bacteria onolivetree phyllosphere: Exploring tissue and cultivar effect. *Microb.*
Ecol. **80**, 145–157. (doi:10.1007/s00248-020-01488-8)

43. Sowndhararajan K, Marimuthu S, Manian S. 2013 Biocontrol potential of
phylloplane bacterium *Ochrobactrum anthropi* BMO-111 against blister blight
disease of tea. *J. Appl. Microbiol.* **114**, 209–218. (doi:10.1111/jam.12026)

44. Zang C, Lin Q, Xie J, Lin Y, Zhao K, Liang C. 2020 The biological control of the
grapevine downy mildew disease using *Ochrobactrum* sp. *Plant Prot. Sci.* **56**,
52–61. (doi:10.17221/87/2019-PPS)

45. Pattemore DE, Goodwin RM, McBrydie HM, Hoyte SM, Vanneste JL. 2014
Evidence of the role of honey bees (*Apis mellifera*) as vectors of the bacterial
plant pathogen Pseudomonas syringae. *Australas. Plant Pathol.* **43**, 571–575.
(doi:10.1007/s13313-014-0306-7)

46. Junker RR, Loewel C, Gross R, Dötterl S, Keller A, Blüthgen N. 2011
Composition of epiphytic bacterial communities differs on petals and leaves.
Plant Biol. **13**, 918–924. (doi:10.1111/j.1438-8677.2011.00454.x)

47. Hayes RA, Rebolleda-Gómez M, Butela K, Cabo LF, Cullen N, Kaufmann N,
O'Neill S, Ashman TL. 2021 Spatially explicit depiction of a floral epiphytic
bacterial community reveals role for environmental filtering within petals.
Microbiologyopen **10**, 1–19. (doi:10.1002/mbo3.1158)

48. Allard SM, Ottesen AR, Brown EW, Micallef SA. 2018 Insect exclusion limits
variation in bacterial microbiomes of tomato flowers and fruit. *J. Appl.*
Microbiol. **125**, 1749–1760. (doi:10.1111/jam.14087)

49.   Billing E. 1980 Fireblight *Erwinia amylovora* and weather: a comparison of
warning systems. *Ann. Appl. Biol.* **95**, 365–377. (doi:10.1111/j.1744-
7348.1980.tb04756.x)

50.   Pietrarelli L, Balestra GM, Varvaro L. 2006 Effects of simulated rain on
Pseudomonas syringae pv. *tomato* populations on tomato plants. *J. Plant Pathol.*
88, 245–251.

51.   Llontop MEM, Hurley K, Tian L, Bernal Galeano VA, Wildschutte HK, Marine
SC, Yoder KS, Vinatzer BA. 2020 Exploring rain as source of biological control
agents for fire blight on apple. *Front. Microbiol.* **11**.
(doi:10.3389/fmicb.2020.00199)

52.   Young JM. 1988 Bacterial blight of kiwifruit in New Zealand. *EPPO Bull.* **18**,
131–140. (doi:10.1111/j.1365-2338.1988.tb00358.x)

53.   Goumas DE, Malathrakis NE, Chatzaki AK. 1999 Characterization of
Pseudomonas viridiflava associated with a new symptom on tomato fruit. *Eur. J.*
Plant Pathol. **105**, 927–932. (doi:10.1023/A:1008725818334)

54.   Eisikowitch D, Lachance MA, Kevan PG, Willis S, Collins-Thompson DL. 1990
The effect of the natural assemblage of microorganisms and selected strains of
the yeast *Metschnikowia reukaufii* in controlling the germination of pollen of the
common milkweed *Asclepias syriaca*. *Can. J. Bot.* **68**, 1163–1165.
(doi:10.1139/b90-147)

55.   Cullen NP, Fetters AM, Ashman TL. 2021 Integrating microbes into pollination.
Curr. Opin. Insect Sci. **44**, 48–54. (doi:10.1016/j.cois.2020.11.002)

56.   Vannette RL, Gauthier MPL, Fukami T. 2013 Nectar bacteria, but not yeast,
weaken a plant– pollinator mutualism. *Proc. R. Soc. B Biol. Sci.* **280**.
(doi:10.1098/rspb.2012.2601)

57.   Sobhy IS *et al.* 2018 Sweet scents: Nectar specialist yeasts enhance nectar
attraction of a generalist aphid parasitoid without affecting survival. *Front. Plant*
Sci. **9**. (doi:10.3389/fpls.2018.01009)

58.   Rowe M, Veerus L, Trosvik P, Buckling A, Pizzari T. 2020 The reproductive
microbiome: An emerging driver of sexual selection, sexual conflict, mating
systems, and reproductive isolation. *Trends Ecol. Evol.* **35**, 220–234.
(doi:10.1016/j.tree.2019.11.004)

Table 1. The results of Blast search of the five dominant ASVs, and their mean percentages in the samples.

ASV	Class	Order	Family	Genus	Percentage
T00020	Actinobacteria	Actinomycetales	Propionibacteriaceae	*Cutibacterium*	17.20
T00030	Alphaproteobacteria	Rhizobiales	Brucellaceae	*Ochrobactrum*	12.36
T00004	Gammaproteobacteria	Enterobacterales	Erwiniaceae	*Erwinia*	8.82
T00007	Gammaproteobacteria	Pseudomonadales	Pseudomonadaceae	*Pseudomonas*	8.60
T00005	Gammaproteobacteria	Pseudomonadales	Pseudomonadaceae	*Pseudomonas*	6.13

604   Figure 1. Experimental design used in this study. a, An open flower and flower buds on
605   an inflorescence of *Alpinia japonica*. b, Floral parts of *Alpinia japonica* in open state
606   (left) and bud state (right). c, Experimental and sampling procedures.

 Figure 3. Association between the proportion of Cluster 5 and predicted probability of

fruit set. (a) Variation in the total proportions of the bacteria among the treatments. (b)

The predicted probabilities of fruit set 6 weeks after flowering under the different

treatments by the GLMM. The thick lines and the boxes indicate the predicted values

and their 95 % confidential intervals. Distribution of flowers that set fruits (top) or were

aborted (bottom) are shown by dots. Colors indicate the five individual plants (see the

caption of Fig. 2).

Table S1. The number of the samples used to monitor fruit set for the five treatments

and to analyze microbial communities.

	Fruit set and seed production					Microbial community				
Treatments	N1	N2	N4	N5	N6	N1	N2	N4	N5	N6
Bud	-	-	-	-	-	5	5	5	5	5
Open	10	26	14	16	19	5	5	5	5	5
Net bag	9	17	15	17	10	5	4	4	4	4
Paper bag	7	9	5	7	9	5	5	5	5	5
Microbe	6	11	5	7	5	5	5	5	5	5
Water	4	3	6	7	7	3	5	4	5	4

Table S2. Characteristics of the 10 clusters. The number of ASVs included, the median

proportion in the samples of the microbial inoculated (MI) and water sprayed (WI)

treatments, and p values of Wilcoxon rank sum tests in the comparisons of the

proportions between MI and WI are shown. The asterisks indicate that the proportions

are significantly different between the treatments (**, $p < 0.001$; ***, $p < 0.0001$).

Cluster 5, which shows significantly higher proportions in MI than in WI, is highlighted

with boldface type.

Cluster ID	No. ASVs included	Median of proportions		p-value of Wilcoxon rank sum test (MI vs. WI)
		Microbe (MI)	Water (WI)	
Cluster 1	7	0.054	0.326	0.0001**
Cluster 2	3	0.000	0.000	0.4535
Cluster 3	2	0.000	0.000	0.0694
Cluster 4	2	0.000	0.000	0.3206
Cluster 5	**5**	**0.825**	**0.208**	**0.0000***
Cluster 6	4	0.027	0.036	0.7899
Cluster 7	1	0.000	0.000	0.1691
Cluster 8	4	0.015	0.075	0.2253
Cluster 9	3	0.000	0.000	0.6284
Cluster 10	3	0.000	0.000	0.2313

Figure S1. Rarefaction curves of the samples. The vertical axis shows the number of observed prokaryotic ASVs. The number of sequences
per sample is shown on the horizontal axis. The three panels show the same graph with different ranges of the horizontal axis.

Figure S2. Variation in the prokaryotic diversity among the treatments and samples. The

diversity was significantly different among the treatments (Kruskal-Wallis test; p =

0.0054). Colors indicate the five individual plants.

Figure S3. Variation of the prokaryotic communities on the flowers under different
treatments. (a) NMDS plots of the microbial communities of flower buds and flowers
under different treatments. In the five panels, all microbial samples are plotted with grey
points, and samples from each plant are overlayed individually. The five treatments are
indicated by different colors. (b) Distribution of the distances from the treatment
centroid. The diversity was significantly different among the treatments (Kruskal-Wallis
test; p = 0.0026). Colors indicate the five individual plants.

Figure S4. Variation in the total proportions of the 10 ASV clusters among the flower bud and flowers under the five treatments. Colors

indicate the five individual plants.