Title: Insulin-dependent and -independent dynamics of insulin receptor trafficking in

muscle cells

Authors: Haoning Howard Cen[1], Jason Rogalski[2], Libin Abraham[3], Michael R. Gold[3],

Leonard J. Foster[2], James D. Johnson[1]*

Affiliation:

1 Department of Cellular and Physiological Sciences, Life Sciences Institute, University of

British Columbia, Vancouver, BC. Canada.

2 Department of Biochemistry & Molecular Biology, and Michael Smith Laboratories,

University of British Columbia, Vancouver, BC, Canada

3 Department of Microbiology and Immunology, Life Life Sciences Institute, University of

British Columbia, Vancouver, BC. Canada

Running title: Muscle cell insulin receptor trafficking and interactions

Keywords: insulin, insulin receptor, caveolin, clathrin, endocytosis

*Address correspondence to: James D. Johnson, Ph.D., Professor. Diabetes Research

Group, Dept. of Cellular and Physiological Sciences & Dept. of Surgery, University of

British Columbia, 5358 Life Sciences Building, 2350 Health Sciences Mall, Vancouver,

BC, Canada, V6T 1Z3

E-mail: James.D.Johnson@ubc.ca; Fax: (604) 822-2316

Twitter: @JimJohnsonSci

Acknowledgments: Spinning disk confocal and STED microscopy were performed in

the LSI IMAGING facility of the Life Sciences Institute of the University of British

Columbia using infrastructure funded by the Canadian Foundation of Innovation and BC

Knowledge Development Fund as well as a Strategic Investment Fund (Faculty of

Medicine, University of British Columbia). We thank Dr. Ivan Robert Nabi and Dr. Guang

Gao of the LSI IMAGING facility for guidance on imaging experiments. Mass

Spectrometry used here was supported by the Canada Foundation for Innovation and

Genome Canada (264PRO). We thank Yuanyuan Amanda Yang for guidance on Python

and R coding and Yi Han Xia for assistance in R coding. The schematics in Figure 7 is

created with BioRender.com.

Contributions: HHC designed the study, performed all experiments, analyzed data, and

wrote the manuscript. JR performed mass spectrometry, analyzed the output data, and

edited the manuscript. LA and MG contributed to the MATLAB scripts and the analysis of

single particle tracking and edited the manuscript. LF contributed to the design of the

mass-spectrometry experiments and edited the manuscript. JDJ designed the study,

supervised the research, and edited the manuscript. JDJ is the ultimate guarantor of this

work.

Funding: This study was funded by a Canadian Institues of Health Research Operating

Grant (P-133692) to JDJ.

Conflict of interest: The authors declare that they have no conflicts of interest with the

contents of this article.

Abstract

Insulin resistance contributes to type 2 diabetes and can be driven by hyperinsulinemia. Insulin receptor (INSR) internalization and cell-surface dynamics at rest and during insulin exposure are incompletely understood in muscle cells. Using surfacing labeling and live-cell imaging, we observed robust basal internalization of INSR in C2C12 myoblasts, without a robust effect of added insulin. Mass-spectrometry analysis of INSR-binding proteins identified potential molecular mechanisms associated with internalization. We confirmed known interactors, including IGF1R, but also identified underappreciated INSR-binding factors such as ANXA2. Protein-protein interaction network mapping suggested links between INSR and caveolin-mediated endocytosis. INSR interacted with both caveolin and clathrin heavy chain (CLTC) in mouse skeletal muscle and C2C12 myoblasts. Whole cell 2D super-resolution imaging revealed that high levels of insulin (20 nM) increased INSR colocalization with CAV1 but decreased its colocalization with CLTC. Single particle tracking confirmed the colocalization of cell-surface INSR with both over-expressed CAV1-mRFP and CLTC-mRFP. INSR tracks that colocalized with CAV1 exhibited longer radii and lifetimes, regardless of insulin exposure, compared to non-colocalized tracks, whereas insulin further increased the lifetime of INSR/CLTC colocalized tracks. Overall, these data suggest that muscle cells utilize both CAV1 and CLTC-dependent pathways for INSR dynamics and internalization.

Introduction

The insulin receptor (INSR) is one of the most studied membrane proteins due to its important roles in insulin resistance and diabetes. However, many fundamental aspects of INSR biology remain unclear, including the mechanisms and kinetics of its internalization. The INSR can be internalized via endocytosis[1,2] and endosomal INSR localization contributes to the phosphorylation of specific downstream targets when

compared with INSR at the cell surface[3,4]. Moreover, ligand-receptor complex

dissociation in acidified endosomes permits insulin degradation while the receptor can be

sorted to recycling or degradation pathways[1,2]. INSR internalization is reduced in

monocytes[5,6] and adipocytes[7] obtained from patients with insulin resistance and type 2

diabetes, suggesting the importance of this process in disease pathogenesis.

Tissue-specific insulin resistance has been implicated in the development of

diabetes[8,9], but it is less clear whether there are tissue-specific INSR internalization

kinetics and/or mechanisms. Interestingly, adipocytes internalize insulin at a lower rate in

the presence of high insulin concentrations, as compared to low insulin conditions[10],

while hepatocytes are the opposite[11]. The consensus view is that INSR is endocytosed

via a clathrin-coated pit mechanism, based on evidence from a variety of cell types,

including human and rat hepatocytes, fibroblasts, and lymphocytes[11-14]. However, other

studies have shown that caveolin-rich membrane lipid rafts, caveolae, are critical for

INSR internalization in other cell types, including adipocytes[15,16]. In pancreatic β-cells, we

showed that caveolin-1 (CAV1) mediates INSR internalization, bypassing clathrin positive

compartments[4]. In muscle cells, previous studies have shown that clathrin-mediated

endocytosis is involved in INSR internalization, but the role of the caveolin-mediated

pathway has not been investigated[17]. The INSR internalization kinetics and routes are

less well characterized in muscle cells.

In the present study, we employed inter-domain tagged INSR with fluorescent

proteins[4] and SNAP-tag[18], surface biotinylation assays, as well as super-resolution and

TIRF imaging, to explore the dynamics of INSR internalization and the effects of ambient

insulin. We also used proteomics to identify INSR interacting proteins and defined the

relationship between INSR and caveolin- or clathrin-mediated internalization pathways.

Together, our data show that INSR exhibited a high degree of insulin-independent

internalization in C2C12 myoblasts.

Results

Effects of insulin on INSR internalization

First, we characterized INSR internalization in C2C12 myoblasts under basal

conditions and in the context of added insulin. A high rate of basal INSR internalization

was observed in high-glucose serum-free DMEM medium when compared with PBS.

INSR internalization over 5 min was not significantly stimulated by addition of insulin to

the DMEM medium (Fig. 1A). Therefore, INSR has a high rate of constitutive

internalization in this cell type, which is enhanced by the nutrients in medium, but that is

largely independent of acute exposure to insulin across a range of physiological doses.

These observations in myoblasts are consistent with our previous measurements in

differentiated C2C12 myotubes[19].

Qualitative analysis of INSR kinetics in the context of insulin doses

We also visualized INSR internalization at single-cell level using

inter-domain-tagged INSR, which we have previously found more faithfully recapitulates

the sub-cellular localization of INSR compared to terminal fusion proteins[4]. C2C12

myoblast cells with stable overexpression of SNAP-tagged INSR isoform A

(INSR-A-SNAP) were labeled using a cell non-permeable SNAP-Surface Alexa Flour 488

dye (Fig. 1B). Live-cell time-lapse imaging of surface-labeled INSR-A-SNAP by spinning

disk confocal microscopy revealed INSR internalization in the context of 0, 0.2, and 20

nM insulin over 15 min (Fig. 1B, Video S1). Consistent with the surface biotinylation

assay results in Figure 1A, we did not observe a clear difference in INSR internalization in

the presence of insulin in these cells. Internalized INSR vesicles exerted dynamic

movement and interaction, which was more evident between 20-30 min (Fig. 1C-F, Video

S2). INSR-containing vesicles exhibited many apparent fusion and fission events

regardless of insulin concentrations. Some smaller vesicles merged into a larger vesicle

that remained in the field undergoing limited and apparently random motion (one example

is shown in Fig. 1F), suggestive of a fusion event. A caveat associated with 2D imaging

and limited Z resolution is that the observed INSR vesicles might not all be on the same

optical plane. Our experiments using surface labeling of SNAP-tagged INSR permitted

the visualization of the dynamic behavior of internalized INSR vesicles. These qualitative

observations motivated the more quantitative studies below.

Quantitative analysis of INSR kinetics in the context of insulin

Lateral movement of tyrosine kinase receptors has been linked to their functional

state. Ligand binding of some monomeric receptors, such as EGFR, promotes their

dimerization and aggregation and decreases their lateral mobility[20]. While INSRs are

thought to already exist as dimers in plasma membrane, aggregation of INSR by

antibodies can stimulate INSR signaling[20], and INSR may need to localize to membrane

lipid rafts for normal signaling[21,22]. Therefore, we thought to quantitatively assessed

whether insulin affects the lateral movement of INSR in or near the plasma membrane

using time-lapse TIRF microscopy of C2C12 myoblast transiently expressing

inter-domain-tagged INSR-A-EGFP. We imaged on the same cells at 0, 6, or 11 min in

the presence of 2 nM insulin, as well as a control group without insulin to control for other

variables such as photobleaching (Fig. 2A). Single particle tracking was conducted on the

INSR-A-EGFP (Fig. 2B), and we calculated their diffusion coefficient (Fig. 2C,D) and

track radius (Fig. 2E,F). Insulin did not affect the track radius or diffusion coefficient at any

time points. Together with the qualitative data reported above, these observations

suggest that the majority of INSR mobility is independent of acute insulin stimulation in

this cell model.

Novel INSR interactors identified by immunoprecipitation and mass spectrometry

To explore the molecular mechanisms associated with INSR internalization, we

conducted co-immunoprecipitation followed by mass-spectrometry to identify INSR

interactors. We also used INSR knockout C2C12 myoblasts as an negative control and
an INSR antibody that does not cross-react with Insulin-like growth factor 1 receptor
(IGF1R)[23] to identify INSR interactors with higher specificity than any previously reported
study. After filtering the detected proteins against the INSR knockout cell control lysates,
we identified INSR and five high-confidence INSR-binding proteins (Fig. 3A). The
immunoprecipitated abundance of these detected interactors was not significantly
changed by the addition of 2 nM insulin (Fig. 4A). INSR is well established to form
heterodimers with IGF1R[24,25], and its immunoprecipitation represented a type of internal
positive control for this study. The other detected proteins are not well-known INSR
interactors.

To discover whether any of the identified INSR-binding proteins might be connected
with endocytosis machinery, protein-protein interaction (PPI) network was constructed
between these co-immunoprecipated proteins and all the proteins under the
"Endocytosis" pathway in Kyoto Encyclopedia of Genes and Genomes (KEGG). We
manually extracted a subnetwork that connected all our identified proteins, except for
HIST1H1B which did not connect with any proteins in this network (Fig. 3B). In the
resultant PPI network, annexin A2 (ANXA2) is connected with CAV1 and CAV3 (Fig. 3B),
and it has been reported to localize in caveolae[26]. Ribosomal protein SA (RPSA) is a
laminin receptor reported to localize in plasma membrane lipid rafts[27]. Heat shock protein
B1 (HSPB1; a.k.a. HSP27) is exclusively found in insulin-responsive tissues and has
been implicated in insulin sensitivity and INSR/IGF1R signalling[28-30]. The PPI network
and cellular localization of these interactors pointed to the possibility that INSR may also
associate with caveolins and/or related proteins in myoblasts.

Insulin-dependent interaction between INSR and CAV3 or CLTC in vivo
We sought to confirm that INSR could interact with both caveolin and clathrin in
skeletal muscle *in vivo*. Skeletal muscle was collected 5 or 10 min after intraperitoneal

injection of insulin (1.5 U/kg) or PBS, timepoints sufficient to measure acute insulin

signalling via both AKT and ERK signalling pathways (Fig. 3C). Acute insulin

administration did not change the amount of INSR that could be immunoprecipitated from

skeletal muscle, nor did it alter the association between INSR and CLTC (Fig. 3D).

However, 10 min after insulin injection, there was a significantly increased interaction

between INSR and caveolin, specifically, the mature skeletal muscle-specific isoform

CAV3[31,32] (Fig. 3D). These data show that acute insulin signaling results in an increased

association between INSR and CAV3 *in vivo*, but does not define the cellular context for

this interaction.

Super-resolution imaging of INSR, CAV1, and CLTC interactions in the presence of

insulin

We next used super-resolution STED imaging to examine colocalization between

INSR and clathrin (CLTC) or caveolin (C2C12 myoblasts only express CAV1 before

differentiation30). C2C12 myoblast cells with stable overexpression of INSR-A-SNAP

were labeled by cell-nonpermeable SNAP substrate, to image only newly internalized

INSR. Cells were incubated with 0, 0.2 or 20 nM insulin for 30 min, and then fixed.

Endogenous CLTC and CAV1 were then labeled using immunofluorescent antibody

staining (Fig. 4A). The colocalization between INSR-SNAP, CAV1, and CLTC was

accessed by Object Pearson Coefficient (Fig. 4B,C). The degree of colocalization

between INSR and CAV1 was similar in the presence of 0 or 0.2 nM insulin but was

higher under 20 nM insulin (Fig. 4B). However, INSR and CLTC colocalized more at 0.2

nM insulin, but less at 20 nM insulin, compared to 0 nM insulin (Fig. 4B). CAV1 and CLTC

do not have known biological interactions; therefore, their low level of colocalization

served as an internal negative control and, indeed, it did not change under different

insulin levels (Fig. 4B). Using the same data, the relative colocalizations between INSR

and CAV1 or CLTC were also considered at different insulin concentrations, further

illustrating how high insulin drives INSR to colocalize with CAV1 (Fig. 4C). Together,

these data revealed the insulin-dependent interaction of INSR with both CAV1 in vivo and

in vitro, alongside the constitutive interaction with CLTC.

Effects of CAV1 and CLTC on INSR mobility

To better understand the interplay between INSR, CAV1, and CLTC, we

co-expressed INSR-B-TagBFP with CAV1-mRFP or CLTC-mRFP in C2C12 myoblasts

and imaged them by TIRF time-lapse microscopy. Colocalization between INSR and

CAV1 or CLTC were observed (Fig. 5A,B). Some colocalized puncta moved or

disappeared together with closely associated intensity, referred to here as colocalized

tracks (examples in Fig. 5C,D). Tracks were considered as colocalized when the distance

between the centers of particle signals was less than 2 pixels for more than 4 continuous

frames. These observations suggested that INSR interacted with CAV1 and CLTC on the

cell surface and might be endocytosed through both caveolin- and clathrin-mediated

pathways.

We also conducted single particle tracking on these INSR puncta to define the

effects of CAV1 or CLTC on INSR movement in the plasma membrane. Similar to our

observations above (Fig. 2), insulin did not alter the diffusion coefficient or track radius of

INSR. Cells with CAV1 co-expression had lower diffusion coefficients and track radius of

INSR compared to cells with CLTC co-expression, regardless of ambient insulin

exposure (Fig. 5E-H). CLTC-expressing cells under 0 nM insulin had shorter INSR

lifetimes than other groups (Fig. 5I,J), which might suggest a more rapid internalization.

These data suggest that CAV1 may restrain INSR movement, relative to CLTC.

Finally, we performed an analysis within the same cells, comparing INSR tracks that

were colocalized or not colocalized with either CAV1 or CLTC tracks. Within cells, INSR

spent more time associated with CAV1, and for longer periods, compared with CLTC

(Fig. 6A,B). There were no clear effects of time on the association between INSR and

CAV1 or CLTC across different moments of their lifetimes together (Fig. 6A,B). Within the

same cells, we compared the colocalized and the non-colocalized INSR tracks, for both

CAV1 and CLTC, and found no significant differences in mean diffusion coefficients (Fig.

6C,D). The track radius of INSR tracks colocalized with CAV1 was larger than

non-colocalized tracks regardless of insulin (Fig. 6E). The track radius of INSR tracks

colocalized with CLTC was larger than non-colocalized tracks, but only in the no insulin

condition (Fig. 6F). The lifetimes of INSR tracks colocalized with CAV1 were longer than

non-colocalized tracks regardless of insulin (Fig. 6G). INSR tracks colocalized with CLTC

also had longer lifetimes than non-colocalized tracks, and insulin further increased the

lifetime of colocalized tracks (Fig. 6H). Overall, our live-cell imaging of INSR confirmed its

separate interactions with either CAV1 or CLTC on the cell surface.

Discussion

The goal of this study was to explore the INSR trafficking dynamics and interactions

in a myoblast cell line. We demonstrated that INSR was constitutively internalized both in

the absence and presence of insulin. We used a high-stringency proteomics approach to

identify specific INSR binding proteins and PPI network modeling to predict a role for

caveolin in INSR internalization. Biochemical experiments, super-resolution imaging, and

TIRF imaging suggested a role for CAV1 in INSR internalization and cell-surface

trafficking (Fig. 7).

The constitutive internalization of INSR has been described previously in

lymphocytes and shown not to require ligand binding and kinase activity, although

insulin-stimulated autophosphorylation of INSR could accelerate internalization[14]. In

differentiated C2C12 myotubes, INSR phosphorylation levels were constitutively high,

with little additional activation by insulin across the physiological range[19], consistent with

the lack of stimulation of INSR internalization by acute insulin treatment in this study.

Surface labeling by SNAP-tag enabled clear visualization of internalized INSR-containing

vesicles, permitting the study of INSR endosomal sorting and traffic. The lateral

movement of INSR in or near the plasma membrane was also not significantly affected by

insulin, consistent with a previous study that reported unchanged INSR diffusion

coefficient with or without insulin in HEK293T, HepG2, and human IR-overexpressing

Rat-1 cells [33]. CAV1 may have a role in decreasing the mobility of INSR. Indeed,

caveolae have restricted lateral movement in the plasma membrane due to the

connection to antin cytoskeleton through Filamin-A[34], so the CAV1-dependent decrease

in INSR mobility might reflect the interaction of INSR with caveolae. Similarly, the

accumulation of ganglioside GM3 in TNFα-induced insulin resistance state was

associated with the elimination of INSR from the caveolae and increased INSR mobility

on the cell surface[21]. Addition studies will be required to fully define the processes that

control INSR movement within the plasma membrane.

The possibility of more than one internalization pathway for INSR has been

understudied. The current study supports the concept that INSR utilizes both clathrin-

and caveolin-mediated endocytosis. Another study confirmed the involvement of the

clathrin-mediated pathway in myoblast by identifying LMBD1 protein as the cargo-specific

adaptor of INSR in clathrin-mediated endocytosis[17]. The combination of dual clathrin and

caveolin endocytosis routes has been demonstrated for IGFR[35], which has a similar

structure and function as INSR and can form heterodimers with INSR[36]. In hepatocytes, it

has also been suggested that INSRs are internalized in two functionally separate

pathways based on the much more rapid recycling of receptors internalized by high

concentrations of insulin (17.2 nM) versus low insulin (0.17 nM), and treatments known to

inhibit coated pit formation blocked internalization of the insulin-INSR complex at low

concentrations of insulin but allowed internalization in response to high concentrations of

insulin[11]. Most existing studies on INSR internalization looked at one pathway at a time.

The mechanisms of coordinated multiple internalization routes, implied by the

insulin-dependent clathrin and caveolin interactions we observed in muscle cells, might

direct INSR to different destinations (e.g. recycle or degradation)[37] and should be

comprehensively characterized in more cell types to assess potential roles in

tissue-specific insulin resistance.

We used stringent proteomics and INSR CRISPR knockout control cells to identify

known, novel, and under-appreciated interactors of INSR. As expected, we identified

IGF1R, an established dimerization partner of INSR. We also demonstrated the specific

binding of HSBP1/HSP27, which is involved in tyrosine kinase signalling pathways

downstream of p38 MAPK, ERK, and MAPKAPK5[38] and has been previously linked to

insulin sensitivity[28-30]. We demonstrated an interaction between INSR and RPSA, which

has been previously co-immunoprecipitated with INSR and insulin receptor substrates in

3T3L1 preadipocytes[39]. In other cell types, RPSA drives ERK signalling and cytoskeletal

organization, with nuclear functions affecting cell growth, survival, migration, and protein

synthesis[27]. ANXA2 is not a well-known interactor of INSR, but one study reported

ANXA2 interaction with INSR and IGF1R[40]. Other studies found that INSR signalling can

tyrosine-phosphorylate[41,42] and SUMOylate[43] AXNA2, which mediates insulin-stimulated

actin rearrangement. Interestingly, INSR internalization seemed to be necessary for

AXNA2 phosphorylation because mutant INSR that was incapable of internalization but

normal for tyrosine kinase signalling failed to phosphorylate AXNA2[41]. On the other hand,

ANXA2 has a strong direct connection with epidermal growth factor receptor (EGFR) in

the PPI network which belongs to the same receptor tyrosine kinase superfamily as

INSR[44]. ANXA2 knockdown halted EGFR endocytosis beyond the early endosome,

which was accompanied by enhanced EGFR and AKT phosphorylation[45]. These data

support that possiblity that ANXA2 may have a profound role in INSR trafficking and

signalling that is worth investigating. Both *Anxa2* and *Rspa* mRNAs were upregulated by

prolonged insulin in C2C12 myotubes[8]. HIST1H1B belongs to the family of linker histones

that is important in chromatin organization[46], but the meaning of the INSR interaction

requires further investigation. This proteomic analysis provides clear evidence for a

limited number of proteins that are likely to be direct interactors with INSR in myoblast

cells, and demonstrates that the majority of these interactors were not significantly

altered by insulin exposure.

There are caveats and unexplained observations in the current study. Although

STED superresolution microscopy might be able to reach 20-nm resolution, primary and

secondary antibody labeling might add ~30-nm distance and reduce the accuracy[47]. TIRF

microscopy has an axal resolution of ~100 nm and and a defraction-limited lateral

resolution (~200 – 300 nm)[48]. Therefore, the colocalization analyses supports the

co-occurance but may not confirm the functional interaciton of the proteins. Nonetheless,

the insulin-dependent colocalization observed in STED microscopy and the track

colocalization over multiple frames observed in TIRF microscopy are evidences that the

proteins had functional interaction instead of random colocalization. The CAV1- or

CLTC-colocalized INSR tracks had longer lifetime than non-colocalized tracks, which

conflicted with the assumption that CAV1 or CLTC internalized INSR, but instead, they

seemed to retain INSR on the membrane. There are multiple possible explanations; for

example, overexpressed CAV1 might stabilize caveolae[49], or CLTC could form stable flat

lattices, termed clathrin plaques[50], which might have novel involvement in INSR

internalization. We did not perform loss-of-function studies to confirm the roles of caveolin

and clathrin in INSR endocytosis. Further research is needed to elucidate the roles of

specific INSR internalization routes on insulin actions. Inhibition of clathrin-mediated

endocytosis of INSR by knocking down LMBD1 in myoblasts increased activation of the

INSR and AKT[17]. We also recently discovered that reducing CAV1-dependent INSR

internalization blocked the activation of ERK, but not AKT, signalling in pancreatic

beta-cells[4]. Cholesterol extraction by methyl-β-cyclodextrin, which is often used as a

caveolae-specific inhibitor but also affects clathrin-coated pit formation[51], inhibited the

phosphorylation of IRS1, SHC, and AKT, but not ERK or INSR phosphorylation, in

adipocytes[16,52]. In contrast, inhibition of both clathrin- and caveolin-mediated endocytosis

by the dynamin K44A dominant-negative mutant[53] reduced the phosphorylation of SHC

and ERK phosphorylation, but not IRS1 or AKT phosphorylation, in rat hepatoma H4IIE

cells[12]. Therefore, INSR internalization enables selective activation of downstream

signalling pathways, and in a tissue-specific manner.

In summary, this study provides important information on the internalization of INSR

in myoblast cells and defines the stringent INSR interactome. We also provide evidence

using a variety of methods that INSR is internalized via both clathrin- and

caveolin-dependent pathways in this cell model. The current study can serve as a

stepping stone for more detailed research on how multiple internalization routes can

provide cell-type customized regulations on INSR internalization.

Materials and Methods

Cell culture

The C2C12 mouse myoblast (ATCC cell line provided by Dr. Brain Rodrigues,

University of British Columbia, Vancouver, Canada) was maintained in Dulbecco's

modified Eagle's medium (DMEM, Gibco) supplemented with 10% (v/v) fetal bovine

serum (FBS, Gibco), and 1% penicillin-streptomycin (100 µg/ml; Gibco). All downstream

experiments were conducted after 6-hour serum starvation in serum-free medium

(DMEM supplemented with 1% penicillin-streptomycin), except as otherwise indicated.

Western blot analyses

C2C12 myotubes or mice skeletal muscle (gastrocnemius) tissues were sonicated

in RIPA buffer (50 mM β-glycerol phosphate, 10 mM HEPES, 1% Triton X-100, 70 mM

NaCl, 2 mM EGTA, 1 mM Na_3VO_4, and 1 mM NaF) supplemented with complete mini

protease inhibitor cocktail (Roche, Laval, QC). Lysates were incubated in Blue Loading

Buffer containing 50 mM DTT (Cell Signaling, Danvers, MA) at 95°C for 5 min and

resolved by SDS-PAGE. Proteins were then transferred to PVDF membranes (BioRad,

CA) and probed with antibodies against p-ERK1/2 (Thr202/Tyr204) (1:1000, Cat. #4370),

ERK1/2 (1:1000, Cat. #4695), p-AKT (Ser473) (1:1000, Cat. #9271), p-AKT (Thr308)

(1:1000, Cat. #9275), AKT (1:1000, Cat. #9272), INSR-β subunit (1:1000, Cat. #3020S),

p-INSRβ (Tyr1150/1151) (1:1000, Cat. #3024), FOXO1 (1:1000, Cat. #2880), p-FOXO1

(Thr24) (1:1000, Cat. #9464), all from Cell Signalling (CST), and β-tubulin (1:2000, Cat.

#T0198, Sigma). The signals were detected by secondary HRP-conjugated antibodies

(anti-mouse Cat. #7076, anti-rabbit Cat. #7074; Cell Signaling) and Pierce ECL

Western Blotting Substrate (Thermo Fisher Scientific). Protein band intensities were

quantified with Image Studio Lite software (LI-COR).

Surface Protein Biotinylation Assay

Biotinylation of surface proteins was performed as previously described[19]. In brief,

cells were incubated with cell-impermeable EZ-Link-NHS-SS-biotin (300 µg/ml in PBS;

Pierce) at 37°C for 2 min and washed with ice-cold 50 mM Tris-buffered saline (TBS) to

remove excess biotin. For detecting internalized proteins, cells were washed with PBS

and incubated in PBS or serum-free medium supplemented with 0, 0.2, 2, or 20 nM

insulin at 37°C to stimulate INSR internalization. After certain time periods, cells were

incubated with glutathione solution (50 mM glutathione, 75 mM NaCl, 1 mM EDTA, 1%

BSA, 75 mM NaOH) to strip the remaining surface biotin lysed in complete RIPA buffer.

Biotinylated internalized proteins in lysates were captured by NeutrAvidin beads (Pierce)

and eluted by incubating in Blue Loading Buffer (50 mM DTT, Cell Signaling, Danvers,

MA) containing 50 mM DTT at 95°C for 5 min. Internalized INSR in eluent and total INSR

in lysates were detected in Western blot analysis.

Plasmids, SNAP-tag cloning and lentivirus infection

CAV1-mRFP and CLTC-mRFP plasmids were generous gifts from Dr. Ivan Robert

Nabi, Vancouver, BC, Canada. INSR-A-TagBFP and INSR-B-TagBFP plasmids are

previously described[4].

PCR amplification of DNA fragments was performed using Q5 High-Fidelity DNA

Polymerase (Cat. #M0491, New England Biolabs, Ipswich, MA, USA) according to the

manufacturer's instructions. Prior ligation with the target vector, all PCR products were

subcloned into pJET1.2 vector using the CloneJET PCR Cloning Kit (Cat. #K1232,

Thermo Scientific, Waltham, MA, USA) according to the manufacturer's instructions.

Subsequently, the PCR products were excised from the pJET1.2 vector, gel purified and

ligated with the digested target vector using T4 DNA ligase. To generate INSR-A-EGFP

and INSR-B-EGFP plasmids, eGFP was PCR amplified using primers 5'-CGT ACG ATC

GAA TTC CCC GTG AGC AAG GGC GAG GAG CTG-3' (forward) and 5'- ACC GGT

ATC TGA TCC GGA CTT GTA CAG CTC GTC CAT GCC GA-3' (reverse) and inserted

into pcDNA3.1(-)-INSR-A-TagRFP and pcDNA3.1(-)-INSR-B-TagRFP (previously

described[4]) to replace TagRFP by EGFP using BsiWI and AgeI restriction sites.

To generate pLenti-INSR-A-SNAPf and pLenti-INSR-B-SNAPf lentiviral plasmids,

SNAP tag was firstly PCR amplified from pSNAPf Vector (Cat. #N9183S, New England

Biolabs, Ipswich, MA, USA) using primers 5'-CGT ACG ATC GAA TTC CCC ATG GAC

AAA GAC TGC GAA ATG AAG CGC ACC ACC-3' (forward) and 5'- ACC GGT ATC TGA

TCC GGA ACC CAG CCC AGG CTT GCC CAG-3' (reverse) and inserted into

pcDNA3.1(-)-INSR-A-TagRFP and pcDNA3.1(-)-INSR-B-TagRFP to replace TagRFP by

SNAP-tag using BsiWI and AgeI restriction sites. Subsequently, INSR-A-SNAP and

INSR-B-SNAP ORFs were PCR amplified from pcDNA3.1(-)-INSR-A/B-SNAPf plasmids

using primers 5'-GAA CCC ACT GCT TAC TGG CT-3' (forward) and 5'-GGT GGT TGT

ACA CTA GAA GGC ACA GTC GAG GC -3' (reverse) and inserted into

pLenti-C-Myc-DDK-IRES-GFP vector to replace the Myc tag, Flag tag, IRES, and

TurboGFP using NheI and BsrGI restriction sites. The PCR program for this long

INSR-SNAPf fragments was optimized to be as follows: initial denaturation 98°C 3min; 30

cycles of 98°C 10 s, 71°C 30 s, 72°C 4min; final extension 72°C 2min; hold 4°C.

SNAP-tag labeling and spinning disk confocal imaging of C2C12 myoblast

Cell non-permeable SNAP-Surface Alexa Fluor 488 substrate (Cat. #S9129S) was

dissolved and diluted to 5 µM concentration in serum-free medium containing 0.5% BSA

according to the manufacture's instruction. Next, stable C2C12 myoblasts cell line

expressing INSR-A-SNAP seeded on glass-bottom dishes (Cat. #P35G-1.5-14-C,

MatTek) were incubated in the substrate for 30 min on ice and washed 3 times with

serum-free medium (0.5% BSA).

C2C12 dish was then set up on the Spinning disk microscope, applied 37°C

FuoroBrite DMEM (Cat. #A1896701, Gibco) serum-free medium containing 0, 0.2 or 20

nM human insulin (Cat. #I9278, SigmaAldrich), and immediately started imaging in a

37°C chamber using a 100x/1.45 oil objective and QuantEM 512SC Photometrics

camera on a spinning disk confocal system based on Zeiss Axiovert 200M microscope

(LSI Imaging Core, UBC). Time-lapse imaging was conducted at 30 s intervals between 0

– 20 min or at 5 s intervals between 20-30 min.

Immunofluorescence staining

Stable C2C12 cell line seeded on glass-bottom dishes were labeled with cell

non-permeable SNAP-Surface Alexa Fluor 488 substrate, incubated with 0, 0.2 or 20

nM human insulin for 30 min, fixed with 4% (w/v) paraformaldehyde (Sigma-Aldrich) for

20 minutes at room temperature, and washed 3 times with PBS. After permeabilized in

0.1% (v/v) Triton X-100 (Sigma-Aldrich) for 15 minutes, primary antibodies targeting

Caveolin1 (1:400, Rabbit mAb, Cat. #3267, Cell Signaling) and CLTC heavy chain

(1:200, Goat pAb, Cat. #sc-6579, Santa Cruz) were incubated overnight at 4°C.

Coverslips were then washed in PBS, blocked in Dako Protein Block (Cat #X0909,

Agilent), and stained by secondary antibodies (1:1000, Cy3 Donkey anti-Rabbit, Cat.

#711-165-152, Jackson ImmunoResearch; 1:500, Alexa Fluor 633 Donkey anti-Goat,

Cat. # A21082, Invitrogen) diluted in Dako Antibody Diluent (Cat. # S0809, Agilent) for 2

hours at room temperature. After washing in PBS, coverslips were mounted with Prolong

Diamond Antifade Mountant (Cat. #P36961, Invitrogen).

STED microscopy and image analysis

Stimulated emission depletion (STED) microscopy was performed with the 100×/1.4

Oil HC PL APO CS2 STED White objective of a Leica TCS SP8 3× STED microscope

(Leica, Wetzlar, Germany) equipped with a white light laser, HyD detectors, and Leica

Application Suite X (LAS X) software (LSI Imaging Core, Life Sciences Institute,

University of British Columbia). Time-gated fluorescence detection was used to further

improve lateral resolution. INSR-SNAP labeled by Alexa Fluor 488 was excited at 488 nm

and depleted using the 592-nm depletion laser. CAV1 labeled by Cy3 was excited at 555

nm and depleted using the 592-nm depletion laser. CLTC labeled by Alex Fluor 633 was

excited at 633 nm and depleted using the 660-nm depletion laser. Sequential acquisition

in the order of AF633/Cy3/AF488 was used to avoid cross-talk.

Huygens Professional software (Scientific Volume Imaging, Hilversum, the

Netherlands) was used to deconvolve STED images and calculate the Object Pearson

correlation coefficient for INSR-SNAP, CAV1, and CLTC.

TIRF microscopy and single particle tracking analysis

Total internal reflection fluorescence (TIRF) microscopy was performed on a Zeiss

Axiovert 200M with a 100x Alpha-PlanFluar NA 1.45 oil objective (Zeiss, Germany) and a

TIRF laser angle modifier (Zeiss). Cells were kept at 37°C in a temperature-controlled

incubation chamber (Harvard Apparatus, Holliston, MA). Images were acquired with a

CoolSNAP HQ2 CCD camera (Photometrics, Tucson, AZ). 405 nm and 561 nm

solid-state diode laser system was used to excite TagBFP and mRFP tagged proteins.

The ImageJ distribution Fiji was used to process microscope images[54]. The local

background of TIRF images was subtracted using the rolling ball method (20-pixel radius)

before single particle tracking.

Single particles (INSR, CAV1, or CLTC) were detected and tracked using Icy

bioimaging analysis software version 1.9.8.1[55]. Particles were detected by the

UnDecimated Wavelet Transform Detector plugin that was set to detect bright spots over

a dark background, exhibit 70% sensitivity to spots <3 pixels, detect particle size between

2 and 3000 pixels. Particles were tracked using the Multiple Hypothesis Tracking method

built into the Spot Tracking plugin that was set to recognize particles that exhibited either

diffusive or directed motion[56,57]. Track radius (maximum displacement of the particle

track) and lifetime of the particles were exported from the Motion Profile Processor of the

Spot Tracking plugin. Diffusion coefficients were calculated in MATLAB as described

previously but without corrections for positional errors and blurring[58]. Colocalization of

tracks was determined by Track Processor Interaction Analysis (distance threshold 2

pixels) in the Spot Tracking plugin. INSR tracks that colocalized with CAV1 or CLTC

tracks for more than 4 continuous frames are determined to be colocalized tracks. The

exported time and duration of colocalization were organized in Python and plotted in R

studio. Diffusion coefficients, track radius, and the lifetime of tracks were plotted in R

studio. Specifically, SuperPlots were used to plot the mean value of the cells and

individual tracks of each cell[59]. Code is available at GitHub deposit

https://github.com/hcen/INSR_tracking.

Skeletal muscle co-immunoprecipitation

Mice received 1.5U/kg insulin or PBS through intraperitoneal injection. Skeletal

muscle (gastrocnemius) was collected after 5 or 10 min of injection. INSR was

immunoprecipitated by incubating 500 μg of protein lysate with 2 μg INSR antibodies

(Rabbit pAb, Cat. #sc-711, Santa Cruz) at 4°C overnight. Subsequently, the solution was

incubated with PureProteome™ Protein G Magnetic Beads (Cat. # LSKMAGG10,

Millipore, Burlington, MA, USA) and washed according to the manufacturer's instructions.

INSR protein complexes were eluted from the beads by incubation in Blue Loading Buffer

(50 mM DTT, Cell Signaling) at 95°C for 5 min and were analyzed by standard western

blot procedures. INSR, CAV1, and CLTC were detected by the following Cell Signaling

primary antibodies: INSRβ (1:1000, Mouse mAb, Cat. #3020S), CLTC (1:1000, Rabbit

mAb, Cat. #4796), Caveolin1 (1:1000, Rabbit mAb, Cat. #3267), and HRP-conjugated

VeriBlot secondary antibodies (1:200, anti-mouse Cat. #ab131368, anti-rabbit ab131366,

Abcam) that do not bind to IgG heavy and light chains to reduce background noise.

Immunoprecipitation-Mass Spectrometry

C2C12 myoblasts were serum-starved for 6 hours and were given 0 or 2 nM insulin

for 15 min before collecting their lysates in RIPA buffer. INSRα antibody (NNC0276-3000,

a gift from NovoNordisk compound sharing program) that does not cross-react with

IGF1R[23] was covalently conjugated to Dynabeads M-270 Epoxy beads (Cat. #14301,

Invitrogen) to immunoprecipitate INSR according to the manufacture's protocol. INSR

protein complexes were eluted from the beads by incubation in Blue Loading Buffer (50

mM DTT, Cell Signaling) at 95°C for 5 min.

Eluted co-immunoprecipitation samples were run into 10% SDS PAGE gel for

clean-up procedure, fixed, stained, and destained in water. Each of the samples was

excised, reduced, alkylated, and digested according to standard in-gel digest protocols[60].

Extracted peptide samples were then cleaned up via STAGE-tip purification, as

previously described[61] .

Samples were reconstituted in 2% ACN, 0.5% formic acid, and the peptides were

analyzed using a quadrupole – time of flight mass spectrometer (Impact II; Bruker

Daltonics) on-line coupled to an EasyLC 1000 HPLC (ThermoFisher Scientific) using a

Captive spray nanospray ionization source (Bruker Daltonics) including Aurora Series

Gen2 (CSI) analytical column, (25cm x 75µm 1.6µm FSC C18, with Gen2 nanoZero and

CSI fitting; Ion Opticks, Parkville, Victoria, Australia), and a µ-Precolumn, 300 µm ID x 5

mm, C18 PepMap, 5 µm, 100 A, Thermo Scientific, Waltham, Massachusetts, United

States). Samples were reconstituted in (0.1% aqueous formic acid and 2 % acetonitrile in

water) and 1ug of sample was loaded, injected in triplicates. A standard 90 min

reversed-phase seperation was employed, using water:acetonitrile gradients.

LC and MS acquisition parameters were standard for peptide sequencing. Acquired

data were then searched by MaxQuant (V1.6.7.) against the Uniprot protein database for

Mouse C2C12 cell line, and LFQ intensities extracted and normalized using the MaxLFQ

algorithm, with 20ppm and 30ppm mass accuracies for precursor and product ion

masses, respectively, and a 1% false discovery rate cut-off. Differential expression was

determined by comparing the mean protein intensities of the triplicate injections between

the groups. Significance ($p<0.05$) was calculated via t-test using default parameters in

Perseus (v1.6.14.0).

Statistics

Data were presented as mean ± SEM in addition to the individual data points. A

significance level of $p < 0.05$ was used throughout. Student's t-tests were used for

comparisons between two groups. Comparisons between more than two groups were

performed using multiple comparison p-values were adjusted using the Tukey method.

Comparisons between multiple groups and two categorical variables were performed

using Two-way ANOVA followed by Tukey post hoc test. Statistical analysis above was

performed using GraphPad Prism (Version 9.0.2). Twe-sided Kolmogorov-Smirnov Tests

were used for comparing cumulative probablility, which was performed with ks.text()

function in R.

References

1.   Knutson VP. Cellular trafficking and processing of the insulin receptor. *FASEB J.*
1991;5(8):2130-2138.

2.   Kahn CR, Crettaz M. Insulin receptors and the molecular mechanism of insulin
action. *Diabetes Metab Rev.* 1985;1(1-2):5-32.

3.   Bevan AP, Burgess JW, Drake PG, Shaver A, Bergeron JJ, Posner BI. Selective
activation of the rat hepatic endosomal insulin receptor kinase. Role for the
endosome in insulin signaling. *The Journal of biological chemistry.*
1995;270(18):10784-10791.

4.   Boothe T, Lim GE, Cen H, et al. Inter-domain tagging implicates caveolin-1 in
insulin receptor trafficking and Erk signaling bias in pancreatic beta-cells. *Mol*
Metab. 2016;5(5):366-378.

5.   Trischitta V, Gullo D, Squatrito S, Pezzino V, Goldfine ID, Vigneri R. Insulin
Internalization into Monocytes Is Decreased in Patients with Type II Diabetes
Mellitus. 1986;62:522-528.

6.   Trischitta V, Brunetti A, Chiavetta A, Benzi L, Papa V, Vigneri R. Defects in
insulin–receptor internalization and processing in monocytes of obese subjects
and obese NIDDM patients. 1989;38:1579–1584.

7.   Jochen AL, Berhanu P, Olefsky JM. Insulin internalization and degradation in
adipocytes from normal and type II diabetic subjects. *J Clin Endocrinol Metab.*
1986;62(2):268-274.

8.   Rask-Madsen C, Kahn CR. Tissue-specific insulin signaling, metabolic syndrome,
and cardiovascular disease. *Arterioscler Thromb Vasc Biol.*
2012;32(9):2052-2059.

9.   Guo S. Insulin signaling, resistance, and the metabolic syndrome: insights from
mouse models into disease mechanisms. *The Journal of endocrinology.*
2014;220(2):T1-T23.

10.   Jochen A, Hays J, Lee M. Kinetics of insulin internalization and processing in
adipocytes: effects of insulin concentration. *J Cell Physiol.* 1989;141(3):527-534.

11.   McClain DA, Olefsky JM. Evidence for two independent pathways of
insulin-receptor internalization in hepatocytes and hepatoma cells. *Diabetes.*
1988;37(6):806-815.

12.   Ceresa BP, Kao AW, Santeler SR, Pessin JE. Inhibition of clathrin-mediated
endocytosis selectively attenuates specific insulin receptor signal transduction
pathways. *Mol Cell Biol.* 1998;18(7):3862-3870.

13.   Carpentier JL, Van Obberghen E, Gorden P, Orci L. Surface redistribution of
125I-insulin in cultured human lymphocytes. *J Cell Biol.* 1981;91(1):17-25.

14.   Paccaud JP, Siddle K, Carpentier JL. Internalization of the human insulin receptor.
600   The insulin-independent pathway. *The Journal of biological chemistry.*
601   1992;267(18):13101-13106.

602   15.   Bickel PE. Lipid rafts and insulin signaling. *Am J Physiol Endocrinol Metab.*
603   2002;282(1):E1-E10.

604   16.   Cohen AW, Combs TP, Scherer PE, Lisanti MP. Role of caveolin and caveolae in
605   insulin signaling and diabetes. *Am J Physiol Endocrinol Metab.*
606   2003;285(6):E1151-1160.

607   17.   Tseng LT, Lin CL, Tzen KY, Chang SC, Chang MF. LMBD1 protein serves as a
608   specific adaptor for insulin receptor internalization. *The Journal of biological*
609   *chemistry.* 2013;288(45):32424-32432.

610   18.   Gautier A, Juillerat A, Heinis C, et al. An engineered protein tag for multiprotein
611   labeling in living cells. *Chem Biol.* 2008;15(2):128-136.

612   19.   Cen H, Botezelli JD, Johnson JD. Modulation of Insr and insulin receptor signaling
613   by hyperinsulinemia in vitro and in vivo. *Biorxiv.* 2021.

614   20.   Jans DA. Lateral Mobility of Polypeptide Hormone Receptors and GTP-Binding
615   Proteins. In: *The Mobile Receptor Hypothesis: The Role of Membrane Receptor*
616   *Lateral Movement in Signal Transduction.* Boston, MA: Springer US; 1997:83-115.

617   21.   Kabayama K, Sato T, Saito K, et al. Dissociation of the insulin receptor and
618   caveolin-1 complex by ganglioside GM3 in the state of insulin resistance. *Proc*
619   *Natl Acad Sci U S A.* 2007;104(34):13678-13683.

620   22.   Ikonen E, Vainio S. Lipid microdomains and insulin resistance: is there a
621   connection? *Sci STKE.* 2005;2005(268):pe3.

622   23.   Ørstrup LH, Slaaby R, Rasch MG, et al. Cross-species reactive monoclonal
623   antibodies against the extracellular domains of the insulin receptor and IGF1
624   receptor. *J Immunol Methods.* 2019;465:20-26.

625   24.   Benyoucef S, Surinya KH, Hadaschik D, Siddle K. Characterization of insulin/IGF
626   hybrid receptors: contributions of the insulin receptor L2 and Fn1 domains and the
627   alternatively spliced exon 11 sequence to ligand binding and receptor activation.
628   *Biochem J.* 2007;403(3):603-613.

629   25.   Moxham CP, Duronio V, Jacobs S. Insulin-like growth factor I receptor
630   beta-subunit heterogeneity. Evidence for hybrid tetramers composed of
631   insulin-like growth factor I and insulin receptor heterodimers. *The Journal of*
632   *biological chemistry.* 1989;264(22):13238-13244.

633   26.   Shu L, Lee L, Chang Y, et al. Caveolar structure and protein sorting are
634   maintained in NIH 3T3 cells independent of glycosphingolipid depletion. *Arch*
635   *Biochem Biophys.* 2000;373(1):83-90.

636   27.   DiGiacomo V, Meruelo D. Looking into laminin receptor: critical discussion
637   regarding the non-integrin 37/67-kDa laminin receptor/RPSA protein. *Biol Rev*
638   *Camb Philos Soc.* 2016;91(2):288-310.

28.   Zoubeidi A, Zardan A, Wiedmann RM, et al. Hsp27 promotes insulin-like growth
factor-I survival signaling in prostate cancer via p90Rsk-dependent
phosphorylation and inactivation of BAD. *Cancer Res.* 2010;70(6):2307-2317.

29.   Muranova LK, Shatov VM, Bukach OV, Gusev NB. Cardio-Vascular Heat Shock
Protein (cvHsp, HspB7), an Unusual Representative of Small Heat Shock Protein
Family. *Biochemistry (Mosc).* 2021;86(Suppl 1):S1-S11.

30.   Yuan H, Wang T, Niu Y, Liu X, Fu L. AMP-activated protein kinase-mediated
expression of heat shock protein beta 1 enhanced insulin sensitivity in the skeletal
muscle. *FEBS Lett.* 2017;591(1):97-108.

31.   Chidlow JH, Sessa WC. Caveolae, caveolins, and cavins: complex control of
cellular signalling and inflammation. *Cardiovasc Res.* 2010;86(2):219-225.

32.   Capozza F, Cohen AW, Cheung MW, et al. Muscle-specific interaction of caveolin
isoforms: differential complex formation between caveolins in fibroblastic vs.
muscle cells. *Am J Physiol Cell Physiol.* 2005;288(3):C677-691.

33.   Chang M, Kwon M, Kim S, et al. Aptamer-based single-molecule imaging of
insulin receptors in living cells. *J Biomed Opt.* 2014;19(5):051204.

34.   Hubert M, Larsson E, Lundmark R. Keeping in touch with the membrane; protein-
and lipid-mediated confinement of caveolae to the cell surface. *Biochem Soc*
Trans. 2020;48(1):155-163.

35.   Martins AS, Ordóñez JL, Amaral AT, et al. IGF1R signaling in Ewing sarcoma is
shaped by clathrin-/caveolin-dependent endocytosis. *PLoS One.*
2011;6(5):e19846.

36.   Pandini G, Frasca F, Mineo R, Sciacca L, Vigneri R, Belfiore A. Insulin/insulin-like
growth factor I hybrid receptors have different biological characteristics depending
on the insulin receptor isoform involved. *The Journal of biological chemistry.*
2002;277(42):39684-39695.

37.   Tomas A, Futter CE, Eden ER. EGF receptor trafficking: consequences for
signaling and cancer. *Trends Cell Biol.* 2014;24(1):26-34.

38.   Kammanadiminti SJ, Chadee K. Suppression of NF-kappaB activation by
Entamoeba histolytica in intestinal epithelial cells is mediated by heat shock
protein 27. *The Journal of biological chemistry.* 2006;281(36):26112-26120.

39.   Ku HC, Chang HH, Liu HC, et al. Green tea (-)-epigallocatechin gallate inhibits
insulin stimulation of 3T3-L1 preadipocyte mitogenesis via the 67-kDa laminin
receptor pathway. *Am J Physiol Cell Physiol.* 2009;297(1):C121-132.

40.   Zhao WQ, Chen GH, Chen H, et al. Secretion of Annexin II via activation of insulin
receptor and insulin-like growth factor receptor. *The Journal of biological*
chemistry. 2003;278(6):4205-4215.

41.   Biener Y, Feinstein R, Mayak M, Kaburagi Y, Kadowaki T, Zick Y. Annexin II is a
novel player in insulin signal transduction. Possible association between annexin
II phosphorylation and insulin receptor internalization. *The Journal of biological*
chemistry. 1996;271(46):29489-29496.

42.   Rescher U, Ludwig C, Konietzko V, Kharitonenkov A, Gerke V. Tyrosine
681   phosphorylation of annexin A2 regulates Rho-mediated actin rearrangement and
682   cell adhesion. *J Cell Sci.* 2008;121(Pt 13):2177-2185.
683   43.   Caron D, Boutchueng-Djidjou M, Tanguay RM, Faure RL. Annexin A2 is
684   SUMOylated on its N-terminal domain: regulation by insulin. *FEBS Lett.*
685   2015;589(9):985-991.
686   44.   Lemmon MA, Schlessinger J. Cell signaling by receptor tyrosine kinases. *Cell.*
687   2010;141(7):1117-1134.
688   45.   de Graauw M, Cao L, Winkel L, et al. Annexin A2 depletion delays EGFR
689   endocytic trafficking via cofilin activation and enhances EGFR signaling and
690   metastasis formation. *Oncogene.* 2014;33(20):2610-2619.
691   46.   Behrends M, Engmann O. Linker histone H1.5 is an underestimated factor in
692   differentiation and carcinogenesis. *Environ Epigenet.* 2020;6(1):dvaa013.
693   47.   Vicidomini G, Bianchini P, Diaspro A. STED super-resolved microscopy. *Nat*
694   *Methods.* 2018;15(3):173-182.
695   48.   Fish KN. Total internal reflection fluorescence (TIRF) microscopy. *Curr Protoc*
696   *Cytom.* 2009;Chapter 12:Unit12.18.
697   49.   Le PU, Guay G, Altschuler Y, Nabi IR. Caveolin-1 is a negative regulator of
698   caveolae-mediated endocytosis to the endoplasmic reticulum. *Journal of*
699   *Biological Chemistry.* 2002;277(5):3371-3379.
700   50.   Humphries AC, Way M. The non-canonical roles of clathrin and actin in pathogen
701   internalization, egress and spread. *Nat Rev Microbiol.* 2013;11(8):551-560.
702   51.   Rodal SK, Skretting G, Garred O, Vilhardt F, van Deurs B, Sandvig K. Extraction
703   of cholesterol with methyl-beta-cyclodextrin perturbs formation of clathrin-coated
704   endocytic vesicles. *Mol Biol Cell.* 1999;10(4):961-974.
705   52.   Parpal S, Karlsson M, Thorn H, Strålfors P. Cholesterol depletion disrupts
706   caveolae and insulin receptor signaling for metabolic control via insulin receptor
707   substrate-1, but not for mitogen-activated protein kinase control. *The Journal of*
708   *biological chemistry.* 2001;276(13):9670-9678.
709   53.   Hussain KM, Leong KL, Ng MM, Chu JJ. The essential role of clathrin-mediated
710   endocytosis in the infectious entry of human enterovirus 71. *The Journal of*
711   *biological chemistry.* 2011;286(1):309-321.
712   54.   Schindelin J, Arganda-Carreras I, Frise E, et al. Fiji: an open-source platform for
713   biological-image analysis. *Nat Methods.* 2012;9(7):676-682.
714   55.   de Chaumont F, Dallongeville S, Chenouard N, et al. Icy: an open bioimage
715   informatics platform for extended reproducible research. *Nat Methods.*
716   2012;9(7):690-696.
717   56.   Chenouard N, Bloch I, Olivo-Marin JC. Multiple hypothesis tracking for cluttered
718   biological image sequences. *IEEE Trans Pattern Anal Mach Intell.*
719   2013;35(11):2736-3750.

57.   Chenouard N, Smal I, de Chaumont F, et al. Objective comparison of particle
tracking methods. *Nat Methods.* 2014;11(3):281-289.

58.   Abraham L, Lu HY, Falcão RC, et al. Limitations of Qdot labelling compared to
directly-conjugated probes for single particle tracking of B cell receptor mobility.
Sci Rep. 2017;7(1):11379.

59.   Lord SJ, Velle KB, Mullins RD, Fritz-Laylin LK. SuperPlots: Communicating
reproducibility and variability in cell biology. *J Cell Biol.* 2020;219(6).

60.   Shevchenko A, Wilm M, Vorm O, Mann M. Mass spectrometric sequencing of
proteins silver-stained polyacrylamide gels. *Anal Chem.* 1996;68(5):850-858.

61.   Rappsilber J, Mann M, Ishihama Y. Protocol for micro-purification, enrichment,
pre-fractionation and storage of peptides for proteomics using StageTips. *Nat
Protoc.* 2007;2(8):1896-1906.

Figure legends

Figure 1. Biochemical and live-cell imaging characterization of the effects of

insulin on INSR.

(A) Internalized to total INSR ratio was quantified using surface biotinylation in

undifferentiated C2C12 myoblasts were incubated in PBS (no insulin) or serum-free

DMEM containing 0, 0.2, 2, or 20 nM insulin for 15 min. **(B)** Western blot showing

SNAP-tagged INSR expressed from lentiviral vector in comparison to wild-type INSR or

in C2C12 myoblasts. **(C,D).** Qualitative snapshots of surface INSR-A-SNAP labeled by

Surface Alexa Flour 488 cell non-permeable dye in 0, 0.2 or 20 nM insulin conditions

were imaged in live undifferentiated C2C12 myoblasts from **(C)** 0 - 15 min or **(D)** after 20

min using a spinning disk confocal microscope. Time-lapse images of INSR vesicle

interactions starting from the selected subregions (white squares) of snapshots in are

shown in the *insets.* (n=3 cells)

Figure 2. Live-cell TIRF imaging of cell-surface INSR.

(A) Experimental design of TIRF microscopy of same cells at 3 timepoints with 2 nM

insulin or no insulin. **(B)** Representative TIRF image of INSR-A-EGFP puncta, binary

image of detected INSR spots, and INSR tracks.   **(C)** Diffusion coefficient of

INSR-A-EGFP tracks in control or insulin group. Data is plotted as SuperPlot with bigger

dots showing mean values of the cells and smaller dots showing individual INSR tracks of

the cells. **(D)** The cumulative probability shows the distribution of diffusion coefficient in

(C). **(E)** Track radius of INSR-A-EGFP tracks in control or insulin group. **(F)** The

cumulative probability shows the distribution of diffusion coefficient in **(E)**. (n=3 cells in

control group, n=9 cells in insulin group)

Figure 3. Identification of INSR interactors.

(A) Proteins that are detected in INSR immunoprecipitation mass-spectrometry, and

significantly different between wild-type and INSR knockout groups. Proteins in green

area were specific interactors that had little detection in knockout cells. **(C)** AKT and ERK

phosphorylation in. **(C)** Western blot of mice skeletal muscle lysates after insulin injection.

Phospho-AKT (Ser473) to total AKT ratio or phospho-ERK1/2 to total ERK ratio were

quantified. (n=6) **(D)** Co-immunoprecipitation of skeletal muscle INSR after PBS or insulin

injection.   Interacting CLTC to INSR ratio or CAV3 to INSR ratio were quantified.

(n=7-10)

Figure 4. Colocalization and interaction between INSR, CAV1 and CLTC under

different insulin concentrations.

(A) Representative STED microscopy images of C2C12 myoblasts expressing

INSR-A-SNAP (surface labeled) that were fixed after stimulated with 0 nM, 0.2 nM, or 20

nM insulin for 30 min and stained for CAV1 and CLTC (Scale bar = 5 μm). **(B,C)**

Colocalizations between INSR-A-SNAP, CAV1, and CLTC were quantified by Object

Pearson Coefficient. Data are plotted to show differences between insulin concentrations

(B) or between protein pairs **(C)**. (n=7-9 images, 1-4 cells per image. *p<0.05, Tukey's

multiple comparison after 2-ANOVA. Box represents median and 25th to 75th

percentiles.)

Figure 5. Live-cell TIRF imaging of cell-surface INSR, CAV1 and CLTC.

(A,B)TIRF microscopy showed colocalization between INSR-B-TagBFP and **(A)**

CAV1-mRFP or **(B)** CLTC-mRFP. **(C,D)**The associated fluorescent intensity of

representative INSR-B-TagBFP and colocalized **(C)** CAV1-mRFP or **(D)** CLTC-mRFP. **(E)**

Diffusion coefficient of INSR-B-TagBFP tracks in cells expressing CAV1-mRFP under 0

nM (n=2 cells, 360 tracks) or 2 nM insulin (n=2 cells, 324 tracks) or expressing

CLTC-mRFP under 0 nM (n=4 cells, 917 tracks) or 2 nM insulin (n=2 cells, 146 tracks).

Data is plotted as SuperPlot, with bigger dots showing mean values of the cells and

smaller dots showing individual INSR tracks of the cells. The cumulative probability

shows the distribution of the diffusion coefficient. **(F)** Track radius of INSR-B-TagBFP

tracks in cells expressing CAV1-mRFP under 0 nM (n=2 cells, 360 tracks) or 2 nM insulin

(n=2 cells, 324 tracks) or expressing CLTC-mRFP under 0 nM (n=4 cells, 917 tracks) or 2

nM insulin (n=2 cells, 146 tracks). The cumulative probability shows the distribution of the

track radius. **(G)** Track radius of INSR-B-TagBFP tracks in cells expressing CAV1-mRFP

under 0 nM (n=2 cells, 360 tracks) or 2 nM insulin (n=2 cells, 324 tracks) or expressing

CLTC-mRFP under 0 nM (n=4 cells, 917 tracks) or 2 nM insulin (n=2 cells, 146 tracks).

The cumulative probability shows the distribution of the diffusion coefficient. (# effect of

co-expression, 2-ANOVA.)   Experiments shown were conducted in Ringer's buffer.

Figure 6. Analysis of INSR tracks colocalized or not colocalized with CAV1 or

CLTC.

(A,B) TIRF microscopy showed colocalization between INSR-B-TagBFP and **(A)**

CAV1-mRFP or **(B)** CLTC-mRFP membrane**(A)** INSR tracks that colocalized with CAV1

under 0 or 2 nM insulin (referred to as CAV1.Y.0nM or CAV1.Y.2nM below). **(B)** INSR

tracks that colocalized with CLTC under 0 or 2 nM insulin (referred to as CLTC.Y.0nM or

CLTC.Y.2nM below). **(C)** Diffusion coefficient of INSR-B-TagBFP tracks that did not

colocalize with CAV1-mRFP tracks under 0 nM insulin (CAV1.N.0nM) or 2 nM insulin

(CAV1.N.2nM) or colocalized with CAV1-mRFP tracks under 0 nM insulin (CAV1.Y.0nM)

or 2 nM insulin (CAV1.Y.2nM). The cumulative probability shows the distribution of

diffusion coefficient. **(D)** Diffusion coefficient of INSR-B-TagBFP tracks that did not

colocalize with CLTC-mRFP tracks under 0 nM insulin (CLTC.N.0nM) or 2 nM insulin

(CLTC.N.2nM) or colocalized with CLTC-mRFP tracks under 0 nM insulin (CLTC.Y.0nM)

or 2 nM insulin (CLTC.Y.2nM). The cumulative probability shows the distribution of

diffusion coefficient. **(E)** Track radius of INSR-B-TagBFP tracks that did not colocalize

with CAV1-mRFP tracks under 0 nM insulin (CAV1.N.0nM) or 2 nM insulin (CAV1.N.2nM)

or colocalized with CAV1-mRFP tracks under 0 nM insulin (CAV1.Y.0nM) or 2 nM insulin

(CAV1.Y.2nM). The cumulative probability shows the distribution of track radius **(F)** Track

radius of INSR-B-TagBFP tracks that did not colocalize with CLTC-mRFP tracks under 0

nM insulin (CLTC.N.0nM) or 2 nM insulin (CLTC.N.2nM) or colocalized with CLTC-mRFP

tracks under 0 nM insulin (CLTC.Y.0nM) or 2 nM insulin (CLTC.Y.2nM). The cumulative

probability shows the distribution of track radius. **(G)** Lifetime of INSR-B-TagBFP tracks

that did not colocalize with CAV1-mRFP tracks under 0 nM insulin (CAV1.N.0nM) or 2 nM

insulin (CAV1.N.2nM) or colocalized with CAV1-mRFP tracks under 0 nM insulin

(CAV1.Y.0nM) or 2 nM insulin (CAV1.Y.2nM). The cumulative probability shows the

distribution of lifetime. **(H)** Lifetime of INSR-B-TagBFP tracks that did not colocalize with

CLTC-mRFP tracks under 0 nM insulin (CLTC.N.0nM) or 2 nM insulin (CLTC.N.2nM) or

colocalized with CLTC-mRFP tracks under 0 nM insulin (CLTC.Y.0nM) or 2 nM insulin

(CLTC.Y.2nM). The cumulative probability showing the distribution of lifetime.

(CAV1.N.0nM, n=2 cells, 125 tracks; CAV1.N.2nM, n=2 cells, 96 tracks; CAV1.Y.0nM,

n=2 cells, 235 tracks; CAV1.Y.2nM, n=2 cells, 228 tracks; CLTC.N.0nM, n= 4 cells,   597

tracks; CLTC.N.2nM, n= 2 cells,   91 tracks; CLTC.Y.0nM, n= 4 cells, 320 tracks;

CLTC.Y.2nM, n= 2 cells, 55 tracks. # effect of colocalization, $ effect of insulin, %

interaction of the 2 variables, 2-ANOVA.) Experiments shown were conducted in Ringer's

buffer.

Figure 7. Summary of the effects of CAV1 and CLTC on INSR tracks. Higher insulin

(20 nM) promoted INSR and CAV1 colocalization, while lower insulin (0.2 nM) promoted

INSR and CLTC colocalization. INSR tracks had lower diffusion coefficient and track

radius in CAV1-overexpressing cells than CLTC-overexpressing cells. Within the same

cells, INSR tracks colocalized with CAV1 had longer lifetime and larger track radius than

non-colocalized tracks. INSR tracks colocalized with CLTC had longer lifetime than

non-colocalized tracks, which is further increased by insulin.

Figure 1. Biochemical and live-cell imaging characterization of the effects of insulin on INSR.

Figure 2. Live-cell TIRF imaging of cell-surface INSR.

Figure 3. Identification of INSR interactors.

Figure 4. Colocalization and interaction between INSR, CAV1 and CLTC under different insulin concentrations.

Figure 5. Live-cell TIRF imaging of cell-surface INSR, CAV1 and CLTC.

Figure 6. Analysis of INSR tracks colocalized or not colocalized with CAV1 or CLTC.

Figure 7. Summary of the effects of CAV1 and CLTC on INSR tracks.