Title: Insulin receptor trafficking and interactions in muscle cells

Authors: Haoning Howard Cen[1,2], Aurora J. Mattison[1,2], Alireza Omidi[3,4], Jason

Rogalski[2,5], Libin Abraham[2,6], Guang Gao[2,7], Michael R. Gold[2,6], Leonard J. Foster[2,3,4],

Jörg Gsponer[3,4], James D. Johnson[1,2,8]*

Affiliation:

1 Department of Cellular and Physiological Sciences, University of British Columbia,

Vancouver, BC. Canada

2 Life Sciences Institute, University of British Columbia, Vancouver, BC. Canada

3 Department of Biochemistry and Molecular Biology, University of British Columbia,

Vancouver, BC, Canada

4 Michael Smith Laboratories, University of British Columbia, Vancouver, BC, Canada

5 The Proteomics and Metabolomics Core Facility, University of British Columbia,

Vancouver, BC, Canada

6 Department of Microbiology and Immunology, University of British Columbia, Vancouver,

BC, Canada

7 Life Sciences Institute Imaging Core Facility, University of British Columbia, Vancouver,

BC, Canada

8 Vancouver Coastal Health Research Institute, Vancouver, BC, Canada

ORCIDs: Haoning Howard Cen 0000-0001-8045-1335; Aurora J. Mattison

0000-0001-5688-5109; Alireza Omidi 0000-0003-3598-9480; Jason Rogalski

0009-0000-5736-2991; Libin Abraham 0000-0002-4725-3502; Guang Gao

0000-0001-7523-2234; Michael R. Gold 0000-0003-1222-3191; Leonard J. Foster

0000-0001-8551-4817; James D. Johnson 0000-0002-7523-9433

Keywords: insulin, insulin receptor, caveolin, clathrin, endocytosis

*Address correspondence to: James D. Johnson, Ph.D., Professor. Diabetes Research Group, Dept. of Cellular and Physiological Sciences & Dept. of Surgery, University of British Columbia, 5358 Life Sciences Building, 2350 Health Sciences Mall, Vancouver, BC, Canada, V6T 1Z3

E-mail: James.D.Johnson@ubc.ca; Fax: (604) 822-2316

Twitter: @JimJohnsonSci

Acknowledgments: Spinning disk confocal and STED microscopy were performed in the LSI IMAGING facility of the Life Sciences Institute of the University of British Columbia using infrastructure funded by the Canadian Foundation of Innovation and BC Knowledge Development Fund as well as a Strategic Investment Fund (Faculty of Medicine, University of British Columbia). We thank Dr. Ivan Robert Nabi at UBC for guidance on imaging experiments and helpful discussions. Mass spectrometry used here was supported by the Canada Foundation for Innovation and Genome Canada (264PRO). We thank Yuanyuan Amanda Yang for guidance on Python and R coding, Yi Han Xia for assistance in R coding, and Jiashuo Aaron Zhang for assistance in data analysis. We thank Michael G. Atser for assistance in western blots. INSR antibody NNC0276-3000 was provided by Novo Nordisk Compound Sharing. The schematics in Figure 7 were created with BioRender.

Contributions: HHC designed the study, performed experiments, analyzed data, and

wrote the manuscript. AJM contributed to STED experiments and analyzed data. AO

performed the AlphaFold-Multimer analysis and wrote part of the manuscript. JR

performed mass spectrometry, analyzed data, and edited the manuscript. LA and MG

contributed to the MATLAB scripts and the analysis of single-particle tracking and edited

the manuscript. GG contributed to the spinning disk confocal and STED microscopy and

edited the manuscript. LF contributed to the design of the mass-spectrometry

experiments and edited the manuscript. JG supervised the AlphaFold-Multimer analysis

and edited the manuscript. JDJ designed the study, supervised the research, and edited

the manuscript. JDJ is the ultimate guarantor of this work.

Funding: This study was funded by a Canadian Institutes of Health Research Operating

Grant (P-133692) to JDJ.

Conflict of interest: The authors declare that they have no conflicts of interest with the

contents of this article.

Abstract

Insulin resistance contributes to type 2 diabetes and can be driven by hyperinsulinemia. Insulin receptor (INSR) internalization and cell-surface dynamics at rest and during insulin exposure are incompletely understood in muscle cells. Using surface labelling and live-cell imaging, we observed robust basal internalization of INSR in C2C12 myoblasts, without an effect of added insulin. Mass-spectrometry using INSR knockout cells as controls, identified high-confidence binding partners, including proteins associated with internalization. We confirmed known interactors, including IGF1R, but also identified underappreciated INSR-binding factors such as ANXA2. AlphaFold-Multimer analysis of these INSR-binding proteins predicted potential INSR binding sites of these proteins. Protein-protein interaction network mapping suggested links between INSR and caveolin-mediated endocytosis. INSR interacted with both caveolin and clathrin heavy chain (CLTC) in mouse skeletal muscle and C2C12 myoblasts. Whole cell 2D super-resolution imaging revealed that high levels of insulin (20 nM) increased INSR colocalization with CAV1 but decreased its colocalization with CLTC. Single particle tracking confirmed the colocalization of cell-surface INSR with both over-expressed CAV1-mRFP and CLTC-mRFP. INSR tracks that colocalized with CAV1 exhibited longer radii and lifetimes, regardless of insulin exposure, compared to non-colocalized tracks, whereas insulin further increased the lifetime of INSR/CLTC colocalized tracks. Overall, these data suggest that muscle cells utilize both CAV1 and CLTC-dependent pathways for INSR dynamics and internalization.

Introduction

The insulin receptor (INSR) is one of the most studied membrane proteins due to its

importance in insulin resistance and diabetes. However, many fundamental aspects of

INSR biology remain unclear, including the mechanisms and kinetics of its internalization.

The INSR can be internalized via endocytosis (1,2), and endosomal INSR localization

contributes to the phosphorylation of specific downstream targets when compared with

INSR at the cell surface (3,4). Moreover, ligand-receptor complex dissociation in acidified

endosomes permits insulin degradation, while the receptor can be sorted to recycling or

degradation pathways (1,2). INSR internalization is reduced in monocytes(5,6) and

adipocytes (7) obtained from patients with insulin resistance and type 2 diabetes,

suggesting the importance of this process in disease pathogenesis.

Tissue-specific insulin resistance has been implicated in the development of

diabetes (8,9), but it is less clear whether there are tissue-specific INSR internalization

kinetics or mechanisms. Interestingly, adipocytes internalize insulin at a lower rate in the

presence of high insulin concentrations, as compared to low insulin conditions (10), while

hepatocytes are the opposite (11). The consensus view is that INSR is endocytosed via a

clathrin-coated pit mechanism, based on evidence from a variety of cell types, including

human and rat hepatocytes, fibroblasts, and lymphocytes (11-15). However, other

studies have shown that caveolin-rich membrane lipid rafts, caveolae, are critical for

INSR internalization in other cell types, including adipocytes (16,17). In pancreatic

β-cells, we showed that caveolin-1 (CAV1) mediates INSR internalization, bypassing

clathrin positive compartments (4). However, INSR internalization kinetics and routes are

less well characterized in muscle cells, despite that muscle is one of the major insulin

target tissues. It is estimated that skeletal muscle is a major glucose depot accounting for

60-70% of glucose uptake in response to insulin during hyperinsulinemic-euglycemic

clamp(18). Under postprandial conditions, muscle disposes around 1/3 of orally derived

glucose(19). In muscle cells, one study has shown that clathrin-mediated endocytosis is

involved in INSR internalization(20), but the role of the caveolin-mediated pathway has

not been investigated.

In the present study, we aim to characterize the dynamics of INSR internalization

and endocytosis routes in muscle cells. We employed inter-domain tagged INSR with

fluorescent proteins (4) and SNAP-tag (21), surface biotinylation assays, as well as

super-resolution and TIRF imaging, to explore the dynamics of INSR internalization and

the effects of ambient insulin. We also used proteomics and AlphaFold-Multimer to

identify INSR interacting proteins and defined the relationship between INSR and

caveolin or clathrin. Together, our data show that INSR exhibited a high degree of

insulin-independent internalization in C2C12 myoblasts.

Results

Effects of insulin on INSR internalization

First, we characterized INSR internalization in C2C12 myoblasts under basal

conditions and in the context of added insulin. Circulating insulin in humans oscillates in a

range between approximately 0.01 nM and 0.75 nM (22-27). As for mice, fed insulin is

~0.2 nM in lean mice and ~3-5 nM in extremely obese mice (28). We used insulin

concentrations of 0.2, 2, or 20 nM to cover a spectrum from physiological to

supraphysiological concentrations. A higher rate of basal INSR internalization was

observed in high-glucose serum-free DMEM medium when compared with PBS. The net

INSR internalization over 15 min under various insulin concentrations in the DMEM

medium did not significantly differ from the basal condition (Fig. 1A). Therefore, INSR has

a high rate of constitutive internalization in this cell type, which may be enhanced by the

nutrients in the medium, but that is largely independent of acute exposure to insulin

across a range of physiological doses. These observations in myoblasts are consistent

with our previous measurements in differentiated C2C12 myotubes, in which INSR

internalization in insulin-free DMEM media is evident; insulin-stimulated internalization

over time is also obvious but does not significantly differ at different insulin doses (29).

Qualitative analysis of INSR kinetics in the context of different insulin doses

We also visualized INSR internalization at single-cell level using

inter-domain-tagged INSR, which we have previously found more faithfully recapitulates

the sub-cellular localization of INSR compared to terminal fusion proteins (4). The

inter-domain-tagged INSR was also shown to have functional AKT and ERK signaling in

C2C12 (Fig. S1 (30)) and other cell lines (4). C2C12 myoblast cells with stable

overexpression of SNAP-tagged INSR isoform A (INSR-A-SNAP; Fig. 1B) were labelled

using a cell non-permeable SNAP-Surface Alexa Flour 488 dye (Fig. 1C). Although a few

INSR vesicles seemed to be internalized during the small gap between labelling and

imaging, and were present at time zero, substantially more internalized INSR vesicles

were observed at later timepoints (Fig. 1C). Live-cell time-lapse imaging of

surface-labelled INSR-A-SNAP at the middle plane of cells by spinning disk confocal

microscopy revealed INSR internalization in the context of 0, 0.2, and 20 nM insulin over

15 min (Fig. 1C, Video S1 (30)). Consistent with the surface biotinylation assay results in

Figure 1A, we did not observe a clear difference in INSR internalization in the presence of

insulin in these cells. Internalized INSR vesicles exerted dynamic movement and

interaction, which was more evident between 20-30 min (Fig. 1D-F, Video S2 (30)).

INSR-containing vesicles exhibited many apparent fusion and fission events regardless

of insulin concentrations. Some smaller vesicles merged into a larger vesicle that

remained in the field, undergoing limited and apparently random motion (one example is

shown in Fig. 1F), suggestive of a fusion event. A caveat associated with 2D confocal

imaging is limited Z resolution, raising the possibility that the observed INSR vesicles are

not all in the same optical plane. Our experiments using surface labelling of SNAP-tagged

INSR permitted the visualization of the dynamic behaviour of internalized INSR vesicles,

and these qualitative observations motivated the more quantitative study below.

Quantitative analysis of INSR kinetics in the context of insulin

Lateral movement of tyrosine kinase receptors has been linked to their functional state. Ligand binding of some monomeric receptors, such as EGFR, promotes their dimerization and aggregation and decreases their lateral mobility (31). Although INSRs are thought to already exist as dimers in plasma membrane, aggregation of INSR by antibodies can stimulate INSR signaling (31), and INSR may need to localize to membrane lipid rafts for normal signaling (32,33). Therefore, we sought to quantitatively assess whether insulin affects the lateral movement of INSR in or near the plasma membrane. To do this, we used time-lapse TIRF microscopy of C2C12 myoblast transiently expressing inter-domain-tagged INSR-A-EGFP. We imaged the same cells at 0, 6, or 11 min in the presence of 2 nM insulin, as well as a control group without insulin to control for other variables such as photobleaching (Fig. 2A). Single-particle tracking of INSR-A-EGFP (Fig. 2B) was used to calculate their diffusion coefficient (Fig. 2C,D) and track radius (Fig. 2E,F). Insulin did not have a big impact on the track radius or diffusion coefficient, despite some very subtle differences in the cumulative probability curves with very close median values (Fig. 2D,F). Together with the qualitative data reported above, these observations suggest that INSR mobility is largely unaffected by acute insulin stimulation in this cell model.

Novel INSR interactors identified by immunoprecipitation and mass spectrometry

To explore the molecular mechanisms associated with INSR internalization, we conducted co-immunoprecipitation followed by mass-spectrometry to identify INSR interactors. We used INSR knockout C2C12 myoblasts as a negative control and an INSR antibody that does not cross-react with Insulin-like growth factor 1 receptor (IGF1R)(34) to identify INSR interactors with higher specificity than any previously reported study. After filtering the detected proteins against the INSR knockout cell control

lysates, we identified INSR and five high-confidence INSR-binding proteins (Fig. 3A). The

immunoprecipitated abundance of these detected interactors was not significantly

changed by the addition of 2 nM insulin (Fig. 3A). INSR is well established to form

heterodimers with IGF1R(35-37), and its immunoprecipitation represented a type of

internal positive control for this study. The other detected proteins are not well-known

INSR interactors. To discover whether any of the identified INSR-binding proteins might

be connected with endocytosis machinery, protein-protein interaction (PPI) network was

constructed between these co-immunoprecipated proteins and all the proteins under the

"Endocytosis" pathway in Kyoto Encyclopedia of Genes and Genomes (KEGG). We

manually extracted a subnetwork that connected all our identified proteins, except for

HIST1H1B which did not connect with any proteins in this network (Fig. 3B). In the

resultant PPI network, annexin A2 (ANXA2) is connected with CAV1 and CAV3 (Fig. 3B),

and it has been reported to localize in caveolae (38). Ribosomal protein SA (RPSA) is a

laminin receptor reported to localize in plasma membrane lipid rafts (39). Heat shock

protein B1 (HSPB1; a.k.a. HSP27) is exclusively found in insulin-responsive tissues and

has been implicated in insulin sensitivity and INSR/IGF1R signalling (40-42). The PPI

network and cellular localization of these interactors pointed to the possibility that INSR

may also associate with caveolins and/or related proteins in myoblasts.

Modelling INSR interactions *in silico* using AlphaFold-Multimer

AlphaFold-Multimer has been used to accurately model protein complexes (43,44).

Recent work showed that AlphaFold-Multimer more accurately predicts interactions of

protein fragments than lengthy full proteins and established a new pipeline to identify

potential interacting fragments/sites (45). We applied this new pipeline to infer whether

and how the putative INSR interactors identified with mass spectroscopy directly bind to

INSR. We included a mitochondrial protein MTCO2 (aka COX2), expected to have no

interaction with INSR, as a negative control in these *in silico* experiments. Several known

or expected interactor of INSR were also tested, including IRS1, AP2M1, CAV1 and

CAV3. First, we ran AlphaFold-Multimer on the cytoplasmic region of INSR and each of

the full-length interactors to predict a protein complex structure. We made five separate

predictions for each pair of proteins, shown as their predicted template modeling (pTM)

and interface-pTM (ipTM) scores (Fig. 3C). Positive interaction with IRS1 and borderline

interactions (ipTM ≥ 0.42 as suggested in (45)) with AP2M1 and IGF1R are predicted (Fig.

3C). CAV1, CAV3 and all other interactors did not have significant scores. The

predictions with full-length proteins allowed us to extract per-residue scores of minimum

expected distance (minD), minimum inter-predicted aligned error (min-iPAE), and the

predicted local distance difference test (pLDDT) for INSR and all interactors (Fig. 3D and

Fig. S2 (30)). minD and min-iPAE scores pinpoint potential interaction sites, with minD

being the most accurate predictor of interaction sites between two proteins (45). pLDDT

is a measure of local confidence and can indicate whether the region of interest is

structured or unstructured. Using the INSR and AP2M1 interaction as an example, we

observed that AlphaFold-Multimer predicts a sharp peak in all scores at residue 1350 of

INSR, suggesting that the region around residue 1350 is in contact with AP2M1 (Fig. 3D).

Yet this is not reflected in their full-length protein predictions (highest ipTM = 0.5, Fig. 3C),

demonstrating that AlphaFold-Multimer may detect the interaction site between two

proteins but is not able to predict the interaction when full-length proteins are used. To

overcome this limitation, proteins can be fragmented into shorter segments, which are

then submitted to AlphaFold-Multimer for prediction. This fragmentation can be guided by

minD peaks that identify potential interaction sites. Thus, we cut both the INSR and the

interactor proteins into fragments of 30 residues surrounding minD peaks (45). Doing so,

we saw a large increase in ipTM scores for IGF1R, IRS1, AP2M1, CAV3, but not CAV1

(Fig. 3E). Interestingly, although we did not find positive predictions for CAV1 direct

interaction, a peak in all scores (minD, min-iPAE and pLDDT) of CAV1 was detected

around a phosphotyrosine residue 25, which might be an interacting site (Fig. S2 (30)).

Predictions via fragmentation also suggest interactions between INSR and ANXA2, H1-5,

HSBP1, and RPSA (Fig. 3E). The ipTM scores of the negative control MTCO2 remained

low, supporting the confidence in the positive predictions (Fig. 3E). Overall, the

AlphaFold-Multimer predictions support the direct interactions between INSR and the

novel interactors and identify potential interacting sites.

Insulin-dependent interaction between INSR and CAV3 or CLTC *in vivo*

INSR has not been shown to utilize both caveolin and clathrin-mediated endocytosis

in the same tissue; therefore, we sought to confirm that INSR could interact with both

caveolin and clathrin in skeletal muscle *in vivo*. Skeletal muscle was collected 5 or 10 min

after intraperitoneal injection of PBS or insulin (1.5 U/kg), an insulin dose and timepoints

sufficient to measure acute insulin signalling via both AKT and ERK signalling pathways

(Fig. 4A). Acute insulin administration did not change the amount of INSR that could be

immunoprecipitated from skeletal muscle, nor did it alter the association between INSR

and CLTC (Fig. 4B). However, 10 min after insulin injection, there was a significantly

increased interaction between INSR and caveolin, specifically, the mature skeletal

muscle-specific isoform CAV3 (46,47) (Fig. 4B). These data show that acute insulin

signalling results in an increased association between INSR and CAV3 *in vivo*, but does

not define the cellular context for this interaction.

Super-resolution imaging of INSR, CAV1, CLTC, and AXNA2 interactions in the

presence of insulin

We next used super-resolution STED imaging to examine colocalization between

INSR and clathrin (CLTC) or caveolin (C2C12 myoblasts only express CAV1 before

differentiation (48)). C2C12 myoblast cells with stable overexpression of INSR-A-SNAP

were labelled using a cell-nonpermeable SNAP substrate to image only

surface-associated and newly internalized INSR. Cells were incubated with 0, 0.2 or 20

nM insulin for 30 min, and then fixed. Endogenous CLTC and CAV1 were then labelled

using immunofluorescent antibody staining (Fig. 4C). The colocalization between

INSR-SNAP, CAV1, and CLTC was quantified by Object Pearson Coefficient (Fig. 4D,E).

The degree of colocalization between INSR and CAV1 was similar in the presence of 0 or

0.2 nM insulin but was higher under 20 nM insulin (Fig. 4D). However, INSR and CLTC

colocalized more at 0.2 nM insulin, but less at 20 nM insulin, compared to 0 nM insulin

(Fig. 4D). CAV1 and CLTC do not have known biological interactions; therefore, their low

level of colocalization served as an internal negative control and, indeed, it did not

change under different insulin levels (Fig. 4D). Using the same data, the relative

colocalizations between INSR and CAV1 or CLTC were also considered at different

insulin concentrations, further illustrating how high insulin concentrations drive INSR to

colocalize with CAV1 (Fig. 4E). At 0.2 nM, insulin promoted INSR interaction with CLTC

whereas 20 nM insulin redirected the INSR from CLTC to CAV1. Together, these data

revealed the insulin-dependent interaction of INSR with both CAV1 and CLTC *in vivo* and

in vitro.

Since our novel interactor ANXA2 is a target of insulin signaling (49,50), we also

examined its insulin-dependent colocalization with INSR using the same SNAP labelling

method and STED microscopy. Surface labeled INSR was allowed to internalize for 15 or

30 min in the presence of 0, 0.2, 2 or 20 nM insulin. Endogenous ANXA2 proteins were

then labelled using antibody immunofluorescence. Colocalized INSR and ANXA2 puncta

were observed in all conditions (Fig. 5A,C). The degree of colocalization quantified by

Object Pearson Coefficient remained to be moderate in most conditions (lower than the

INSR/CAV1 or INSR/CLTC interactions above) (Fig. 5B,D), except that 0.2 nM insulin

seemed to decrease their colocalization at 30 min (Fig. 5D). This observation suggests

that INSR and ANXA2 interactions are moderate and may be negatively regulated by

insulin signalling.

Effects of CAV1 and CLTC on INSR mobility

To better understand the interplay between INSR, CAV1, and CLTC, we co-expressed INSR-B-TagBFP with CAV1-mRFP or CLTC-mRFP in C2C12 myoblasts and imaged them by TIRF time-lapse microscopy. Colocalizations between INSR and CAV1 or CLTC were observed (Fig. 6A,B). Some colocalized puncta moved or disappeared together with closely associated intensity, referred to here as colocalized tracks (examples in Fig. 6C,D). Tracks were considered as colocalized when the distance between the centers of particle signals was less than 2 pixels for more than 4 continuous frames. These observations suggested that INSR interacted with both CAV1 and CLTC on the cell surface.

To define the effects of CAV1 or CLTC over-expression on INSR movement, we first conducted single particle tracking on all INSR puncta. Similar to our observations above (Fig. 2), insulin did not alter the diffusion coefficient or track radius of INSR. Cells with CAV1 co-expression had lower diffusion coefficients and track radius of INSR compared to cells with CLTC co-expression, regardless of ambient insulin exposure (differences in medians are at least ~2 fold for diffusion coefficients and ~1.5 fold for track radius; Fig. 6E,F). CLTC-expressing cells under 0 nM insulin had higher diffusion coefficients than those under 2 nM insulin (~1.6 fold difference in median; Fig. 6E), as well as shorter INSR lifetimes than all other groups (>1.5 fold difference in median; Fig. 6G), which might suggest a more rapid internalization. Together, these data suggest that CAV1 may restrain INSR movement, relative to CLTC.

Finally, we performed an analysis within the same cells, comparing INSR tracks that were colocalized or not colocalized with either CAV1 or CLTC tracks, to further dissect the effects of CAV1 or CLTC. INSR appeared to spend more time associated with CAV1, and for longer periods, compared with CLTC (Fig. 7A,B). There were no clear effects of time on the association between INSR and CAV1 or CLTC across different moments of

their lifetimes together (Fig. 7A,B). We compared the colocalized and the non-colocalized

INSR tracks, for both CAV1 and CLTC, and found no significant differences in mean

diffusion coefficients (Fig. 7C,D), except that in CLTC-expressing cells without insulin,

non-colocalized INSR had higher diffusion coefficients than CLTC-colocalized INSR,

based on the cumulative probability curve (Fig.7D). The track radius of INSR colocalized

with CAV1 was larger than non-colocalized tracks regardless of insulin (Fig. 7E), but

colocalization with CLTC had significant effects (Fig. 7F). Strikingly, the lifetimes of INSR

tracks colocalized with CAV1 were much longer than non-colocalized tracks regardless of

insulin (3.8~4 fold differences in medians; Fig. 7G). INSR tracks colocalized with CLTC

also had longer lifetimes than non-colocalized tracks (~2.5 fold difference in median), and

insulin also further increased the lifetime of colocalized tracks (Fig. 7H). Overall, our

live-cell imaging of INSR confirmed its distinct interactions with either CAV1 or CLTC

around the cell surface.

Discussion

The goal of this study was to explore the INSR trafficking dynamics and interactions

in a muscle cell model. We found that INSR was constitutively internalized both in the

absence and presence of insulin. We used high-stringency proteomics and

AlphaFold-Multimer to identify specific INSR binding proteins in this cell type and PPI

network analysis to predict a role for caveolin in INSR internalization. Biochemical

experiments, super-resolution imaging, and TIRF imaging suggested both caveolin and

clathrin are involved in INSR internalization and cell-surface trafficking (Fig. 8).

The constitutive internalization of INSR has been described in lymphocytes and

shown not to require ligand binding or kinase activity (although insulin-stimulated

autophosphorylation of INSR could accelerate this process (14,51)). In differentiated

C2C12 myotubes, INSR phosphorylation levels were constitutively high, with little

additional activation by insulin across the physiological range (29), consistent with the

lack of stimulation of INSR internalization by acute insulin treatment in this study. Surface

labeling by SNAP-tag enabled clear visualization of internalized INSR-containing vesicles,

permitting the study of INSR endosomal sorting and traffic. The lateral movement of INSR

in or near the plasma membrane was also not dramatically affected by insulin, consistent

with a previous study that reported unchanged INSR diffusion coefficient with or without

insulin in HEK293T, HepG2, and human IR-overexpressing Rat-1 cells (52). CAV1 may

have a role in decreasing the mobility of INSR, perhaps due to their restricted lateral

movement in the plasma membrane owing to actin cytoskeleton connections through

Filamin-A (53). Accumulation of ganglioside GM3 in TNFα-induced insulin resistance was

associated with the elimination of INSR from the caveolae and increased INSR mobility

on the cell surface (32).

The possibility of more than one internalization pathway for INSR in some cell types

has been understudied. Dual clathrin and caveolin endocytosis routes have been

suggested for IGFR in Ewing sarcoma cells (54), which has a similar structure and

function as INSR and can form heterodimers with INSR (55). The current study supports

the concept that INSR may also utilize both clathrin- and caveolin-mediated endocytosis

in the same cell type. In hepatocytes, it has been suggested that INSRs are internalized

in two functionally separate pathways based on the much more rapid recycling of

receptors internalized by high concentrations of insulin (17.2 nM) versus low insulin (0.17

nM); treatments known to inhibit coated pit formation blocked internalization of the

insulin-INSR complex at low but not high insulin concentrations (11). Most studies of

INSR internalization examined only one route. The mechanisms of coordinated multiple

internalization routes, implied by the insulin-dependent clathrin and caveolin interactions

we observed in muscle cells, might direct INSR to different destinations (e.g. recycle or

degradation)(56) and should be comprehensively characterized in more cell types to

assess potential roles in tissue-specific insulin resistance. Generally, we expect our

results will translate to other muscle cell models and human muscle, given the strong

evolutionary conservation of insulin signaling (57), but our results should be validated in

human muscle in future studies.

We used stringent proteomics and INSR CRISPR knockout control cells to identify

known, novel, and under-appreciated interactors of INSR. As expected, we identified

IGF1R, an established hetero-dimerization partner of INSR. We also demonstrated the

specific binding of HSBP1 (HSP27), which is involved in tyrosine kinase signalling

pathways downstream of p38 MAPK, ERK, and MAPKAPK5 (58) and has been

previously linked to insulin sensitivity (40-42). We demonstrated an interaction between

INSR and RPSA, which has been previously co-immunoprecipitated with INSR and

insulin receptor substrates in 3T3L1 preadipocytes (59). In other cell types, RPSA drives

ERK signalling and cytoskeletal organization, with nuclear functions affecting cell growth,

survival, migration, and protein synthesis (39). ANXA2 is not a known interactor of muscle

INSR, but it has been reported in NIH3T3 fibroblasts (60). Other studies found that INSR

signalling can tyrosine-phosphorylate (49,50) and SUMOylate (61) AXNA2, which

mediates insulin-stimulated actin rearrangement. Interestingly, internalized INSR seemed

necessary for AXNA2 phosphorylation because mutant INSR that was incapable of

internalization but normal for tyrosine kinase signalling failed to phosphorylate AXNA2

(49). AXNA2 also interacts with endosomal and adaptor prortein APPL2, which interacts

with INSR (62,63). The PPI network showed that ANXA2 has direct connection with the

epidermal growth factor receptor (EGFR) which belongs to the same receptor tyrosine

kinase superfamily as INSR (64). ANXA2 knockdown halted EGFR endocytosis beyond

the early endosome, which was accompanied by enhanced EGFR and AKT

phosphorylation (65). These data support the possibility that ANXA2 may have a role in

INSR trafficking and signalling that is worth investigating. Interestingly, evidence

compiled on the Common Metabolic Disease Knowledge Portal shows that ANXA2

genetic variation is strongly associated with fasting C peptide levels, pointing to a causal

role, direct or indirect, in hyperinsulinemia/insulin resistance. Both *Anxa2* and *Rpsa*

mRNAs were upregulated by prolonged insulin in C2C12 myotubes (8). HIST1H1B

belongs to the family of linker histones that is important in chromatin organization (66),

but the significance of this INSR interaction requires further investigation. Our stringent

proteomic analysis, together with AlphaFold-Multimer modelling, provides evidence for a

limited number of proteins that are likely to be direct interactors with INSR in myoblast

cells, and demonstrates that the majority of these interactors were not significantly

altered by insulin exposure.

There are limitations and unexplained observations in the current study. Although

our inter-domain-tagged INSR has functional signalling and internalization, it tends to

have higher pro-INSR bands (Fig. 1B)(4), which is indicative of a potential processing

issue. Nonetheless, higher pro-INSR could also occur in wildtype C2C12 myoblasts (Fig.

S1A (30)), possibly due to stress (67), which may be a co-culprit in the processing issue

of the overexpressed tagged INSR. STED superresolution microscopy usually has a

resolution of 40-50 nm and might be able to reach 20 nm resolution in the best scenario,

and primary and secondary antibody labelling might also add ~30nm distance and reduce

the accuracy (68). TIRF microscopy has diffraction-limited lateral resolution (~200 – 300

nm) with the thickness of the excitation depth around 100 nm (69). Therefore, the

colocalization analyses may not confirm the interactions of the proteins. Nonetheless, the

insulin-dependent colocalization observed in STED microscopy and the track

colocalization over multiple frames observed in TIRF microscopy are evidence that the

proteins had interactions instead of random colocalization. This is further supported by

the AlphaFold-Multimer predictions. The CAV1- or CLTC-colocalized INSR tracks had

longer lifetimes than non-colocalized tracks, which conflicted with the assumption that

CAV1 or CLTC internalized INSR, but instead, they seemed to retain INSR on the

membrane. There are multiple possible explanations; for example, overexpressed CAV1

might stabilize caveolae (70), or CLTC could form stable flat lattices, termed clathrin

plaques (71), which might have novel involvement in INSR internalization or signalling.

As we did not perform loss-of-function studies to confirm the roles of caveolin and clathrin

in INSR endocytosis, further research is needed to elucidate the roles of specific INSR

internalization routes on insulin actions. For example, inhibition of clathrin-mediated

endocytosis of INSR by knocking down LMBD1 in myoblasts increased activation of the

INSR and AKT (20). Our lab discovered that reducing CAV1-dependent INSR

internalization blocked the activation of ERK, but not AKT, signalling in pancreatic

beta-cells (4). Inhibition of INSR internalization in adipocytes (72-74) or hepatocytes

selectively blunted different signaling branches (12,75). More recent advances in INSR

endocytosis mechanisms and consequences are reviewed in (76,77). For example, INSR

endocytosis is found to be regulated by the spindle checkpoint proteins (MAD2, CDC20,

BUBR1 and p31^{comet})(78,79) and SHP1-MAPK pathway (80) in the liver and by the

inceptor protein in beta cells (81,82); decreased insulin-stimulated endocytosis or

increased insulin-independent endocytosis by manipulating these proteins can promote

or impair insulin signaling, respectively. Therefore, INSR internalization enables selective

activation of downstream signalling pathways, and in a tissue-specific manner. Future

study could be done on human muscle cell lines or biopsies to confirm the interactions

and their functional roles.

In summary, this study provides important information on the internalization of INSR

in myoblast cells and defines the stringent INSR interactome. We also provide evidence

using a variety of methods that INSR interacts with both clathrin and caveolins in this cell

model. The current study can serve as a stepping stone for more detailed research on

how multiple internalization routes can provide cell-type customized regulations on INSR

internalization.

Materials and Methods

Cell culture

The C2C12 mouse myoblast (ATCC cell line provided by Dr. Brain Rodrigues, University of British Columbia, Vancouver, Canada) was maintained in Dulbecco's modified Eagle's medium (DMEM, Gibco) supplemented with 10% (v/v) fetal bovine serum (FBS, Gibco), and 1% penicillin-streptomycin (100 µg/ml; Gibco). All downstream experiments were conducted after 6-hour serum starvation in serum-free medium (DMEM supplemented with 1% penicillin-streptomycin), except as otherwise indicated.

Western blot analyses

C2C12 myotubes or mice skeletal muscle (gastrocnemius) tissues were sonicated in RIPA buffer (50 mM β-glycerol phosphate, 10 mM HEPES, 1% Triton X-100, 70 mM NaCl, 2 mM EGTA, 1 mM Na_3VO_4, and 1 mM NaF) supplemented with complete mini protease inhibitor cocktail (Roche, Laval, QC). Lysates were incubated in Blue Loading Buffer containing 50 mM DTT (Cell Signaling, Danvers, MA) at 95°C for 5 min and resolved by SDS-PAGE. Proteins were then transferred to PVDF membranes (BioRad, CA) and probed with antibodies against p-ERK1/2 (Thr202/Tyr204) (1:1000, Cat. #4370, RRID: AB_2315112), ERK1/2 (1:1000, Cat. #4695, RRID: AB_390779), p-AKT (Ser473) (1:1000, Cat. #9271, RRID: AB_329825), p-AKT (Thr308) (1:1000, Cat. #9275, RRID: AB_329828), AKT (1:1000, Cat. #9272, RRID: AB_329827), INSR-β subunit (1:1000, Cat. #3020S, RRID: AB_2249166), p-INSRβ (Tyr1150/1151) (1:1000, Cat. #3024, RRID: AB_331253), FOXO1 (1:1000, Cat. #2880, RRID: AB_2106495), p-FOXO1 (Thr24) (1:1000, Cat. #9464, RRID: AB_329842), all from Cell Signalling (CST), and β-tubulin (1:2000, Cat. #T0198, Sigma, RRID: AB_477556). The signals were detected by secondary HRP-conjugated antibodies (anti-mouse, Cat. #7076, RRID: AB_330924; anti-rabbit, Cat. #7074, RRID: AB_2099233; Cell Signaling) and Pierce ECL Western

Blotting Substrate (Thermo Fisher Scientific). Protein band intensities were quantified

with Image Studio Lite software (LI-COR).

Surface Protein Biotinylation Assay

Biotinylation of surface proteins was performed as previously described(29). In brief,

cells were incubated with cell-impermeable EZ-Link-NHS-SS-biotin (300 µg/ml in PBS;

Pierce) at 37°C for 2 min and washed with ice-cold 50 mM Tris-buffered saline (TBS) to

remove excess biotin. For detecting internalized proteins, cells were washed with PBS

and incubated in PBS or serum-free medium supplemented with 0, 0.2, 2, or 20 nM

insulin at 37°C to stimulate INSR internalization. After certain time periods, cells were

incubated with glutathione solution (50 mM glutathione, 75 mM NaCl, 1 mM EDTA, 1%

BSA, 75 mM NaOH) to strip the remaining surface biotin lysed in complete RIPA buffer.

Biotinylated internalized proteins in lysates were captured by NeutrAvidin beads (Pierce)

and eluted by incubating in Blue Loading Buffer (50 mM DTT, Cell Signaling, Danvers,

MA) containing 50 mM DTT at 95°C for 5 min. Internalized INSR in eluent and total INSR

in lysates were detected in Western blot analysis.

Plasmids, SNAP-tag cloning and lentivirus infection

CAV1-mRFP and CLTC-mRFP plasmids were generous gifts from Dr. Ivan Robert

Nabi, Vancouver, BC, Canada. INSR-A-TagBFP and INSR-B-TagBFP plasmids are

previously described(4).

PCR amplification of DNA fragments was performed using Q5 High-Fidelity DNA

Polymerase (Cat. #M0491, New England Biolabs, Ipswich, MA, USA) according to the

manufacturer's instructions. Prior ligation with the target vector, all PCR products were

subcloned into pJET1.2 vector using the CloneJET PCR Cloning Kit (Cat. #K1232,

Thermo Scientific, Waltham, MA, USA) according to the manufacturer's instructions.

Subsequently, the PCR products were excised from the pJET1.2 vector, gel purified and

ligated with the digested target vector using T4 DNA ligase. To generate INSR-A-EGFP

and INSR-B-EGFP plasmids, eGFP was PCR amplified using primers 5'-CGT ACG ATC

GAA TTC CCC GTG AGC AAG GGC GAG GAG CTG-3' (forward) and 5'- ACC GGT

ATC TGA TCC GGA CTT GTA CAG CTC GTC CAT GCC GA-3' (reverse) and inserted

into   pcDNA3.1(-)-INSR-A-TagRFP   and   pcDNA3.1(-)-INSR-B-TagRFP   (previously

described(4)) to replace TagRFP by EGFP using BsiWI and AgeI restriction sites.

To generate pLenti-INSR-A-SNAPf and pLenti-INSR-B-SNAPf lentiviral plasmids,

SNAP tag was firstly PCR amplified from pSNAPf Vector (Cat. #N9183S, New England

Biolabs, Ipswich, MA, USA) using primers 5'-CGT ACG ATC GAA TTC CCC ATG GAC

AAA GAC TGC GAA ATG AAG CGC ACC ACC-3' (forward) and 5'- ACC GGT ATC TGA

TCC GGA ACC CAG CCC AGG CTT GCC CAG-3' (reverse) and inserted into

pcDNA3.1(-)-INSR-A-TagRFP and pcDNA3.1(-)-INSR-B-TagRFP to replace TagRFP by

SNAP-tag using BsiWI and AgeI restriction sites. Subsequently, INSR-A-SNAP and

INSR-B-SNAP ORFs were PCR amplified from pcDNA3.1(-)-INSR-A/B-SNAPf plasmids

using primers 5'-GAA CCC ACT GCT TAC TGG CT-3' (forward) and 5'-GGT GGT TGT

ACA CTA GAA GGC ACA GTC GAG GC -3' (reverse) and inserted into

pLenti-C-Myc-DDK-IRES-GFP vector to replace the Myc tag, Flag tag, IRES, and

TurboGFP using NheI and BsrGI restriction sites. The PCR program for this long

INSR-SNAPf fragments was optimized to be as follows: initial denaturation 98°C 3min; 30

cycles of 98°C 10 s, 71°C 30 s, 72°C 4min; final extension 72°C 2min; hold 4°C.

SNAP-tag labelling and spinning disk confocal imaging of C2C12 myoblast

Cell non-permeable SNAP-Surface Alexa Fluor 488 substrate (Cat. #S9129S, New

England Biolabs) was dissolved and diluted to 5 µM concentration in serum-free medium

containing 0.5% BSA according to the manufacturer's instruction. Next, stable C2C12

myoblasts cell line expressing INSR-A-SNAP seeded on glass-bottom dishes (Cat.

#P35G-1.5-14-C, MatTek) were incubated in the substrate for 30 min on ice and washed

3 times with cold FuoroBrite DMEM (Cat. #A1896701, Gibco) serum-free medium.

C2C12 dish was then set up on the Spinning disk microscope, applied 37°C

FuoroBrite DMEM serum-free medium containing 0, 0.2 or 20 nM human insulin (Cat.

#I9278, Sigma-Aldrich), and immediately started imaging in a 37°C chamber using a

100×/1.45 oil objective and QuantEM 512SC Photometrics camera on a spinning disk

confocal system based on Zeiss Axiovert 200M microscope (LSI Imaging Core, UBC). A

focal plane in the middle of the cell was acquired to avoid capturing the top and bottom of

the cell. Time-lapse imaging was conducted at 30 s intervals between 0 – 20 min or at 5 s

intervals between 20-30 min.

Immunofluorescence staining

Stablly-transfected C2C12 cell lines were seeded on #1.5 glass coverslip (Cat. #

72230-01, Electron Microscopy Sciences) and labelled with cell non-permeable

SNAP-Surface Alexa Fluor 488 substrate, incubated with 0, 0.2 or 20 nM human insulin

for 30 min, fixed with 4% (w/v) paraformaldehyde (Sigma-Aldrich) for 20 minutes at room

temperature, and washed 3 times with PBS. After permeabilized in 0.1% (v/v) Triton

X-100 (Sigma-Aldrich) for 15 minutes, primary antibodies targeting Caveolin1 (1:400,

Rabbit mAb, Cat. #3267, Cell Signaling, RRID: AB_2275453) and CLTC heavy chain

(1:200, Goat pAb, Cat. #sc-6579, Santa Cruz, RRID:AB_2083170) were incubated

overnight at 4°C. Similarly, for AXNA2 experiment, cells labeled with SNAP-Surface

Alexa Fluor 488 substrate were incubated with insulin for 15 or 30 min, and labeled with

Annexin A2 primary antibody (1:150, Rabbit mAb, Cat. # 8235S, Cell Signaling,

RRID:AB_11129437). Coverslips were then washed in PBS, blocked in Dako Protein

Block (Cat #X0909, Agilent), and stained by secondary antibodies (1:1000, Cy3 Donkey

anti-Rabbit, Cat. #711-165-152, Jackson ImmunoResearch, RRID:AB_2307443; 1:500,

Alexa Fluor 633 Donkey anti-Goat, Cat. # A21082, Thermo Fisher Scientific,

RRID:AB_2535739) diluted in Dako Antibody Diluent (Cat. # S0809, Agilent) for 2 hours

at room temperature. After washing in PBS, coverslips were mounted with Prolong

Diamond Antifade Mountant (Cat. #P36961, Invitrogen).

STED microscopy and image analysis

Stimulated emission depletion (STED) microscopy was performed with the 100×/1.4

Oil HC PL APO CS2 STED White objective of a Leica TCS SP8 3× STED microscope

(Leica, Wetzlar, Germany) equipped with a white light laser, HyD detectors, and Leica

Application Suite X (LAS X) software (LSI Imaging Core, Life Sciences Institute,

University of British Columbia). Time-gated fluorescence detection was used to further

improve lateral resolution. INSR-SNAP labelled by Alexa Fluor 488 was excited at 488

nm and depleted using the 592-nm depletion laser. CAV1 or AXNA2 labelled by Cy3 was

excited at 555 nm and depleted using the 660 nm depletion laser. CLTC labelled by Alex

Fluor 633 was excited at 633 nm and depleted using the 775-nm depletion laser.

Sequential acquisition in the order of AF633/Cy3/AF488 was used to avoid cross-talk.

Huygens Professional software (Scientific Volume Imaging, Hilversum, the

Netherlands) was used to deconvolve STED images and calculate the Object Pearson

Coefficients for the colocalizations between INSR-SNAP, CAV1, CLTC, or ANXA2.

TIRF microscopy and single particle tracking analysis

Total internal reflection fluorescence (TIRF) microscopy was performed on a Zeiss

Axiovert 200M with a 100x Alpha-PlanFluar NA 1.45 oil objective (Zeiss, Germany) and a

TIRF laser angle modifier (Zeiss). Cells were kept at 37°C in a temperature-controlled

incubation chamber (Harvard Apparatus, Holliston, MA) and imaged in Ringer's buffer

supplemented with 20 mM glucose, as previously described(4). For insulin stimulations,

cells were in the Ringer's (20 mM glucose) buffer containing 0 (control) or 2 nM of insulin.

Images were acquired with a CoolSNAP HQ2 CCD camera (Photometrics, Tucson, AZ).

405 nm and 561 nm solid-state diode laser system was used to excite TagBFP and

mRFP tagged proteins.

The ImageJ distribution Fiji was used to process microscope images(83). The local

background of TIRF images was subtracted using the rolling ball method (20-pixel radius)

before single particle tracking.

Single particles (INSR, CAV1, or CLTC) were detected and tracked using Icy

bioimaging analysis software version 1.9.8.1(84). Particles were detected by the

UnDecimated Wavelet Transform Detector plugin that was set to detect bright spots over

a dark background, exhibit 70% sensitivity to spots <3 pixels, detect particle size between

2 and 3000 pixels. Particles were tracked using the Multiple Hypothesis Tracking method

built into the Spot Tracking plugin that was set to recognize particles that exhibited either

diffusive or directed motion(85,86). Track radius (maximum displacement of the particle

track) and lifetime of the particles were exported from the Motion Profile Processor of the

Spot Tracking plugin. Diffusion coefficients were calculated in MATLAB as described

previously but without corrections for positional errors and blurring(87). Colocalization of

tracks was determined by Track Processor Interaction Analysis (distance threshold 2

pixels) in the Spot Tracking plugin. INSR tracks that colocalized with CAV1 or CLTC

tracks for more than 4 continuous frames are determined to be colocalized tracks. The

exported time and duration of colocalization were organized in Python and plotted in R

studio. Diffusion coefficients, track radius, and the lifetime of tracks were plotted in R

studio. Specifically, SuperPlots were used to plot the mean value of the cells and

individual tracks of each cell(88). Code is available at GitHub deposit

https://github.com/hcen/INSR_tracking.

Skeletal muscle co-immunoprecipitation

Animal protocols were approved by the University of British Columbia Animal Care

Committee in accordance with national guidelines. Mice received 1.5U/kg insulin (a dose

commonly used in insulin tolerance test) or PBS through intraperitoneal injection. Skeletal

muscle (gastrocnemius) was collected after 5 or 10 min of injection. INSR was

immunoprecipitated by incubating 500 μg of protein lysate with 2 μg INSR antibodies

(Rabbit pAb, Cat. #sc-711, Santa Cruz, RRID: AB_631835) at 4°C overnight.

Subsequently, the solution was incubated with PureProteome™ Protein G Magnetic

Beads (Cat. # LSKMAGG10, Millipore, Burlington, MA, USA) and washed according to

the manufacturer's instructions. INSR protein complexes were eluted from the beads by

incubation in Blue Loading Buffer (50 mM DTT, Cell Signaling) at 95°C for 5 min and

were analyzed by standard western blot procedures. INSR, CAV1, and CLTC were

detected by the following Cell Signaling primary antibodies: INSRβ (1:1000, Mouse mAb,

Cat. #3020S, RRID: AB_2249166), CLTC (1:1000, Rabbit mAb, Cat. #4796,

RRID:AB_10828486), Caveolin1 (1:1000, Rabbit mAb, Cat. #3267, RRID: AB_2275453),

and HRP-conjugated VeriBlot secondary antibodies (1:200, anti-mouse Cat. #ab131368,

RRID:AB_2895114; anti-rabbit ab131366, RRID:AB_2892718; Abcam) that do not bind

to IgG heavy and light chains to reduce background noise.

Immunoprecipitation-Mass Spectrometry

C2C12 myoblasts were serum-starved for 6 hours and were given 0 or 2 nM insulin

for 15 min before collecting their lysates in RIPA buffer (50 mM β-glycerol phosphate, 10

mM HEPES, 1% Triton X-100, 70 mM NaCl, 2 mM EGTA, 1 mM Na3VO4, and 1 mM

NaF) supplemented with complete mini protease inhibitor cocktail (Roche, Laval, QC).

INSRα antibody (NNC0276-3000, a gift from Novo Nordisk compound sharing program)

that does not cross-react with IGF1R(34) was covalently conjugated to Dynabeads M-270

Epoxy beads (Cat. #14301, Invitrogen) to immunoprecipitate INSR according to the

manufacturer's protocol. INSR protein complexes were eluted from the beads by

incubation in Blue Loading Buffer (50 mM DTT, Cell Signaling) at 95°C for 5 min.

Eluted co-immunoprecipitation samples were run into 10% SDS PAGE gel for

clean-up procedure, fixed, stained, and destained in water. Each of the samples was

excised, reduced, alkylated, and digested according to standard in-gel digest

protocols(89). Extracted peptide samples were then cleaned up via STAGE-tip

purification, as previously described(90).

Samples were reconstituted in 2% ACN, 0.5% formic acid, and the peptides were

analyzed using a quadrupole – time of flight mass spectrometer (Impact II; Bruker

Daltonics) on-line coupled to an EasyLC 1000 HPLC (ThermoFisher Scientific) using a

Captive spray nanospray ionization source (Bruker Daltonics) including Aurora Series

Gen2 (CSI) analytical column, (25cm x 75µm 1.6µm FSC C18, with Gen2 nanoZero and

CSI fitting; Ion Opticks, Parkville, Victoria, Australia), and a µ-Precolumn, 300 µm ID x 5

mm, C18 PepMap, 5 µm, 100 A, Thermo Scientific, Waltham, Massachusetts, United

States). Samples were reconstituted in (0.1% aqueous formic acid and 2 % acetonitrile in

water) and 1ug of sample was loaded, injected in triplicates. A standard 90 min

reversed-phase separation was employed, using water:acetonitrile gradients.

LC and MS acquisition parameters were standard for peptide sequencing. Acquired

data were then searched by MaxQuant (V1.6.7.) against the Uniprot protein database for

Mouse C2C12 cell line, and LFQ intensities extracted and normalized using the MaxLFQ

algorithm, with 20ppm and 30ppm mass accuracies for precursor and product ion

masses, respectively, and a 1% false discovery rate cut-off. Differential expression was

determined by comparing the mean protein intensities of the triplicate injections between

the groups. Significance (p<0.05) was calculated via t-test using default parameters in

Perseus (v1.6.14.0).

AlphaFold-Multimer and minD

Analysis of direct interactions between INSR and interactors was carried out using

AlphaFold-Multimer and minD method(44,45). We downloaded all the protein sequences

from UniProt (IDs: INSR - P15208, RPSA - P14206, HSPB1 - P14602, ANXA2 - P07356,

HIST1H1B - P43276 and IGF1R - E9QNX9, CAV1 - P49817, CAV3 - P51637, AP2M1 -

P84091, IRS1 - P35569, MTCO2 - P00405). We then followed the previously described

protocol(45) to extract potential binding sites and do an in-silico evaluation of the

interactions between binding sites. Briefly, we first paired INSR with other proteins and

ran AlphaFold-Multimer on those pairs. We used full-length protein sequences to extract

minD scores. minD is defined as the minimum expected distance of one residue of one

protein chain from the other protein chain in the pairs. This metric is extracted from the

Distogram head of AlphaFold-Multimer and has been shown to peak when a potential

binding site is detected. After the binding sites on both proteins are identified, we then cut

the region surrounding them. This results in a handful of fragments for each pair. We then

pair

fragments for a second round of AlphaFold-Multimer runs. Finally, ipTM scores are

extracted to filter out non-binding pairs. A threshold of 0.42 is used to discriminate

non-interacting and interacting fragment pairs(45). The code for this analysis is available

at https://github.com/alirezaomidi/AFminD.

Statistics

Data were presented as mean ± SEM in addition to the individual data points. A

significance level of $p < 0.05$ was used throughout. Student's t-tests were used for

comparisons between two groups. Comparisons between more than two groups were

performed using one-way ANOVA and multiple comparison p-values adjusted using the

Tukey method. Comparisons between multiple groups and two categorical variables were

performed using Two-way ANOVA followed by Tukey post hoc test. The statistical

analysis above was performed using GraphPad Prism (Version 9.0.2). Two-sided

Kolmogorov-Smirnov Tests were used for comparing cumulative probability, which was

performed with ks.text() function in R.

Data availability

Data are contained within the article or can be shared upon request.

References

1. Knutson VP. Cellular trafficking and processing of the insulin receptor. *FASEB J.* 1991;5(8):2130-2138.
2. Kahn CR, Crettaz M. Insulin receptors and the molecular mechanism of insulin action. *Diabetes Metab Rev.* 1985;1(1-2):5-32.
3. Bevan AP, Burgess JW, Drake PG, Shaver A, Bergeron JJ, Posner BI. Selective activation of the rat hepatic endosomal insulin receptor kinase. Role for the endosome in insulin signaling. *The Journal of biological chemistry.* 1995;270(18):10784-10791.
4. Boothe T, Lim GE, Cen H, Skovsø S, Piske M, Li SN, Nabi IR, Gilon P, Johnson JD. Inter-domain tagging implicates caveolin-1 in insulin receptor trafficking and Erk signaling bias in pancreatic beta-cells. *Mol Metab.* 2016;5(5):366-378.
5. Trischitta V, Gullo D, Squatrito S, Pezzino V, Goldfine ID, Vigneri R. Insulin Internalization into Monocytes Is Decreased in Patients with Type II Diabetes Mellitus. 1986;62:522-528.
6. Trischitta V, Brunetti A, Chiavetta A, Benzi L, Papa V, Vigneri R. Defects in insulin–receptor internalization and processing in monocytes of obese subjects and obese NIDDM patients. 1989;38:1579–1584.
7. Jochen AL, Berhanu P, Olefsky JM. Insulin internalization and degradation in adipocytes from normal and type II diabetic subjects. *J Clin Endocrinol Metab.* 1986;62(2):268-274.
8. Rask-Madsen C, Kahn CR. Tissue-specific insulin signaling, metabolic syndrome, and cardiovascular disease. *Arterioscler Thromb Vasc Biol.* 2012;32(9):2052-2059.
9. Guo S. Insulin signaling, resistance, and the metabolic syndrome: insights from mouse models into disease mechanisms. *J Endocrinol.* 2014;220(2):T1-T23.
10. Jochen A, Hays J, Lee M. Kinetics of insulin internalization and processing in adipocytes: effects of insulin concentration. *J Cell Physiol.* 1989;141(3):527-534.
11. McClain DA, Olefsky JM. Evidence for two independent pathways of insulin-receptor internalization in hepatocytes and hepatoma cells. *Diabetes.* 1988;37(6):806-815.
12. Ceresa BP, Kao AW, Santeler SR, Pessin JE. Inhibition of clathrin-mediated endocytosis selectively attenuates specific insulin receptor signal transduction pathways. *Mol Cell Biol.* 1998;18(7):3862-3870.
13. Carpentier JL, Van Obberghen E, Gorden P, Orci L. Surface redistribution of 125I-insulin in cultured human lymphocytes. *J Cell Biol.* 1981;91(1):17-25.

14. Paccaud JP, Siddle K, Carpentier JL. Internalization of the human insulin receptor.
The insulin-independent pathway. *The Journal of biological chemistry*.
1992;267(18):13101-13106.
15. Jose M, Biosca JA, Trujillo R, Itarte E. Characterization of the hepatic insulin
receptor undergoing internalization through clathrin-coated vesicles and
endosomes. *FEBS Lett*. 1993;334(3):286-288.
16. Nystrom FH, Chen H, Cong LN, Li Y, Quon MJ. Caveolin-1 interacts with the
insulin receptor and can differentially modulate insulin signaling in transfected
Cos-7 cells and rat adipose cells. *Mol Endocrinol*. 1999;13(12):2013-2024.
17. Fagerholm S, Ortegren U, Karlsson M, Ruishalme I, Str√·lfors P. Rapid
insulin-dependent endocytosis of the insulin receptor by caveolae in primary
adipocytes. *PLoS ONE*. 2009;4(6):e5985.
18. Knowler WC, Fowler SE, Hamman RF, Christophi CA, Hoffman HJ, Brenneman
AT, Brown-Friday JO, Goldberg R, Venditti E, Nathan DM, Group DPPR. 10-year
follow-up of diabetes incidence and weight loss in the Diabetes Prevention
Program Outcomes Study. *Lancet*. 2009;374(9702):1677-1686.
19. Moore MC, Cherrington AD, Wasserman DH. Regulation of hepatic and
peripheral glucose disposal. *Best Pract Res Clin Endocrinol Metab*.
2003;17(3):343-364.
20. Tseng LT, Lin CL, Tzen KY, Chang SC, Chang MF. LMBD1 protein serves as a
specific adaptor for insulin receptor internalization. *The Journal of biological
chemistry*. 2013;288(45):32424-32432.
21. Gautier A, Juillerat A, Heinis C, Corrêa IR, Kindermann M, Beaufils F, Johnsson K.
An engineered protein tag for multiprotein labeling in living cells. *Chem Biol*.
2008;15(2):128-136.
22. Melmed S, Polonsky KS, Larsen PR, Kronenberg HM. Williams Textbook of
Endocrinology. 13th ed: Elsevier
Saunders.
23. Timmons JA, Atherton PJ, Larsson O, Sood S, Blokhin IO, Brogan RJ, Volmar CH,
Josse AR, Slentz C, Wahlestedt C, Phillips SM, Phillips BE, Gallagher IJ, Kraus
WE. A coding and non-coding transcriptomic perspective on the genomics of
human metabolic disease. *Nucleic Acids Res*. 2018;46(15):7772-7792.
24. McAuley KA, Williams SM, Mann JI, Walker RJ, Lewis-Barned NJ, Temple LA,
Duncan AW. Diagnosing insulin resistance in the general population. *Diabetes
Care*. 2001;24(3):460-464.
25. Hengist A, Edinburgh RM, Davies RG, Walhin JP, Buniam J, James LJ, Rogers PJ,
Gonzalez JT, Betts JA. Physiological responses to maximal eating in men. *Br J
Nutr*. 2020;124(4):407-417.
26. Polonsky KS, Given BD, Hirsch LJ, Tillil H, Shapiro ET, Beebe C, Frank BH,
Galloway JA, Van Cauter E. Abnormal patterns of insulin secretion in
non-insulin-dependent diabetes mellitus. *N Engl J Med*. 1988;318(19):1231-1239.

27.   Hamley S, Kloosterman D, Duthie T, Dalla Man C, Visentin R, Mason SA, Ang T,
Selathurai A, Kaur G, Morales-Scholz MG, Howlett KF, Kowalski GM, Shaw CS,
Bruce CR. Mechanisms of hyperinsulinaemia in apparently healthy non-obese
young adults: role of insulin secretion, clearance and action and associations with
plasma amino acids. *Diabetologia*. 2019;62(12):2310-2324.

28.   Le Marchand-Brustel Y, Jeanrenaud B, Freychet P. Insulin binding and effects in
isolated soleus muscle of lean and obese mice. *Am J Physiol*.
1978;234(4):E348-358.

29.   Cen HH, Hussein B, Botezelli JD, Wang S, Zhang JA, Noursadeghi N, Jessen N,
Rodrigues B, Timmons JA, Johnson JD. Human and mouse muscle
transcriptomic analyses identify insulin receptor mRNA downregulation in
hyperinsulinemia-associated insulin resistance. *FASEB J*. 2022;36(1):e22088.

30.   Cen HH, Johnson JD. Insulin receptor trafficking and interactions in muscle cells -
Supplemental Materials. Zenodo. 2026; https://doi.org/10.5281/zenodo.18272839

31.   Jans DA. Lateral Mobility of Polypeptide Hormone Receptors and GTP-Binding
Proteins. The Mobile Receptor Hypothesis: The Role of Membrane Receptor
Lateral Movement in Signal Transduction. Boston, MA: Springer US; 1997:83-115.

32.   Kabayama K, Sato T, Saito K, Loberto N, Prinetti A, Sonnino S, Kinjo M, Igarashi
Y, Inokuchi J. Dissociation of the insulin receptor and caveolin-1 complex by
ganglioside GM3 in the state of insulin resistance. *Proc Natl Acad Sci U S A*.
2007;104(34):13678-13683.

33.   Ikonen E, Vainio S. Lipid microdomains and insulin resistance: is there a
connection? *Sci STKE*. 2005;2005(268):pe3.

34.   Ørstrup LH, Slaaby R, Rasch MG, Rasmussen N, Lund S, Brandt J, Schluckebier
G, Wang Z, Lützen A, Pedersen T, Hvid H, Hansen BF, Blume N. Cross-species
reactive monoclonal antibodies against the extracellular domains of the insulin
receptor and IGF1 receptor. *J Immunol Methods*. 2019;465:20-26.

35.   Benyoucef S, Surinya KH, Hadaschik D, Siddle K. Characterization of insulin/IGF
hybrid receptors: contributions of the insulin receptor L2 and Fn1 domains and the
alternatively spliced exon 11 sequence to ligand binding and receptor activation.
Biochem J. 2007;403(3):603-613.

36.   Moxham CP, Duronio V, Jacobs S. Insulin-like growth factor I receptor
beta-subunit heterogeneity. Evidence for hybrid tetramers composed of
insulin-like growth factor I and insulin receptor heterodimers. *The Journal of
biological chemistry*. 1989;264(22):13238-13244.

37.   Treadway JL, Morrison BD, Goldfine ID, Pessin JE. Assembly of
insulin/insulin-like growth factor-1 hybrid receptors in vitro. *J Biol Chem*.
1989;264(36):21450-21453.

38.   Shu L, Lee L, Chang Y, Holzman LB, Edwards CA, Shelden E, Shayman JA.
Caveolar structure and protein sorting are maintained in NIH 3T3 cells

independent of glycosphingolipid depletion. *Arch Biochem Biophys*.
2000;373(1):83-90.
39. DiGiacomo V, Meruelo D. Looking into laminin receptor: critical discussion
regarding the non-integrin 37/67-kDa laminin receptor/RPSA protein. *Biol Rev*
Camb Philos Soc. 2016;91(2):288-310.
40. Zoubeidi A, Zardan A, Wiedmann RM, Locke J, Beraldi E, Fazli L, Gleave ME.
Hsp27 promotes insulin-like growth factor-I survival signaling in prostate cancer
via p90Rsk-dependent phosphorylation and inactivation of BAD. *Cancer Res*.
2010;70(6):2307-2317.
41. Muranova LK, Shatov VM, Bukach OV, Gusev NB. Cardio-Vascular Heat Shock
Protein (cvHsp, HspB7), an Unusual Representative of Small Heat Shock Protein
Family. *Biochemistry (Mosc)*. 2021;86(Suppl 1):S1-S11.
42. Yuan H, Wang T, Niu Y, Liu X, Fu L. AMP-activated protein kinase-mediated
expression of heat shock protein beta 1 enhanced insulin sensitivity in the skeletal
muscle. *FEBS Lett*. 2017;591(1):97-108.
43. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O,
Tunyasuvunakool K, Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl
SAA, Ballard AJ, Cowie A, Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T,
Petersen S, Reiman D, Clancy E, Zielinski M, Steinegger M, Pacholska M,
Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW, Kavukcuoglu K,
Kohli P, Hassabis D. Highly accurate protein structure prediction with AlphaFold.
Nature. 2021;596(7873):583-589.
44. Zhu W, Shenoy A, Kundrotas P, Elofsson A. Evaluation of AlphaFold-Multimer
prediction on multi-chain protein complexes. *Bioinformatics*. 2023;39(7).
45. Omidi A, Møller MH, Malhis N, Bui JM, Gsponer J. AlphaFold-Multimer accurately
captures interactions and dynamics of intrinsically disordered protein regions.
Proc Natl Acad Sci U S A. 2024;121(44):e2406407121.
46. Chidlow JH, Sessa WC. Caveolae, caveolins, and cavins: complex control of
cellular signalling and inflammation. *Cardiovasc Res*. 2010;86(2):219-225.
47. Capozza F, Cohen AW, Cheung MW, Sotgia F, Schubert W, Battista M, Lee H,
Frank PG, Lisanti MP. Muscle-specific interaction of caveolin isoforms: differential
complex formation between caveolins in fibroblastic vs. muscle cells. *Am J*
Physiol Cell Physiol. 2005;288(3):C677-691.
48. Schöneich C, Dremina E, Galeva N, Sharov V. Apoptosis in differentiating C2C12
muscle cells selectively targets Bcl-2-deficient myotubes. *Apoptosis*.
2014;19(1):42-57.
49. Biener Y, Feinstein R, Mayak M, Kaburagi Y, Kadowaki T, Zick Y. Annexin II is a
novel player in insulin signal transduction. Possible association between annexin
II phosphorylation and insulin receptor internalization. *The Journal of biological*
chemistry. 1996;271(46):29489-29496.

50.   Rescher U, Ludwig C, Konietzko V, Kharitonenkov A, Gerke V. Tyrosine
phosphorylation of annexin A2 regulates Rho-mediated actin rearrangement and
cell adhesion. *J Cell Sci.* 2008;121(Pt 13):2177-2185.

51.   Carpentier JL. Insulin-induced and constitutive internalization of the insulin
receptor. *Horm Res.* 1992;38(1-2):13-18.

52.   Chang M, Kwon M, Kim S, Yunn NO, Kim D, Ryu SH, Lee JB. Aptamer-based
single-molecule imaging of insulin receptors in living cells. *J Biomed Opt.*
2014;19(5):051204.

53.   Hubert M, Larsson E, Lundmark R. Keeping in touch with the membrane; protein-
and lipid-mediated confinement of caveolae to the cell surface. *Biochem Soc*
Trans. 2020;48(1):155-163.

54.   Martins AS, Ordóñez JL, Amaral AT, Prins F, Floris G, Debiec-Rychter M,
Hogendoorn PC, de Alava E. IGF1R signaling in Ewing sarcoma is shaped by
clathrin-/caveolin-dependent endocytosis. *PLoS One.* 2011;6(5):e19846.

55.   Pandini G, Frasca F, Mineo R, Sciacca L, Vigneri R, Belfiore A. Insulin/insulin-like
growth factor I hybrid receptors have different biological characteristics depending
on the insulin receptor isoform involved. *The Journal of biological chemistry.*
2002;277(42):39684-39695.

56.   Tomas A, Futter CE, Eden ER. EGF receptor trafficking: consequences for
signaling and cancer. *Trends Cell Biol.* 2014;24(1):26-34.

57.   Viola CM, Frittmann O, Jenkins HT, Shafi T, De Meyts P, Brzozowski AM.
Structural conservation of insulin/IGF signalling axis at the insulin receptors level
in Drosophila and humans. *Nature Communications.* 2023;14(1):6271.

58.   Kammanadiminti SJ, Chadee K. Suppression of NF-kappaB activation by
Entamoeba histolytica in intestinal epithelial cells is mediated by heat shock
protein 27. *The Journal of biological chemistry.* 2006;281(36):26112-26120.

59.   Ku HC, Chang HH, Liu HC, Hsiao CH, Lee MJ, Hu YJ, Hung PF, Liu CW, Kao YH.
Green tea (-)-epigallocatechin gallate inhibits insulin stimulation of 3T3-L1
preadipocyte mitogenesis via the 67-kDa laminin receptor pathway. *Am J Physiol*
Cell Physiol. 2009;297(1):C121-132.

60.   Zhao WQ, Chen GH, Chen H, Pascale A, Ravindranath L, Quon MJ, Alkon DL.
Secretion of Annexin II via activation of insulin receptor and insulin-like growth
factor receptor. *The Journal of biological chemistry.* 2003;278(6):4205-4215.

61.   Caron D, Boutchueng-Djidjou M, Tanguay RM, Faure RL. Annexin A2 is
SUMOylated on its N-terminal domain: regulation by insulin. *FEBS Lett.*
2015;589(9):985-991.

62.   Urbanska A, Sadowski L, Kalaidzidis Y, Miaczynska M. Biochemical
characterization of APPL endosomes: the role of annexin A2 in APPL membrane
recruitment. *Traffic.* 2011;12(9):1227-1241.

63.   Ryu J, Galan AK, Xin X, Dong F, Abdul-Ghani MA, Zhou L, Wang C, Li C, Holmes
BM, Sloane LB, Austad SN, Guo S, Musi N, DeFronzo RA, Deng C, White MF, Liu

F, Dong LQ. APPL1 potentiates insulin sensitivity by facilitating the binding of
IRS1/2 to the insulin receptor. *Cell Rep*. 2014;7(4):1227-1238.

64.   Lemmon MA, Schlessinger J. Cell signaling by receptor tyrosine kinases. *Cell*.
2010;141(7):1117-1134.

65.   de Graauw M, Cao L, Winkel L, van Miltenburg MH, le Dévédec SE, Klop M, Yan
K, Pont C, Rogkoti VM, Tijsma A, Chaudhuri A, Lalai R, Price L, Verbeek F, van de
Water B. Annexin A2 depletion delays EGFR endocytic trafficking via cofilin
activation and enhances EGFR signaling and metastasis formation. *Oncogene*.
2014;33(20):2610-2619.

66.   Behrends M, Engmann O. Linker histone H1.5 is an underestimated factor in
differentiation and carcinogenesis. *Environ Epigenet*. 2020;6(1):dvaa013.

67.   Brown M, Dainty S, Strudwick N, Mihai AD, Watson JN, Dendooven R, Paton AW,
Paton JC, Schröder M. Endoplasmic reticulum stress causes insulin resistance by
inhibiting delivery of newly synthesized insulin receptors to the cell surface. *Mol*
Biol Cell. 2020;31(23):2597-2629.

68.   Vicidomini G, Bianchini P, Diaspro A. STED super-resolved microscopy. *Nat*
Methods. 2018;15(3):173-182.

69.   Fish KN. Total internal reflection fluorescence (TIRF) microscopy. *Curr Protoc*
Cytom. 2009;Chapter 12:Unit12.18.

70.   Le PU, Guay G, Altschuler Y, Nabi IR. Caveolin-1 is a negative regulator of
caveolae-mediated endocytosis to the endoplasmic reticulum. *Journal of*
Biological Chemistry. 2002;277(5):3371-3379.

71.   Humphries AC, Way M. The non-canonical roles of clathrin and actin in pathogen
internalization, egress and spread. *Nat Rev Microbiol*. 2013;11(8):551-560.

72.   Rodal SK, Skretting G, Garred O, Vilhardt F, van Deurs B, Sandvig K. Extraction
of cholesterol with methyl-beta-cyclodextrin perturbs formation of clathrin-coated
endocytic vesicles. *Mol Biol Cell*. 1999;10(4):961-974.

73.   Parpal S, Karlsson M, Thorn H, Strålfors P. Cholesterol depletion disrupts
caveolae and insulin receptor signaling for metabolic control via insulin receptor
substrate-1, but not for mitogen-activated protein kinase control. *The Journal of*
biological chemistry. 2001;276(13):9670-9678.

74.   Cohen AW, Combs TP, Scherer PE, Lisanti MP. Role of caveolin and caveolae in
insulin signaling and diabetes. *Am J Physiol Endocrinol Metab*.
2003;285(6):E1151-1160.

75.   Hussain KM, Leong KL, Ng MM, Chu JJ. The essential role of clathrin-mediated
endocytosis in the infectious entry of human enterovirus 71. *The Journal of*
biological chemistry. 2011;286(1):309-321.

76.   Wu J, Park SH, Choi E. The insulin receptor endocytosis. *Prog Mol Biol Transl Sci*.
2023;194:79-107.

77.   Chen Y, Huang L, Qi X, Chen C. Insulin Receptor Trafficking: Consequences for
Insulin Sensitivity and Diabetes. *Int J Mol Sci*. 2019;20(20).

78.   Choi E, Zhang X, Xing C, Yu H. Mitotic Checkpoint Regulators Control Insulin
Signaling and Metabolic Homeostasis. *Cell*. 2016;166(3):567-581.

79.   Park J, Hall C, Hubbard B, LaMoia T, Gaspar R, Nasiri A, Li F, Zhang H, Kim J,
Haeusler RA, Accili D, Shulman GI, Yu H, Choi E. MAD2-Dependent Insulin
Receptor Endocytosis Regulates Metabolic Homeostasis. *Diabetes*.
2023;72(12):1781-1794.

80.   Choi E, Kikuchi S, Gao H, Brodzik K, Nassour I, Yopp A, Singal AG, Zhu H, Yu H.
Mitotic regulators and the SHP2-MAPK pathway promote IR endocytosis and
feedback regulation of insulin signaling. *Nat Commun*. 2019;10(1):1473.

81.   Ansarullah, Jain C, Far FF, Homberg S, Wißmiller K, von Hahn FG, Raducanu A,
Schirge S, Sterr M, Bilekova S, Siehler J, Wiener J, Oppenländer L, Morshedi A,
Bastidas-Ponce A, Collden G, Irmler M, Beckers J, Feuchtinger A, Grzybek M,
Ahlbrecht C, Feederle R, Plettenburg O, Müller TD, Meier M, Tschöp MH, Coskun
Ü, Lickert H. Inceptor counteracts insulin signalling in β-cells to control glycaemia.
Nature. 2021;590(7845):326-331.

82.   Siehler J, Bilekova S, Chapouton P, Dema A, Albanese P, Tamara S, Jain C, Sterr
M, Enos SJ, Chen C, Malhotra C, Villalba A, Schomann L, Bhattacharya S, Feng J,
Akgün Canan M, Ribaudo F, Ansarullah, Burtscher I, Ahlbrecht C, Plettenburg O,
Kurth T, Scharfmann R, Speier S, Scheltema RA, Lickert H. Inceptor binds to and
directs insulin towards lysosomal degradation in β cells. *Nat Metab*.
2024;6(12):2374-2390.

83.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T,
Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein
V, Eliceiri K, Tomancak P, Cardona A. Fiji: an open-source platform for
biological-image analysis. *Nat Methods*. 2012;9(7):676-682.

84.   de Chaumont F, Dallongeville S, Chenouard N, Hervé N, Pop S, Provoost T,
Meas-Yedid V, Pankajakshan P, Lecomte T, Le Montagner Y, Lagache T, Dufour A,
Olivo-Marin JC. Icy: an open bioimage informatics platform for extended
reproducible research. *Nat Methods*. 2012;9(7):690-696.

85.   Chenouard N, Bloch I, Olivo-Marin JC. Multiple hypothesis tracking for cluttered
biological image sequences. *IEEE Trans Pattern Anal Mach Intell*.
2013;35(11):2736-3750.

86.   Chenouard N, Smal I, de Chaumont F, Maška M, Sbalzarini IF, Gong Y, Cardinale
J, Carthel C, Coraluppi S, Winter M, Cohen AR, Godinez WJ, Rohr K, Kalaidzidis
Y, Liang L, Duncan J, Shen H, Xu Y, Magnusson KE, Jaldén J, Blau HM,
Paul-Gilloteaux P, Roudot P, Kervrann C, Waharte F, Tinevez JY, Shorte SL,
Willemse J, Celler K, van Wezel GP, Dan HW, Tsai YS, Ortiz de Solórzano C,
Olivo-Marin JC, Meijering E. Objective comparison of particle tracking methods.
Nat Methods. 2014;11(3):281-289.

87.   Abraham L, Lu HY, Falcão RC, Scurll J, Jou T, Irwin B, Tafteh R, Gold MR,
Coombs D. Limitations of Qdot labelling compared to directly-conjugated probes
for single particle tracking of B cell receptor mobility. *Sci Rep*. 2017;7(1):11379.
88. Lord SJ, Velle KB, Mullins RD, Fritz-Laylin LK. SuperPlots: Communicating
reproducibility and variability in cell biology. *J Cell Biol*. 2020;219(6).
89. Shevchenko A, Wilm M, Vorm O, Mann M. Mass spectrometric sequencing of
proteins silver-stained polyacrylamide gels. *Anal Chem*. 1996;68(5):850-858.
90. Rappsilber J, Mann M, Ishihama Y. Protocol for micro-purification, enrichment,
pre-fractionation and storage of peptides for proteomics using StageTips. *Nat*
Protoc. 2007;2(8):1896-1906.

1007

1008

Figures

Figure 1. Biochemical and live-cell imaging characterization of the effects of

insulin on INSR.

(A) Internalized to total INSR ratio was quantified using surface biotinylation in

undifferentiated C2C12 myoblasts. Cells were incubated in PBS (no insulin) or

serum-free DMEM containing 0, 0.2, 2, or 20 nM insulin for 15 min. The ratios are

normalized to the 0nM group of each gel. (p>0.05 when not specified.) **(B)** Western blot

showing SNAP-tagged INSR expressed from lentiviral vector in comparison to wild-type

INSR in C2C12 myoblasts. **(C,D)** Representative images of INSR-A-SNAP -labelled

using an Alexa Fluor 488 cell non-permeable dye in 0, 0.2 or 20 nM insulin conditions in

live undifferentiated C2C12 myoblasts from **(C)** 0 - 15 min or **(D)** after 20 min using a

spinning disk confocal microscope. Time-lapse images of INSR vesicle interactions

starting from the selected subregions (white squares) of snapshots are shown in the

insets. (n=3 cells)

1023

Figure 2. Live-cell TIRF imaging of cell-surface INSR.

(A) Experimental design of TIRF microscopy of the same cells at 3 timepoints with 2 nM

insulin or no insulin (control). **(B)** Representative TIRF image of INSR-A-EGFP puncta,

binary image of detected INSR spots, and INSR tracks. **(C)** Diffusion coefficient of

INSR-A-EGFP tracks in control or insulin group. Data is plotted as SuperPlot with larger

dots showing mean values of the cells and smaller dots showing individual INSR tracks of

the cells. **(D)** The cumulative probability shows the distribution of diffusion coefficients in

(C). **(E)** Track radius of INSR-A-EGFP tracks in control or insulin group. **(F)** The

cumulative probability shows the distribution of diffusion coefficient in **(E)**. (n=3 cells in

control group, n=9 cells in insulin group)

1034

1035

1036

1037

Figure 3. Identification of INSR interactors.

(A) Proteins that are detected in INSR immunoprecipitation mass-spectrometry (IP-MS),

and significantly different between wild-type and INSR knockout groups. Proteins in

green area were specific interactors that had little detection in knockout cells. The other 3

proteins are also significantly different but not lower or absent in INSR KO cells. (n=3) **(B)**

Protein-protein interaction network of INSR interactors and endocytosis proteins. **(C)**

Modeling the interactions between the cytoplasmic region of INSR (residues 968 - 1372)

and full-length interactor (target) proteins using AlphaFold-Multimer. pTM scores

measure the accuracy of the overall structure of the protein complex and is relatively

insensitive to localized inaccuracies. ipTM measures the accuracy of the interacting

subunits of the complex, and ipTM > 0.42 (dash line) predicts direct interaction. The

target proteins are classified as IP-MS identified novel interactors (Unknown ●), the

known or expected interactors (positive **+**), and an expected non-interactor protein as a

negative control (negative **✖**). The cytoplasmic region of IGF1R (residues 961-1369) is

used. (n = 5 predictions.) **(D)** Potential interaction sites based on per-residue scores

along the INSR cytoplasmic region (residues 968 to 1372; top panel) and interactor

(target) proteins Ap2m1 (bottom). minD (more accurate) and min-iPAE scores detect

interaction sites. pLDDT shows whether the region is structured or unstructured. Shades

show confidence errors of 99%. **(E)** ipTM scores for fragment-protein interactions. Thirty

residues around the minD peaks (potential interacting sites) in **(D)** and **Fig. S2** (30) are

analyzed. Each box relates to the interaction of the fragments of first protein vs the

full-lenght second protein. For example, title "Insr vs Mtco2" means we have cut Insr into

fragments and we have run AlphaFold-Multimer on each of those fragments vs full-length

Mtco2. Each dot is the top ipTM score for one fragment (we do 5 predictions for each

fragment vs protein). The protein residue positions of the interacting fragments (ipTM >

0.42) are labeled.

1064

Figure 4. Interaction and colocalization between INSR, CAV3, CAV1 and CLTC

under different insulin stimulations.

(A) Western blot of mice skeletal muscle lysates after insulin injection. Phospho-AKT

(Ser473) to total AKT ratio or phospho-ERK1/2 to total ERK ratio were quantified.

(n=6) **(B)** Co-immunoprecipitation of skeletal muscle INSR after PBS or insulin injection.

CLTC to INSR ratio or CAV3 to INSR ratio were quantified. (n=7-10) **(C)** Representative

STED microscopy images of C2C12 myoblasts expressing INSR-A-SNAP (surface

labeled) that were fixed after stimulation with 0 nM, 0.2 nM, or 20 nM insulin for 30 min

and stained for CAV1 and CLTC (Scale bar = 5 µm). **(D,E)** Colocalizations between

INSR-A-SNAP, CAV1, and CLTC were quantified by Object Pearson Coefficient. Data

are plotted to show differences between insulin concentrations **(D)** or between protein

pairs **(E)**. (n=7-9 images, 1-4 cells per image. *$p<0.05$, Tukey's multiple comparison after

2-ANOVA. Box represents median and 25th to 75th percentiles.)

1078

Figure 5. Colocalization and interaction between INSR and ANXA2 under different

insulin concentrations.

(A,B) Representative STED microscopy images of C2C12 myoblasts expressing

INSR-A-SNAP (surface labeled) that were fixed after stimulation with 0 nM, 0.2 nM, or 20

nM insulin for (A)15 or (B)30 min and stained for ANXA2 (Scale bar = 5 µm). **(C,D)**

Colocalization between INSR-A-SNAP and ANXA2 at (C)15 or (D)30 min of

insulin-stimulated were quantified by Object Pearson Coefficient. (n=4-13 images, 1- 3

cells per image. $^{\#}$p<0.05, 1-ANOVA of all groups. Box represents median and 25th to

75th percentiles.)

1088

Figure 6. Live-cell TIRF imaging of cell-surface INSR, CAV1 and CLTC.

(A,B)TIRF microscopy (2s/frame) showed colocalization between INSR-B-TagBFP and

(A) CAV1-mRFP or **(B)** CLTC-mRFP. **(C,D)**The associated fluorescent intensity of

representative INSR-B-TagBFP and colocalized **(C)** CAV1-mRFP or **(D)** CLTC-mRFP.

(E-G) The (E) diffusion coefficient, (F) track radius, and (G) lifetime of INSR-B-TagBFP

tracks in cells expressing CAV1-mRFP under 0 nM (n=2 cells, 360 tracks) or 2 nM insulin

(n=2 cells, 324 tracks) or expressing CLTC-mRFP under 0 nM (n=4 cells, 917 tracks) or 2

nM insulin (n=2 cells, 146 tracks). Data is plotted as SuperPlot, with bigger dots showing

mean values of the cells and smaller dots showing individual INSR tracks of the cells. The

cumulative probability shows the distribution of the diffusion coefficient, with the dots and

numbers showing the median values. (2-ANOVA: # effect of co-expression;   post-hoc

Tukey test: *p<0.05; Kolmogorov-Smirnov (KS) test: *p<0.05, **p<0.005, ***p<0.0005.)

Experiments shown were conducted in Ringer's buffer.

1102

Figure 7. Analysis of INSR tracks colocalized or not colocalized with CAV1 or

CLTC.

(A) INSR tracks that colocalized with CAV1 under 0 or 2 nM insulin (referred to as

CAV1.Y.0nM or CAV1.Y.2nM below). **(B)** INSR tracks that colocalized with CLTC under 0

or 2 nM insulin (referred to as CLTC.Y.0nM or CLTC.Y.2nM below). **(C)** Diffusion

coefficient of INSR-B-TagBFP tracks that did not colocalize with CAV1-mRFP tracks

under 0 nM insulin (CAV1.N.0nM) or 2 nM insulin (CAV1.N.2nM) or colocalized with

CAV1-mRFP tracks under 0 nM insulin (CAV1.Y.0nM) or 2 nM insulin (CAV1.Y.2nM).

The cumulative probability shows the distribution of the measurements, with the dots and

numbers showing the median values (same for other cumulative probability plots). **(D)**

Diffusion coefficient of INSR-B-TagBFP tracks that did not colocalize with CLTC-mRFP

tracks under 0 nM insulin (CLTC.N.0nM) or 2 nM insulin (CLTC.N.2nM) or colocalized

with CLTC-mRFP tracks under 0 nM insulin (CLTC.Y.0nM) or 2 nM insulin (CLTC.Y.2nM).

(E) Track radius of INSR-B-TagBFP tracks that did not colocalize with CAV1-mRFP

tracks under 0 nM insulin (CAV1.N.0nM) or 2 nM insulin (CAV1.N.2nM) or colocalized

with CAV1-mRFP tracks under 0 nM insulin (CAV1.Y.0nM) or 2 nM insulin (CAV1.Y.2nM).

(F) Track radius of INSR-B-TagBFP tracks that did not colocalize with CLTC-mRFP

tracks under 0 nM insulin (CLTC.N.0nM) or 2 nM insulin (CLTC.N.2nM) or colocalized

with CLTC-mRFP tracks under 0 nM insulin (CLTC.Y.0nM) or 2 nM insulin (CLTC.Y.2nM).

(G) Lifetime of INSR-B-TagBFP tracks that did not colocalize with CAV1-mRFP tracks

under 0 nM insulin (CAV1.N.0nM) or 2 nM insulin (CAV1.N.2nM) or colocalized with

CAV1-mRFP tracks under 0 nM insulin (CAV1.Y.0nM) or 2 nM insulin (CAV1.Y.2nM). **(H)**

Lifetime of INSR-B-TagBFP tracks that did not colocalize with CLTC-mRFP tracks under

0 nM insulin (CLTC.N.0nM) or 2 nM insulin (CLTC.N.2nM) or colocalized with

CLTC-mRFP tracks under 0 nM insulin (CLTC.Y.0nM) or 2 nM insulin (CLTC.Y.2nM).

(CAV1.N.0nM, n=2 cells, 125 tracks; CAV1.N.2nM, n=2 cells, 96 tracks; CAV1.Y.0nM,

n=2 cells, 235 tracks; CAV1.Y.2nM, n=2 cells, 228 tracks; CLTC.N.0nM, n= 4 cells,   597

tracks; CLTC.N.2nM, n= 2 cells, 91 tracks; CLTC.Y.0nM, n= 4 cells, 320 tracks;

CLTC.Y.2nM, n= 2 cells, 55 tracks.) (2-ANOVA: # effect of colocalization, $ effect of

insulin, % interaction of the 2 variables; Kolmogorov-Smirnov (KS) test: *$p < 0.05$,

***$p < 0.0005$.) Experiments shown were conducted in Ringer's buffer.

1134

Figure 8. Summary of the effects of CAV1 and CLTC on INSR tracks. Higher insulin

(20 nM) promoted INSR and CAV1 colocalization, while lower insulin (0.2 nM) promoted

INSR and CLTC colocalization. INSR tracks had lower diffusion coefficient and track

radius in CAV1-overexpressing cells than CLTC-overexpressing cells. Within the same

cells, INSR tracks colocalized with CAV1 had longer lifetimes and larger track radius than

non-colocalized tracks. INSR tracks colocalized with CLTC had longer lifetimes than

non-colocalized tracks, which is further increased by insulin.

1142

1143

1144

1145

1146

1147

1148

1149

1150

1151

1152

A

p=
p=

DMEM

Insulin 15 min

Int. INSR

INSR

α-tubulin

B

C2C12 WT   C2C12 *INSR-A*-SNAP

← pro-INSR
← INSRβ-SNAP
← INSRβ

C

	0 min	5 min	10 min	15 min
0 nM				
0.2 nM				
20 nM				

10 µm

D
0 nM insulin
After 20 min
10 µm

+0s +5s +10s +15s
+20s +25s +30s +35s

E
0.2 nM insulin
After 20 min
10 µm

+0s +5s +10s +15s +20s
+25s +30s +35s +40s +50s

F
20 nM insulin
After 20 min
10 µm

+0s +5s +10s +15s +20s +25s
+30s +35s +40s +50s +55s +60s

A

PBS 1.5U/kg insulin
5 min | 5 min | 10min

p=
p

p-AKT

AKT

p-ERK

ERK *p=*
p=

α-tubulin

B

PBS 1.5U/kg insulin
5 min | 5 min | 10min

PBS 1.5U/kg insulin
5 min | 5 min | 10min

INSR-β

CLTC   IP: InsR

CAV3

INSR-β

CLTC   Input

CAV3

p

p=   *p=*

C

INSR-A-SNAP   CAV1   CLTC   Merge

0 nM insulin

0.2 nM insulin

20 nM insulin

D

☐ 0 nM insulin
☐ 0.2 nM insulin
■ 20 nM insulin

E

☐ INSR and CAV1
☐ INSR and CLTC
☐ CAV1 and CLTC