Title: Insulin-dependent and -independent dynamics of insulin receptor trafficking in

muscle cells

Authors: Haoning Howard Cen[1,2], Aurora J. Mattison[1,2], Alireza Omidi[3,4], Jason

Rogalski[2,5], Libin Abraham[2,6], Guang Gao[2,7], Michael R. Gold[2,6], Leonard J. Foster[2,3,4],

Jörg Gsponer[3,4], James D. Johnson[1,2,8]*

Affiliation:

1 Department of Cellular and Physiological Sciences, University of British Columbia,

Vancouver, BC. Canada

2 Life Sciences Institute, University of British Columbia, Vancouver, BC. Canada

3 Department of Biochemistry and Molecular Biology, University of British Columbia,

Vancouver, BC, Canada

4 Michael Smith Laboratories, University of British Columbia, Vancouver, BC, Canada

5 The Proteomics and Metabolomics Core Facility, University of British Columbia,

Vancouver, BC, Canada

6 Department of Microbiology and Immunology, University of British Columbia, Vancouver,

BC, Canada

7 Life Sciences Institute Imaging Core Facility, University of British Columbia, Vancouver,

BC, Canada

8 Vancouver Coastal Health Research Institute, Vancouver, BC, Canada

ORCIDs: Haoning Howard Cen 0000-0001-8045-1335; Aurora J. Mattison

0000-0001-5688-5109; Alireza Omidi 0000-0003-3598-9480; Jason Rogalski

0009-0000-5736-2991; Libin Abraham 0000-0002-4725-3502; Guang Gao

0000-0001-7523-2234; Michael R. Gold 0000-0003-1222-3191; Leonard J. Foster

0000-0001-8551-4817; James D. Johnson 0000-0002-7523-9433

Running title: Muscle cell insulin receptor trafficking and interactions

Keywords: insulin, insulin receptor, caveolin, clathrin, endocytosis

*Address correspondence to: James D. Johnson, Ph.D., Professor. Diabetes Research

Group, Dept. of Cellular and Physiological Sciences & Dept. of Surgery, University of

British Columbia, 5358 Life Sciences Building, 2350 Health Sciences Mall, Vancouver,

BC, Canada, V6T 1Z3

E-mail: James.D.Johnson@ubc.ca; Fax: (604) 822-2316

Twitter: @JimJohnsonSci

Acknowledgments: Spinning disk confocal and STED microscopy were performed in

the LSI IMAGING facility of the Life Sciences Institute of the University of British

Columbia using infrastructure funded by the Canadian Foundation of Innovation and BC

Knowledge Development Fund as well as a Strategic Investment Fund (Faculty of

Medicine, University of British Columbia). We thank Dr. Ivan Robert Nabi at UBC for

guidance on imaging experiments and helpful discussions. Mass spectrometry used here

was supported by the Canada Foundation for Innovation and Genome Canada

(264PRO). We thank Yuanyuan Amanda Yang for guidance on Python and R coding, Yi

Han Xia for assistance in R coding, and Jiashuo Aaron Zhang for assistance in data

analysis. We thank Michael G. Atser for assistance in western blots. INSR antibody

NNC0276-3000 was provided by Novo Nordisk Compound Sharing. The schematics in

Figure 7 were created with BioRender.

Contributions: HHC designed the study, performed experiments, analyzed data, and

wrote the manuscript. AJM contributed to STED experiments and analyzed data. AO

performed the AlphaFold-Multimer analysis and wrote part of the manuscript. JR

performed mass spectrometry, analyzed data, and edited the manuscript. LA and MG

contributed to the MATLAB scripts and the analysis of single-particle tracking and edited

the manuscript. GG contributed to the spinning disk confocal and STED microscopy. LF

contributed to the design of the mass-spectrometry experiments and edited the

manuscript. JG supervised the AlphaFold-Multimer analysis and edited the manuscript.

JDJ designed the study, supervised the research, and edited the manuscript. JDJ is the

ultimate guarantor of this work.

Funding: This study was funded by a Canadian Institutes of Health Research Operating

Grant (P-133692) to JDJ.

Conflict of interest: The authors declare that they have no conflicts of interest with the

contents of this article.

Synopsis: We aimed to determine the INSR internalization routes and dynamics in

muscle cells. Robust basal INSR internalization in the absence of insulin was observed,

and insulin did not significantly affect INSR internalization and cell-surface dynamics.

Through mass-spectrometry, protein-protein interaction network analysis, and

AlphaFold-Multimer, we identified new interactors of INSR and links between INSR and

caveolae. Super-resolution microscopy, co-immunoprecipitation, TIRF microscopy, and

single-particle tracking suggest that muscle cells utilize both clathrin and caveolin for

INSR internalization and dynamics.

Abstract

Insulin resistance contributes to type 2 diabetes and can be driven by

hyperinsulinemia. Insulin receptor (INSR) internalization and cell-surface dynamics at

rest and during insulin exposure are incompletely understood in muscle cells. Using

surfacing labelling and live-cell imaging, we observed robust basal internalization of INSR

in C2C12 myoblasts, without an effect of added insulin. Mass-spectrometry using INSR

knockout cells as controls, identified high-confidence binding partners, including proteins

associated with internalization. We confirmed known interactors, including IGF1R, but

also identified underappreciated INSR-binding factors such as ANXA2.

AlphaFold-Multimer analysis of these INSR-binding proteins predicted potential INSR

binding sites of these proteins. Protein-protein interaction network mapping suggested

links between INSR and caveolin-mediated endocytosis. INSR interacted with both

caveolin and clathrin heavy chain (CLTC) in mouse skeletal muscle and C2C12

myoblasts. Whole cell 2D super-resolution imaging revealed that high levels of insulin (20

nM) increased INSR colocalization with CAV1 but decreased its colocalization with CLTC.

Single particle tracking confirmed the colocalization of cell-surface INSR with both

over-expressed CAV1-mRFP and CLTC-mRFP. INSR tracks that colocalized with CAV1

exhibited longer radii and lifetimes, regardless of insulin exposure, compared to

non-colocalized tracks, whereas insulin further increased the lifetime of INSR/CLTC

colocalized tracks. Overall, these data suggest that muscle cells utilize both CAV1 and

CLTC-dependent pathways for INSR dynamics and internalization.

Introduction

The insulin receptor (INSR) is one of the most studied membrane proteins due to its importance in insulin resistance and diabetes. However, many fundamental aspects of INSR biology remain unclear, including the mechanisms and kinetics of its internalization. The INSR can be internalized via endocytosis[1,2], and endosomal INSR localization contributes to the phosphorylation of specific downstream targets when compared with INSR at the cell surface[3,4]. Moreover, ligand-receptor complex dissociation in acidified endosomes permits insulin degradation, while the receptor can be sorted to recycling or degradation pathways[1,2]. INSR internalization is reduced in monocytes[5,6] and adipocytes[7] obtained from patients with insulin resistance and type 2 diabetes, suggesting the importance of this process in disease pathogenesis.

Tissue-specific insulin resistance has been implicated in the development of diabetes[8,9], but it is less clear whether there are tissue-specific INSR internalization kinetics or mechanisms. Interestingly, adipocytes internalize insulin at a lower rate in the presence of high insulin concentrations, as compared to low insulin conditions[10], while hepatocytes are the opposite[11]. The consensus view is that INSR is endocytosed via a clathrin-coated pit mechanism, based on evidence from a variety of cell types, including human and rat hepatocytes, fibroblasts, and lymphocytes[11-15]. However, other studies have shown that caveolin-rich membrane lipid rafts, caveolae, are critical for INSR internalization in other cell types, including adipocytes[16,17]. In pancreatic β-cells, we showed that caveolin-1 (CAV1) mediates INSR internalization, bypassing clathrin positive compartments[4]. However, INSR internalization kinetics and routes are less well characterized in muscle cells. In muscle cells, one study has shown that clathrin-mediated endocytosis is involved in INSR internalization[18], but the role of the caveolin-mediated pathway has not been investigated.

In the present study, we employed inter-domain tagged INSR with fluorescent

proteins[4] and SNAP-tag[19], surface biotinylation assays, as well as super-resolution and

TIRF imaging, to explore the dynamics of INSR internalization and the effects of ambient

insulin. We also used proteomics and AlphaFold-Multimer to identify INSR interacting

proteins and defined the relationship between INSR and caveolin or clathrin. Together,

our data show that INSR exhibited a high degree of insulin-independent internalization in

C2C12 myoblasts.

Results

Effects of insulin on INSR internalization

First, we characterized INSR internalization in C2C12 myoblasts under basal

conditions and in the context of added insulin. A high rate of basal INSR internalization

was observed in high-glucose serum-free DMEM medium when compared with PBS.

INSR internalization over 15 min was not significantly stimulated by the addition of insulin

to the DMEM medium (Fig. 1A). Therefore, INSR has a high rate of constitutive

internalization in this cell type, which may be enhanced by the nutrients in the medium,

but that is largely independent of acute exposure to insulin across a range of

physiological doses. These observations in myoblasts are consistent with our previous

measurements in differentiated C2C12 myotubes[20].

Qualitative analysis of INSR kinetics in the context of different insulin doses

We also visualized INSR internalization at single-cell level using

inter-domain-tagged INSR, which we have previously found more faithfully recapitulates

the sub-cellular localization of INSR compared to terminal fusion proteins[4]. The

inter-domain-tagged INSR was also shown to have functional AKT and ERK signaling in

C2C12 (Fig. S1) and other cell lines[4]. C2C12 myoblast cells with stable overexpression of

SNAP-tagged INSR isoform A (INSR-A-SNAP; Fig. 1B) were labelled using a cell

non-permeable SNAP-Surface Alexa Flour 488 dye (Fig. 1C). Although a few INSR

vesicles seemed to be internalized during the small gap between labelling and imaging,

and were present at time zero, substantially more internalized INSR vesicles were

observed at later timepoints (Fig. 1C). Live-cell time-lapse imaging of surface-labelled

INSR-A-SNAP at the middle plane of cells by spinning disk confocal microscopy revealed

INSR internalization in the context of 0, 0.2, and 20 nM insulin over 15 min (Fig. 1C,

Video 1). Consistent with the surface biotinylation assay results in Figure 1A, we did not

observe a clear difference in INSR internalization in the presence of insulin in these cells.

Internalized INSR vesicles exerted dynamic movement and interaction, which was more

evident between 20-30 min (Fig. 1D-F, Video 2). INSR-containing vesicles exhibited

many apparent fusion and fission events regardless of insulin concentrations. Some

smaller vesicles merged into a larger vesicle that remained in the field undergoing limited

and apparently random motion (one example is shown in Fig. 1F), suggestive of a fusion

event. A caveat associated with 2D confocal imaging is limited Z resolution, raising the

possibility that the observed INSR vesicles are not all in the same optical plane. Although

our experiments using surface labelling of SNAP-tagged INSR permitted the visualization

of the dynamic behaviour of internalized INSR vesicles.

Quantitative analysis of INSR kinetics in the context of insulin

Lateral movement of tyrosine kinase receptors has been linked to their functional

state. Ligand binding of some monomeric receptors, such as EGFR, promotes their

dimerization and aggregation and decreases their lateral mobility[21]. Although INSRs are

thought to already exist as dimers in plasma membrane, aggregation of INSR by

antibodies can stimulate INSR signaling[21], and INSR may need to localize to membrane

lipid rafts for normal signaling[22,23]. Therefore, we sought to quantitatively assess whether

insulin affects the lateral movement of INSR in or near the plasma membrane. To do this,

we used time-lapse TIRF microscopy of C2C12 myoblast transiently expressing

inter-domain-tagged INSR-A-EGFP. We imaged the same cells at 0, 6, or 11 min in the

presence of 2 nM insulin, as well as a control group without insulin to control for other

variables such as photobleaching (Fig. 2A). Single-particle tracking of INSR-A-EGFP

(Fig. 2B) was used to calculate their diffusion coefficient (Fig. 2C,D) and track radius (Fig.

2E,F). Insulin did not have a big impact on the track radius or diffusion coefficient, despite

some very subtle differences in the cumulative probability curves with very close median

values (Fig. 2D,F). Together with the qualitative data reported above, these observations

suggest that INSR mobility is largely unaffected by acute insulin stimulation in this cell

model.

Novel INSR interactors identified by immunoprecipitation and mass spectrometry

To explore the molecular mechanisms associated with INSR internalization, we

conducted co-immunoprecipitation followed by mass-spectrometry to identify INSR

interactors. We used INSR knockout C2C12 myoblasts as a negative control and an

INSR antibody that does not cross-react with Insulin-like growth factor 1 receptor

(IGF1R)[24] to identify INSR interactors with higher specificity than any previously reported

study. After filtering the detected proteins against the INSR knockout cell control lysates,

we identified INSR and five high-confidence INSR-binding proteins (Fig. 3A). The

immunoprecipitated abundance of these detected interactors was not significantly

changed by the addition of 2 nM insulin (Fig. 3A). INSR is well established to form

heterodimers with IGF1R[25,26], and its immunoprecipitation represented a type of internal

positive control for this study. The other detected proteins are not well-known INSR

interactors. To discover whether any of the identified INSR-binding proteins might be

connected with endocytosis machinery, protein-protein interaction (PPI) network was

constructed between these co-immunoprecipated proteins and all the proteins under the

"Endocytosis" pathway in Kyoto Encyclopedia of Genes and Genomes (KEGG). We

manually extracted a subnetwork that connected all our identified proteins, except for

HIST1H1B which did not connect with any proteins in this network (Fig. 3B). In the

resultant PPI network, annexin A2 (ANXA2) is connected with CAV1 and CAV3 (Fig. 3B),

and it has been reported to localize in caveolae[27]. Ribosomal protein SA (RPSA) is a

laminin receptor reported to localize in plasma membrane lipid rafts[28]. Heat shock protein

B1 (HSPB1; a.k.a. HSP27) is exclusively found in insulin-responsive tissues and has

been implicated in insulin sensitivity and INSR/IGF1R signalling[29-31]. The PPI network

and cellular localization of these interactors pointed to the possibility that INSR may also

associate with caveolins and/or related proteins in myoblasts.

Modelling INSR interactions *in silico* using AlphaFold-Multimer

AlphaFold-Multimer has been used to accurately model protein complexes[32,33].

Recent work showed that AlphaFold-Multimer more accurately predicts interactions of

protein fragments than lengthy full proteins and established a new pipeline to identify

potential interacting fragments/sites[34]. We applied this new pipeline to infer whether and

how the putative INSR interactors identified with mass spectroscopy directly bind to

INSR. We included a mitochondrial protein MTCO2 (aka COX2), expected to have no

interaction with INSR, as a negative control in these *in silico* experiments. Several known

or expected interactor of INSR were also tested, including IRS1, AP2M1, CAV1 and

CAV3. First, we ran AlphaFold-Multimer on the cytoplasmic region of INSR and each of

the full-length interactors to predict a protein complex structure. We made five separate

predictions for each pair of the proteins, shown as their predicted template modeling

(pTM) and interface-pTM (ipTM) scores (Fig. 3C). Positive interaction with IRS1 and

borderline interactions (ipTM ≥ 0.42 as suggested in [34]) with AP2M1 and IGF1R are

predicted (Fig. 3C). CAV1, CAV3 and all other interactors did not have significant scores.

The predictions with full-length proteins allowed us to extract per-residue scores of

minimum expected distance (minD), minimum inter-predicted aligned error (min-iPAE),

and the predicted local distance difference test (pLDDT) for INSR and all interactors (Fig.

3D and Fig. S2). minD and min-iPAE scores pinpoint potential interaction sites, with minD

being the most accurate predictor of interaction sites between two proteins[34]. pLDDT is a

measure of local confidence and can indicate whether the region of interest is structured

or unstructured. Using the INSR and AP2M1 interaction as an example, we observed that

AlphaFold-Multimer predicts a sharp peak in all scores at residue 1350 of INSR,

suggesting that the region around residue 1350 is in contact with AP2M1 (Fig. 3D). Yet

this is not reflected in their full-length protein predictions (highest ipTM = 0.5, Fig. 3C),

demonstrating that AlphaFold-Multimer may detect the interaction site between two

proteins but is not able to predict the interaction when full-length proteins are used. To

overcome this limitation, proteins can be fragmented into shorter segments, which are

then submitted to AlphaFold-Multimer for prediction. This fragmentation can be guided by

minD peaks that identify potential interaction sites. Thus, we cut both the INSR and the

interactor proteins into fragments of 30 residues surrounding minD peaks[34]. Doing so, we

saw a large increase in ipTM scores for IGF1R, IRS1, AP2M1, CAV3, but not CAV1 (Fig.

3E). Interestingly, although we did not find positive predictions for CAV1 direct interaction,

a peak in all scores (minD, min-iPAE and pLDDT) of CAV1 was detected around a

phosphotyrosine residue 25, which might be an interacting site (Fig. S2). Predictions via

fragmentation also suggest interactions between INSR and ANXA2, H1-5, HSBP1, and

RPSA (Fig. 3E). The ipTM scores of the negative control MTCO2 remained low,

supporting the confidence in the positive predictions (Fig. 3E). Overall, the

AlphaFold-Multimer predictions support the direct interactions between INSR and the

novel interactors and identify potential interacting sites.

Insulin-dependent interaction between INSR and CAV3 or CLTC *in vivo*

INSR has not been shown to utilize both caveolin and clathrin-mediated endocytosis

in the same tissue; therefore, we sought to confirm that INSR could interact with both

caveolin and clathrin in skeletal muscle *in vivo*. Skeletal muscle was collected 5 or 10 min

after intraperitoneal injection of insulin (1.5 U/kg) or PBS, timepoints sufficient to measure

acute insulin signalling via both AKT and ERK signalling pathways (Fig. 4A). Acute insulin

administration did not change the amount of INSR that could be immunoprecipitated from

skeletal muscle, nor did it alter the association between INSR and CLTC (Fig. 4B).

However, 10 min after insulin injection, there was a significantly increased interaction

between INSR and caveolin, specifically, the mature skeletal muscle-specific isoform

CAV3[35,36] (Fig. 4B). These data show that acute insulin signalling results in an increased

association between INSR and CAV3 *in vivo*, but does not define the cellular context for

this interaction.

Super-resolution imaging of INSR, CAV1, CLTC, and AXNA2 interactions in the

presence of insulin

We next used super-resolution STED imaging to examine colocalization between

INSR and clathrin (CLTC) or caveolin (C2C12 myoblasts only express CAV1 before

differentiation[37]). C2C12 myoblast cells with stable overexpression of INSR-A-SNAP

were labelled using a cell-nonpermeable SNAP substrate to image only

surface-associated and newly internalized INSR. Cells were incubated with 0, 0.2 or 20

nM insulin for 30 min, and then fixed. Endogenous CLTC and CAV1 were then labelled

using immunofluorescent antibody staining (Fig. 4C). The colocalization between

INSR-SNAP, CAV1, and CLTC was quantified by Object Pearson Coefficient (Fig. 4D,E).

The degree of colocalization between INSR and CAV1 was similar in the presence of 0 or

0.2 nM insulin but was higher under 20 nM insulin (Fig. 4D). However, INSR and CLTC

colocalized more at 0.2 nM insulin, but less at 20 nM insulin, compared to 0 nM insulin

(Fig. 4D). CAV1 and CLTC do not have known biological interactions; therefore, their low

level of colocalization served as an internal negative control and, indeed, it did not

change under different insulin levels (Fig. 4D). Using the same data, the relative

colocalizations between INSR and CAV1 or CLTC were also considered at different

insulin concentrations, further illustrating how high insulin concentrations drive INSR to

colocalize with CAV1 (Fig. 4E). At 0.2 nM, insulin promoted INSR interaction with CLTC

whereas 20 nM insulin redirected the INSR from CLTC to CAV1. Together, these data

revealed the insulin-dependent interaction of INSR with both CAV1 and CLTC *in vivo* and

in vitro.

Since our novel interactor ANXA2 is a target of insulin signaling[38,39], we also

examined its insulin-dependent colocalization with INSR using the same SNAP labelling

method and STED microscopy. Surface labeled INSR was allowed to internalize for 15 or

30 min in the presence of 0, 0.2, 2 or 20 nM insulin. Endogenous ANXA2 proteins were

then labelled using antibody immunofluorescence. Colocalized INSR and ANXA2 puncta

were observed in all conditions (Fig. 5A,C). The degree of colocalization quantified by

Object Pearson Coefficient remained to be moderate in most conditions (lower than the

INSR/CAV1 or INSR/CLTC interactions above) (Fig. 5B,D), except that 0.2 nM insulin

seemed to decrease their colocalization at 30 min (Fig. 5D). This observation suggests

that INSR and ANXA2 interactions are moderate and may be negatively regulated by

insulin signalling.

Effects of CAV1 and CLTC on INSR mobility

To better understand the interplay between INSR, CAV1, and CLTC, we

co-expressed INSR-B-TagBFP with CAV1-mRFP or CLTC-mRFP in C2C12 myoblasts

and imaged them by TIRF time-lapse microscopy. Colocalizations between INSR and

CAV1 or CLTC were observed (Fig. 6A,B). Some colocalized puncta moved or

disappeared together with closely associated intensity, referred to here as colocalized

tracks (examples in Fig. 6C,D). Tracks were considered as colocalized when the distance

between the centers of particle signals was less than 2 pixels for more than 4 continuous

frames. These observations suggested that INSR interacted with both CAV1 and CLTC

on the cell surface.

To define the effects of CAV1 or CLTC over-expression on INSR movement, we first

conducted single particle tracking on all INSR puncta. Similar to our observations above

(Fig. 2), insulin did not alter the diffusion coefficient or track radius of INSR. Cells with

CAV1 co-expression had lower diffusion coefficients and track radius of INSR compared

to cells with CLTC co-expression, regardless of ambient insulin exposure (differences in

medians are at least ~2 fold for diffusion coefficients and ~1.5 fold for track radius; Fig.

6E,F). CLTC-expressing cells under 0 nM insulin had higher diffusion coefficients than

those under 2 nM insulin (~ 1.6 fold difference in median; Fig. 6E), as well as shorter

INSR lifetimes than all other groups (>1.5 fold difference in median; Fig. 6G), which might

suggest a more rapid internalization. Together, these data suggest that CAV1 may

restrain INSR movement, relative to CLTC.

Finally, we performed an analysis within the same cells, comparing INSR tracks that

were colocalized or not colocalized with either CAV1 or CLTC tracks, to further dissect

the effects of CAV1 or CLTC. INSR appeared to spend more time associated with CAV1,

and for longer periods, compared with CLTC (Fig. 7A,B). There were no clear effects of

time on the association between INSR and CAV1 or CLTC across different moments of

their lifetimes together (Fig. 7A,B). We compared the colocalized and the non-colocalized

INSR tracks, for both CAV1 and CLTC, and found no significant differences in mean

diffusion coefficients (Fig. 7C,D), except that in CLTC-expressing cells without insulin,

non-colocalized INSR had higher diffusion coefficients than CLTC-colocalized INSR,

based on the cumulative probability curve (Fig.7D). The track radius of INSR colocalized

with CAV1 was larger than non-colocalized tracks regardless of insulin (Fig. 7E), but

colocalization with CLTC had significant effects (Fig. 7F). Strikingly, the lifetimes of INSR

tracks colocalized with CAV1 were much longer than non-colocalized tracks regardless of

insulin (3.8~4 fold differences in medians; Fig. 7G). INSR tracks colocalized with CLTC

also had longer lifetimes than non-colocalized tracks (~2.5 fold difference in median), and

insulin also further increased the lifetime of colocalized tracks (Fig. 7H). Overall, our

live-cell imaging of INSR confirmed its distinct interactions with either CAV1 or CLTC

around the cell surface.

Discussion

The goal of this study was to explore the INSR trafficking dynamics and interactions

in a muscle cell model. We demonstrated that INSR was constitutively internalized both in

the absence and presence of insulin. We used a high-stringency proteomics approach

and AlphaFold-Multimer to identify specific INSR binding proteins and PPI network

analysis to predict a role for caveolin in INSR internalization. Biochemical experiments,

super-resolution imaging, and TIRF imaging suggested both caveolin and clathrin are

involved in a different manner in INSR internalization and cell-surface trafficking (Fig. 8).

The constitutive internalization of INSR has been described previously in

lymphocytes and shown not to require ligand binding and kinase activity, although

insulin-stimulated autophosphorylation of INSR could accelerate internalization[14]. In

differentiated C2C12 myotubes, INSR phosphorylation levels were constitutively high,

with little additional activation by insulin across the physiological range[20], consistent with

the lack of stimulation of INSR internalization by acute insulin treatment in this study.

Surface labeling by SNAP-tag enabled clear visualization of internalized INSR-containing

vesicles, permitting the study of INSR endosomal sorting and traffic. The lateral

movement of INSR in or near the plasma membrane was also not dramatically affected

by insulin, consistent with a previous study that reported unchanged INSR diffusion

coefficient with or without insulin in HEK293T, HepG2, and human IR-overexpressing

Rat-1 cells[40]. CAV1 may have a role in decreasing the mobility of INSR. Indeed, caveolae

have restricted lateral movement in the plasma membrane due to the connection to actin

cytoskeleton through Filamin-A[41], so the CAV1-dependent decrease in INSR mobility

might reflect the association of INSR with caveolae. Consistently, the accumulation of

ganglioside GM3 in the TNFα-induced insulin resistance state was associated with the

elimination of INSR from the caveolae and increased INSR mobility on the cell surface[22].

Additional studies will be required to fully define the processes that control INSR

movement within the plasma membrane.

The possibility of more than one internalization pathway for INSR in some cell types

has been understudied. The dual clathrin and caveolin endocytosis routes have been

demonstrated for IGFR in Ewing sarcoma cells[42], which has a similar structure and

function as INSR and can form heterodimers with INSR[43]. The current study supports the

concept that INSR may also utilize both clathrin- and caveolin-mediated endocytosis in

the same cell. In hepatocytes, it has also been suggested that INSRs are internalized in

two functionally separate pathways based on the much more rapid recycling of receptors

internalized by high concentrations of insulin (17.2 nM) versus low insulin (0.17 nM), and

treatments known to inhibit coated pit formation blocked internalization of the

insulin-INSR complex at low but not high insulin concentrations[11]. However, most studies

on INSR internalization looked at one route at a time. The mechanisms of coordinated

multiple internalization routes, implied by the insulin-dependent clathrin and caveolin

interactions we observed in muscle cells, might direct INSR to different destinations (e.g.

recycle or degradation)[44] and should be comprehensively characterized in more cell

types to assess potential roles in tissue-specific insulin resistance.

We used stringent proteomics and INSR CRISPR knockout control cells to identify

known, novel, and under-appreciated interactors of INSR. As expected, we identified

IGF1R, an established hetero-dimerization partner of INSR. We also demonstrated the

specific binding of HSBP1 (HSP27), which is involved in tyrosine kinase signalling

pathways downstream of p38 MAPK, ERK, and MAPKAPK5[45] and has been previously

linked to insulin sensitivity[29-31]. We demonstrated an interaction between INSR and RPSA,

which has been previously co-immunoprecipitated with INSR and insulin receptor

substrates in 3T3L1 preadipocytes[46]. In other cell types, RPSA drives ERK signalling and

cytoskeletal organization, with nuclear functions affecting cell growth, survival, migration,

and protein synthesis[28]. ANXA2 is not a well-known interactor of INSR. One study

reported ANXA2 interaction with INSR and IGF1R[47]. Other studies found that INSR

signalling can tyrosine-phosphorylate[38,39] and SUMOylate[48] AXNA2, which mediates

insulin-stimulated actin rearrangement. Interestingly, internalized INSR seemed to be

necessary for AXNA2 phosphorylation because mutant INSR that was incapable of

internalization but normal for tyrosine kinase signalling failed to phosphorylate AXNA2[38].

AXNA2 also interacts with endosomal and adaptor prortein APPL2, which interacts with

INSR[49,50]. On the other hand, the PPI network showed that ANXA2 has a strong direct

connection with the epidermal growth factor receptor (EGFR) which belongs to the same

receptor tyrosine kinase superfamily as INSR[51]. ANXA2 knockdown halted EGFR

endocytosis beyond the early endosome, which was accompanied by enhanced EGFR

and AKT phosphorylation[52]. These data support the possibility that ANXA2 may have a

role in INSR trafficking and signalling that is worth investigating. Interestingly, evidence

compiled on the Common Metabolic Disease Knowledge Portal shows that ANXA2

genetic variation is strongly associated with fasting C peptide levels, pointing to a causal

role, direct or indirect, in hyperinsulinemia/insulin resistance. Both *Anxa2* and *Rpsa*

mRNAs were upregulated by prolonged insulin in C2C12 myotubes[8]. HIST1H1B belongs

to the family of linker histones that is important in chromatin organization[53], but the

meaning of this INSR interaction requires further investigation. This proteomic analysis,

together with AlphaFold-Multimer modelling, provides clear evidence for a limited number

of proteins that are likely to be direct interactors with INSR in myoblast cells, and

demonstrates that the majority of these interactors were not significantly altered by insulin

exposure.

There are caveats and unexplained observations in the current study. Although our

inter-domain-tagged INSR has functional signalling and internalization, it tends to have

higher pro-INSR bands (Fig. 1B)[4], which is indicative of a potential processing issue.

Nonetheless, higher pro-INSR could also occur in wildtype C2C12 myoblasts (Fig. S1A

and [54]), possibly due to stress[54], which may be a co-culprit in the processing issue of the

overexpressed tagged INSR. STED superresolution microscopy usually has a resolution

of 40-50 nm and might be able to reach 20 nm resolution in the best scenario, and

primary and secondary antibody labelling might also add ~30nm distance and reduce the

accuracy[55]. TIRF microscopy has diffraction-limited lateral resolution (~200 – 300 nm)

with the thickness of the excitation depth around 100 nm[56]. Therefore, the colocalization

analyses may not confirm the interactions of the proteins. Nonetheless, the

insulin-dependent colocalization observed in STED microscopy and the track

colocalization over multiple frames observed in TIRF microscopy are evidence that the

proteins had interactions instead of random colocalization. This is further supported by

the AlphaFold-Multimer predictions. The CAV1- or CLTC-colocalized INSR tracks had

longer lifetimes than non-colocalized tracks, which conflicted with the assumption that

CAV1 or CLTC internalized INSR, but instead, they seemed to retain INSR on the

membrane. There are multiple possible explanations; for example, overexpressed CAV1

might stabilize caveolae[57], or CLTC could form stable flat lattices, termed clathrin

plaques[58], which might have novel involvement in INSR internalization or signalling. As

we did not perform loss-of-function studies to confirm the roles of caveolin and clathrin in

INSR endocytosis, further research is needed to elucidate the roles of specific INSR

internalization routes on insulin actions. For example, inhibition of clathrin-mediated

endocytosis of INSR by knocking down LMBD1 in myoblasts increased activation of the

INSR and AKT[18]. Our lab discovered that reducing CAV1-dependent INSR internalization

blocked the activation of ERK, but not AKT, signalling in pancreatic beta-cells[4].

Cholesterol extraction by methyl-β-cyclodextrin, which is often used as a

caveolae-specific inhibitor but also affects clathrin-coated pit formation[59], inhibits the

phosphorylation of IRS1, SHC, and AKT, but not ERK or INSR phosphorylation, in

adipocytes[60,61]. In contrast, inhibition of both clathrin- and caveolin-mediated endocytosis

by the dynamin K44A dominant-negative mutant[62] reduced the phosphorylation of SHC

and ERK phosphorylation, but not IRS1 or AKT phosphorylation, in rat hepatoma H4IIE

cells[12]. More recent advances in INSR endocytosis mechanisms and consequences are

reviewed in [63,64]. For example, INSR endocytosis is found to be regulated by the spindle

checkpoint proteins (MAD2, CDC20, BUBR1 and p31[comet])[65,66] and SHP1-MAPK

pathway[67] in the liver and by the inceptor protein in β cells[68,69]; decreased

insulin-stimulated endocytosis or increased insulin-independent endocytosis by

manipulating these proteins can promote or impair insulin signaling, respectively.

Therefore, INSR internalization enables selective activation of downstream signalling

pathways, and in a tissue-specific manner.

In summary, this study provides important information on the internalization of INSR

in myoblast cells and defines the stringent INSR interactome. We also provide evidence

using a variety of methods that INSR interacts with both clathrin and caveolins in this cell

model. The current study can serve as a stepping stone for more detailed research on

how multiple internalization routes can provide cell-type customized regulations on INSR

internalization.

Materials and Methods

Cell culture

The C2C12 mouse myoblast (ATCC cell line provided by Dr. Brain Rodrigues,

University of British Columbia, Vancouver, Canada) was maintained in Dulbecco's

modified Eagle's medium (DMEM, Gibco) supplemented with 10% (v/v) fetal bovine

serum (FBS, Gibco), and 1% penicillin-streptomycin (100 µg/ml; Gibco). All downstream

experiments were conducted after 6-hour serum starvation in serum-free medium

(DMEM supplemented with 1% penicillin-streptomycin), except as otherwise indicated.

Western blot analyses

C2C12 myotubes or mice skeletal muscle (gastrocnemius) tissues were sonicated

in RIPA buffer (50 mM β-glycerol phosphate, 10 mM HEPES, 1% Triton X-100, 70 mM

NaCl, 2 mM EGTA, 1 mM Na_3VO_4, and 1 mM NaF) supplemented with complete mini

protease inhibitor cocktail (Roche, Laval, QC). Lysates were incubated in Blue Loading

Buffer containing 50 mM DTT (Cell Signaling, Danvers, MA) at 95°C for 5 min and

resolved by SDS-PAGE. Proteins were then transferred to PVDF membranes (BioRad,

CA) and probed with antibodies against p-ERK1/2 (Thr202/Tyr204) (1:1000, Cat. #4370),

ERK1/2 (1:1000, Cat. #4695), p-AKT (Ser473) (1:1000, Cat. #9271), p-AKT (Thr308)

(1:1000, Cat. #9275), AKT (1:1000, Cat. #9272), INSR-β subunit (1:1000, Cat. #3020S),

p-INSRβ (Tyr1150/1151) (1:1000, Cat. #3024), FOXO1 (1:1000, Cat. #2880), p-FOXO1

(Thr24) (1:1000, Cat. #9464), all from Cell Signalling (CST), and β-tubulin (1:2000, Cat.

#T0198, Sigma). The signals were detected by secondary HRP-conjugated antibodies

(anti-mouse Cat. #7076, anti-rabbit Cat. #7074; Cell Signaling) and Pierce ECL

Western Blotting Substrate (Thermo Fisher Scientific). Protein band intensities were

quantified with Image Studio Lite software (LI-COR).

Surface Protein Biotinylation Assay

Biotinylation of surface proteins was performed as previously described[20]. In brief,

cells were incubated with cell-impermeable EZ-Link-NHS-SS-biotin (300 µg/ml in PBS;

Pierce) at 37°C for 2 min and washed with ice-cold 50 mM Tris-buffered saline (TBS) to

remove excess biotin. For detecting internalized proteins, cells were washed with PBS

and incubated in PBS or serum-free medium supplemented with 0, 0.2, 2, or 20 nM

insulin at 37°C to stimulate INSR internalization. After certain time periods, cells were

incubated with glutathione solution (50 mM glutathione, 75 mM NaCl, 1 mM EDTA, 1%

BSA, 75 mM NaOH) to strip the remaining surface biotin lysed in complete RIPA buffer.

Biotinylated internalized proteins in lysates were captured by NeutrAvidin beads (Pierce)

and eluted by incubating in Blue Loading Buffer (50 mM DTT, Cell Signaling, Danvers,

MA) containing 50 mM DTT at 95°C for 5 min. Internalized INSR in eluent and total INSR

in lysates were detected in Western blot analysis.

Plasmids, SNAP-tag cloning and lentivirus infection

CAV1-mRFP and CLTC-mRFP plasmids were generous gifts from Dr. Ivan Robert

Nabi, Vancouver, BC, Canada. INSR-A-TagBFP and INSR-B-TagBFP plasmids are

previously described[4].

PCR amplification of DNA fragments was performed using Q5 High-Fidelity DNA

Polymerase (Cat. #M0491, New England Biolabs, Ipswich, MA, USA) according to the

manufacturer's instructions. Prior ligation with the target vector, all PCR products were

subcloned into pJET1.2 vector using the CloneJET PCR Cloning Kit (Cat. #K1232,

Thermo Scientific, Waltham, MA, USA) according to the manufacturer's instructions.

Subsequently, the PCR products were excised from the pJET1.2 vector, gel purified and

ligated with the digested target vector using T4 DNA ligase. To generate INSR-A-EGFP

and INSR-B-EGFP plasmids, eGFP was PCR amplified using primers 5'-CGT ACG ATC

GAA TTC CCC GTG AGC AAG GGC GAG GAG CTG-3' (forward) and 5'- ACC GGT

ATC TGA TCC GGA CTT GTA CAG CTC GTC CAT GCC GA-3' (reverse) and inserted

into pcDNA3.1(-)-INSR-A-TagRFP and pcDNA3.1(-)-INSR-B-TagRFP (previously

described[4]) to replace TagRFP by EGFP using BsiWI and AgeI restriction sites.

To generate pLenti-INSR-A-SNAPf and pLenti-INSR-B-SNAPf lentiviral plasmids,

SNAP tag was firstly PCR amplified from pSNAPf Vector (Cat. #N9183S, New England

Biolabs, Ipswich, MA, USA) using primers 5'-CGT ACG ATC GAA TTC CCC ATG GAC

AAA GAC TGC GAA ATG AAG CGC ACC ACC-3' (forward) and 5'- ACC GGT ATC TGA

TCC GGA ACC CAG CCC AGG CTT GCC CAG-3' (reverse) and inserted into

pcDNA3.1(-)-INSR-A-TagRFP and pcDNA3.1(-)-INSR-B-TagRFP to replace TagRFP by

SNAP-tag using BsiWI and AgeI restriction sites. Subsequently, INSR-A-SNAP and

INSR-B-SNAP ORFs were PCR amplified from pcDNA3.1(-)-INSR-A/B-SNAPf plasmids

using primers 5'-GAA CCC ACT GCT TAC TGG CT-3' (forward) and 5'-GGT GGT TGT

ACA CTA GAA GGC ACA GTC GAG GC -3' (reverse) and inserted into

pLenti-C-Myc-DDK-IRES-GFP vector to replace the Myc tag, Flag tag, IRES, and

TurboGFP using NheI and BsrGI restriction sites. The PCR program for this long

INSR-SNAPf fragments was optimized to be as follows: initial denaturation 98°C 3min; 30

cycles of 98°C 10 s, 71°C 30 s, 72°C 4min; final extension 72°C 2min; hold 4°C.

SNAP-tag labelling and spinning disk confocal imaging of C2C12 myoblast

Cell non-permeable SNAP-Surface Alexa Fluor 488 substrate (Cat. #S9129S, New

England Biolabs) was dissolved and diluted to 5 μM concentration in serum-free medium

containing 0.5% BSA according to the manufacturer's instruction. Next, stable C2C12

myoblasts cell line expressing INSR-A-SNAP seeded on glass-bottom dishes (Cat.

#P35G-1.5-14-C, MatTek) were incubated in the substrate for 30 min on ice and washed

3 times with cold FuoroBrite DMEM (Cat. #A1896701, Gibco) serum-free medium.

C2C12 dish was then set up on the Spinning disk microscope, applied 37°C

FuoroBrite DMEM serum-free medium containing 0, 0.2 or 20 nM human insulin (Cat.

#I9278, Sigma-Aldrich), and immediately started imaging in a 37°C chamber using a

100×/1.45 oil objective and QuantEM 512SC Photometrics camera on a spinning disk

confocal system based on Zeiss Axiovert 200M microscope (LSI Imaging Core, UBC). A

focal plane in the middle of the cell was acquired to avoid capturing the top and bottom of

the cell. Time-lapse imaging was conducted at 30 s intervals between 0 – 20 min or at 5 s

intervals between 20-30 min.

Immunofluorescence staining

Stablly-transfected C2C12 cell lines were seeded on #1.5 glass coverslip (Cat. #

72230-01, Electron Microscopy Sciences) and labelled with cell non-permeable

SNAP-Surface Alexa Fluor 488 substrate, incubated with 0, 0.2 or 20 nM human insulin

for 30 min, fixed with 4% (w/v) paraformaldehyde (Sigma-Aldrich) for 20 minutes at room

temperature, and washed 3 times with PBS. After permeabilized in 0.1% (v/v) Triton

X-100 (Sigma-Aldrich) for 15 minutes, primary antibodies targeting Caveolin1 (1:400,

Rabbit mAb, Cat. #3267, Cell Signaling) and CLTC heavy chain (1:200, Goat pAb, Cat.

#sc-6579, Santa Cruz) were incubated overnight at 4°C. Similarly, for AXNA2

experiment, cells labeled with SNAP-Surface Alexa Fluor 488 substrate were incubated

with insulin for 15 or 30 min, and labeled with Annexin A2 primary antibody (1:150,

Rabbit mAb, Cat. # 8235S, New England Biolabs). Coverslips were then washed in PBS,

blocked in Dako Protein Block (Cat #X0909, Agilent), and stained by secondary

antibodies (1:1000, Cy3 Donkey anti-Rabbit, Cat. #711-165-152, Jackson

ImmunoResearch; 1:500, Alexa Fluor 633 Donkey anti-Goat, Cat. # A21082, Invitrogen)

diluted in Dako Antibody Diluent (Cat. # S0809, Agilent) for 2 hours at room temperature.

After washing in PBS, coverslips were mounted with Prolong Diamond Antifade Mountant

(Cat. #P36961, Invitrogen).

STED microscopy and image analysis

Stimulated emission depletion (STED) microscopy was performed with the 100×/1.4

Oil HC PL APO CS2 STED White objective of a Leica TCS SP8 3× STED microscope

(Leica, Wetzlar, Germany) equipped with a white light laser, HyD detectors, and Leica

Application Suite X (LAS X) software (LSI Imaging Core, Life Sciences Institute,

University of British Columbia). Time-gated fluorescence detection was used to further

improve lateral resolution. INSR-SNAP labelled by Alexa Fluor 488 was excited at 488

nm and depleted using the 592-nm depletion laser. CAV1 or AXNA2 labelled by Cy3 was

excited at 555 nm and depleted using the 660 nm depletion laser. CLTC labelled by Alex

Fluor 633 was excited at 633 nm and depleted using the 775-nm depletion laser.

Sequential acquisition in the order of AF633/Cy3/AF488 was used to avoid cross-talk.

Huygens Professional software (Scientific Volume Imaging, Hilversum, the

Netherlands) was used to deconvolve STED images and calculate the Object Pearson

Coefficients for the colocalizations between INSR-SNAP, CAV1, CLTC, or ANXA2.

587   TIRF microscopy and single particle tracking analysis

Total internal reflection fluorescence (TIRF) microscopy was performed on a Zeiss

Axiovert 200M with a 100x Alpha-PlanFluar NA 1.45 oil objective (Zeiss, Germany) and a

TIRF laser angle modifier (Zeiss). Cells were kept at 37°C in a temperature-controlled

incubation chamber (Harvard Apparatus, Holliston, MA) and imaged in Ringer's buffer

supplemented with 20 mM glucose, as previously described[4]. For insulin stimulations,

cells were in the Ringer's (20 mM glucose) buffer containing 0 (control) or 2 nM of insulin.

Images were acquired with a CoolSNAP HQ2 CCD camera (Photometrics, Tucson, AZ).

405 nm and 561 nm solid-state diode laser system was used to excite TagBFP and

mRFP tagged proteins.

The ImageJ distribution Fiji was used to process microscope images[70]. The local

background of TIRF images was subtracted using the rolling ball method (20-pixel radius)

before single particle tracking.

Single particles (INSR, CAV1, or CLTC) were detected and tracked using Icy

bioimaging analysis software version 1.9.8.1[71]. Particles were detected by the

UnDecimated Wavelet Transform Detector plugin that was set to detect bright spots over

a dark background, exhibit 70% sensitivity to spots <3 pixels, detect particle size between

2 and 3000 pixels. Particles were tracked using the Multiple Hypothesis Tracking method

built into the Spot Tracking plugin that was set to recognize particles that exhibited either

diffusive or directed motion[72,73]. Track radius (maximum displacement of the particle

track) and lifetime of the particles were exported from the Motion Profile Processor of the

Spot Tracking plugin. Diffusion coefficients were calculated in MATLAB as described

previously but without corrections for positional errors and blurring[74]. Colocalization of

tracks was determined by Track Processor Interaction Analysis (distance threshold 2

pixels) in the Spot Tracking plugin. INSR tracks that colocalized with CAV1 or CLTC

tracks for more than 4 continuous frames are determined to be colocalized tracks. The

exported time and duration of colocalization were organized in Python and plotted in R

studio. Diffusion coefficients, track radius, and the lifetime of tracks were plotted in R

studio. Specifically, SuperPlots were used to plot the mean value of the cells and

individual tracks of each cell[75]. Code is available at GitHub deposit

https://github.com/hcen/INSR_tracking.

Skeletal muscle co-immunoprecipitation

Animal protocols were approved by the University of British Columbia Animal Care

Committee in accordance with national guidelines. Mice received 1.5U/kg insulin or PBS

through intraperitoneal injection. Skeletal muscle (gastrocnemius) was collected after 5

or 10 min of injection. INSR was immunoprecipitated by incubating 500 μg of protein

lysate with 2 μg INSR antibodies (Rabbit pAb, Cat. #sc-711, Santa Cruz) at 4°C

overnight. Subsequently, the solution was incubated with PureProteome™ Protein G

Magnetic Beads (Cat. # LSKMAGG10, Millipore, Burlington, MA, USA) and washed

according to the manufacturer's instructions. INSR protein complexes were eluted from

the beads by incubation in Blue Loading Buffer (50 mM DTT, Cell Signaling) at 95°C for 5

min and were analyzed by standard western blot procedures. INSR, CAV1, and CLTC

were detected by the following Cell Signaling primary antibodies: INSRβ (1:1000, Mouse

mAb, Cat. #3020S), CLTC (1:1000, Rabbit mAb, Cat. #4796), Caveolin1 (1:1000, Rabbit

mAb, Cat. #3267), and HRP-conjugated VeriBlot secondary antibodies (1:200,

anti-mouse Cat. #ab131368, anti-rabbit ab131366, Abcam) that do not bind to IgG heavy

and light chains to reduce background noise.

Immunoprecipitation-Mass Spectrometry

C2C12 myoblasts were serum-starved for 6 hours and were given 0 or 2 nM insulin for 15 min before collecting their lysates in RIPA buffer. INSRα antibody (NNC0276-3000, a gift from Novo Nordisk compound sharing program) that does not cross-react with IGF1R[24] was covalently conjugated to Dynabeads M-270 Epoxy beads (Cat. #14301, Invitrogen) to immunoprecipitate INSR according to the manufacturer's protocol. INSR protein complexes were eluted from the beads by incubation in Blue Loading Buffer (50 mM DTT, Cell Signaling) at 95°C for 5 min.

Eluted co-immunoprecipitation samples were run into 10% SDS PAGE gel for clean-up procedure, fixed, stained, and destained in water. Each of the samples was excised, reduced, alkylated, and digested according to standard in-gel digest protocols[76]. Extracted peptide samples were then cleaned up via STAGE-tip purification, as previously described[77].

Samples were reconstituted in 2% ACN, 0.5% formic acid, and the peptides were analyzed using a quadrupole – time of flight mass spectrometer (Impact II; Bruker Daltonics) on-line coupled to an EasyLC 1000 HPLC (ThermoFisher Scientific) using a Captive spray nanospray ionization source (Bruker Daltonics) including Aurora Series Gen2 (CSI) analytical column, (25cm x 75µm 1.6µm FSC C18, with Gen2 nanoZero and CSI fitting; Ion Opticks, Parkville, Victoria, Australia), and a µ-Precolumn, 300 µm ID x 5 mm, C18 PepMap, 5 µm, 100 A, Thermo Scientific, Waltham, Massachusetts, United States). Samples were reconstituted in (0.1% aqueous formic acid and 2 % acetonitrile in water) and 1ug of sample was loaded, injected in triplicates. A standard 90 min reversed-phase seperation was employed, using water:acetonitrile gradients.

LC and MS acquisition parameters were standard for peptide sequencing. Acquired data were then searched by MaxQuant (V1.6.7.) against the Uniprot protein database for Mouse C2C12 cell line, and LFQ intensities extracted and normalized using the MaxLFQ algorithm, with 20ppm and 30ppm mass accuracies for precursor and product ion

masses, respectively, and a 1% false discovery rate cut-off. Differential expression was

determined by comparing the mean protein intensities of the triplicate injections between

the groups. Significance (p<0.05) was calculated via t-test using default parameters in

Perseus (v1.6.14.0).

AlphaFold-Multimer and minD

Analysis of direct interactions between INSR and interactors was carried out using

AlphaFold-Multimer and minD method[33,34]. We downloaded all the protein sequences

from UniProt (IDs: INSR - P15208, RPSA - P14206, HSPB1 - P14602, ANXA2 - P07356,

HIST1H1B - P43276 and IGF1R - E9QNX9, CAV1 - P49817, CAV3 - P51637, AP2M1 -

P84091, IRS1 - P35569, MTCO2 - P00405). We then followed the previously described

protocol[34] to extract potential binding sites and do an in-silico evaluation of the

interactions between binding sites. Briefly, we first paired INSR with other proteins and

ran AlphaFold-Multimer on those pairs. We used full-length protein sequences to extract

minD scores. minD is defined as the minimum expected distance of one residue of one

protein chain from the other protein chain in the pairs. This metric is extracted from the

Distogram head of AlphaFold-Multimer and has been shown to peak when a potential

binding site is detected. After the binding sites on both proteins are identified, we then cut

the region surrounding them. This results in a handful of fragments for each pair. We then

pair

fragments for a second round of AlphaFold-Multimer runs. Finally, ipTM scores are

extracted to filter out non-binding pairs. A threshold of 0.42 is used to discriminate

non-interacting and interacting fragment pairs[34]. The code for this analysis is available at

https://github.com/alirezaomidi/AFminD.

Statistics

Data were presented as mean ± SEM in addition to the individual data points. A

significance level of p < 0.05 was used throughout. Student's t-tests were used for

comparisons between two groups. Comparisons between more than two groups were

performed using one-way ANOVA and multiple comparison p-values adjusted using the

Tukey method. Comparisons between multiple groups and two categorical variables were

performed using Two-way ANOVA followed by Tukey post hoc test. The statistical

analysis above was performed using GraphPad Prism (Version 9.0.2). Two-sided

Kolmogorov-Smirnov Tests were used for comparing cumulative probability, which was

performed with ks.text() function in R.

Data availability

Data are contained within the article or can be shared upon request.

References

1.   Knutson VP. Cellular trafficking and processing of the insulin receptor. *FASEB J.*
1991;5(8):2130-2138.
2.   Kahn CR, Crettaz M. Insulin receptors and the molecular mechanism of insulin
action. *Diabetes Metab Rev.* 1985;1(1-2):5-32.
3.   Bevan AP, Burgess JW, Drake PG, Shaver A, Bergeron JJ, Posner BI. Selective
activation of the rat hepatic endosomal insulin receptor kinase. Role for the
endosome in insulin signaling. *The Journal of biological chemistry.*
1995;270(18):10784-10791.
4.   Boothe T, Lim GE, Cen H, et al. Inter-domain tagging implicates caveolin-1 in
insulin receptor trafficking and Erk signaling bias in pancreatic beta-cells. *Mol*
Metab. 2016;5(5):366-378.
5.   Trischitta V, Gullo D, Squatrito S, Pezzino V, Goldfine ID, Vigneri R. Insulin
Internalization into Monocytes Is Decreased in Patients with Type II
Diabetes Mellitus. 1986;62:522-528.
6.   Trischitta V, Brunetti A, Chiavetta A, Benzi L, Papa V, Vigneri R. Defects in
insulin–receptor internalization and processing in monocytes
of obese subjects and obese NIDDM patients. 1989;38:1579–1584.

7. Jochen AL, Berhanu P, Olefsky JM. Insulin internalization and degradation in
adipocytes from normal and type II diabetic subjects. *J Clin Endocrinol Metab.*
1986;62(2):268-274.
8. Rask-Madsen C, Kahn CR. Tissue-specific insulin signaling, metabolic syndrome,
and cardiovascular disease. *Arterioscler Thromb Vasc Biol.*
2012;32(9):2052-2059.
9. Guo S. Insulin signaling, resistance, and the metabolic syndrome: insights from
mouse models into disease mechanisms. *The Journal of endocrinology.*
2014;220(2):T1-T23.
10. Jochen A, Hays J, Lee M. Kinetics of insulin internalization and processing in
adipocytes: effects of insulin concentration. *J Cell Physiol.* 1989;141(3):527-534.
11. McClain DA, Olefsky JM. Evidence for two independent pathways of
insulin-receptor internalization in hepatocytes and hepatoma cells. *Diabetes.*
1988;37(6):806-815.
12. Ceresa BP, Kao AW, Santeler SR, Pessin JE. Inhibition of clathrin-mediated
endocytosis selectively attenuates specific insulin receptor signal transduction
pathways. *Mol Cell Biol.* 1998;18(7):3862-3870.
13. Carpentier JL, Van Obberghen E, Gorden P, Orci L. Surface redistribution of
125I-insulin in cultured human lymphocytes. *J Cell Biol.* 1981;91(1):17-25.
14. Paccaud JP, Siddle K, Carpentier JL. Internalization of the human insulin receptor.
The insulin-independent pathway. *The Journal of biological chemistry.*
1992;267(18):13101-13106.
15. Jose M, Biosca JA, Trujillo R, Itarte E. Characterization of the hepatic insulin
receptor undergoing internalization through clathrin-coated vesicles and
endosomes. *FEBS Lett.* 1993;334(3):286-288.
16. Nystrom FH, Chen H, Cong LN, Li Y, Quon MJ. Caveolin-1 interacts with the
insulin receptor and can differentially modulate insulin signaling in transfected
Cos-7 cells and rat adipose cells. *Mol Endocrinol.* 1999;13(12):2013-2024.
17. Fagerholm S, Ortegren U, Karlsson M, Ruishalme I, Str$\sqrt{}$·lfors P. Rapid
insulin-dependent endocytosis of the insulin receptor by caveolae in primary
adipocytes. *PLoS ONE.* 2009;4(6):e5985.
18. Tseng LT, Lin CL, Tzen KY, Chang SC, Chang MF. LMBD1 protein serves as a
specific adaptor for insulin receptor internalization. *The Journal of biological*
chemistry. 2013;288(45):32424-32432.
19. Gautier A, Juillerat A, Heinis C, et al. An engineered protein tag for multiprotein
labeling in living cells. *Chem Biol.* 2008;15(2):128-136.
20. Cen H, Botezelli JD, Johnson JD. Modulation of Insr and insulin receptor signaling
by hyperinsulinemia in vitro and in vivo. *Biorxiv.* 2021.
21. Jans DA. Lateral Mobility of Polypeptide Hormone Receptors and GTP-Binding
Proteins. In: *The Mobile Receptor Hypothesis: The Role of Membrane Receptor*
Lateral Movement in Signal Transduction. Boston, MA: Springer US; 1997:83-115.

22. Kabayama K, Sato T, Saito K, et al. Dissociation of the insulin receptor and
caveolin-1 complex by ganglioside GM3 in the state of insulin resistance. *Proc*
Natl Acad Sci U S A. 2007;104(34):13678-13683.
23. Ikonen E, Vainio S. Lipid microdomains and insulin resistance: is there a
connection? *Sci STKE.* 2005;2005(268):pe3.
24. Ørstrup LH, Slaaby R, Rasch MG, et al. Cross-species reactive monoclonal
antibodies against the extracellular domains of the insulin receptor and IGF1
receptor. *J Immunol Methods.* 2019;465:20-26.
25. Benyoucef S, Surinya KH, Hadaschik D, Siddle K. Characterization of insulin/IGF
hybrid receptors: contributions of the insulin receptor L2 and Fn1 domains and the
alternatively spliced exon 11 sequence to ligand binding and receptor activation.
Biochem J. 2007;403(3):603-613.
26. Moxham CP, Duronio V, Jacobs S. Insulin-like growth factor I receptor
beta-subunit heterogeneity. Evidence for hybrid tetramers composed of
insulin-like growth factor I and insulin receptor heterodimers. *The Journal of*
biological chemistry. 1989;264(22):13238-13244.
27. Shu L, Lee L, Chang Y, et al. Caveolar structure and protein sorting are
maintained in NIH 3T3 cells independent of glycosphingolipid depletion. *Arch*
Biochem Biophys. 2000;373(1):83-90.
28. DiGiacomo V, Meruelo D. Looking into laminin receptor: critical discussion
regarding the non-integrin 37/67-kDa laminin receptor/RPSA protein. *Biol Rev*
Camb Philos Soc. 2016;91(2):288-310.
29. Zoubeidi A, Zardan A, Wiedmann RM, et al. Hsp27 promotes insulin-like growth
factor-I survival signaling in prostate cancer via p90Rsk-dependent
phosphorylation and inactivation of BAD. *Cancer Res.* 2010;70(6):2307-2317.
30. Muranova LK, Shatov VM, Bukach OV, Gusev NB. Cardio-Vascular Heat Shock
Protein (cvHsp, HspB7), an Unusual Representative of Small Heat Shock Protein
Family. *Biochemistry (Mosc).* 2021;86(Suppl 1):S1-S11.
31. Yuan H, Wang T, Niu Y, Liu X, Fu L. AMP-activated protein kinase-mediated
expression of heat shock protein beta 1 enhanced insulin sensitivity in the skeletal
muscle. *FEBS Lett.* 2017;591(1):97-108.
32. Jumper J, Evans R, Pritzel A, et al. Highly accurate protein structure prediction
with AlphaFold. *Nature.* 2021;596(7873):583-589.
33. Evans R, O'Neill M, Pritzel A, et al. Protein complex prediction with AlphaFold-Multimer. *bioRxiv.*
2022.
34. Omidi A, Møller MH, Malhis N, Bui JM, Gsponer J. AlphaFold-Multimer accurately
captures interactions and dynamics of intrinsically disordered protein regions.
Proc Natl Acad Sci U S A. 2024;121(44):e2406407121.
35. Chidlow JH, Sessa WC. Caveolae, caveolins, and cavins: complex control of
cellular signalling and inflammation. *Cardiovasc Res.* 2010;86(2):219-225.

36.   Capozza F, Cohen AW, Cheung MW, et al. Muscle-specific interaction of caveolin
803   isoforms: differential complex formation between caveolins in fibroblastic vs.
804   muscle cells. *Am J Physiol Cell Physiol.* 2005;288(3):C677-691.

805   37.   Schöneich C, Dremina E, Galeva N, Sharov V. Apoptosis in differentiating C2C12
806   muscle cells selectively targets Bcl-2-deficient myotubes. *Apoptosis.*
807   2014;19(1):42-57.

808   38.   Biener Y, Feinstein R, Mayak M, Kaburagi Y, Kadowaki T, Zick Y. Annexin II is a
809   novel player in insulin signal transduction. Possible association between annexin
810   II phosphorylation and insulin receptor internalization. *The Journal of biological
811   chemistry.* 1996;271(46):29489-29496.

812   39.   Rescher U, Ludwig C, Konietzko V, Kharitonenkov A, Gerke V. Tyrosine
813   phosphorylation of annexin A2 regulates Rho-mediated actin rearrangement and
814   cell adhesion. *J Cell Sci.* 2008;121(Pt 13):2177-2185.

815   40.   Chang M, Kwon M, Kim S, et al. Aptamer-based single-molecule imaging of
816   insulin receptors in living cells. *J Biomed Opt.* 2014;19(5):051204.

817   41.   Hubert M, Larsson E, Lundmark R. Keeping in touch with the membrane; protein-
818   and lipid-mediated confinement of caveolae to the cell surface. *Biochem Soc
819   Trans.* 2020;48(1):155-163.

820   42.   Martins AS, Ordóñez JL, Amaral AT, et al. IGF1R signaling in Ewing sarcoma is
821   shaped by clathrin-/caveolin-dependent endocytosis. *PLoS One.*
822   2011;6(5):e19846.

823   43.   Pandini G, Frasca F, Mineo R, Sciacca L, Vigneri R, Belfiore A. Insulin/insulin-like
824   growth factor I hybrid receptors have different biological characteristics depending
825   on the insulin receptor isoform involved. *The Journal of biological chemistry.*
826   2002;277(42):39684-39695.

827   44.   Tomas A, Futter CE, Eden ER. EGF receptor trafficking: consequences for
828   signaling and cancer. *Trends Cell Biol.* 2014;24(1):26-34.

829   45.   Kammanadiminti SJ, Chadee K. Suppression of NF-kappaB activation by
830   Entamoeba histolytica in intestinal epithelial cells is mediated by heat shock
831   protein 27. *The Journal of biological chemistry.* 2006;281(36):26112-26120.

832   46.   Ku HC, Chang HH, Liu HC, et al. Green tea (-)-epigallocatechin gallate inhibits
833   insulin stimulation of 3T3-L1 preadipocyte mitogenesis via the 67-kDa laminin
834   receptor pathway. *Am J Physiol Cell Physiol.* 2009;297(1):C121-132.

835   47.   Zhao WQ, Chen GH, Chen H, et al. Secretion of Annexin II via activation of insulin
836   receptor and insulin-like growth factor receptor. *The Journal of biological
837   chemistry.* 2003;278(6):4205-4215.

838   48.   Caron D, Boutchueng-Djidjou M, Tanguay RM, Faure RL. Annexin A2 is
839   SUMOylated on its N-terminal domain: regulation by insulin. *FEBS Lett.*
840   2015;589(9):985-991.

49. Urbanska A, Sadowski L, Kalaidzidis Y, Miaczynska M. Biochemical
characterization of APPL endosomes: the role of annexin A2 in APPL membrane
recruitment. *Traffic.* 2011;12(9):1227-1241.
50. Ryu J, Galan AK, Xin X, et al. APPL1 potentiates insulin sensitivity by facilitating
the binding of IRS1/2 to the insulin receptor. *Cell Rep.* 2014;7(4):1227-1238.
51. Lemmon MA, Schlessinger J. Cell signaling by receptor tyrosine kinases. *Cell.*
2010;141(7):1117-1134.
52. de Graauw M, Cao L, Winkel L, et al. Annexin A2 depletion delays EGFR
endocytic trafficking via cofilin activation and enhances EGFR signaling and
metastasis formation. *Oncogene.* 2014;33(20):2610-2619.
53. Behrends M, Engmann O. Linker histone H1.5 is an underestimated factor in
differentiation and carcinogenesis. *Environ Epigenet.* 2020;6(1):dvaa013.
54. Brown M, Dainty S, Strudwick N, et al. Endoplasmic reticulum stress causes
insulin resistance by inhibiting delivery of newly synthesized insulin receptors to
the cell surface. *Mol Biol Cell.* 2020;31(23):2597-2629.
55. Vicidomini G, Bianchini P, Diaspro A. STED super-resolved microscopy. *Nat*
Methods. 2018;15(3):173-182.
56. Fish KN. Total internal reflection fluorescence (TIRF) microscopy. *Curr Protoc*
Cytom. 2009;Chapter 12:Unit12.18.
57. Le PU, Guay G, Altschuler Y, Nabi IR. Caveolin-1 is a negative regulator of
caveolae-mediated endocytosis to the endoplasmic reticulum. *Journal of*
Biological Chemistry. 2002;277(5):3371-3379.
58. Humphries AC, Way M. The non-canonical roles of clathrin and actin in pathogen
internalization, egress and spread. *Nat Rev Microbiol.* 2013;11(8):551-560.
59. Rodal SK, Skretting G, Garred O, Vilhardt F, van Deurs B, Sandvig K. Extraction
of cholesterol with methyl-beta-cyclodextrin perturbs formation of clathrin-coated
endocytic vesicles. *Mol Biol Cell.* 1999;10(4):961-974.
60. Parpal S, Karlsson M, Thorn H, Strålfors P. Cholesterol depletion disrupts
caveolae and insulin receptor signaling for metabolic control via insulin receptor
substrate-1, but not for mitogen-activated protein kinase control. *The Journal of*
biological chemistry. 2001;276(13):9670-9678.
61. Cohen AW, Combs TP, Scherer PE, Lisanti MP. Role of caveolin and caveolae in
insulin signaling and diabetes. *Am J Physiol Endocrinol Metab.*
2003;285(6):E1151-1160.
62. Hussain KM, Leong KL, Ng MM, Chu JJ. The essential role of clathrin-mediated
endocytosis in the infectious entry of human enterovirus 71. *The Journal of*
biological chemistry. 2011;286(1):309-321.
63. Wu J, Park SH, Choi E. The insulin receptor endocytosis. *Prog Mol Biol Transl Sci.*
2023;194:79-107.
64. Chen Y, Huang L, Qi X, Chen C. Insulin Receptor Trafficking: Consequences for
Insulin Sensitivity and Diabetes. *Int J Mol Sci.* 2019;20(20).

65.   Choi E, Zhang X, Xing C, Yu H. Mitotic Checkpoint Regulators Control Insulin
883   Signaling and Metabolic Homeostasis. *Cell.* 2016;166(3):567-581.
884   66.   Park J, Hall C, Hubbard B, et al. MAD2-Dependent Insulin Receptor Endocytosis
885   Regulates Metabolic Homeostasis. *Diabetes.* 2023;72(12):1781-1794.
886   67.   Choi E, Kikuchi S, Gao H, et al. Mitotic regulators and the SHP2-MAPK pathway
887   promote IR endocytosis and feedback regulation of insulin signaling. *Nat*
888   *Commun.* 2019;10(1):1473.
889   68.   Ansarullah, Jain C, Far FF, et al. Inceptor counteracts insulin signalling in β-cells
890   to control glycaemia. *Nature.* 2021;590(7845):326-331.
891   69.   Siehler J, Bilekova S, Chapouton P, et al. Inceptor binds to and directs insulin
892   towards lysosomal degradation in β cells. *Nat Metab.* 2024;6(12):2374-2390.
893   70.   Schindelin J, Arganda-Carreras I, Frise E, et al. Fiji: an open-source platform for
894   biological-image analysis. *Nat Methods.* 2012;9(7):676-682.
895   71.   de Chaumont F, Dallongeville S, Chenouard N, et al. Icy: an open bioimage
896   informatics platform for extended reproducible research. *Nat Methods.*
897   2012;9(7):690-696.
898   72.   Chenouard N, Bloch I, Olivo-Marin JC. Multiple hypothesis tracking for cluttered
899   biological image sequences. *IEEE Trans Pattern Anal Mach Intell.*
900   2013;35(11):2736-3750.
901   73.   Chenouard N, Smal I, de Chaumont F, et al. Objective comparison of particle
902   tracking methods. *Nat Methods.* 2014;11(3):281-289.
903   74.   Abraham L, Lu HY, Falcão RC, et al. Limitations of Qdot labelling compared to
904   directly-conjugated probes for single particle tracking of B cell receptor mobility.
905   *Sci Rep.* 2017;7(1):11379.
906   75.   Lord SJ, Velle KB, Mullins RD, Fritz-Laylin LK. SuperPlots: Communicating
907   reproducibility and variability in cell biology. *J Cell Biol.* 2020;219(6).
908   76.   Shevchenko A, Wilm M, Vorm O, Mann M. Mass spectrometric sequencing of
909   proteins silver-stained polyacrylamide gels. *Anal Chem.* 1996;68(5):850-858.
910   77.   Rappsilber J, Mann M, Ishihama Y. Protocol for micro-purification, enrichment,
911   pre-fractionation and storage of peptides for proteomics using StageTips. *Nat*
912   *Protoc.* 2007;2(8):1896-1906.

Figure legends

Figure 1. Biochemical and live-cell imaging characterization of the effects of insulin on INSR.

(A) Internalized to total INSR ratio was quantified using surface biotinylation in undifferentiated C2C12 myoblasts. Cells were incubated in PBS (no insulin) or serum-free DMEM containing 0, 0.2, 2, or 20 nM insulin for 15 min. (p>0.05 when not specified.) **(B)** Western blot showing SNAP-tagged INSR expressed from lentiviral vector in comparison to wild-type INSR in C2C12 myoblasts. **(C,D).** Representative images of INSR-A-SNAP -labelled using an Alexa Fluor 488 cell non-permeable dye in 0, 0.2 or 20 nM insulin conditions in live undifferentiated C2C12 myoblasts from **(C)** 0 - 15 min or **(D)** after 20 min using a spinning disk confocal microscope. Time-lapse images of INSR vesicle interactions starting from the selected subregions (white squares) of snapshots are shown in the *insets*. (n=3 cells)

Figure 2. Live-cell TIRF imaging of cell-surface INSR.

(A) Experimental design of TIRF microscopy of the same cells at 3 timepoints with 2 nM insulin or no insulin (control). **(B)** Representative TIRF image of INSR-A-EGFP puncta, binary image of detected INSR spots, and INSR tracks. **(C)** Diffusion coefficient of INSR-A-EGFP tracks in control or insulin group. Data is plotted as SuperPlot with larger dots showing mean values of the cells and smaller dots showing individual INSR tracks of

the cells. **(D)** The cumulative probability shows the distribution of diffusion coefficients in

(C). **(E)** Track radius of INSR-A-EGFP tracks in control or insulin group. **(F)** The

cumulative probability shows the distribution of diffusion coefficient in **(E)**. (n=3 cells in

control group, n=9 cells in insulin group)

Figure 3. Identification of INSR interactors.

(A) Proteins that are detected in INSR immunoprecipitation mass-spectrometry (IP-MS),

and significantly different between wild-type and INSR knockout groups. Proteins in

green area were specific interactors that had little detection in knockout cells. The other 3

proteins are also significantly different but not lower or absent in INSR KO cells. (n=3) **(B)**

Protein-protein interaction network of INSR interactors and endocytosis proteins. **(C)**

Modeling the interactions between the cytoplasmic region of INSR (residues 968 - 1372)

and full-length interactor (target) proteins using AlphaFold-Multimer. pTM scores

measure the accuracy of the overall structure of the protein complex and is relatively

insensitive to localized inaccuracies. ipTM measures the accuracy of the interacting

subunits of the complex, and ipTM > 0.42 (dash line) predicts direct interaction. The

target proteins are classified as IP-MS identified novel interactors (Unknown ●), the

known or expected interactors (positive **+**), and an expected non-interactor protein as a

negative control (negative **×**). The cytoplasmic region of IGF1R (residues 961-1369) is

used. (n = 5 predictions.) **(D)** Potential interaction sites based on per-residue scores

along the INSR cytoplasmic region (residues 968 to 1372; top panel) and interactor

(target) proteins Ap2m1 (bottom). minD (more accurate) and min-iPAE scores detect

interaction sites. pLDDT shows whether the region is structured or unstructured. Shades

show confidence errors of 99%. **(E)** ipTM scores for fragment-protein interactions. Thirty

residues around the minD peaks (potential interacting sites) in **(D)** and **Fig. S2** are

analyzed. Each box relates to the interaction of the fragments of first protein vs the

full-lenght second protein. For example, title "Insr vs Mtco2" means we have cut Insr into

fragments and we have run AlphaFold-Multimer on each of those fragments vs full-length

Mtco2. Each dot is the top ipTM score for one fragment (we do 5 predictions for each

fragment vs protein). The protein residue positions of the interacting fragments (ipTM >

0.42) are labeled.

Figure 4. Interaction and colocalization between INSR, CAV3, CAV1 and CLTC

under different insulin stimulations.

(A) Western blot of mice skeletal muscle lysates after insulin injection. Phospho-AKT

(Ser473) to total AKT ratio or phospho-ERK1/2 to total ERK ratio were quantified.

(n=6) **(B)** Co-immunoprecipitation of skeletal muscle INSR after PBS or insulin injection.

CLTC to INSR ratio or CAV3 to INSR ratio were quantified. (n=7-10) **(C)** Representative

STED microscopy images of C2C12 myoblasts expressing INSR-A-SNAP (surface

labeled) that were fixed after stimulation with 0 nM, 0.2 nM, or 20 nM insulin for 30 min

and stained for CAV1 and CLTC (Scale bar = 5 µm). **(D,E)** Colocalizations between

INSR-A-SNAP, CAV1, and CLTC were quantified by Object Pearson Coefficient. Data

are plotted to show differences between insulin concentrations **(D)** or between protein

pairs **(E)**. (n=7-9 images, 1-4 cells per image. *p<0.05, Tukey's multiple comparison after

2-ANOVA. Box represents median and 25th to 75th percentiles.)

Figure 5. Colocalization and interaction between INSR and ANXA2 under different

insulin concentrations.

(A,B) Representative STED microscopy images of C2C12 myoblasts expressing

INSR-A-SNAP (surface labeled) that were fixed after stimulation with 0 nM, 0.2 nM, or 20

nM insulin for (A)15 or (B)30 min and stained for ANXA2 (Scale bar = 5 µm). **(C,D)**

Colocalization between INSR-A-SNAP and ANXA2 at (C)15 or (D)30 min of

insulin-stimulated were quantified by Object Pearson Coefficient. (n=4-13 images, 1- 3

cells per image. *p<0.05, 1-ANOVA. Box represents median and 25th to 75th

percentiles.)

1002

Figure 6. Live-cell TIRF imaging of cell-surface INSR, CAV1 and CLTC.

(A,B)TIRF microscopy (2s/frame) showed colocalization between INSR-B-TagBFP and

(A) CAV1-mRFP or **(B)** CLTC-mRFP. **(C,D)**The associated fluorescent intensity of

representative INSR-B-TagBFP and colocalized **(C)** CAV1-mRFP or **(D)** CLTC-mRFP.

(E-G) The (E) diffusion coefficient, (F) track radius, and (G) lifetime of INSR-B-TagBFP

tracks in cells expressing CAV1-mRFP under 0 nM (n=2 cells, 360 tracks) or 2 nM insulin

(n=2 cells, 324 tracks) or expressing CLTC-mRFP under 0 nM (n=4 cells, 917 tracks) or 2

nM insulin (n=2 cells, 146 tracks). Data is plotted as SuperPlot, with bigger dots showing

mean values of the cells and smaller dots showing individual INSR tracks of the cells. The

cumulative probability shows the distribution of the diffusion coefficient, with the dots and

numbers showing the median values. (2-ANOVA: # effect of co-expression;   post-hoc

Tukey test: *p<0.05; Kolmogorov-Smirnov (KS) test: *p<0.05, **p<0.005, ***p<0.0005.)

Experiments shown were conducted in Ringer's buffer.

1016

Figure 7. Analysis of INSR tracks colocalized or not colocalized with CAV1 or

CLTC.

(A) INSR tracks that colocalized with CAV1 under 0 or 2 nM insulin (referred to as

CAV1.Y.0nM or CAV1.Y.2nM below). **(B)** INSR tracks that colocalized with CLTC under 0

or 2 nM insulin (referred to as CLTC.Y.0nM or CLTC.Y.2nM below). **(C)** Diffusion

coefficient of INSR-B-TagBFP tracks that did not colocalize with CAV1-mRFP tracks

under 0 nM insulin (CAV1.N.0nM) or 2 nM insulin (CAV1.N.2nM) or colocalized with

CAV1-mRFP tracks under 0 nM insulin (CAV1.Y.0nM) or 2 nM insulin (CAV1.Y.2nM).

The cumulative probability shows the distribution of the measurements, with the dots and

numbers showing the median values (same for other cumulative probability plots). **(D)**

Diffusion coefficient of INSR-B-TagBFP tracks that did not colocalize with CLTC-mRFP

tracks under 0 nM insulin (CLTC.N.0nM) or 2 nM insulin (CLTC.N.2nM) or colocalized

with CLTC-mRFP tracks under 0 nM insulin (CLTC.Y.0nM) or 2 nM insulin (CLTC.Y.2nM).

(E) Track radius of INSR-B-TagBFP tracks that did not colocalize with CAV1-mRFP

tracks under 0 nM insulin (CAV1.N.0nM) or 2 nM insulin (CAV1.N.2nM) or colocalized

with CAV1-mRFP tracks under 0 nM insulin (CAV1.Y.0nM) or 2 nM insulin (CAV1.Y.2nM).

(F) Track radius of INSR-B-TagBFP tracks that did not colocalize with CLTC-mRFP

tracks under 0 nM insulin (CLTC.N.0nM) or 2 nM insulin (CLTC.N.2nM) or colocalized

with CLTC-mRFP tracks under 0 nM insulin (CLTC.Y.0nM) or 2 nM insulin (CLTC.Y.2nM).

(G) Lifetime of INSR-B-TagBFP tracks that did not colocalize with CAV1-mRFP tracks

under 0 nM insulin (CAV1.N.0nM) or 2 nM insulin (CAV1.N.2nM) or colocalized with

CAV1-mRFP tracks under 0 nM insulin (CAV1.Y.0nM) or 2 nM insulin (CAV1.Y.2nM). **(H)**

Lifetime of INSR-B-TagBFP tracks that did not colocalize with CLTC-mRFP tracks under

0 nM insulin (CLTC.N.0nM) or 2 nM insulin (CLTC.N.2nM) or colocalized with

CLTC-mRFP tracks under 0 nM insulin (CLTC.Y.0nM) or 2 nM insulin (CLTC.Y.2nM).

(CAV1.N.0nM, n=2 cells, 125 tracks; CAV1.N.2nM, n=2 cells, 96 tracks; CAV1.Y.0nM,

n=2 cells, 235 tracks; CAV1.Y.2nM, n=2 cells, 228 tracks; CLTC.N.0nM, n= 4 cells,   597

tracks; CLTC.N.2nM, n= 2 cells, 91 tracks; CLTC.Y.0nM, n= 4 cells, 320 tracks;

CLTC.Y.2nM, n= 2 cells, 55 tracks.) (2-ANOVA: # effect of colocalization, $ effect of

insulin, % interaction of the 2 variables; Kolmogorov-Smirnov (KS) test: *p<0.05,

***p<0.0005.) Experiments shown were conducted in Ringer's buffer.

1048

Figure 8. Summary of the effects of CAV1 and CLTC on INSR tracks. Higher insulin

(20 nM) promoted INSR and CAV1 colocalization, while lower insulin (0.2 nM) promoted

INSR and CLTC colocalization. INSR tracks had lower diffusion coefficient and track

radius in CAV1-overexpressing cells than CLTC-overexpressing cells. Within the same

cells, INSR tracks colocalized with CAV1 had longer lifetimes and larger track radius than

non-colocalized tracks. INSR tracks colocalized with CLTC had longer lifetimes than

non-colocalized tracks, which is further increased by insulin.

1056

1057 **Supplementary Figures and Videos**

1058

Figure S1. Insulin signalling of C2C12 myoblast overexpressing

inter-domain-tagged INSR-A-EGFP.

(A) Representative western blot image of C2C12 cells after 5 min of insulin stimulation.

Control lanes were samples without overexpressing tagged INSR. # indicates unspecific

band, which is previously reported in C2C12 cells[54]. It does not always appear[54]; in our

hands, it didn't appear in INSR-A-EGFP lanes or other samples in Fig. 1B. A lot more

proteins were unfortunately loaded for control samples, shown by α-tubulin bands as

loading control. Total AKT and ERK were not successfully blotted due to technical issues.

(B) Percentage overexpression of INSR-A-EGFP, showing a moderate overexpression of

~35%. It is calculated by the intensity of the INSRβ-EGFP band divided by the

endogenous INSRβ of the same lane. (n = 2 biological replicates.) **(C, D)** pAKT and

pERK normalized by **(C)** tubulin or **(D)** INSR (INSRβ band or the sum of INSRβ and

INSRβ-EGFP bands). Due to unequal loading, the intensities of WT and INSR-A-EGFP

groups are less directly comparable (as the chemiluminescence of western blot is only

semi-quantitative with a limited linear range); therefore, each group is normalized by the

mean of its own 0 nM insulin condition to make the signalling more comparable. Overall,

the pAKT and pERK activations in cells overexpressing INSR-A-EGFP were similar to or

no less than the controls. (n = 2 biological replicates.)

1077

1078 **Figure S2. Potential interaction sites based on per-residue scores along the**

1079 **interactor (target) proteins against the INSR cytoplasmic region (residues 968 to**

1080 **1372).**

1081   minD (more accurate) and min-iPAE scores detect interaction sites. pLDDT shows

1082   whether the region is structured or unstructured. Shades show confidence errors of 99%.

1083

1084   **Video 1. Live-cell imaging of surface-labeled INSR-A-SNAP from 0 to 15 min.**

1085

1086   **Video 2. Live-cell imaging of surface-labeled INSR-A-SNAP from 20 to 30 min.**

Figure 1. Biochemical and live-cell imaging characterization of the effects of insulin on INSR.

Figure 2. Live-cell TIRF imaging of cell-surface INSR.

Figure 3. Identification of INSR interactors.

Figure 4. Interaction and colocalization between INSR, CAV3, CAV1 and CLTC under different insulin stimulations.

Figure 5. Colocalization between INSR and ANXA2 under different insulin concentrations.

Figure 6. Live-cell TIRF imaging of cell-surface INSR, CAV1 and CLTC.

Figure 7. Analysis of INSR tracks colocalized or not colocalized with CAV1 or CLTC.

Figure 8. Summary of the effects of CAV1 and CLTC on INSR tracks.

Supplementary Figures

Figure S1. Insulin signalling of C2C12 myoblast overexpressing inter-domain-tagged INSR-A-EGFP.
(A) Representative western blot image of C2C12 cells after 5 min of insulin stimulation. Control lanes were samples without overexpressing tagged INSR. # indicates unspecific band, which is previously reported in C2C12 cells[54]. It does not always appear[54]; in our hands, it didn't appear in INSR-A-EGFP lanes or other samples in Fig. 1B. A lot more proteins were unfortunately loaded for control samples, shown by α-tubulin bands as loading control. Total AKT and ERK were not successfully blotted due to technical issues. **(B)** Percentage overexpression of INSR-A-EGFP, showing a moderate overexpression of ~35%. It is calculated by the intensity of the INSRβ-EGFP band divided by the endogenous INSRβ of the same lane. (n = 2 biological replicates.) **(C, D)** pAKT and pERK normalized by **(C)** tubulin or **(D)** INSR (INSRβ band or the sum of INSRβ and INSRβ-EGFP bands). Due to unequal loading, the intensities of WT and INSR-A-EGFP groups are less directly comparable (as the chemiluminescence of western blot is only semi-quantitative with a limited linear range); therefore, each group is normalized by the mean of its own 0 nM insulin condition to make the signalling more comparable. Overall, the pAKT and pERK activations in cells overexpressing INSR-A-EGFP were similar to or no less than the controls. (n = 2 biological replicates.)

Figure S2. Potential interaction sites based on per-residue scores along the interactor (target) proteins against the INSR cytoplasmic region (residues 968 to 1372).
minD (more accurate) and min-iPAE scores detect interaction sites. pLDDT shows whether the region is structured or unstructured. Shades show confidence errors of 99%.

Figure S1. Insulin signaling of C2C12 myoblast overexpressing inter-domain-tagged INSR-A-EGFP.

Figure S2. Potential interaction sites based on per-residue scores along the interactor (target) proteins against the INSR cytoplasmic region (residues 968 to 1372).