**Interacting effects of cold snaps, rain, and agriculture on the fledging success of a declining**

**aerial insectivore**

Daniel R. Garrett*[1,2], Fanie Pelletier[1], Dany Garant[1], and Marc Bélisle[1]

Author's Affiliations

[1]Département de biologie, Université de Sherbrooke, 2500 Boulevard de l'Université,

Sherbrooke, Québec J1K2R1 Canada

[2]daniel.garrett88@gmail.com

*Corresponding author: Université de Sherbrooke, 2500 boulevard de l'Université, Sherbrooke,

Qc, Canada, J1K 2R1, daniel.garrett88@gmail.com

Running Head: Weather, agriculture, and avian fitness

**Abstract**

Climate change predicts the increased frequency, duration, and intensity of inclement weather

periods, such as unseasonably low temperatures (i.e., cold snaps) and prolonged precipitation.

Many migratory species have advanced the phenology of important life history stages, and as a

result will likely be exposed to these periods of inclement spring weather more often, thus

risking reduced fitness and population growth. For declining avian species, including aerial

insectivores, anthropogenic landscape changes such as agricultural intensification are another

driver of population declines. These landscape changes may affect the foraging ability of food

provisioning parents, and reduce the survival of nestlings exposed to inclement weather, through

for example pesticide exposure impairing thermoregulation and punctual anorexia. Breeding in

agro-intensive landscapes may thus exacerbate the negative effects of inclement weather under

climate change. We observed that a significant reduction in the availability of insect prey

occurred when daily maximum temperatures fell below 18.3°C, and thereby defined any day

where the maximum temperature fell below this value as a day witnessing a cold snap. We then

combined daily information on the occurrence of cold snaps and measures of precipitation to

assess their impact on the fledging success of Tree Swallows (*Tachycineta bicolor*) occupying a

nest box system placed across a gradient of agricultural intensification. Estimated fledging

success of this declining aerial insectivore was 36.2% lower for broods experiencing four cold

snap days during the 12 days post hatching period versus broods experiencing none, and this

relationship was worsened when facing more precipitation. We further found that the overall

negative effects of a brood experiencing periods of inclement weather was exacerbated in more

agro-intensive landscapes. Our results indicate that two of the primary hypothesized drivers of

many avian population declines may interact to further increase the rate of declines in certain

landscape contexts.

Key words: aerial insectivores; agricultural intensification; climate change; fledging success;

precipitation; severe weather events; *Tachycineta bicolor*; Tree Swallows.

## Introduction

Several avian groups are experiencing dramatic population declines (Rosenberg et al. 2019). Population estimates of many North American farmland and grassland birds, including aerial insectivores, suggest declines of over 50% since the 1970s (Stanton et al. 2018, Rosenberg et al. 2019). The spatio-temporal occurrence of these declines has led to the hypothesis that they are driven by a combination of global climate change and large-scale landscape modifications (Stanton et al. 2018, Spiller and Dettmers 2019), and particularly the process of agricultural intensification (Stanton et al. 2018).

In the northern hemisphere, global climate change has resulted in earlier spring temperatures (Hartmann et al. 2013). For instance, in USA the first fall frost has advanced by nearly 5.5 days and the last spring frost occurs nearly 7 days sooner when compared to the previous century (Kukal and Irmak 2018). The swift rate of change in timing of temperatures may lead to migrating species being unable to compensate for changes in thermally suitable habitats through dispersal, plasticity, or evolution (Crick 2004, Visser and Gienapp 2019). The inability of species to respond to such changes may generate or accentuate phenological mismatches between peaks of seasonal food resources and peaks in resource requirements of migratory species, and as a result, cause fitness decreases due to overall mistimed breeding schedules (Visser and Gienapp 2019). Furthermore, global climate change predicts increases in the frequency and intensity of inclement spring weather, such as swift changes in ambient temperature (e.g., cold snaps) and prolonged periods of precipitation (Rahmstorf and Coumou 2011, Wuebbles et al. 2014). Though often difficult to define (van de Pol et al. 2017), extremes in weather (e.g., storms, blizzards, and oceanic heatwaves) can have significant impacts on ecosystem functions (reviewed in Ummenhofer and Meehl 2017). In birds, even slightly lower than average daily temperatures or

greater than average precipitation can lower fitness (Tuero et al. 2018, de Zwaan et al. 2019,

Skwarska et al. 2021). Individuals experiencing such periods of inclement weather may be

subjected to the reduction of temperature-dependent food resources, as well as to the additional

consequence of the direct effects of poor weather (Pipoly et al. 2013, Moreno et al. 2015,

Arbeiter et al. 2016, Marcelino et al. 2020). Therefore, migrating species adapting to global

changes through advancing their spring migration, may be at greater risk of experiencing periods

of inclement weather (Both et al. 2010, Visser and Gienapp 2019). The phenological

advancement of several declining species has recently been observed (Dunn and Winkler 1999,

Møller et al. 2006, Townsend et al. 2013, Bourret et al. 2015). For example, two Tree Swallow

(*Tachycineta bicolor*) populations located respectively in Ontario, Canada, and New York State,

USA, exhibit both chronic population declines and advanced clutch initiation dates (Dunn and

Winkler 1999, Shutler et al. 2012). During the same period of declines, a significant increase in

the number of poor weather days (e.g., unseasonably low temperatures or prolonged

precipitation) was observed and resulted in elevated nestling mortality (Cox et al. 2019, Shipley

et al. 2020).

The declines of farmland and grassland birds are correlated with not only climate change but also

with the process of agricultural intensification (Stanton et al. 2018, Spiller and Dettmers 2019).

Hypothesized agricultural drivers of declines are divided into direct (e.g., habitat loss,

mechanization resulting in nest destruction, or acute pesticide exposure) and indirect effects,

principally through the reduction in prey availability or sublethal pesticide exposure (Benton et

al. 2003, Stanton et al. 2018). Certain agricultural practices are observed to alter insect

populations and communities through changes in abundance, phenology, or species composition

and interactions (Grüebler et al. 2008, Pisa et al. 2015, Wagner 2020). Insectivorous birds

breeding in modern agricultural landscapes, including aerial insectivores, could thereby be

subjected to reduced availability of prey items (Poulin et al. 2010, Nocera et al. 2012, Garrett et

al. 2021a).

Agricultural drivers of population declines may also impact the physiology of birds in ways that

influence their capacity to respond adaptively to periods of inclement weather. For instance,

sublethal exposure to organophosphate and carbamate insecticides may induce short-lived

hypothermia, likely due to the impairment of thermoregulation (Grue et al. 1997). Moreover,

sublethal exposure to neonicotinoid insecticides may result in anorexia (Eng et al. 2019), thereby

aggravating reduced thermoregulatory capacity and other challenges posed by reduced prey

availability (Garrett et al. 2021a). Birds breeding within more agro-intensive landscapes are at

elevated risk of exposing themselves or their offspring to such agents (DiBartolomeis et al. 2019,

Malaj et al. 2020, Sigouin et al. 2021) and may subsequently be less likely to survive through

periods of poor weather. Furthermore, landscapes dominated by row crop monocultures express

landscape simplification in which large swaths of areas are occupied by only a handful of

habitats (Benton et al. 2003). This phenomenon may make finding and exploiting suitable food

resources more difficult for animals relying upon "residual" marginal habitats, as landscape

simplification lowers the functional connectivity of agricultural landscapes (Hinsley 2000,

Bélisle 2005, Rainho and Palmeirim 2011). Parents may compensate for poor breeding landscape

quality by increasing foraging effort, but such increases may result in reduced body condition

(Hinsley 2000, Olsson et al. 2008). Periods of inclement weather may thus exacerbate the added

stressors stemming from breeding within more agro-intensive landscapes (Stanton et al. 2016,

Staggenborg et al. 2017, Evens et al. 2018, Garrett et al. 2021b), and may even influence the

capacity for parental care and investment (de Zwaan et al. 2019). Therefore, given that selection

favors earlier breeding events (Sheldon et al. 2003, Porlier et al. 2012, Marrot et al. 2017), it

seems imperative to evaluate how the expected increase of inclement weather events may

interact with the consequences of breeding within more agro-intensive landscapes.

Here we present the results of an 11-year study monitoring the breeding success of a Tree

Swallow population experiencing a wide range of spring temperatures and precipitation. This

population breeds within a nesting box system placed along a gradient of agricultural

intensification in southern Québec, Canada. We first identified a critical temperature in which

the availability of their main prey, namely Diptera (Bellavance et al. 2018), changed

significantly, using an approach similar to Winkler et al. (2013). We then defined days in which

ambient temperature fell below this critical temperature as a day representing a cold snap. We

then evaluated the interacting roles that cold snaps, prolonged precipitation and the gradient of

agricultural intensification had on fledging success within this population. As observed by Cox et

al. (2019) and Shipley et al. (2020), we expected cold snaps to reduce fledging success, and the

effect size to increase with both the increasing number of cold snaps and the total amount of

precipitation experienced by broods. Finally, we expected the severity of these relationships to

increase as breeding landscapes became increasingly composed of agro-intensive cultures.

**Methods**

**Study area and nest box system**

In 2004, 10 nest boxes, spaced approximately 50 m apart, were placed along the field margins of

40 separate farms dispersed across a gradient of agricultural intensification in southern Québec,

Canada (Fig. 1; see Ghilain and Bélisle 2008 for details). In 2006 we started intensively

monitoring the breeding activity of these nest boxes and continued through the breeding season

of 2016. This monitoring protocol resulted in 400 nest boxes being actively monitored

throughout this period. The gradient of agricultural intensification was characterized by an east-

west shift of agricultural production. The eastern portion of the system was composed primarily

of pastures and forage crops (e.g., hay, alfalfa (*Madicago sativa*), and clover (*Trifolium spp.*)),

requiring less agricultural inputs and interspersed with large expanses of forest cover. The west

was dominated by large-scale row-crop monocultures (principally corn (*Zea mays*), soybean

(*Glycine max*) and wheat (*Triticum spp.*)) and was denuded of forest cover (Jobin et al. 2003,

Ruiz and Domon 2009). Increased use of monocultures has resulted in a near reliance on

fertilizers, pesticides, mechanization and drainage of surface waters and wetlands (Jobin et al.

2003). Between 2011 to 2019, nearly 100% of the corn and 60% of the soybean were sown as

neonicotinoid-coated seeds in our study area (MDDELCC 2015). As a consequence, the rate at

which several pesticides were detected above the levels deemed safe for the chronic exposure for

aquatic life has increased in the surface waters of this region (Giroux 2019).

**Nest box monitoring**

Starting prior to the arrival of Tree Swallows, we monitored the breeding activity in nest boxes

every two days throughout each breeding season. This monitoring schedule allowed us to

estimate the dates of laying, incubation, hatching, and fledging, as well as to record the number

of eggs, nestlings, and fledglings of each breeding event. On average, first-clutch-occupancy by

Tree Swallows was 60% ranging between 51% and 74% throughout the 11-year data set. We

caught and banded adult females during incubation and adult males during food provisioning.

We were 99% and 80% successful, respectively, at capturing targeted adults. We applied an

aluminum US Geological Survey (USGS) band containing a unique identification code to

unbanded adults upon capture, as individuals can breed within this system over multiple breeding

seasons.

**Insect sampling and local prey availability**

Throughout the study, 2 insect traps were placed on each of the 40 farms (N=80). Traps were spaced ~ 250 m apart along the central portion of each nest box transect. Trap content was collected every two days throughout each breeding season (see Garrett et al. 2021a for details). The processing of insect samples focused on the period between 1 June and 15 July, covering the nestling rearing period of 96% of the monitored breeding attempts. The content of traps included a wide range of insects stemming from many of the surrounding habitat types, and that overlap significantly the diet of nestlings (Laplante 2013, Bellavance et al. 2018). In this system, 74% of the biomass of the food boluses provided to nestlings were Diptera (Bellavance et al. 2018), a pattern shared with other Tree Swallows study systems (Winkler et al. 2020). We thus used the biomass of Diptera within each sample as a proxy of local prey availability. It should be noted that, in a separate analysis of fledging success using this same data set, models incorporating prey availability measures represented by Diptera outperformed ones in which prey availability was represented as all invertebrates (Appendix from Garrett et al. 2021a). Once extracted, Diptera samples were placed in an oven at 60ºC for over 24 hours, ensuring no further change in biomass occurred and were then weighed without delay to the nearest 0.0001 g. The dried biomass of Diptera from these samples (hereafter referred to as Diptera biomass) thus represents an estimate of local prey availability on a specific farm and year over the two days prior to the sample collection date.

**Landscape composition**

Landscape composition (i.e., the relative coverage of habitats composing a given landscape) focused on habitats influential to Tree Swallows and ones composing landscapes throughout our study system [e.g., tree cover, open habitats, and cover of surface water, (Rendell and Robertson

1989, 1990, Winkler et al. 2020)]. The relative cover of habitats was calculated within 500 m of

nest boxes, a spatial scale in which ~80% of food provisioning adult female Tree Swallows

spend their time (Elgin et al. 2020, Garrett et al. 2021b). Tree Swallows are often associated with

open bodies of water and will travel distances greater than 500 m to forage over them (Elgin et

al. 2020, Garrett et al. 2021b). The relative cover of this habitat within 500 m of nest boxes and

in the agricultural contexts covered by this study is low (0.66% $\pm$ 1.07%). Recognizing this

habitat as important to Tree Swallows (Berzins et al. 2021), we calculated its relative cover at 3

km from nest boxes following methods presented in Garrett et al. (2021a). Calculations used

vector layers acquired from the Canadian National Hydro Network (NHN, 2020) and the sf

(Pebesma 2018) and rgeos (Bivand and Rundel 2019) packages in R version 3.6.2 (R Core Team

2019). Characterization within 500 m of nest boxes occurred separately for each farm at the end

of each breeding season to facilitate crop identification. Parcels representing different habitats

and agricultural fields were first delineated using orthophotos (scale 1:40,000) in QGIS (version

3.16) (QGIS 2020), and then characterized *in situ*. We determined which cultures, if any, were in

agricultural fields and then reclassified them into forested, corn and soybean, forage fields

(including hay, other grasses, alfalfa, clover, and pastures), and cereals (principally wheat, and to

a lesser extent, oat (*Avena spp*.) and barley (*Hordeum spp*.)). We then calculated the mean

percent cover of these habitats across the 10 nest boxes on each farm and for each year

independently.

We desired an integrative measure of the percent cover of all habitat groups observed for each

year and farm combination, defined as the landscape context. We therefore used a robust

principal components analysis (PCA) for compositional data on the entire landcover data set to

assign "site scores" to each of the farms during each year (Filzmoser et al. 2009). Site scores

were the year and farm specific values along the first two components of the resulting

compositional PCA. The single PCA was fitted using the robCompositions package (Templ et al.

2011) in R, resulting in the calculation of 440 different landscape contexts (i.e., 40 farms x 11

years). Site scores were assigned to each breeding attempt and insect sample and used in all

subsequent analyses.

The PCA's first component (Comp.1) explained 80.34% of the variance in landcover, correlating

positively with corn and soybean and negatively with both forage fields and forest cover. The

second component (Comp.2) explained 14.69% and correlated negatively with forage fields and

positively with forest cover (Fig. 2). To avoid overly complex models, we included only these

two components to represent the landscape context. Landscapes expressed by minimizing

Comp.1 and Comp.2 values represent ones for which there is an abundance of forage fields and

above average forest cover and are referred to as forage landscapes.

**Weather variables**

We monitored hourly temperature with iButtons (model DS1922L; Embedded Data Systems,

Lawrenceburg, Kentucky, USA) attached to the underside of a single nest box on each farm.

Temperatures were recorded throughout each breeding season and started prior to the spring

arrival of Tree Swallows. We derived daily summaries of the minimum, mean, and maximum

daily temperatures between daytime hours (05:00 to 20:00). We measured total farm-specific

precipitation every two days throughout the breeding season using a single pluviometer placed

on each farm and recorded measurements to the nearest 0.5 ml during each farm visit.

**Statistical analyses**

*Cold-snap days*

Cold snap values were grounded in the functional response of local prey availability to maximum

daily temperature, as we expected insects to require a threshold temperature before they are

active (Williams 1961, Grüebler et al. 2008, Roitberg and Mangel 2016). We calculated the

mean maximum daily temperature between the day of and the day prior to sample collection

(mean maximum daily temperature). We then used a similar approach to that of Winkler et al.

(2013) using generalized additive mixed effects models (GAMMs) to model the relationship

between Diptera biomass and mean maximum daily temperature to determine the temperature at

which the instantaneous rate of biomass change was greatest. The cold snap threshold

temperature was thus calculated as the temperature corresponding to the maximum of the first

derivative of the relationship between Diptera biomass and mean maximum daily temperature.

Peak rate in change was chosen because temperatures around this critical threshold likely result

in dramatic differences in prey availability (Williams 1961, Roitberg and Mangel 2016). We

undertook an extensive modeling exercise to determine the treatment of the functional

relationship between these variables, including evaluating the hypothesis that the critical

threshold temperature varies along the agricultural intensification gradient (full details in

Appendix S1). Modeling used the "bam" function from the mgcv package in R (Wood 2015).

Diptera biomass was modeled with a Gamma distribution with a log link function. The influence

of mean maximum daily temperature was treated as a tensor product smooth, and in order to

evaluate overall trends, we kept the basis dimension of this predictor low (k=10). In all models,

we included the effect of Julian date of sample collection as a tensor product smooth and

precipitation as a linear predictor to account for phenological variations in biomass. We further

included the year and farm as random effects to control for the hierarchical structure of the

sampling design (Pedersen et al. 2019). All correlations and pairwise functional responses are

provided in Appendix S1 (Fig. S1).

*Fledging success*

We limited analyses of fledging success to first breeding attempts (90% of all attempts and

ranging between 80% and 95% across years), as reproductive success varies greatly between first

and second attempts, and second attempts occur near exclusively following the failure of a nest

during laying or incubation (Robertson and Rendell 2001). The date at which insect samples

started to be processed placed a constraint on the breeding data we could use. Most notably, very

early breeding attempts were omitted due to hatching occurring prior to the start of the

processing window (~1% omitted). It should be highlighted that during the 11-year data set

presented, the likelihood of a brood being omitted because its hatching date occurred prior to

insect sampling was irrespective of the breeding season. Fledging success was modeled as the

ratio between the number of fledglings and the number of nestlings (i.e., number of successes

over trials) via generalized linear mixed effects models (GLMMs) with a binomial distribution

and logit link function. The year, farm, and nest box IDs were included as random factors to

account for the hierarchical structure of the data set (i.e., nest box nested within farm nested

within year). Model covariate summaries can be found in Table 1 and were averaged across a 12-

day window post-hatching. This period is when Tree Swallow nestlings become homeothermic,

reach peak body mass, and experience the greatest rate of mortality, thus representing the period

where resource availability is presumably most crucial (McCarty and Winkler 1999, Houle et al.

2020). Cold temperatures and precipitation may act both directly on nestlings through

thermoregulation and indirectly through a reduction in resource availability. To control for

trophic mediated indirect effects of weather on fledging success, we derived estimates of local

prey availability during the nestling rearing period of each breeding attempt. Prey availability

was represented by predictions from generalized additive models (GAMs) in which raw values

of Diptera biomass were regressed against the Julian date of sample collection for each farm and

year separately. Modeled Diptera availability captures its general phenology throughout each

season and avoids biases caused by more punctual or local disturbances (e.g., capturing an insect

swarm). From these predictions, we calculated an estimate of the mean prey availability during

the respective 12-day window of each breeding attempt. GAMs were fitted as a tensor product

smoother and an identical degree of smoothness (k=10).

We took an information theoretic and multimodel approach to assess how weather influenced

fledging success and if its effect varied with landscape context (Burnham and Anderson 2002).

We compared a set of competing models, including a Null model with only random effects.

These data were previously used to determine the combined influence of prey availability and

landscape context within our study area on fledging success, and as a result a most predictive and

parsimonious model had then been determined (Garrett et al. 2021a). We thus compared all

subsequent models to this base model (Base). This model included the brood size, age of the

breeding female [second year (SY) vs. after second year (ASY); (Robertson and Rendell 2001)],

hatching date, percent cover of open water within 3 km of the breeding attempt, the interaction

between site scores (Comp.1 and Comp.2) and the prey availability estimate. To these variables

we also added the number of cold snaps and the mean daily precipitation during the 12 days

following hatching, including the interaction between these variables, as these factors are

identified as influential to Tree Swallows from other study systems (Cox et al. 2019, Shipley et

al. 2020). It should be explicitly stated that our measure of cold snap and precipitation

occurrence cannot measure the consecutiveness or coincidence of these events. Instead, these

measures represent a coarse proxy of the weather that broods experienced during this time

window. All model terms within Base were included in all subsequent models of the candidate

set. Principal interests were in the interactive effects of cold snaps, precipitation, and the gradient

of agricultural intensification on fledging success. We included models with interaction terms

between Comp.1 and the number of cold snaps (Base + Land*Snap), between Comp.1 and the

mean precipitation (Base + Land*Rain), or both of these interaction terms (Base + Land*Snap +

Land*Rain). We further predicted these factors may interact with one another and included a

model with a three-way interaction between these three variables (Base + Land*Snap*Rain). A

summary table of all continuous covariates can be seen in Table 1 and the correlations and

functional relationships between all these variables are provided in Appendix S2 (Fig. S1).

It should further be noted that we explored our choice of temperature threshold and post hatching

window in an extensive modeling exercise (Appendix S3). This modeling endeavor did not

highlight any qualitative differences from the following results. We therefore continued to utilize

the methodology presented thus far in an attempt to build models based on sound biological

variables influential to the behavior and physiology of Tree Swallows (Burnham and Anderson

2002, Harrison et al. 2017).

In models of fledging success, the effects of key individual variables, including interactions,

were estimated via multi-model inference whereby predictions were calculated by model-

averaging with shrinkage and shown with their 95% unconditional confidence intervals

(Burnham and Anderson 2002). All quantitative covariates were Z-transformed, $R$-squares

calculated following Nakagawa and Schielzeth (2013), and analyses performed in R using the

glmmTMB (Brooks et al. 2017) and AICcmodavg (Mazerolle 2020) packages. Model validation,

including evaluation of normally distributed residuals (simulated), heteroscedasticity, and

collinearity checks with variance inflation factors, followed Zuur et al. (2009) and used the

DHARMa package (Hartig 2020).

**Results**

**Effects of temperature on local prey availability**

We collected and processed 15,916 insect samples from 8,614 farm visits. Mean Diptera biomass

($\pm$ SD) was $0.030 \pm 0.044$ g (per trap), ranging between 0.019 g and 0.037 g across years.

Overall mean maximum daily temperature ($\pm$ SD) was 25.7°C $\pm$ 4.1 ranging between 24.1°C and

26.6°C across years. The greatest instantaneous rate of change in Diptera biomass occurred at

18.3°C (Fig. 3, A and B). Around this critical threshold, a two degree decrease or increase in

maximum daily temperature led to a 13.6% drop or 15.0% rise in Diptera biomass, respectively.

Based on this threshold, 8.2% of farm visits followed at least one cold snap (Appendix S2: Fig.

S2). A second, less steep maximum of the first derivative was noted at 29.2°C resulting in a

marked decrease in Diptera biomass (Appendix S1: Fig. S2). We further observed that the daily

likelihood to observe a cold snap has increased slightly over the course of this data set (Appendix

S4: Table S1 and Fig. S1), and that they were more likely to occur in less agro-intensive

landscapes (Appendix S4: Table S2 and Fig. S2).

**Effects of inclement weather on fledging success**

We monitored the breeding activity of 1,897 broods across the 40 farms and 11 breeding

seasons. Overall mean ($\pm$ SD) fledging success (i.e., ratio of the brood that fledged) was $0.74 \pm$

0.38, varying between 0.63 and 0.88 among years. We observed that the average hatching date

advanced by ~1.1 days over the study period (Appendix S4: Table S3 and Fig. S3). Overall mean

number of cold snap days ($\pm$ SD) experienced by a brood during its first 12 days was $0.9 \pm 1.1$

days and varied between 0.2 and 2.3 days among years (Fig. 4). At least 52% of broods

experienced at least one cold snap day throughout the duration of the study; 25%, 11%, and 5%

of broods experienced at least 2, 3, and 4 cold snap days, respectively. Mean daily precipitation

during the first 12 days ($\pm$ SD) was 7.4 $\pm$ 4.9 ml, ranging between 2.4 and 13.6 ml across years.

The correlation between the number of cold snap days and mean daily precipitation experienced

by a brood was relatively low ($r$=0.18). For a more detailed look at the appearance of cold snaps

and precipitation along the agricultural gradient of this data set see Appendix S2 (Fig. S3).

Models incorporating an interaction between Comp.1 and the number of cold snap days

(combined AICc $w$ > 0.99) were substantially supported over Base ($\Delta$AICc > 16.15, Appendix

S2: Table S1). Fledging success decreased with the number of cold snap days (Fig. 5). Fledging

success was on average 36.2% lower when broods experienced four cold snap days and average

precipitation in their first 12 days instead of none. Increasing precipitation had a negative effect

on fledging success, especially when broods were reared during windows containing both several

cold snap days and high average daily amounts of precipitation (Appendix S2: Fig. S4). Broods

subjected to four cold snap days versus none showed up to a 46.0% reduction in fledging success

if exposed to the 95[th] percentile of observed values for mean daily precipitation. The effects of

cold snap days varied with landscape context (Fig. 5). For broods experiencing increased cold

snap days, higher levels of agro-intensive cover was associated with further reductions in

fledging success. When broods incurred the average number of cold snap days and mean daily

precipitation there was no observable difference in fledging success along the agricultural

gradient. However, this pattern changed to fledging success being 43.6% lower in agro-intensive

landscapes when broods experienced four cold snap days and average precipitation (Appendix

S2: Fig. S5). The synergistic negative effect of the number of cold snap days and high amount of

precipitation on fledging success was, however, far greater in agro-intensive landscapes than in

forage landscapes. In the absence of precipitation, the negative effect of increasing cold snap

days was substantial only in agro-intensive landscapes, but similarly detrimental over the entire

intensification gradient under high precipitation levels (Fig. 6). Lastly, fledging success

increased with local prey availability and percent cover of surface water within 3 km, it

decreased with increasing hatching day and brood size, and was lower for SY than ASY females

(Fig. 5).

**Discussion**

An outcome of climate change is the increased frequency and duration of weather events

potentially reducing the availability of thermally sensitive prey resources (Rahmstorf and

Coumou 2011, Wuebbles et al. 2014). These reductions, when occurring during critical life

history stages (e.g., nestling periods), reduce the fitness of avian aerial insectivores (Pipoly et al.

2013, Moreno et al. 2015, Arbeiter et al. 2016). Though climate change influences a vast array of

species and their interactions, another large-scale continued threat to biodiversity is

anthropogenic reductions in habitat amounts and quality, notably through forestry and agriculture

(Maxwell et al. 2016). We contended that the use of certain habitats may pose a risk to an

animal's ability to breed or survive through periods of reduced food availability and/or poor

weather. We provide evidence that the negative impact of weather conditions expected from

climate change on the annual breeding success of an aerial insectivore are exacerbated in habitats

created by agricultural intensification.

The relationship between prey availability and mean maximum temperature, between the day of

and day prior to sample collection, revealed the rate of change in prey availability was

maximized at a temperature (18.3°C) similar to that deemed critical in past works (18.5°C;

Winkler et al. 2013, Cox et al. 2019, Shipley et al. 2020). Such similarity may highlight a

generality to temperatures impactful to Tree Swallows and their prey. We further found that this

temperature threshold did not vary across the large-scale gradient of agricultural intensification

studied here.

In accordance with previous studies, the appearance and frequency of cold snaps during the 12-

day post hatching period of each brood led to reduced fledging success of Tree Swallows.  This

effect increased when multiple cold snaps and increased rain occurred during the 12-day post-

hatching period. However, the overall negative effect of experiencing increasing precipitation

appeared to be much weaker than increasing number of cold snaps. The impact of cold snaps

likely acts on fledging success both through variation in resource availability, and by increased

thermoregulatory demand (Sauve et al. 2021). While precipitation, because nests are protected

within nest boxes, likely only impacts nestling survival through the former mechanism. This

result is further evidenced by the role of the agricultural gradient in this relationship. Most

notably, the negative effect of poor weather was stronger for breeding attempts in more agro-

intensive landscapes. Such a result suggests features associated to these landscapes exacerbate

the consequences of broods experiencing poor weather.

It should be mentioned that under very dry and warm conditions, fledging success appears to be

slightly better in the most agro-intensive landscapes (7 % increase along the Comp.1 axis in Fig.

1 with 0 ml of rain). This relationship may highlight that agricultural landcover is only influential

to fledging success during certain weather conditions as highlighted by the significant support of

models incorporating interaction terms representing such a relationship. There is no clear

explanation for this increase in fledging success in agro-intensive areas under these conditions.

However, one potential hypothesis is that hay harvesting, which causes insect mortality and

which timing depends on weather conditions, may affect the differences in quality between

agricultural habitats (Humbert et al. 2009). However, food provisioning rates appear to be the

lowest within the most agro-intensive parts of this study system (Garrett et al. 2021b). If the

lowered food provisioning in these conditions are indicative of greater costs for food

provisioning parents, then these costs may carry over and influence the survival and lifetime

reproductive success of these parents. Therefore, under optimal weather conditions, fledging

success may be slightly greater in more agro-intensive landscapes, and yet negatively impact the

survival of adults when compared to forage landscapes. It should be stated that in our data set,

broods experiencing 0 cold snap days and mean precipitation predominantly occurred within

more agro-intensive landscapes, and that the daily likelihood to experience a cold snap was

greatest in more forage landscapes. This is likely due to the elevational change occurring along

the agricultural gradient. Taken together this may indicate that we underestimate the

consequences of rearing young within agro-intensive landscapes, as environmental conditions

from this data set appeared spatially worse for those broods in forage focused landscapes.

We propose two non-mutually exclusive mechanisms for the finding that broods experiencing

inclement weather succumbed to greater mortality in more agro-intensive landscapes compared

to forage focused landscapes: the structural characteristics of agro-intensive landscapes and the

likelihood of exposure to agrochemical agents. Successfully acquiring food during periods of

poor weather potentially result in greater costs for food provisioning parents, inasmuch as the

increased foraging effort needed to overcome the reduced availability of prey likely results in

reduced body condition (Hinsley 2000, Evens et al. 2018). Parents may possibly forgo foraging

during poor weather periods in order to brood nestlings and provide food at greater rates once

poor weather has subsided (Cox et al. 2019). Given the homogeneity of intensively-managed

agricultural landscapes, prey patches may be fewer or farther apart therein (Benton et al. 2003,

Fahrig et al. 2011). Furthermore, agro-intensive landscapes tend to be denuded of structural

habitats [e.g., woodlots and hedgerows; Fig. 1; (Benton et al. 2003, Fahrig et al. 2011)]. In

addition to being prey habitat, these features may block wind, and their removal may limit prey

aggregation (Evans et al. 2003, Grüebler et al. 2008) and facilitate heat dissipation of nests and

adults (Heenan 2013, Tapper et al. 2020a). Therefore, the impacts of experiencing periods of

inclement weather in forage dominated landscapes may be less harsh than in more agro-intensive

landscapes.

Evidence suggests that agrochemical agents such as herbicides (e.g., glyphosate, atrazine, S-

metolachlor, etc.), neonicotinoid insecticides (e.g., clothianidin, thiacloprid, thiametoxam, etc.),

and fertilizers can alter avian physiology in ways that potentially hinder an individual's capacity

to respond to poor weather (Mayne et al. 2005, Mineau and Palmer 2013, Gibbons et al. 2014,

Lopez-Antia et al. 2015), most notably through a reduction in thermoregulatory capacity (Grue et

al. 1997). These same agents may also influence the ability for food provisioning parents to

optimize foraging strategies by inducing anorexia and impairing both locomotor and navigational

function (Eng et al. 2019). These factors illustrate that within more agro-intensive landscapes,

food provisioning parents may present longer foraging bouts that potentially come at a cost to

their own body condition, while also reducing the time spent brooding nestlings (Garrett et al.

2021b).

Our conclusions, as well as predicted impacts of climate change on aerial insectivores, assume

food provisioning adults are unable or less capable of foraging during periods of poor weather.

However, fine scale movements of small passerines are becoming less difficult to observe (Elgin

et al. 2020, Garrett et al. 2021b). Understanding such movements may further elucidate the

combined roles of breeding habitat quality and predictions of climate change on animal behavior.

For instance, the increased food provisioning of nestlings by adults, following periods of poor

weather (Cox et al. 2019), may be facilitated by foraging in more profitable habitat patches

(Elgin et al. 2020, Geary et al. 2020, Garrett et al. 2021b). In the context of climate change, the

combined roles of poor weather and landscape structure on foraging responses is still unknown.

Validation of these assumptions will be imperative as the effectiveness of food provisioning,

during periods of poor weather, likely varies across a gradient of breeding landscape quality.

Finally, a focus here was on the effects of cold snaps, yet another growing concern of climate

change are periods of high ambient temperature [i.e., heat waves; (Perkins-Kirkpatrick and

Lewis 2020)]. The energetic expenditure of food provisioning Tree Swallows is potentially

constrained by an upper thermal limit (Tapper et al. 2020b). Moreover, prey availability

decreased sharply above 30ºC (Appendix S1: Fig. S2) in our study system. Food provisioning

parents may thus be hindered and the effects of breeding within more agro-intensive landscapes

may further be exacerbated during heat waves. Future work should focus on evaluating this

hypothesis, as reduced food provisioning potentially results in reduced nestling growth and

survival (Cox et al. 2019, Garrett et al. 2021b).

Finally, our focus on the relationship between a threshold temperature and nestling survival

utilized a single temperature and assumed that experiencing days falling below this value

impacts both the amount of food resources coming into nest boxes and the thermoregulatory

capacity of nestlings. However, our threshold temperature of 18.3°C occurs multiple times

during a breeding season and thus selective processes would likely favor more robust capacity of

nestlings to survive such temperatures. Yet, we found strong support that nestling mortality

increased when broods experienced increasing number of cold snaps. We propose that there is

likely two threshold temperatures (Appendix S3: Fig. S1), one influential to the availability of

prey resources and another influential to nestling's thermoregulator capacity (Stodola et al. 2010,

Pérez et al. 2016, de Zwaan et al. 2019), as proposed by Winkler et al (2013). Future work needs

to explicitly evaluate this hypothesis across similar gradients of agricultural intensification.

**Conclusion and applications**

Aerial insectivore populations are declining in several parts of their North American and

European breeding grounds (Spiller and Dettmers 2019, Rosenberg et al. 2019, Bowler et al.

2019). Here we provided evidence that two of the primary hypotheses explaining these

reductions, especially for aerial insectivores, may interact to further intensify declines. We

propose that future work investigating insectivorous bird population declines should explicitly

focus not only on climate change or anthropogenic reductions or alterations of breeding habitats,

but also on how these factors may interact with one another. Furthermore, understanding the

various mechanisms that lead to increased nestling mortality during inclement weather or within

more agro-intensive landscapes will be critical as they potentially identify drivers under

anthropogenic control.

**Acknowledgments**

We are indebted to the farm owners who kindly accepted to partake in our long-term study since

2004. Sincere thanks to the many graduate students, field and lab assistants who helped collect,

enter, and proof data throughout the years, notably with respect to insect sample processing. This

work was conducted under the approval of the animal care committee of the Université de

Sherbrooke and was financially supported by Natural Sciences and Engineering Research

Council of Canada (NSERC) discovery grants to FP, DG and MB, two team research grants from

the Fonds de recherche du Québec—Nature et technologies (FRQNT) to FP, DG and MB, by the

**Literature cited**

Agriculture and Agri-Food Canada (AAFC). 2018. Annual Crop Inventory.

https://open.canada.ca/data/en/dataset/.

Arbeiter, S., M. Schulze, P. Tamm, and S. Hahn. 2016. Strong cascading effect of weather

conditions on prey availability and annual breeding performance in European bee-eaters

Merops apiaster. Journal of Ornithology 157:155–163.

Bélisle, M. 2005. Measuring landscape connectivity: the challenge of behavioral landscape

ecology. Ecology 86:1988–1995.

Bellavance, V., M. Bélisle, J. Savage, F. Pelletier, and D. Garant. 2018. Influence of agricultural

intensification on prey availability and nestling diet in Tree Swallows (Tachycineta bicolor).

Canadian Journal of Zoology 96:1053–1065.

Benton, T. G., J. A. Vickery, and J. D. Wilson. 2003. Farmland biodiversity: is habitat

heterogeneity the key? Trends in Ecology & Evolution 18:182–188.

Berzins, L. L., A. K. Mazer, C. A. Morrissey, and R. G. Clark. 2021. Pre-fledging quality and

recruitment in an aerial insectivore reflect dynamics of insects, wetlands and climate.

Oecologia.

Bivand, R. S., and C. Rundel. 2019. Package 'rgeos' - Interface to Geometry Engine - Open

Source ('GEOS').

Both, C., C. A. M. Van Turnhout, R. G. Bijlsma, H. Siepel, A. J. Van Strien, and R. P. B.

Foppen. 2010. Avian population consequences of climate change are most severe for long-

distance migrants in seasonal habitats. Proceedings of the Royal Society B: Biological

Sciences 277:1259–1266.

Bourret, A., M. Bélisle, F. Pelletier, and D. Garant. 2015. Multidimensional environmental

influences on timing of breeding in a tree swallow population facing climate change.

Evolutionary Applications 8:933–944.

Bowler, D. E., H. Heldbjerg, A. D. Fox, M. Jong, and K. Böhning☐Gaese. 2019. Long☐term

declines of European insectivorous bird populations and potential causes. Conservation

Biology 33:1120–1130.

Brooks, M. E., K. Kristensen, K. J. van Benthem, A. Magnusson, C. W. Berg, A. Nielsen, H. J.

Skaug, M. Mächler, and B. M. Bolker. 2017. glmmTMB balances speed and flexibility

among packages for zero-inflated generalized linear mixed modeling. R Journal.

Burnham, K. P., and D. R. Anderson. 2002. Model Selection and Multimodel Inference. second.

Springer.

Canada, N. R. 2020. National Hydro Network - NHN - GeoBase Series.

https://www.nrcan.gc.ca/science-and-data/science-and-research/earth-

sciences/geography/topographic-information/geobase-surface-water-program-

geeau/national-hydrographic-network/21361.

Cox, A. R., R. J. Robertson, Á. Z. Lendvai, K. Everitt, and F. Bonier. 2019. Rainy springs linked

to poor nestling growth in a declining avian aerial insectivore (Tachycineta bicolor).

Proceedings of the Royal Society B: Biological Sciences 286:20190018.

Crick, H. Q. P. 2004. The impact of climate change on birds. Ibis 146:48–56.

DiBartolomeis, M., S. Kegley, P. Mineau, R. Radford, and K. Klein. 2019. An assessment of

acute insecticide toxicity loading (AITL) of chemical pesticides used on agricultural land in

the United States. PLOS ONE 14:e0220029.

Dunn, P. O., and D. W. Winkler. 1999. Climate change has affected the breeding date of tree

swallows throughout North America. Proceedings of the Royal Society of London. Series

B: Biological Sciences 266:2487–2490.

Elgin, A. S., R. G. Clark, and C. A. Morrissey. 2020. Tree Swallow selection for wetlands in

agricultural landscapes predicted by central-place foraging theory. The Condor 122:1–12.

Eng, M. L., B. J. M. Stutchbury, and C. A. Morrissey. 2019. A neonicotinoid insecticide reduces

fueling and delays migration in songbirds. Science 365:1177–1180.

Evans, K. L., R. B. Bradbury, and J. D. Wilson. 2003. Selection of hedgerows by Swallows

Hirundo rustica foraging on farmland: the influence of local habitat and weather. Bird Study

50:8–14.

Evens, R., N. Beenaerts, T. Neyens, N. Witters, K. Smeets, and T. Artois. 2018. Proximity of

breeding and foraging areas affects foraging effort of a crepuscular, insectivorous bird.

Scientific Reports 8:1–11.

Fahrig, L., J. Baudry, L. Brotons, F. G. Burel, T. O. Crist, R. J. Fuller, C. Sirami, G. M.

Siriwardena, and J.-L. Martin. 2011. Functional landscape heterogeneity and animal

biodiversity in agricultural landscapes. Ecology Letters 14:101–112.

Filzmoser, P., K. Hron, and C. Reimann. 2009. Principal component analysis for compositional

data with outliers. Environmetrics 20:621–632.

Garrett, D. R., F. Pelletier, D. Garant, and M. Bélisle. 2021a. Combined influence of food

availability and agricultural intensification on a declining aerial insectivore.

bioRxiv:10.1101/2021.02.02.427782.

Garrett, D. R., F. Pelletier, D. Garant, and M. Bélisle. 2021b. Negative effects of agricultural

intensification on energetic gain rates of a declining aerial insectivore.

bioRxiv:10.1101/2021.03.24.436831.

Geary, B., P. L. Leberg, K. M. Purcell, S. T. Walter, and J. Karubian. 2020. Breeding Brown

Pelicans Improve Foraging Performance as Energetic Needs Rise. Scientific Reports 10:1–

9.

Gibbons, D., C. A. Morrissey, and P. Mineau. 2014. A review of the direct and indirect effects of

neonicotinoids and fipronil on vertebrate wildlife. Environmental Science and Pollution

Research 22:103–118.

Giroux, I. 2019. Présence de pesticides dans l'eau au Québec : Portrait et tendances dans les

zones de maïs et de soya – 2015 à 2017, Québec, ministère de l'Environnement et de la

Lutte contre les changements climatiques, Direction générale du suivi de l'état de l'environ.

Grue, C. E., P. L. Gibert, and M. E. Seeley. 1997. Neurophysiological and Behavioral Changes

in Non-Target Wildlife Exposed to Organophosphate and Carbamate Pesticides:

Thermoregulation, Food Consumption, and Reproduction. American Zoologist 37:369–388.

Grüebler, M. U., M. Morand, and B. Naef-Daenzer. 2008. A predictive model of the density of

airborne insects in agricultural environments. Agriculture, Ecosystems & Environment

123:75–80.

Harrison, X. A., L. Donaldson, M. E. Correa-Cano, J. Evans, D. N. Fisher, C. Goodwin, B.

Robinson, D. J. Hodgson, and R. Inger. 2017. Best practice in mixed effects modelling and

multi-model inference in ecology. PeerJ Preprints 5:e3113v1-48.

Hartig, F. 2020. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)
Regression Models. R package version 0.2.0.

Hartmann, D. L., A. M. G. Klein Tank, M. Rusticucci, L. V. Alexander, S. Brönnimann, Y. A.-
R. Charabi, F. J. Dentener, E. J. Dlugokencky, D. R. Easterling, A. Kaplan, B. J. Soden, P.
W. Thorne, M. Wild, and P. Zhai. 2013. Observations: Atmosphere and Surface. Pages
159–254 *in* T. F. Stocker, D. Qin, G.-K.Plattner, M. Tignor, S. K. Allen, J. Boschung, A.
Nauels, Y. Xia, V. Bex, and P. M. Midgley, editors. Climate Change 2013 the Physical
Science Basis: Working Group I Contribution to the Fifth Assessment Report of the
Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge,
United Kingdom and New York, NY, USA.

Heenan, C. B. 2013. An Overview of the Factors Influencing the Morphology and Thermal
Properties of Avian Nests. Avian Biology Research 6:104–118.

Hinsley, S. A. 2000. The costs of multiple patch use by birds. Landscape Ecology 15:765–775.

Houle, C., F. Pelletier, M. Bélisle, and D. Garant. 2020. Impacts of environmental heterogeneity
on natural selection in a wild bird population. Evolution 74:1142–1154.

Humbert, J.-Y., J. Ghazoul, and T. Walter. 2009. Meadow harvesting techniques and their
impacts on field fauna. Agriculture, Ecosystems & Environment 130:1–8.

Jobin, B., J. Beaulieu, M. Grenier, L. Bélanger, C. Maisonneuve, D. Bordage, and B. Filion.
2003. Landscape changes and ecological studies in agricultural regions, Québec, Canada.
Landscape Ecology 18:575–590.

Kukal, M. S., and S. Irmak. 2018. U.S. Agro-Climate in 20th Century: Growing Degree Days,

First and Last Frost, Growing Season Length, and Impacts on Crop Yields. Scientific

Reports 8:6977.

Laplante, L. 2013. Influence de l'hétérogénéité et de la composition du paysage agricole sur la

communauté de diptères schizophores du sud du québec (canada). Université de

Sherbrooke.

Lopez-Antia, A., M. E. Ortiz-Santaliestra, F. Mougeot, and R. Mateo. 2015. Imidacloprid-treated

seed ingestion has lethal effect on adult partridges and reduces both breeding investment

and offspring immunity. Environmental Research 136:97–107.

Malaj, E., L. Freistadt, and C. A. Morrissey. 2020. Spatio-Temporal Patterns of Crops and

Agrochemicals in Canada Over 35 Years. Frontiers in Environmental Science 8:1–12.

Marcelino, J., J. P. Silva, J. Gameiro, A. Silva, F. C. Rego, F. Moreira, and I. Catry. 2020.

Extreme events are more likely to affect the breeding success of lesser kestrels than average

climate change. Scientific Reports 10:7207.

Marrot, P., D. Garant, and A. Charmantier. 2017. Multiple extreme climatic events strengthen

selection for earlier breeding in a wild passerine. Philosophical Transactions of the Royal

Society B: Biological Sciences 372:20160372.

Maxwell, S. L., R. A. Fuller, T. M. Brooks, and J. E. M. Watson. 2016. Biodiversity: The

ravages of guns, nets and bulldozers. Nature 536:143–145.

Mayne, G. J., C. A. Bishop, P. A. Martin, H. J. Boermans, and B. Hunter. 2005. Thyroid

Function in Nestling Tree Swallows and Eastern Bluebirds Exposed to Non-Persistent

Pesticides and p, p-DDE in Apple Orchards of Southern Ontario, Canada. Ecotoxicology

14:381–396.

Mazerolle, M. J. 2020. AICcmodavg: model selection and multimodel inference based on

(Q)AIC(c).

McCarty, J. P., and D. W. Winkler. 1999. Relative importance off environmental variables in

determining the growth off nestling Tree Swallows Tachycineta bicolor. Ibis 141:286–296.

MDDELCC. 2015. Ministère du Développement durable, de l'Environnement et de la Lutte

contre les changements climatiques. Stratégie québécoise sur les pesticides 2015-2018.

http://www.mddelcc.gouv.qc.ca/.

Mineau, P., and C. Palmer. 2013. The impact of the nation's most widely used insecticides on

birds. Cambridge University Press.

Møller, A. P., E. Flensted-Jensen, and W. Mardal. 2006. Rapidly advancing laying date in a

seabird and the changing advantage of early reproduction. Journal of Animal Ecology

75:657–665.

Moreno, J., S. González-Braojos, and R. Ruiz-de-Castañeda. 2015. A spring cold snap is

followed by an extreme reproductive failure event in a mountain population of Pied

Flycatchers Ficedula hypoleuca. Bird Study 62:466–473.

Nakagawa, S., and H. Schielzeth. 2013. A general and simple method for obtaining R 2 from

generalized linear mixed-effects models. Methods in Ecology and Evolution 4:133–142.

Nocera, J. J., J. M. Blais, D. V. Beresford, L. K. Finity, C. Grooms, L. E. Kimpe, K. Kyser, N.

Michelutti, M. W. Reudink, and J. P. Smol. 2012. Historical pesticide applications

coincided with an altered diet of aerially foraging insectivorous chimney swifts.

Proceedings of the Royal Society B: Biological Sciences 279:3114–3120.

Olsson, O., J. S. Brown, and K. L. Helf. 2008. A guide to central place effects in foraging.
Theoretical Population Biology 74:22–33.

Pebesma, E. J. 2018. Simple Features for R: Standardized Support for Spatial Vector Data. The
R Journal 10:439.

Pedersen, E. J., D. L. Miller, G. L. Simpson, and N. Ross. 2019. Hierarchical generalized
additive models in ecology: an introduction with mgcv. PeerJ 7:e6876.

Pérez, J. H., J. S. Krause, H. E. Chmura, S. Bowman, M. McGuigan, A. L. Asmus, S. L. Meddle,
K. E. Hunt, L. Gough, N. T. Boelman, and J. C. Wingfield. 2016. Nestling growth rates in
relation to food abundance and weather in the Arctic. The Auk 133:261–272.

Perkins-Kirkpatrick, S. E., and S. C. Lewis. 2020. Increasing trends in regional heatwaves.
Nature Communications 11:3357.

Pipoly, I., V. Bókony, G. Seress, K. Szabó, and A. Liker. 2013. Effects of Extreme Weather on
Reproductive Success in a Temperate-Breeding Songbird. PLoS ONE 8:e80033.

Pisa, L. W., V. Amaral-Rogers, L. P. Belzunces, J. M. Bonmatin, C. A. Downs, D. Goulson, D.
P. Kreutzweiser, C. Krupke, M. Liess, M. McField, C. A. Morrissey, D. A. Noome, J.
Settele, N. Simon-Delso, J. D. Stark, J. P. Van der Sluijs, H. Van Dyck, and M. Wiemers.
2015. Effects of neonicotinoids and fipronil on non-target invertebrates. Environmental
Science and Pollution Research 22:68–102.

van de Pol, M., S. Jenouvrier, J. H. C. Cornelissen, and M. E. Visser. 2017. Behavioural,
ecological and evolutionary responses to extreme climatic events: challenges and directions.

Philosophical Transactions of the Royal Society B: Biological Sciences 372:20160134.

Porlier, M., A. Charmantier, P. Bourgault, P. Perret, J. Blondel, and D. Garant. 2012. Variation

in phenotypic plasticity and selection patterns in blue tit breeding time: between- and

within-population comparisons. Journal of Animal Ecology 81:1041–1051.

Poulin, B., G. Lefebvre, and L. Paz. 2010. Red flag for green spray: adverse trophic effects of Bti

on breeding birds. Journal of Applied Ecology 47:884–889.

QGIS. 2020. QGIS Geographic Information System. QGIS Association. http://www.qgis.org.

R Core Team. 2019. R: A language and environment for statistical computing.

Rahmstorf, S., and D. Coumou. 2011. Increase of extreme events in a warming world.

Proceedings of the National Academy of Sciences 108:17905–17909.

Rainho, A., and J. M. Palmeirim. 2011. The Importance of Distance to Resources in the Spatial

Modelling of Bat Foraging Habitat. PLoS ONE 6:e19227.

Rendell, W. B., and R. J. Robertson. 1989. Nest-Site Characteristics, Reproductive Success and

Cavity Availability for Tree Swallows Breeding in Natural Cavities. The Condor 91:875.

Rendell, W. B., and R. J. Robertson. 1990. Influence of Forest Edge on Nest-Site Selection by

Tree Swallows. Wilson Journal of Ornithology 102:634–644.

Robertson, R. J., and W. B. Rendell. 2001. A long-term study of reproductive performance in

tree swallows: the influence of age and senescence on output. Journal of Animal Ecology

70:1014–1031.

Roitberg, B. D., and M. Mangel. 2016. Cold snaps, heatwaves, and arthropod growth. Ecological

Entomology 41:653–659.

Rosenberg, K. V., A. M. Dokter, P. J. Blancher, J. R. Sauer, A. C. Smith, P. A. Smith, J. C.

Stanton, A. Panjabi, L. Helft, M. Parr, and P. P. Marra. 2019. Decline of the North

American avifauna. Science 366:120–124.

Ruiz, J., and G. Domon. 2009. Analysis of landscape pattern change trajectories within areas of

intensive agricultural use: case study in a watershed of southern Québec, Canada.

Landscape Ecology 24:419–432.

Sauve, D., V. L. Friesen, and A. Charmantier. 2021. The Effects of Weather on Avian Growth

and Implications for Adaptation to Climate Change. Frontiers in Ecology and Evolution

9:1–20.

Sheldon, B. C., L. E. B. Kruuk, and J. Merila. 2003. Natural selection and inheritance of

breeding time and clutch size in the collared flycatcher. Evolution 57:406–420.

Shipley, J. R., C. W. Twining, C. C. Taff, M. N. Vitousek, A. Flack, and D. W. Winkler. 2020.

Birds advancing lay dates with warming springs face greater risk of chick mortality.

Proceedings of the National Academy of Sciences 117:25590–25594.

Shutler, D., D. J. T. Hussell, D. R. Norris, D. W. Winkler, R. J. Robertson, F. Bonier, W. B.

Rendell, M. Bélisle, R. G. Clark, R. D. Dawson, N. T. Wheelwright, M. P. Lombardo, P. A.

Thorpe, M. a. Truan, R. Walsh, M. L. Leonard, A. G. Horn, C. M. Vleck, D. Vleck, A. P.

Rose, L. a. Whittingham, P. O. Dunn, K. a. Hobson, and M. T. Stanback. 2012.

Spatiotemporal Patterns in Nest Box Occupancy by Tree Swallows Across North America.

Avian Conservation and Ecology 7:art3.

Sigouin, A., M. Bélisle, D. Garant, and F. Pelletier. 2021. Agricultural pesticides and

ectoparasites: potential combined effects on the physiology of a declining aerial insectivore.

Conservation Physiology 9:1–16.

Skwarska, J., A. Podstawczyńska, M. Bańbura, M. Glądalski, A. Kaliński, M. Markowski, J.

Wawrzyniak, P. Zieliński, and J. Bańbura. 2021. Effects of ambient temperature during the

nestling stage on a stress indicator in nestling pied flycatchers Ficedula hypoleuca.

International Journal of Biometeorology.

Spiller, K. J., and R. Dettmers. 2019. Evidence for multiple drivers of aerial insectivore declines

in North America. The Condor 121:1–13.

Staggenborg, J., H. M. Schaefer, C. Stange, B. Naef-Daenzer, and M. U. Grüebler. 2017. Time

and travelling costs during chick-rearing in relation to habitat quality in Little Owls Athene

noctua. Ibis 159:519–531.

Stanton, R., C. A. Morrissey, and R. G. Clark. 2016. Tree Swallow (Tachycineta bicolor)

foraging responses to agricultural land use and abundance of insect prey. Canadian Journal

of Zoology 94:637–642.

Stanton, R., C. A. Morrissey, and R. G. Clark. 2018. Analysis of trends and agricultural drivers

of farmland bird declines in North America: A review. Agriculture, Ecosystems &

Environment 254:244–254.

Stodola, K. W., D. A. Buehler, D. H. Kim, K. E. Franzreb, and E. T. Linder. 2010. Biotic and

Abiotic Factors Governing Nestling-period Length in the Ovenbird (Seiurus aurocapilla).

The Auk 127:204–211.

Tapper, S., J. J. Nocera, and G. Burness. 2020a. Heat dissipation capacity influences

reproductive performance in an aerial insectivore. The Journal of Experimental Biology

223:jeb222232.

Tapper, S., J. J. Nocera, and G. Burness. 2020b. Experimental evidence that hyperthermia limits

offspring provisioning in a temperate-breeding bird. Royal Society Open Science

7:10.1098/rsos.201589.

Templ, M., K. Hron, and P. Filzmoser. 2011. robCompositions: An R-package for Robust

Statistical Analysis of Compositional Data. Pages 341–355 Compositional Data Analysis.

John Wiley & Sons, Ltd.

Townsend, A. K., T. S. Sillett, N. K. Lany, S. A. Kaiser, N. L. Rodenhouse, M. S. Webster, and

R. T. Holmes. 2013. Warm Springs, Early Lay Dates, and Double Brooding in a North

American Migratory Songbird, the Black-Throated Blue Warbler. PLoS ONE 8:e59467.

Tuero, D. T., A. E. Jahn, M. S. Husak, D. V. Roeder, D. A. Masson, F. M. Pucheta, T. J.

Michels, A. Quickle, J. Q. Vidoz, M. Domínguez, and J. C. Reboreda. 2018. Ecological

determinants of Tyrannus flycatcher nestling growth at north- and south-temperate latitudes.

The Auk 135:439–448.

Ummenhofer, C. C., and G. A. Meehl. 2017. Extreme weather and climate events with ecological

relevance: a review. Philosophical Transactions of the Royal Society B: Biological Sciences

372:20160135.

Visser, M. E., and P. Gienapp. 2019. Evolutionary and demographic consequences of

phenological mismatches. Nature Ecology & Evolution 3:879–885.

Wagner, D. L. 2020. Insect Declines in the Anthropocene. Annual Review of Entomology

65:457–480.

Williams, C. 1961. Studies in the effect of weather conditions on the activity and abundance of

insect populations. Philosophical Transactions of the Royal Society of London. Series B,

Biological Sciences 244:331–378.

Winkler, D. W., K. K. Hallinger, D. R. Ardia, R. J. Robertson, B. J. M. Stutchbury, and R. R.

Cohen. 2020. Tree Swallow (Tachycineta bicolor). Page *in* A. F. Poole, editor. Birds of the

World. Cornell Lab of Ornithology.

Winkler, D. W., M. K. Luo, and E. Rakhimberdiev. 2013. Temperature effects on food supply

and chick mortality in tree swallows (Tachycineta bicolor). Oecologia 173:129–138.

Wood, S. N. 2015. Package 'mgcv.'

Wuebbles, D. J., K. Kunkel, M. Wehner, and Z. Zobel. 2014. Severe Weather in United States

Under a Changing Climate. Eos, Transactions American Geophysical Union 95:149–150.

Zuur, A. F., E. N. Ieno, N. Walker, A. A. Saveliev, and G. M. Smith. 2009. Mixed effects models

and extensions in ecology with R. Page Smart Society. Springer New York.

de Zwaan, D. R., A. F. Camfield, E. C. MacDonald, and K. Martin. 2019. Variation in offspring

development is driven more by weather and maternal condition than predation risk.

Functional Ecology 33:447–456.

	**Tables and Figures**

771	**Table 1**: Mean (±SD) of covariates used to model fledging success of 1,897 Tree Swallow

772	broods between 2006 and 2016. Covariates were averaged over a 12-day period after hatching.

773	Variable groups identify which covariates were present within models found in Appendix S2:

774	Table S1.

| Covariate | Acronym | Unit | Mean ± SD | Variable group |
|---|---|---|---|---|
| Site score Comp.1 | Comp.1 | - | 0.20 ± (1.80) | Base/Land |
| Site score Comp.2 | Comp.2 | - | -0.18 ± (0.91) | Base/Land |
| Cold snap days | CSD | (day) | 0.93 ± (1.14) | Base/Snap |
| Precipitation | RN | (ml) | 7.44 ± (4.92) | Base/Rain |
| Open water | WT | (%) | 0.01 ± (0.01) | Base |
| Prey availability | PA | (g) | 0.03 ± (0.02) | Base |
| Hatching day | HD | (Julian day) | 159.91 ± (5.76) | Base |
| Brood size | NN | (nestlings) | 4.83 ± (1.27) | Base |

 **Figure legends**

**Fig. 1**: Location of the 40 farms (white dots) monitored across a gradient of agricultural

intensification within southern Québec, Canada. Underlying image represents the agricultural

gradient derived from the Annual crop inventory of 2013 (AAFC, 2018). Each pixel is either one

of the five higher order land cover categories or open water. The box within inset map denotes

the extent of the main map.

**Fig. 2**: Robust compositional principal components analysis (PCA) of the landscape habitat

composition surrounding each of the 40 farms monitored between 2006 and 2016. Arrows

indicate the eigenvalue and loadings of each higher order land cover. Points within the

background represent the site scores assigned to each farm and year combination and used to

define the landscape context within 500 m of nest boxes (N=440 farms*years).

**Fig. 3**: Functional relationship between Diptera biomass and the mean maximum temperature

between the day of and day prior to sample collection (A) and the first derivative values of this

relationship (B). Diptera biomass was derived with GAMMs using a Gamma distribution and a

log link function. Vertical dashed line indicates the temperature at which the maximum of the

first derivative was observed. Model controlled for, as tensor product smoothed terms, the Julian

date and the interaction between sites score values (Comp.1 * Comp.2), and the total

precipitation two days prior to insect collection (N=15,916 samples). Model also included year

and farm IDs as random effects.

**Fig. 4**: Number of cold snap days (jittered) experienced by Tree Swallow broods (N=1,897)

according to their hatching dates (1 June and 15 July are Julian 152 and 196, respectively)

between 2006 and 2016. Cold snap days were days on which maximum temperature on the farm

did not surpass 18.3°C. Point color represents the mean daily precipitation volumes (ml) that

broods experienced over a 12-day period after hatching. Black vertical line in the background is

the mean hatching date of that breeding season.

**Fig. 5.** Standardized coefficient estimates and 95% confidence intervals of the three top ranking

models, with combined $w > 0.99$ and max $\Delta$AICc $= 1.2$. Models quantified the effect of

covariates on the fledging success of Tree Swallow broods (N=1,897) raised across a gradient of

agricultural intensification within southern Québec, Canada, between 2006 and 2016. See Table

1 for acronyms and summary statistics of fixed effects and Appendix S2: Table S1 for outcome

of model selection. The GLMM with binomial response and logit link function included year,

farm, and nest box IDs as random effects. Model weights of the top three supported models

suggest they are comparable and thus are presented together.

**Fig. 6**: Unconditional model predictions (including unconditional 95% confidence intervals) of

the effect of landscape context (Comp.1 in Fig. 2) on the fledging success of Tree Swallow

broods (N=1,897) exposed to different numbers of cold snap days (days which maximum

temperature $< 18.3$°C) and, in each panel, the mean daily precipitation volumes (ml) over 12

days post hatching. Predictions were made keeping all other covariates at their mean.

**Fig. 1**

818 **Fig. 2**

821 **Fig. 3**

823 **Fig. 4**

825    **Fig. 5**

827 **Fig. 6**